



PATIENT NAME: Test, Font 10

DOB: 10/10/1910

DATE OF STUDY: 3/3/2019

FILE # : 101010

REFERRING PHYSICIAN: Dr. ReferringPhysician2 Exio

COMPARISON: None

TECHNIQUE: Axial CT images obtained of the spine with sagittal and coronal reconstructions. This is the technique section. This is the technique section. This is the technique section. This is the technique section. This is the technique section. This is the technique section. This is the technique section. This is the technique section. This is the technique section. This is the technique section.

FINDINGS: No acute fracture, dislocation or abnormal lesion is shown. There is loss of the cervical lordosis. Osteophyte is noted at C4-5, and C5-6 levels with moderate disc space narrowing. The spinal cord is normal. No Chiari malformation. No extradural soft tissue masses or paraspinal soft tissue masses. Upper thoracic spine is normal. This is the findings section. This is the findings section. This is the findings section. This is the findings section. This is the findings section. This is the findings section. This is the findings section. This is the findings section. This is the findings section. This is the findings section. The previous lines are all one paragraph and should not have any breaks. The next several lines are one line (one paragraph) per spine section.

C2-3: No disc herniation, central stenosis or neural foraminal stenosis.

C3-4: Shallow central and right paracentral disc herniation. No central stenosis or neural foraminal stenosis.

C4-5: Diffuse disc bulge and right posterior lateral osteophyte with uncovertebral joint hypertrophy and moderate bilateral neural foraminal stenosis. No central stenosis.

C5-6: Shallow central and right posterior lateral disc herniation with mass-effect along the cervical cord, right lateral recess stenosis and bilateral neural foraminal stenosis, moderate on the left and mild on the right. No central stenosis.

C6-7: No disc herniation, central stenosis or neural foraminal stenosis.

C7-T1: No disc herniation, central stenosis or neural foraminal stenosis.

IMPRESSION:

1. Moderate degenerative disc disease with loss of cervical lordosis. This is line 1 of the impression.
2. C3-4 level with shallow central and right paracentral disc herniation but no stenosis. This is line 2 of the impression.
3. C4-5 level with diffuse disc bulge, right posterior lateral osteophyte and uncovertebral joint hypertrophy with moderate bilateral neural foraminal stenosis. This is line 3 of the impression.
4. C5-6 level with shallow central and right posterior lateral disc herniation, right lateral recess stenosis and bilateral neural foraminal stenosis, moderate on the left greater than right. This is line 4 of the impression.

Thank you for the courtesy of this referral.

Sample Exio Signature

Review Exio1

Font 10 Test
3/3/2019
MRI L-Spine

Page 2 of 2

Signed: 3/3/2019 4:28:39 PM by: Review Exi01 [Electronic Signature]
Certified Radiologist of the American Board of Radiology