



Proposal for a Clinical Approach to Treat Bipolar Disorders Mixed States

Giuseppe Tavormina*

Department of Psychiatry, President of "Psychiatric Studies Center", Italy

Abstract

It is essential not to forget that the depression is only one phase of a broader bipolar mood disorder, and this has to be the illness to be treated by psychiatrists and, generally, by clinicians managing an appropriate poly therapy with mood-stabilisers and antidepressants. Very often clinicians meet great difficulties in making a correct diagnosis of mood disorders which they are assessing, above all when mixed states are present: this because the patients mainly focus on their own symptoms of depressive uneasiness; mixed symptoms can insidiously infiltrate into the mood and life of the patients causing a chronic and worsening clinical state.

Keywords: Bipolar disorders; Psychiatric; Dysphoria

Introduction

It is essential to remark once again what has been described in my previous papers: that the "instability of mood", more than the "depression", is the main issue which the clinician needs to deal with in a patient with mood disorder; this relates to the important notion, that the depressive episode is only one phase of a broader "bipolar spectrum of mood" [1-5].

When the mood quickly "swings" between depression and euphoria-irritability-hypomania or there is overlap between these conditions, we are dealing with a mixed state, even if the depressive symptoms seem to be prevalent: this then requires the clinician to carefully consider an appropriate pharmacotherapy.

Very often clinicians meet great difficulties in making a correct diagnosis of mood disorders which they are assessing, above all when mixed states are present: this because the patients mainly focus on their own symptoms of depressive uneasiness (inducing the clinicians to frequently prescribe antidepressant drugs alone or together with benzodiazepines), inducing them to prescribe these inadequate treatments and not take note of the real problem of increasing dysphoria consequent on these treatments.

The above reasons are why mixed symptoms can insidiously infiltrate into the mood and life of the patients causing a chronic and worsening clinical state.

The symptoms to note carefully on diagnosing mixed states are (at least two of these to be present at the same time) [4,5]: overlapping depressed mood and irritability, presence of agitation and restlessness, irritability and aggression and impulsivity, reduced ability to concentrate and mental over activity, high internal and muscular tension, gastritis, colitis, headaches, or other somatic symptoms (for ex: increasing of eczema or psoriasis), co-morbidity with anxiety disorders (PAD, GAD, Social phobia, OCD), insomnia (mainly fragmentary sleep and/or low quality of sleep), disorders of appetite, a sense of despair and suicidal ideation, hyper/hypo-sexual activity, substance abuse (alcohol and/or drugs), antisocial behaviour.

The "mixity" of depressive phases (that are the most insidious symptoms of overlapped depression-restlessness-irritability) can cause increased risk of suicidality [6]; besides, the co-presence of various types of somatisation symptoms [7], as well as the abuse of substances, should suggest the possibility of a "mixed state" of the bipolar spectrum [4,5].

The "G.T. Mixed States Rating Scale", or "G.T. MSRS" [5], a self-administered rating scale structured with 11 items (7 among them present also sub-items) can help the clinician to make a diagnosis of mixed state; if a patient is positive on the "G.T. MSRS", this will suggest a "generic" diagnosis for a mixed state in the bipolar spectrum, based on the full-spectrum scheme described by Akiskal [8,9] or Tavormina [2,4]. Subsequently the clinician will need to carefully make a correct

OPEN ACCESS

*Correspondence:

Giuseppe Tavormina, Department of Psychiatry, President of "Psychiatric Studies Center", Piazza Portici, 11 - 25050 Provaglio d'Iseo (BS)-Italy, Tel: 39 0309882061;

E-mail: dr.tavormina.g@libero.it

Received Date: 28 Jun 2016

Accepted Date: 10 Oct 2016

Published Date: 02 Nov 2016

Citation:

Tavormina G. Proposal for a Clinical Approach to Treat Bipolar Disorders Mixed States. *Remed Open Access*. 2016; 1: 1022.

Copyright © 2016 Tavormina G. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1: Mixed states diagnosis: percentage of evidence of the symptoms.

	Irritable Cyclothymia	Rapid cycling bipolarity	Mixed Disphoria	Agitated Depression	Cyclothymic Temperament
Euphoria with apathy	X	X	X		X
Depress. with irritability	XXX	XXX	XXX	XX	X
Substance abuse	XX	XX	XX		
Dis. of appetite		X	XX	X	
Suicidal ideation	XX	XXX	XX		
Chronic anhedonia			XX	XXX	
Delus./allucinat	X				
Hyper-hypo sexual act	XX	XX	XX	XX	
Hynsomnia/hypersomnia	XXX	XXX	XXX	XX	X
Mental overactivity	XXX	XXX	XXX	XXX	X
Somatic symptoms	XX	XX	XX	XX	X

Table 2: Mixed states diagnosis: percentage of intensity of the level of "GT-MSRS".

	Irritable Cyclothymia	Rapid cycling bipolarity	Mixed Disphoria	Agitated Depression	Cyclothymic Temperament
Level of "GT-MSRS"	High	High	High / Medium	Medium/High	Light

sub-diagnosis of the sub-groups of mixed state.

The pharmacological therapy of mixed affective states consists of a poly therapy with mood-stabilisers (mainly: lithium, carbamazepine, valproate, gabapentin, oxcarbazepine, lamotrigine, topiramate, olanzapine, asenapine, loxapine, pipamperone) and antidepressants (mainly: SSRIs, SNRIs and Bupropione) [3]. The clinician should never use antidepressants alone without mood-stabilisers when treating mixed states (or benzodiazepine alone), in order to avoid an increase in dysphoria [3,10-12].

Steps for the Clinicians

First step

It is essential at the beginning of the clinical interview to evaluate the present clinical situation which led the patient to consult the psychiatrist, and to assess what had led up to the present situation, including when the first symptoms of mood disturbance started, even though the first symptoms might have been very attenuated. In order to make a correct diagnosis of disturbance of mood within the bipolar spectrum it is essential to evaluate carefully the longitudinal psychiatric history of the patient, with particular attention to any sub-threshold symptoms and careful evaluation of the patient's temperament and the family psychiatric history [2,4,13,14].

Second step

It is essential to use the "G.T. Mixed States Rating Scale" (or "G.T. MSRS") administering it to the patient, to conduct (or not) the clinician to a "generic" diagnosis for a mixed state in the bipolar spectrum [15]. Within the full-spectrum already described in my past papers [2,4,16], the dysphoric-mixed component of unstable mood is usually present in Irritable Cyclothymia (following from, and/or developing to, rapid cycling bipolarity), in Mixed Dysphoria (typical depressive mixed state), in Agitated Depression and in the Cyclothymic Temperament.

The full-spectrum of the mood has been structured putting acute mania and unipolar depression in opposite sides of a chart, and between them all the different typologies of instability of mood, with all the fluctuations of the mood-waves, described as the following sub-types: Bipolar I, Bipolar II, Cyclothymia, Irritable Cyclothymia

Table 3: Mood stabilisers to use.

<i>Anticonvulsants</i>	Valproate
	Carbamazepine
	Gabapentin
	Oxcarbazepine
	Lamotrigine
	Topiramate
Lithium	
<i>Atypical</i>	Olanzapine
	Asenapine
	Loxapine
	Pipamperone

(or rapid cycling bipolarity), Mixed Dysphoria (or depressive mixed state), Agitated depression, three temperaments (Cyclothymic, Hyperthymic and Depressive temperament), Brief recurrent depression, and Unipolar depression [2,4].

In the Table 1 there are depicted the main symptoms (corresponding to the items of the "GT-MSRS") present in the mixed states diagnosis (data followed from the paper "Clinical utilisation of the G.T.-MSRS, [15]; furthermore, the Table 2 shows the percentage of intensity of the level of "GT-MSRS" (data followed from the same paper "Clinical utilisation of the G.T.-MSRS").

Third step

It is essential to have defined the diagnosis of the mixed state, and thence to choose the adequate mood stabiliser/s (Table 3), adding a small dosage of antidepressant in consequence of the intensity of the depressive symptoms measured both by the GT-MSRS and the clinical interview. The paper "A long term clinical diagnostic-therapeutic evaluation of 30 case reports of bipolar spectrum mixed states" Tavormina et al. [3] showed that small dosages of antidepressants (in this study the following antidepressants has been used: Escitalopram, Sertraline, Paroxetine, Venlafaxine) are important, together with

Table 4: Mixed states treatment: steps of choice of the mood stabilisers.

	1 st step	2 nd step	3 rd step	4 th step
Cyclothymic Temperament	Valproate (or Gabapentin)	Gabapentin + Valproate		
Agitated Depression	Gabapentin	Gabapentin + Valproate	Gabapentin + Valproate + Olanzapine	
Mixed Disphoria	Carbamazepine (or Valproate)	Carbamazepine (or Valproate) + Gabapentin	Carbamazepine + Valproate + Gabapentin	adding Olanzapine (or other atypical)
Rapid cycling bipolarity	Carbamazepine (or Valproate) + Gabapentin	Carbamazepine + Valproate + Gabapentin	adding Olanzapine (or other atypical)	
Irritable Cyclothymia	Carbamazepine (or Valproate) + Lithium	Carbamazepine + Valproate + Lithium	adding Olanzapine (or other atypical)	

one or more mood stabilisers, to allow a good mood balance of the patients. The presence of emotive lability, or sadness or apathy suggests the need for small dosages of antidepressants, which when in presence of mood regulators will not prolong mixed states but only reduce the emotive lability and the sadness [3].

The Table 4 shows the steps to use the mood stabilisers in the different types of mixed states. Eventual side effects, which may that force us to change drug, will induce the clinician to replace it with another drug of the same group; the paper “A long term clinical diagnostic-therapeutic evaluation of 30 case reports of bipolar spectrum mixed states” [3] described all other different properties of the safety of the mood stabilisers.

Fourth step

In consequence of the diagnosis of mixed states and their symptoms presented in the patients, the clinician has to select the mood stabiliser more indicated to be used. The high presence of severe somatic symptoms (such as colitis and gastritis) may induce the clinician to prescribe Gabapentin as first choice: Gabapentin has already been successfully used in chronic pelvic pain and irritable bowel syndrome [17,18]; Gabapentin has been used as a mood regulator with efficacy and tolerability [19,20], and in my clinical experience with efficacy and tolerability also in presence of somatic symptoms [3]. The addition of Valproate and/or Carbamazepine is useful if there is a high presence of mental overactivity and confusion with reduced ability to concentrate [3]; finally, adding also Olanzapine if there is a serious insomnia with a long term utilisation of BDZ (this, to be progressively reduced until it is discontinued, and to use subsequently only occasionally when it is necessary) [3]. It is known that all mood regulators (and above all the anticonvulsants) have to be used carefully in women of childbearing age.

Eventual side effects, which may cause us to change any kind of medication, will induce the psychiatrist to replace the drug with another one of the same group [3].

Concluding Remarks

It is essential not to forget that the depression is only one phase of a broader bipolar mood disorder, which is the fundamental illness to be treated by psychiatrists by clinicians managing an appropriate polytherapy with mood-stabilisers and antidepressants. Despite some international papers having written some reflections about the usefulness of the antidepressants on mixed states, my clinical experience [3,5] and the first study to validate the rating scale on mixed states, the “G.T. MSRS” [15], can confirm that small dosages of antidepressants added to more than one mood stabiliser will allow the patients to achieve a good mood balance when there is the presence of emotive lability or sadness.

The consequences of the lack of recognition and treatment of a mood disorder can lead to a higher risk of suicide, reduction in the

expectation and/or the quality of life (personal, family and work), increased loss of working days, increased use of health care resources, including for concurrent diseases [6,21,22]; and to the mood disorder becoming chronic and the clinical picture becoming worse.

Future studies are in project to give additional details and scientific evidence on these points.

References

1. Akiskal HS. The prevalent clinical spectrum of bipolar disorders: beyond DSM-IV. *J Clin Psychopharmacol.* 1996; 16: 4-14.
2. Akiskal HS, Pinto O. The evolving bipolar spectrum: Prototypes I, II, III, IV. *Psychiatr Clin North Am.* 1999; 22: 517-534.
3. Akiskal HS. Targeting suicide prevention to modifiable risk factors: has bipolar II been overlooked? *Acta Psychiatr Scand.* 2007; 116: 395-402.
4. Agius M, Tavormina G, Murphy CL, Win A, Zaman R. Need to improve diagnosis and treatment for bipolar disorder. *Br J Psych.* 2007; 190: 189-191.
5. Agius M, Zaman R, Ho J, Singh S. Mixed affective states: a study within a community mental health team with treatment recommendations. *Psychiatria Danubina.* 2011; 23: 25-28.
6. Carta MG, Hardoy MC, Hardoy MJ, Grunze H, Carpiniello B. The clinical use of gabapentin in bipolar spectrum disorders. *J Affect Dis.* 2003; 75: 83-91.
7. Carta MG, Cabras PL, Hardoy MJ, Hardoy MC. Clinical Experience with gabapentin in patients with Bipolar or Schizoaffective Disorder. *J Clin Psychiatry.* 1999; 60: 245-248.
8. Lee K J, Kim JH, Cho SW. Gabapentin reduces rectal mechanosensitivity and increases rectal compliance in patients with diarrhoea-predominant irritable bowel syndrome. *Alimentary Pharmacology and Therapeutics.* 2005; 22: 981-988.
9. Sator-Katzenschlager SM, Scharbert G, Kress HG, Frickey N, Ellend A, Gleiss A, et al. Chronic pelvic pain treated with gabapentin and amitriptyline: a randomized, controlled pilot study. *Wien Klin Wochenschr.* 2005; 117: 761-768.
10. Tavormina G, Agius M. The high prevalence of the bipolar spectrum in private practice. *J Bipolar Dis: Rev & Comm.* 2007; 3: 19.
11. Tavormina G, Agius M. A study of the incidence of bipolar spectrum disorders in a private psychiatric practice. *Psychiatria Danubina.* 2007; 19: 370-374.
12. Rihmer Z, Akiskal HS, Rihmer A, Akiskal HS. Current research on affective temperaments. *Current Opinion in Psychiatry.* 2010; 23: 12-18.
13. Tavormina G. The bipolar spectrum diagnosis: the role of the temperaments. *Psychiatria Danubina.* 2009; 21: 160-161.
14. Tavormina G. The temperaments and their role in early diagnosis of bipolar spectrum disorders. *Psychiatria Danubina.* 2010; 22: 15-17.
15. Tavormina G. The temperaments: its knowledge is a crucial way in early diagnosis of bipolar disorders. *European Psychiatry.* 2011; 26: P01-199.

16. Agius M, Zaman R. Mixed affective states: a study within a community mental health team with treatment recommendations. *European Psychiatry*. 2011; 26: P01-145.
17. Tavormina G. Are somatisations symptoms important evidence for an early diagnosis of bipolar spectrum mood disorders? *Psychiatria Danubina*. 2011; 23: 13-14.
18. Tavormina G, Agius M. An approach to the diagnosis and treatment of patients with bipolar spectrum mood disorders, identifying temperaments. *Psychiatria Danubina*. 2012; 24: 25-27.
19. Tavormina G. An early diagnosis of bipolar spectrum disorders needs of valuing the somatisation symptoms of patients. *J Int Clin Psychopharmacol*. 2012; 28: e59-e60.
20. Tavormina G. A long term clinical diagnostic-therapeutic evaluation of 30 case reports of bipolar spectrum mixed states. *Psychiatria Danubina*. 2013; 25: 190-193.
21. Tavormina G. An introduction to the bipolar spectrum. The management of bipolar spectrum disorders. Summer. 2013; CEPiP: 3-6.
22. Tavormina G. Treating the bipolar spectrum mixed states: a new rating scale to diagnose them. *Psychiatria Danubina*. 2014; 26: 6-9.
23. Tavormina G. Clinical utilisation of the "G.T. MSRS", the rating scale for mixed states: 35 cases report. *Psychiatria Danubina*. 2015; 27: 155-159.