

Use of Output from the New England SPARROW Model to Estimate Concentrations of Total Nitrogen in Estuaries

Edward H. Dettmann¹, Richard B. Moore², Keith W. Robinson²,
Henry A. Walker¹, and Jaime B. Palter³

¹U.S. Environmental Protection Agency (USEPA), ORD,
National Health and Environmental Effects Research Laboratory,
Atlantic Ecology Division, Narragansett, Rhode Island

²U.S. Geological Survey (USGS), New Hampshire/Vermont District,
Pembroke, New Hampshire

³Duke University, Department of Environment and Earth Sciences,
Division of Earth and Ocean Sciences, Durham, North Carolina

EMAP Symposium, May 3-7, 2004, Newport, Rhode Island



Purpose of This Work

- Link USEPA Estuary Nitrogen Model (ENM) to SPARROW Model for Application to:
 - National Coastal Assessment
 - Development of TMDLs for Estuaries
 - Development of Nutrient Criteria for Estuaries

Presentation Outline

Background

Description of Estuary Nitrogen Model (ENM)

Discuss Compatibility of SPARROW Model & ENM

Application of SPARROW & ENM to:

- o Narragansett Bay
- o Boston Harbor
- o Great Bay Estuary

Summary

The Estuary Nitrogen Model.

Dettmann (2001)

$$\frac{dN}{dt} = L_{land} + L_{sea} - E - \alpha N$$

N Loading

Export **Internal Losses**

The diagram shows the differential equation for nitrogen change in an estuary. The equation is $\frac{dN}{dt} = L_{land} + L_{sea} - E - \alpha N$. Above the equation, the text "N Loading" is written in blue. Two blue arrows point downwards from "N Loading" to the L_{land} and L_{sea} terms. Below the equation, the text "Export" and "Internal Losses" are written in blue. Two blue arrows point upwards from "Export" to the $-E$ term, and two blue arrows point upwards from "Internal Losses" to the $-\alpha N$ term.

Assumptions:

Model deals with long-term (e.g. annual or multi-year averages).

Approximate steady state at scale of yearly cycle, i.e.

$$\frac{dN}{dt} = 0$$

The Estuary Nitrogen Model.

Dettmann (2001)

Final Equations:

$$F_{ex} = \frac{1}{1 + \alpha \tau}$$

$$[N] = \left(\frac{L_{land} \tau}{V} + [N_{sea}] \right) \frac{1}{1 + \alpha \tau}$$

τ = freshwater residence time

$\alpha \simeq 0.3 \text{ mo}^{-1}$ (*nonlinear least squares estimate for 11 estuaries*)

Nitrogen Budgets for 11 Estuaries. Internal N Losses are Function of Residence Time

Denitrification

Export




Fig. 2. The fraction of upland nitrogen input that is exported from 11 estuaries versus freshwater residence time (logarithmic time scale).





Fig. 3. The fraction of upland nitrogen input that is denitrified versus freshwater residence time. The solid model line ($\gamma = 0.69$, $r^2 = 0.85$) is the fit to all the data. The dashed model lines correspond to the 95% confidence limits for γ (0.57 and 0.80). If Chesapeake Bay is excluded from the fit (see text), the model line ($\gamma = 0.76$, $r^2 = 0.97$) lies between those for $\gamma = 0.69$ and $\gamma = 0.80$.

Predicted vs. Observed [TN] for 17 Side-Embayments of Buzzards Bay



Characteristics of Nitrogen Output from Sparrow Model

Annual Loads Only

TN Only (No Components)

Loads Only to Nontidal Streams

Loads Can Be Partitioned by Source Category

Sources of Riverine TN Loading

≈ 68% of loading to Narragansett Bay

	<u>% TN / yr</u>
• Atmospheric:	17.4%
• Urban:	18.4%
• Agriculture:	2.6%
• Point Sources:	61.2%

*(point sources discharging directly
into estuary not included)*

Summary of Data Requirements of the Estuary Nitrogen Model

Annual Loads of Total Nitrogen to Estuary from:


- o Watershed
- o Atmosphere
- o Point sources

Average Annual Freshwater Residence Time (τ)

Estuary Volume

Background Nitrogen Concentration from Transport
Across Seaward Boundary ($[N_{\text{sea}}]$)

Estimated TN Input to Narragansett Bay from Rivers & Streams (NE SPARROW Model)



TN Loading to Narragansett Bay


	<u>kg N y⁻¹</u>
Sparrow (30 tributaries)	6,227,261
Nixon et al. (1995)	6,120,928
TN loading from SPARROW	6,227,261
Direct Atmospheric Deposition*	420,201
<u>Sewage Treatment Plants*</u>	<u>2,563,226</u>
Total TN Loading	9,210,688

*(Nixon et al., 1995)

Riverine TN loading to Narragansett Bay from New England SPARROW Model is 68% of total.

Building a scientific foundation for sound environmental decisions

Measured [TN] (1985—1986 SINBADD Cruises)



Sakonnet River excluded from calculations

Figure adapted from Hunt et al. (1987a)

Average TN Concentration in Narragansett Bay

$$L_{\text{land}} = 766,766 \text{ kg N mo}^{-1}$$

$$\tau = 26 \text{ d} = 0.855 \text{ mo}$$

$$V = 2.821 \times 10^9 \text{ m}^3$$

$$V_{\text{sw}} = 2.584 \times 10^9 \text{ m}^3$$

$$[N_{\text{b}}] = 0.201 \text{ mg L}^{-1}$$

$$[N_{\text{sea}}] = 0.184 \text{ mg L}^{-1}$$

$\alpha = 0.3 \text{ mo}^{-1}$ (*permanent loss to denitrification & burial*)

$$[N] = \left(\frac{L_{\text{land}} \tau}{V} + [N_{\text{sea}}] \right) \frac{1}{1 + \alpha \tau}$$


Model-Calculated [TN] = $(0.232 + 0.185)/1.2565 = 0.332 \text{ mg L}^{-1}$

Measured [TN] (1985—1986 SINBADD Cruises)* = 0.358 mg L^{-1}

* (estimate, based on weighted average of TN, (Hunt et al., 1987))

Calculated [TN] is within 7.3% of measured concentration.

Estimated TN Input to Boston Harbor from Rivers & Streams (NE SPARROW Model)



TN Loading to Boston Harbor Early 1990s (prediversion)

	<u>kg N y⁻¹</u>
Sparrow (all tributaries)	1,305,245
All rivers and urban runoff *	893,000
TN loading from SPARROW	1,305,245
Direct Atmospheric Deposition*	307,000
Sewage Treatment Plants*	11,350,000
Groundwater*	93,000
<hr/>	
Total TN Loading	13,055,345

*Alber and Chan (1994)

Riverine TN loading to Boston Harbor from New England
SPARROW Model is 10% of total.

Complicating Factors

Largest source of TN is on the seaward boundary.

Residence time for TN discharged by Deer Island WWTF is smaller than that of Boston Harbor as a whole.

Average TN Concentration in Boston Harbor (prediversion)

$$L_{land} = 108,770 \text{ kg N mo}^{-1}$$

$$V = 612.5 \times 10^6 \text{ m}^3$$

$$V_{sw} = 603.9 \times 10^6 \text{ m}^3$$

$$[N_b] = 0.233 \text{ mg L}^{-1}$$

$$[N_{sea}] = 0.230 \text{ mg L}^{-1}$$

$$\tau = 10 \text{ d} = 0.33 \text{ mo}$$

$$\alpha = 0.3 \text{ mo}^{-1}$$

Calculated [TN] (model with full loading
and $\tau = 10 \text{ d}$ for Deer I. effluent) = 0.745 mg L⁻¹

Calculated [TN] (model with reduced loading
and $\tau = 5 \text{ d}$ for Deer I. effluent) = 0.481 mg L⁻¹

Measured [TN] (mean of MWRA data) = 0.475 mg L⁻¹

Calculated [TN] is within 1.3% of measured concentration.

Average TN Concentration in Boston Harbor (postdiversion)

$$L_{land} = 142,104 \text{ kg N mo}^{-1}$$

$$V = 612.5 \times 10^6 \text{ m}^3$$

$$V_{sw} = 603.9 \times 10^6 \text{ m}^3$$

$$[N_b] = 0.233 \text{ mg L}^{-1}$$

$$[N_{sea}] = 0.230 \text{ mg L}^{-1}$$

$$\tau = 10 \text{ d} = 0.33 \text{ mo}$$

$$\alpha = 0.3 \text{ mo}^{-1}$$

Calculated [TN] (model with no loading
from WWTFs) =


0.295 mg L⁻¹

Measured [TN] (rough estimate of mean
of MWRA data) =

0.280 mg L⁻¹

This is within 5.4% of measured concentration.

Estimated TN Input to Great Bay Estuary from Rivers & Streams (NE SPARROW Model)



TN Loading to Great Bay Piscataqua Estuary

	<u>kg N y⁻¹</u>
Sparrow (all tributaries)	1,268,612
TN loading from SPARROW	1,268,612
Direct Atmospheric Deposition*	69,853
Sewage Treatment Plants into Estuary*	<u>172,982</u>
Total TN Loading	1,511,447

*Jones (2000)

Riverine TN loading to Great Bay/Piscataqua Estuary from
New England SPARROW Model is 84% of total.

Average Calculated [TN] in Great Bay/Piscataqua Estuary

$$L_{land} = 125,954 \text{ kg N mo}^{-1}$$

$$V = 198 \times 10^6 \text{ m}^3 *$$

$$V_{sw} = x \times 10^6 \text{ m}^3 *$$

$$\alpha = 0.3 \text{ mo}^{-1}$$

$$[N_b] = 0.19 \text{ mg L}^{-1}$$

$$[N_{sea}] = 0.16 \text{ mg L}^{-1}$$

$$\tau = 22 \text{ d} = 0.72 \text{ mo} *$$

For lower estuary,
 $\tau = 1 \text{ d} = 0.033 \text{ mo} *$

* Brown & Arellano (1980)

$$\text{Calculated [TN]} = (0.001 + 0.354 + 0.133) \text{ mg L}^{-1} = 0.488 \text{ mg L}^{-1}$$

Portsmouth & Kittery

All Other Loads

Background



Summary & Conclusions

- SPARROW Model TN loads are directly useable by the Estuary Nitrogen Model
- SPARROW loads must be supplemented with other loads to the estuary (e.g. point sources, atmospheric deposition, etc.)
- Three tests of this approach calculated TN concentrations that agree to within a few percent with measured values

Summary & Conclusions (cont.)

- This approach calculates annual average nitrogen concentrations, although it may be possible to infer seasonal values
- This approach readily permits analysis of the relative magnitudes of nitrogen loading from individual source classes

Acknowledgment


- Wendy Leo - MWRA

References

- Alber, M and AB Chan. 1994. Sources of contaminants to Boston Harbor: Revised loading estimates. Massachusetts Water Resources Authority, Environmental Quality Department Technical Report Series No. 94-1, Boston, Massachusetts.
- Brown, WS & E Arellano. 1980 The application of a segmented tidal mixing model to the Great Bay Estuary, N.H. *Estuaries* 3(4):248-257.
- Dettmann, EH 2001. Effect of water residence time on annual export and denitrification of nitrogen in estuaries: A Model analysis. *Estuaries* 24(4):481-490.
- Hunt, CD et al. 1987a-d. Narragansett Bay water quality monitoring and source strength measurements (Four volumes describing SINBADD Cruises 1-4). Marine Ecosystems Research Laboratory, Graduate School of Oceanography, University of Rhode Island, Narragansett, RI 02882.
- Jones, SH. 2000. A technical characterization of estuarine and coastal New Hampshire. New Hampshire Estuaries Project.
- Nixon, SW, SL Granger, & BL Nowicki. 1995. An assessment of the annual mass balance of carbon, nitrogen, and phosphorus in Narragansett Bay. *Biogeochemistry* 31:15-61.


Backup Slides

Seasonal TN Concentration Ratios



Building a scientific foundation for sound environmental decisions

Seasonal DIN Concentration Ratios




Building a scientific foundation for sound environmental decisions



Predicted Nitrogen Yield

New England SPARROW Model


Sources of Nitrogen




Base from U.S. Geological Survey Digital line graph, 1:2,000,000 scale, 1990-94




Atmospheric



Urban



Agriculture



Point Source