# \*\*\*DRAFT\*\*

## Fire Regime Condition Class (FRCC) Interagency Handbook Reference Conditions

Modeler: Ron Masters Date: 5/28/04 PNVG Code: SEOK4

Potential Natural Vegetation Group: Xeric Pine-Oak Woodlands, Western

**Geographic Area:** Interior highlands and uplands of Arkansas, eastern Oklahoma, southern Missouri and uplands of east Texas.

**Description**: Potential natural vegetation group common to the Interior Highlands and xeric upland sites to the south and west of the Mississippi River to east Texas. In Highlands occupying all but steep north slopes at all elevations. In the northern part of this geographic area shortleaf pine (*Pinus echinata*), xeric oaks and some hickory dominate the overstory with a high percentage of oak on steep north slopes and on post oak (*Quercus stellata*) flats. In the southern part loblolly pine (*P. taeda*) is an associate. Associated species include post oak, blackjack oak (*Quercus marylandica*), mockernut hickory (*Carya tomentosa*) on drier sites and to the west black hickory (*Carya texana*). Pine often emergent on upper slopes. Stand density increases with available moisture. Various bluestems often dominate the understory.

**Fire Regime Description:** Fire regime group I, with frequent surface fires. Area fire frequency 3-4 year mean fire interval (range=1-12 years) from Masters et al. (1995).

Class*	Percent of	Description		
	Landscape			
A: post replacement	17	Pine and oak reproduction to 15' tall. Herbaceous community dominated by bluestems and forbs. More persistent on shallow soils. Openings may be small to extensive and have scattered live trees.		
<b>B</b> : mid-seral closed	6	Mid-seral with closed canopy (>70%; on mountainous sites >60%) shortleaf and loblolly pine and pole-sized oak with little or no herbaceous understory.		
C: mid- seral open	33	Mid-seral open woodland/savanna pine and oak overstory with bluestem grasses and forbs. Shrub layer may be prevalent on some sites and dominated by various oak sprouts and a few shrub species. Prevalence highly dependant on time since burned. Cover <70%; on mountainous sites cover <60%.		
D: late- seral open	40	Late-seral woodland/savanna pine and oak overstory with bluestem grasses and forbs. Shrub layer may be prevalent on some sites and dominated by various oak sprouts and a few shrub species. Prevalence highly dependant on time since burned. Shrub layer may be absent on other sites, particularly on shallow soils. Cover <70%; on mountainous sites cover <60%.		
E: late- seral closed	5	Late-seral, closed canopy (>70%; on mountainous sites >60%) pine-oak dominated overstory community. No herbaceous cover and few shrubs.		
Total	100			

#### **Vegetation Type and Structure**

\*Formal codes for classes A-E are: AESP, BMSC, CMSO, DLSO, and ELSC, respectively.

#### **Fire Frequency and Severity**

	Fire Frequency	Probability	Percent,	Description
Fire Severity	(yrs)	-	All Fires	-
Replacement Fire	104	.01	4	Late growing season fire that occurs in drought years
Non-Replacement Fire	4.5	.22	96	Primarily surface fire in all classes. Some mosaic fire.
All Fire Frequency*	4.3	.23	100	

\*All Fire Probability = sum of replacement fire and non-replacement fire probabilities. All Fire Frequency = inverse of all fire probability (previous calculation).

#### References

Albert, Lois E. 1981. Five Thousand Years of Environmental Change in Southeastern Oklahoma. Okla. Arch. Survey, No. 7;.

Bragg, D. C. 2002. Reference conditions for old-growth pine forests in the Upper West Gulf Coastal Plain. Journal of the Torrey Botanical Society 129(4):261-288.

Brissette, J. C. and J. P. Barnett (eds). 1992. Proceedings: Shortleaf Pine Regeneration Workshop. So. For. Exp. Sta., GTR-SO-90.

Burger, George V., et al. (eds). Proceedings of the Oak Woods Management Workshop. E. III. Univ.; 1991.

Brown, J. K. and J. K. Smith editors. Wildland Fire in Ecosystems: effects of fire on flora. Gen, Tech. Rep. RMRS-GTR-42-vol. 1-2. USDA, Forest Service, Rocky Mountain Research Station. 2000.

Cain, M. D. and M. G. Shelton. 2003. Effects of Alternative Thinning Regimes and Prescribed Burning in Natural, Even-aged Loblolly-Shortleaf Pine Stands: 25 year results. Southern Journal of Applied Forestry 27(1):

Cain, M. D. and M. G. Shelton. 2000. Survival and growth of Pinus and Quercus seedlings in response to simulated summer and winter prescribed burns. Canadian Journal of Forest Resources 30:.

Campbell, J.J.N., et al. 1991. Floristic and Historical Evidence of Fire Maintained, Grassy Pineoak Barrens before Settlement in Southeastern Kentucky. In Proc: Fire and the Environment: Ecological and Cultural Perspectives. SE For. Exp. Sta.

Cram, D. S., R. E. Masters, F. S. Guthery, D. M. Engle, and W. G. Montague. 2002. Northern Bobwhite Population and Habitat Response to Pine-grassland Restoration. Journal of Wildlife Management 66:1031-1039.

Delcourt, H. R. and P. A. Delcourt. 1991. Late Quaternary Vegetation History of the Interior Highlands of Missouri, Arkansas, and Oklahoma. In D. Henderson and L. D. Hedrick, editors. Proc: Restoration of Old Growth Forests of the Interior Highlands of Arkansas and Oklahoma. Winrock International. Morrilton, Ark.

Foti, T. and S. Glenn. 1991. The Ouachita Mountains Landscape at the Time of Settlement. In D. Henderson and L. D. Hedrick, editors. Proc.: Conference on Restoring Old Growth Forest in the Interior Highlands of Arkansas and Oklahoma. Winrock International, Morrilton, Ark.

Fryar, Roger D. 1991. Old Growth Stands of the Ouachita National Forest. In D. Henderson and L. D. Hedrick, editors. Proc: Restoration of Old Growth Forest in the Interior Highlands of Arkansas and Oklahoma. Winrock International. Morrilton, Ark.

Glitzenstein, J. S., P. A. Harcomb, and D. R. Streng. 1986. Disturbances, succession, and maintenance of species diversity in an east Texas forest. Ecological Monographs 56:243-258.

Guyette, R. P. and E. A. McGinnes, Jr. 1982. Fire History of an Ozark Glade in Missouri. Trans. Mo. Acad. Sci.16:85-93.

Guyette, R. P and B. E. Cutter. 1991. Tree-ring analysis of fire history of a post-oak savanna in the Missouri Ozarks. Natural Areas Journal 11(2): 93-99.

Guyette, R. P. and M. A. Spetich 2003. Fire history in oak-pine forests in the Lower Boston Mountains, Arkansas, USA. Forest and Ecology Management 180:463-474.

Henderson, D. and L. D. Hedricks (eds). 1991. Restoration of Old Growth Forests in the Interior Highlands of Arkansas and Oklahoma. Conf. Proc., Winrock International. Morrilton, Ark.

Hessl, A. and S. Spackman. 1995. Effects of Fire on Threatened and Endangered Plants: An annotated Bibliography. USDOI, National biological Service.

Honess, C. W. 1923. Geology of the southern Ouachita Mountains of Oklahoma. Bulletin 32, Parts I and II. Oklahoma Geological Survey. Norman Okla. 354 pp.

Jansma, J. and H. H, Jansma. George Engelmann in Arkansas Territory. Ark. Hist. Quart., pp. 225-248.

Johnson, F. L. and G. D. Schnell. 1985. Wildland Fire History and the Effects of Fire on Vegetative Communities at Hot Springs National Park, Arkansas. Rep. To NPS, Santa Fe, NM., Okla. Biological Survey., University of Oklahoma, Norman, Okla. 49 pp.

Jurney, D. R. Evans, J. Ippolito, and V. Bergstrom. 2004. The role of Wildland fire in portions of southeastern North America. Pages IN PRESS in R. T. Engstrom and W. J. de Groot (eds). 22<sup>nd</sup> Tall Timbers Fire Ecology Conf. Proceedings. Kanaskas, Alberta.

Kreiter, S. D. 1994. Dynamics and Spatial Patterns of a Virgin Old-growth Hardwood-pine Forest in the Ouachita Mountains, Oklahoma, from 1896-1994. M.S. Thesis. Oklahoma State University, Stillwater.

Langevede, F. V., C. A. D. M. Vande Vijver, L. Kumar, H. Van De Koppel, N. De Ridder, J. Van Andel, A. K. Skidmore, J. W. Hearne, L. Stroosnijder, W. J. Bond, H. H. T. Prins, and M. Rietkerk. 2003. Effects of Fire and Herbivory on the Stability of Savanna Ecosystems. Ecology 84(2):337-350.

Lewis, A. 1924. La Harpe's first expedition in Oklahoma, 1718-1719. Chron. Oklahoma 2(4):331-349.

MacCleery, D. 1994. Understanding the Role the Human Dimension Played in Shaping America's Forest and Grassland Landscapes: Is there a landscape archaeologist in the house? Eco-watch 2: .

Masters, R. E. 1991. Effects of timber harvest and prescribed fire on wildlife habitat and use in the Ouachita Mountains of eastern Oklahoma. Ph.D. Thesis, Oklahoma State Univ. Stillwater. 351 pp.

Masters, R. E., J. E. Skeen, and J. A. Garner. 1989. Red-cockaded woodpecker in Oklahoma; an update of Wood's 1974-77 Study. Proc. Okla. Acad. Sci. 69:27-31.

Masters, R. E., J. E. Skeen, and J. Whitehead. 1995. Preliminary fire history of McCurtain County Wilderness Area and implications for red-cockaded woodpecker management. Pages 290-302 *in* D. L. Kulhavy, R. G. Hooper, and R. Costa. (eds.). Red-cockaded woodpecker: Species recovery, ecology and management. Center for Applied Studies, Stephen F. Austin University, Nacogdoches, TX.

Masters, R. E. 1991. Effects of fire and timber harvest on vegetation and cervid use on oak -pine sites in Oklahoma Ouachita Mountains. Pages 168-176. In S. C. Nodvin and T. A. Waldrop, (eds.). Fire and the environment: ecological and cultural perspectives. Proc. of an international symposium. USDA For. Serv. Gen. Tech. Rep. SE-69. Southeast For. Exp. Sta., Asheville, N.C.

Masters, R. E., and D. M. Engle. 1994. BEHAVE-evaluated for prescribed fire planning in mountainous oak-shortleaf pine habitats. Wildlife Society Bulletin 22:184-191.

Masters, R. E., D. M. Engle, and R. Robinson. 1993. Effects of timber harvest and periodic fire on soil chemical properties in the Ouachita Mountains. Southern Journal of Applied Forestry 17:139-145.

Masters, R. E., R. L. Lochmiller, and D. M. Engle. 1993. Effects of timber harvest and periodic fire on white-tailed deer forage production. Wildlife Society Bulletin 21:401-411.

Masters, R. E., R. L. Lochmiller, S. T. McMurry, and G. A. Bukenhofer. 1998. Small mammal response to pine-grassland restoration for red-cockaded woodpeckers. Wildlife Society Bulletin 28:148-158.

Masters, R. E., C. W. Wilson, G. A. Bukenhofer, and M. E. Payton. 1996. Effects of pinegrassland restoration for red-cockaded woodpeckers on white-tailed deer forage production. Wildlife Society Bulletin 24:77-84.

Masters, R. E., C. W. Wilson, D. S. Cram, G. A. Bukenhofer, and R. L. Lochmiller. 2002. Influence of ecosystem restoration for red-cockaded woodpeckers on breeding bird and small mammal communities. Pages 73-90 in W. M. Ford, K. R. Russell, and C. E. Moorman, editors. In The role of fire in non-game wildlife management and community restoration: traditional uses and new directions: proceedings of a special workshop. Annual Meeting of The Wildlife Society, Nashville, Tenn. USDA For. Ser. Northeast Research Station. General Technical Report NE-288.

Nelson, J. C. 1997. Presettlement Vegetation Patterns along the 5<sup>th</sup> Principal Meridian, Missouri Territory, 1815. Am. Midl. Nat. 137:70-94.

Palmer, E. J. 1921. The Forest Flora of the Ozark Region. J. Arnold Arbor. 2:.

Palmer, E. J. 1924. The Ligneous Flora of Rich Mountain, Arkansas and Oklahoma. J. Arnold Arbor. 5:

Panzer, R.. Compatibility of Prescribed Burning with Conservation of Insects in Small, Isolated Prairie Preserves. Conservation Biology, Vol. 16, no. 5, 2002, pp1296-1307.

Perttula, T. K. 1993. The Long Term Consequences and Effects of the de Soto Entrada on Aboriginal Caddoan Populations. In Proc: De Soto Sym, 1988 and 1990;

Rebertus, A. J. and B. R. Burns. 1997. The Importance of gap processes in the development and maintenance of oak savannas and dry forests. Journal of Ecology 85:633-645.

Rudis, Victor A. and Thomas V. Skinner. Fire's Importance in South Central U.S. Forests: distribution of fire evidence.

Smith, B. A., R. J. Tyrl, and R. E. Masters. 1997. Floristic inventory of the McCurtain County Wilderness Area (Oklahoma). Okla. Acad. Sci. 77:99-102.

Smith, K. L. 1986. Sawmill: The story of cutting the last great virgin forest east of the Rockies. Univ. Ark. Press.

Smith, K. G. and J. C. Neal. 1992. Pre-settlement Birds and Mammals of the Interior Highlands. In D. Henderson and L. D. Hedrick, editors. Proc.: Conference on Restoring Old Growth Forest in the Interior Highlands of Arkansas and Oklahoma. Winrock International, Morrilton, Ark.

Sparks, J. C. 1996. Growing-Season and Dormant-Season Fire Behavior and Effects on Vegetation in the Ouachita Mountains, Arkansas. M.S. Thesis. Oklahoma State University, Stillwater. 186 pp.

Sparks, J. C., and R. E. Masters. 1996. Fire seasonality effects on vegetation in mid-, tall-, and southeastern pine-grassland communities: a review. Trans. No. Am. Wildlife and Natur. Res. Conf. 61:230-239.

Sparks, J. C., R. E. Masters, D. M. Engle, and G. A. Bukenhofer. 2002. Season of burn influences fire behavior and fuel consumption in restored shortleaf pine-grassland communities. Restoration Ecology 10:714-722.


Sparks, J. C., R. E. Masters, D. M. Engle, M. Palmer and G. A. Bukenhofer. 1998. Effects of late growing-season and late dormant-season prescribed fire on herbaceous vegetation in restored pine-grassland communities. Journal of Vegetation Science 9: 133-142.


Williams, G. W. 2002. Aboriginal Use of Fire: are there any "natural " plant communities?. Wilderness and Political Ecology: Aboriginal Land Management-Myths and Reality (Charles E. Kay and Randy T. Simmons (eds). Univ. of Utah Press.


Wilson, C. W., R. E. Masters, and G. A. Bukenhofer. 1995. Breeding bird response to pinegrassland community restoration for red-cockaded woodpeckers. Journal of Wildlife Manage. 59:56-67.

### **VDDT File Documentation**

Include screen captures (print-screens) from any of the VDDT graphs that were used to develop reference conditions.









Reference conditions circa 1900.

Reference conditions circa 1925.




Reference conditions circa 1920, near Beavers Bend, Oklahoma (from Honess 1923).



A. Post-replacement – early.



A. Post-replacement – early.



B. Mid-seral closed.



B. Mid-seral closed.



C. Mid-seral open.



C. Mid-seral open.



D. Late-seral open.



D. Late-seral open.



E. Late-seral closed

E. Late-seral closed.



E. Late-seral closed landscape, Ouachita Highlands.