such use is in accordance with good agricultural or manufacturing practices.

Acetic acid, sodium salt	127–09–3 None 9000–40–2
Carob gum (locust bean gum) Castor oil Castor oil, Castor oil, Cellulose Cellulose acetate	
Castor oil	9000-40-2
Castor oil, hydrogenated	
Cellulose	8001-79-4
Cellulose acetate	8001-78-3
	9004-34-6
	9004-35-7
Cellulose, carboxy methyl ether, sodium salt	9004-32-4
Cellulose, 2-hydroxyethyl ether	9004-62-0
Cellulose, 2-hydroxypropyl ether	9004-64-2
Cellulose, 2-hydroxypropyl methyl ether	9004-65-3
Cellulose, methyl ether	9004-67-5
Cellulose, mixture with cellulose carboxymethyl	
ether, sodium salt	51395-75-6
Cellulose, pulp	65996-61-4
Cellulose, regenerated	68442-85-3
Citric acid	77-92-9
Citric acid, calcium salt	7693-13-2
Citric acid, calcium salt (2:3)	813-94-5
Citric acid, dipotassium salt	3609-96-9
Citric acid, disodium salt	144-33-2
Citric acid, monohydrate	5949-29-1
Citric acid, monopotassium salt	866-83-1
Citric acid, monosodium salt	18996-35-5
Citric acid, potassium salt	7778-49-6
Citric acid, tripotassium salt	866-84-2
Citric acid, tripotassium salt, monohydrate	6100-05-6
Citric acid, sodium salt	994-36-5
Citric acid, trisodium salt	68-04-2
Citric acid, trisodium salt, dihydrate	6132-04-3
Citric acid, trisodium salt, pentahydrate	6858-44-2
Coffee grounds	68916-18-7
Dextrins	9004-53-9
Fumaric acid	110-17-8
Guar gum	9000-30-0
Lactic acid, n-butyl ester	138-22-7
Lactic acid, ethyl ester	97-64-3
Lanolin	8006-54-0
Lecithins	8002-43-5
Lecithins, soya	8030-76-0
Licorice Extract	68916-91-6
Maltodextrin	9050-36-6
Paper	None
Potassium chloride	7447-40-7
Silica, amorphous, fumed (crystalline free)	112945-52-
	5
Silica, amorphous, precipitated and gel	7699-41-4
Silica gel	63231-67-4
Silica gel, precipitated, crystalline-free	112926-00-
•	8
Silica, hydrate	10279-57-9
Silica, vitreous	60676-86-0
Soapbark (Quillaja saponin)	1393-03-9
Sodium alginate	9005-38-3
Codium oblarida	7647-14-5
Sodium chloride	57-13-6
Urea	
	11138-66-2

[67 FR 36537, May 24, 2002, as amended at 67 FR 56229, Sept. 3, 2002; 67 FR 78718, Dec. 26, 2002; 68 FR 16437, Apr. 4, 2003; 68 FR 18552, Apr. 16, 2003]

$\S\,180.960$ Polymers; exemptions from the requirement of a tolerance.

Residues resulting from the use of the following substances, that meet the definition of a polymer and the criteria specified for defining a low-risk polymer in 40 CFR 723.250, as an inert ingredient in a pesticide chemical formulation, including antimicrobial pesticide chemical formulations, are exempted from the requirement of a tolerance under FFDCA section 408, if such use is in accordance with good agricultural or manufacturing practices.

manaracearing practices:	
Polymer	CAS No.
Acetic acid ethenyl ester, polymer with ethenol and (α)-2-propenyl-(ω)-hydroxypoly (oxy-1,2-ethanediyl) minimum number average molecular weight (in amu), 15,000	137091– 12–4
Acrylic acid, polymerized, and its ethyl and methyl esters	None
Acrylic acid-sodium acrylate-sodium-2- methylpropanesulfonate copolymer, minimum average molecular weight (in amu), 4,500	97953– 25–8
Acrylic acid-stearyl methacrylate copolymer, min- imum number average molecular weight (in amu), 2,500	27756– 15–6
Acrylic acid, styrene, α-methyl styrene copolymer, ammonium salt, minimum number average mo- lecular weight (in amu), 1,250	89678– 90–0
Acrylic acid terpolymer, partial sodium salt, min- imum number average molecular weight (in amu), 2,400	151006– 66–5
Acrylic polymers composed of one or more of the following monomers: Acrylic acid, methyl acrylate, ethyl acrylate, butyl acrylate, hydroxyethyl acrylate, hydroxyptopyl acrylate, hydroxyptopyl acrylate, hydroxybutyl acrylate, carboxyethyl acrylate, methacrylate, butyl methacrylate, ethyl methacrylate, butyl methacrylate, isobutyl methacrylate, hydroxyptopyl methacrylate, hydroxyptopyl methacrylate, hydroxyptopyl methacrylate, isobutyl methacrylate, lauryl methacrylate, and stearyl methacrylate; with none and/or one or more of the following monomers: Acrylamide, N-methyl acrylamide, N,N-dimethyl acrylamide, maleic anhydride, maleic acid, monoethyl maleate, and their corresponding sodium, potassium, ammonium, isopropylamine, triethylamine, monoethanolamine, and/or triethanolamine salts; the resulting polymer having a minimum number average molecular weight (in amu), 1,200	None
Acrylonitrile-butadiene copolymer conforming to 21 CFR 180.22, minimum average molecular weight (in amu), 1,000.	9003–18– 3
Acrylonitrile-styrene-hydroxypropyl methacrylate copolymer, minimum number average molecular weight (in amu), 447,000	None

Environmental Protection Agency

Polymer	CAS No.
$\alpha\text{-alkyl}$ $(C_{12}\text{-}C_{15})$ - $\omega\text{-hydroxypol-y(oxypropylene)poly(oxyethylene)copolymers}$ (where the poly(oxypropylene) content is 3–60 moles and the poly(oxyethylene) content is 5–80 moles), the resulting ethoxylated propoxylated (C12–C15) alcohols having a minimum molecular weight (in amu), 1,500	68551– 13–3
Alkyl (C ₁₂ -C ₂₀) methacrylate-methacrylic acid copolymer, minimum molecular weight (in amu), 11,900	None
1,3 Benzene dicarboxylic acid, 5-sulfo-, 1,3-dimethyl ester, sodium salt, polymer with 1,3-benzene dicarboxylic acid, 1,4-benzene dicarboxylac acid, 11,4-benzene dicarboxylate and 1,2-ethanediol, minimum number average molecular weight (in amu), 2,580	212842– 88–1
3,5-Bis(6-isocyanatohexyl)-2H-1,3,5-oxadiazine- 2,4,6-(3H,5H)-trione, polymer with diethylenetriamine, minimum number average molecular weight (in amu), 1,000,000	87823– 33–4
Butadiene-styrene copolymer	None
1,4-Butanediol-methylenebis(4-phenylisocyanate)- poly(tetramethylene glycol) copolymer, min- imum molecular weight (in amu) 158,000	9018– 04–6
Butene, homopolymer minimum number average molecular weight (in amu), 1,330	9003–29– 6
2-Butenedioic acid (Z)-, polymer with ethenol and ethenyl acetate, sodium salt, minimum number average molecular weight (in amu), 75,000	139871– 83–3
Butyl acrylate-vinyl acetate-acrylic acid copolymer, minimum number average molecular weight (in amu), 18,000	65405– 40–5
$\alpha\textsc{-Butyl-}\omega\textsc{-hydroxypoly(oxypropylene)}$ block polymer with poly(oxyethylene); molecular weight (in amu), 2,400–3,500	None
Castor oil, polyoxyethylated; the poly(oxyethylene) content averages 5–54 moles	None
Chlorinated polyethylene	64754– 90–1
Cross-linked nylon-type polymer formed by the reaction of a mixture of sebacoyl chloride and polymethylene polyphenylisocycanate with a mixture of ethylenediamine and diethylenetriamine	None
Cross-linked polyurea-type encapsulating polymer	None
Dimethylpolysiloxane minimum number average molecular weight (in amu), 6,800	63148– 62–9
Dimethyl silicone polymer with silica, minimum number average molecular weight (in amu), 1,100,000	67762– 90–7
Docosyl methacrylate-acrylic acid copolymer, or docosyl methacrylate-octadecyl methacrylate- acrylic acid copolymer, minimum number aver- age molecular weight (in amu), 3,000	None

3	100.700
Polymer	CAS No.
1,12-Dodecanediol dimethacrylate polymer, minimum molecular weight (in amu), 100,000	None
 2-Ethanediamine, polymer with methyl oxirane and oxirane, minimum number average molec- ular weight (in amu), 1,100 	26316- 40-5
Ethylene glycol dimethyacrylate-lauryl methacry- late copolymer, minimum molecular weight (in amu), 100,000	None
Ethylene glycol dimethacrylate polymer, minimum molecular weight (in amu), 100,000	None
Fumaric acid-isophthalic acid-styrene-ethylene/ propylene glycol copolymer, minimum average molecular weight (in amu), 1 x 10 ¹⁸	None
Hexadecyl acrylate-acrylic acid copolymer, hexadecyl acrylate-butyl acrylate-acrylic acid copolymer, or hexadecyl acrylate-dodecyl acry- late-acrylic acid copolymer, minimum number average molecular weight (in amu), 3,000	None
Hexamethyl disilizane, reaction product with sili- ca, minimum number average molecular weight (in amu), 645,000	68909– 20–6
1,6-Hexanediol dimethyacrylate polymer, min- imum molecular weight (in amu), 100,000	None
α-Hydro-ω-hydroxy-poly(oxyethylene) C8 alkyl ether citrates, poly(oxyethylene) content is 4– 12 moles, minimum number average molecular weight (in amu) 1,300.	330977- 00-9
α-Hydro-ω-hydroxy-poly(oxyethylene) C10–C16- alkyl ether citrates, poly(oxyethylene) content is 4–12 moles, minimum number average molec- ular weight (in amu) 1,100.	330985– 58–5
α-Hydro-ω-hydroxy-poly(oxyethylene) C16–C18- alkyl ether citrates, poly(oxyethylene) content is 4–12 moles, minimum number average molec- ular weight (in amu) 1,300.	330985– 61–0
α-Hydro-ω-hydroxypoly(oxyethylene), minimum molecular weight (in amu), 100,000	None
α-Hydro-ω-hydroxypoly(oxyethylene)poly (oxypropylene) poly(oxyethylene) block copolymer; the minimum poly(oxypropylene) content is 27 moles and the minimum molecular weight (in amu) is 1,900	None
$\begin{array}{ll} \alpha\text{-Hydro-}\omega\text{-hydroxypoly(oxypropylene)}; & \text{minimum} \\ & \text{molecular weight (in amu) 2,000} \end{array}$	None
12-Hydroxystearic acid-polyethylene glycol co- polymer, minimum number average molecular weight (in amu), 3,690	70142– 34–6
Lauryl methacrylate-1,6-hexanediol dimethacrylate copolymer, minimum molecular weight (in amu), 100,000	None
Maleic acid-butadiene copolymer	None
Maleic acid monobutyl ester-vinyl methyl ether copolymer, minimum average molecular weight (in amu), 52,000	25119- 68-0

40 CFR Ch. I (7-1-03 Edition)

§ 180.960

Polymer	CAS No.
Maleic acid monoethyl ester-vinyl methyl ether copolymer, minimum average molecular weight (in amu), 46,000	25087- 06-3
Maleic acid monoisopropyl ester-vinyl methyl ether copolymer, minimum average molecular weight (in amu), 49,000	31307– 95–6
Maleic anhydride-diisobutylene copolymer, so- dium salt, minimum number average molecular weight (in amu) 5,0007–18,000	37199– 81–8
Maleic anhydride-methylstyrene copolymer so- dium salt, minimum number average molecular weight (in amu), 15,000	60092– 15–1
Maleic anhydride-methyl vinyl ether, copolymer, average molecular weight (in amu), 250,000	None
Methacrylic acid-methyl methacrylate-poly- ethylene glycol methyl ether methacrylate co- polymer, minimum number averge molecular weight (in amu), 3,700	100934- 04-1
Methacrylic copolymer, minimum number average molecular weight (in amu), 15,000	63150– 03–8
Methyl methacrylate-methacrylic acid- monomethoxypolyethylene glycol methacrylate copolymer,) minimum number average molec- ular weight (in amu), 2,730	119724– 54–8
Methyl methacrylate-2-sulfoethyl methacrylate- dimethylaminoethylmethacrylate-glycidyl meth- acrylate-styrene-2-ethylhexyl acrylate graft co- polymer, minimum average molecular weight (in amu), 9,600	None
Methyl vinyl ether-maleic acid copolymer), min- imum number average molecular weight (in amu), 75,000	25153– 40–6
Methyl vinyl ether-maleic acid copolymer, calcium sodium salt, minimum number average molecular weight (in amu), 900,000	62386– 95–2
Monophosphate ester of the block copolymer α -hydro- α -hydroxypoly(oxyethylene) poly(oxypropylene) poly(oxypthylene); the poly(oxypropylene) content averages 37–41 moles, average molecular weight (in amu), 8,000	None
α -(ρ -Nonylphenyl- ω -hydroxypoly(oxypropylene) block polymer with poly(oxyethylene); polyoxypropylene content of 10–60 moles; polyoxyethylene content of 10–80 moles; molecular weight (in amu), 1,200–7,100.	None
α-(ρ-Nonylphenyl)poly(oxypropylene) block poly- mer with poly(oxyethylene); poly oxyethylene content 30 to 90 moles; molecular weight (in amu) averages 3,000	None
Octadecanoic acid, 12-hydroxy-, homopolymer, octadecanoate minimum number average molecular weight (in amu), 1,370	58128– 22–6),
α-cis-9-Octadecenyl-ω-hydroxypoly(oxyethylene); the octadecenyl group is derived from oleyl al- cohol and the poly(oxyethylene) content aver- ages 20 moles	None

Polymer	CAS No
Octadecyl acrylate-acrylic acid copolymer, octa- decyl acrylate-dodecyl acrylate-acrylic acid co- polymer, octadecyl methacrylate-butyl acrylate- acrylic acid copolymer, octadecyl methacrylate- hexyl acrylate-acrylic acid copolymer, octadecyl methacrylate-dodecyl acrylate-acrylic acid co- polymer, or octadecyl methacrylate-dodecyl methacrylate-acrylic acid copolymer, minimum number average molecular weight (in amu) 3,000	None
Oleic acid diester of α -hydro- ω -hydroxypoly(oxyethylene); the poly(oxyethylene), average molecular weight (in amu), 2,300	None
Oxirane, methyl-, polymer with oxirane, mono[2- (2-butoxyethoxy) ethyl] ether, minimum number average molecular weight (in amu), 2,500	85637– 75–8
Polyamide polymer derived from sebacic acid, vegetable oil acids with or without dimerization, terephthalic acid and/or ethylenediamine	None
Polyethylene glycol-polyisobutenyl anhydride-tall oil fatty acid copolymer, minimum number aver- age molecular weight (in amu), 2,960	68650- 28-2
Polyethylene, oxidized, minimum number average molecular weight (in amu), 1,200	None
Polymethylene polyphenylisocyanate, polymer with ethylene diamine, diethylene triamine and sebacoyl chloride, cross-linked; minimum number average molecular weight (in amu), 100,000	None
Polyoxyethylated primary amine $(C_{14}$ – $C_{18})$; the fatty amine is derived from an animal source and contains 3% water; the poly(oxyethylene) content averages 20 moles	None
Polyoxyethylated sorbitol fatty acid esters; the polyoxyethylated sorbitol solution containing 15% water is reacted with fatty acids limited to C ₁₂ , C ₁₄ , C ₁₆ , and C ₁₈ , containing minor amounts of associated fatty acids; the poly(oxyethylene) content averages 30 moles.	
Polyoxyethylated sorbitol fatty acid esters; the sorbitol solution containing up to 15% water is reacted with 20–50 moles of ethylene oxide and aliphatic alkanoic and/or alkenoic fatty acids C_8 through C_{22} with minor amounts of associated fatty acids; the resulting polyoxyethylene sorbitol ester having a minimum molecular weight (in amu), 1,300	None
Poly(oxyethylene/oxypropylene) monoalkyl ($C_{\rm c-}$ $C_{\rm 10}$) ether sodium fumarate adduct, minimum number average molecular weight (in amu), 1,900	102900- 02-7
Polyoxymethylene copolymer, minimum number average molecular weight (in amu), 15,000	None
Poly(oxypropylene) block polymer with poly(oxyethylene), molecular weight (in amu), 1,800-16,000	None
Poly(phenylhexylurea), cross-linked, minimum average molecular weight (in amu), 36,000	None

Environmental Protection Agency

Polymer	CAS No.
Polypropylene	9003–07– 0
Polystyrene, minimum number average molecular weight (in amu), 50,000	9003–53– 6
Polytetrafluoroethylene	9002–84– 0
Polyvinyl acetate, copolymer with maleic anhydride, partially hydrolyzed, sodium salt, minimum number average molecular weight (in amu), 53,000	None
Polyvinylpyrrolidone butylated polymer, minimum number average molecular weight (in amu), 9,500	26160– 96–3
Polyvinyl acetate, minimum molecular weight (in amu), 2,000	None
Polyvinyl acetatepolyvinyl alcohol copolymer, minimum number average molecular weight (in amu), 50,000	25213– 24–5
Polyvinyl alcohol	9002–89– 5
Polyvinyl chloride	None
Polyvinyl chloride, minimum number average mo- lecular weight (in amu), 29,000	9002–86–
Poly(vinylpyrrolidone), minimum number average molecular weight (in amu), 4,000	9003–39–
Poly(vinylpyrrolidone-1-eicosene), minimum average molecular weight (in amu), 3,000	28211- 18-9
Poly(vinylpyrrolidone-1-hexadecene), minimum average molecular weight (in amu), 4,700	63231- 81-2
2-Propene-1-sulfonic acid sodium salt, polymer with ethenol and ethenyl acetate, number average molecular weight (in amu) 6,000-12,000	None
2-Propenoic acid, polymer with 2-propenamide, sodium salt, minimum number average molecular weight (in amu), 18,000	25085- 02-3
2-Propenoic acid, sodium salt, polymer with 2-propenamide, minimum number average molecular weight (in amu), 18,000	25987– 30–8
Silane, dichloromethyl- reaction product with silica minimum number average molecular weight (in amu), 3,340,000	68611– 44–9
Sodium polyflavinoidsulfonate, consisting chiefly of the copolymer of catechin and leucocyanidin	None
Stearyl methacrylate-1,6-hexanediol dimethacrylate copolymer, minimum molecular weight (in amu), 100,000	None

Polymer	CAS No.
Styrene, copolymers with acrylic acid and/or methacrylic acid, with none and/or one or more of the following monomers: Acrylamidopropyl methyl sulfonic acid, methallyl sulfonic acid, 3-sulfopropyl acrylate, 3-sulfopropyl methacrylate, hydroxypropyl acrylate, hydroxypropyl acrylate, hydroxypropyl acrylate, and/or hydroxyethyl acrylate; and its sodium, potassium, ammonium, monoethanolamine, and triethanolamine salts; the resulting polymer having a minimum number average molecular weight (in amu), 1,200	None
Styrene, 2-ethylhexyl acrylate, butyl acrylate co- polymer, minimum number average molecular weight (in amu), 4,200	30795– 23–4
Styrene-2-ethylhexyl acrylate-glycidyl methacry- late-2-acrylamido-2-methylpropanesulfonic acid graft copolymer, minimum number average mo- lecular weight (in amu), 12,500	None
Styrene-maleic anhydride copolymer	None
Styrene-maleic anhydride copolymer, ester derivative	None
Tetradecyl acrylate-acrylic acid copolymer, min- imum number average molecular weight (in amu), 3,000	None
Tetraethoxysilane, polymer with hexamethyldisiloxane, minimum number average molecular weight (in amu), 6,500	104133– 09–7
$\begin{array}{lll} \alpha\text{-}[p\text{-}(1,1,3,3\text{-Tetramethylbutyl}) phenyl] \\ & poly(oxypropylene) & block & polymer & with \\ & poly(oxyethylene); & the poly(oxypropylene) & content averages 25 moles, & the poly(oxyethylene) \\ & content & averages & 40 & moles, & the & molecular \\ & weight (in amu) & averages & 3,400 \\ \end{array}$	None
$\alpha\text{-}[2,4,6\text{-}Tris[1\text{-}(phenyl)ethyl]phenyl]-}\omega\text{-}hydroxy poly(oxyethylene) poly(oxypropylene) copolymer, the poly(oxypropylene) content averages 2–8 moles, the poly(oxyethylene) content averages 16–30moles, average molecular weight (in amu), 1,500$	None
Urea-formaldehyde copolymer, minimum average molecular weight (in amu), 30,000	9011–05– 6
Vinyl acetate-allyl acetate-monomethyl maleate copolymer, minimum average molecular weight (in amu), 20,000	None
Vinyl acetate-ethylene copolymer, minimum num-	24937-

Polymer	CAS No.
Vinyl acetate polymer with none and/or one or more of the following monomers: Ethylene, propylene, N-methyl acrylamide, acrylamide, monoethyl maleate, diethyl maleate, monooctyl maleate, dioctyl maleate, maleic anhydride, maleic acid, octyl acrylate, butyl acrylate, ethyl acrylate, methyl acrylate, butyl acrylate, ethyl methacrylate, methyl methacrylate, ethyl methacrylate, methyl methacrylate, methacrylate, and diallyl phthalate; and their corresponding sodium, potassium, ammonium, isopropylamine, triethylamine, monoethanolamine and/or triethanolamine salts; the resulting polymer having a minimum number average molecular weight (in amu), 1,200	None
Vinyl acetate-vinyl alcohol-alkyl lactone copoly- mer, minimum number average molecular weight (in amu), 40,000; minimum viscosity of 18 centipoise	None
Vinyl alcohol-disodium itaconate copolymer, min- imum average molecular weight (in amu), 50,290	None
Vinyl alcohol-vinyl acetate copolymer, benz- aldehyde-o-sodium sulfonate condensate, min- imum number average molecular weight (in amu), 20,000	None
Vinyl alcohol-vinyl acetate-monomethyl maleate, sodium salt-maleic acid, disodium salt-γ-butyro-lactone acetic acid, sodium salt copolymer, minimum number average molecular weight (in amu), 20,000	None
Vinyl chloride-vinyl acetate copolymers	None
Vinyl pyrrolidone-acrylic acid copolymer, minimum number average molecular weight (in amu), 6,000	28062- 44-4
Vinyl pyrrolidone-dimethylaminoethylmethacrylate copolymer, minimum number average molecular weight (in amu), 20,000	30581- 59-0
Vinyl pyrrolidone-styrene copolymer	25086- 29-7

[67 FR 36528, May 24, 2002, as amended at 68 FR 8850, Feb. 26, 2003; 68 FR 10986, Mar. 7, 2003; 68 FR 15967, Apr. 2, 2003; 68 FR 23072, Apr. 30, 2003]

§ 180.1001 Exemptions from the requirement of a tolerance.

- (a) An exemption from a tolerance shall be granted when it appears that the total quantity of the pesticide chemical in or on all raw agricultural commodities for which it is useful under conditions of use currently prevailing or proposed will involve no hazard to the public health.
- (b) When applied to growing crops, in accordance with good agricultural practice, the following pesticide chemicals are exempt from the requirement of a tolerance:
 - (1) [Reserved]
- (2) N-Octylbicyclo(2,2,1)-5-heptene-2,3-dicarboximide.
 - (3) Petroleum oils.
 - (4) Piperonyl butoxide.
 - (5) [Reserved]
 - (6) Pyrethrum and pyrethrins.
 - (7) Rotenone or derris or cube roots.
 - (8) Sabadilla.

These pesticides are not exempted from the requirement of a tolerance when applied to a crop at the time of or after harvest.

(c) Residues of the following materials are exempted from the requirement of a tolerance when used in accordance with good agricultural practice as inert (or occasionally active) in gredients in pesticide formulations applied to growing crops or to raw agricultural commodities after harvest:

Inert ingredients	Limits	Uses
Acetic acid		Catalyst
Acetic anhydride		Solvent, cosolvent
Acetone		Do.
Alkanoic and alkenoic acids, mono- and diesters of α -hydro- ω -hydroxypoly (oxyethylene) with molecular weight (in amu) range of 200 to 6.000.		Emulsifiers
Alkyl (C ₈ -C ₂₄) benzenesulfonic acid and its ammonium, calcium, magnesium, potassium, sodium, and zinc salts.		Surfactants, related adjuvants of surfactants
α -Alkyl (C ₉ -C ₁₈ - ω -hydroxypoly(oxyethylene) with poly(oxyethylene) content of 2-30 moles.		Solvent, cosolvent, surfactant, and related adjuvants of surfactants
α-(p-Alkylphenyl)-ω-hydroxypoly(oxyethylene) produced by the condensation of 1 mole of alkylphenol (alkyl is a mixture of propylene tetramer and pentamer isomers and averages C ₁₋₁) with 6 moles of ethylene oxide.		Surfactants, related adjuvants of surfactants