

MUC Glossary of Terms

This glossary provides definitions, decision criteria, and examples for the land cover types outlined in the Modified UNESCO Classification (MUC) System. The land cover types are organized numerically in the same order as the classes appear in the MUC System Table. Miscellaneous terms used in the glossary are defined in the section following the numbered MUC definitions.

The MUC Glossary of Terms contains four columns of information:

1. **MUC Class** – the number used to classify each land cover type.
2. **MUC Name** – the name used to describe each land cover type.
3. **MUC Level** – the hierarchical level of the MUC System for each MUC Class from 1 (general classes) to 4 (detailed classes).
4. **Definitions** – definitions, decision criteria, and examples used to define each MUC Class.

References

- A land use and land cover classification system for use with remote sensor data. J.R. Anderson, E.E. Hardy, J.T. Roach, and R.E. Witmer. U.S. Geol. Surv. Professional Paper, 1976.
- The Atmosphere: An Introduction to Meteorology*, 6th Ed. Lutgens, Fredrick K. & Tarbuck, Edward J. Englewood Cliffs, NJ: Prentice Hall, Inc., 1995.
- Biology of Plants*, 5th Ed. Raven, Peter H., Evert, Ray F., & Eichhorn, Susan E. New York, NY: Worth Publishers, 1992.
- Classification of wetlands and deepwater habitats of the United States*. L.M. Cowardin, V. Carter, F.C. Golet, and E.T. LaRoe. U.S. Fish and Wildlife Service. FWS/OBS-79/31, 1979.
- International classification and mapping of vegetation*. United Nations Educational, Scientific and Cultural Organization. Switzerland: UNESCO, 1973.
- NOAA Coastal Change Analysis Program (C-CAP): Guidance for Regional Implementation. J.E. Dobson et al. NOAA Technical Report NMFS 123, 1995.

MUC (Modified UNESCO Classification) System Glossary

| MUC Class | MUC Name | MUC Level | Definitions |
|-----------|-----------------------------------|-----------|---|
| 0 | Closed Forest | level 1 | Formed by trees at least 5 meters tall with their crowns (i.e. branches) interlocking. The tree canopy covers at least 40% of the ground. |
| 01 | Mainly Evergreen | level 2 | Within <i>Closed Forest</i> (0). The canopy is never without green foliage. At least 50% of the trees that reach the canopy are evergreen. Individual trees may shed their leaves. |
| 011 | Tropical Wet (Rain) | level 3 | Within <i>Mainly Evergreen Closed Forest</i> (01). Often called a tropical rain forest. Consisting mainly of broad-leaved evergreen trees, neither cold nor drought resistant. Truly evergreen, i.e. the forest canopy remains green all year though a few individual trees may be leafless for a few weeks. Leaves of many species have “drip tips.” |
| 0111 | Lowland | level 4 | Within <i>Tropical Wet Mainly Evergreen Closed Forest</i> (011). Consists usually of numerous species of fast growing trees, many exceeding 50 meters tall, generally with smooth, often thin bark, some with buttresses. Emergent trees or at least a very uneven canopy often present. Undergrowth is sparse, composed mainly of tree seedlings. Palms and other tuft trees usually are rare. Crustose lichens and green algae are present, and climbing vines are usually only abundant in extremely humid regions (e.g., Sumatra, Atrato Valley, Columbia). |
| 0112 | Submontane | level 4 | Within <i>Tropical Wet Mainly Evergreen Closed Forest</i> (011). Emergent trees are largely absent and the canopy is relatively even. Forbs are common in the undergrowth. Vascular epiphytes and vines are abundant. E.g., Atlantic slopes of Costa Rica. |
| 0113 | Montane | level 4 | Within <i>Tropical Wet Mainly Evergreen Closed Forest</i> (011). Trees are less than 50 meters tall, have crowns that extend relatively far down the stem, and often have rough bark. Undergrowth abundant, often with ferns, herbs, mosses, and small palms. E.g., Sierra de Talamanca, Costa Rica. |
| 0114 | Subalpine | level 4 | Within <i>Tropical Wet Mainly Evergreen Closed Forest</i> (011). Occurs at elevations above montane forests, with characteristic vegetation, which is dependent on latitude. |
| 0115 | Cloud | level 4 | Within <i>Tropical Wet Mainly Evergreen Closed Forest</i> (011). Trees are gnarled, have rough bark and are rarely greater than 20 meters tall. Tree crowns, branches, and trunks are burdened with epiphytes, mainly chamaephytic bryophytes. Also, the ground is covered with hygromorphic chamaephytes such as <i>Selaginella</i> and ferns. E.g., Blue Mountains, Jamaica. |
| 012 | Tropical and Subtropical Seasonal | level 3 | Within <i>Mainly Evergreen Closed Forest</i> (01). Consisting mainly of broad-leaved evergreen trees. Foliage reduction during the dry season is noticeable, often as partial shedding. Transitional between Tropical Wet Forest and Tropical and Subtropical Semi-deciduous. |
| 0121 | Lowland | level 4 | Within <i>Tropical and Subtropical Evergreen Seasonal Closed Forest</i> (012). Consists of fast growing trees, many exceeding 50 meters tall and usually forming an uneven canopy. Undergrowth is sparse, lichen and green algae are present, and climbing vines are absent. |
| 0122 | Submontane | level 4 | Within <i>Tropical and Subtropical Evergreen Seasonal Closed Forest</i> (012). Trees form an even canopy. Forbs are common in the undergrowth. Vascular epiphytes and vines are abundant. |
| 0123 | Montane | level 4 | Within <i>Tropical and Subtropical Evergreen Seasonal Closed Forest</i> (012). Trees are less than 50 meters tall, have crowns that extend relatively far down the stem and have rough bark. There are no tree ferns; instead, evergreen shrubs are most common. |

| | | | |
|------|---|---------|--|
| 0124 | Subalpine | level 4 | Within <i>Tropical and Subtropical Evergreen Seasonal Closed Forest</i> (012). This forest resembles the Winter-rain Evergreen Broad-leaved Sclerophyllous dry forest and usually occurs above the cloud forest. Trees are mostly evergreen sclerophyllous trees, smaller than 20 meters with little or no undergrowth, few climbing vines, and few epiphytes, except lichens. |
| 013 | Tropical and Subtropical Semi-Deciduous | level 3 | Within Mainly <i>Evergreen Closed Forest</i> (01). Most of the upper canopy trees are drought-deciduous; many of the understory trees and shrubs are evergreen and more or less sclerophyllous. However, evergreen and deciduous woody plants and shrubs are not always separated by layers; they may occur mixed within the same layer, or shrubs may be primarily deciduous and trees evergreen. Nearly all trees have bud protection and leaves without “drip tips.” Trees have rough bark, except some bottle trees, which may be present. |
| 0131 | Lowland | level 4 | Within Mainly <i>Evergreen Tropical and Subtropical Semi-deciduous Closed Forest</i> (013). The taller trees may be bottle trees (e.g., <i>Catiba</i>). There are practically no epiphytes present. The undergrowth is composed of shrubs and seedlings. Succulents such as thin-stemmed caespitose cacti may be present. Vines occur occasionally. A sparse layer of herbaceous vegetation may also be present. |
| 0133 | Montane and Cloud | level 4 | Within Mainly <i>Evergreen Tropical and Subtropical Semi-deciduous Closed Forest</i> (013). This forest is similar to a Semideciduous Lowland Forest, however, the canopy is lower and covered with xerophytic epiphytes such as <i>Tillandsia usneoides</i> . |
| 014 | Subtropical Wet | level 3 | Within Mainly <i>Evergreen Closed Forest</i> (01). Present only locally and in small fragmentary stands, because the subtropical climate typically has a dry season. It usually grades into Tropical Wet Forest (e.g., Queensland, Australia and Taiwan). Some shrubs may grow in the understorey. Seasonal temperature change occurs between summer and winter. There is a more pronounced temperature difference between summer and winter than the (Tropical Wet) Montane Forest (0113). |
| 0141 | Lowland | level 4 | Within Mainly <i>Evergreen Subtropical Wet Closed Forest</i> (014). Consists usually of numerous species of fast growing trees, many exceeding 50 meters tall, generally with smooth, often thin bark, some with buttresses. Emergent trees or at least a very uneven canopy often present. Undergrowth is sparse, composed mainly of tree seedlings. Palms and other tuft trees usually are rare. Crustose lichens and green algae are present, and climbing vines are usually only abundant in extremely humid regions. |
| 0142 | Submontane | level 4 | Within Mainly <i>Evergreen Subtropical Wet Closed Forest</i> (014). Emergent trees are largely absent and the canopy is relatively even. Forbs are common in the undergrowth. Vascular epiphytes and vines are abundant. |
| 0143 | Montane | level 4 | Within Mainly <i>Evergreen Subtropical Wet Closed Forest</i> (014). Trees are less than 50 meters tall, have crowns that extend relatively far down the stem, and often have rough bark. Undergrowth abundant, often with ferns, herbs, mosses, and small palms. |
| 0144 | Subalpine | level 4 | Within Mainly <i>Evergreen Subtropical Wet Closed Forest</i> (014). Occurs at elevations above montane forests, with characteristic vegetation, which is dependent on latitude. |
| 0145 | Cloud | level 4 | Within Mainly <i>Evergreen Subtropical Wet Closed Forest</i> (014). Trees are gnarled, have rough bark and are rarely greater than 20 meters tall. Tree crowns, branches, and trunks are burdened with epiphytes, mainly chamaephytic bryophytes. Also, the ground is covered with hygromorphic chamaephytes (e.g., <i>Selaginella</i> and herbaceous ferns). |
| 015 | Temperate or Subpolar Wet | level 3 | Within Mainly <i>Evergreen Closed Forest</i> (01). Occurs only in the extremely oceanic, nearly frost-free climates of the southern hemisphere, mainly in Chile. Consisting mostly of truly evergreen hemisclerophyllous trees and shrubs. Rich in epiphytic mosses, liverworts, and lichens that grow on trees, and in ground-rooted herbaceous ferns. |
| 0151 | Temperate | level 4 | Within Mainly <i>Evergreen Temperate or Subpolar Wet Closed Forest</i> (015). Trees are generally greater than 10 meters tall. Vascular epiphytes and vines may be present. |

| | | | |
|------|---|---------|--|
| 0152 | Subpolar | level 4 | Within <i>Mainly Evergreen Temperate or Subpolar Wet Closed Forest</i> (015). Trees are generally less than 10 meters tall and often have reduced leaf size. There are few vascular epiphytes present. E.g., beech forests of New Zealand. |
| 016 | Temperate with Broad-Leaved Deciduous | level 3 | Within <i>Mainly Evergreen Closed Forest</i> (01). Requires adequate summer rainfall. This is a mixed evergreen-deciduous class. The dominant trees are mainly hemisclerophyllous evergreen trees (more than 50% of the canopy) and shrubs, and the subdominant trees are deciduous broad-leaved trees and shrubs (more than 25% of the canopy). Rich in perennial herbaceous plants. Very few or no vascular epiphytes and vines. |
| 0161 | Lowland | level 4 | Within <i>Temperate Deciduous Broad-Leaved Mainly Evergreen Closed Forest</i> (016). Consists usually of numerous species of fast growing trees, many exceeding 50 meters tall, generally with smooth, often thin bark, some with buttresses. Emergent trees or at least a very uneven canopy often present. Undergrowth is sparse, composed mainly of tree seedlings. Palms and other tuft trees usually are rare. Crustose lichens and green algae are present, and climbing vines are usually only abundant in extremely humid regions. |
| 0162 | Submontane | level 4 | Within <i>Temperate Deciduous Broad-Leaved Mainly Evergreen Closed Forest</i> (016). Emergent trees are largely absent and the canopy is relatively even. Forbs are common in the undergrowth. Vascular epiphytes and vines are abundant. |
| 0163 | Montane | level 4 | Within <i>Temperate Deciduous Broad-Leaved Mainly Evergreen Closed Forest</i> (016). Trees are less than 50 meters tall, have crowns that extend relatively far down the stem, and often have rough bark. Undergrowth abundant, often with ferns, herbs, mosses, and small palms. |
| 0164 | Subalpine | level 4 | Within <i>Temperate Deciduous Broad-Leaved Mainly Evergreen Closed Forest</i> (016). Occurs at elevations above montane forests, with characteristic vegetation, which is dependent on latitude. |
| 017 | Winter-Rain Broad-Leaved Sclerophyllous | level 3 | Within <i>Mainly Evergreen Closed Forest</i> (01). Often understood as Mediterranean, but present also in southwestern Australia, Chile, and other locations. The climate has a pronounced summer drought. Consisting mainly of sclerophyllous evergreen trees and shrubs, most of which have rough bark. There is very little herbaceous undergrowth. No vascular and few non-flowering epiphytes and lichens, but evergreen woody vines are present. |
| 0171 | Lowland and Submontane >50m | level 4 | Within <i>Winter-Rain Evergreen Broad-Leaved Sclerophyllous Closed Forest</i> (017). Dominated by trees over 50 meters tall (at least 50% of the canopy) such as giant eucalyptus (e.g., <i>Eucalyptus regnans</i>) in Victoria, Australia and <i>E. diversicolor</i> in Western Australia). |
| 0172 | Lowland and Submontane <50m | level 4 | Within <i>Winter-Rain Evergreen Broad-Leaved Sclerophyllous Closed Forest</i> (017). Dominated by trees less than 50 meters tall (more than 50% of the canopy). E.g., Californian live-oak forests. |
| 018 | Tropical and Subtropical Needle-Leaved | level 3 | Within <i>Mainly Evergreen Closed Forest</i> (01). Consisting mainly of needle-leaved or scale-leaved evergreen trees (more than 50% of the canopy). Broad-leaved trees may be present. Vascular epiphytes and vines rarely present. Species typical of the tropical/subtropical zone. |
| 0181 | Lowland and Submontane | level 4 | Within <i>Tropical and Subtropical Needle-Leaved Mainly Evergreen Closed Forest</i> (018). E.g., the pine forests of Honduras and Nicaragua. |
| 0182 | Montane and Subalpine | level 4 | Within <i>Tropical and Subtropical Needle-Leaved Mainly Evergreen Closed Forest</i> (018). E.g., the pine forests of the Philippines and southern Mexico. |
| 019 | Temperate and Subpolar Needle-Leaved | level 3 | Within <i>Mainly Evergreen Closed Forest</i> (01). Consisting mainly of needle-leaved or scale-leaved evergreen trees (more than 50% of the canopy), but broad-leaved trees may be present. Vascular epiphytes and vines are rarely present. Species typical of the temperate/subpolar zone. |
| 0191 | Giant (> 50m) | level 4 | Within <i>Temperate and Subpolar Needle-Leaved Mainly Evergreen Closed Forest</i> (019). Dominated by trees (at least 50% of the canopy) greater than 50 meters tall (e.g., <i>Sequoia</i> and <i>Pseudo-tsuga</i>) forest in the Pacific West of North America). |

| | | | |
|------|--|---------|--|
| 0192 | Irregularly Rounded Crowns | level 4 | Within <i>Temperate and Subpolar Needle-Leaved Mainly Evergreen Closed Forest</i> (019). Dominated by trees 45-50 meters tall (more than 50% of the canopy), with broad, irregularly rounded crowns (e.g., <i>Pinus</i> spp.). |
| 0193 | Conical Crowns | level 4 | Within <i>Temperate and Subpolar Needle-Leaved Mainly Evergreen Closed Forest</i> (019). Dominated by trees 45-50 meters tall (more than 50% of the canopy), with conical crowns (like most <i>Picea</i> and <i>Abies</i>). E.g., California red fir forests. |
| 0194 | Cylindrical Crowns | level 4 | Within <i>Temperate and Subpolar Needle-Leaved Mainly Evergreen Closed Forest</i> (019). Dominated by trees 45-50 meters tall (more than 50% of the canopy), with crowns with very short branches and therefore a narrow cylindrical shape. |
| 02 | Mainly Deciduous | level 2 | Within <i>Closed Forest</i> (0). The majority of trees (more than 50% of the canopy) shed their foliage simultaneously in connection with the unfavorable season (drought or cold). |
| 021 | Tropical and Subtropical Drought-Deciduous | level 3 | Within <i>Mainly Deciduous Closed Forest</i> (02). The unfavorable season is mainly characterized by drought, in most cases by winter-drought. Foliage is shed regularly every year. Most trees have relatively thick, fissured bark. |
| 0211 | Broad-Leaved Lowland and Submontane | level 4 | Within <i>Tropical and Subtropical Drought-Deciduous Closed Forest</i> (021). Practically no evergreen plants in stratum except some succulents. Woody and herbaceous vines and deciduous bottle-trees are present occasionally. Sparse herbaceous vegetation present in the undergrowth. E.g., the broad-leaved deciduous forests of northwestern Costa Rica. |
| 0212 | Montane and Cloud | level 4 | Within <i>Tropical and Subtropical Drought-Deciduous Closed Forest</i> (021). Some evergreen species are present in the understory. Drought resistant epiphytes are present or abundant, often of the bearded form (e.g., <i>Usnea</i> or <i>Tillandsia usneoides</i>). This formation is not frequent, but well developed. E.g., in northern Peru. |
| 022 | Cold-Deciduous with Evergreens | level 3 | Within <i>Mainly Deciduous Closed Forest</i> (02). The unfavorable season is mainly characterized by winter frost. Deciduous broad-leaved trees are dominant (more than 50% of the canopy), but evergreen species are present (more than 25% of the canopy) as part of the main canopy or the understory. Climbers and vascular epiphytes are scarce or absent. |
| 0221 | With Evergreen Broad-Leaved Trees and Climbers | level 4 | Within <i>Cold-Deciduous with Evergreens Closed Forest</i> (022). Rich in epiphytes, including mosses. Vascular epiphytes may be present at the base of tree stems. Climbing vines may be common on flood plains. Ex. <i>Ilex aquifolium</i> and <i>Hedera helix</i> in western Europe and <i>Magnolia</i> spp. in North America. |
| 0222 | With Evergreen Needle-Leaved Trees | level 4 | Within <i>Cold-Deciduous with Evergreens Closed Forest</i> (022). With evergreen needle-leaved trees such as hemlock (<i>Tsuga</i>) and pine (<i>Pinus</i>). E.g., the maple-hemlock or oak-pine forests of Northeastern, U.S.A. |
| 023 | Cold-Deciduous without Evergreen Trees | level 3 | Within <i>Mainly Deciduous Closed Forest</i> (02). Deciduous trees are absolutely dominant (more than 75% of the canopy). Evergreen herbs and some evergreen shrubs (less than 2 meters tall) may be present. Climbers insignificant but may be common on flood plains. Vascular epiphytes are absent (except occasionally at the lower base of the tree). Mosses, liverworts and particularly lichens are always present. |
| 0231 | Temperate Lowland and Submontane Broad-Leaved | level 4 | Within <i>Cold-Deciduous without Evergreen Trees Closed Forest</i> (023). Trees are up to 50 meters tall. Epiphytes are primarily algae and crustose lichens. E.g., the Mixed Mesophytic Forest of U.S.A. |
| 0232 | Montane and Boreal | level 4 | Within <i>Cold-Deciduous without Evergreen Trees Closed Forest</i> (023). Trees may be up to 50 meters tall, but in montane or boreal forest normally not taller than 30 meters. Epiphytes are primarily lichens and bryophytes. This class includes lowland or submontane in topographic positions with high atmospheric humidity. |
| 0233 | Subalpine and Subpolar | level 4 | Within <i>Cold-Deciduous without Evergreen Trees Closed Forest</i> (023). Trees are not taller than 20 meters and tree trunks are frequently gnarled. Epiphytes are lichens and bryophytes, and are more abundant than in the Montane or Boreal class (0232). This class often grades into woodland. |

| | | | |
|----------|-------------------------------------|----------------|---|
| 03 | Extremely Xeromorphic (Dry) | level 2 | Within <i>Closed Forest</i> (0). Dense stands of trees adapted to dry conditions, such as bottle trees, tuft trees with succulent leaves and stem succulents. Undergrowth has shrubs adapted to dry conditions, succulent perennial herbs and annual and perennial herbaceous plants. Often grades into woodland. |
| 031 | Sclerophyllous-Dominated | level 3 | Within <i>Extremely Xeromorphic Closed Forest</i> (03). There is a predominance of sclerophyllous trees, many of which have bulbous stem bases largely embedded in the soil. |
| 032 | Thorn-Dominated | level 3 | Within <i>Extremely Xeromorphic Closed Forest</i> (03). Species with thorns are dominant (more than 50% of the canopy). |
| 0321 | Mixed Deciduous-Evergreen | level 4 | Within <i>Extremely Xeromorphic Thorn-Dominated Closed Forest</i> (032). Both deciduous and evergreen thorn species are more than 25% of the tree canopy. |
| 0322 | Purely Deciduous | level 4 | Within <i>Extremely Xeromorphic Thorn-Dominated Closed Forest</i> (032). Deciduous thorn species are absolutely dominant (more than 75% of the canopy). |
| 033 | Mainly Succulent | level 3 | Within <i>Extremely Xeromorphic Closed Forest</i> (03). Tree-formed (scapose) and shrub-formed (caespitose) succulents are very frequent (more than 50% of the canopy), but other trees and shrubs adapted to dry conditions are usually present as well. |
| 1 | Woodland | level 1 | Comprised of open stands of trees at least 5 meters tall with crowns not interlocking. The tree canopy covers at least 40% of the ground. Definitions for Mainly Evergreen Woodland, Mainly Deciduous Woodland, and Extremely Xeromorphic Woodland are similar to forest definitions, with sparser stocking of individual trees. |
| 11 | Mainly Evergreen | level 2 | Within <i>Woodland</i> (1). The canopy is never without green foliage. At least 50% of the trees that reach the canopy are evergreen. Individual trees may shed their leaves. |
| 111 | Broad-Leaved | level 3 | Within <i>Mainly Evergreen Woodland</i> (11). Mainly sclerophyllous broad-leaved trees and shrubs, with no epiphytes. |
| 112 | Needle-Leaved | level 3 | Within <i>Mainly Evergreen Woodland</i> (11). Trees are mainly needle- or scale-leaved (more than 50% of the canopy). Crowns of many trees extend to the base of the stem or are very branched. |
| 1121 | Irregularly Rounded Crowns | level 4 | Within <i>Mainly Evergreen Needle-Leaved Woodland</i> (112). Dominated by trees (more than 50% of the canopy) with broad, irregularly rounded crowns (e.g., <i>Pinus</i>). |
| 1122 | Conical Crowns | level 4 | Within <i>Mainly Evergreen Needle-Leaved Woodland</i> (112). Dominated by trees (more than 50% of the canopy) with conical crowns. Mostly in subalpine areas. |
| 1123 | Cylindrical Crowns | level 4 | Within <i>Mainly Evergreen Needle-Leaved Woodland</i> (112). Dominated by trees (more than 50% of the canopy) with crowns with very short branches and therefore a narrow cylindrical shape (e.g., <i>Picea</i> in the boreal regions). |
| 12 | Mainly Deciduous | level 2 | Within <i>Woodland</i> (1). The majority of trees (more than 50% of the canopy) shed their foliage simultaneously in connection with the unfavorable season (drought or cold). |
| 121 | Drought-Deciduous | level 3 | Within <i>Mainly Deciduous Woodland</i> (12). The unfavorable season is mainly characterized by drought, in most cases by winter-drought. Foliage is shed regularly every year. Most trees have relatively thick, fissured bark. |
| 1211 | Broad-Leaved Lowland and Submontane | level 4 | Within <i>Drought-Deciduous Woodland</i> (121). Practically no evergreen plants in any stratum except some succulents. Woody and herbaceous vines and deciduous bottle-trees are present. Sparse herbaceous vegetation present in the undergrowth. |
| 1212 | Montane and Cloud | level 4 | Within <i>Drought-Deciduous Woodland</i> (121). Some evergreen species are present in the understory. Drought resistant epiphytes are present or abundant, often of the bearded form (e.g., <i>Usnea</i> or <i>Tillandsia usneoides</i>). This formation is not frequent, but well developed. E.g., in northern Peru. |

| | | | |
|------|--|---------|---|
| 122 | Cold-Deciduous with Evergreens | level 3 | Within Mainly Deciduous Woodland (12). The unfavorable season is mainly characterized by winter frost. Deciduous broad-leaved trees are dominant (more than 50% of the canopy), but evergreen species are present (more than 25% of the canopy) as part of the main canopy or the understory. Climbers and vascular epiphytes are scarce or absent. |
| 1221 | With Evergreen Broad-Leaved Trees and Climbers | level 4 | Within Cold-Deciduous with Evergreens Woodland (122). Rich in epiphytes, including mosses. Vascular epiphytes may be present at the base of tree stems. Climbing vines may be common on flood plains. <i>Ilex aquifolium</i> and <i>Hedera helix</i> in western Europe and <i>Magnolia</i> spp. in North America are examples of this class type. |
| 1222 | With Evergreen Needle-Leaved Trees | level 4 | Within Cold-Deciduous with Evergreens Woodland (122). With evergreen needle-leaved trees such as hemlock (<i>Tsuga</i>) and pine (<i>Pinus</i>). E.g., the maple-hemlock or oak-pine woodlands of Northeastern, U.S.A. |
| 123 | Cold-Deciduous without Evergreen Trees | level 3 | Within Mainly Deciduous Woodland (12). Cold-deciduous tree species are absolutely dominant (more than 75% of the canopy). Evergreen herbs and some evergreen shrubs (less than 2 meters tall) may be present. Climbers insignificant but may be common on flood plains. Vascular epiphytes are absent (except occasionally at the lower base of the tree). Mosses, liverworts and particularly lichens are always present. Most frequent in the subarctic region, elsewhere only in swamps or bogs. |
| 1231 | Broad-Leaved | level 4 | Within Cold-Deciduous without Evergreen Trees Woodland (123). Broad-leaved deciduous species are absolutely dominant (more than 75% of the canopy). |
| 1232 | Needle-leaved | level 4 | Within Cold-Deciduous without Evergreen Trees Woodland (123). Needle-leaved deciduous species are absolutely dominant (more than 75% of the canopy). |
| 1233 | Mixed | level 4 | Within Cold-Deciduous without Evergreen Trees Woodland (123). Both broad-leaved and needle leaved deciduous species provide more than 25% of the canopy. |
| 13 | Extremely Xeromorphic (Dry) | level 2 | Within Woodland (1). Stands of trees and shrubs adapted to dry conditions, such as bottle trees, tuft trees with succulent leaves and stem succulents. Undergrowth has shrubs adapted to dry conditions, succulent perennial herbs and annual and perennial herbaceous plants. Woodlands may grade into forests. |
| 131 | Sclerophyllous-Dominated | level 3 | Within Extremely Xeromorphic Woodland (13). There is a predominance of sclerophyllous trees, many of which have bulbous stem bases largely embedded in the soil. |
| 132 | Thorn-Dominated | level 3 | Within Extremely Xeromorphic Woodland (13). Species with thorns are dominant (more than 50% of the canopy). |
| 1321 | Mixed Deciduous-Evergreen | level 4 | Within Extremely Xeromorphic Thorn-Dominated Woodland (132). Both deciduous species and evergreen species are more than 25% of the tree canopy. See definitions of Mainly Evergreen Woodland, class 11 and Mainly Deciduous Woodland (MUC Class 12). |
| 1322 | Purely Deciduous | level 4 | Within Extremely Xeromorphic Thorn-Dominated Woodland (132). Deciduous thorn species are absolutely dominant (more than 75% of the canopy). See definition of Mainly Deciduous Woodland (MUC Class 12). |
| 133 | Mainly Succulent | level 3 | Within Extremely Xeromorphic Woodland (13). Tree-formed (scapose) and shrub-formed (caespitose) succulents are very frequent (more than 50% of the tree canopy), but other trees and shrubs adapted to dry conditions are usually present as well. |
| 2 | Shrubland or Thicket | level 1 | The shrub canopy covers at least 40% of the ground and is composed of matted, clumped or clustered woody plants 0.5 to 5 meters tall. Shrublands of the individual shrubs are not touching each other; often with grass growing between shrubs. Shrublands are also further defined (like Forests and Woodlands) as Evergreen Broad-leaved, Evergreen Needle-leaved, Mainly Deciduous, etc. Thickets individual shrub branches are interlocked. |

| | | | |
|------|--|---------|---|
| 21 | Mainly Evergreen | level 2 | Within Shrubland or Thicket (2). The canopy is never without green foliage. At least 50% of the shrubs that reach the canopy are evergreen. Individual shrubs may shed their leaves. |
| 211 | Broad-Leaved | level 3 | Within Mainly Evergreen Shrubland or Thicket (21). Evergreen broad-leaved species are dominant (more than 50% of the canopy). |
| 2111 | Low Bamboo | level 4 | Within Mainly Evergreen Broad-Leaved Shrubland or Thicket (211). Bamboo species are dominant. (Lignified creeping graminoid nano- or microphanerophytes). |
| 2112 | Tuft-Tree | level 4 | Within Mainly Evergreen Broad-Leaved Shrubland or Thicket (211). Composed of small trees and woody shrubs. E.g., Mediterranean dwarf palms shrubland or Hawaiian tree fern thicket or shrubland. |
| 2113 | Broad-Leaved Hemi-Sclerophyllous | level 4 | Within Mainly Evergreen Broad-Leaved Shrubland or Thicket (211). Matted or clumped shrubs and plants with large soft leaves (caespitose, creeping or lodged nano- or microphanerophytes). E.g., subalpine <i>Rhododendron</i> thickets, or <i>Hibiscus tiliaceus</i> matted thickets of Hawaii. |
| 2114 | Broad-Leaved Sclerophyllous | level 4 | Within Mainly Evergreen Broad-Leaved Shrubland or Thicket (211). Dominated by broad-leaved sclerophyllous shrubs and immature trees (e.g., chappatal or macchia). May often merge with parkland, grassland or heath. |
| 2115 | Suffruticose | level 4 | Within Mainly Evergreen Broad-Leaved Shrubland or Thicket (211). Stand of semi-lignified nanophanerophytes that in dry years may shed part of their shoot systems (e.g., <i>Cistus</i> heath). |
| 212 | Needle-Leaved or Microphyllous | level 3 | Within Mainly Evergreen Shrubland or Thicket (21). Dominant species (more than 50% of the canopy) have either needle leaves or small leaves. |
| 2121 | Needle-Leaved | level 4 | Within Mainly Evergreen Needle-Leaved or Microphyllous Shrubland or Thicket (212). Composed of creeping or lodged needle-leaved shrubs (e.g., <i>Pinus</i> mughus, "Krummhölz"). |
| 2122 | Microphyllous | level 4 | Within Mainly Evergreen Needle-Leaved or Microphyllous Shrubland or Thicket (212). Evergreen species have small leaves, (e.g., desert plants) or leaves with a single unbranched vein. Mostly in tropical subalpine belts. |
| 22 | Mainly Deciduous | level 2 | Within Shrubland or Thicket (2). The majority of shrubs (more than 50% of the canopy) shed their foliage simultaneously in connection with the unfavorable season (cold or drought). |
| 221 | Drought-Deciduous with Evergreen Woody Plants | level 3 | Within Mainly Deciduous Shrubland or Thicket (22). Drought-deciduous shrubs are dominant (greater than 50% of the canopy) and are mixed with at least 25% evergreen woody plants. The unfavorable season is mainly characterized by drought. |
| 222 | Drought-Deciduous without Evergreen Woody Plants | level 3 | Within Mainly Deciduous Shrubland or Thicket (22). Drought-deciduous shrubs are absolutely dominant (more than 75% of the canopy). The unfavorable season is mainly characterized by drought. |
| 223 | Cold-Deciduous | level 3 | Within Mainly Deciduous Shrubland or Thicket (22). The unfavorable season is mainly characterized by winter frost. Deciduous shrubs are dominant (more than 50% of the canopy). |
| 2231 | Temperate | level 4 | Within Cold-Deciduous Shrubland or Thicket (223). Composed of dense scrub without, or with very little herbaceous undergrowth. Very few to no cryptogams. |
| 2232 | Subalpine and Subpolar | level 4 | Within Cold-Deciduous Shrubland or Thicket (223). Composed of upright or lodged matted shrubs with great vegetative regeneration capacity and usually covered by snow for at least half a year. |
| 23 | Extremely Xeromorphic (Subdesert) Shrubland | level 2 | Within Shrubland or Thicket (2). Very open stands of shrubs with various adaptations to dry conditions, such as: extremely thickened, hardened foliage; very reduced leaves; green branches without leaves; or succulent stems, some of them with thorns. |

| | | | |
|------|--|---------|--|
| 231 | Mainly Evergreen | level 3 | Within <i>Extremely Xeromorphic Shrubland</i> (23). The canopy is never without green foliage. At least 50% of the shrubs that reach the canopy are evergreen. In extremely dry years some leaves and shoot portions may be shed. |
| 2311 | Purely Evergreen | level 4 | Within <i>Mainly Evergreen Extremely Xeromorphic Shrubland</i> (231). Composed of broad-leaved mostly sclerophyllous shrubs (e.g., mulga scrub in Australia) leafless green-stemmed plants (e.g. <i>Retama retam</i>) or succulents dominated by variously branched stem and leaf succulents. |
| 2312 | Semi-Deciduous | level 4 | Within <i>Mainly Evergreen Extremely Xeromorphic Shrubland</i> (231). May consist of either facultatively deciduous shrubs (e.g., <i>Atriplex-Kochia</i> saltbush in Australia and North America) or a combination of evergreen and deciduous shrubs (i.e. evergreen shrubs are dominant, deciduous shrubs cover more than 25%). |
| 232 | Mainly Deciduous | level 3 | Within <i>Extremely Xeromorphic Shrubland</i> (23). The majority of shrubs (more than 50% of the canopy) shed their foliage simultaneously in connection with the unfavorable season (cold or drought). |
| 2321 | Without Succulents | level 4 | Within <i>Mainly Deciduous Extremely Xeromorphic Shrubland</i> (232). Succulents cover less than 25% of the ground. |
| 2322 | With Succulents | level 4 | Within <i>Mainly Deciduous Extremely Xeromorphic Shrubland</i> (232). Succulents cover at least 25% of the ground. |
| 3 | Dwarf-Shrubland or Dwarf-Thicket | level 1 | Shrubs rarely exceed 50 cm in height (sometimes called heaths or heathlike formations). The shrub canopy covers at least 40% of the ground. The shrub cover density distinguishes between Dwarf-Shrubland and Dwarf-Thicket classes. |
| | | | Dwarf-Shrubland: individual dwarf-shrubs are isolated or in clumps. Dwarf-Thicket: individual shrub branches are interlocked. |
| 31 | Mainly Evergreen | level 2 | Within <i>Dwarf-Shrubland or Dwarf-Thicket</i> (3). The canopy is never without green foliage. At least 50% of the shrubs that reach the canopy are evergreen. Individual shrubs may shed their leaves. |
| 311 | Dwarf-Thicket | level 3 | Within <i>Mainly Evergreen Dwarf-Shrubland or Dwarf-Thicket</i> (31). Composed of densely closed dwarf-shrub cover, which dominates the landscape. |
| 3111 | Caespitose | level 4 | Within <i>Mainly Evergreen Dwarf-Thicket</i> (311). Shrub branches stand upright and are often occupied by lichens (foliose). Cushion-shaped mosses, lichens and other herbaceous plants are often found on the ground (e.g., <i>Calluna</i> heath). |
| 3112 | Creeping | level 4 | Within <i>Mainly Evergreen Dwarf-Thicket</i> (311). Shrub branches creep along the ground. Variously combined with shrubs (e.g., thallochamaephytes) with branches that may be embedded (e.g., <i>Loiseleuria</i> heath). |
| 312 | Dwarf-Shrubland | level 3 | Within <i>Mainly Evergreen Dwarf-Shrubland or Dwarf-Thicket</i> (31). Open or less dense cover of dwarf-shrubs. Shrub canopies are not interlocked. Herbaceous vegetation (i.e. grasses and forbs) covers less than 25% of the ground. |
| 3121 | Cushion | level 4 | Within <i>Mainly Evergreen Dwarf-Shrubland</i> (312). Shrubs are isolated in clumps forming dense cushions and are often thorny (e.g., <i>Astragalus</i> - and <i>Acantholimon</i> "porcupine" heath of the East Mediterranean mountains). |
| 313 | Mixed Evergreen and Herbaceous Dwarf-Shrubland | level 3 | Within <i>Mainly Evergreen Dwarf-Shrubland or Dwarf-Thicket</i> (31). Shrub canopies are not interlocked. Evergreen shrubs are mixed with herbaceous vegetation (at least 25% of the ground). |
| 3131 | True Evergreen and Herbaceous Mixed | level 4 | Within <i>Mixed Evergreen and Herbaceous Dwarf-Shrubland</i> (313). True Evergreen individuals do not seasonally shed parts of their shoot systems. E.g., <i>Nardus</i> <i>Calluna</i> -heath. |
| 3132 | Partial Evergreen and Herbaceous Mixed | level 4 | Within <i>Mixed Evergreen and Herbaceous Dwarf-Shrubland</i> (313). Many individuals shed parts of their shoot systems during the dry season (e.g., <i>Phrygana</i> in Greece). |
| 32 | Mainly Deciduous | level 2 | Within <i>Dwarf-Shrubland or Dwarf-Thicket</i> (3). The majority of shrubs (more than 50% of the canopy) shed their foliage simultaneously in connection with the unfavorable season (cold or drought). |

| | | | |
|------|---|---------|--|
| 321 | Facultative Drought-Deciduous | level 3 | Within Mainly Deciduous Dwarf-Shrubland or Dwarf-Thicket (32). Dwarf-shrubs shed their foliage only in extremely dry years. |
| 322 | Obligate Drought-Deciduous | level 3 | Within Mainly Deciduous Dwarf-Shrubland or Dwarf-Thicket (32). Densely closed dwarf-shrubs lose all or at least part of their leaves in the dry season. |
| 3221 | Caespitose Dwarf-Thicket | level 4 | Within Obligate Drought-Deciduous Dwarf-Shrubland or Dwarf-Thicket (322). Shrub branches stand upright and are often occupied by lichens (foliose). Cushion-shaped mosses, lichens and other herbaceous plants are often found on the ground (e.g., <i>Calluna</i> heath). |
| 3222 | Creeping Dwarf-Thicket. | level 4 | Within Obligate Drought-Deciduous Dwarf-Shrubland or Dwarf-Thicket (322). Shrub branches creep along the ground. Variously combined with shrubs (i.e. thallochamaephytes) with branches that may be embedded (e.g., <i>Loiseleuria</i> heath). |
| 3223 | Cushion Dwarf-Shrubland | level 4 | Within Obligate Drought-Deciduous Dwarf-Shrubland or Dwarf-Thicket (322). Shrubs are isolated in clumps forming dense cushions and are often thorny. |
| 3224 | Mixed Dwarf-Shrubland | level 4 | Within Obligate Drought-Deciduous Dwarf-Shrubland or Dwarf-Thicket (322). Deciduous and evergreen dwarf-shrubs, caespitose herbaceous plants, succulent perennial herbs, and other species intermixed. |
| 323 | Cold-Deciduous | level 3 | Within Mainly Deciduous Dwarf-Shrubland or Dwarf-Thicket (32). Densely closed dwarf-shrubs shed foliage at the beginning of a cold season. Richer in mosses and ferns than the Obligate Drought-deciduous Dwarf Thicket or Shrubland class (322). |
| 3231 | Caespitose Dwarf-Thicket | level 4 | Within Cold-deciduous Dwarf-Shrubland or Dwarf-Thicket (323). Shrub branches stand upright and are often occupied by lichens (foliose). Cushion-shaped mosses, lichens and other herbaceous plants are often found on the ground. |
| 3232 | Creeping Dwarf-Thicket | level 4 | Within Cold-deciduous Dwarf-Shrubland or Dwarf-Thicket (323). Shrub branches creep along the ground; combined with shrubs with branches that may be embedded. |
| 3233 | Cushion Dwarf-Shrubland | level 4 | Within Cold-deciduous Dwarf-Shrubland or Dwarf-Thicket (323). Shrubs are isolated in clumps forming dense cushions and are often thorny. |
| 3234 | Mixed Dwarf-Shrubland | level 4 | Within Cold-deciduous Dwarf-Shrubland or Dwarf-Thicket (323). Deciduous and evergreen dwarf-shrubs, caespitose herbaceous plants, succulent perennial herbs, and other species intermixed. |
| 33 | Extremely Xeromorphic (Subdesert) Dwarf-Shrubland | level 2 | Within Dwarf-Shrubland or Dwarf-Thicket (3). Composed of open formations of dwarf-shrubs, succulents, and herbaceous plants adapted to survive or to avoid a long dry season. Mostly subdesertic. |
| 331 | Mainly Evergreen | level 3 | Within Extremely Xeromorphic Dwarf-Shrubland (33). The canopy is never without green foliage. At least 50% of the shrubs that reach the canopy are evergreen. In extremely dry years some leaves and shoot portions may be shed. |
| 3311 | Purely Evergreen | level 4 | Within Mainly Evergreen Extremely Xeromorphic Dwarf-Shrubland (331). Composed of broad-leaved mostly sclerophyllous shrubs, leafless green-stemmed plants, or succulents dominated by variously branched stem and leaf succulents. |
| 3312 | Semi-Deciduous | level 4 | Within Mainly Evergreen Extremely Xeromorphic Dwarf-Shrubland (331). May consist of either facultatively deciduous shrubs or a combination of evergreen and deciduous shrubs (i.e. evergreen shrubs are dominant, deciduous shrubs cover more than 25%). |
| 332 | Mainly Deciduous | level 3 | Within Extremely Xeromorphic Dwarf-Shrubland (33). The majority of shrubs (more than 50% of the canopy) shed their foliage simultaneously in connection with the unfavorable season (cold or drought). |
| 3321 | Without Succulents | level 4 | Within Mainly Deciduous Extremely Xeromorphic Dwarf-Shrubland (332). Succulents cover less than 25% of the ground. |

| | | | |
|------|------------------------------------|---------|---|
| 3322 | With Succulents | level 4 | Within Mainly Deciduous Extremely Xeromorphic Dwarf-Shrubland (332). Succulents cover at least 25% of the ground. |
| 34 | Tundra | level 2 | Within <i>Dwarf-Shrubland or Dwarf-Thicket</i> (3). Slowly growing, low formations, consisting mainly of dwarf-shrubs, graminoids, mosses, liverworts and lichens, found beyond the subpolar tree line. Often showing plant patterns caused by freezing movements of the soil. Except in boreal regions, dwarf-shrub formations above the mountain tree line should not be called tundra, because they are, as a rule, richer in dwarf-shrubs and grasses, and grow taller due to greater solar radiation in lower latitudes. |
| 341 | Mainly Bryophyte | level 3 | Within <i>Tundra Dwarf-Shrubland or Dwarf-Thicket</i> (34). Dominated by mats or small cushions of chamaephytic mosses (more than 50% of the vegetative cover). Groups of dwarf-shrubs are as a rule scattered irregularly and are not very dense. The general aspect is more or less dark green, olive green or brownish. |
| 3411 | Caespitose | level 4 | Within <i>Mainly Bryophyte Tundra Dwarf-Shrubland or Dwarf-Thicket</i> (341). Clumped or clustered dwarf-shrubs are present. |
| 3412 | Creeping | level 4 | Within <i>Mainly Bryophyte Tundra Dwarf-Shrubland or Dwarf-Thicket</i> (341). Creeping or matted dwarf-shrubs are present. |
| 342 | Mainly Lichen | level 3 | Within <i>Tundra Dwarf-Shrubland or Dwarf-Thicket</i> (34). Mats of fruticose lichens dominate (more than 50% of the vegetative cover), giving the formation a more or less pronounced gray aspect. Mostly evergreen, creeping or cushion-shaped dwarf-shrubs are present. |
| 4 | Herbaceous Vegetation | level 1 | Dominated by herbaceous growth of two major types: graminoids and forbs . Graminoids include all herbaceous grasses and grass-like plants such as sedges (Carex) , rushes (Juncus) and cattails (Typha) . Forbs are broad-leaved herbaceous plants such as clover (Trifolium) , sunflowers (Helianthus) , ferns , and milkweeds (Asclepias) . Total ground coverage must be greater than 60% herbaceous vegetation. |
| 41 | Tall Graminoid | level 2 | Within <i>Herbaceous Vegetation</i> (4). Plant community consists of dominant grasses over 2 meters tall when flowering or mature (more than 50% of the herbaceous vegetation). Forbs may be present but comprise less than 50% of herbaceous vegetation. |
| 411 | With Trees Covering 10-40% | level 3 | Within <i>Tall Graminoid Herbaceous Vegetation</i> (41). May be with or without shrubs. This is somewhat like a very open woodland with more or less continuous ground cover (over 60%) of tall graminoids. |
| 4110 | Trees: Needle-Leaved Evergreen | level 4 | Within <i>Tall Graminoid Herbaceous Vegetation with Trees Covering 10-40%</i> (411). Needle-leaved evergreen species are greater than 50% of the tree canopy. |
| 4111 | Trees: Broad-Leaved Evergreen | level 4 | Within <i>Tall Graminoid Herbaceous Vegetation with Trees Covering 10-40%</i> (411). Broad-leaved evergreen species are greater than 50% of the tree canopy. |
| 4112 | Trees: Broad-Leaved Semi-Evergreen | level 4 | Within <i>Tall Graminoid Herbaceous Vegetation with Trees Covering 10-40%</i> (411). Broad-leaved evergreen species are greater than 50% of the tree canopy and broad-leaved deciduous trees. |
| 4113 | Trees: Broad-Leaved Deciduous | level 4 | Within <i>Tall Graminoid Herbaceous Vegetation with Trees Covering 10-40%</i> (411). Broad-leaved species are greater than 50% of the tree canopy. The area is seasonally flooded. E.g., Northeast Bolivia. |
| 412 | With Trees Covering <10% | level 3 | Within <i>Tall Graminoid Herbaceous Vegetation</i> (41). Grassland with trees covering less than 10% of the ground, with or without shrubs. |
| 4120 | Trees: Needle-Leaved Evergreen | level 4 | Within <i>Trees Covering <10% Tall Graminoid Herbaceous Vegetation</i> (412). Needle-leaved evergreen species are greater than 50% of the tree canopy. |
| 4121 | Trees: Broad-Leaved Evergreen | level 4 | Within <i>Trees Covering <10% Tall Graminoid Herbaceous Vegetation</i> (412). Broad-leaved evergreen species are greater than 50% of the tree canopy. |

| | | | |
|------|--|---------|--|
| 4122 | Trees: Broad-Leaved Semi-Evergreen | level 4 | Within Trees Covering <10% Tall Graminoid Herbaceous Vegetation (412). Trees present are at least 25% each of broad-leaved evergreen and broad-leaved deciduous trees. |
| 4123 | Trees: Broad-Leaved Deciduous | level 4 | Within Trees Covering <10% Tall Graminoid Herbaceous Vegetation (412). Broad-leaved species are greater than 50% of the tree canopy. The area is seasonally flooded. |
| 4124 | Tropical and Subtropical with Trees and Shrubs in Tufts on Termite Nests | level 4 | Within Trees Covering <10% Tall Graminoid Herbaceous Vegetation (412). Tropical or subtropical tall grassland with trees and/or shrubs growing in tufts on termite nests. Also called termite savannah. |
| 413 | With Shrubs | level 3 | Within Tall Graminoid Herbaceous Vegetation (41). The shrub canopy must cover more than 25% of the ground. |
| 4130 | Shrubs: Needle-Leaved Evergreen | level 4 | Within Tall Graminoid Herbaceous Vegetation with Shrubs (413). Needle-leaved evergreen species are greater than 50% of the shrub canopy. |
| 4131 | Shrubs: Broad-Leaved Evergreen | level 4 | Within Tall Graminoid Herbaceous Vegetation with Shrubs (413). Broad-leaved evergreen species are greater than 50% of the shrub canopy. |
| 4132 | Shrubs: Broad-Leaved Semi-Evergreen | level 4 | Within Tall Graminoid Herbaceous Vegetation with Shrubs (413). Shrubs present are at least 25% each of broad-leaved evergreen and broad-leaved deciduous. |
| 4133 | Shrubs: Broad-Leaved Deciduous | level 4 | Within Tall Graminoid Herbaceous Vegetation with Shrubs (413). Broad-leaved species are greater than 50% of the shrub canopy. The area is seasonally flooded. |
| 4134 | Tropical and Subtropical with Trees and Shrubs in Tufts on Termite Nests | level 4 | Within Tall Graminoid Herbaceous Vegetation with Shrubs (413). Tropical or subtropical tall grassland with trees and/or shrubs growing in tufts on termite nests. Also called termite savannah. |
| 414 | With Tuft Plants | level 3 | Within Tall Graminoid Herbaceous Vegetation (41). The canopy of the tuft plants (usually palms) must cover more than 25% of the ground. |
| 4141 | Tropical with Palms | level 4 | Within Tall Graminoid Herbaceous Vegetation with Tuft Plants (414). Tropical grasslands with palms. E.g., the palm savannas of Aroconia totai and Attalea princeps north of Santa Cruz de la Sierra, Bolivia. |
| 415 | Without Woody Synusia | level 3 | Within Tall Graminoid Herbaceous Vegetation (41). Grasslands without trees or shrubs. |
| 4151 | Tropical | level 4 | Within Tall Graminoid Herbaceous Vegetation Without Woody Synusia (415). Tropical grassland as in various low-latitude regions of Africa. Often seasonally flooded (e.g., Compos de Varzea of the lower Amazon Valley), (e.g., Papyrus swamps of the upper Nile Valley). |
| 42 | Medium Tall Graminoid | level 2 | Within Herbaceous Vegetation (4). The dominant grasses are 50 cm to 2 m tall when flowering or mature (greater than 50% of the herbaceous vegetation). Forbs may be present but comprise less than 50% of the herbaceous vegetation. |
| 421 | With Trees Covering 10-40% | level 3 | Within Medium Tall Graminoid Herbaceous Vegetation (42). May be with or without shrubs. This is somewhat like a very open woodland with more or less continuous ground cover of medium tall graminoids. |
| 4210 | Trees: Needle-Leaved Evergreen | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with Trees Covering 10-40% (421). Needle-leaved evergreen species are greater than 50% of the tree canopy. |
| 4211 | Trees: Broad-Leaved Evergreen | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with Trees Covering 10-40% (421). Broad-leaved evergreen species are greater than 50% of the tree canopy. |

| | | | |
|------|--|---------|--|
| 4212 | Trees: Broad-Leaved Semi-Evergreen | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Trees Covering 10-40%</i> (421). Trees present are at least 25% each of broad-leaved evergreen and broad-leaved deciduous trees. |
| 4213 | Trees: Broad-Leaved Deciduous | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Trees Covering 10-40%</i> (421). Broad-leaved species are greater than 50% of the tree canopy. The area is seasonally flooded. |
| 422 | With Trees Covering <10% | level 3 | Within Medium Tall Graminoid Herbaceous Vegetation (42). Grassland with trees covering less than 10% of the ground, with or without shrubs. |
| 4220 | Trees: Needle-Leaved Evergreen | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Trees Covering <10%</i> (422). Needle-leaved evergreen species are greater than 50% of the tree canopy. |
| 4221 | Trees: Broad-Leaved Evergreen | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Trees Covering <10%</i> (422). Broad-leaved evergreen species are greater than 50% of the tree canopy. |
| 4222 | Trees: Broad-Leaved Semi-Evergreen | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Trees Covering <10%</i> (422). Broad-leaved evergreen species are greater than 50% of the tree canopy. |
| 4223 | Trees: Broad-Leaved Deciduous | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Trees Covering <10%</i> (422). Broad-leaved species are greater than 50% of the tree canopy. The area is seasonally flooded. |
| 4224 | Tropical and Subtropical with Trees and Shrubs in Tufts on Termite Nests | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Trees Covering <10%</i> (422). Tropical or subtropical medium tall grassland with trees and/or shrubs growing in tufts on termite nests. Also called termite savannah. |
| 423 | With Shrubs | level 3 | Within Medium Tall Graminoid Herbaceous Vegetation (42). The shrub canopy must cover more than 25% of the ground. |
| 4230 | Shrubs: Needle-Leaved Evergreen | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Shrubs (423)</i> . Needle-leaved evergreen species are greater than 50% of the shrub canopy. |
| 4231 | Shrubs: Broad-Leaved Evergreen | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Shrubs (423)</i> . Broad-leaved species are greater than 50% of the shrub canopy. |
| 4232 | Shrubs: Broad-Leaved Semi-Evergreen | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Shrubs (423)</i> . Shrubs present are at least 25% each of broad-leaved evergreen and broad-leaved deciduous. |
| 4233 | Shrubs: Broad-Leaved Deciduous | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Shrubs (423)</i> . Broad-leaved evergreen species are greater than 50% of the shrub canopy. The area is seasonally flooded. |
| 4234 | Tropical and Subtropical with Trees and Shrubs in Tufts on Termite Nests | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Shrubs (423)</i> . Tropical or subtropical medium tall grassland with trees and/or shrubs growing in tufts on termite nests. Also called termite savannah. |
| 4235 | Woody Symusia of Deciduous Thorny Shrubs | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation with <i>Shrubs (423)</i> . Consists of deciduous thorny shrubs covering at least 25% of the ground. E.g., the tropical thorn bush savannah of the Sahel region in Africa with <i>Acacia tortilis</i> , <i>A. senegal</i> and other species. |
| 424 | Open Symusia of Tuft Plants | level 3 | Within Medium Tall Graminoid Herbaceous Vegetation (42). The canopy of the tuft plants (usually palms) must cover more than 25% of the ground. |

| | | | |
|------|--|---------|--|
| 4241 | Subtropical with Open Palm Groves | level 4 | Within <i>Open Synusia of Tuft Plants</i> Medium Tall Graminoid Herbaceous Vegetation (424). Medium tall grassland with open groves of palms (e.g., Corrientes, Argentina). Some areas are seasonally flooded (e.g., Mauritia palm groves in the Colombian and Venezuelan llanos). |
| 425 | Without Woody Synusia | level 3 | Within Medium Tall Graminoid Herbaceous Vegetation (42). Medium tall grasslands without trees or shrubs (less than 25% of the ground). |
| 4251 | Mainly Sod Grasses | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation Without Woody Synusia (425). Perennial, highly branched, creeping grass, which binds the sand or soils with its root system. E.g., St. Augustine grass (<i>Stenotaphrum secundatum</i>), the tall-grass prairie in eastern Kansas, or on sandy soil or dunes, such as the communities of <i>Andropogon hallii</i> in the Nebraska Sand Hills. In some locations the grassland is wet or flooded most of the year (e.g., <i>Typha</i> swamps). If that is the case classify as a wetland. See MUC class 6. |
| 4252 | Mainly Bunch Grasses | level 4 | Within Medium Tall Graminoid Herbaceous Vegetation Without Woody Synusia (425). Grasses that chiefly grow in tufts forming an irregular textured surface. E.g., the hard tussock (<i>Festuca novae-zelandiae</i>) grasslands in New Zealand. |
| 43 | Short Graminoid | level 1 | Within Herbaceous Vegetation (4). The dominant grasses are less than 50 cm tall when flowering or mature (more than 50% of the herbaceous vegetation). Forbs may be present but they comprise less than 50% of the herbaceous vegetation. |
| 431 | With Trees Covering 10-40% | level 3 | Within Short Graminoid Herbaceous Vegetation (43). May be with or without shrubs. This is somewhat like a very open woodland with more or less continuous ground cover of short graminoids. |
| 4310 | Trees: Needle-Leaved Evergreen | level 4 | Within Short Graminoid Herbaceous Vegetation with Trees Covering 10-40% (431). Needle-leaved evergreen species are greater than 50% of the tree canopy. |
| 4311 | Trees: Broad-Leaved Evergreen | level 4 | Within Short Graminoid Herbaceous Vegetation with Trees Covering 10-40% (431). Broad-leaved evergreen species are greater than 50% of the tree canopy. |
| 4312 | Trees: Broad-Leaved Semi-Evergreen | level 4 | Within Short Graminoid Herbaceous Vegetation with Trees Covering 10-40% (431). Trees present are at least 25% each of broad-leaved evergreen and broad-leaved deciduous trees. |
| 4313 | Trees: Broad-Leaved Deciduous | level 4 | Within Short Graminoid Herbaceous Vegetation with Trees Covering 10-40% (431). Broad-leaved species are greater than 50% of the tree canopy. The area is seasonally flooded. |
| 432 | With Trees Covering <10% | level 3 | Within Short Graminoid Herbaceous Vegetation (43). Grassland with trees covering less than 10% of the ground, with or without shrubs. |
| 4320 | Trees: Needle-Leaved Evergreen | level 4 | Within Short Graminoid Herbaceous Vegetation with Trees Covering <10% (432). Needle-leaved evergreen species are greater than 50% of the tree canopy. |
| 4321 | Trees: Broad-Leaved Evergreen | level 4 | Within Short Graminoid Herbaceous Vegetation with Trees Covering <10% (432). Broad-leaved evergreen species are greater than 50% of the tree canopy. |
| 4322 | Trees: Broad-Leaved Semi-Evergreen Deciduous | level 4 | Within Short Graminoid Herbaceous Vegetation with Trees Covering <10% (432). Trees present are at least 25% each of broad-leaved evergreen and broad-leaved deciduous trees. |
| 4323 | Trees: Broad-Leaved Deciduous | level 4 | Within Short Graminoid Herbaceous Vegetation with Trees Covering <10% (432). Broad-leaved species are greater than 50% of the tree canopy. The area is seasonally flooded. |
| 4324 | Tropical and Subtropical with Trees and Shrubs in Tufts on Termite Nests | level 4 | Within Short Graminoid Herbaceous Vegetation with Trees Covering <10% (432). Tropical or subtropical short grassland with trees and/or shrubs growing in tufts on termite nests. Also called termite savannah. |

| | | | |
|------|--|---------|---|
| 433 | With Shrubs | level 3 | Within Short Graminoid Herbaceous Vegetation (43). The shrub canopy must cover more than 25% of the ground. |
| 4330 | Shrubs: Needle-Leaved Evergreen | level 4 | Within Short Graminoid Herbaceous Vegetation with Shrubs (433). Needle-leaved evergreen species are greater than 50% of the shrub canopy. |
| 4331 | Shrubs: Broad-Leaved Evergreen | level 4 | Within Short Graminoid Herbaceous Vegetation with Shrubs (433). Broad-leaved evergreen species are greater than 50% of the shrub canopy. |
| 4332 | Shrubs: Broad-Leaved Semi-Evergreen | level 4 | Within Short Graminoid Herbaceous Vegetation with Shrubs (433). Shrubs present are at least 25% each of broad-leaved evergreen and broad-leaved deciduous. |
| 4333 | Shrubs: Broad-Leaved Deciduous | level 4 | Within Short Graminoid Herbaceous Vegetation with Shrubs (433). Broad-leaved species are greater than 50% of the shrub canopy. The area is seasonally flooded. |
| 4334 | Tropical and Subtropical with Trees and Shrubs in Tufts on Termite Nests | level 4 | Within Short Graminoid Herbaceous Vegetation with Shrubs (433). Tropical or subtropical short grassland with trees and/or shrubs growing in tufts on termite nests. Also called termite savannah. |
| 4335 | Woody Symusia of Deciduous Thorny Shrubs | level 4 | Within Short Graminoid Herbaceous Vegetation with Shrubs (433). Consists of deciduous thorny shrubs covering at least 25% of the ground. |
| 434 | Open Symusia of Tuft Plants | level 3 | Within Short Graminoid Herbaceous Vegetation (43). The canopy of the tuft plants (usually palms) must cover more than 25% of the ground. |
| 4341 | Subtropical with Open Palm Groves | level 4 | Within Open Symusia of Tuft Plants Short Graminoid Herbaceous Vegetation (434). Short grassland with open groves of palms. The canopy of palms must cover more than 25% of the ground. |
| 435 | Mainly Bunch Grasses with Woody Symusia | level 3 | Within Short Graminoid Herbaceous Vegetation (43). Grasses that grow in tufts, with woody plants interspersed. |
| 4351 | Tropical Alpine with Tuft Plants | level 4 | Within Mainly Bunch Grasses with Woody Symusia Short Graminoid Herbaceous Vegetation (435). This grassland often contains <i>Espeletia</i> , <i>Lobelia</i> , <i>Senecio</i> , microphyllous dwarf-shrubs, and cushion plants (often with woolly leaves). Above the timberline in low latitudes. E.g., Paramo and related vegetation types without snow in the alpine regions of Kenya, Colombia, Venezuela, etc. |
| 4352 | Tropical Alpine without Tuft Plants | level 4 | Within Mainly Bunch Grasses with Woody Symusia Short Graminoid Herbaceous Vegetation (435). Similar to Tropical Alpine with Tuft Plants (4351) but very open and without tuft plants. In these grasslands there is frequent nocturnal snowfall (though the snow is gone by 9 a.m.). E.g., the Super-Paramo (i.e. above Paramo) of J. Cuatrescasas. |
| 4353 | Tropical and Subtropical Alpine with Open Stands of Evergreens | level 4 | Within Mainly Bunch Grasses with Woody Symusia Short Graminoid Herbaceous Vegetation (435). This grassland may also have deciduous shrubs and dwarf shrubs. E.g., Puna south of Oruro, Bolivia. |
| 4354 | With Dwarf Shrubs | level 4 | Within Mainly Bunch Grasses with Woody Symusia Short Graminoid Herbaceous Vegetation (435). Consists of bunch grass with varying coverage of dwarf shrubs. Cushion plants may also grow in this grassland, and may be locally more important than the dwarf-shrubs. E.g., Puna south of Oruro, Bolivia. |
| 436 | Without Woody Symusia | level 3 | Within Short Graminoid Herbaceous Vegetation (43). Short grasslands without trees or shrubs. |

| | | | |
|------|---|---------|---|
| 4361 | Short-Grass Communities | level 4 | Within <i>Short Graminoid Herbaceous Vegetation Without Woody Synusia</i> (436). These communities may fluctuate in structure and floristic composition due to greatly fluctuating precipitation of the semi-arid climate. E.g., short-grass (<i>Bouteloua gracilis</i> and <i>Buchloe dactyloides</i>) prairie of eastern Colorado. |
| 4362 | Bunch-Grass Communities | level 4 | Within <i>Short Graminoid Herbaceous Vegetation Without Woody Synusia</i> (436). E.g., blue tussock (<i>Poa cloanoides</i>) communities of New Zealand, and alpine dry Puna with <i>Festuca orthophylla</i> of northern Chile and southern Bolivia. |
| 437 | Short to Medium Tall Mesophytic Communities | level 3 | Within <i>Short Graminoid Herbaceous Vegetation</i> (43). Plants growing in or adapted to a moderately moist environment. |
| 4371 | Sod Grass Communities | level 4 | Within <i>Short to Medium Tall Mesophytic Communities Short Graminoid Herbaceous Vegetation</i> (437). The grassland is often rich in forbs, and occurs in lower altitudes with a cool, humid climate in North America and Eurasia. Many plants may remain at least partly green during the winter, even below the snow in the higher latitudes. |
| 4372 | Alpine and Subalpine Meadows | level 4 | Within <i>Short to Medium Tall Mesophytic Communities Short Graminoid Herbaceous Vegetation</i> (437). These grasslands are usually moist much of the summer due to snow melt water. May be rich in forbs (e.g., Olympic Peninsula, Washington); rich in dwarf-shrubs (e.g., the Rocky Mountains of Colorado); snow-bed communities rich in small forbs and/or forb-like dwarf-shrubs (e.g., <i>Salix herbacea</i>); or avalanche meadows, occurring as narrow strips of grassland between forests on steep slopes of high mountains where avalanches, descending annually in spring, prevent forest growth. |
| 44 | Forb Vegetation | level 2 | Within <i>Herbaceous Vegetation</i> (4). Broad-leaved herbaceous plants dominate the plant community, such as clover, sunflowers (<i>Helianthus</i>), ferns, and milkweeds (<i>Asclepias</i>) (all plants except grasses). Forbs cover at least 50% of the herbaceous area. Grasses may be present but often less than (often much less than) 50%. |
| 441 | Tall Communities | level 3 | Within <i>Forb Herbaceous Vegetation</i> (44). The dominant forb growth forms are more than 1 meter tall when fully developed. |
| 4411 | Fern Thickets | level 4 | Within <i>Tall Forb Communities Herbaceous Vegetation</i> (441). Ferns occur sometimes in nearly pure stands, especially in humid climates (e.g., <i>Pteridium aquilinum</i>). |
| 4412 | Mainly Annual | level 4 | Within <i>Tall Forb Communities Herbaceous Vegetation</i> (441). Annual forbs, which germinate in the beginning and die at the end of each growing season, are the dominant form (greater than 50% of forb vegetation). |
| 4413 | Mainly Perennial Flowering Forbs and Ferns | level 4 | Within <i>Tall Forb Communities Herbaceous Vegetation</i> (441). Some part of the plant is alive all year round. |
| 442 | Low Communities | level 3 | Within <i>Forb Herbaceous Vegetation</i> (44). These communities are dominated by forbs less than 1 meter tall when fully developed. |
| 4421 | Mainly Perennial Flowering Forbs and Ferns | level 4 | Within <i>Low Forb Communities Herbaceous Vegetation</i> (442). Some part of the plant is alive all year round. |

| | | | |
|------|----------------------|---------|--|
| 4422 | Mainly Annual | level 4 | Within Low Forb Communities Herbaceous Vegetation (442). Annual forbs, which germinate in the beginning and die at the end of each growing season, are the dominant form (greater than 50% of forb vegetation). There are several types of low annual forbs: <i>Ephemeral forb communities in tropical and subtropical regions:</i> Forbs grow with very little precipitation where, from autumn to spring, clouds moisten vegetation and soil. The dry season aspect is desert-like. E.g., the coastal hills of Peru and northern Chile <i>Ephemeral or episodical forb communities of arid regions:</i> The “flowering desert” consists of mostly fast growing forbs, sometimes concentrated in depressions where water can accumulate in shrub or dwarf shrub formations of arid regions. E.g., the Sonoran Desert. |
| 5 | Barren Land | level 1 | Land with less than 40% vegetative cover. Barren land has a limited ability to support life, and is usually made up of thin soil, sand, or rocks. |
| 51 | Dry Salt Flats | level 2 | Occur on flat floored bottoms of interior desert basins. High concentrations of salts are present due to extensive water evaporation. |
| 52 | Sandy Areas | level 2 | Accumulations of sand/gravel (e.g., beaches or dunes). |
| 53 | Bare Rock | level 2 | Exposed bedrock, desert pavement, scarpas, talus slides, volcanic material, rock glaciers and other accumulations of rock without vegetative cover. |
| 54 | Perennial Snowfields | level 2 | Accumulations of snow and ice that did not entirely melt during the previous summer, occurring where the daily average temperature is 0°C (32°F) in the warmest summer months. |
| 55 | Glaciers | level 2 | Snow compacted into firm and finally to ice under weight of successive annual accumulations. Re-frozen melt water contributes to increasing density of the glacial ice mass. All glaciers exhibit evidence of present or past motions (moraines, crevasses, etc.). |
| 56 | Other | level 2 | Dirt, gravel, other loose rock, etc. |
| 6 | Wetland | level 1 | Marsches, swamps, bogs and other types of wetlands that are periodically or constantly saturated during the growing season. This periodic or constant saturation produces soils with special chemical characteristics and vegetation specifically adapted to wet conditions. The area must have at least 40% vegetative cover to be classified as a wetland. |
| 61 | Riverine | level 2 | Wetlands adjacent to a fresh water river channel (riparian wetlands). |
| 62 | Palustrine | level 2 | Wetlands dominated by trees, shrubs, persistent emergents (plants), mosses, lichens, etc. The wetlands surround water that is less than 1 hectare in size, has no active channel or tide, is less than 2 meters deep, and has low salinity. The water should be included as part of the wetland. |
| 63 | Estuarine | level 2 | Wetlands occurring adjacent to a tidal channel, or in and adjacent to the intertidal zone. An estuary is a water passage where the tide meets the current of a stream. Deepwater tidal habitats and adjacent tidal wetlands are usually semi-enclosed by land but have open, partially obstructed, or sporadic access to ocean water (at least occasionally diluted by freshwater runoff from the land). |
| 64 | Lacustrine | level 2 | Wetlands surrounding open water (e.g., ponds and lakes) that are greater than 1 hectare in size and greater than 2 meters deep. |
| 7 | Open Water | level 1 | Lakes, ponds, rivers and oceans. The surface of the land is continually submerged by water greater than 2 meters deep and at least one hectare in size; or continually submerged in an actively flowing channel or subtidal zone. Water should cover greater than 60% of the area. |

| | | | |
|----------|----------------------------|----------------|---|
| 71 | Freshwater | level 2 | Lakes, ponds, and rivers with low salinity. |
| 72 | Marine | level 2 | Open ocean overlying the continental shelf or an actively flowing tidal channel. |
| 8 | Cultivated Land | level 1 | The ground is covered by greater than 60% non-native cultivated species (e.g., agricultural crops, cultivated short grasses, and lawns) and usually can be distinguished by the regular geometric patterns created by the lawns and fields. |
| 81 | Agriculture | level 2 | Land is used for growing crops, orchards, horticulture, feeding livestock, and other agriculture. |
| 811 | Row Crop and Pasture | level 3 | Examples include corn, wheat, cow pastures, fallow fields, cultivated cranberry bogs, and rice fields. |
| 812 | Orchard and Horticulture | level 3 | Examples include apple orchards, vineyards, and tree nurseries. |
| 813 | Confined Livestock Feeding | level 3 | These areas are found on large farms and are used for feeding beef cattle, dairy cows (with confined feedlots), hogs and poultry. |
| 814 | Other Agriculture | level 3 | Examples include corrals and breeding and training facilities on horse farms. |
| 82 | Non-Agriculture | level 2 | Land is used for parks, playing fields, cemeteries, and golf courses. |
| 821 | Parks and Athletic Fields | level 3 | Examples include baseball diamonds, soccer fields, play grounds, and parks. |
| 822 | Golf Courses | level 3 | Golf Courses |
| 823 | Cemeteries | level 3 | Cemeteries |
| 824 | Other Non-Agriculture | level 3 | Any other non-agricultural cultivated areas that do not fit into classes 821, 822 or 823 (parks and playing fields, golf courses, or cemeteries). |
| 9 | Urban | level 1 | Areas developed for residential, commercial, industrial, or transportation uses. Must be greater than 40% urban land cover. |
| 91 | Residential | level 2 | Greater than 50% of the urban land cover consists of residential property (e.g., apartments, private dwellings) |
| 92 | Commercial and Industrial | level 2 | Greater than 50% of the urban land cover consists of commercial or industrial property (e.g., businesses, factories, warehouses) |
| 93 | Transportation | level 2 | Greater than 50% of the urban land cover consists of transportation routes (e.g., roads, highways, railroads, and airport runways). |
| 94 | Other | level 2 | At least 50% of the urban land cover consists of developed areas that do not fit into residential, commercial, or transportation categories. |

| Miscellaneous Definitions | |
|---------------------------|--|
| Annual Plant | Live and grows for only one year or season. |
| Aspect | View or appearance; a side facing a particular direction. |
| Boreal | Also called cold temperate zone has a climate with cool wet summers and cold winters lasting more than six months. |

| | |
|--------------------|--|
| Bryophyte | Non-flowering plants (mosses & liverworts) characterized by rhizoids rather than true roots. |
| Buttresses | Flanges of tissue protruding from the trunk of a tree, tapering outward at the base to give support. Common among large tropical trees. |
| Caespitose | Arranged or combined in a thick mat or clumps, having a low stem forming a dense turf or sod, growing in clusters. |
| Canopy | Uppermost layer of vegetation detected by satellite sensors. |
| Chamaephyte | A perennial plant that has its winter buds placed very close to the soil surface. |
| Cold-Deciduous | Plants that shed leaves during the cold season. |
| Crustose Lichens | Lichens that are encrusting. E.g., <i>Caloplaca saxicola</i> . |
| Cultivated Land | Landscaped yards, playing fields, cemeteries, golf courses, and other cultivated vegetated areas should be classified as cultivated land (class 8) if non-native cultivated species is greater than 60% coverage. If the buildings, roads and unnatural structures (bridges, etc.) cover greater than 40% of the land, the area should be classified as urban. If wooded residential neighborhoods have greater than 40% trees covering the ground, the area would be considered forest or woodlands (see classes 0 and 1). If it is difficult to decide upon a cover type, try to determine what the satellite would see. Compare similar areas with the satellite image you receive of your school's location. |
| Deciduous | Vegetation that sheds its leaves at the end of the growing period or in association with the unfavorable season (drought, cold). |
| Drip Tips | Extended slender tips of tropical leaves that allow water to roll off the leaf surface. |
| Drought-Deciduous | Plants that shed their leaves during the dry season. |
| Emergent | Aquatic plant with the lower part submerged and the upper part extending above the water. |
| Epiphytes | Plants not connected with the soil, that grow on another plant (upon which it depends for mechanical support) but not for receiving food and water from it, such as certain orchids or ferns. |
| Facultative | Organisms able to live and thrive under more than one set of conditions. |
| Firm | Snow that has been partially consolidated, or compacted, by thawing and freezing but not yet converted to glacial ice. |
| Forb | A broad-leaved herbaceous plant other than a grass such a clover, sunflowers, ferns, and milkweeds. |
| Fruticose Lichens | Lichens which appear shrubby or hair-like, especially in form. |
| Graminoid | Grasses and grass-like plants. |
| Herbaceous | Pertaining to or characteristic of an herb as distinguished from a woody plant. Vascular plant rooted in the ground with foliage that dies back annually. The meristem (stem growth tip) is located just above or below the ground. |
| Hygromorphic | The form of the plant is altered due to changes in moisture in the plant. E.g., hygromorphic chamaephytes <i>Selaginella</i> and herbaceous ferns. |
| Hemisclerophyllous | Vegetation with slightly thickened foliage, with large soft leaves, that is resistant to water loss. E.g., subalpine <i>Rhododendron</i> thickets, or <i>Hibiscus tiliaceus</i> matted thickets of Hawaii. |
| Lichen | Plant made up of an alga and a fungus living in a symbiotic relationship. Specifically, any of a numerous plants consisting of a fungus, usually of the class <i>Ascomycetes</i> , in close combination with certain of the green or blue-green algae, characteristically forming a crustlike, scaly, or branching growth on rocks or tree trunks. |

| | |
|--------------------|---|
| Lignified | Woody, hardened. Has formed or turned into wood through the formation and deposit of lignin in the cell walls. |
| Lowland | An area of land that is low in relation to the surrounding country. It may be necessary to consult local resources to determine the specific classification. Vegetation will vary depending on both the latitude and the altitude. |
| Mesophytic | Growing in, or adapted to, a moderately moist environment. |
| Microphanerophytes | Small flowering plants. |
| Microphyllous | Having small leaves with a single unbranched vein (e.g., desert plants). |
| Montane | Of, growing in, or inhabiting mountain areas. It may be necessary to consult local resources to determine the specific classification. Vegetation will vary depending on both the latitude and the altitude. |
| Nanophanerophytes | Very small flowering plants. |
| Obligate | Organisms restricted to a particular condition of life (that condition is essential for survival). |
| Overstory | Uppermost layer of vegetation detected by satellite sensors. |
| Perennial Plant | Has a life span of more than two years. |
| Polar | In polar climates, the mean temperature of the warmest month is below 10°C and there is low precipitation distributed over the entire year. There is a short, wet, nightless summer and a very long, cold, dark winter. Generally, the climate is too cold to support the growth of trees. |
| Saturated | Soaked with moisture - the maximum water holding capacity of a soil. |
| Scapose | Having a leafless flower stalk growing directly from the ground. E.g., agave/century plant. |
| Sclerophyllous | Vegetation with thickened, hardened foliage that is resistant to water loss (sclerophyll). E.g., plants of the chaparral (semi-arid Mediterranean) such as toyon, ironwood, manzanita, coyote bush, mountain mahogany, and black sage. |
| Subalpine | Of, designating, or growing or living in mountainous regions just below the timberline. It may be necessary to consult local resources to determine the specific classification. Vegetation will vary depending on both the latitude and the altitude. |
| Submontane | Located under or at the base of a mountain or mountain range. It may be necessary to consult local resources to determine the specific classification. Vegetation will vary depending on both the latitude and the altitude. |
| Subpolar | Transitional between the cold temperate zone and the polar zone. It may be necessary to consult local resources to determine the specific classification. Vegetation will vary depending on both the latitude and the altitude. |
| Subtropical | From the edge of the tropical zone toward the poles, in the region of the descending air masses, which get warmer as it descends and becomes very dry. Rainfall is very low, and the daytime temperatures are very high because of intense solar radiation. In the winter months, however, the temperature may sink to zero at night as a result of the greater net loss of heat energy in outgoing radiation. This is the hot desert zone. |
| Succulent | Having thickened, juicy, fleshy tissues (leaves or stems), more or less soft in texture, that conserve moisture. E.g., a sedum or a cactus. |
| Suffruticose | Has a woody stem or base and is somewhat shrubby. |
| Synusia | A layer or stratum of a community. A structural unit of a major ecological community characterized by relative uniformity of life form or of height and usually constituting a particular stratum of that community. |

| | |
|-------------|---|
| Temperate | <p>Temperate zones show greater seasonal temperature changes and can be broken down as follows:</p> <p><i>Warm temperate</i>: mild or no winter and extremely wet, especially in summer.</p> <p><i>Typical temperate</i>: cold, short winters or a winter free of frost and with very cool summers (near the ocean) (e.g., central European or coastal northeastern U.S.A.).</p> <p><i>Arid temperate</i>: large temperature contrasts between summer and winter, and little precipitation.</p> <p><i>Boreal or cold temperate</i>: cool wet summers and cold winters lasting more than six months.</p> |
| Tropical | <p>Lies 40 degrees to the north and south of the equator. A certain seasonal variation in the mean daily temperature is noticeable. Rainfall reaches a maximum in the summer, with a dry season in the cool months. The duration of the cool season increases as the distance from the equator becomes greater, and at the same time the annual rainfall decreases.</p> |
| Understory | <p>Layer of vegetation that grows beneath the overstory consisting of smaller trees and shrubs.</p> |
| Wet | <p>Vegetation or environments capable of withstanding or thriving in the presence of much rain.</p> |
| Xeromorphic | <p>Climatic conditions favorable for the development of vegetation that is adapted to, thrives in, or tolerates an environment that is poor in available moisture.</p> |
| Xerophyte | <p>A plant which is adapted to and thrives in dry conditions.</p> |