

Historical Research

by Jerry Cox, Ph.D., Science Chair at Carlsbad High School, New Mexico

The first European beef animals to enter the New World were introduced by ship and unloaded in southern Mexico during 1521 (Wagoner 1952). Cattle and sheep were introduced into the semi-desert grasslands in the Chihuahuan and Sonoran Deserts as the Spanish began exploring the northern frontier. Livestock distribution continued to expand as Father Kino and Spanish explorers established missions and trading posts in Arizona and New Mexico. Between 1770 and 1827, Spanish cattlemen were frequently forced from the frontier by Indians, malaria and drought (Bahre 1977). Well over 500,000 cattle freely roamed the frontier, and many were killed for food and hides between 1830 and 1840 (Humphrey 1958).

American botanists and military personnel (Bartlett 1854, Hinton 1890, Barnes 1936) traveling in the valley bottoms of the semi-desert grasslands noted lush sacaton (*Sporobolus airoides* and *Sporobolus wrightii*) and gramagrasses (*Bouteloua spp.*). Upland mesas and canyons were relatively brush-free and dominated by gramagrasses and bluestems (*Andropogon spp.*). Brushy species such as mesquite (*Prosopis spp.*), catclaw and whitethorn (*Acacia spp.*), creosotebush (*Larrea tridentata*) and tarbush (*Flourensia cernua*) were present but limited in numbers. Early explorers consistently write about: (1) their inability to find firewood between Midland TX and Wilcox AZ, and (2) the constant burning of grasslands by Indians in southwest TX, southern NM and southeastern AZ (Humphrey 1958).

After the American exploration and the control of most Indian tribes, American and Mexican cattlemen freely grazed the semi-desert grasslands (Wagoner 1952). Railroad expansion, following the American Civil War and the revolutions in Mexico, provided rapid transportation and movement of agricultural and mining products (Hastings and Turner 1965). As human population increased so did the demand for meat, vegetables and fiber. Cattle, sheep, goats and pigs were driven or shipped from eastern Texas and northeastern Mexico into southern New Mexico to graze the mesas and canyons; while fertile lowlands were plowed and rivers re-channeled to provide irrigation for crops. Most water sources dried up in 1893, resulting in the death of 50 to 75 percent of the livestock population. The drought was over by 1895, but the combined effects of overgrazing the uplands, drought and flooding had resulted in accelerated sheet and gully erosion and the loss of shallow water tables in valley bottoms (Bryan 1925).

Griffith (1901) reviewed the deterioration of southwestern rangelands in 1899 and published statements of area ranchers. H. C. Hooker, owner of the Sierra Bonita Ranch in southeastern Arizona, described the destructive events: “In 1870 there were large beds of sacaton and grama grasses and the river ran shallow with banks covered with grass, shrubs and cottonwood trees. In 1900 the river bed had dropped 20 feet and the vegetation had been removed by grazing, farming and flooding.” Commenting on livestock numbers, Hooker stated: “There were fully 50,000 head of cattle at the upper

end of the valley in 1890. In 1900 there were not more than one-half that number, and they are doing poorly.” C. J. Bayless, another Arizona-New Mexico rancher, stated: *“Beaver dams checked water flow in 1885, but the trappers exterminated the population, and within 5 years the channel was from 3 to 20 feet deep.”* Bayless grazed 40,000 cattle in 1888, but in 1901 there was insufficient forage for 3,000. Other observations (Bartlett 1854, Hinton 1890, Barnes 1925, Bryan 1925) indicate similar vegetation changes throughout the Chihuahuan and Sonoran Deserts.

Historical records clearly illustrate that southern New Mexico and southeastern Arizona desert grasslands were overgrazed and deteriorated rapidly between 1880 and 1930 (Bahre 1977, Bartlett 1854, Cooke and Reeves 1976, Humphrey 1958, York and Dick-Peddie 1969, Herbel *et al.* 1973, Hastings and Turner 1965). Griffith (1901) and Cox *et al.* (1983) documented changes in vegetation, corresponding livestock losses, and reductions in livestock populations.

Cooperative studies to restore desert grassland ecosystems were initiated in the early 1900s by the Division of Agrostology (USDA) and the State Experimentation Stations at Las Cruces, NM and Tucson, AZ. Between 1890 and 1980, the seed of more than 300 forb, grass, shrub and tree species were sown at 400 planting sites within the Chihuahuan and Sonoran Deserts (Cox *et al.* 1982). Fourteen of the species were widely adapted in both deserts. Eleven of the adapted species were introduced perennial grasses, two were native perennial grasses, and one was a native perennial shrub.

Because native plants were more difficult to establish from seed, state and Federal scientists, and land managers usually seeded introduced plants. This trend began to change in the 1970s, when it was recognized that introduced plants often replaced native plant populations. Between 1970 and 1995, a series of scientific investigations were initiated to compare establishment characteristics of both native and introduced plants, and to develop revegetation techniques to aid in the establishment of native plants (Cox *et al.* 1982, Friseur *et al.* 1985, Cox *et al.* 1987).

Plant establishment from seed requires a viable seed and a favorable environment for seedling development. Seed often germinate, but seedlings fail to survive because inadequate soil moisture has limited the development of a root system capable of supporting the plant through later periods of less favorable soil moisture conditions. Available soil moisture was a factor in the establishment of blue grama (*Bouteloua gracilis*), crested wheatgrass (*Agropyron desertorum*) and Russian wildrye (*Elymus junceus*) in cold deserts (Briske and Wilson 1977, 1978, 1980, Wilson and Briske 1979; Hassanyar and Wilson 1978), and sideoats grama (*Bouteloua curtipendula*), Lehmann lovegrass (*Eragrostis lehmanniana*), Boer lovegrass (*Eragrostis curvula*) and blue panicgrass (*Panicum antidotale*) in hot deserts (Frasier *et al.* 1984, 1985). Water stress during germination and seedling development was detrimental to the survival of cool- and warm-season grasses, shrubs and trees from Arizona to Texas (Knipe 1968, 1973; Piatt 1976; Kruse 1970; Scifres and Brock 1969).

Wilson and Briske (1979b), Frasier *et al.* (1984) and Cox *et al.* (1993) found that grasses, shrubs and trees required four to six days of moist soil conditions and soil temperatures between 15 and 35°C to germinate and initiate seminal root growth. To insure adventitious root growth and plant establishment, the first wet period must be followed by a second wet period after two to eight weeks.

In southeastern New Mexico and north central Mexico (the Chihuahuan Desert), the amount and distribution of late summer, and fall-winter precipitation needed to insure native seed germination and seedling establishment can be expected in one of ten years (Herbel *et al.* 1973, Mexico Climatic Reporting Stations 1992, FAO-UNESCO 1975). Because precipitation probabilities are low Cox *et al.* (1989) and Houser (1983) suggested transplanting techniques to establish native perennial plants.

To increase the success of plant establishment, foresters (Tinus 1978), landscape architects (Aratani 1976), mine reclamationists (Howard *et al.* 1978) and wildlife biologists (Springfield 1972, Ferguson *et al.* 1975) have transplanted containerized greenhouse grown seedlings and provided supplemental irrigation during establishment.

Transplant survival was dependent upon a continuous supply of soil moisture for approximately 30 days (Cox *et al.* 1989). Therefore, transplanting dates should follow the expected initiation of the summer rainy season when plants would have the greatest likelihood of success.

Cox and Madrigal (1988) transplanted, and Thacker and Cox (1992) seeded perennial native forage plants on rangelands and abandoned farmland in the Chihuahuan and Sonoran Deserts in North America. The establishment potential on both land types was similar, but the costs associated with clearing rangeland was 10 to 15 times greater than planting and irrigating at abandoned farmland sites, and the survival of transplants was 250 times greater than for seeded plants.

Literature Cited

Aratani, T. 1976. Effectiveness of tubelings -A dryland planting technique. Report FHWA-HI-HWY 73 - 1. Department of Transportation, Honolulu, HI, 15 p.

Bahre, C. J. 1977. Land-use history of the research ranch, Elgin AZ. J. Arizona Academy Science 12: 1 - 32.

Barnes, W. C. 1936. Herds of the San Simon Valley. American Forests 42: 456–457.

Bartlett, J. R. 1854. Personal narrative of explorations and incidences in Texas, New Mexico and California, Sonora and Chihuahua. Volume I. Appleton and Co., NY.

Briske, D. D., and A. M. Wilson. 1977. Temperature effects on adventitious root development in blue grama seedlings. J. Range Management 30: 276 - 280.

- Briske, D. D., and A. M. Wilson. 1978.** Moisture and temperature requirements for adventitious root development in blue grama seedlings. *J. Range Management* 31: 174-178.
- Briske, D. D., and A. M. Wilson. 1980.** Drought tolerance on adventitious root development in blue grama seedlings. *J. Range Management* 33: 323 - 327.
- Bryan, K. 1925.** Date of channel trenching (arroyo cutting) in the arid southwest. *Science* 62: 338 - 344.
- Cooke, R. U., and R. W. Reeves. 1976.** Arroyos and environmental change in the American Southwest. Clarendon Press, Oxford, England.
- Cox, J. R., H. L. Morton, T. N. Johnsen, Jr., G. L. Jordan, S. C. Martin and L. C. Fierro. 1982.** Vegetation restoration in the Chihuahuan and Sonoran Deserts of North America. UDSA-ARS Agricultural Reviews and Manuals. ARM-W-28. Oakland, CA.
- Cox, J. R., H. L. Morton, J. T. LaBaume and K. G. Renard. 1983.** Reviving Arizona's rangelands. *J. Soil and Water Conservation* 38: 342 - 345.
- Cox, J. R., M. H. Martin-R., F. A. Ibarra-F. and H. L. Morton. 1987.** Establishment of range grasses on various seedbeds at Creosotebush (*Larrea tridentata*) sites in Arizona, USA and Chihuahua, Mexico. *J. Range Management* 39: 540 - 546.
- Cox, J. R. and R. M. Madrigal. 1988.** Establishing perennial grasses on abandoned farmland in southeastern Arizona. *Applied Agricultural Research* 3: 36 - 43.
- Cox, J. R., R. M. Madrigal and G. W. Frasier. 1989.** Survival of perennial grass transplants in the Sonoran Desert of the southwestern USA. *Arid Soil Research and Rehabilitation* 1: 77 - 87.
- Cox, J. R., A. Alba-Avila, R. W. Rice and J. N. Cox. 1993.** Biological and physical factors influencing *Acacia constricta* and *Prosopis velutina* establishment in the Sonoran Desert. *J. Range Management* 46: 43 - 48.
- FAO-UNESCO. 1975.** Food and Agricultural Organization of the United Nations. United Nations Educational, Scientific and Cultural Organization. Volume III. Soil Map and Precipitation trends for Mexico and Central America.
- Ferguson, R. B., R. A. Ryker and E. D. Ballard. 1975.** Portable oscilloscope techniques for detecting dormancy in nursery stock. USDA-Forest Service General Technical Report INT-26. Intermountain Forest and Range Experiment Station. Ogden, UT.
- Frasier, G. W., D. A. Woolhiser and J. R. Cox. 1984.** Emergence and seedling

survival of two warm-season grasses as influenced by the timing of precipitation: A greenhouse study. *J. Range Management* 37: 7 - 11.

Frasier, G. W., J. R. Cox and D. A. Wollhiser. 1985. Emergence and survival responses of seven grasses for six wet-dry sequences. *J. Range Management* 38: 372 - 377.

Griffith, D. 1901. Range improvements in Arizona. USDA-Bureau of Plant Industry, Bulletin #4. 12p.

Hassanyar, A. S. and A. M. Wilson. 1978. Drought tolerance of seminal lateral root apices in crested wheatgrass and Russian wildrye. *J. Range Management* 31: 254 - 258.

Hastings, J. R. and R. M. Turner. 1965. The changing mile. University of Arizona Press, Tucson, AZ.

Herbel, C. H., G. H. Abernathy, C. Y. Yarbrough and D. K. Gardner. 1973. Rootplowing and seeding arid rangelands in the southwest. *J. Range Management* 26: 193 - 197.

Hinton, R. J. 1890. Irrigation in the United States. Volume I, Report of the Special Committee of the United States Senate on Irrigation and Reclamation of Arid Lands. Senate Report # 928, 51st Congress, 1st Session. US Govt. Printing Office, Washington, DC.

Houser, V. L. 1983. Grass establishment by bandoleers, transplants, and germinated seeds. *Transactions American Society Agricultural Engineers* 26: 74 - 80.

Howard, G. S., F. Rauzi and G. E. Schuman. 1978. Woody plant trials at six mine reclamation sites in Wyoming and Colorado. USDA-Production Research Report #177. US Govt. Printing Office, Washington, DC.

Humphrey, R. R. 1958. The desert grasslands. *Botanical Review* 24: 193 - 252.

Knipe, O. D. 1968. Effects of moisture stress on germination of alkali sacaton, galleta, and blue grama. *J. Range Management* 21: 3 - 4.

Knipe, O. D. 1973. Western wheatgrass germination as related to temperature, light, and moisture stress. *J. Range Management* 26: 68 - 69.

Kruse, W. H. 1970. Temperature and moisture stress affect germination of *Gutierrezia Sarothrae* (broom snakeweed). *J. Range Management* 23: 143 -144.

Mexico Climatic Reporting Stations. 1992. Climatography of Mexico. Govt. Printing Office, Mexico City, MX.

- Piatt, J. R. 1976.** Effects of water stress and temperature on germination of true mountain mahogany. *J. Range Management* 29: 138 - 140.
- Scifres, C. J. and J. H. Brock. 1969.** Moisture-temperature interrelations in germination and early seedling development of mesquite. *J. Range Management* 22: 334 - 337.
- Springfield, H. W. 1972.** Mulching improves survival and growth of *Cercocarpus* transplants. USDA-Forest Service Research Note RM-200. Rocky Mountain Forest and Range Experiment Station. Fort Collins, CO.
- Thacker, G. W. and J. R. Cox. 1992.** How to establish a permanent vegetation cover on farmland. USDA-Cooperative Extension, College of Agriculture, University of Arizona, Bulletin 191051. Tucson, AZ.
- Tinus, R. W. 1978.** Production of container-grown hardwoods. *Tree Planter's Notes* 29: 3 - 9.
- Wilson, A. M. and D. D. Briske. 1979a.** Seminal and adventitious root growth of blue grama seedlings on the central plains. *J. Range Management* 32: 209 - 213.
- Wilson, A. M. and D. D. Briske. 1979b.** Drought and temperature effects on the establishment of blue grama seedlings. In: *Proceeding 1st International Rangeland Congress*. Society for Range Management. Denver, CO.
- Wagoner, J. J. 1952.** History of the cattle industry in southern Arizona, 1540 - 1940. *University of Arizona Social Sciences Bulletin # 20*, Tucson, AZ.
- York, J. C. and W. A. Dick-Peddie. 1969.** Vegetation changes in southern New Mexico during the past 100 years. In: *Aridlands in perspective*. University of Arizona Press, Tucson, AZ.