

# USAID Country Health Statistical Report

# **Brazil**

## May 2007



### **Table of Contents**

### 1. About the Report

### 2. Country Profile

### I. Statistical Overview

Demographic Indicators

Socioeconomic Indicators

Family Planning Indicators

Maternal Health Indicators

Child Survival Indicators

HIV/AIDS Prevention Indicators

Infectious Diseases Control Indicators

### II. Trends in Health Indicators

Current and Projected Population by Age and Sex (Figure 1 and Table 1.1)
Population Estimates and Projections (Table 1.2)
Estimated Total Fertility Rate per Woman (Figure 2 and Table 2)
Infant Mortality Rates / Under-5 Mortality Rates (Figure 3 and Table 3)
Vaccination Coverage (DPT3) Trends

(Figure 4 and Table 4)
Contraceptive Prevalence Rates - Married Wome

Contraceptive Prevalence Rates - Married Women (Figure 5 and Table 5)

### 3. Data Notes

### 4. Sources

### **About the Report**

### Released May 2007

This report is one of a series of Country Health Statistical Reports produced on behalf of the United States Agency for International Development (USAID) by the Analysis, Information Management and Communication Activity (AIM) Project. Each profile contains statistical data on current health conditions, population dynamics, health and family planning behavior, and health and population trends in a given developing country. Information is compiled from AIM's health statistics database, which draws data from a diverse range of sources listed at the end of this profile.

Hard copy editions of USAID's profiles are available from AIM'S publications department. Reports are available on the USAID Global Health Web site in Portable Document Format (pdf). Any feedback you have on the content or presentation of this report would be greatly appreciated. We would also appreciate receiving any more recent, more accurate, or more representative information. Contact us at the following:

**Analysis, Information Management** 

ጼ

**Communications Activity (AIM)** 

655 15th Street, NW, Suite 450 Washington, DC 20005

> Phone: 202-661-8600 Fax: 202-661-8601

E-mail: dolphn@aimglobalhealth.org

The Analysis, Information Management & Communications (AIM) Activity, a USAID contract managed by MasiMax Resources, Inc, with John Snow, Inc., ORC Macro, and Insight Systems Corporation prepared this document under task order number GPO-M-00-05-00043-00, with the Bureau for Global Health, USAID.

# **I. Statistical Overview**

\*see data notes

Demographic Indicators				
Indicator	Value	Data Unit	Year	Source
Total Population	188,078,227		2006	BUCEN-IDB-2006
Population Growth Rate	1	%	2006	BUCEN-IDB-2006
Percent Urban	84	%	2004	World Bank/WDI-2006
Women, 15-19	8,412,520		2006	BUCEN-IDB-2006
Women, 15-49	53,053,877		2006	BUCEN-IDB-2006
Life Expectancy at Birth	72		2006	BUCEN-IDB-2006
Crude Birth Rate	16.6	per 1,000	2006	BUCEN-IDB-2006
Crude Death Rate	6.2	per 1,000	2006	BUCEN-IDB-2006
Number of Live Births	3,114,575		2006	BUCEN-IDB-2006
Healthy Life Expectancy: Female	62.4		2002	WHO World Health Report-2004
Healthy Life Expectancy: Male	57.2		2002	WHO World Health Report-2004
Population: Percent < Age 15	25.3	%	2007	BUCEN-IDB-2007

Socioeconomic Indicators				
Indicator	Value	Data Unit	Year	Source
GNI per Capita (PPP)	8,230	\$	2005	World Bank/WDI-2006
Health Expenditure as Percentage of GDP	7.6	%	2003	World Bank/WDI-2006
Physicians per 1,000 People	2.1	per 1,000	1997-2004	World Bank/WDI-2006
Adult Literacy Rate	88.6	%	2004	World Bank/WDI Database-2007
Adult Literacy Rate, Female	89 1	%	2002	World Bank/WDI-2006
Adult Literacy Rate, Male	88 2	%	2002	World Bank/WDI-2006
Gross Enrollment Ratio - Primary School	141	%	2003	UNESCO-2007
Gender Parity Index - Net Enrollment Ratio	1.2		2001	UNESCO-2005
Access to an Improved Water Source (Rural)	58	%	2002	World Bank/WDI-2005
Access to an Improved Water Source (Urban)	96	%	2002	World Bank/WDI-2005
Access to Improved Sanitation Facilities (Rural)	35	%	2002	World Bank/WDI-2005
Access to Improved Sanitation Facilities (Urban)	83	%	2002	World Bank/WDI-2005
Population Living Below \$1 a Day	7.5	%	2006	Human Development Report-2006
Human Development Index	0.79		2006	Human Development Report-2006
Population Below Poverty Line	22	%	1998	CIA World Factbook-2006
Real GDP (Growth) Rate	3.1	%	2006	CIA World Factbook-2007
Income Inequality (Gini index)	56.7		2005	CIA World Factbook-2007
Unemployment Rate	9.6	%	2006	CIA World Factbook-2007
Inflation Rate	4.2	%	2006	CIA World Factbook-2007
Airports	4,276		2006	CIA World Factbook-2007
Airports with paved runways	714		2006	CIA World Factbook-2007

<sup>&</sup>lt;sup>1</sup> Based on survey data.<sup>2</sup> Based on survey data

Family Planning Indicators							
Indicator	Value	Data Unit	Year	Source			
Total Fertility Rate (BUCEN)	1.9		2006	BUCEN-IDB-2006			
Total Fertility Rate (DHS)	2.5		1996	Brazil DHS-1996			
Contraceptive Prevalence Rate, Modern Methods, All Women	51	%	1996	Brazil DHS-1996			
Contraceptive Prevalence Rate, Modern Methods, Married Women	Data Not Available						
Median Age of Sexual Debut Among Women, Ages 25-49	19.5		1996	Brazil DHS-1996			
Mean Ideal Family Size	2.3		1996	Brazil DHS-1996			
Women 20-24 Who Gave Birth Before Age 20	32	%	1996	Brazil DHS-1996			
Total Fertility Rate, Rural	3.5		1996	DHS STATcompiler as of January 2007-2007			
Unmet Need for Family Planning: Rural	12.5	%	1996	DHS STATcompiler as of January 2007-2007			
Unmet Need for Family Planning: Urban	6	%	1996	DHS STATcompiler as of January 2007-2007			
Contraceptive Prevalence Rate, All Methods	76.7	%	1996	DHS STATcompiler as of March 2007-2007			
Equity in use of modern contraception	Data Not Available						
Percent of need satisfied by modern methods of family planning	Data Not Available						

Maternal Health Indicators						
Indicator	Value	Data Unit	Year	Source		
Maternal Mortality Ratio (WHO/Hill)	260	Per 100,000 live births	2000	WHO/Hill-2004		
Maternal Mortality Ratio (DHS)		Data Not Ava	ailable			
Antenatal Care (at least 1 visit)	85.7	%	1996	Brazil DHS-1996		
Antenatal Care (2+ visits)	84.5	%	1996	Brazil DHS-1996		
Antenatal Care (4+ visits)	76.9	%	1996	Brazil DHS-1996		
Assisted Delivery by a Health Professional	87.6	%	1996	Brazil DHS-1996		
Assisted Delivery by Doctor	77.6	%	1996	Brazil DHS-1996		
Assisted Delivery by Other Health Professional	10	%	1996	Brazil DHS-1996		
Anemia prevalence among women of reproductive age	Data Not Available					
Equity in skilled attendance at delivery	quity in skilled attendance at delivery  Data Not Available					

Child Survival Indicators									
Indicator	Value	Data Unit	Year	Source					
Acute Respiratory Infection (ARI) and Oral Rehydration Therapy (ORT)									
ARI Care Seeking - Children Under 5	46.1	%	1996	Brazil DHS-1996					
ORT Use Rate (ORS, RHS, or Increased Fluids)	73.4	%	1996	Brazil DHS-1996					
Mortality Indicators									
Infant Mortality Rate (BUCEN)	28.6	per 1,000 live births	2006	BUCEN-IDB-2006					
Infant Mortality Rate (DHS)	39.3	per 1,000 live births	1996	Brazil DHS-1996					
Infant Mortality Rate, Females (BUCEN)	24.7	per 1,000 live births	2006	BUCEN-IDB-2006					
Infant Mortality Rate, Males (BUCEN)	32.3	per 1,000 live births	2006	BUCEN-IDB-2006					
Under-5 Mortality Rate (BUCEN)	32.6	per 1,000 live births	2006	BUCEN-IDB-2006					
Neonatal Mortality Rate	19	per 1000	1996	DHS STATcompiler as of January 2007- 2007					
Under-5 Mortality Rate (UNICEF)	33	per 1,000 live births	2005	UNICEF-2007					
Under-5 Mortality Rate (DHS)	48.8	per 1,000 live births	1996	Brazil DHS-1996					
Nutrition Indicators									
Exclusive Breastfeeding (under 4 mos.)	40.3	%	1996	Brazil DHS-1996					
Exclusive Breastfeeding (under 6 mos.)			Data Not	Available					
Stunted (height-for-age)	10.5	%	1996	Brazil DHS-1996					
Underweight (weight-for-age)	5.7	%	1996	Brazil DHS-1996					
Wasted (weight-for-height)	2.3	%	1996	Brazil DHS-1996					
Vaccination Coverage									
DPT3 Vaccination Rate (DHS)	80.8	%	1996	Brazil DHS-1996					
DPT3 Vaccination Rate (WHO)	96	%	2004	WHO/Global Summary-2005					
Measles Vaccination Rate (DHS)	87.2	%	1996	Brazil DHS-1996					
Measles Vaccination Rate (WHO)	99	%	2004	WHO/Global Summary-2005					
Polio Vaccination Rate (DHS)	80.7	%	1996	Brazil DHS-1996					
Polio Vaccination Rate (WHO)	98	%	2004	WHO/Global Summary-2005					
Tetanus Toxoid Vaccination	45.1	%	1996	Brazil DHS-1996					
Equity in DPT3 coverage (DHS)			Data Not	Available					
Improved sanitation/hygiene practices			Data Not	Available					
Vitamin A supplementation coverage among children 6-59 months			Data Not	Available					

HIV/AIDS Prevention Indicators											
Indicator	Value	Data Unit	Year	Source							
HIV Prevalence proportion: Adults (15–49 years)	0.5	%	2005	UNAIDS 2006-2006							
Estimated number of people living with HIV: Adults and Children	620,000		2005	UNAIDS 2006-2006							
Estimated number of people living with HIV: Women (15+ years)	220,000		2005	UNAIDS 2006-2006							
Estimated number of people living with HIV: Children (0–14 years)	Data Not Available										
Males Reporting Condom Use With Last Non-Regular Partner  Data Not Available											
Females Reporting Condom Use With Last Non-Regular Partner  Data Not Available											

Infectious Diseases Control Indicators							
Indicator	Value	Data Unit	Year	Source			
TB Estimated Number of Cases	86,881		2004	WHO/TB Control Report-2006			
TB Case Detection Rate	46.7	%	2004	WHO/TB Control Report-2006			
TB (DOTS) Treatment Success Rate	83.2	%	2003	WHO/TB Control Report-2006			
Malaria Reported Number of Cases	379,551	Reported # of cases	2003	WHO/RBM-2005			
DOTS Coverage	52	%	2004	WHO/TB Control Report-2006			
HIV-Infected with Active TB	16.9	%	2006	WHO/TB Control Report-2006			
Avian Influenza: Cumulative Number of Confirmed Human Cases		Dat	ta Not Ava	ailable			
Avian Influenza: Cumulative Number of Confirmed Human Deaths	Data Not Available						
Child Sleeps Under Insecticide-Treated Net	Data Not Available						
Equity in Bednet (ITN) Ownership		Dat	ta Not Ava	ailable			

### **II. Trends in Health Indicators**




Figure 1: Current and Projected Population by Age and Sex (in thousands)

Table 1.1: Current and Projected Population by Age and Sex


Source: BUCEN-IDB Age Group Male - 2000 Male - 2020 Female - 2000 Female - 2020 0-4 8,464,596 7,217,289 8,131,962 6,917,575 5-9 8,435,011 7,399,141 8,115,714 7,104,105 10-14 8,896,482 7,607,692 8,580,957 7,315,437 15-19 8,956,122 7,875,877 8,690,058 7,596,549 20-24 8,588,098 8,308,868 8,414,374 8,063,252 25-29 7,896,446 8,234,181 7,804,041 8,053,269 30-34 7,293,533 8,591,324 7,274,230 8,490,646 35-39 6,440,717 8,550,804 6,537,662 8,564,757 5,402,122 8,107,037 8,245,404 40-44 5,576,433 45-49 4,452,219 7,333,091 4,688,854 7,573,692 3,524,425 6,595,813 3,813,385 6,952,034 50-54 55-59 2,689,316 5,587,806 3,022,689 6,107,148 60-64 2,106,410 4,397,654 2,490,390 5,034,272 65-69 1,566,991 3,301,863 1,977,744 4,017,825 70-74 2,270,967 1,086,785 1,524,820 3,003,423 75-79 660,617 2,070,264 1,395,893 1,037,736 472,243 1,237,428 939,589 2,385,337 80+

**TABLE 1.2: Population Estimates and Projections** 

Source: BUCEN-IDB

1960	1970	1980	1990	2000	2010	2020	2030	2040	2050
71,694,810	95,684,297	122,958,132	151,083,809	175,552,771	195,579,661	211,507,717	222,838,366	228,453,853	228,426,737

Figure 2: Estimated Total Fertility Rate per Woman



**Table 2: Estimated Total Fertility Rate per Woman** 

Source: BUCEN-IDB

1950	1960	1970	1980	1990	2000	2010	2020	2030	2040	2050
5.9	6.1	5.3	4.1	2.6	2.1	1.8	1.7	1.7	1.7	1.7

Figure 3: Infant Mortality Rates / Under-5 Mortality Rates




Table 3: Infant Mortality Rates / Under-5 Mortality Rates

Source: BUCEN-IDB 2006

Indicator	1960	1970	1980	1990	2000	2010	2020	2030	2040	2050
Infant Mortality Rate (BUCEN)	116.0	97.9	73.7	49.9	35.2	24.9	17.7	12.9	9.7	7.5
Under-5 Mortality Rate (BUCEN)		128.0	89.9	57.4	40.2	28.3	20.0	14.5	10.9	8.5

Figure 4: Vaccination Coverage (DPT3) Trends




Table 4: Vaccination Coverage (DPT3) Trends

Source: WHO/Global Summary

1985	1990	1995	2000
66.0	66.0	84.0	95.0

Figure 5: Contraceptive Prevalence Rates, Modern Methods, Married Women
Data Not Available

Table 5: Contraceptive Prevalence Rates, Modern Methods, Married Women
Data Not Available

### **Data Notes**

\*Indicator definitions that differ from those in the data notes section are identified by footnotes accompanying the indicator tables.

Demod		hi.	India	-4
Dellio	ırav	nic i	maic	ลเบาร

Total Population The number of people in a given area (i.e., country) in a particular time period (usually a midyear

estimate).

Population Growth Rate The average annual growth rate is the rate of natural increase in a population plus the net migration

rate. The rate of natural increase is the difference between the birth rate and the death rate, but it is

conventionally measured in percentage terms (per hundred rather than per thousand).

Percent Urban The percentage of the midyear population living in areas defined as urban in each country and reported

to the United Nations.

Women, 15-19 The number of women between the ages of 15-19 in the midyear population.

Women, 15-49 The number of women between the ages of 15-49 in the midyear population.

Life Expectancy at Birth The average number of years that a person at age 0 will live if age-specific death rates remain constant.

Life expectancy at birth is highly affected by rates of infant and child death.

Crude Birth Rate The number of births per thousand of the population. The product of the number of live births, divided by

the midpoint population, multiplied by 1,000.

Crude Death Rate The number of deaths per thousand of the population. The product of the number of deaths divided by

the midpoint population, multiplied by 1,000.

Number of Live Births The number of live births, annually, within a country.

Healthy Life Expectancy:

Female

The number of years of in full health that a female newborn can expect to live based on current rates of ill-health and mortality. HALE is based on life expectancy at birth but includes an adjustment for time spent in poor health.

**Healthy Life Expectancy:** 

Male

The number of years of in full health that a male newborn can expect to live based on current rates of ill-health and mortality. HALE is based on life expectancy at birth but includes an adjustment for time spent

in poor health.

Population: Percent < Age 15

Percent of population less than 15 years of age.

#### Socioeconomic Indicators

**GNI per Capita (PPP)** 

The gross national income (GNI) converted to international dollars using purchasing power parity (PPP) rates. An international dollar has the same purchasing power over GNI as a U.S. dollar has in the United States.

Health Expenditure as Percentage of GDP

Total health expenditures is the sum of public and private health expenditures figured as a percentage of a country's gross domestic product (GDP). It covers the provision of health services (preventive and curative), family planning activities, nutrition activities, and emergency aid designated for health but does not include provision of water and sanitation.

Physicians per 1,000 People

Physicians are defined as graduates of any faculty or school of medicine who are working in the country in any medical field (practice, teaching, research).

**Adult Literacy Rate** 

The percentage of people ages 15 and over who can, with understanding, read and write a short, simple statement about their everyday life.

Adult Literacy Rate, Female

The percentage of women ages 15 and over who can, with understanding, read and write a short, simple statement about their everyday life.

Adult Literacy Rate, Male

The percentage of men ages 15 and over who can, with understanding, read and write a short, simple statement about their everyday life.

Gross Enrollment Ratio - Primary School

Total enrolment in primary level of education, regardless of age, expressed as a percentage of the population in the official age group corresponding to primary level of education. The GER can exceed 100% due to late entry or/and repetition.

Gender Parity Index - Net Enrollment Ratio

The ratio of the female-to-male values (or male to female, in certain cases) of net primary school enrollment rates (NER). NER measures the number of pupils in the official age group for a given level of ecudation, expressed as a percentage of the population in that age group.

Access to an Improved Water Source (Rural)

Access to an improved water source in rural areas. Refers to the percentage of the rural population with reasonable access to an adequate amount of water from an improved source, such as a household connection, public standpipe, borehole, protected well or spring, or rainwater collection. Unimproved sources include vendors, tanker trucks, and unprotected wells and springs.

Access to an Improved Water Source (Urban)

Access to an improved water source in urban areas. Refers to the percentage of the urban population with reasonable access to an adequate amount of water from an improved source, such as a household connection, public standpipe, borehole, protected well or spring, or rainwater collection. Unimproved sources include vendors, tanker trucks, and unprotected wells and springs.

Access to Improved Sanitation Facilities (Rural)

Access to improved sanitation facilities in rural areas. Refers to the percentage of the rural population with access to at least adequate excreta disposal facilities (private or shared, but not public) that can effectively prevent human, animal, and insect contact with excreta.

Access to Improved Sanitation Facilities (Urban)

Access to improved sanitation facilities in urban areas. Refers to the percentage of the urban population with access to at least adequate excreta disposal facilities (private or shared, but not public) that can effectively prevent human, animal, and insect contact with excreta.

Population Living Below \$1 a Day

The percentage of the population living below the specified poverty line of \$1 a day at 1985 international prices (equivalent to \$1.08 at 1993 international prices), adjusted for purchasing power parity.

**Human Development Index**  The HDI is a composite index that measures the average achievements in a country in three basic dimensions of human development: a long and healthy life, as measured by life expectancy at birth; knowledge, as measured by the adult literacy rate and the combined gross enrollment ratio for primary, secondary, and tertiary schools; and a decent standard of living, as measured by gross domestic product (GDP) per capita in purchasing power parity (PPP) US dollars.

Population Below Poverty Line

National estimates of the percentage of the population falling below the poverty line are based on surveys of sub-groups, with the results weighted by the number of people in each group. Definitions of poverty vary considerably among nations. For example, rich nations generally employ more generous standards of poverty than poor nations.

Real GDP (Growth) Rate

This entry gives the gross domestic product (GDP) growth on an annual basis adjusted for inflation and expressed as a percent.

Income Inequality (Gini index)

This index measures the degree of inequality in the distribution of family income in a country. It is calculated from the Lorenz curve, in which cumulative family income is plotted against the number of families arranged from the poorest to the richest. The index is the ratio of (a) the area between a country's Lorenz curve and the 45 degree helping line to (b) the entire triangular area under the 45 degree line. The more nearly equal a country's income distribution, the closer its Lorenz curve to the 45 degree line and the lower its Gini index, e.g., a Scandinavian country with an index of 25. The more unequal a country's income distribution, the higher its Gini index, e.g., a Sub-Saharan country with an index of 50. If income were distributed with perfect equality, the index would be zero; if income were distributed with perfect inequality, the index would be 100.

**Unemployment Rate** 

This entry contains the percent of the labor force that is without jobs. Substantial underemployment might be noted.

**Inflation Rate** 

This entry furnishes the annual percent change in consumer prices compared with the previous year's consumer prices.

#### **Airports**

This entry gives the total number of airports or airfields recognizable from the air. The runway(s) may be paved (concrete or asphalt surfaces) or unpaved (grass, earth, sand, or gravel surfaces) but may include closed or abandoned installations. Airports or airfields that are no longer recognizable (overgrown, no facilities, etc.) are not included. Note that not all airports have accommodations for refueling, maintenance, or air traffic control.

# Airports with paved runways

This entry gives the total number of airports with paved runways (concrete or asphalt surfaces) by length. For airports with more than one runway, only the longest runway is included according to the following five groups - (1) over 3,047 m, (2) 2,438 to 3,047 m, (3) 1,524 to 2,437 m, (4) 914 to 1,523 m, and (5) under 914 m. Only airports with usable runways are included in this listing. Not all airports have facilities for refueling, maintenance, or air traffic control.

### Family Planning Indicators

Total Fertility Rate (BUCEN)

The number of children a woman between ages 15-49 would have during her reproductive life, if, for all of her childbearing years she were to experience the age-specific birth rates for that given year.

**Total Fertility Rate (DHS)** 

The number of children a woman between ages 15-49 would have during her lifetime if she were to bear children at the currently observed rates.

Contraceptive Prevalence Rate, Modern Methods, All Women Percentage of all women ages 15-49 currently using a modern method of contraception. Modern methods include oral contraceptives, IUDs, injectables, female and male sterilization, all emergency contraception, and barrier methods (diaphragm, foam, jelly, male and female condom).

Contraceptive Prevalence Rate, Modern Methods, Married Women Percent of currently married women ages 15-49 currently using a modern method of contraception. Modern methods include oral contraceptives, IUDs, injectables, female and male sterilization, all emergency contraception, and barrier methods (diaphragm, foam, jelly, male and female condom).

Median Age of Sexual Debut Among Women, Ages 25-49 Median age of first sexual intercourse for women ages 25-49.

Mean Ideal Family Size

Mean ideal number of children for all women, according to number of living children.

Women 20-24 Who Gave Birth Before Age 20

Percentage of women ages 20-24 who have given birth before age 20.

**Total Fertility Rate, Rural** 

In rural areas only, the number of children a woman between ages 15-49 would have during her lifetime if she were to bear children at the currently observed rates.

Unmet Need for Family Planning: Rural

In rural areas, percentage of married women ages 15-49 with unmet need for family planning. Unmet need for spacing includes pregnant women whose pregnancy was mistimed, amenorrheic women who are not using family planning and whose last birth was mistimed, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and say they want to wait 2 or more years for their next birth. Also included are fecund women who are not using any method of family planning and say they are unsure whether they want another child or who want another child but are unsure when to have the birth unless they say it would not be a problem if they discovered they were pregnant in the next few weeks. Unmet need for limiting refers to pregnant women whose pregnancy was unwanted, amenorrheic women whose last child was unwanted, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and want no more children.

**Unmet Need for Family Planning: Urban** 

In urban areas, percentage of married women ages 15-49 with unmet need for family planning. Unmet need for spacing includes pregnant women whose pregnancy was mistimed, amenorrheic women who are not using family planning and whose last birth was mistimed, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and say they want to wait 2 or more years for their next birth. Also included are fecund women who are not using any method of family planning and say they are unsure whether they want another child or who want another child but are unsure when to have the birth unless they say it would not be a problem if they discovered they were pregnant in the next few weeks. Unmet need for limiting refers to pregnant women whose pregnancy was unwanted, amenorrheic women whose last child was unwanted, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and want no more children.

Contraceptive Prevalence Rate, All Methods Percentage of currently married women ages 15-49 currently using ANY method of contraception.

Equity in use of modern contraception

Contraceptive prevalence rate in the lowest wealth quintile/contraceptive prevalence rate in the highest wealth quintile.

Percent of need satisfied by modern methods of family planning Number of women in union who are fecund and are currently using modern contraceptive methods to stop or postpone the next pregnancy/number of women in union who are fecund and who desire to either terminate childbearing or to postpone their next birth for 2 years or more.

#### Maternal Health Indicators

Maternal Mortality Ratio (WHO/Hill)

The estimated number of women who die as a result of pregnancy or childbirth per 100,000 live births arrived through Hill et al., adjustment procedure depending on the nature of data used. The margins of uncertainty associated with the estimated maternal mortality ratios are very large and the estimates should not be used to monitor trends in the short term (including comparisons between 1995 and 2000 estimates). In addition, cross-country comparisons should be treated with considerable circumspection because different strategies are used to derive the estimates for different countries rendering comparisons fraught with difficulty. For further information please refer to the source.

Maternal Mortality Ratio (DHS)

The estimated number of women who die as a result of pregnancy or childbirth per 100,000 live births, arrived mostly through the "sisterhood method". The data are aggregated based on different time periods ranging from four to ten years preceding the survey. Thus, the data may not be suitable for trend analysis due to over-lapping year ranges of estimates. For further information on methodology please refer to: Stanton, Cynthia, Noureddine Abderrahim, and Kenneth Hill. 1997. DHS Maternal Mortality Indicators: An assessment of Data Quality and Implications for Data Use. Calverton: Macro International Inc., or individual country DHS reports.

Antenatal Care (at least 1 visit)

Percentage of women of reproductive age (15-49) who receive at least one antenatal care visit during pregnancy (in the three-year period preceding the survey).

Antenatal Care (2+ visits)

Percentage of women of reproductive age (15-49) who receive at least two antenatal care visits during pregnancy (in the three-year period preceding the survey).

**Antenatal Care (4+ visits)** 

Percentage of women of reproductive age (15-49) who receive four or more antenatal care visits during pregnancy (in the three-year period preceding the survey).

Assisted Delivery by a Health Professional

The percentage of births/deliveries that occur with the assistance of any trained health professional during the five-year period preceding the survey. May include doctors, nurses, or midwives.

Assisted Delivery by Doctor

The percentage of live births/deliveries that occur with the assistance of a doctor during the five-year period preceding the survey.

Assisted Delivery by Other Health Professional The percentage of live births/deliveries that occur with the assistance of a nurse or midwife during the five-year period preceding the survey.

Anemia prevalence among women of reproductive age

Number of women of reproductive age (15-49 years) with anemia/number of women of reproductive age (15-49 years).

Equity in skilled attendance at delivery

Percentage of births in the lowest wealth quintile attended by a doctor, nurse, or trained midwife/ percentage of births in the highest wealth quintile attended by a doctor, nurse, or trained midwife.

#### **Child Survival Indicators**

ARI Care Seeking - Children Under 5

Percentage of children under five years who were ill with an acute respiratory infection (ARI), which is associated with cough, rapid breathing and a high fever, during the two weeks preceding the survey, for whom care was sought from a health facility or provider.

ORT Use Rate (ORS, RHS, or Increased Fluids)

Oral Rehydration Therapy (ORT). Percentage of children under age 5 with diarrhea two weeks prior to survey who received increased fluids, oral rehydration solution (ORS) or recommended home solution (RHS).

Infant Mortality Rate (BUCEN)

The estimated annual number of deaths of infants under 12 months in a given year per 1,000 live births in that same year.

Infant Mortality Rate The estimated annual number of deaths of infants under 12 months in a given year per 1,000 live births (DHS) in that same year (five-year period preceding survey). **Infant Mortality Rate,** The estimated annual number of deaths of female infants under 12 months in a given year per 1,000 Females (BUCEN) live births in that same year. **Infant Mortality Rate,** The estimated annual number of male infants under 12 months in a given year per 1,000 live births in Males (BUCEN) that same year. **Under-5 Mortality Rate** Annual number of deaths that occur in children 0-4 years old per 1,000 births. (BUCEN) **Neonatal Mortality Rate** The estimated number of infant deaths in the first month of life per 1,000 live births in the five-year period preceding the survey. **Under-5 Mortality Rate** Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period (UNICEF) preceding survey). **Under-5 Mortality Rate** Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period (DHS) preceding survey). **Exclusive Breastfeeding** Percentage of children under 4 months exclusively breastfed. Exclusive breastfeeding is defined as (under 4 mos.) providing no food or liquid other than breast milk to the child during the 24-hour period before the survev. **Exclusive Breastfeeding** Percentage of children under 6 months exclusively breastfed. Exclusive breastfeeding is defined as (under 6 mos.) providing no food or liquid other than breast milk to the child during the 24-hour period before the survey. Stunted (height-for-age) Percentage of children under age five whose height-for-age is below minus-two standard deviations from the median of the reference population. Underweight (weight-for-Percentage of children under age five whose weight-for-age is below minus-two standard deviations age) from the median of the reference population. Wasted (weight-for-Percentage of children under age five whose weight-for-height is below minus-two standard deviations height) from the median of the reference population. **DPT3 Vaccination Rate** Proportion of living children 12-23 months old who have received three complete doses of vaccines (DHS) against diphtheria, pertussis, and tetanus. **DPT3 Vaccination Rate** Proportion of living children 12-23 months old who have received three complete doses of vaccines (WHO) aginst diphtheria, pertussis, and tetanus. **Measles Vaccination** Percentage of living children 12-23 months old who have received one dose of Measles Containing Rate (DHS) Vaccince **Measles Vaccination** Percentage of living children 12-23 months old who have received one dose of Measles Containing Rate (WHO) Vaccince. **Polio Vaccination Rate** Proportion of living children 12-23 months who have received three doses of polio vaccine. (DHS) **Polio Vaccination Rate** Proportion of living children 12-23 months who have received three doses of polio vaccine. (WHO) **Tetanus Toxoid** Percentage of pregnant women receiving two (or more) doses of tetanus toxoid (three-year period Vaccination preceding survey). **Equity in DPT3 coverage** DPT3 coverage in the lowest wealth quintile/DPT3 coverage in the highest wealth quintile. (DHS) Improved sanitation/ Number of households with child under 5 years of age whose youngest child's feces was safely hygiene practices disposed of the last time he/she passed stool/total number of households with a child under 5 years of age. Vitamin A Number of children aged 6-59 months who received a vitamin A supplement in the last 6 months/total supplementation number of children aged 6-59 months surveyed. coverage among children

6-59 months

HIV Prevalence proportion: Adults (15–49 years)

The adult HIV prevalence proportion is the estimated number of adults living with HIV divided by the adult population (aged 15–49).

Estimated number of people living with HIV: Adults and Children

Estimated number of adults and children living with HIV. Adults are 15 years and over. Children are defined as those aged 0–14 years.

Estimated number of people living with HIV: Women (15+ years)

Estimated number of women (aged 15 and over) living with HIV.

Estimated number of people living with HIV: Children (0–14 years)

Estimated number of children under age 15 living with HIV.

Males Reporting Condom Use With Last Non-Regular Partner Among men who have had high risk sex in the last year, percentage who used condoms during their last sexual intercourse with a non-regular partner/non-cohabitating partner.

Females Reporting Condom Use With Last Non-Regular Partner Among women who have had high risk sex in the last year, percentage who used condoms during their last sexual intercourse with a non-regular partner/non-cohabitating partner.

#### Infectious Diseases Control Indicators

TB Estimated Number of Cases

The estimated number of cases (all forms).

**TB Case Detection Rate** 

TB Case Detection Rate is defined as the percentage of the annual new smear-positive notifications of the estimated annual new smear-positive incidence.

TB (DOTS) Treatment Success Rate

The proportion of smear-positive patients who were cured plus the proportion who completed treatment.

Malaria Reported Number of Cases

Reported number of malaria cases during the year for which data was reported.

**DOTS Coverage** 

The percentage of the national population living in areas where health services have adopted DOTS. "Areas" are the lowest administrative or management units in the country (townships, districts, counties, etc). If an area is considered a DOTS area, then all cases registered and reported in that area are considered DOTS cases and the population living within the boundaries of that area counts toward national DOTS coverage. As a measure of patient access to diagnosis and treatment under DOTS, coverage is an approximation, and usually an overestimate.

HIV-Infected with Active TB

The TB incidence rate in HIV-infected people 15-49 divided by the incidence rate in HIV-uninfected people 15-49.

Avian Influenza: Cumulative Number of Confirmed Human Cases The cumulative number of confirmed human cases of Avian Influenza A/(H5N1) reported to WHO. The total number of cases includes number of deaths. WHO reports only laboratory-confirmed cases.

Avian Influenza: Cumulative Number of Confirmed Human Deaths The cumulative number of confirmed human deaths from Avian Influenza A/(H5N1) reported to WHO. WHO reports only laboratory-confirmed cases and deaths.

Child Sleeps Under Insecticide-Treated Net Percentage of children under five years of age who slept under an Insecticde Treated Net (ITN) the night before the survey. An ITN is (1) a permanent net that does not require any treatment, or (2) a pretreated net obtained within the last six months, or (3) a net that has been soaked with insecticide within the past six months.

Equity in Bednet (ITN)
Ownership

Proportion of households in the lowest income quintile with an ITN / proportion of households in the highest income quintile with an ITN.

## **Statistical Sources For Brazil**

WHO/Hill 2004

Brazil DHS 1996	Sociedade Civil Bem-Estar Familiar no Brasil and Macro International Inc., Brasil Pesquisa Nacional Sobre Demografia e Saude 1996, Calverton, Maryland, USA: March 1997.
BUCEN-IDB 2005	United States Census Bureau (BUCEN), International Programs Center. International Database. May 2006.
BUCEN-IDB 2006	United States Census Bureau (BUCEN), International Programs Center. International Database. August 2006.
BUCEN-IDB 2007	United States Census Bureau (BUCEN), International Programs Center. International Database. March 2007.
CIA World Factbook 2006	CIA World Factbook. 2006. Available at: https://cia.gov/cia//publications/factbook/docs/profileguide.html. Last accessed December, 2006.
CIA World Factbook 2007	CIA World Factbook. 2007. Available at: https://cia.gov/cia//publications/factbook/docs/profileguide.html. Last accessed February, 2007.
DHS STATcompiler as of January 2007 2007	DHS STATcompiler as of January 2007 http://www.statcompiler.com/statcompiler/index.cfm
DHS STATcompiler as of March 2007 2007	DHS STATcompiler as of March 2007 http://www.statcompiler.com/statcompiler/index.cfm
Human Development Report 2006	Human Development Report 2006. New York, United Nations Development Programme, 2006. Available at: http://hdr.undp.org/hdr2006/. Last accessed February, 2007.
UNAIDS 2006 2006	2006 Report on the Global AIDS Epidemic. Geneva: UNAIDS, 2006.
UNESCO 2005	EFA Global Monitoring Report. Education for All: The Quality Imperative. United Nations Educational, Scientific and Cultural Organization, Paris.
UNESCO 2007	EFA Global Monitoring Report 2007. Confronting the Challenges of Gender Equality and Fragile States. United Nations Educational, Scientific and Cultural Organization, Paris.
UNICEF 2007	UNICEF. Progress since the World Summit for Children. http://www.childinfo.org/areas/childmortality/u5data.php. Data accessed on Web site in April 2007.
WHO World Health Report 2004	The World Health Report 2004: Changing history. Geneva, World Health Organization, 2004. Available at: http://www.who.int/whr/2004/en/index.html. Last accessed February, 2007.
WHO/Global Summary 2002	WHO Vaccine-Preventable Diseases Monitoring System. 2002 Global Summary. World Health Organization, Geneva.
WHO/Global Summary 2005	WHO Vaccine-Preventable Diseases: Monitoring System. 2005 Global Summary. World Health Organization, Geneva.

developed by WHO, UNICEF and UNFPA, 2004.

Kenneth Hill, Carla Abou Zahr & Tessa Wardlaw: Maternal mortality in 2000. Estimates

WHO/RBM 2005	World Health Organization. Global Health Atlas. http://www.who.int/globalatlas/default. asp. Data accessed on web site in September 2006.
WHO/TB Control Report 2006	Global Tuberculosis Control: Surveillance, Planning, Financing, WHO Report 2006. World Health Organization, Geneva.
World Bank/WDI 2005	World Bank Development Indicators 2005, CD-Rom. Washington, DC: The World Bank,

2005.

World Bank/WDI 2006

World Bank Development Indicators 2006. Washington, DC: The World Bank, 2005.

Available at: http://devdata.worldbank.org/wdi2006/contents/index2.htm. Last accessed April, 2007.

World Bank/WDI Database 2007
World Bank Development Indicators Database 2007. Washington, DC: The World Bank, 2007. Available at: http://devdata.worldbank.org/data-query/. Last accessed April, 2007.