

**PLANT MATERIALS**

**2006**

**IDAHO EVALUATION SUMMARIES**

**FIELD, DSI and DEMONSTRATION PLANTINGS**

## IDAHO DIVISION I PLANT MATERIALS PLANTINGS

### **FIELD OFFICE: BONNERS FERRY**

**ID99005 Paul Headings** Regar meadow brome - Field Plantings (2). Materials ordered February 22, 1999.

Field 1—pure stand of Regar. Field 2—mixed stand of Regar and alfalfa. Purpose – demonstration planting to document growth patterns, production, and forage quality. Site characteristics – MLRA E43b, silt loam soils, 5-10 percent slopes, north aspect, 2300 feet elevation, 24 inch precipitation zone, non-irrigated, T62N R1E NW ¼ Section 2. FY99 planted spring 1999. FY00 due to dry years 1999 and 2000 stand establishment was slow, but excellent stands in each field are establishing. Plantings average 3 tons per acre. FY01 Planting 1 - The “pure” stand of Regar Brome planting averaged 2 ton/acre. A forage analysis indicted the crude protein to be 8.75%. The forage grass for hay is fine leaves and stems. The hay feeds well to animals. In hot dry weather, the “windrows” have to be carefully harvested and cured to avoid damaging brittle leaves and stems. The crop can be “pulverized” easily. The average bale weight was 103 pounds. The owner applied 110 lbs. 40-0-0 to enhance production and will increase application rates up to 200 lbs/acre 40-0-0. There were no second cuttings since the field was planted three years ago due to poor to fair moisture conditions.

Planting 2 - The Regar/Agate alfalfa mixture established well. The first cutting has grass present and makes great cattle feed. The second cutting has very little grass within the alfalfa due to slow recovery. This may be due to dry weather conditions. Also, this may be a good attribute for the producer who can sell hay with grass and no grass. FY01 Planting 1 - The "pure" stand of Regar has an excellent stand with 5 plants per square foot, good vigor, and 4000 pounds per acre production. Landowner applied 220 lbs. 40-0-0 in early spring. Planting 2 - Regar/alfalfa mixture has a good stand with 2 Regar/5 alfalfa plants per square foot, fair to good vigor, and 7000 pounds per acre production.

**FY06** The field is still in production. It has been an excellent hay crop averaging 4 tons/ac dryland over the 10 year period with one year producing 5.5 tons/ac. Landowner applies 300 pounds/ac of Nitrogen each spring. This grass needs to be managed for harvesting - cut and windrow at 50% cured. 80% cured results in loss of leaves because it is too brittle because of fine leaves. Cooperator also reports Regar also does well when planted with alfalfa, is easy to manage and he is very happy with it.

**ID00016 Boundary Creek WRP** – cropland area planted to permanent perennial species field planting. A mix of Alkar tall wheatgrass, Greenar intermediate wheatgrass, Ranger alfalfa, birdsfoot trefoil, red clover, Sherman big bluegrass, tufted hairgrass, orchardgrass, and timothy at critical area planting rates was dormant planted on 1000 acres in late fall 1999. A 42 feet air-seeder with fertilizer attachment planted mix with 2000 units per acre of nitrogen, phosphorus, potassium, and sulfur applied 1 inch below and to side of seed. FY00 excellent stand is establishing with some species as tall as 3-4 feet by early July. In October wild oats were present throughout stand. FY01 The permanent wildlife planting mixture established well utilizing the 42-foot air seeder. The drill was calibrated with the producer based upon 14.2 lbs. PLS/acre. A “flush” of wild oats occurred the first year. The stand was seeded the 1<sup>st</sup> week of November 1999. The “so called dormant planting” resulted in some sprouting of clovers due to a warmer than normal late fall. As a result, some mortality occurred in the clovers. An excellent stand of Alkar tall wheatgrass, Greenar intermediate wheatgrass, birdsfoot trefoil, Ranger alfalfa, Latar orchardgrass, timothy and clover exists. The Sherman big bluegrass is “spotty” due to becoming overpowered by the other species in the mix. There are some ridges in the field with quackgrass, which is good cover. The IDF&G is actively spot spraying the Canadian thistle. They plan to obtain a boom sprayer in order to treat the acreage more uniformly. FY02 The overall stand is good to excellent with the primary species including Alkar tall wheatgrass, Greenar intermediate wheatgrass, Latar orchardgrass and redtop. Some birdsfoot trefoil, clover, timothy, and alfalfa are present in scattered locations. Tufted hairgrass and Sherman big bluegrass were not found. FY04 prescribed burned fall 2004 (15 acres) to rejuvenate existing stand – resulted in excellent response in plant vigor. Stand is primarily Latar, Alkar, Greenar, and alfalfa – general overall stand is predominately wheatgrasses and orchardgrass. Providing excellent wildlife nesting and escape cover. **Next evaluation 2007.**

**ID04002 Dave Wattenburger** Field Planting. Delar small burnet ordered August 19, 2003. Planting seeded fall 2004. No report FY05 and FY06.

**ID06007 Idaho Fish and Game** – Field planting for wildlife winter nesting habitat. Blackwell switchgrass and 905439 switchgrass seed ordered March 9, 2006. Site characteristics: Farnhampton silt loam soil, 0-2 percent slopes, south aspect, elevation 1760 feet, 24 inch precipitation, non-irrigated, T65N R2W Sections 23 and 25. FY06 no report.

**ID06008 Bernie Heinemann** – Riparian Field Planting. Coyote willow (50); 9067541 Peachleaf willow (20); 9067546 Peachleaf willow (20); 9067549 Peachleaf willow (20); 9067375 Peachleaf willow (20); 9067376 Peachleaf willow (20); and 9067560 Peachleaf willow (20) cuttings ordered February 2006. Site characteristics: Porthill silt loam soil, 5 percent slopes, 2000 feet elevation, 24 inch precipitation, non-irrigated, T65N R1W, Section 10. FY06 no report.

**FIELD OFFICE: COUER D'ALENE**

None

**FIELD OFFICE: PLUMMER**

None

**FIELD OFFICE: SANDPOINT**

**ID96029 Lee Johnson** wood fiber mulch, Niner sideoats grama, Alma blue grama, annual rye, Durar hard fescue, Durar hard fescue/clover, prairie junegrass, and alpine bluegrass field plantings - tree nursery ground cover trial. Site loam soil (low to mod. permeability/high erosion potential), 5-10% slopes on SE exposure. FY96 planted 5/31/96. 1. Wood mulch is doing excellent job of weed control and no rodent activity to date - mulch was about 10 inches deep when applied 2. Excellent stand of annual rye established, Durar hard fescue plants are very small and establishing beneath cover crop 3. Many young Durar hard fescue plants were establishing, but very few clover plants - soil may have been too loose when seeded and clover seed may be too deep 4. Excellent initial stand of sideoats and blue grama establishing - could not tell which species was doing the best 5. Very few prairie junegrass plants establishing - appears some germination is occurring this fall 6. A lot of alpine bluegrass seedlings - appears germination did not occur until fall. FY97 and FY98 no evaluations. FY99 Treatment 1: Control no cover and normal weed control - 0 percent desirable cover with 50-80 weeds. Treatment 2: Cedar bark mulch 6-8 inches thick – 100 percent desirable cover in rows with 5 percent weeds invading mulch and some evidence of rodents in mulch. Trees near cedar mulch are more chlorotic than other treatments. Treatment 3: Durar hard fescue and annual ryegrass – 50-70 percent desirable cover with up to 20 percent weeds. Fescue blends provide more biomass than other seedings and good cover – almost 100 percent cover if mowed. Treatment 4: Durar hard fescue and Berseem annual clover – 60-80 percent desirable cover and up to 15 percent weeds. Treatment 5: blue grama and sideoats grama – 20-50 percent desirable cover with 30-80 percent weeds. Clearly the worst treatment in trial. Treatment 6: Prairie junegrass – 60-80 percent desirable cover and 10-15 percent weeds. A good alternative since this is a low growing cover. Treatment 7: Alpine bluegrass – 50-80 percent cover with 5-10 percent weeds. Less biomass produced than fescue or prairie junegrass. The alpine bluegrass produced more of a thick sod with seedheads 6-8 inches tall. This would be a better choice for nurseries that are concerned with the shading effect of taller grasses on lower branches. It also covers the ground better once established, especially in shady areas. One potential problem is its ability to spread, including into the tree rows. **This planting was canceled.**

**ID00004 Paul Jayo** Regar meadow brome field planting – irrigated/non-irrigated and hay/grazing trial. Seed ordered January 21, 2000 for delivery in early April. Site is 30-acre field with Hoodoo silt loam soil, 0-1 percent slopes, 32-inch rainfall zone, and 2485 feet elevation. FY00 planting was delayed due to dry spring weather. Cooperator plans to plant fall 2000. FY01 - FY05 no evaluations. **This planting was canceled.**

## IDAHO DIVISION II PLANT MATERIALS PLANTINGS

### FIELD OFFICE: GRANGEVILLE

**ID02002 Teresa Seloske** Forest Field Planting. Lind Douglas fir (30 plants) and Yakima Douglas fir (13 plants) ordered July 16, 2001. Plants delivered to FO April 3, 2002 by WAPMC. FY02 Planting completed April 6, 2002. Lind Douglas fir 10 percent survival with poor vigor. Yakima Douglas fir 15 percent survival with fair vigor. Survival effected by extremely dry conditions. FY03 very hot dry summer resulted in failure of this planting. File was cancelled at end of 2003. **FY06** field determination indicated the Yakima ecotype failed to establish, but the Lind ecotype is still alive. Lind ecotype has not grown much, but there is good survival of this ecotype during field evaluation in July 2006.

**ID04009 Carl Skyrman** demonstration planting. Anatone bluebunch wheatgrass and Secar Snake River wheatgrass. Seed ordered March 8, 2004. Site characteristics: Chard sandy loam soil, northwest aspect, 1820 feet elevation, 16-22 inch precipitation, non-irrigated, T26N R1E NW1/4 Section 13. FY04 – Secar and Anatone were planted side by side in the spring of 2004. Good stands for each with > 5 plants per square foot establishing and it is hard to differentiate between plantings. Anatone plants were a bit more robust than Secar plants during evaluation 7/22/04. FY05 no evaluation. **FY06** (4/25/06) good established stand, Carl will spray with Sencore for cheatgrass and ventenata control.

**ID04010 Marcia Heaton** riparian planting. 9023733 redosier dogwood, 9023739 redosier dogwood, 9023740 redosier dogwood, Laurel willow, White willow, Coyote willow, and Golden willow. Cuttings ordered March 5, 2004. Site characteristics: Wilkems silt loam soil, 2980 feet elevation, 24 inch precipitation, non-irrigated, T31N R3E NE1/4 Section 34. FY04 – approximately 60% survival for all willow species and about 20% survival for dogwood species. FY05 no evaluation. **FY06** (4/26/06) very poor survival – **Cancel**.

**ID05003 Steve Hunter** – starthistle control project. Rush intermediate wheatgrass, Tegmar intermediate wheatgrass and Newhy hybrid wheatgrass were ordered February 4, 2005. Site characteristics: 3 acres, MLRA B9, Bluesprin skeletal loam soil, 20 percent slopes, southwest aspect, elevation 2700 feet, 18 inch precipitation zone, non-irrigated, T30, R3, NW ¼ section 36. FY05 not planted. **FY06** planted May 22, 2006 into poorly prepared seedbed. Fair stand establishing with about 0.25 plants/ft<sup>2</sup> and fair vigor

**ID05004 Tony Carson (combined with ID04004)** – field planting. Anatone bluebunch wheatgrass, Magnar basin wildrye, Nezpar Indian ricegrass, High Plains Sandberg bluegrass, Rosana western wheatgrass, Sherman big bluegrass, Snake River Plains fourwing saltbush and Northern Cold Desert winterfat were ordered February 4, 2005. Site characteristics: 1 acre, Lickskillett – Tannahill silt loam soil complex, 45 percent slopes, south aspect, elevation 1960 feet, 16 inch precipitation, T28N, R1E, NE ¼ section 12. FY05 seeding was completed in early spring 2005. Half of the seeded area was treated with a straw pellet mulch. Above average spring rainfall resulted in very encouraging initial stand establishment with positively identified plants of Northern Cold Desert winterfat, Rosana western wheatgrass, Nezpar Indian ricegrass, Anatone bluebunch wheatgrass and many small seedlings present on July 13, 2005. **FY06** April 25, 2006 excellent stand establishing, primarily seedlings, but also includes a few established grasses and fourwing saltbush. Good soil moisture during evaluation and cooperators will irrigate in 2-3 weeks if no additional rains occur.

**ID05005 Tim Boddine** – medusahead control project. Newhy hybrid wheatgrass, Bannock thickspike wheatgrass, Bozoisky Russian wildrye and Rosana western wheatgrass were ordered February 4, 2005. Site characteristics: 3 acres, MLRA B9, Ferdinand loamy soil, 5 percent slopes, south aspect, elevation 3080 feet, 20 inch precipitation zone, non-irrigated, T31N, R2E, SE ¼ section 10. FY05 site was sprayed with Plateau herbicide on 4/6/05, two weeks after planting. Plant establishment is very poor for all species and it is thought that the herbicide application may have interfered with seed germination, or herbicide may have killed seedlings as they emerged. Recommend additional evaluation next year to verify these preliminary conclusions. **FY06** medusahead was too much competition and planting failed – **cancel**.

**ID05006 Gary Crea (combined with ID04008)** – Feedlot species adaptation trial. (1<sup>st</sup> planting P27 Siberian wheatgrass, Sodar streambank wheatgrass, Topar pubescent wheatgrass, Vavilov Siberian wheatgrass, Rush intermediate wheatgrass, Rosana western wheatgrass, Durar hard fescue and Alkar tall wheatgrass) (2<sup>nd</sup> planting Newhy hybrid wheatgrass, Critana thickspike wheatgrass, and Rosana western wheatgrass) Seed was ordered on February 4, 2005. Site characteristics: 0.5 acres, MLRA B9, Ferdinand-Flybow-Riggins soil complex, 2-8 percent slopes, west to southwest aspect, 20-24 inch precipitation, non-irrigated, T31N, R1E, SW of SW ¼ of section 27. FY04 overall the stand establishment is excellent due to good rainfall this year. Stand establishment exceeds 2 plants per square foot for all species except Alkar. The grasses are suppressing weeds in the feedlot. FY05 stand is spotty possibly due to excessive weed competition during establishment. **FY06** no evaluation

**ID05007 Les Killgore** – field planting. Covar sheep fescue, Durar hard fescue, Bannock thickspike wheatgrass and Rosana western wheatgrass seed was ordered on February 4, 2005. Site characteristics: 1.5 acres, MLRA E43a, loamy skeletal soil, 10 percent slopes, east aspect, elevation 2200 feet, 18 inch precipitation, non-irrigated, T28N, R1E NE ¼ section 33. FY05 and FY06 not planted.

**ID05008 Ray Stowers** – Deer Creek project. Newhy hybrid wheatgrass, Bannock thickspike wheatgrass and Rosana western wheatgrass were ordered on February 4, 2005. Site characteristics: 0.25 acres, MLRA E43a, Tanahill loam soil, 2-7 percent slopes, south aspect, elevation 2700 feet, 18 inch precipitation, non-irrigated, T27N R1E SW ¼ section 8. FY05 this seeding germinated well in the spring of 2005 with good spring rain, but went dormant and/or withered and died with no rainfall from early June through September. Spring 2006 evaluation will determine seeding success. **FY06** seedbed preparation was poor and seeding failed due to excessive competition from reed canarygrass – **cancel**.

**ID05013 Barney Chambers** – Hanley Creek riparian planting. Coyote willow, golden willow and Laurel willow were ordered February 4, 2005 for shipment in early April. FY05 - golden willow 80% survival with good vigor and 4 feet plant height. Coyote willow 50% survival with good vigor and 4 feet plant height (note coyote willow was accidentally sprayed with herbicide causing the partial plant death). Laurel willow 80% survival with good vigor and 3 feet plant height. **FY06** golden willow 60% survival with good vigor and 4 feet height; Laurel willow 60% survival with good vigor and 3 feet height; coyote willow 30% survival with good vigor and 4 feet height. Cooperator destroyed planting in late fall – **cancel**.

**ID06005 Tony Carlson** – Field planting of Rush intermediate wheatgrass, Bozoisky Russian wildrye, Magnar basin wildrye, Nezpar Indian ricegrass, and Sherman big bluegrass. Seed ordered February 21, 2006. Site characteristics: silt loam soil, 2 percent slopes, east aspect, 2100 feet elevation, 14-15 inch precipitation, non-irrigated, T28N R1E NE 1/4 Section 12. **FY06** seeded spring of 2006 (4/25/06) excellent stand establishing and seeding will be sprayed for broadleaf weed control.

#### **FIELD OFFICE: LEWISTON**

**ID82001 Richardson** Starthistle control field planting. Covar sheep fescue planted in early 1980's. FY01 good to excellent stand with 2 plants per foot squared average, excellent vigor, fair spread for bunch grass. Plants are 10 inches tall with seedheads averaging 14 inches tall and 6-inch diameter plants. Overall Covar is providing good starthistle control. Starthistle is present in plot, but not reproducing seed. Where Covar has 4 plants per foot squared, starthistle is not present. Covar is moving slowly downslope into starthistle dominated area. FY04 excellent stand of Covar with excellent vigor, 7 inch height and light infestation of yellow starthistle. FY05 and FY06 no evaluation.

**ID95028 Dau** Bannock thickspike wheatgrass and Rush intermediate wheatgrass field planting. Seed ordered 4/3/95. FY95 - FY99 no evaluations. FY00 40 plants per foot squared of Rush intermediate wheatgrass. Bannock thickspike wheatgrass failed. FY01 40 seedheads per foot squared, 4.5 feet tall, 3000 pounds per acre, estimate 500 pounds per acre seed production and stand is weed free. FY04 good stand with good vigor. This stand is suppressing yellow starthistle fairly well and also providing excellent erosion control. FY05 and FY06 no evaluation.

**ID98016 Fred Kaufman** Hycrest crested wheatgrass, Vavilov Siberian wheatgrass and Sherman big bluegrass field planting. FY98 and FY99 no evaluations. FY00 excellent stands of Hycrest and Vavilov established. FY02 excellent stand with excellent vigor for each cultivar. Hycrest crested wheatgrass suppressing cheatgrass better than Vavilov Siberian wheatgrass. FY04 excellent stand and vigor of Vavilov, Hycrest and Sherman. Stands are doing good job of suppressing weeds, providing erosion control and very good habitat for upland game birds (pheasants and quail). FY05 and FY06 no evaluation.

**ID04014 City of Lewiston – Mike Bowman** Delar small burnet field planting. Seed ordered April 6, 2004. Site characteristics: MLRA B9, 4 acres, Tainey silt loam soil, 5-10 percent slope, west to north aspect, 3000 feet elevation, 26-28 inch precipitation zone, non-irrigated. FY04 – FY05 no evaluation. FY06 Mike Bowman, Lewiston City Forester retired and is not returning phone calls. The planting may or not have occurred and if it did the planting location is not known. **Cancel**

**FIELD OFFICE: MOSCOW**

**ID06001A Lee and Roxanne Carrick** riparian field planting. Cuttings ordered August 9, 2005. Cuttings to be shipped mid-late October 2005. Site characteristics: MLRA B9, Hampson silt loam soil, 0-3% slopes, NW aspect, 2600 feet elevation, 24 inch precipitation, non-irrigated, T41N R3W NW ¼ Section 3. 85 each of Rivar Mackenzie willow, Curlew Drummond willow, and Silvar coyote willow will be dormant fall planted 2005. **FY06** planted November 4, 2006.

**ID06001B Lee and Roxanne Carrick** riparian field planting. Plants to be shipped early-mid April 2006. Site characteristics: MLRA B9, Hampson silt loam soil, 0-3% slopes, NW aspect, 2600 feet elevation, 24 inch precipitation, non-irrigated, T41N R3W NW ¼ Section 3. 60 each of Blanchard blue elderberry, Okanogan snowberry, St Maries mockorange and 125 Cheney redosier dogwood will be spring planted in 2006. **FY06** planted May 25, 2006.

**FIELD OFFICE: NEZPERCE**

**ID05009 William Stillman** – Weed control project. Rush intermediate wheatgrass, Topar pubescent wheatgrass, Bannock thickspike wheatgrass, Paiute orchardgrass, Delar small burnet and Sherman big bluegrass seed was ordered on February 4, 2005. Site characteristics: 12 acres, MLRA B9, Jacket-Larkin silt loam soil, 20 percent slopes, south aspect, elevation 2900 feet, 23 inch precipitation, non-irrigated, T36N R1E SW ¼ section 19. FY05 no evaluation. **FY06** plots were broadcast seeded March 12, 2005 and trampled by cattle and sprayed with Roundup. All plots contain intermediate wheatgrass from prior planting making identification of planted wheatgrass species difficult. Scattered patches Paiute were observed. Small burnet was observed across all plots. No Sherman was observed. Heavy competition from weeds such as starthistle, medusahead and ventenata is present.

**FIELD OFFICE: OROFINO**

**ID99010 Ray Geidl** field planting. Species include Coyote willow, Geyer 435 willow, Geyer 448 willow, Geyer 483 willow, Geyer 491 willow, Snowberry, Elderberry, Dogwood 733, Dogwood 740, and Chokecherry. FY99 and FY00 and FY01 no evaluations. FY02 Plantings are located in area with heavy reed canarygrass competition. Good survival for all willow and dogwood accessions with 4 of 5 cuttings for each still surviving, fair vigor for each, 40 inch height for all willows and 20 inches height for all dogwoods. Snowberry, Elderberry and chokecherry failed. FY03 – FY06 no evaluations.

**ID04011 Clearwater County Riparian Project.** 9067541 peachleaf willow, 9067546 peachleaf willow, 9067 549 peachleaf willow, 9067568 black cottonwood, 9067569 black cottonwood, 9023 733 redosier dogwood, 9023739 redosier dogwood, 9023740 redosier dogwood and Okanogan snowberry. Cuttings ordered March 5, 2004. Site characteristics: Cobbly soil, flat aspect, 1100 feet elevation, 26 inch precipitation, non-irrigated. FY04 – FY06 no evaluations.

**ID04012 Ray Geidl Project.** 9067541 peachleaf willow, 9067546 peachleaf willow, 9067549 peachleaf willow, 9023733 redosier dogwood, 9023739 redosier dogwood, 9023740 redosier dogwood and Okanogan snowberry. Cuttings ordered March 5, 2004. Site characteristics: fine loamy soil, flat aspect, 3000 feet elevation, 35 inch precipitation, non-irrigated (naturally sub-irrigated). FY04 – FY06 no evaluations.

**ID04013 Paul Schroder Project.** 9067541 peachleaf willow, 9067546 peachleaf willow, 9067 549 peachleaf willow, 9023733 redosier dogwood, 9023739 redosier dogwood, 9023740 redosier dogwood and Okanogan snowberry. Cuttings ordered March 5, 2004. Site characteristics: Fine loamy soil, flat aspect, 3000 feet elevation, 35 inch precipitation, non-irrigated (naturally sub-irrigated). FY04 – FY06 no evaluations.

## IDAHO DIVISION III PLANT MATERIALS PLANTINGS

### FIELD OFFICE: CALDWELL

**ID99006 Jacy Gibbs-cooperator will complete evaluations** for demo plots. Site characteristics: very warm dry summers, Cencove fine sandy loam soil, 0-2 percent slopes, about 2200 feet elevation, 8-10 inch precipitation, T3N R5W NE1/4 Section 10. Seed ordered February 24, 1999. Aberdeen accessions: Bannock thickspike wheatgrass, Sodar streambank wheatgrass, Goldar bluebunch wheatgrass, Appar blue flax, Magnar basin wildrye, Nezpar Indian ricegrass, Richfield Selection firecracker penstemon, Clearwater Selection alpine penstemon, Snake River Plain fourwing saltbush. Bridger accessions: Trailhead basin wildrye, Rimrock Indian ricegrass, M1 Nevada bluegrass, PI434231 plains bluegrass, 9005460 alpine bluegrass, High Plains Sandberg bluegrass, Shoshone beardless wildrye, 9019219 bottlebrush squirreltail, Critana thickspike wheatgrass, Wytana fourwing saltbush. Meeker accessions: Summit Louisiana sagewort, Timp Utah sweetvetch, Bandera Rocky Mountain penstemon, 9040187-bottlebrush squirreltail, 9040189 bottlebrush squirreltail, 9043501 Salina wildrye, Maybell antelope bitterbrush. Pullman accessions Secar Snake River wheatgrass, Covar sheep fescue, Canbar Canby bluegrass, Sherman big bluegrass, Whitmar beardless wheatgrass, and Schwendimar thickspike wheatgrass. FY99 no evaluation. FY00 Nezpar has excellent seedling vigor, easy to transplant, remains green, and is an attractive landscape plant. Schwendimar is best thickspike wheatgrass, remains green longer, best regrowth, responds well after mowing, good dryland and limit irrigation. Goldar and Whitman stands are very poor due to cheatgrass competition. Basin wildrye, Sherman, Secar mix good weed competition. Basin wildrye, Sherman, Covar, Secar are all good landscape plants. Using Covar along one side of property for firebreak – it will be excellent. Penstemon species are very slow growing, remain green and will be good landscape plants. Appar can be a nuisance and is not very shade tolerant. Maybell is slow growing. Timp is a preferred species by rabbits resulting in difficulty establishing stand. Summary of best plants – Grasses: Secar Snake River wheatgrass, Magnar basin wildrye, Sherman big bluegrass, Nezpar Indian ricegrass, Covar sheep fescue, sand dropseed, Bannock thickspike wheatgrass, and Schwendimar thickspike wheatgrass. Forbs: western yarrow, Drummond phlox, white evening primrose, scarlet globemallow, silky lupine, Louisiana sagewort, Rocky Mountain iris, and Appar blue flax. Shrubs: native fourwing saltbush, native basin big sagebrush, Maybell bitterbrush, curleaf mountain mahogany, Saskatoon serviceberry, Woods rose, almond, and Drummond willow. Trees: Idaho hybrid poplar, and Rocky Mountain juniper. FY01 - FY04 no evaluations. **FY05** all plants are under some type of supplemental irrigation. Both the basin wildrye accessions are doing fine and are good landscape plants. The Secar bluebunch wheatgrass is doing well and is a good landscape plant. Covar sheep fescue has done well near the edges of walks and driveways where a little additional moisture is available and the soils are deep. It did poorly in an area that had topsoil removed and the soil may be somewhat compacted and is a good xeriscape plant. The accessions of Indian ricegrass that germinated and survived are doing well. There was poor germination and survival in general. They are good xeriscape plants. Appar Lewis flax is doing well with partial irrigation and is a good xeriscape plant. Scarlet globemallow seed was collected by cooperator and seeded. It did well for a couple of years, but died after 3-4 years. A good looking xeriscape plant. All accessions of penstemon, mostly Eaton or firecracker have done well under partial irrigation and are good xeriscape plants. The curleaf mountain mahogany has done well and is a good landscaping plant. It receives some extra water. Serviceberry is doing well and is near full irrigation. Utah sweetvetch came up well, but received heavy use from rabbits and only one plant has survived. It is doing well. Golden current came into our yard through birds. Where it has volunteered, it is doing well and is a good xeriscape shrub. It needs to be pruned to make it denser since it is pretty leggy without pruning. It receives some additional water. I planted 3 or 4 silver buffaloberry shrubs and they have been slow in developing and are still quite small. They may do ok. The Maybell bitterbrush is doing fine. It receives some additional water and is a good xeriscape plant. Western yarrow was planted. It receives some additional run-on water where it was planted. It has spread readily to other areas with partial irrigation and some areas of full irrigation. In many ways this is a weed at my house. Rocky mountain iris has done fair in a place that I can fully irrigate and with no competition. Woods rose has done well and is a good landscaping plant. It is under full irrigation and should be pruned, similar to other roses to keep it under control. Western clematis started very slowly. This was from seed and not from Pullman PMC. Once it got going, after about 3 years, it has grown quite well and in fact needs to be controlled for spread. The almond we harvested near Brownlee reservoir has done very well under full irrigation. It has very good vigor, has good shape and in favorable springs has produced good seed crops. It is a medium sized tree, about 20 ft. high now. Herbaceous sagewort or Louisiana sage was planted from seed and receives partial irrigation. It does very well where it was planted and if the water was spread to a larger area, it would

expand from rhizomes into that area. Mulberry trees volunteered through the help of birds to our yard. They do very well here and are good looking medium sized trees. They get partial to full irrigation and are not producing seed yet at 3 years old and 18 ft. tall. The ones down near the river do produce seed.

**ID05010 Jacy Gibbs** – shrub test plots. Prospector common snowberry, Trapper western snowberry, St Marie's mockorange (plants) and Colfax mockorange (plants) were ordered February 4, 2005. Site characteristics: MLRA 11, silt loam to loam soil, 1 percent slope, elevation 2250 feet, 8 inch precipitation, irrigated. **Cooperator will complete evaluations. FY05** Mockorange received in April in good condition. The accessions were not marked. NOTE: this was indicated on receipt that was sent back to Pullman PMC. Wayne Crowder called shortly after this and no resolution on how to identify was made. No way to evaluate them separately. They were planted shortly after they arrived. Four mockorange received partial irrigation and had moderate competition from grass. Six mockorange received full irrigation and had little or no grass competition. Most of the partially irrigated plants died. The others have low vigor and have grown little for full irrigation. There should be better first year growth with full irrigation to be a good landscape plant. They are planted in partial shade and may come on in 2006. Seed of the snowberry accessions were received in April. Warm-cold stratification occurred per instructions. The seed was planted 1/03/06. **FY06** snowberry failed. Mockorange was not sent with separate identification so accessions can not be determined. Mockorange is still alive, but growth rates are very slow.

**ID06002 CB River Ranch** WRP upland planting. Seeding mixture includes Bozoisky Russian wildrye, Pryor slender wheatgrass, Vavilov Siberian wheatgrass, Magnar basin wildrye, Nezpar Indian ricegrass and Snake River Plain fourwing saltbush. Seed was ordered September 26, 2005 and planting date is scheduled for May 2006. Site Characteristics: Feltham loamy fine sand soil, 3-12 percent slope, NE aspect, 11 inch precipitation and site will be irrigated for establishment. **FY06** no evaluation.

**FIELD OFFICE: EMMETT**

**ID04016 Richard Zamzow** WRP upland field planting. Vavilov Siberian wheatgrass, Sodar streambank wheatgrass, Bannock thickspike wheatgrass and Magnar basin wildrye. Seed ordered July 2003. Site characteristics: fine sandy loam soil, 2100 feet elevation, 10-12 inch precipitation, aspect-flat. Planting planned for fall 2003. **FY04 – FY05** no evaluations. **FY04- FY06** no evaluation.

**FIELD OFFICE: MARSING/GRANDVIEW**

**None**

**FIELD OFFICE: MERIDIAN**

**ID07002 Doug Austin** field planting. Regar meadow brome, orchardgrass and alfalfa field planting. Seed ordered August 28, 2006. Seed will be planted in late summer – early fall and irrigated for establishment. Site characteristics: silt loam soil, 0-2 percent slope, 2800 feet elevation and irrigated. Seed was planted in early September 2006 and irrigated for establishment.

**ID07001 Wayne Newbill** field planting. Regar meadow brome and Cache meadow brome irrigated forages trial. Seed ordered August 10, 2006. Seed will be planted in late summer – early fall and surface irrigated (furrows) for establishment. Seed was delivered on 8/18/06. **FY06** weed control using 2 pints Roundup per acre was applied on 8/30/06 followed by discing, corrugating, pre-irrigation, harrowing, seeding with drill – 7 inch spacing on 9/9/06 and final corrugation. Regar is located in west field and Cache is located in east field. Fields were irrigated following planting.

**OTHER PLANTINGS MANAGED BY PMS**

**ID06003 Rebecca Laramie** Field Planting. Roadcrest crested wheatgrass and Ephraim crested wheatgrass low moisture lawn trial. Seed shipped March 2005. Seedbed preparation included roto-tilling and hand raking. Lawn was seeded on September 1, 2005. Seed was broadcast at about 600 seeds per square foot, lightly raked and the entire area was mulched with dry grass clippings and then watered. Irrigation the first month was 3 times per week. On September 7, 2005 seedlings could be seen coming through the mulch. On September 25, 2005, no noticeable difference could be seen between Roadcrest and Ephraim establishment. **FY06** - the grass seeding is looking good, did spot seeding in a few spots where stand wasn't too thick.



**ID07005 Gwen Denton** field planting. Regar meadow brome and Rush intermediate wheatgrass interseeding trial. Seed was ordered November 13, 2006. Planting will be dormant fall (Dec. – Jan.).

**FIELD OFFICE: MOUNTAIN HOME**

None

**FIELD OFFICE: PAYETTE**

None

**FIELD OFFICE: WEISER**

**ID91029 Grafe** Bannock and Critana thickspike wheatgrass field planting. Site is a sandy loam soil, non-irrigated, 12-14 inch ppt, 2500 feet elevation, and 4-8% slopes on west exposure. FY92 estimate 20% stand. FY93 survival is 90% for both species. The existing plants are healthy and holding their own with competition. Neither species is as vigorous as Oahe on same sites. FY94 survival is 95% for each species, good stands, and excellent vigor. This trial continues to improve; the stands are spreading and filling in open ground. Both species appear well adapted to site even considering the extended drought conditions. Total forage production is less than adjacent intermediate wheatgrass, but is more palatable. Plants are producing seed this year. The stands are starting to provide competition for annual weeds, grasses and cereal rye. I am now starting to see the value of these plants on some of our most droughty and limiting sites. FY95 Good stands for both Bannock and Critana (95% survival). Both species continue to improve over time. Cereal rye is not affecting growth. Neither thickspike wheatgrass is producing as well as Oahe intermediate wheatgrass. Both species would fit well with similar palatability grasses in mixture (suggest Goldar or Secar bluebunch wheatgrass). FY96 good stands of both with 6 plants/ft<sup>2</sup> of each and excellent vigor. Growth of both species is still very good and weed competition is light. Total production continues to be less than adjacent intermediate wheatgrass. FY97 good stands (5 plants per foot), survival, and vigor for both Bannock and Critana. Growth and vigor for both does not reflect the excellent moisture year we had and stands are maintaining or declining slightly. FY98 no evaluation. FY99 good stands of both species with 90 percent survival and good vigor. Producing between 500 and 1000 pounds per acre in an extremely dry April through November year. Bannock is slightly taller at 18 inches than Critana at 16 inches. Heavy grasshopper damage this year. Cheatgrass invasion is slight. FY00 no evaluation. FY01 stands of both Bannock and Critana were rated poor, with 1 plant per square foot, fair vigor and 200 pounds of production per acre. Two years of drought has heavily impacted this planting and cheatgrass is invading. FY04 – plots continue to be plagued by drought conditions and severe cheatgrass infestations. They are adapted to site, but suppressed due to these factors. FY05 no evaluation. FY06 no evaluation.

**ID94025 Eckhardt** Ephraim crested wheatgrass, Magnar basin wildrye, Mankota Russian wildrye, Trailhead basin wildrye, P27 Siberian wheatgrass, Manska pubescent wheatgrass, Reliant intermediate wheatgrass, Bannock thickspike wheatgrass, Schwendimar thickspike wheatgrass, Greenar intermediate wheatgrass, Sherman big bluegrass, Secar Snake River wheatgrass, Goldar bluebunch wheatgrass, Bozoisky Russian wildrye, Hycrest crested wheatgrass, Rush intermediate wheatgrass demo plots. Site is clay loam soil, non-irrigated, 10-12 inch ppt, 3000 feet elevation, and 5% slopes on NE exposure. Seed ordered July 1994. FY94 and FY95 due to drought conditions, seeding planned for spring 96. FY96 planted April 9, 1996 by hand planting and raking plots to control bulbous bluegrass competition. June 19, 1996 evaluation for establishment: Mankota poor, Manska good, Sherman very poor, Greenar good, Trailhead fair, Reliant good, Bozoisky good, Bannock good. July 8, 1996 establishment: Mankota fair, Manska good, Sherman poor, Greenar good, Trailhead fair, Reliant good, Bozoisky good, Bannock good, Goldar good, Rush excellent, Secar fair. Rush has the best stand establishment to date with Goldar next. FY97 no evaluation. FY98 first set of plots; Reliant is out producing all other plots, Greenar is second in production, Sherman hand planted plot is third in production, Sherman broadcast plot failed, T6633-P is fourth in production. Second set of plots; Bozoisky performed the best with Mankota second, and trailhead the poorest. The wildryes, thickspike wheatgrasses and intermediate wheatgrasses have shown adaptation to this area and could play a roll in revegetating local rangelands. FY99 plots were grazed this spring and grazing preference was evaluated. Plots: Greenar and Reliant were grazed the heaviest, followed by Mankota and Bozoisky Russian wildrye. This was uniform for all replications. Thickspike wheatgrasses and all other varieties had slight utilization. Basin wildryes were not utilized. Grazing preference for the larger plantings: Bozoisky Russian wildrye was used the heaviest, followed by Goldar bluebunch wheatgrass, and Rush intermediate wheatgrass used the least. Cattle are grazing Fourwing saltbush. The producer is very happy with results from these plots and uses the information to make his planting decisions. Cattle in mid May grazed FY00 the small plot species. Grazing preference was for Goldar, Bozoisky, and the intermediate wheatgrasses. The intermediate wheatgrasses are spreading into adjacent plots. Moderate use was made on Magnar and Trailhead. Sherman was used only slightly. Fourwing saltbush

was utilized and continues to get taller (20 inches tall). In the large acre sized plots adjacent to a Hycrest planting, grazing preference (mid May) in order are: 1) Goldar, 2) Bozoisky, 3) Rush, and 4) Secar. Use of Goldar was similar too slightly heavier than the Hycrest. FY01 all plots are grazed this year. Utilization was heaviest on Greenar intermediate wheatgrass and Reliant intermediate wheatgrass plots. The larger plantings showed grazing preference was highest for Bozoisky Russian wildrye, then Goldar bluebunch wheatgrass, followed by Rush intermediate wheatgrass. FY03 plots were grazed this fall at time of evaluation. FY04 – Cattle preference (cows were moved into field 4 days prior to evaluation on 10/5/04). Most preferred species during this period was Bozoisky-Select Russian wildrye which was grazed very close. Second most preferred species was Goldar bluebunch wheatgrass which was grazed to a uniform 2 inch stubble height. Secar Snake River wheatgrass and Rush intermediate wheatgrass were not utilized. FY05 and FY06 no evaluation.

**ID94026 Weber** Goldar bluebunch wheatgrass, Rush intermediate wheatgrass, Luna pubescent wheatgrass, Secar Snake River wheatgrass, Greenar intermediate wheatgrass, Schwendimar thickspike wheatgrass, Bozoisky Russian wildrye, Bannock thickspike wheatgrass, Delar small burnet, Firecracker and Alpine penstemon, Sherman big bluegrass, Wytana fourwing saltbush, and Rincon fourwing saltbush demo plots. Site is stony clay loam soil, non-irrigated, 16 inch ppt, 3200 feet elevation, 0-2% slopes. Seed ordered July 1994. FY94, FY95, and FY96 due to drought conditions, seeding not planted. FY97 seeded May 16, 1997 with good rains following planting. Weed competition is high. In general initial establishment was good for wheatgrasses, fair for wildryes and poor for forbs. FY98 rainfall was 150 percent of average this year resulting in a flush of weeds. All plots except forbs were sprayed for broadleaf weed control and were shredded to reduce overstory competition. The most successful plants include: GRASSES Rush is by far the superior plot from standpoint of vigor, total growth, and total production. Luna is rated second and Reliant is rated third. Other grasses are only marginally successful to non-existent due to possibly saturated soils and weed competition during the establishment year. FORBS Delar is doing very well and appears very hardy and adapted to wet soil conditions. Penstemons and Lupine did not establish. SHRUBS Rincon is taller (10-15 inches) than Wytana (4-6 inches). FY98 no evaluations. FY99 Weeds and saturated soils are a problem on this site. Most successful plants – grasses: Rush intermediate wheatgrass followed by Luna pubescent wheatgrass, and Reliant intermediate wheatgrass, with others only marginally successful; Forbs: Delar small burnet is performing very well and no other forbs established; Shrubs: Rincon fourwing saltbush is superior to Wytana fourwing saltbush on this site. FY00 no evaluation. FY01 following two years of extreme drought Greenar intermediate wheatgrass was the most productive and vigorous followed by Reliant intermediate wheatgrass and Luna pubescent wheatgrass. Rush intermediate wheatgrass, Mankota Russian wildrye, and Manska pubescent wheatgrass did not grow much this year. Magnar basin wildrye was superior to Trailhead basin wildrye in production and survivability. Thickspike wheatgrass and Russian wildrye accessions grew very slowly. Delar small burnet plants are not handling drought well and are dying. Rincon fourwing saltbush is better than Wytana fourwing saltbush with some plants to 18 inches in height. Weeds are infesting site. FY02 was a very dry growing season. Intermediate wheatgrasses - Greenar is producing more forage than any other species, Greenar is not spreading as fast as Rush or Reliant which is probably an advantage on this droughty site, Luna is the best pubescent wheatgrass, but not producing as much as Greenar. Basin wildryes - Magnar and Trailhead are nearly identical in production with Magnar slightly higher with more vigor than trailhead. Russian wildrye - Bozoisky is by far the best performer of the R. wildryes. Small burnet - Delar is no longer present. Fourwing Saltbush - Rincon is a little better than Wytana, but they lack vigor. Thickspike wheatgrass - all accessions are barely surviving. FY06 no evaluation. **Next evaluation scheduled for 2007.**

**ID96024 Howard Sutton** Rush intermediate wheatgrass, Luna pubescent wheatgrass, and Oahe intermediate wheatgrass field planting. Site is loam soil, non-irrigated, 15-17-inch ppt, 3320 feet elevation, 1-4% slope on south exposure. Seed ordered March 14, 1996. FY96 planted in May into good seedbed with good weed control. Good stand establishing with about 3 plants per foot squared, each species was planted with alfalfa in alternate rows and alternating sections. FY97 good stands with excellent vigor of each cultivar. The Oahe/alfalfa stand was cut for hay and produced 1.5 tons/acre. Because of topography the Rush/alfalfa and Luna/alfalfa were not cut for hay. The entire field was grazed; grazing was uniform across all trials so preferences could not be determined. Producer is very happy with all three from standpoint of production potential when seeded with alfalfa. FY98 good stands and vigor for each species with about 7 plants per square foot. Yield for all species was about 5000 pounds per acre or about 3 AUMs per acre. Cattle are selecting Luna as first choice, then go to Rush before Oahe. The Rush was more mature than Luna when steers were put in pasture which may account for selection choices. FY99 good stands and vigor of all three species. Entire 84 acre seeding provided 135 AUMs or 1.6 AUMs/ac. Due to later season of use; cattle prefer Luna and Oahe to Rush. Rush initiates growth earlier and is more mature when cattle are turned into pasture, which probably accounts for this preference. FY00 similar report to last year. FY01 good stands and vigor for all species. Grazing preference

continues to be for Oahe, followed by Luna, and the Rush. Production is about the same for all species although reduced this year due to two years of extreme drought. FY02 good stand, and vigor with greatly reduced production this drought year for all accessions. Produced 0.5-0.7 AUM/Acre for each accession, less than 50% of the normal precipitation year. Grazing is slowing spread of these species. FY04 – good stands with good vigor for all species. Production was approximately 0.7 AUMs per acre. FY05 and FY06 no evaluation.

**ID02010 Hugh Pangman - New Meadows Riparian Planting.** 9067541 Peachleaf willow - Baker source and Golden willow. 50 cuttings ordered February 11, 2002 for shipment in early May 2002. To be planted with waterjet stinger. FY02 willows were planted through cobbly site using a backhoe to watertable located at 5-6 feet depth. 95 survival of each species. Peachleaf willows are 18-20 inches tall and Golden willows are 24 inches tall. Golden willows are more vigorous with more stem growth. FY03 Peachleaf willow 95 percent survival with 36-48 inch height. Golden willow local cuttings also have 95 percent survival with 48 inch plus height. Producer is please with this planting. FY04 no evaluation. FY05 end of 4<sup>th</sup> growing season - peachleaf willow 90% survival with excellent vigor, 15 feet plant height, 10 feet crown width, 3 inch DBH. Golden willow 90% survival with very good vigor, 10 feet plant height, 6 feet crown width and 2 inch DBH. Peachleaf plants are more vigorous than golden willow, but in a slightly better site based on soil and moisture availability. Plants are protected from grazing by domestic livestock. FY06 no evaluation.

**ID02011 Tom Vogel - Paddock Riparian Planting.** 9067546 Peachleaf willow - Burns source and local coyote willow. 50 cuttings ordered February 11, 2002 for shipment in late March 2002. To be planted with waterjet stinger. FY02 willows were planted on April 3, 2002 using the waterjet stinger. Stream was dry for most of July and August. Peachleaf willows have about 75 percent survival with some leader growth up to 36 inches. Coyote willow has about 60% survival. FY03 - FY06 no evaluations.

**ID02014 Mink Land and Livestock Riparian Planting.** 9067549 Peachleaf willow - Prairie City source and local source coyote willow, 2002 for shipment in late March 2002. To be planted with waterjet stinger. FY02 Peachleaf willow survival 50% and Coyote willow survival 10%. Planting depth (soils were very dry for most of season) was probably too shallow and plant perhaps should have been completed sooner. FY03 Peachleaf willow 80 percent survival with 48 to 96 inch height. Coyote willow local cuttings have 65 percent survival with 24 to 36 inch heights. FY04 – FY06 no evaluations.

**ID02017 Jim Eckhardt Field Planting - Plateau Herbicide Trial** (4 oz, 8 oz, 12 oz, Control 4 oz, 8 oz, 12 oz). Seed ordered March 20, 2002 for shipment in early October. Species include: Magnar basin wildrye, Trailhead basin wildrye, Bozoisky Russian wildrye, Mankota Russian, Bannock thickspike wheatgrass, Critana thickspike wheatgrass, Goldar bluebunch wheatgrass, High Plains Sandberg bluegrass, Vavilov Siberian wheatgrass, CD-II crested wheatgrass and Hycrest crested wheatgrass. Site Characteristics: MLRA B10, Deshler-Devon silty clay loam soil, 2-5 percent slope, south aspect, 2600 feet elevation, 12 inch rainfall zone, T11N R6W NE 1/4 NW1/4 Section 1. FY02 Plateau was applied (4, 8 and 12 ounce rates) March 27, 2002 by Joe Vollmer. Did not control salsify, fiddleneck or sunflower. Planted November 4, 2002 under dry/cold conditions with a rangeland drill at 12-inch spacing. FY03 three planted species established this year: 1) Vavilov Siberian wheatgrass had the best stand and was the most vigorous. It did not grow in the untreated control plot – established well in the 4 and 8 ounce treatments – did not establish in the 12 ounce treatment; 2) CD-II crested wheatgrass was not as vigorous as Vavilov and had fewer plants established. It had no establishment in the no treatment - some establishment in the 4 ounce treatment – good establishment in the 8 ounce treatment – no establishment in the 12 ounce treatment; 3) Hycrest crested wheatgrass was the least vigorous of the establishing species with 30-35 percent fewer plants than Vavilov and CD-II. It had no establishment in the untreated plot - spotty establishment in the 4 and 8 ounce plots – no establishment in the 12 ounce plots. At this evaluation the 8 ounce treatment appears to be the best rate for Plateau herbicide. FY04 – The best stands include: Vavilov Siberian wheatgrass with good stand with good vigor; CD-II crested wheatgrass with fair stand with fair vigor; Hycrest crested wheatgrass with fair stand with fair vigor. All other planted species appear to have failed. The best cheatgrass control rate was 8 ounces/acre of Plateau herbicide. 4 ounces is not enough and 12 ounces effects perennial plant growth. The Plateau application has helped existing bottlebrush squirreltail. Conservationist would not recommend this method seedbed preparation because he does not feel the additional expense warrants the limited vegetation produced. FY05 and FY06 no evaluation.

## IDAHO DIVISION IV PLANT MATERIALS PLANTINGS

### **FIELD OFFICE: BURLEY**

**ID94003 Bronson** Bozoisky Russian wildrye, Mankota Russian wildrye, Trailhead basin wildrye, Magnar basin wildrye, Goldar bluebunch wheatgrass (firebreaks and winter grazing). Site is sandy loam soil (weakly saline), 9-10" ppt, partially irrigated, 4800 feet elevation, 0-2% slopes. Species seeded in fall of 1994 with good seedbed. FY95 good stands of Mankota, Magnar and Trailhead; fair stands of Bozoisky and Goldar. All seedings are establishing well except in weedy areas. No seed production during establishment year. FY96 good stand of Goldar, fair stand of Mankota and Magnar, and very poor stand of Trailhead and Bozoisky. All plants that are present look good and are producing seed. There are weeds present including cheatgrass, tumble mustard, Russian thistle, broom snakeweed and sagebrush. FY97 Goldar full stand, Trailhead has improved and is spreading, Magnar is very thin, and both Russian wildryes are adapted with thin stands. FY98 good stands of Bozoisky and Goldar and fair stands of Mankota, Trailhead and Magnar. Stands are grazed in winter. FY99 Good stand and vigor of all species. All species are in same pasture and the Bozoisky is grazed closer than the other species. FY00 fair to good stand of all species. Cooperator is very pleased with all species and prefers them over crested wheatgrass varieties. Site was grazed in spring. Cooperator states that livestock make good use of Bozoisky and Mankota in spring, Trailhead in winter, and Magnar in fall and winter. Magnar stays greener than Trailhead. FY01 this site is suffering from two years of drought. Mankota Russian wildrye has 36-inch height, fair to good stand and good vigor. Bozoisky has 20-inch height, fair stand with fair vigor. Magnar has 30-inch height and Trailhead has 20-inch height and both have fair to poor stands with fair to good vigor. Goldar has 24-inch height, fair to poor stand with good vigor. FY02 Survival/Plant Height - Mankota 75%/26 inch, Magnar 80%/40 inch, Trailhead 80%/36 inch, Bozoisky 75%/30 inch, Goldar 30%/26 inch. Magnar and Trailhead are only lightly grazed and are showing very little effect from grazing. Bozoisky and Mankota stands are heavily grazed and stand are beginning to decline. Goldar stand is also heavily grazed and stand has declined significantly. Producer comments indicate that Goldar is always the first species to be grazed in this pasture followed by the Russian wildrye. FY03 - FY04 no evaluation. FY05 Mankota good stand and vigor with 24 inch plant height; Bozoisky good stand and vigor with 36 inch plant height; Goldar good stand and vigor with 26 inch plant height; Magnar poor stand with good vigor and 60 inch plant heights; Trailhead poor stand with good vigor and 60 inch plant heights. Cooperator states that Goldar is the first plant grazed each season and then Bozoisky and Mankota are utilized. Magnar and Trailhead are the last grasses utilized each season, but calves do utilized the basin wildrye stands for thermal cover. Both basin wildrye accessions are spreading into other plots. **Next evaluation FY08.**

**ID96012 Poulton** Garrison field planting for plug nursery. Seed ordered 12/8/96. FY96 no evaluations. FY97 field has full stand with 2 plus plants/ft<sup>2</sup>. Plants have height of 36 inches and no weeds. Stand is gravity irrigated and was fertilized with 80 pounds of N in early June. FY98 excellent stand that has improved significantly in the last year. The stand was hayed this year. FY99 good to excellent stand. The stand was 36 inches tall when swathed for hay and had 6 inches of regrowth in early September. Cooperator is very pleased with this grass. Elk are utilizing planting. FY00 planting was cut for hay and elk are utilizing it heavily due to drought conditions. FY01 due to drought conditions, this planting was hayed earlier than normal and has been heavily grazed. Production was below normal. Stand is solid with no bare spots or invading species. FY02 same comments as last year. FY03 - FY04 no evaluation. FY05 cooperator indicated that yields are up over previous years due to better rainfall this spring with 24-30 inch vegetative heights and seedheads up to 48 inches in height. The wetter areas of the field are primarily Garrison even in areas where it was not originally planted. The original planting was irrigated, but is no longer irrigated today and Garrison is going out of this area. Cooperator like Garrison as a forage species and would like to have in more of his pastures. Note: Garrison creeping foxtail requires full moisture either through irrigation and/or sub moisture conditions. It is very productive and a very high quality forage species if fully irrigated and if fertilized. You might consider recommending a fertility program to Mike if he wants to increase production. This planting is providing good information and should be maintained. **Next evaluation FY08.**

**ID97006 Gary Jones** Field planting of Garrison creeping foxtail. Site is silt loam soil, irrigated, 5000 feet elevation, and 0-3% slope on south exposure. Seed ordered 10/17/96. FY97 new seeding and very difficult to determine establishment. FY98 poor stand establishing with .5 plants per foot<sup>2</sup>. FY99 good stand with about 4 plants per square foot and 4000 pounds per acre production. Fertilizer would benefit stand and reduce weeds. FY00 good stand with

excellent vigor. Planting was hayed this year. FY01 this is a good planting. It was cut earlier than usual for hay due to shortage of irrigation water. Yield was down this year, but cooperators were satisfied with yield given the droughty conditions. FY02 landowner is enthused about Garrison production/performance and plans to plant additional field to this species. FY03 - FY04 no evaluation. FY05 Garrison is probably about 50% of the stand throughout field. Cooperator likes Garrison and said it is an excellent hay and grazable forage. Note: Garrison creeping foxtail requires full moisture either through irrigation and/or sub moisture conditions. It is very productive and a very high quality forage species if fully irrigated and if fertilized. You might consider recommending a fertility program to Gary if he wants to increase production. This planting is providing good information and should be maintained. **Next evaluation FY08.**

#### **FIELD OFFICE: GOODING/FAIRFIELD**

**ID00005 Camas SCD (Koonce)** formerly ID86010 Koonce multiple species demo plots. FY99 field evaluation determined these plots to be contaminated and planting was destroyed, site cleaned-up and fallowed during 1999, and was replanted in the spring of 2000. Plots replanted May 1, 2000. Plots will be irrigated the first growing season. **FY00** plots were irrigated until mid June, and then discontinued. Most of the wheatgrasses sprouted in the central and northern portions of the plot, but remained small at evaluation time due to dry season. Plot remains relatively weed-free except the southernmost 15 feet of the plot (sheep fescue area) which is a solid stand of globe mallow. The fescue is sprouted underneath the large mallow leaves. This is a particularly difficult weed to control once established. Special attention needs to be directed here in spring 2001. **FY01** the plots have been subjected to two seasons of unfavorable plant growth (dry springs) and one of the lowest winter snow pack recorded on the Camas Prairie. Still, all varieties exhibit some level of success except for the following varieties which could not be found for observation: Durar hard fescue, Nezpar Indian ricegrass, 9043501 Salina wildrye, and Thurber's needlegrass. These varieties did not establish at all or remain yet as dormant seed due to drought. Some of the absent species may have germinated but died unnoticed due to drought. Weed competition most likely is not a factor of establishment difficulties in the plot. Possible exceptions may be in the Covar sheep fescue area that had significant amounts of common mallow in 2000 but is now under control due to spot spraying. Scouringrush is invading in the Bighorn sheep fescue and Magnar basin wildrye areas and may be a factor there. The entire demo plot was spot-sprayed in 2001 twice (last of June and first of August) with 2, 4-D/Banvel. At the time of this evaluation the plot did not contain weed problems significant to grass establishment. The wheatgrasses are performing the best. The highest performing wheatgrasses include Rush and Reliant intermediate wheatgrasses, Manska and Luna pubescent wheatgrasses, CDII and Nordan crested wheatgrasses, Bannock thickspike wheatgrass, and Pryor slender wheatgrass. Weak wheatgrass performance was observed with Arriba western, Whitmar beardless wildrye, San Luis slender wheatgrass, Critana thickspike wheatgrass, Ephraim crested wheatgrass, Douglas crested wheatgrass, and P27 Siberian wheatgrass. Bozoisky and Mankota Russian wildrye performed moderately, but the other wildryes either did poorly (Volga Mammoth and Magnar) or did not establish (Salina and Trailhead). Manchar and Liso smooth bromes have done well considering the drought with moderate performances, but Garnet and Bromar mountain bromes and Regar meadow brome did not fare so well and have overall weak ratings. The fescues, needlegrasses, orchardgrasses, ricegrasses, timothy, and foxtail are currently performing weakly or did not establish. Sherman big bluegrass had low establishment density but the existing plants have good vigor with many seedheads produced. **FY02** drought continues. Excellent plots include: Rush, Greenar, Reliant, Topar, Manska, Luna, Bozoisky, CD-II, Hycrest, and Nordan. Good plots include: Rosana, Manchar, Regar, Alkar, Jose, Liso, Oahe, Tegmar, 238, Goldar, P-7, Mankota, Secar, Pryor, Bannock, Schwendimar, Sodar, Sherman, Vavilov, and Magnar. Fair plots include: Latar, Garrison, Arriba, Climax, Covar, Volga, Whitmar, San Luis, Critana, Ephraim, Douglas, P-27, Rimrock, High Plains, and Trailhead. Poor plots include: Paiute, Garnet, Bromar, Durar, 902484, and 9040137. Failed plots include: Salina and Nezpar. **FY03** plants with best density, vigor and seed production include: Rush, Reliant, Manska, Bozoisky, CD-II, Nordan, Arriba, Greenar, Topar, P7, Mankota, Hycrest, Vavilov, Alkar, Jose, Oahe, Tegmar, Luna, Ephraim, and P27. Generally, the wheatgrasses are out performing the fescues, wildryes, needlegrasses, bromes, bluegrasses, timothy and orchardgrass. Plants that have failed include: Paiute, Rimrock, 9040137 needlegrass, Nezpar, Volga, 9043501 Salina wildrye, Bighorn sheep fescue. **FY04** wheatgrasses as a group dominate as the best adapted species for this site. Intermediate (Rush and Reliant) wheatgrass, pubescent (Manska and Luna) wheatgrass, Sherman big bluegrass and Trailhead basin wildrye improved over last year despite unfavorable conditions. All other plots remained static or declined in performance. Garnet and Bromar mountain brome and San Luis slender wheatgrasses (all short-lived perennials) died out this past year. **FY05** cooperators would like to **cancel** this plot due to maintenance issues – SCD will look for another plot location.

**ID00006 Bill Simon** Bannock thickspike wheatgrass District Seed Increase. Seed ordered February 10, 2000 for mid April delivery. FY00 this new Bannock seeding in spring 2000 was installed adjacent and south of existing Bannock field under file ID98020. Bannock was drilled at 3 pounds per acre PLS on 24-inch centers. The field was helicopter sprayed with 2, 4-D the third week of June. Where helicopter missed, Russian thistle prevailed this year but should diminish next year. At evaluation time on November 1, 2000, the stand was well on its way to establishment considering the dry year. FY01 unfavorable moisture year - 200 pounds per acre seed production. FY02 unfavorable moisture year - 110 pounds per acre seed production. FY03 good stand and vigor – field produced 43 pounds per acre probably due to spring frost (May 19 - 16<sup>o</sup>, May 20 – 21<sup>o</sup>, June 23 – 26<sup>o</sup>), low precipitation, and very hot summer. FY04 - good stand and vigor with field producing 90 pounds of clean seed per acre. Producer feels too much vegetative growth was produced this year due to spring rains hurt seed production. Producer plans to maintain stand for seed production one more year. FY05 Wet spring, dry summer and fall. Bill said about 150 pounds per acres clean seed on this field, up from 90 pounds last year. Sprayed with 2 pints/ac 2,4-D in mid June. No fertilizer. Bannock is first seed harvested on the Camas Prairie, about August 1. FY05 wet spring – harvested about 150 lb/ac clean seed on August 1. Field was sprayed with 2 pints/ac 2,4-D in mid June, field was not fertilized. FY06 no evaluation.

**ID01007 Spring Cove Ranch – Butler** demonstration plantings of Magnar basin wildrye, Snake River Plain fourwing saltbush, and Northern Cold Desert winterfat. Seed ordered March 16, 2001. Site characteristics: Planting 1. Vertisol soil, 11-inch rainfall, irrigated, 3300 feet elevation, south of Pioneer Reservoir. Planting 2. Sodic soil, 12-inch rainfall, irrigated, 3500 feet elevation, near Clover Creek – Hill City Road – southern base of Bennett Mountain foothills. FY01 - FY04 seed not planted due to extreme drought. Cooperator plans to plant fall 2004. FY05 Planting Site 1: Seed again not planted. Dan said he still wants to drill the Magnar next spring (2006) in the planned site (Planting Site 1). Said site in 2005 was too dry. As of 1/10/06 site is under flood waters. Moisture should be good for spring 2006 planting. He said he will drill seed in spring 2006.

Planting Site 2: Seeding cancelled, property ownership change. Winterfat and Four-wing seed retained by Dan Butler.

**ID02015 Bob Josaitis** Field Planting. 905439 switchgrass (Bridger PMC) and Blackwell switchgrass (Manhattan PMC) were ordered March 15, 2002 for shipment about April 1, 2002. Purpose: portion of seed mix for wildlife nesting cover. Site Characteristics: MLRA 11a, Harsand fine sandy loam soil, 0-2 percent slope, 3700 feet elevation, 11 inches precipitation, full irrigation, T6S R15E Section 4. FY02 - FY04 seed not planted due to drought and field change. Seeding planned for spring 2005. FY05 905439 and Blackwell switchgrass seeded on different halves of same field via Brillion drill on July 4, 2006. Late drilling was to favor establishment of warm season switchgrass over the cool season grasses which dominate the mix. Due to round smooth small seeds both switchgrass varieties added by hand into small seed box on Brillion as field was being drilled. Large seed box contained mixture of Bannock thickspike wheatgrass, Alkar tall wheatgrass, Rush intermediate wheatgrass, Eski sainfoin, and Delar small burnet. Site was smooth and firm prior to drilling. Much seed after drilling was not adequately covered by Brillion press wheels. Field was therefore harrowed after drilling, all seed thereafter covered. Irrigated immediately and for remainder of season. Young redroot pigweeds dominating by end of July, killed with 0.5 pint/ac LV-4 (2,4-D). Some sainfoin germinated by then but not affected by LV-4. Sainfoin well represented by end of season. Bannock germinated within 7 days of seeding, was main planned grass at end of season. Other species in mix including switchgrass appeared to remain dormant this first season. Site dominated by bristlegrass this first year (Setaria), should diminish or disappear in 2006. Evaluate in 2006 growing season. Site was seeded for permanent nesting habitat.

**ID07006 Hugh Koonce** field planting. Sodar streambank wheatgrass was ordered November 13, 2006 for critical area planting seeding. Seed will be dormant broadcast planted between November and January and harrowed to incorporate seed. Site characteristics: clay loam to silt loam soil, 5,000 feet elevation 2-6 percent slopes, 12 – 16 inch precipitation.

**FIELD OFFICE: JEROME**

None

**FIELD OFFICE: RUPERT**

None

**FIELD OFFICE: SHOSHONE/HAILEY**

None

**FIELD OFFICE: TWIN FALLS**

**ID00007 Twin Falls SWCD/Twin Falls Highway District** Drought tolerant landscape-weed control demonstration plantings. Seed ordered March 1, 2000 for late March delivery. Planting 1: Vavilov Siberian wheatgrass, Bozoisky Russian wildrye, and Ladak alfalfa. Planting 2: Hycrest crested wheatgrass, Bozoisky Russian wildrye, and Ladak alfalfa. Planting 3: Secar Snake River wheatgrass, Critana thickspike wheatgrass, Trailhead basin wildrye, Rimrock Indian ricegrass, and Wytana fourwing saltbush. Planting 4: Secar Snake River wheatgrass, Bannock thickspike wheatgrass, Magnar basin wildrye, Nezpar Indian ricegrass, and Snake River Plain fourwing saltbush. Site characteristics: MLRA B11A, Portneuf silt loam soil, 0-2 percent slopes, north exposure, 3800 feet elevation, 10-12 inch precipitation, irrigated for establishment only, T11S R18E SW1/4 of SW1/4 of Section 13. FY00 due to very dry spring the planting was delayed until better planting conditions occur. FY01 site was planted in mid to late April and sprinkler irrigated in May to assist with plant establishment. Site was also mowed several times during growing season for weed control. Because of mowing, species identification was not possible – estimated initial stand establishment for all plantings are fair with good plant vigor. FY02 introduced plantings are well established - native plantings failed. Introduced seed of Vavilov Siberian wheatgrass (15 lb) and Bozoisky Russian wildrye (5 lb) was ordered on September 15, 2002 to replant failed portion. Planting completed for October 25, 2002 (dormant planting). FY03 field observation determined that little establishment has occurred this year due to drought conditions. FY04 stands were mowed in June and inadequate moisture was available for regrowth. Wytana fourwing saltbush and Snake River Plains fourwing saltbush are becoming more evident with scattered plants throughout plantings 3 and 4. Mowing is keeping the fourwing saltbush short, but does not appear to be killing the shrubs. FY05 this is the first year of above normal spring moist since plantings were installed. Planting 1: good stand of Vavilov (2-3 plants/ft<sup>2</sup>), Bozoisky (2 plants/ft<sup>2</sup>) and alfalfa (< 1 plant/ft<sup>2</sup>) and good vigor for grasses and poor vigor for alfalfa. Planting 2: good stand of Hycrest (3 plants/ft<sup>2</sup>) and thickspike (2 plants/ft<sup>2</sup>). Wytana fourwing saltbush and Snake River Plains fourwing saltbush are becoming more evident and plants are larger than last year throughout the older plantings 3 and 4. FY06 no evaluation.

**ID02009 Shoshone Creek Riparian Planting – Rob Rogerson.** 9067541 Peachleaf willow - Baker source, 9067549 Peachleaf willow - Prairie City source, and 9067560 Peachleaf willow - Deer Creek source. Cuttings ordered February 11, 2002 for shipment April 1, 2002. FY02 - 9067549 60 percent survival with good vigor - 9067541 76 percent survival with good to excellent vigor - 9067560 50 percent survival with fair vigor, native Planeleaf willow 100 percent survival with excellent vigor. Death loss can primarily be related to livestock damage when cattle were place in field for 5 days. FY03 no evaluation. FY04 9067549 peachleaf willow failed, 9067541 peachleaf willow 24 percent survival with fair vigor, 9067560 peachleaf willow not evaluated, native willows 100 percent survival with good vigor. FY05 9067541 28% survival with good vigor and 18 inch height; 9067549 10% survival with good vigor and 24 inch height; 9067560 failed; native Planeleaf willow 100% survival with good vigor and 24 inch height. FY06 no evaluation.

**ID03001 Walt Coiner Field Planting.** Purpose: Field Planting - windbreak interspace perennial cover/weed control study - irrigated-semi irrigated-dryland trials. Seed was ordered on September 17, 2002. Approximately 1 acre per species - broadcast seeding rates - Aberdeen PMC broadcast planters were used for seeding - dormant fall planting completed November 4 and 5, 2002. **Irrigated species:** Durar hard fescue; Sherman big bluegrass; Foothills Canada bluegrass, and Talon Canada bluegrass. **Semi Irrigated species:** Covar sheep fescue; Sodar streambank wheatgrass; Paiute orchardgrass; Ephraim crested wheatgrass; Sherman big bluegrass; Roadcrest crested wheatgrass; and Quatro sheep fescue. **Dryland species:** Vavilov Siberian wheatgrass; Rosana western wheatgrass and Bozoisky Russian wildrye. FY03 initial evaluation August 20, 2003. FY04 evaluation September 13 2004. FY05 evaluation August 11, 2005 following well above average spring moisture.

Species	<u>Stand</u>				<u>Vigor</u>			
	2003	2004	2005	2006	2003	2004	2005	2006
<u>Irrigated Perennial Cover</u>								
Sherman big bluegrass	good	fair	fair	fair	exc.	fair	exc.	exc.
Talon Canada bluegrass	good	exc.	exc.	exc.	exc.	exc.	exc.	exc.
Foothills C. bluegrass	exc.	exc.	exc.	exc.	exc.	exc.	exc.	exc.
Durar hard fescue	fair	exc.	fair	good	exc.	exc.	fair	good
<u>Semi-Irrigated Perennial Cover</u>								
Covar sheep fescue	poor	fair	good	good	fair	good	exc.	exc.
Quatro sheep fescue	poor	good	exc.	good	fair	good	exc.	exc.

Newhy hybrid wheatgrass	poor	failed	fair	fair	fair	v. poor	good	good
Roadcrest c. wheatgrass	good	fair	poor	poor	good	good	good	fair
Ephraim c. wheatgrass	exc.	fair	exc.	exc.	good	fair.	exc.	exc.
Sodar s. wheatgrass	good	poor	poor	poor	fair	poor	poor	poor
Paiute orchardgrass	fair	fair	fair	fair	fair	fair	fair	fair

#### Dryland Perennial Cover

Vavilov S. wheatgrass	good	exc.	exc.	good	good	exc.	exc.	good
Bozoisky R. wildrye	poor	v. poor	good	good	fair	poor	good	good
Sherman big bluegrass	v. poor	v. poor	good	good	poor	v. poor	good	good
Rosana w. wheatgrass	fair	good	exc.	exc.	good	good	exc.	exc.

#### Recommendations based on four evaluation years

Irrigated – Talon Canada bluegrass and Foothills Canada bluegrass are best fully irrigated choices.

Semi-irrigated – Ephraim crested wheatgrass and the good dryland species Vavilov, Rosana, Bozoisky.

Dryland – Vavilov Siberian wheatgrass mixed with Rosana western wheatgrass or Bozoisky.

**ID04003 Steve Schuyler** field planting – windbreak. Siouland poplar, Carolina poplar, Golden willow and Laurel willow cuttings. Cuttings ordered January 12, 2004. Site characteristics: 0-1 percent slope, north aspect, 8-10 inch precipitation zone, irrigated-gravity, Portneuf silt loam soil. Planted April 10, 2004 – weed barrier fabric was installed – planting protected with snow fence along west edge. FY04 survival and height - 91 percent – 35 inches Laurel willow, 42 percent – 6 inches Carolina poplar, 82 percent – 42 inches Golden willow, 0 percent Siouland poplar. FY05 replacements ordered February 22<sup>nd</sup> 10 golden willow, 25 Carolina poplar, and 5 Laurel willow. Evaluation August 11, 2005- Laurel willow 94% survival with excellent vigor, 8 feet height and 5 feet crown width; Carolina poplar 58% survival with excellent vigor, 9.3 feet height and 7.5 feet crown width; Golden willow 82% survival with excellent vigor, 9.5 feet height and 11 feet crown width.; Siouland poplar failed.

**ID04006 Dickenson 319** riparian woody planting. Laurel willow, golden current, Wood’s Rose, redosier dogwood, Siberian peashrub, coyote willow, golden willow, chokecherry, blue spruce, and Austrian pine. Plantings are protected from grazing and grass is mowed around pines, spruce, juniper and sumac. FY04 planted in May 2004. Plantings are protected from grazing with a fence and arranged in clumps (copses) for natural appearance. Laurel willow 92 percent survival, excellent vigor, 24-36 inch height. Golden current 100 percent survival, excellent vigor, and 18-24 inch height. Wood’s rose 100 percent survival, excellent vigor, and 18-24 inch height. Redosier dogwood 60 percent survival, fair vigor, and 18-24 inch height. Siberian peashrub 100 percent survival, excellent vigor, and 18-24 inch height. Coyote willow 80 percent survival, good vigor and 12-48 inch height. Golden willow 100 percent survival, excellent vigor and 72 inch height. Chokecherry 23 percent survival, poor vigor and 36 inch height. Blue spruce 73 percent survival, good vigor and 36 inch height. Austrian pine 100 percent survival, excellent vigor and 36 inch height. FY05 evaluation August 11, 2005- Laurel willow 100% survival, excellent vigor, 4-8 feet height and 2 feet crown width; Golden current 92% survival, excellent vigor, 4 feet height and 2.5 feet crown width; Wood’s rose 100% survival, excellent vigor, 2.5 feet height and 3 feet crown width; Redosier dogwood 83% survival, excellent vigor, 4 feet height and 2 feet crown width; Siberian peashrub 12% survival, very poor vigor; Coyote willow 33% survival, good vigor, 5 feet height and 0.5 feet crown width; Golden willow 90% survival, excellent vigor, and 6 feet height; chokecherry 27% survival, fair vigor and 4.4 feet height; blue spruce 73% survival, fair vigor and 4.5 feet height; Austrian pine 100% survival, excellent vigor and 4.6 feet height; Rocky Mountain juniper 100% survival, excellent vigor and 14 inch height; Skunkbush sumac 80% survival, good vigor and 2 feet height.

**ID05002 Perinne Coulee 319 Project** riparian planting. Redosier dogwood (accessions 9023733, 9023739 and 9023740), Laurel willow and Peachleaf willow (accessions (9067375, 9067376, 9067541, 9067546, 9067549 and 9067560) cuttings were ordered February 4, 2005. Planted spring 2005. Survival and identification difficult in 2005.

**ID05011 Twin Falls County – Rock Creek Park** critical area field planting. Regar meadow brome, Topar pubescent wheatgrass, Bannock thickspike wheatgrass, Snake River Plains fourwing saltbush, Rosana western wheatgrass, pryor slender wheatgrass and Garrison creeping foxtail were ordered March 14, 2005. Site characteristics: silt loam soil, 2



percent slopes, NW aspect, 3600 feet elevation, 12 in rainfall zone, non-irrigated (riparian – sub irrigated), T10S R17E SW ¼ Section 8. Twin Falls county coordinator said only a small amount of seed was planted in 2005.

**ID06006 Twin Falls Co. Riparian Projects.** Woody field planting. Simon poplar (30), Peachleaf willow – Caribou Source (20), Peachleaf willow – Pocatello Source (20), Coyote willow (120), Golden willow (20), Laurel willow (20), White willow (20), Redosier dogwood – Harrington Source (50), Redosier Dogwood – Cheney Source (50) and Redosier dogwood – Wallowa Source (50). Cuttings ordered February 2, 2006.

## **IDAHO DIVISION V PLANT MATERIALS PLANTINGS**

### **FIELD OFFICE: AMERICAN FALLS/ABERDEEN**

**None**

### **FIELD OFFICE: BLACKFOOT**

**ID02006 Paul Ricks** Demonstration Planting. Seed ordered February 11, 2002 for shipment to Aberdeen PMC by March 4, 2002. FY02 Planting completed in May 2002. August 27, 2002 initial evaluation indicated at least some establishment of all seed plots. FY03 evaluated 12/9/03. FY04 evaluated and clipped 6/23/04. FY05 and FY06 no evaluation. See attached tables at end of this section.

### **FIELD OFFICE: FORT HALL**

**ID03002 Shoshone-Bannock Tribe** Demonstration Planting. Nezpar Indian ricegrass, Goldar bluebunch wheatgrass, Magnar basin wildrye, Sodar streambank wheatgrass, High Plains Sandberg bluegrass, and Sherman big bluegrass seed was ordered September 30, 2002. Planting completed early November 2002. FY03 no evaluation. FY04 Nezpar Indian ricegrass excellent stand and vigor with 24 inch height. Goldar bluebunch wheatgrass excellent stand and vigor with 24 inch height. Magnar basin wildrye excellent stand with good vigor and 36 inch height. Sodar streambank wheatgrass excellent stand and vigor with 30 inch height. Sherman big bluegrass good stand with good vigor and 30 inch height. Very poor stand and vigor with 8 inch height (only 3 plants came up – seed may have been buried too deep). FY05 evaluation June 15, 2005. Magnar excellent stand, 98% survival, excellent vigor and 48 inch height; Nezpar good stand, 90% survival, good vigor and 36 inch height; Goldar good stand, 95% survival, good vigor and 42 inch height; Sodar excellent stand, 98% survival, excellent vigor and 42 inch height; Sherman excellent stand, 98% survival, excellent vigor and 38 inch height; High Plains failed and will be replanted next year and irrigated for establishment. **FY06** excellent stands of Goldar bluebunch wheatgrass and Magnar basin wildrye, good stands of Sodar streambank wheatgrass, and Sherman big bluegrass, fair stand of Nezpar Indian ricegrass and High Plains Sandberg bluegrass failed.

**ID03005 ShoBan High School** field planting. Common Camas bulbs. Bulbs ordered from Corvallis PMC January 14, 2003. Site - MLRA B11b, 10-12 inch precipitation, sub-irrigated wet to semiwet bottomlands, non-irrigated. FY04 Camas bulbs were planted in the fall of 2003 and no evaluation has been completed. FY05 could not find any camas plants – recommend evaluation one additional year to ensure this planting is a failure – might recommend collecting camas bulbs from more local location for future studies and evaluations. FY06 planting failed. **Cancel**

**ID06010 Shoshone Bannock Tribe – ShoBan High School** common camas field planting. FY06 Tribal members collected bulbs from the Camas Prairie near Fairfield in Camas County. The bulbs were planted about 4 inches deep in the native garden behind the field office and also near the constructed wetland at the ShoBan Jr./Sr. High School. Plantings are being heavily watered. FO– 1 plant/ft<sup>2</sup>; School– 1 plant/hole approximately 10-15 feet apart.

### **FIELD OFFICE: MALAD**

**ID04005 Hybrid poplar study – Don Buhler** field planting. Robust poplar, Carolina poplar, Siouxland poplar, Simon poplar, OP367 poplar and 52-225 poplar cuttings were ordered March 5, 2004. Site Characteristics: Zukom silt loam soil, 7.4-8.4 soil pH, very wet site in early spring, 0-1% slopes, south aspect, 18-20 inch precipitation, non-irrigated, 5180 feet elevation. FY04 - Robust poplar 20 percent survival with fair vigor. Carolina poplar 13 percent survival with fair vigor. Siouxland poplar failed. Simon poplar 53 percent survival with good vigor. OP367 poplar 20 percent survival with poor vigor. 52-225 poplar 13 percent survival with poor vigor. FY05 and FY06 no evaluation.

### **FIELD OFFICE: MONTPELIER**

**None**

### **FIELD OFFICE: POCATELLO**

**None**

**FIELD OFFICE: PRESTON**

**ID95036 Franklin County** Bannock thickspike wheatgrass and Sodar streambank wheatgrass critical area planting. Site is landfill, Wheelon/Collonston soil, non-irrigated, 14-15 inch ppt, 5000 feet elevation, 12-20% slopes on north exposure. Seed ordered 5/5/95. FY95 seed planted 5/17/95 in good clean seedbed. Fall evaluation indicated good stand establishing for both species. FY96 good stands of both species with 3 plants/ft<sup>2</sup> and spreading. Species are providing good erosion control. FY97 and FY98 no evaluations. FY99 good stand of each specie with 3-4 plants per square foot, good vigor, good ability to spread, and good erosion control under these conditions. Weed infestation of planting is very low. FY00 Bannock and Sodar stands are good with good vigor and 4 plants per square foot. FY01-FY05 no evaluations. **Cancel**

**FIELD OFFICE: SODA SPRINGS**

**ID05001 Michael Tingey** – Irrigated forages Demonstration Plots. Latar orchardgrass, Regar meadow brome, Cache meadow brome, Paiute orchardgrass, Garrison creeping foxtail, Rush intermediate wheatgrass, Bozoisky Russian wildrye, 905439 switchgrass, Blackwell switchgrass and Lutana cicer milkvetch seed was ordered February 4, 2005. SCD/Cooperator Supplies the following: Paddock meadow brome, Forager alfalfa, Kemal festolium, Potomic orchardgrass, Rebound meadow brome, Fuego tall fescue, Tekapo orchardgrass, Mara perennial ryegrass, Barliza timothy, Pradel meadow fescue, Barloex tall fescue, Bariane tall, fescue, Barcell tall fescue, Baridana orchardgrass, Hakari Alaska brome, Birdsfoot trefoil, Sainfoin, Sorgam, Grazing corn, Lakota prairie brome and Alice white clover. Site characteristics: 0.8 acres, MLRA B13, Rexburg-Ririe silt loam soil complex, 1-4 percent slopes, north aspect, elevation 5140 feet, 12-14 inch precipitation, irrigated, T11S R41E SW ¼ section 19. Planted in late spring 2005 due to persistent rainfall that did not allow earlier final land preparation and planting. FY06 evaluated 10/27/06.

Species	Percent Stand			Vigor			Height		
	2005	2006	2007	2005	2006	2007	2005	2006	2007
Kura Clover	failed	---	---	---	---	---	---	---	---
Forager alfalfa	exc.	fair		good	good		24"	grazed	
Lutana cicer milkvetch	good	poor		good	good		6"	grazed	
Alice white clover	poor	poor		fair	fair		4-6"	grazed	
Birdsfoot trefoil	fair	fair		good	good		3"	grazed	
Eski sainfoin	good	good		good	good		12"	grazed	
Baridana orchardgrass	poor	good		good	good		16"	grazed	
Tekapo orchardgrass	fair	good		good	good		12"	grazed	
Paiute orchardgrass	poor	poor		fair	fair		12"	grazed	
Latar orchardgrass	poor	poor		fair	fair		12"	grazed	
Potomic orchardgrass	fair	fair		good	good		12"	grazed	
Satin orchardgrass	poor	poor		good	good		8"	grazed	
Renegade orchardgrass	fair	good		good	good		18"	grazed	
Rebound meadow brome	good	good		good	good		24"	grazed	
Cache meadow brome	fair	fair		good	good		30"	grazed	
Regar meadow brome	fair	fair		good	good		12"	grazed	
Lakota prairie brome	good	v.poor		exc.	v.poor		36"	grazed	
Hakari Alaska brome	exc.	good		exc.	exc.		12"	grazed	
Seine tall fescue	poor	fair		good	good		24"	grazed	
Johnstone tall fescue	poor	fair		good	good		18"	grazed	
Bronson tall fescue	fair	good		good	good		24"	grazed	
Bariane tall fescue	fair	fair		good	good		12"	grazed	
Dovy tall fescue	fair	fair		good	good		18"	grazed	
Pradel tall fescue	fair	fair		good	good		12"	grazed	
Garrison creeping foxtail	v.poor	poor		fair	fair		12"	grazed	
Rush intermediate whtgrs	fair	good		fair	fair		6"	grazed	
Bozoisky Russian wildrye	fair	fair		poor	good		4"	grazed	
Kemal festolium	exc.	exc.		exc.	exc.		24"	grazed	
Mara perennial ryegrass	exc.	exc.		good	exc.		8"	grazed	
Barliza timothy	poor	poor		poor	poor		4"	grazed	
Outlaw timothy	fair	v.poor		poor	poor		8"	grazed	

Blackwell switchgrass	v.poor	poor	fair	poor	18"	grazed
9005439(MT) switchgrass	v.poor	v.poor	fair	fair	8"	grazed
Garrison sorgum-sudan	exc.	--	good	--	54"	--

**ID05012 Don Ayers** – herbaceous windbreak field planting. Magnar basin wildrye seed ordered March 15, 2005. Site Characteristics: Lantonia-Chinahat silt loam soil, 1-4 percent slopes, 5983 feet elevation, 14-16 inch precipitation, non-irrigated, T8S R41E NW ¼ Section 24. FY05 Two of the four rows had good emergence and two rows had very poor emergence. Ground preparation was much better in rows that the best emergence. Plants that emerged have grown well and look very healthy - fair stand with 4 plants/ft<sup>2</sup>, good vigor and 4 inch height. Several more plants emerged in the fall. **FY06** planting was accidentally tilled and destroyed – cooperators plans to replant.

## **IDAHO DIVISION VI PLANT MATERIALS PLANTINGS**

### **FIELD OFFICE: ARCO**

**ID03003 Hill-Freeman** Snake River Plain fourwing saltbush field planting. Seed ordered October 18, 2002. FY03 one half pound of Snake River Plains fourwing saltbush was included in a five acre marginal pastureland seeding adjacent to Warm Springs Creek on Barton Flat (South Custer County). The entire seeding area of 13.3 acres included a three and a half acre stand of decadent crested wheatgrass. A seed mix of Vavilov Siberian wheatgrass (1.2 lbs/ac), Bannock thickspike wheatgrass (2.0 lbs/ac), Bozoisky Russian wildrye (1.2 lbs/ac), Rincon fourwing saltbush ( 0.25 lbs/ac), and Bighorn skunkbush sumac (0.25 lbs/ac) was broadcast over the seeding area. The area was then rolled to obtain seed to soil contact on a firm weed free seedbed. FY04- FY06 no evaluations.

### **FIELD OFFICE: DRIGGS**

**ID91006 Fair Grounds** Multiple Species Demo Plots. FY92 planted spring 1992 excellent survival on all species except trefoil, mountain brome and cicer milkvetch which will have to be replanted. FY93 Remont, Bromar, Lutana planted spring of 1993. Remont is not tolerant of frequent irrigation. Bozoisky exhibits poor seedling vigor, Goldar has poor plant vigor, Canbar not recommended for pure stands, Magnar not adapted to shallow soils, Newhy lacks seedling vigor, Manchar exhibits poor summer regrowth, Whitmar is not tolerant of excessive moisture, and Garrison adapted to wet soils. Magnar, Bromar, Rush, and Lutana are all doing poorly. Ordered Rush, P27, Magnar, Canbar, and Bozoisky on 3/17/94 to be included in plots. FY94 all plots good to excellent stand except Lutana, Remont and Delar. These plots are all irrigated so evaluations for drought, flood, salt and acid tolerance not possible. This planting does provide excellent trials for irrigated varieties in high mountain valleys. FY95 best performers are Hycrest, Critana, Alkar, Tegmar, Luna, Greenar, Topar, Rush, Regar, Manchar, Latar, Paiute, Sodar, Newhy, Durar, Sherman, Canby and Delar. Complete evaluations are available on request. FY96 not evaluated. FY97 Durar and Delar good to excellent stands with high vigor; Regar, Amur, Manchar, Latar, Paiute good stands with excellent vigor; Rush fair stand with fair vigor; Sodar, Goldar, Cascade, Appar poor stands with fair vigor; Hycrest, Critana, Alkar, Tegmar, Luna, Greenar, Topar, Lutana, Garrison, Whitmar, Secar, P27, Bromar, Magnar, Bozoisky, Canbar, Sherman, Kalo, very poor to failed stands. All plots are subject to turfgrass encroachment. February 9, 1998 ordered Hycrest, CD-II (Hycrest II), Sherman, Newhy, Critana, Bannock, Garrison, and Bozoisky for plots. FY98 species with good to excellent stands include Amur, Rush, Manchar, Latar, Durar, Cascade, and Delar. Species with poor to fair stands include Alkar, Luna, Topar, P27, Bromar, Paiute, Magnar, Appar, and Bozoisky. Failed stands include Hycrest, Critana, Tegmar, Greenar, Secar, Whitmar, Garrison, Lutana, Regar, Sodar, Newhy, Kalo, Sherman, Canbar, and Goldar. FY99- FY05 no evaluations.

**Cancel**

**ID99018 SCD** field planting – leafy spurge competition study. Species include Rush intermediate wheatgrass, Luna pubescent wheatgrass, Regar meadow brome, Bromar mountain brome, Durar hard fescue, Bozoisky Russian wildrye, and Climax timothy. Seed ordered April 28, 1999 for shipment about May 17, 1999. FY99 Roundup was applied on June 10<sup>th</sup> to leafy spurge plots with up to 200 stems per 9.6 square foot hoop. Grass was drilled into plots on July 1, 1999 using a Brillion drill. Evaluation of germination and establishment will be performed in the spring of 2000. Replicated plots will be installed in May of 2000. FY00- FY05 no evaluations. **Planting failed - Cancel**

**ID02019 Lowel Curtis** field planting. Species include Garrison creeping foxtail, Regar meadow brome and Johnstone tall fescue. Seed ordered April 8, 2002. FY02- FY05 no evaluations. **Never installed - Cancel**

## **FIELD OFFICE: IDAHO FALLS**

**ID94020 Winterfeld** Magnar basin wildrye and Trailhead basin wildrye vegetative terraces field planting. Seed ordered 3/94. FY94 planted 5/94. Good initial stand establishment with good vigor. FY95 excellent stand establishment with over 3 plants/ft<sup>2</sup>. Plants average 24" height. Grouse are using basin wildrye for nesting cover. Working well for erosion control. FY96 excellent stands with excellent vigor Trailhead and good vigor Magnar. Excellent wildlife use by game birds, deer, owls, and coyotes. Both species are very good for snow catchment and field windbreaks. FY97 100% survival, Trailhead spreading a little faster than Magnar. Plant height about 96 inches for each. Cooperator notes that Trailhead is more drought tolerant and Magnar is more robust. FY98 100 percent survival for both species. Cut for seed this year with 140 pounds of clean seed per acre. FY99 excellent stands: Magnar 96 inches tall with little to no spread; Trailhead 84 inches tall with good spread via seed shatter. FY00 excellent stands with excellent vigor for both Magnar and Trailhead. Magnar is more robust with 96 inches height. Trailhead is spreading rapidly, is more drought tolerant, and approximately 84 inches tall. FY01 excellent stand and vigor with 96 inch height. Seed production was approximately 100 pounds per acre. Straw yield was 1.6 tons per acre. FY02 Trailhead plowed out. Magnar excellent stand with excellent vigor, 72 inch height, and 4000 pounds per acre production. FY03 no seed crop due to insect damage. FY04 – excellent stands with excellent vigor and each accession was approximately 96 inches tall this year. Trailhead is spreading beyond original planting. **FY06 plowed out – cancel.**

**ID95046 Winterfeld** Venus penstemon and Firecracker penstemon District Seed Increase. Seed sent 8/95. FY95 planted fall 1995. FY96 poor stand establishing for Alpine and no emergence for Firecracker, no seed production. FY97 Alpine slow establisher and susceptible to frost, no seed production. FY98 fair stand of both Firecracker and Alpine penstemon (1 plant per foot 2). Stands for both species are getting better each year. FY99 fair stands in unfavorable moisture year and no seed production. FY00 Firecracker penstemon died due to drought and short-lived character. Alpine penstemon has good stand with good vigor and stands 24 inches tall. Seed production was unknown at evaluation date. FY01 firecracker penstemon came back, excellent stands and vigor for both species. Seed production estimated at 600 pound per acre bulk. FY02 - Venus - fair stand with excellent vigor, 24 inch height, and 100 pounds per acre bulk production. Firecracker - fair stand with excellent vigor, but slower establishment, 24 inch height, and 100 pounds per acre bulk production. FY03 Firecracker penstemon stand is going out – no production. Venus penstemon produced 80 pounds of seed. FY04 – excellent stand and vigor for each accession. No seed production reported. FY05 Venus penstemon – good stand and good vigor – no seed production reported. Firecracker penstemon stand was plowed out. FY06 excellent stand and vigor.

**ID03007 Winterfeld** San Juan penstemon - Demonstration planting. Seed ordered February 10, 2003. Seed shipped February 18, 2003. FY03 not planted. FY04 planted with poor stand establishing. FY05 replanted October 26, 2005. FY06 fair stand establishing with 2 plants/ft<sup>2</sup>.

**ID04015 Winterfeld** Maple Grove Lewis flax for seed increase. Seed shipped April 19, 2004. FY04 excellent stand with excellent vigor establishing. Plants are about 5 inches tall. FY05 good stand, good vigor and plants are about 24 inches tall. Lighter in color than Appar and not as good a competitor with severe weed competition in stand. No seed production reported. FY06 good stand with good vigor – seed production was not reported.

**ID05015 Winterfeld** Pryor slender wheatgrass for seed increase. Seed shipped May 2, 2005. FY05 no evaluation. FY06 excellent stand with excellent vigor - seed production not reported.

**ID07003 Winterfeld** Appar blue flax for seed increase. Seed shipped September 26, 2006.

**ID07004 Winterfeld** Sodar streambank wheatgrass for Foundation seed increase.

## **FIELD OFFICE: REXBURG**

**ID89015 Wagoner** Luna pubescent wheatgrass, P-27 Siberian wheatgrass, Sodar streambank wheatgrass, Greenar intermediate wheatgrass, Delar small burnet, Trevois alfalfa field planting on rangeland. Site is gravelly loam soil with a pan at 5-6 inches, non-irrigated, 12-inch ppt, 6300 feet elevation, and 3% slopes on NE exposure. FY89 ripped rangeland in spring and seeded mix in fall of 1990. FY91 excellent stand establishing with production about 1400 lbs/ac. FY92 clipping data: No Treatment - 318 lbs/ac., chisel only treatment (native species) - 495 lbs/ac., chisel/disc/seed treatment - 1110 lbs/ac. Clipped 7/9/92. FY93 Clipped plots resulted in production of 1200-2000 lbs/ac. FY94 production of about 800 lbs/ac in extremely droughty year. Non treated rangeland producing about 100 lbs/ac this year. FY95 excellent stand Luna and Greenar, Good stand P-27, Sodar and Travois and Poor stand of Delar.

Stand produced 1400+ lbs/acre this year. High antelope use of stand was noted. Stand was grazed 3 weeks in spring and 4 weeks in fall with good management. FY96 excellent stand of Trevois and good stands of Luna, P27, Sodar, and Greenar. Very poor stand of Delar. Considered 90% stand overall. Produced 1000 lbs/ac in very poor moisture year. Stand is doing great under good management. FY03 Disc-Seed treatment – near fence good stand of natives – primarily crested wheatgrass in seeding with 5-6 percent sagebrush and 600 pounds per acre production in very dry year. Ripped-Disc-No Seed treatment – sagebrush very heavy with forage producing about 200 pounds per acre and brush producing about 200 pounds per acre in very dry year. Ripped-Disc-Seed treatment – excellent stand of primarily Bozoisky wildrye, Nordan crested wheatgrass, P27 Siberian wheatgrass and some Trevois alfalfa. Very little intermediate wheatgrass left in stand. Production is about 1000 pounds per acre in very dry year. FY05 There is a good stand of native bluebunch wheatgrass, Sandberg bluegrass and Indian ricegrass near west fence-line producing about 750 pounds per acre. The disced and seeded stand near west fence has a good stand of crested wheatgrass with about 5 percent sagebrush invasion and producing about 1000 pounds per acre. The ripped, disced and seeded area has an excellent stand of primarily Nordan crested wheatgrass and Bozoisky Russian wildrye with 3-4 plants per square foot, excellent vigor and producing about 1300 pounds per acre this year. P27 Siberian wheatgrass, greenar intermediate wheatgrass and Trevois alfalfa are present, but in much lower amounts. **Next evaluation 2008.**

**ID90025 Wagoner** Rush intermediate wheatgrass field planting on rangeland. Site is gravelly loam soil with a pan at 5-6 inches, non-irrigated, 12-inch ppt, 6300 feet elevation, and 3% slopes on NE exposure. FY89 ripped rangeland. FY90 planted April 1990. FY91 excellent stand establishing with no weeds. Production is 1400 lbs/ac. FY92 stand excellent with 1200 lbs/ac production. FY93 excellent stand producing 2000+ lbs/ac. Grazing value - appears to be a highly preferred/selected species according to cooperator. FY94 excellent stand producing 800 lbs/ac in very droughty year. FY95 excellent stand producing 1800+ lbs/acre. Rush is the most productive species in all range trials. FY96 excellent stand with 5-10 plants/ft<sup>2</sup> producing 1000-lbs/ac and good vigor in very low rainfall year. FY03 good to excellent stand with 3 plants per square foot and good to excellent vigor. Producing 700 pounds per acre in very dry year – produces about 1400 pounds per acre in average to favorable years. Sagebrush invasion is about 1-5 percent of plant community. No weeds in stand. **Next evaluation 2008.**

**ID90035 Wagoner** Bozoisky Russian wildrye field planting on rangeland. Site is gravelly loam soil, non-irrigated, 12-inch ppt, 6200 feet elevation, and 2% slopes on NE exposure. FY90 planted April. FY91 good stand establishing. FY92 excellent stand producing 1100 lbs/ac. FY93 90% + stand and up to 4' tall, estimated production 1200-1400 lbs/ac. FY94 good stand producing about 600 lbs/ac in very droughty year and only 50% of plants produced seedheads this year. FY95 good stand producing 1200+ lbs/acre. This species is doing very well and is well adapted to site. FY96 good stand with 4-5 plants/ft<sup>2</sup> and 1200-lbs/ac production in very low summer rainfall year. FY03 good stand of P27 Siberian wheatgrass and Bozoisky Russian wildrye with 3 plants per square foot and good to excellent vigor. Stand is producing about 800 pounds per acre in a very dry year. Estimate 1400-1600 pounds per acre in an average to favorable moisture year. FY05 the Bozoisky Russian wildrye stand is maintaining very well with approximately 3 plants per square foot, excellent vigor and production about 1200 pounds per acre. Cattle seek out this species year around according to cooperator. **Next evaluation 2008.**

**ID92013 Webster** Regar meadow brome, Bozoisky Russian wildrye, Luna pubescent wheatgrass, Critana thickspike wheatgrass field planting on rangeland. Site is gravelly silt loam soil, non-irrigated, 14-inch ppt, 6000 feet elevation, and 4% slopes on SE exposure. FY92 site sprayed for weed control, but too dry to seed. FY93 seeding not completed. FY94 very poor moisture conditions, planting not installed. FY95 good stand of all species establishing with good spring moisture. FY96 good stand of all species with 2-4 plants/ft<sup>2</sup> and good vigor on all except Regar has fair vigor. Stand had low production and is still establishing. FY97 good stands for all species with 60% stands and good vigor - they have been slow to establish on this tough site. FY99 Bozoisky and Luna good stands, Regar and Critana fair stands. FY03 good to excellent stand of Bozoisky Russian wildrye and Regar meadow brome with 3 plants per square foot (70% Bozoisky – 30% Regar), good vigor and about 1500 pounds per acre production in a very dry year. Good to excellent stand of Bozoisky Russian wildrye and Trevois alfalfa with 3 plants per square foot (70% Bozoisky – 30% Trevois), good vigor and about 1500 pounds per acre production in a very dry year. Fair to good stand of Critana thickspike wheatgrass with 9 plants per square foot, poor vigor and about 400 pounds per acre production in a very dry year. Good to excellent stand of Luna pubescent wheatgrass with 5 plants per square foot, good vigor and about 1500 pounds per acre production in a very dry year. Bozoisky is heavily grazed (80-90 percent utilization) by cattle and elk and stands are maintaining very well. FY05 Plot 1: good stand with 2 plants per square foot - Bozoisky Russian wildrye 100% survival, Regar meadow brome failed, Trevois alfalfa 50% survival; stand producing about 1300 pounds per acre. Cattle and elk are utilizing the stand at about 60 percent utilization on Bozoisky and 30 percent utilization on

alfalfa. Plot 2: excellent stand with 3 plants per square foot – Bozoisky 100 percent survival and Trevois 50 percent survival; stand is producing about 1700 pounds per acre; Cattle and elk are utilizing stand with about 85 percent utilization on Bozoisky and 30 percent utilization on alfalfa. Plot 3; fair stand of Critana thickspike wheatgrass with 9 plants per square foot and fair vigor; stand is producing about 700 pounds per acre. Cattle and elk are not utilizing this plot. Plot 4: good stand of Luna pubescent wheatgrass with good vigor and 5 plant per square foot; stand is producing about 1700 pounds per acre; Cattle and elk are not utilizing this stand. **Next evaluation 2008.**

#### **FIELD OFFICE: RIGBY/TERRETON**

**ID96019a Mud Lake** Willows and cottonwood demo planting Laurel, Coyote, White, Robusta poplar, Siouxland poplar, and Carolina poplar. Cuttings ordered 2/20/96. Planted May 8, 1996 using fabric mulch material and drip irrigation. FY96 Water application, started July 5th with willows receiving 7 gallons/week and poplars receiving 12 gallons/week. Flood irrigation by Park officials resulted in over-irrigation and drip system was cut back. 100% survival of all species except coyote which had 70% survival. Good vigor for all species except Carolina poplar which had fair vigor. Growth: Carolina 3.2 feet; Siouxland 5.7 feet; Robust 5.5 feet; Laurel 2.7 feet; White 3.7 feet; Coyote 4.0 feet. FY97 Irrigation: 3 gallons/tree from May through September. Survival/Vigor/Height: Carolina poplar 75%/good/10.5 feet; Siouxland poplar 100%/excellent/14 feet; Robust poplar 100%/fair/7 feet; Laurel willow 100%/excellent/7.5 feet; White willow 100%/excellent/9 feet; Coyote willow 67%/fair/ 4.5 feet. FY98 Survival/ Vigor/Height: Carolina poplar 75%/good/15 feet; Siouxland poplar 100%/excellent/ 20 feet; Robust poplar 100%/fair/12 feet; Laurel willow 100%/excellent/10.5 feet; White willow 100%/good/14 feet; Coyote willow 70%/good/6.5 feet. FY99 Carolina poplar 75% survival with good vigor and 21.2 feet height. Siouxland poplar 100% survival with excellent vigor and 26.4 feet height. Robust poplar 100% survival with poor vigor (yellow leaves) and 16.6 feet height – seedlings are vigorous with good color and suspect Aberdeen stock may have disease. Laurel willow 100% survival with good vigor and 12.4 feet height. White willow 100% survival with good vigor and 18.5 feet height. Coyote willow 70% survival with fair vigor and 6.9 feet height. FY00 Flood irrigated every two weeks with drip irrigation 6-10 gal/week. Carolina poplar 75 percent survival with excellent vigor and 320 inch height. Siouxland poplar 100 percent survival with excellent vigor and 354 inch height. Robust poplar 100 percent survival with poor vigor (disease) and 216 inch height. Laurel willow 100 percent survival with excellent vigor and 180 inch height. White willow 100 percent survival with fair vigor and 240 inch height. Coyote willow 66 percent survival with fair vigor and 90 inch height. FY01 6-year-old planting was flood irrigated every two week this year. Carolina poplar (10-15 feet spacing recommended) - 75% survival, excellent vigor, 36 feet height, 16 feet crown width, and 5.5 inch DBH. Siouxland poplar (10-15 feet spacing recommended) – 100% survival, excellent vigor, 38 feet height, 15 feet crown width, and 5 inch DBH. Robust poplar (10-15 feet spacing recommended) – 100% survival, poor vigor, 25 feet height, 9 feet crown width, and 3.5 inch DBH. Laurel willow (8-10 feet spacing recommended) – 100% survival, good vigor, 17 feet height, 12.5 feet crown width, and 2 inch DBH. White willow (10-12 feet spacing recommended) – 100% survival, fair vigor, 20 feet height, 12 feet crown width, and 2 inch DBH. Coyote willow (3-5 feet spacing recommended) – 70% survival, fair vigor, 8 feet height, and 3 feet crown width. FY02 Carolina poplar 75% survival, excellent vigor, 439 inch height, and 5.75 dbh. Siouxland poplar 100% survival, excellent vigor, 455 inch height, and 17.5 inch dbh. Robusta poplar 100% survival, fair vigor, 319 inch height, and 4 inch dbh. Laurel willow 100% survival, good vigor, 211 inch height, and 2.25 dbh. White willow 100% survival, good vigor, 235 inch height, and 2.25 inch dbh. Coyote willow 66% survival fair vigor, and 139 inch height. FY03 100 percent survival of Carolina poplar (good vigor – 40 feet height), Souixland poplar (good vigor – 44 feet height), Robust poplar (fair-good vigor – 25-25 feet height), Laurel willow (good vigor – 22 feet height – lower limbs dieing), and White willow (excellent vigor – 16 feet height – good density). 50 percent survival of Coyote willow (fair-good vigor – 21 feet height). Souixland best choice of poplars – White willow best choice of willows. **Next evaluation 2007.**

**ID96019b Rigby** Cottonwood demo planting - Carolina, Siouxland, Robusta. Planted April 29th using fabric mulch and drip irrigation. FY96 Water application 10-14 gallons per week. Growth Carolina 2.0 feet; Siouxland 3.2 feet; Robust 4.0 feet. FY97 100% survival for all poplars. Good vigor for Carolina and Siouxland / poor vigor for Robust. Height 8-9 feet Carolina and Siouxland / 3 feet Robust. FY98 Survival/Vigor/Height: Carolina poplar 100%/good/15 feet; Siouxland poplar 100%/ excellent/18 feet; and Robust poplar 100%/poor/5.5 feet. FY99 Carolina poplar 100% survival with fair vigor and 21 feet height. Siouxland poplar 100% survival with fair vigor and 21 feet height. Robust poplar 100% survival with very poor vigor and 7 feet height. Note – Robust poplars from Lawyers Nursery are thriving, so suspect Aberdeen cuttings may be carrying a disease. FY00 Drip irrigated (14 gal/week) – Carolina poplar 100 percent survival with fair vigor and 240 inch height; Siouxland poplar 100 percent survival with fair vigor and 252 inch height; Robust poplar 100 percent survival with poor vigor and 84 inch height. FY01 6-year-old planting is irrigated with drip irrigation system at 7 gallons per week. Carolina poplar – 100% survival, poor vigor 22 feet height, 7 feet crown width, and 2.5 inch DBH. Siouxland poplar – 100% survival, poor vigor, 24 feet height, 6 feet crown



width, and 3 inch DBH. Robust poplar – 100% survival, very poor vigor, 7 feet height, 4 feet crown width, and 1 inch DBH. Drought stress is evident and drip irrigation system is probably not fully functioning with plugged emitters, need for additional emitters, and need for longer watering sets. FY02 Carolina poplar 100% survival, very poor vigor, 300 inch height, and 2.5 inch dbh. Siouxland poplar 100% survival, fair vigor, 330 inch height, and 2.75 dbh. Robusta poplar 100% survival, very poor vigor, 92 inch height, and 1 inch dbh. Irrigation system problems were repaired and irrigation sets have been extended - expect improvement next year. FY03 100 percent survival of Carolina poplar (fair to good vigor – 10 feet height – some winter die back), Souixland poplar (good vigor – 28 feet height) and Robusta poplar (very poor vigor – 8 feet height). Best choice Souixland poplar. **Next evaluation 2007.**

**ID98013 Jefferson County Landfill** Field planting 1) Ephraim crested wheatgrass, Sodar streambank wheatgrass, and Bannock thickspike wheatgrass; 2) Covar sheep fescue, Schwendimar thickspike wheatgrass, and Secar Snake River wheatgrass. Seed ordered Feb 9, 1998. Site is silty clay loam soil, 0-1 % slope, east aspect, 4785 feet elevation, 10-12 inch ppt, non-irrigated, T6N R33E SE1/4 Section 14. FY98 initial evaluation showed very poor to no establishment of Covar, Schwendimar, Secar, Sodar, and poor to very poor establishment of Ephraim and Bannock. The clay soil portions of the seeding crusted and the sandy soil portion of the seeding may have been too dry. Site should be evaluated one more season before a decision to reseed is made. FY99 Covar – fair stand with poor vigor and .2 plants per square foot. Schwendimar – very poor stand with poor vigor and .1 plants per square foot. Secar – very poor stand with poor vigor and .1 plant per square foot. Bannock fair stand with poor vigor and 1 plant per square foot. Sodar – poor stand with poor vigor and .1 plants per square foot. Ephraim – fair stand with fair vigor and 1 plant per square foot. FY00 Planting Mix 1 – fair stand of Ephraim/Sodar/Bannock is establishing with fair vigor and stand is limiting weed growth. Planting Mix 2 – poor stand of Covar/Swendimar/Secar is establishing with fair vigor. Secar and Schwendimar failed in planting for the most part, but Covar is establishing slowly. Stand is dominated by kochia weed. Planting 3 – Bannock has good stand with fair vigor. Windbreak planting (drip irrigated) is irrigated once per week for 12-16 hours, is doing very well, and trees are uniform – Russian Olive 5-8 feet height with 5 feet crown width; Rocky Mountain Juniper 3-5 feet height with 3 feet crown width; Siberian Peashrub 4-7 feet height with 4 feet crown width. FY01 the Ephraim-Bannock-Sodar mix and Bannock only plantings are increasing and spreading. Covar in the Covar-Swendimar-Secar mix is also increasing. Grass densities of 2+ plants per foot squared occur on more favorable sandy soils. The hard packed clayey areas have few grass seedlings established. The windbreak planting is doing very well with 100% survival and very good maintenance for water (drip irrigation system) and weed control. Russian olive is averaging 9 feet tall and 7 feet crowns on sandier soils and 5-6 feet tall with 5 feet crowns on clayey hard packed soils. Junipers and Siberian peashrub are not affected as much by varied soil conditions with Junipers averaging 5 feet tall with 4 feet crowns on sandy soils and 4.5 feet tall with 4 feet crowns on clayey soils. The Siberian peashrub is averaging 6 feet tall with 5 feet crowns on sandy soils and 5.5 feet tall with 5 feet crowns on clayey soils. FY02 grass planting are doing very well and spreading with over 3 plants per square foot. FY03 planting is doing well. **FY06** mix 1- Covar, Schwendimar, Secar - fair stand with 1-2 plants/ft2 and poor vigor; mix 2- Ephraim, Sodar, Bannock – good stand with 2-3 plants/ft2 and fair vigor – Ephraim stands out as the best performing grass at this site; mix 3- Bannock – fair stand with 1-2 plants/ft2 and fair vigor. Windbreak evaluation – excellent survivability considering site and lack of irrigation. Rocky Mountain juniper – 6-7 feet tall with excellent survival; Russian olive – 6-9 feet with excellent survival; Siberian peashrub – 5-8 feet tall and drought stressed in portions of row (leaves curling). Recommend this evaluation be **cancelled**.

**ID98014 Calvin Moser** Rush intermediate wheatgrass pasture trial. Seed ordered 2/9/98. Site is sandy loam soil, 0-2 % slope, west aspect, 4795 feet elevation, 10-12 inch ppt, irrigated, T4N R38E SE1/4 Section 29. FY98 two acres of Rush were seeded at the end of March with oats as a cover crop (15 lbs/acre oats). The oats were harvested in mid-September and the Rush is responding with average of one foot tall and 2 plants/ft2 at the end of October. FY99 Rush - excellent stand with excellent vigor, 9000 pounds per acre production, 4 to 6 feet height, and 3+ plants per square foot. Regar – not planted. FY00 good stand with fair vigor and 5400 pounds production. Production lower due to heat and severe drought conditions. FY01 good stand with 3 plants per square feet and good vigor. Stand produced about 4000 pounds per acre this year with two flood irrigation applications. Stand probably would have produced more if cooperater had fertilized planting. FY02 good stand with good vigor - planting produced about 2 tons per acre. **FY06** excellent stand of Rush with 4-5 plants/ft2 and excellent vigor. Plant height is 40- 60 inches and production is 2.5- 3.0 tons/acre. Stand is now 8 years old and is thick and healthy.

**FIELD OFFICE: SALMON/CHALLIS**

**ID80100 IDL Bradbury Flat** Multiple Adaptation Evaluation. Planted March 25, 1980. Evaluations 8/7/84, 8/6/86, 7/12/89, 7/7/92, 11/14/95, and 9/99. FY03 evaluated May 21, 2003 by Dan Ogle and Mark Olson - **Next evaluation FY07.**

Accession	Stand	Plants/ft2	Vigor	Comments
B1574 crested wheatgrass	70%	1.0	good-exc.	
P27 Siberian wheatgrass	65%	0.5	good	
Sodar streambank wheatgrass	65%	1.5	good	
AB447 crested wheatgrass	60%	0.5	good	
Secar Snake River wheatgrass	60%	0.25	fair-good	high residue problems
Hatch winterfat	50%	0.5	good-exc.	
AB764 winterfat	50%	0.5	good-exc.	
AB922 fourwing saltbush	1%	< 0.1	fair-good	
AB942 fourwing saltbush	1%	< 0.1	fair-good	

Nezpar Indian ricegrass, Luna pubescent wheatgrass, Goldar bluebunch wheatgrass, Magnar basin wildrye, Topar pubescent wheatgrass, Appar blue flax, NM1143 Firecracker penstemon, Bandera R.M. penstemon, Cedar Palmer penstemon, NM1123 Venus penstemon, AB555 aster, R885a black-eyed susan, Delar small burnet, Immigrant forage kochia, Ladac alfalfa, buckwheat species, and arrowleaf balsamroot failed.

**ID80101 IDL Bradbury Flat** Multiple Adaptation Evaluation. Planted November 7, 1981. Evaluations 8/7/84, 8/6/86, 7/12/89, 7/7/92, 11/14/95, and 9/99. FY03 evaluated May 21, 2003 by Dan Ogle and Mark Olson - **Next evaluation FY07.**

Accession	Stand	Plants/ft2	Vigor	Comments
B1574 crested wheatgrass	50%	0.5	good	
P27 Siberian wheatgrass	60%	0.75	excellent	
Sodar streambank wheatgrass	80%	1.25	excellent	
AB447 crested wheatgrass	65%	0.5	good-exc.	
Secar Snake River wheatgrass	50%	0.25	good-exc.	High residue problems
AB764 winterfat	20%	0.15	poor	
AB585 winterfat	1%	<0.1	very poor	
AB922 fourwing saltbush	3%	0.1	very poor	
AB942 fourwing saltbush	2%	<0.1	very poor	
Immigrant forage kochia	3%	0.1	fair-good	
Bozoisky Russian wildrye	70%	0.5	excellent	
Vinall Russian wildrye	70%	0.7	excellent	

Nezpar Indian ricegrass, Luna pubescent wheatgrass, Goldar bluebunch wheatgrass, Magnar basin wildrye, Topar pubescent wheatgrass, Appar blue flax, NM1143 firecracker penstemon, Bandera R.M. penstemon, Cedar Palmer penstemon, NM1123 Venus penstemon, Delar small burnet, Lodorm green needlegrass, Blair smooth brome, and Paiute orchardgrass failed

**ID82101 BLM Hole In Rock** Multiple Adaptation Evaluation. Planted late October 1982. Evaluations 8/7/84, 7/28/86, 7/13/89, 7/7/92, 9/95 and 9/99. **Access to site is very difficult and future evaluations will be cancelled - maintain file for reference.**

**ID83100 FS Nip & Tuck** Multiple Adaptation Evaluation. Evaluations 7/6/92. 9/95 and 7/02. Site has deteriorated to point future evaluations would provide little future value. **Cancel future evaluations, but maintain file for reference.**

**ID82102 BLM Centennial** Multiple Adaptation Evaluation. Planted late October 1982. Evaluations 8/7/84, 7/28/86, 7/13/89, 6/26/92, 6/20/95. FY99 not evaluated. FY03 evaluated May 21, 2003 by Dan Ogle and Mark Olson - **Next evaluation FY07.**

Accession	Stand	Plants/ft2	Vigor	Comments
GP52 alfalfa	10%	0.1	fair-good	
BC79 alfalfa	3%	0.05	fair	
RS1 wheatgrass cross	25%	0.5	good	
RS2 wheatgrass cross	15%	0.25	fair	
Newhy hybrid wheatgrass	75%	1.0	good	

Scarlet globemallow	1%	<0.1	fair-good
Ephraim crested wheatgrass	85%	1.25	fair-good
Barton western wheatgrass	5%	0.25	poor-fair
Topar pubescent wheatgrass	1%	<0.1	very poor
Whitmar beardless wheatgrass	25%	0.25	fair-good
Goldar bluebunch wheatgrass	25%	0.5	fair-good
Secar Snake River wheatgrass	50%	0.75	fair-good
Vinall Russian wildrye	60%	0.75	good-exc.
Bozoisky Russian wildrye	45%	0.25	excellent
U7881 alfalfa	1%	<0.1	very poor
Nordan crested wheatgrass	70%	0.75	good

Lutana cicer milkvetch, Canbar Canby bluegrass, Immigrant forage kochia, Bandera R.M. penstemon, Cedar Palmer penstemon, Appar blue flax, Paiute orchardgrass, P27 Siberian wheatgrass, Nezpar Indian ricegrass, Magnar basin wildrye, and yellow sweetclover failed

**ID82103 BLM Spud Alluvial** Multiple Adaptation Evaluation. Planted late October 1982. Evaluations 8/7/84, 7/28/86, 7/13/89, 6/25/92, 11/14/95 and 9/99. FY03 evaluated May 20, 2003 by Dan Ogle and Mark Olson - **Next evaluation FY07.**

Accession	Stand	Plants/ft2	Vigor	Comments
RS1 wheatgrass cross	85%	1.5	fair	
RS2 wheatgrass cross	85%	1.5	fair	
Fairway crested wheatgrass	85%	1.5	fair	
Immigrant forage kochia	50%	2.0	excellent	many young plants
Ephraim crested wheatgrass	75%	1.0	good	
Barton western wheatgrass	<5%	0.1	poor	
Whitmar beardless wheatgrass	70%	1.0	fair	
P27 Siberian wheatgrass	90%	1.5	good	
Goldar bluebunch wheatgrass	30%	0.3	poor	
Secar Snake River wheatgrass	80%	0.75	fair-good	
Vinall Russian wildrye	70%	1.0	good-exc.	
Bozoisky Russian wildrye	85%	0.75	excellent	

BC79 Synthetic alfalfa, GP52 Synthetic alfalfa, scarlet globemallow, Cedar Palmer penstemon, Appar blue flax, Paiute orchardgrass, Topar pubescent wheatgrass, Nezpar Indian ricegrass, Magnar basin wildrye, and yellow sweetclover failed.

**ID82104 BLM Jeff's Flat** Multiple Adaptation Evaluation. Planted late October 1982. Evaluations 8/7/84, 7/28/86, 7/13/89, 6/26/92. 1995 no evaluation, and 9/99. FY03 evaluated May 19, 2003 by Dan Ogle and Mark Olson - **Next evaluation FY07.**

Accession	Stand	Plants/ft2	Vigor	Comments
GP52 Synthetic alfalfa	1-5%	<0.25	fair	
BC79 Synthetic alfalfa	1-5%	<0.25	fair	
Manchar smooth brome	50%	4	good	
Baylor smooth brome	50%	4	good	
Durar hard fescue	75%	3	good-exc.	
Covar sheep fescue	45%	2	good	
Nordan crested wheatgrass	25%	0.5	fair-good	
P27 Siberian wheatgrass	40%	0.75	good	
Greenar intermediate wheatgrass	65%	4	excellent	
Magnar basin wildrye	5%	0.1	fair	
Vinall Russian wildrye	3%	0.1	poor	
Bozoisky Russian wildrye	5%	0.1	fair	

RS1 wheatgrass cross, RS2 wheatgrass cross, Hycrest crested wheatgrass, Delar small burnet, Lutana cicer milkvetch, Cedar Palmer penstemon, Appar blue flax, Paiute orchardgrass, Sherman big bluegrass, yellow sweetclover failed.

**ID82105 BLM Round Valley Multiple Adaptation Evaluation.** Planted late October 1982. Evaluations 8/7/84, 8/6/86, 7/12/89, 6/25/92, 11/13/95 and 9/99. FY03 evaluated May 19, 2003 by Dan Ogle and Mark Olson - **Next evaluation FY07.**

Accession	Stand	Plants/ft2	Vigor	Comments
RS1 wheatgrass cross	1%	<0.1	fair	
RS2 wheatgrass cross	1%	<0.1	fair	
Immigrant forage kochia	2%	<0.1	fair-good	
Scarlet globemallow	1%	<0.1	fair	
Nordan crested wheatgrass	70%	1.0	good	
P27 Siberian wheatgrass	70%	1.0	good-exc.	
Vinall Russian wildrye	30%	0.5	good	
Bozoisky Russian wildrye	75%	1.5	excellent	
Nordan crested wheatgrass	60%	1.0	fair-good	

GP52 synthetic alfalfa, BC79 synthetic alfalfa, Critana thickspike wheatgrass, Bandera R.M. penstemon, Cedar Palmer penstemon, Appar blue flax, Paiute orchardgrass, Goldar bluebunch wheatgrass, Secar Snake River wheatgrass, Barton western wheatgrass, Topar pubescent wheatgrass, Whitmar beardless wildrye, Nezpar Indian ricegrass, Magnar basin wildrye, yellow sweetclover failed.

**ID82106 BLM Gooseberry/Sheep Creek Multiple Adaptation Evaluation.** Evaluations 7/7/92. FY03 evaluated May 19, 2003 by Dan Ogle and Mark Olson - **Next evaluation FY07.**

Accession	Stand	Plants/ft2	Vigor	Comments
Nordan crested wheatgrass	5%	0.1	fair-good	
Bozoisky Russian wildrye	10%	0.2	poor-fair	
Vinall Russian wildrye	10%	0.3	fair	
Sherman big bluegrass	95%	1.5	fair-good	
Greenar intermediate wheatgrass	2%	<0.1	very poor	
P27 Siberian wheatgrass	1%	<0.1	very poor	
Ephraim crested wheatgrass	3%	<0.1	poor	
Durar hard fescue	85%	2	good	
Covar sheep fescue	80%	2	fair-good	
Manchar smooth brome	50%	0.5	fair	
Baylor smooth brome	20%	0.25	fair	
Fairway crested wheatgrass	5%	0.1	fair	

Magnar basin wildrye, Appar blue flax, Paiute orchardgrass, Cedar Palmer penstemon, Bandera R.M. penstemon, Lutana cicer milkvetch, Delar small burnet, RS2 wheatgrass cross, RS1 wheatgrass cross, BC79 synthetic alfalfa, and GP52 synthetic alfalfa failed.

#### **FIELD OFFICE: ST. ANTHONY**

**ID06009 John Taft – Henrys Lake WRP.** Field planting of 100 Engelmann spruce and 100 quaking aspen. Planting completed June 20 and 21, 2006 at 7 separate locations. One gallon potted plants; Engelmann spruce - 3 feet tall and quaking aspen (4- 5 feet tall); were planted using a skid steer with mounted 10 inch auger to dig holes. Each tree was planted by hand, pressed in by foot pressure and watered with bucket following planting. Hydrology – soil moisture varied from saturated locations near ponds at south end of project to field capacity at ponds at north end of project.