



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

UCRL-JRNL-155811

# Laboratory Observation of Secondary Shock Formation Ahead of a Strongly Radiative Blast Wave

*J. F. Hansen, M. J. Edwards, D. Froula, G.  
Gregori, A. Edens, and T. Ditmire*

**December 10, 2003**

Physical Review Letters

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

# Laboratory observation of secondary shock formation ahead of a strongly radiative blast wave

J.F. Hansen, M.J. Edwards, and D.H. Froula

*Lawrence Livermore National Laboratory, Livermore, CA 94550, USA*

G. Gregori

*LRC - Rutherford Appleton Laboratory, Chilton, Didcot OX11 0QX, UK*

A. Edens and T. Ditmire

*University of Texas at Austin, Austin, TX 78712, USA*

## Abstract

High Mach number blast waves were created by focusing a laser pulse on a solid pin, surrounded by nitrogen or xenon gas. In xenon, the initial shock is strongly radiative, sending out a supersonic radiative heat wave far ahead of itself. The shock propagates into the heated gas, diminishing in strength as it goes. The radiative heat wave also slows, and when its Mach number drops to two with respect to the downstream plasma, the heat wave drives a second shock ahead of itself to satisfy mass and momentum conservation in the heat wave reference frame; the heat wave becomes subsonic behind the second shock. For some time both shocks are observed simultaneously. Eventually the initial shock diminishes in strength so much that it can no longer be observed, but the second shock continues to propagate long after this time. This sequence of events is a new phenomenon that has not previously been discussed in literature. Numerical simulation clarifies the origin of the second shock, and its position is consistent with an analytical estimate.

We have conducted experiments comparing the shock expansion in blast waves in which radiative effects are observed to be very different, and we report here on our findings, including a new phenomenon that has not previously been observed. The motivation behind this experiment and many other experiments in laboratories around the world [1–9] is an interest in astrophysical shocks which have high Mach numbers and which may be radiative [10]. Astrophysical shocks originating in supernova (SN) explosions [3, 11–14] are the most studied, but other astrophysical phenomena (e.g., T Tauri stars [15] and stellar winds [16]) also generate their own shocks. Interstellar space consists of a tenuous plasma capable of propagating shocks over great distances, so shocks are important to understand as they mix up interstellar matter and thus affect mass-loading, stellar formation [17–19] and the history of the Milky Way and other galaxies.

The radiative nature of a shock, coupled with the optical opacity of its surroundings, largely determines the evolution of the shock and its rate of expansion. A SN shock expanding through interstellar space loses energy through radiation, although some energy may be recovered as the shock sweeps up interstellar material. The energy loss rate for the shock can be quantified [20] by the parameter  $\varepsilon = -\frac{dE}{dt} (2\pi\rho_0)^{-1} r_s^{-2} \left(\frac{dr_s}{dt}\right)^{-3}$ , where  $E$  is the total energy content of the shock,  $\rho_0$  is the density of the ambient interstellar gas,  $r_s$  is shock radius, and  $t$  is time. In a fully radiative case, in which radiation escapes to infinity, the incoming kinetic energy swept up by a shock is entirely radiated away and the shocked material collapses to a thin shell directly behind the shock. The denominator is then precisely the rate at which kinetic energy is accumulated, and  $\varepsilon = 1$ . For an adiabatic case  $\varepsilon = 0$  and once the shock has swept up more mass than what was initially present, the shock could be regarded as without characteristic length or time scales, and one would expect the well-known self-similar motion of a Sedov-Taylor blast wave [21–24],  $r_s \propto t^\alpha$ , where the exponent  $\alpha = \frac{2}{5}$ . (This assumes that there are no density gradients in the swept region of space.) In a case where radiation removes energy from the shock in an optically thin environment, analytical and numerical studies predict a slower shock expansion, such as  $\alpha = \frac{2}{7}$  (the "pressure-driven snowplow"),  $\alpha = \frac{1}{4}$  (the "momentum-driven snowplow"; the shock is simply coasting) [10, 25], and  $\frac{2}{7} < \alpha < \frac{2}{5}$  (the thermal energy of the shocked gas is not completely radiated away) [20, 24].

In a case where the environment is not optically thin, which is the case for many experiments including ours, radiation is reabsorbed in the upstream material and if the shock is

travelling fast enough a supersonic, radiative heat wave (RHW) breaks away from the shock in a situation analogous to a supercritical shock wave [23]. It has been shown that the shock and RHW will coexist and eventually propagate as  $r \propto t^\alpha$  where  $\alpha$  is larger for the shock [26]. This means that the shock would eventually catch RHW, after which a second state is obtained in which RHW is of the ablative type and the shock moves in a classical Sedov-Taylor trajectory with  $\alpha = \frac{2}{5}$ . In this manuscript we report on the additional possibility that prior to the shock catching RHW, the latter enters a transonic regime, stalls, and generates a second shock. Although our motivation for this experiment was an interest in astrophysical shocks, we should point out that the new phenomenon we describe has not been observed in astrophysical shocks and may not occur there.

We create spherically expanding blast waves in the following fashion: a high-power infrared pulsed laser (1064 nm wavelength) is focused onto the tip of a solid (stainless steel) pin surrounded by an ambient gas (nitrogen or xenon) typically at a pressure of about 1 kPa (density  $\rho_0 \sim 10^{-5}$  g/cm<sup>3</sup>). The laser pulse is 5 ns in duration with energy ranging from 10 J to 200 J (we see no qualitatively different behavior in the blastwave evolution over this energy range). The laser pulse ablates the pin and rapid expansion of ablated material shocks the ambient gas. The initial shock travels radially outward from the pin; most of the shocked gas is concentrated in a shell immediately behind the shock front. We use the term blast wave to refer to shock and shell. The blast wave velocity drops as more and more of the ambient gas is accumulated and set in motion by the passing shock. When the mass of swept up material is much larger than the initially ablated material, the motion of the blast wave becomes self-similar. We can conservatively estimate the radius at which this happens, assuming that all laser energy is deposited in pin material, heating it uniformly. (Loss of laser energy, e.g., due to misalignment between the laser beam and the pin, and non-uniform heating will lower our estimated radius.) For example, if the ambient gas is xenon and we employ a 10 J laser pulse, the amount of ablated iron is such that when the blast wave has traveled 0.4 mm it will have swept up a much larger (ten times larger) xenon mass.

To image a blast wave on spatial scales up to  $\approx 5$  cm, we use two lenses in a telescope configuration and a gated, single-frame, high-speed CCD camera (2 ns gate), along with a low energy, green laser pulse ( $\lambda = 532$  nm wavelength, 15 ns duration) as a backlighter. Blast wave radius as a function of time is obtained by employing a schlieren technique so that




FIG. 1: Blast wave expansion through ambient nitrogen gas (1.3 kPa) at times  $t = 150$  ns to  $6 \mu\text{s}$  after an ablative laser pulse (energy  $E = 10$  J, duration 5 ns) is focused on a solid pin (visible in images, pointed down). The laser pulse was incident from the left. The shock is spherical (except on the laser side due to laser-plasma interaction) and its growth is consistent with a Sedov-Taylor blast wave. The image to the right ( $t = 150$  ns to  $6 \mu\text{s}$ ) is a composite of four images (with overlapping pin locations).

only laser light perturbed by the plasma is imaged; light which has not been deflected is removed at the telescope focal point by a small ( $\approx 500 \mu\text{m}$ ) beam block. With this method, image brightness corresponds to the spatial derivative of plasma electron density, and the blast wave structure is readily seen. Schlieren images were obtained from 5 ns up to  $35 \mu\text{s}$  after the initial, ablative laser pulse.

Examples of images using nitrogen as the ambient gas can be seen in Fig. 1. In each image, the laser ablating the target pin is incident from the left. The pin is clearly visible, as is the expanding blast wave. [Note that the shock is not exactly centered on the pin and also is not perfectly spherical; deviation from sphericity is evident on the laser side of the pin. A 2D computer simulation (computed by the CALE code [27], an arbitrary Lagrangian-Eulerian code including both hydrodynamic and radiative effects) shows how laser-plasma

interaction causes pre-ionization and preheating of the plasma on the laser side. The shock then propagates through a non-uniform plasma, leading to a non-uniform shock expansion. This effect is further discussed by Edens *et al.* [28].] The shock expansion settles (to within measurement error) into the Sedov-Taylor relationship for a blast wave  $r_s \propto t^{2/5}$  after an initial, brief, non-self similar phase. This is expected for a shock in which the net radiative effects are small. (It also appeared that  $r_s \propto (E/\rho_0)^{1/5}$ , where the total energy  $E$  is the laser energy, also consistent with a Sedov-Taylor blast wave.)

Examples of images using xenon as the ambient gas can be seen in Fig. 2. With its higher atomic number, xenon radiates more strongly than nitrogen, and there are notable differences in the images pointing to more influential radiative effects: (a) plasma emission from pre-heated gas, that is gas heated by the radiation from the shock, is clearly visible as a glow surrounding the shock at early times ( $t \lesssim 400$  ns). (b) The shock expansion is slower than in nitrogen, even when accounting for the  $r_s \propto \rho^{-1/5}$  factor from the higher density of xenon. We also observe that the shock weakens ( $t \approx 1 \mu\text{s}$  to  $4 \mu\text{s}$ ) and gets increasingly difficult to detect. Before the initial shock becomes undetectable ( $t \approx 8 \mu\text{s}$ ), a second shock forms out ahead of the initial shock. This second shock is a phenomenon that has not previously been discussed in literature in this context. It is not surrounded by a glow of pre-heated gas, and it continues to propagate long after the initial shock can no longer be detected. Examples of shocks in xenon at higher laser energies are shown in Fig. 3. The same qualitative evolution occurs at these higher laser energies. A plot of shock radius versus time, shown in Fig. 4, clearly shows a "step" where the second shock forms. This step must be made by a second shock forming; the initial shock cannot suddenly jump ahead.

To help interpret our experimental results we ran a 1D numerical simulation using the LASNEX code [29, 30], see Fig. 5. Radiation was treated in the multigroup diffusion approximation, which is valid when the radiation field is nearly isotropic. In this experiment this means having a large enough optical depth in the RHW, a condition that is satisfied reasonably early in the experiment (see below). We also saw that more sophisticated radiation treatments gave very similar answers for the entire evolution. Opacities were calculated in-line using a screened hydrogenic (SH) approximation which reproduces the average degree of ionization reported for very similar conditions using a more sophisticated model [31] to within  $\sim 25\%$ . The early time laser interaction region was treated with a time-dependent non local thermodynamic equilibrium (non-LTE) version of the SH model. At later times it

made little difference to the overall energetics and evolution of the blast wave if LTE was assumed or not. The equation of state (EOS) was computed using either QEOS [32] in LTE, or from the non-LTE population distribution in non-LTE. More sophisticated models may alter the details of the predictions, but based on a number of calculations in which we have varied the EOS and opacities, we do not expect the qualitative picture to change. In any event, the simulation captures the observed blast wave behavior rather well.

Qualitatively the simulation follows the discussion of this process given by Reinicke and Meyer-ter-vehn [26], but also includes the birth of the second shock wave. Before presenting a more detailed description of the computational results we summarize them here. At early times, the initial shock (which we will refer to as  $s_1$ ) is fast enough to radiate very strongly. The radiation mean free path in the cold ambient gas ahead of  $s_1$  is relatively short, resulting in the formation of a supersonic RHW, which propagates in advance of  $s_1$ . (Note that if the surrounding gas is optically thin, the energy is instead lost, transported to "infinity," and RHW would not form.) As  $s_1$  continually slows down, it radiates less and less. The radiated power soon drops below the rate at which  $s_1$  sweeps up energy from gas heated by the RHW, and eventually  $s_1$  slows below the minimum velocity at which a shock could have created a supersonic radiative heat wave in the first place. At this time RHW is still far ahead of  $s_1$ , but its velocity has also been diminishing rapidly because of expansion and a rapidly weakening driving source. Before  $s_1$  is able to catch it, RHW becomes transonic, and eventually gives birth to a second shock wave (which we will refer to as  $s_2$ ). RHW then falls behind  $s_2$ , which itself is too slow to be radiative. By the time  $s_2$  is formed  $s_1$  is relatively weak (Mach number  $M \approx 2$ ), and it continues to weaken as it propagates in the downstream material of  $s_2$ . After  $s_2$  has roughly doubled its radius, it is no longer influenced by the details of how it was formed, and the shock trajectory closely assumes that of a self-similar Sedov-Taylor blast wave. We will now elaborate on this sequence of events, beginning with the motion of  $s_1$ .

At the end of the laser pulse ( $t = 5$  ns), motion is still strongly influenced by the details of the initial conditions.  $s_1$  is traveling in excess of 60 km / s and is strongly radiative. This is consistent with the results of Bouquet et al [9], who find that shocks at this speed (and at higher densities) are strongly radiative. To help quantify the importance of radiation we have calculated the (inverse) Boltzmann number  $Bo^{-1} = \sigma T^4 / v_s \rho_0 c_v T$  for this time from the numerical simulation. The denominator in this equation is the flux required to



heat the upstream material (flow speed  $v_s = \frac{dr_s}{dt}$ , density  $\rho_0$ ) to the temperature  $T$  of the downstream plasma, while the numerator would be approximately the flux delivered from the hot, downstream plasma to the upstream material. When these two fluxes are equal, the shock is said to be critical, and the upstream material is heated by radiation to the temperature of the downstream plasma, the "critical" temperature  $T = T_c$ . When the radiative flux exceeds this value, the shock is said to be supercritical. In this case, the upstream plasma is still heated to the temperature  $T > T_c$  of the downstream plasma, and the "excess" flux is used to increase the extent of the radiative precursor. The flux  $F_R$  actually emitted through the shock by the downstream plasma in this case is calculated to be somewhat less than  $\sigma T^4$  because the plasma is too small to be optically thick. Nevertheless, the calculated value of  $F_R$  still exceeds the calculated value of the denominator  $v_s \rho_0 c_v T$  by  $\sim 50$ , and the shock is supercritical at this time. We note that the temperature directly behind the shock spikes above the downstream temperature  $T$  we refer to above, but rapidly relaxes toward  $T$ . An excellent discussion of radiating shock structure, including many details omitted here, can be found in Zel'dovich and Raizer [23], in which the radiation field is formulated in the diffusion approximation to make the problem tractable. Other discussions presenting additional insights and extensions can be found in the excellent texts of Mihalas and Mihalas [33] and Castor [34].

The strongly radiative s1 drives a highly supersonic RHW far out ahead of itself (RHW radius  $r_h \approx 2.5$  mm); this heat wave is driven by radiation, not thermal electrons, as the radiative conductivity is at least two orders of magnitude greater than the electron thermal conductivity (e.g., at  $T \sim 2.5$  eV we calculate  $\sim 5400$  W / (m  $\times$  K) for the radiation and 7.3 W / (m  $\times$  K) for the electrons) and additionally we see no noticeable difference in the numerical simulation if electron heat conduction is explicitly omitted. The s1 compression  $\eta$  is very high ( $\eta > 20$ ), a feature consistent with strongly radiating shocks. By  $t \approx 20$  ns, memory of the details of the initial conditions is no longer apparent. s1 has expanded to  $r_s \approx 1.4$  mm and slowed to  $\sim 30$  km / s, but is still strongly radiative ( $\varepsilon \approx 0.8$ ). However, only about one eighth of the total energy now resides inside s1 with the remainder in RHW which extends to  $r_h \approx 4$  mm. (As a result, the subsequent evolution of RHW cannot be much affected by s1.)

As s1 continues to slow down, its ability to radiate diminishes rapidly. Eventually s1 sweeps up energy from the material heated by RHW faster than it radiates and thereafter

the energy inside s1 gradually increases ( $\varepsilon < 0$ ). This should result in a slight acceleration of the blast wave, but the experimental measurement is not accurate enough to verify this. By  $t = 300$  ns, s1 has expanded to  $r_s \approx 4$  mm and its velocity has fallen to  $\sim 6$  km/s, or twice the speed of sound in the RHW plasma into which s1 propagates. The s1 compression is  $\eta \approx 3$ , consistent with a relatively weak shock wave (this is just like the point explosion with counter pressure discussed by Zel'dovich and Raizer [23]). We calculate that a shock born at 6 km/s in cold xenon gas would be too slow to be strongly radiative. Also, a shock at this speed but free of the significant pressure of the preheated RHW plasma would have a compression  $\eta \approx 7$  (where the additional compression above the  $\gamma=5/3$  strong shock limit of  $\eta = 4$  is simply caused by ionization losses in the xenon).

We turn now to details of the RHW. At early time the plasma inside the RHW is optically thin and the radiation wave is essentially a bleaching wave. However, as RHW expands and cools, the optical depth inside the wave rapidly increases, reaching 10 by  $t \approx 200$  ns. By  $t \approx 300$  ns its radius is about twice the shock radius ( $r_h \approx 9$  mm) and its velocity has dropped to that of s1. From here on s1 gradually makes ground on the stalling RHW, but before s1 can catch RHW the latter enters the transonic regime (relative to the sound speed in the hot plasma behind RHW) and begins to drive a non-linear disturbance which eventually ( $t \approx 1.2 \mu\text{s}$ ) breaks into a shock (which we will refer to as s2). At this time RHW drops behind s2.

Because the formation of s2 is a new result we have gone to some effort to ensure that our interpretation is correct. Most importantly we collected a large amount of experimental data around the transition time (including shocks in xenon-nitrogen mixes that we will expand upon at a later date), e.g., Fig. 4. It categorically shows s1 and s2 simultaneously and follows their evolution. We also performed two confirmatory LASNEX calculations. In the first we took the RHW temperature distribution well before the formation of s2 and imposed it in a stationary, uniform xenon gas. As expected, RHW advanced and s2 formed just as in the full calculation. In the second calculation, we used a spherical piston to drive a shock with the same trajectory as in the full calculation (where the shock was driven by the laser). This second calculation produced nearly identical results to the full calculation, demonstrating that the observed dynamics result from the radiative nature of s1, and are not substantially influenced by radiation from the target. Any heating of the gas as a result of target radiation would be prompt, and we see no evidence of this. The results of the

second calculation were confirmed experimentally by a limited set of shots on graphite pins, which yielded the same results. Also, what makes our interpretation unambiguous is that the radiation from a blast wave in xenon drives secondary blast waves off objects placed some distance away, before the original blast wave reaches the objects. (The objects are also well out of the way of the laser beam.) This does not happen when we use nitrogen gas. We have seen this in several experiments at two different laser facilities.

Finally, we compared the observed formation location of s2 to the following simple, analytical estimate for where s2 forms:

Consider the simple 1D fluid equations for conservation of mass  $\rho_1 u_1 = \rho_2 u_2$  and momentum  $p_1 + \rho_1 u_1^2 = p_2 + \rho_2 u_2^2$  in the lab frame of RHW, where subscript 1 denotes the region ahead of RHW, and subscript 2 denotes the region behind RHW. Assuming an ideal gas (so that  $p = \rho c^2$  where  $c$  is the speed of sound) we combine these to obtain

$$\frac{\rho_2}{\rho_1} = \frac{c_1^2 + u_1^2 \pm \sqrt{(c_1^2 + u_1^2)^2 - 4c_2^2 u_1^2}}{2c_2^2}. \quad (1)$$

A supersonic ( $u_1 > c_1$ ) RHW and a real compression  $\eta \equiv \rho_2/\rho_1$  requires  $u_1 \geq c_2 + \sqrt{c_2^2 - c_1^2} \approx 2c_2$  (where the approximation is valid because the temperature behind RHW is much higher than the temperature before it), i.e., requires the mixed Mach number  $M \equiv u_1/c_2 \geq 2$ . Once the Mach number drops to 2, RHW can no longer fulfill Eq. 1, and a shock (s2) forms at RHW. This is a standard result in heat front physics and is analogous to when a blast wave forms ahead of a fireball [23, 33, 35]. s2 immediately moves ahead of RHW and acts to slow down  $u_1$  so that RHW is now subsonic (satisfying  $u_1 \leq c_2 - \sqrt{c_2^2 - c_1^2} \approx c_1^2/2c_2$ ). To estimate when RHW slows to Mach 2 and what its radius  $r_h$  then is, we can assume a radiative conductivity of the ambient gas of the form  $\chi = \chi_0 \rho^a T^b$  and use Barenblatt's solution for an instantaneous point release of energy [26, 36]. (A point source is a reasonable estimate [26] because most of the energy that s1 can lose through radiation is lost at an early stage when there is a large separation between s1 and RHW.) Using values for our experiment in xenon,  $E = 10$  J,  $\rho_0 = 78$  g/m<sup>3</sup>,  $\gamma = 1.2$ , and  $\chi = 10^{-44} \rho^{-2.2} T^{10}$  in SI-units, we find that the RHW Mach number drops to Mach 2 when  $r_h \approx 10$  mm. In the experiment we first observed s2 with  $r_h \approx 12$  mm, in reasonable agreement with the analytical estimate, particularly considering that the heat front is not sharp and it takes some time before s2 forms and becomes observable.

From all the above, we summarize the blast wave evolution in xenon in the following

steps: (1) The laser energy is deposited in pin material which then becomes very hot. (2) The heated pin material expands rapidly, pushing at the surrounding gas, setting up a strong, radiative s1. (3) Radiation from s1 heats the surrounding gas. The flux is high enough that s1 is supercritical, driving a supersonic RHW that travels rapidly outward, leaving a large separation between s1 and RHW. (4) s1 sweeps up enough material that the details of its initial conditions become unimportant. (5) s1 slows and its ability to radiate efficiently quickly decreases. (6) s1 is traveling into the counter pressure of hot RHW plasma, which is becoming comparable to the ram pressure; the Mach number drops rapidly, and the post-shock compression reduces correspondingly. (7) When the Mach number for RHW reaches  $\sim 2$ , RHW stalls and creates s2. (8) s1 continues to weaken until it dissipates. (9) s2 is essentially non-radiative and once it has swept up enough mass (doubled its initial radius), it propagates like  $r_s \propto t^{2/5}$ , provided it remains strong.

We thank Dwight Price and the staff at the Janus facility (where the experiments were conducted) for their valuable assistance.

This work was performed under the auspices of the U. S. Department of Energy by the University of California, Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48.

- 
- [1] J. C. Bozier, G. Thiell, J. P. LeBreton, S. Azra, M. Decroisette, and D. Schirmann, *Phys. Rev. Lett.* **57**, 1304 (1986).
  - [2] J. Grun, J. Stamper, C. Manka, J. Resnick, R. Burris, J. Crawford, and B. H. Ripin, *Phys. Rev. Lett.* **66**, 2738 (1991).
  - [3] B. A. Remington, D. Arnett, R. P. Drake, and H. Takabe, *Science* **284**, 1488 (1999).
  - [4] D. Ryutov, R. P. Drake, J. Kane, E. Liang, B. A. Remington, and W. M. Wood-Vasey, *Astrophys. J.* **518**, 821 (1999).
  - [5] K. Shigemori, T. Ditmire, B. A. Remington, V. Yanovksy, D. Ryutov, K. G. Estabrook, M. J. Edwards, A. J. MacKinnon, A. M. Rubenchik, K. A. Keilty, et al., *Astrophys. J.* **533**, 159 (2000).
  - [6] H. F. Robey, J. O. Kane, B. A. Remington, R. P. Drake, O. A. Hurricane, H. Louis, R. J. Wallace, J. Knauer, P. Keiter, D. Arnett, et al., *Phys. of Plasmas* **8**, 2446 (2001).

- [7] P. A. Keiter, R. P. Drake, T. S. Perry, H. F. Robey, B. A. Remington, C. A. Iglesias, R. J. Wallace, and J. Knauer, *Phys. Rev. Lett.* **89**, 165003 (2002).
- [8] X. Fleury, S. Bouquet, C. Stehle, M. Koenig, D. Batani, A. Benuzzi-Mounaix, J.-P. Chieze, N. Grandjouan, J. Grenier, T. Hall, et al., *Laser and Particle Beams* **20**, 263 (2002).
- [9] S. Bouquet, C. Stehle, M. Koenig, J.-P. Chieze, A. Benuzzi-Mounaix, D. Batani, S. Leygnac, X. Fleury, H. Merdji, C. Michaut, et al., *Phys. Rev. Lett.* **92**, 225001 (2004).
- [10] J. M. Blondin, E. B. Wright, K. J. Borkowski, and S. P. Reynolds, *Astrophys. J.* **500**, 342 (1998).
- [11] E. Müller, B. Fryxell, and D. Arnett, *Astron. Astrophys.* **251**, 505 (1991).
- [12] J. I. Reed, J. J. Hester, A. C. Fabian, and P. F. Winkler, *Astrophys. J.* **440**, 706 (1995).
- [13] G. Sonneborn, C. S. J. Pun, R. A. Kimble, T. R. Gull, P. Lundqvist, R. McCray, P. Plait, A. Bogges, C. W. Bowers, A. C. Danks, et al., *Astrophys. J. Lett.* **492**, L139 (1998).
- [14] N. Bartel, M. F. Bietenholz, M. P. Rupen, A. J. Beasley, D. A. Graham, V. I. Altunin, T. Venturi, G. Umama, W. H. Cannon, and J. E. Conway, *Science* **287**, 112 (2000).
- [15] B. Reipurth and G. Sandell, *Astron. Astrophys.* **150**, 307 (1985).
- [16] J. R. Jokipii, C. P. Scott, and M. S. Giampapa, eds., Cosmic Winds and the Heliosphere (University of Arizona Press, 1997).
- [17] C. F. McKee and B. T. Draine, *Science* **252**, 397 (1991).
- [18] D. A. Allen and M. G. Burton, *Nature* **363**, 54 (1993).
- [19] R. I. Klein and D. T. Woods, *Astrophys. J.* **497**, 777 (1998).
- [20] E. Cohen, T. Piran, and R. Sari, *Astrophys. J.* **509**, 717 (1998).
- [21] G. I. Taylor, *Proc. R. Soc. London A* **201**, 159 (1950).
- [22] L. I. Sedov, Similarity and Dimensional Methods in Mechanics (Academic, New York, 1959).
- [23] Y. B. Zeldovich and Y. P. Raizer, Physics of Shock Waves and High-Temperature Hydrodynamic Phenomena (Academic, New York, 1966).
- [24] E. Liang and K. Keilty, *Astrophys. J.* **533**, 890 (2000).
- [25] C. F. McKee and J. P. Ostriker, *Astrophys. J.* **218**, 148 (1977).
- [26] P. Reinicke and J. M. ter Vehn, *Phys. Fluids A* **3**, 1807 (1991).
- [27] R. T. Barton, in Numerical Astrophysics, ed. J. M. Centrella, J. M. LeBlanc, and R. L. Bowers (Jones and Bartlett, Boston, 1985).
- [28] A. D. Edens, T. Ditmire, J. F. Hansen, M. J. Edwards, R. G. Adams, P. Rambo, L. Ruggles,

- I. Smith, and J. L. Porter, *Phys. Plasmas* **11**, 4968 (2004).
- [29] G. B. Zimmerman and W. L. Kruer, *Comments Plasma Phys. Controlled Fusion* **2**, 51 (1975).
- [30] G. B. Zimmerman and R. M. More, *J. Quant. Spectrosc. Radiat. Transfer* **23**, 517 (1980).
- [31] J. M. Laming and J. Grun, *Phys. Rev. Lett.* **89**, 125002 (2002).
- [32] R. M. More, K. H. Warren, D. A. Young, and G. B. Zimmerman, *Phys. Fluids* **31**, 3059 (1988).
- [33] D. Mihalas and B. W. Mihalas, Foundations of Radiation Hydrodynamics (Oxford University Press, Oxford, 1984).
- [34] J. I. Castor, Radiation Hydrodynamics (Cambridge University Press, 2004).
- [35] S. P. Hatchett, Tech. Rep., Lawrence Livermore National Laboratory, CA (1991), uCRL-JC-108348: Ablation Gas Dynamics of Low-Z Materials Illuminated by Soft X-Rays. Copies may be obtained from the National Technical Information Service, Springfield, VA 22161.
- [36] G. I. Barenblatt, Similarity, Self-Similarity and Intermediate Asymptotics (Consultants Bureau, New York, 1979).




FIG. 2: Blast wave expansion through ambient Xe gas (1.3 kPa) at times  $t = 50$  ns to  $30 \mu\text{s}$  after an ablative laser pulse (energy  $E = 10$  J, duration 5 ns) is focused on a solid pin (visible in images, pointed down; pin location in bottom row of images is at the left edge of each image). The laser pulse was incident from the left. The initial shock is strongly radiative (super-critical) and preheats the ambient gas. At  $t = 150$  ns both the initial shock and the preheated gas ahead of it are clearly visible. At  $t \approx 1 \mu\text{s}$  the initial shock begins to dissipate, and the shock front is no longer obviously sharp. At  $t \approx 4 \mu\text{s}$  a second shock appears (located at tip of arrow), ahead of the initial shock. The second shock continues to expand while the initial shock gradually becomes undetectable. The final image ( $t = 8 \mu\text{s}, 30 \mu\text{s}$ ) is a composite of two images (with overlapping pin locations).




FIG. 3: Blast wave expansion through ambient Xe gas ( $1.3 \text{ kPa}$ ) at times  $t = 1.0 \mu\text{s}$  to  $4.5 \mu\text{s}$ . Ablative laser pulse energy  $E \approx 120 \text{ J}$ ; duration  $5 \text{ ns}$ . At  $t = 3.5 \mu\text{s}$  both initial and second shocks are clearly visible.






FIG. 4: Measured shock radius versus time in experimental images of shocks in xenon. Ablative laser energy  $E \approx 100 - 200$  J; duration 5 ns. Note the step in radius around the transition time ( $\sim 4 \mu\text{s}$ ) when both shocks are visible.




FIG. 5: Compression and electron temperature versus radius in the Xe plasma at four different stages of the blast wave evolution. Note the initial shock dissipating and the second shock being born at the radiative heat wave front. Also illustrated in the plot (at  $t = 0.3 \mu\text{s}$ ) is the ion temperature which is in equilibrium with the electron temperature except in the thin relaxation layer behind the shock front. This layer is not fully resolved in the simulation shown, and the temperature excursions are not as high as they would be in reality. This is partly responsible for the apparent continuous nature of the electron temperature at the shock. In reality the electrons are essentially adiabatic at the shock, and do undergo a smaller "jump" in temperature [23]. In a simulation of the first few hundred nanoseconds with ten times the spatial resolution, we resolve the relaxation layer much better, yet the other details (and conclusions from those) remain unchanged.