


**ALASKA PRODUCTIVITY SURVEYS
OF GEESE, SWANS, AND BRANT
2005**



Compiled by
Deborah J. Groves

Migratory Bird Management
U.S. Fish and Wildlife Service
3000 Vintage Blvd., Suite 240
Juneau, Alaska 99801
debbie_groves@fws.gov

April 2006

TITLE:

Waterfowl Productivity Surveys for Alaska – 2005

SPECIES SURVEYED:

Pacific Brant (*Branta bernicla nigricans*)
Trumpeter Swan (*Cygnus buccinator*)
Emperor Goose (*Chen canagica*)
Dusky Canada Goose (*Branta canadensis occidentalis*)

CONTRIBUTORS:

U.S. Fish and Wildlife Service (USFWS)
Arctic National Wildlife Refuge
Innoko National Wildlife Refuge
Izembek National Wildlife Refuge
Kanuti National Wildlife Refuge
Kenai National Wildlife Refuge
Koyukuk/Nowitna National Wildlife Refuge Complex
Migratory Bird Management, Anchorage
Migratory Bird Management, Fairbanks
Migratory Bird Management, Juneau
Migratory Bird Management, Laurel, MD
Migratory Bird Management, Mare Island, CA
Migratory Bird Management, Soldotna
Tetlin National Wildlife Refuge
Yukon Flats National Wildlife Refuge
National Park Service
Denali National Park and Preserve
Gates of the Arctic National Park and Preserve
Wrangell-St. Elias National Park and Preserve
Yukon-Charley National Park and Preserve
U.S Forest Service (USFS)
Cordova Ranger District
Yakutat Ranger District
U.S. Geological Survey (USGS)
Alaska Science Center
Alaska Department of Fish and Game
Division of Wildlife Conservation, Statewide Waterfowl Program
Canadian Wildlife Service
University of Alaska Fairbanks
Yakutat Tlingit Tribe
John Liston - Alaska
Comox Valley Naturalists Society – British Columbia
Graeme Fowler – British Columbia
Russ Canniff – Washington
James Wynbrandt – New York

REPORT COMPILED BY:

Deborah J. Groves – Wildlife Biologist, USFWS, Migratory Bird Management, Juneau, AK

ABSTRACT:

Productivity surveys were conducted by several agencies and individuals during late summer, fall, and/or winter of 2005 and early 2006 to estimate juvenile-to-adult age ratios for Pacific brant (*Branta bernicla nigricans*), the Pacific Coast population of trumpeter swans (*Cygnus buccinator*), emperor geese (*Chen canagica*), and dusky Canada geese (*Branta canadensis occidentalis*). The results of these surveys appear in the tables of this report, along with short narratives in the Results section. No productivity data were reported for Cackling cackling geese (*Branta hutchinsii minima*) in 2005.

The following productivity measures were estimated for 2005:

Species	Type of Year	Productivity Estimate	% Change From 2004	% Change From Mean
Pacific Brant	Above Average			
Fall % Juv.		33.3%	+83%	+49%
Fall Juv./Fam.		2.89	+16%	+10%
Winter % Juv.		15.6%	+34%	+27%
Trumpeter Swan	Above Average			
Late Summer Brood Size		3.1	+3%	-3%
Late Summer % Juv.		27.2%	+30%	+7%
Late Summer % Prs. w/ Brd		33.9%	+10%	+6%
Winter % Juv.		20.8%	+22%	+12%
Winter Juv./Fam.		2.35	-6%	+2%
Emperor Goose	Above Average			
Fall % Juv.				
From ground counts		28.6%	+31%	+23%
From aerial photos		18.5%	+67%	+1%
Fall Juv./Fam.		2.79	+15%	+0%
Cackling Cackling Goose	No Report			
Dusky Canada Goose	Below Average			
Late Summer % Juv.		11.8%	-58%	-38%

METHODS:

Fall and winter productivity appraisals generally followed procedures developed by Lynch (1969) and outlined in the Standard Operating Procedures for Productivity Surveys of Geese, Swans and Brant (Draft) 1977. Additional survey methods included late-summer aerial surveys of trumpeter swans (King 1973) and dusky Canada geese (Petrula 2005), analysis of aerial photographs of emperor geese (Anderson et al. 2006), and ocular sightings from the ground (e.g. Audubon Christmas Bird Counts).

RESULTS:

Pacific Brant:

Fall Productivity: Table 1.

Kristine Sowl of Izembek National Wildlife Refuge (NWR) reported that 33.3% juveniles were estimated from a sample of 25,361 brant during ground surveys conducted at Izembek Lagoon, Alaska in September and October. She also estimated a mean of 2.89 juveniles per family group from a sample of 89 families. The proportion of juveniles was 83% higher than 2004 and 49% above the 42-year mean. The mean family group size was 16% higher than 2004 and 10% above the 39-year mean.

Winter Productivity: Table 2.

Russ Canniff collected data from Padilla and Samish Bays, WA on 20 January 2006. He estimated 18.5% juveniles from a sample of 119 brant. He also reported a mean of 2.08 juveniles per family group from a sample of 12 families. He noted that nearly all (97%) of the brant he observed were the gray-bellied variety from the western high arctic population.

Dave Pitkin of Oregon Coast NWR Complex reported that 67 brant were aged in Netarts Bay, Oregon in 2006. From that sample, 7, or 10.4%, were juveniles.

Combining data from both winter survey areas resulted in an estimate of 15.6% juveniles from a sample of 186 brant. This was 34% higher than 2004 and 27% above the 21-year mean.

In summary, Pacific brant experienced above-average production in 2005.

Trumpeter Swan:

Late-Summer Productivity: Tables 3 and 4.

A complete aerial census of trumpeter swans in Alaska was conducted 20 July – 8 September by USFWS and other cooperating agencies. Results indicated a mean brood size of 3.1 (n=2,084 broods), 27.2% juveniles in the population (n=23,692 total swans), and 33.9% pairs with brood (n=5,970 pairs) (Table 3). The mean brood size was 3% higher than 2004 but was 3% below the 29-year mean (Table 4). The proportion of juveniles was 30% higher than 2004 and 7% above the mean. The percentage of pairs with a brood was 10% higher than 2004 and 6% above the mean.

Winter Productivity: Table 5.

In Alaska, Nate Catterson of USFS, Yakutat Ranger District, reported that 15.2% juveniles were observed from a total of 646 swans during an aerial population survey of the Yakutat Forelands on 2 March 2006.

On Vancouver Island, British Columbia, Graeme Fowler reported the results of swan surveys conducted from November 2005 through February 2006 by the Comox Valley Naturalists Society. The mean % juvenile was 20.0% ($n = 419$) in November, 21.0% ($n = 1,814$) in December, 20.5% ($n = 2,323$) in January, and 21.0% ($n = 2,305$) in February. Note that only the February figure was included in Table 5.

This year CWS conducted a winter swan survey of southwestern British Columbia, including all of Vancouver Island and the Fraser Valley between Hope and the Fraser River delta. The survey was conducted on 19 January in the Fraser Valley and 8-10 February on Vancouver Island. Sean Boyd reported that of 7570 total swans recorded, 25.1% were identified as juveniles. Note that to maintain consistency in the report table these results are not included in Table 5.

In northwest Washington, Russ Canniff recorded age ratios for trumpeter swans in Skagit Valley and Port Susan in February 2006. He found that 21.6% were juveniles from a sample of 3,519 swans. He also collected data on family group size from November 2005 through February 2006 and found a mean of 2.35 juveniles per family group from a sample of 28 families.

Data from all winter survey areas combined (excluding the CWS survey) resulted in an estimate of 20.8% juveniles from a sample of 6,470 swans. This was 22% higher than 2004 and 12% above the 28-year mean. The mean family group size was 2.35 from a sample of 28 families. This was 6% lower than 2004 but 2% above the 23-year mean.

In summary, trumpeter swans experienced above-average production in 2005.

Emperor Goose: Tables 6 and 7.

Kristine Sowl reported that 28.6% juveniles were estimated from a sample of 3,983 emperor geese during ground surveys conducted at Izembek Lagoon, Alaska in September and October (Table 6). She also estimated a mean of 2.79 juveniles per family group from a sample of 131 families. The proportion of juveniles was 31% higher than 2004 and 23% above the 38-year mean. The mean family group size was 15% higher than 2004 and equal to the mean.

Paul Anderson and Bob Stehn of USFWS, Migratory Bird Management Anchorage reported the results of aerial photo work on the Alaska Peninsula conducted in late September. They estimated the proportions of juveniles in seven major lagoons from aerial photos and then weighted the proportions by the population counts of those lagoons from an independent aerial population survey. The result was a weighted-mean estimate of 18.5% juveniles for the 2005 fall population, 67% higher than 2004 and 1% above the 20-year mean (Table 7).

In summary, emperor geese experienced above-average production in 2005.

Cackling Cackling Goose: No Report.

Dusky Canada Goose: Table 8.

Tom Rothe of the Alaska Department of Fish and Game reported the results of an aerial production survey that was flown over the west Copper River Delta on 17 July. Of a total count of 5,364 geese, 11.8% were identified as juveniles. The proportion of juveniles was 58% lower than 2004 and 38% below the 34-year mean.

In summary, dusky Canada geese experienced below-average production in 2005.

LITERATURE CITED

- Anderson, P.D., W.W. Larned, E.J. Mallek, C.P. Dau, and R.A. Stehn. 2006. Monitoring emperor goose populations by aerial counts and fall age-ratio. Annual progress report, U.S. Fish and Wildlife Service, Anchorage, AK.
- King, J.G. 1973. The use of small airplanes to gather swan data in Alaska. Wildfowl 24: 15-20.
- Lynch, J.J. 1969. Appraisals of annual productivity and mortality among geese, swans, and other birds. Annual Report, Part II and Appendix A. U.S. Fish and Wildlife Service. 26 pp.
- Petrula, M.J. 2005. Dusky Canada goose production survey – 2005. Final report to Pacific Flyway Study Committee. Alaska Department of Fish and Game, Anchorage, AK. 6pp.
- U.S. Fish and Wildlife Service, 1977. Standard Operating Procedures for Productivity Surveys of Geese, Swans, and Brant (Draft). 52 pp.

ACKNOWLEDGEMENTS

The following agencies and individuals are gratefully acknowledged for their contributions to this report:

U.S. Fish and Wildlife Service

Arctic National Wildlife Refuge

Alan Brackney

Roger Kaye

Innoko National Wildlife Refuge

Robin Corcoran

Izembek National Wildlife Refuge

Kristine Sowl

Kanuti National Wildlife Refuge

Mike Spindler

Chris Harwood

Kenai National Wildlife Refuge

Liz Jozwiak

Koyukuk/Nowitna National Wildlife Refuge Complex

Jenny Bryant

Brad Scotton

Joe Huhndorf

Melanie Hans

Migratory Bird Management, Anchorage

Paul Anderson

Chris Dau

Bill Eldridge

Julian Fischer

Tamara Mills

Russ Oates

Bob Stehn

Heather Wilson

Migratory Bird Management, Fairbanks

Ed Mallek

Migratory Bird Management, Juneau

Bruce Conant

Jack Hodges

Deb Groves

Jim King

Migratory Bird Management, Laurel, MD

Karen Bollinger

Migratory Bird Management, Mare Island, CA

Rod King

Migratory Bird Management, Soldotna

Bill Larned

Tetlin National Wildlife Refuge

Gail Collins

Jim Ellis

Susy Grimes
Hank Timm
Yukon Flats National Wildlife Refuge
Mark Bertram
Mike Vivion
Jim Akaran
Colette Buchholtz
National Park Service
Denali National Park and Preserve
Carol McIntyre
Gates of the Arctic/Yukon-Charley National Parks and Preserves
Nikki Guldager
Wrangell-St. Elias National Park and Preserve
Mason Reid
Tom Betts
Miranda Terwilliger
U.S Forest Service
Cordova Ranger District
Dan Logan
Yakutat Ranger District
Susan Oehlers
Nate Catterson
U.S. Geological Survey
Alaska Science Center
Dave Ward
Alaska Department of Fish and Game
Division of Wildlife Conservation, Statewide Waterfowl Program
Tom Rothe
Mike Petrula
Canadian Wildlife Service
Pacific Wildlife Research Center
Sean Boyd
University of Alaska Fairbanks
Josh Schmidt
Yakutat Tlingit Tribe
Eileen Henniger
Others
John Liston – Yakutat, Alaska
Comox Valley Naturalists Society – Courtenay, British Columbia
Graeme Fowler – Comox, British Columbia
Russ Canniff – Snohomish, Washington
James Wynbrandt – New York, New York

Table 1. Historical fall productivity records for Pacific brant at Izembek Lagoon, AK, 1963-2005.^a

Year	Grouped Birds			Family Associations		
	Adults	Juveniles	% Juv.	Families	Juveniles	Juv./Family
1963	3968	1243	23.9			
1964	13324	4577	25.6			
1965	21210	5050	19.2			
1966	9927	7134	41.8	195	557	2.86
1967	15219	3081	16.8	359	926	2.58
1968	15110	3117	17.1	145	377	2.60
1969	12829	3577	21.8	293	780	2.66
1970	12104	6256	34.1	148	476	3.22
1971	4820	1953	28.8	295	716	2.43
1972	6599	3698	35.9	153	416	2.72
1973	12025	4999	29.4	327	938	2.87
1974	13118	632	4.6	105	239	2.28
1975	9396	5452	36.7	189	543	2.87
1976	7962	4340	35.3	237	674	2.84
1977	8856	4092	31.6	240	603	2.51
1978	10696	1842	14.7	110	326	2.96
1979	13674	2349	14.7	146	361	2.47
1980	9618	3341	25.8	177	489	2.76
1981	4109	936	18.6	154	431	2.80
1982	11509	1213	9.5	89	237	2.66
1983	6149	1947	24.0	173	515	2.98
1984	9451	1499	13.7	192	564	2.94
1985	12032	1915	13.7	624	1538	2.46
1986	15621	2823	15.3	137	352	2.57
1987	17411	7882	31.2	948	2587	2.73
1988	16138	3847	19.2	263	633	2.41
1989	13654	4281	23.9	303	914	3.02
1990	24215	5750	19.2	349	894	2.56
1991	31432	12127	27.8	415	1066	2.57
1992	55795	11044	16.5	404	1127	2.79
1993	103254	31942	23.6	979	2727	2.79
1994	21371	2808	11.6	353	735	2.08
1995	26964	15240	36.1	78	218	2.79
1996	15148	4201	21.7	50	152	3.04
1997	15216	3105	16.9	40	106	2.65
1998	8214	2836	25.7	220	488	2.22
1999	12500	3450	21.6	111	254	2.29
2000	6669	2982	30.9	91	202	2.22
2001	14829	1198	7.5	68	167	2.46
2002	18441	4751	20.5	92	222	2.41
2003	27517	4371	13.7	197	446	2.26
2004	19715	4384	18.2	129	322	2.50
2005	16906	8455	33.3	89	257	2.89
Mean ^b		22.3			2.64	
% Change from:						
2004		83%			16%	
Mean		49%			10%	

^a Data supplied by Izembek National Wildlife Refuge and USGS Alaska Science Center.^b Mean excludes 2005.

Table 2. Historical winter productivity records for brant in the Pacific Flyway, 1983-2005.

Year ^a	Padilla/Samish Bays, WA ^b			Willapa Bay, WA ^c			Olympic Peninsula, WA ^d			Oregon Coast ^e			Combined Productivity % Juv.
	Ad.	Juv.	% Juv.	Ad.	Juv.	% Juv.	Ad.	Juv.	% Juv.	Ad.	Juv.	% Juv.	
1983				982	166	14.5							14.5
1984				2605	251	8.8							8.8
1985													
1986	3731	292	7.3	1925	186	8.8	217	11	4.8				7.7
1987	3110	1242	28.5	997	196	16.4	1540	306	16.6				23.6
1988	2003	297	12.9	1167	184	13.6	1544	311	16.8				14.4
1989	4928	622	11.2	982	88	8.2	2231	232	9.4				10.4
1990	3047	837	21.5				2013	88	4.2				15.5
1991	2464	336	12.0	1189	126	9.6	913	123	11.9				11.4
1992	6294	669	9.6	944	88	8.5	839	46	5.2				9.0
1993	3032	1074	26.2				1299	265	16.9				23.6
1994	3771	197	5.0	937	97	9.4	1034	26	2.5				5.3
1995	1083	185	14.6				634	15	2.3				10.4
1996	1964	530	21.3	70	12	14.6	793	20	2.5				16.6
1997	1660	189	10.2				779	50	6.0				8.9
1998	2573	466	15.3	125	19	13.2							15.2
1999	1199	349	22.5				386	29	7.0				19.3
2000	877	337	27.8	1818	183	9.1	430	32	6.9				15.0
2001	1089	11	1.0				361	24	6.2				2.4
2002							368	28	7.1				7.1
2003	752	48	6.0				551	25	4.3	476	51	9.7	6.5
2004	647	85	11.6										11.6
2005	97	22	18.5							60	7	10.4	15.6
Mean ^f		14.7		11.2			7.9			9.7		12.2	
% Change from:													
2004		59%		N/A			N/A			N/A		34%	
Mean		26%		N/A			N/A			7%		27%	

^a Surveys conducted some time between November of the stated year and February of the next year.

^b Data supplied by Russ Canniff and Washington Department of Wildlife. A high proportion of these birds are the "gray-bellied" variety.

^c Data supplied by Willapa National Wildlife Refuge and Washington Department of Wildlife.

^d Data supplied by Washington Maritime National Wildlife Refuge Complex.

^e Data supplied by Oregon Coast National Wildlife Refuge Complex.

^f Mean excludes 2005.

Table 3. Results of late-summer 2005 complete census of trumpeter swans in Alaska.^a

Area	Number of 1:63,360 Maps Surveyed	Date(s) Surveyed	Adults and Subadults				Cygnets	Total Swans	Broods	Mean Brood Size	% Juv.	% Pairs w/ Brood
			In Pairs	As Singles	In Flocks	Subtotal						
Gulf Coast	44	8/1-9/1	800	85	474	1359	459	1818	141	3.3	25.2	34.5
Copper Canyon	8	8/6	56	3	33	92	24	116	7	3.4	20.7	21.4
Gulkana	81	8/2-8/25	2440	252	510	3202	1228	4430	393	3.1	27.7	31.1
Kenai	23	8/16-8/22	282	20	91	393	172	565	52	3.3	30.4	36.2
Cook Inlet	97	8/3-8/27	1470	196	310	1976	694	2670	216	3.2	26.0	28.4
Lower Tanana	128	8/2-8/31	3054	305	1040	4399	1786	6185	607	2.9	28.9	38.7
Kuskokwim	89	8/8-9/8	1016	69	338	1423	535	1958	186	2.9	27.3	35.4
Koyukuk	81	8/2-9/4	950	104	467	1521	460	1981	163	2.8	23.2	33.1
Yukon Flats	142	8/5-9/8	632	40	374	1046	324	1370	103	3.1	23.6	32.0
S.E. Mainland	21	7/20-7/29	76	10	56	142	70	212	22	3.2	33.0	50.0
Upper Tanana	66	8/11-8/31	1164	73	455	1692	695	2387	194	3.6	29.1	33.0
Total	780		11940	1157	4148	17245	6447	23692	2084	3.1	27.2	33.9

^a Data supplied by USFWS Migratory Bird Management in Juneau, Anchorage, Soldotna, Fairbanks, and Region 9, Arctic NWR, Kanuti NWR, Koyukuk/Nowitna NWR, Tetlin NWR, Yukon Flats NWR, Denali N.P., Gates of the Arctic N.P., Wrangell-St. Elias N.P., Yukon-Charley N.P., USFS Cordova Ranger District, and USFS Yakutat Ranger District.

Table 4. Historical late-summer productivity records for trumpeter swans in Alaska, 1968-2005.^a

Year	Number of 1:63,360 Maps Surveyed	Adults and Subadults				Total Swans	Broods	Mean Brood Size	% Juv.	% Pairs w/ Brood
		In Pairs	As Singles	In Flocks	Subtotal					
1968	181	1320	108	496	1924	923	2847	257	3.6	32.4
1975	285	2102	151	740	2993	1177	4170	378	3.1	28.2
1978	13	284	36	130	450	116	566	37	3.1	20.5
1979	13	264	26	229	519	164	683	46	3.6	24.0
1980	297	3324	169	1766	5259	2437	7696	683	3.6	31.7
1981	19	632	23	673	1328	547	1875	136	4.0	29.2
1982	36	1164	97	443	1704	421	2125	138	3.1	19.8
1983	46	1260	69	488	1817	903	2720	230	3.9	33.2
1984	43	1358	125	780	2263	755	3018	230	3.3	25.0
1985	425	5120	449	2204	7773	1686	9459	588	2.9	17.8
1986	113	2560	184	678	3422	1349	4771	438	3.1	28.3
1987	73	1640	108	760	2508	1030	3538	294	3.5	29.1
1988	54	1610	103	1203	2916	1087	4003	322	3.4	27.2
1989	63	1150	105	295	1550	488	2038	158	3.1	23.9
1990	625	7056	647	2039	9742	3595	13337	1124	3.2	27.0
1991	61	1968	123	936	3027	923	3950	322	2.9	23.4
1992	80	1592	119	819	2530	825	3355	270	3.1	24.6
1993	76	1766	127	663	2556	1080	3636	341	3.2	29.7
1994	69	1982	128	1094	3204	1196	4400	374	3.2	27.2
1995	674	7946	859	3184	11989	3834	15823	1218	3.1	24.2
1996	50	1624	116	1042	2782	814	3596	256	3.2	22.6
1997	46	1212	72	566	1850	584	2434	189	3.1	24.0
1998	51	1702	104	740	2546	976	3522	281	3.5	27.7
1999	27	508	36	212	756	228	984	71	3.2	26.0
2000	733	9986	899	3049	13934	3223	17157	1149	2.8	18.8
2001	22	1164	66	491	1721	531	2252	168	3.2	23.6
2002	35	1118	111	521	1750	488	2238	165	3.0	21.8
2003	55	2066	206	844	3116	1212	4328	407	3.0	28.0
2004	39	1086	118	792	1996	529	2525	177	3.0	21.0
2005	780	11940	1157	4148	17245	6447	23692	2084	3.1	27.2
Mean ^b								3.2	25.4	32.0
% Change from:										
2004								3%	30%	10%
Mean								-3%	7%	6%

^a Complete statewide censuses were conducted in 1968, 1975, 1980, 1985, 1990, 1995, 2000, and 2005 (shaded in gray). In other years, surveys were conducted by various agencies to meet local objectives.

^b Mean excludes 2005.

Table 5. Historical winter productivity records for trumpeter swans in the Pacific Flyway, 1977-2005.

Year ^a	Alaska ^b					SE Vancouver Island, BC ^c			Skagit Valley/Port Susan, WA ^d				Combined Productivity	
	Ad.	Juv.	% Juv.	No. Fam.	Juv./Fam.	Ad.	Juv.	% Juv.	Ad.	Juv.	% Juv.	No. Fam.	Juv./Fam.	% Juv.
1977									214	70	24.6			24.6
1978						384	134	25.9	218	76	25.9			25.9
1979	431	129	23.0	15	2.60	459	158	25.6	273	82	23.1			24.1
1980	167	65	28.0	27	2.41	499	211	29.7	310	127	29.1	45	2.82	29.2
1981									316	92	22.5	41	2.24	22.5
1982	110	35	24.1	14	2.50				339	56	14.2	24	2.33	16.9
1983	115	29	20.1	4	1.50	533	113	17.5	330	94	22.2	39	2.41	19.4
1984	109	79	42.0	5	2.40	1101	216	16.4	359	62	14.7	29	2.14	18.5
1985	95	14	12.8	1	2.00	1336	98	6.8	340	44	11.5	22	2.00	8.1
1986	146	40	21.5	7	1.29	1228	280	18.6	356	113	24.1	49	2.31	20.0
1987	146	52	26.3	20	2.60	1081	334	23.6	347	133	27.7	49	2.71	24.8
1988	164	52	24.1			1353	304	18.3	473	111	19.0	48	2.31	19.0
1989	239	55	18.7			1209	194	13.8	568	128	18.4			15.8
1990	266	57	17.6	14	2.21	1553	295	16.0	678	111	14.1			15.6
1991	696	267	27.7	21	2.67	1049	165	13.6	810	155	16.1	64	2.42	18.7
1992	578	169	22.6	19	2.53	1639	149	8.3	905	94	9.4	45	2.09	11.7
1993	667	322	32.6	30	2.70	1801	530	22.7	762	233	23.4	167	2.40	25.1
1994	562	190	25.3	15	3.27	1543	536	25.8	927	242	20.7	112	2.41	24.2
1995	294	61	17.2			1427	398	21.8	1187	239	16.8	83	2.46	19.4
1996						1307	195	13.0	1774	312	15.0	93	2.31	14.1
1997						1540	272	15.0	1569	249	13.7	102	2.23	14.4
1998	272	35	11.4			1427	286	16.7	2180	381	14.9	76	2.34	15.3
1999	338	59	14.9			1380	198	12.5	2384	336	12.4	67	2.03	12.6
2000	585	118	16.8			1612	275	14.6	2256	355	13.6	84	2.04	14.4
2001	191	79	29.3			1763	204	10.4	1936	366	15.9	53	2.19	14.3
2002	76	17	18.3			1659	263	13.7	2256	521	18.8	149	2.31	16.7
2003	580	151	20.7			1479	339	18.6	4158	912	18.0	210	2.19	18.4
2004	508	84	14.2			1886	377	16.7	3301	706	17.6	106	2.50	17.0
2005	548	98	15.2			1820	485	21.0	2758	761	21.6	28	2.35	20.8
Mean ^e		22.1	2.36				17.4			18.5		2.31	18.6	
% Change from:														
2004		7%	N/A				26%			23%		-6%	22%	
Mean		-31%	N/A				21%			17%		2%	12%	

^a Surveys conducted between November of the given year and February of the next year.^b Data supplied by AK Dept. of Fish and Game, USFS Cordova and Yakutat, AK, USFWS Region 7 Migratory Bird Management, Peter Walsh, and Paul Meyers.^c Data supplied by British Columbia Ministry of Environment, Land, and Parks, Comox Valley Naturalists Society, and Graeme Fowler.^d Data supplied by Russ Canniff.^e Mean excludes 2005.

Table 6. Historical fall productivity records (from ground counts) for emperor geese at Izembek Lagoon, AK, 1966-2005.^a

Year	Grouped Birds			Family Associations ^b		
	Adults	Juveniles	% Juv.	Families	Juveniles	Juv./Family
1966	699	265	27.5	132	331	2.51
1967	1457	585	28.6	66	215	3.26
1968	1195	585	32.9	40	112	2.80
1969	4149	2980	41.8	161	530	3.29
1970	9722	4933	33.7	383	1115	2.91
1971	8142	3458	29.8	484	1318	2.72
1972	4680	2270	32.7	210	641	3.05
1973						
1974	2025	377	15.7	50	130	2.60
1975	744	405	35.2	51	149	2.92
1976	1923	324	14.4	207	567	2.74
1977	996	683	40.7	108	302	2.80
1978	1395	495	26.2	62	188	3.03
1979	841	113	11.8	117	329	2.81
1980	1446	454	23.9	40	93	2.33
1981	1527	747	32.8	235	750	3.19
1982	1653	140	7.8	32	85	2.66
1983	1326	543	29.1	192	612	3.19
1984	2753	795	22.4	80	230	2.88
1985	2245	503	18.3	125	354	2.83
1986	3283	1381	29.6	266	794	2.98
1987	1706	808	32.1	305	993	3.26
1988	3884	1242	24.2	200	616	3.08
1989	3811	1136	23.0	145	455	3.14
1990	4002	1068	21.1	97	309	3.19
1991	8599	2882	25.1	147	480	3.27
1992	9291	1347	12.7	151	451	2.99
1993	13976	2176	13.5	161	441	2.74
1994	4658	792	14.5	301	702	2.33
1995	6434	1618	20.1	99	319	3.22
1996	3128	631	16.8	125	330	2.64
1997	1345	144	9.7	43	114	2.65
1998	1595	432	21.3	97	239	2.46
1999	2395	527	18.0	82	200	2.44
2000	1870	410	18.0	93	192	2.06
2001	1232	228	15.6	42	103	2.45
2002	4789	1842	27.8	260	696	2.68
2003	5744	785	12.0	218	439	2.01
2004	4600	1288	21.9	235	568	2.42
2005	2844	1139	28.6	131	365	2.79
Mean ^c		23.2			2.80	
% Change from:						
2004		31%			15%	
Mean		23%			0%	

^a Data supplied by Izembek National Wildlife Refuge, USGS Alaska Science Center, and USFWS Region 7 Migratory Bird Management.

^b 1979, 1981, and 1987 data include Izembek Lagoon and Alaska Peninsula; 1984-1995 data include Izembek Lagoon and Nelson Lagoon.

^c Mean excludes 2005.

Table 7. Historical fall productivity records (from aerial photos) for emperor geese on the Alaska Peninsula, 1985-2005.^a

Year	No. Photos	No. Birds Aged in Photos	% Juvenile ^b
1985	155	3193	16.5
1986	311	6380	25.4
1987	703	10177	22.8
1988	483	11180	24.4
1989	390	12718	21.9
1990	474	13541	24.1
1991	412	14569	23.2
1992	403	14832	15.5
1993	255	5735	24.2
1994	479	16881	22.8
1995	361	11664	25.5
1996	182	10793	17.8
1997	205	11138	11.1
1998	336	16544	11.8
1999	392	13489	17.8
2000	263	7748	11.2
2001	365	11186	11.5
2002	402	6458	17.8
2003	421	8686	9.3
2004	370	6237	11.1
2005	500	6563	18.5
Mean ^c			18.3
% Change from:			
2004			67%
Mean			1%

^a Data supplied by USFWS Migratory Bird Management, Anchorage and Fairbanks, AK.

^b Mean of % juvenile in each of 7 lagoons from photo samples, weighted by the population counts of those lagoons from an independent aerial survey.

^c Mean excludes 2005.

Table 8. Historical productivity data for dusky Canada geese on the Copper River Delta, AK, from July aerial surveys, 1971-2005.^a

Year	% Juvenile	No. Geese Sampled
1971	16.2	5717
1972	10.6	8193
1973	36.0	5873
1974	51.4	8199
1975	17.9	8990
1976	24.2	7092
1977	44.3	----
1978	24.8	----
1979	16.0	12700
1980	23.7	7500
1981	17.9	8740
1982	23.7	8473
1983	15.0	7740
1984	18.3	11913
1985	3.7	13780
1986	10.7	13309
1987	9.8	12448
1988	22.5	6917
1989	8.6	6114
1990	23.5	5530
1991	21.5	7098
1992	23.1	7633
1993	5.0	4542
1994	5.7	6977
1995	3.9	5818
1996	21.7	6329
1997	10.5	6253
1998	11.7	4919
1999	14.7	4156
2000	24.1	4397
2001	25.4	3165
2002	30.5	3708
2003	7.2	5929
2004	27.8	5678
2005	11.8	5364
Mean ^b	19.2	
% Change from:		
2004	-58%	
Mean	-38%	

^a Data supplied by Alaska Department of Fish and Game.

^b Mean excludes 2005.