

15001 - 15006

Deep Drill Core

242 cm




Figure 1: Photo of Apollo 15 CDR setting up the deep drill. NASA AS15-87-11847.

Introduction

The Apollo 15 deep drill (15001 – 15006) was taken at the ALSEP site – about 120 meters from the LM. There were no small craters in the vicinity (figure 1). The regolith is on top of a mare surface, but the highlands are only about 5 – 10 km away. The combined weight of the core was 1333 grams, and it was found to contain materials from both the mare and the highlands.

Trench soils 15030, 15040 and 15014 were collected about 10 meters away, and are better analyzed. 15030 and 15014 from the bottom of the trench (depth about 30 cm) might be comparable to material from 15005, which 15040 should be compared with 15006. Pre-mission, it was expected that the deep drill might obtain material from the apparent rays from Autolycus and/or Aristillus as well as detritus from the degradation of the Apennine Front.

This was the first use of the rotary-percussion drilling mechanism, but it obtained a core 2.42 meters long (2.04 cm diameter). However, it was very difficult to extract from its hole. Also segments 15001, 15002 and 15003 would not separate on the Moon, so they were brought back as one long segment and broken down in the LRL.

The main result of this core is that it has remained in place, slowing cooking in neutrons, for about half a billion years. The bottom of the core could not have been deposited more than 750 m.y. ago nor less than 420 m.y. ago (Curtis and Wasserburg 1977). That being said, it is also clear that the top of the core has become mature due to micrometeorite bombardment, because of the abundance of agglutinate particles. However, cosmic ray track studies, solar wind implanted rare gasses and other studies indicate that core material had previous exposure to solar wind and cosmic rays, and the record that can be read from this core still requires more work and interpretation.




Figure 2: Maturity as function of depth in Apollo 15 deep drill core as determined by percentage agglutinate and I_s/FeO (from Heiken et al. 1976). Maturity data is tabulated in Morris (1976). Note that the top of the core is very mature, while the bottom is very immature.

This core was found to have significant Pb contamination (Silver 1972). There is also a concern that the ends of the core segments were exposed to moisture.

Petrography

Heiken et al. (1973, 1976) give a complete modal analysis of 34 samples along the length of the Apollo 15 deep drill. They found that the agglutinate content varied from 65% at the top to about 6% at depth. This was found to be consistent with the maturity index I_s/FeO (figure 2). If you ignore the agglutinates and re-average their mode you get the percentages of rock types in table B. Morris (1976) gives the magnetic maturity data and Morris (1978) discusses the evidence for gardening at the top of the deep drill core.

Walker and Papike (1981) and Papike et al. (1982) studied the thin sections prepared from 15003 and found that the deep drill was enriched in highlands materials relative to drive tube 15011 – 15010.

Basu and Bower (1976, 1977) studied fragments of KREEP basalt and determined the overall provenance of the materials in the Apollo 15 deep drill core. They concluded that the core has highland and mare materials in a ratio approximately 60:40, that the mare component generally increases from bottom to top in the core, quartz-normative basalts are nearly twice as abundant as olivine basalts, there were multiple sources of highland material, the highland component is similar to the highland regolith, KREEP basalt flow units were from depth, excavated and delivered as rays and that the green glass may be of impact origin.

Drake (1974) produced a catalog of the rock particles extracted from this core during dissection. Lindstrom et al. (1977) studied the mineralogy and chemistry of some of these particles.

Modal analysis was performed by Heiken et al. (1973 and 1975), and in detail for thin sections of 15003, by Walker and Papike (1981). These are summarized in

Table A: Petrologic mode of A15 drill core.*(90-150 microns) from Heiken et al. 1973*

	15001	15001	15002	15002	15003	15003	15004	15004	15005	15005	15006	15006
depth cm	242	241	203	202	162	161	123	122	82	82	40	39
agglutinates	25	15	35	40	27	25	46	34	24	34	51	41
breccias												
vitric	4.2	8.6	3.1	7.5	5.6	9	0.5	3.6	7.9	9	4.5	5.2
meta.	7.8	5.2	4.1	3.5	4.2	3.5	2.5	2.1	2.1	4.9	5.7	1.7
Basalt	10	5.7	14	6	2.8	5.5	6	3.6	7.9	4.9	2.5	6.4
Anorthosite							0.5	0.7	0.5		0.6	0.6
plagioclase	6.6	12.6	8.2	11.5	5.6	9	5.5	7.1	9.9	8.2	5.7	4
orthopyx.	17	13	10	8	10	15	11	14	9.9	6.6	11	11
clinopyx.	17	22	11	13	18	20	19	24	28	20	13	14
ilmenite		0.6	1	0.5		0.5		2.1		0.8		1.2
Glass												
brown	1.8	6.3	5.2	3	7	3.5	1.5	4.3	2.1	3.3		3.5
colorless	5.4	9.2	4.1	4	16	9.5	4	4.3	7.3	5.6	1.9	10.4
green	2.4		3.1	2							2.5	

Table B: Average components of A15 drill core.*from Heiken et al. 1976*

	Average (%)	Range (%)
Basalt		
coarse-grained	9.8	4 to 15
intergranular	3.9	0 to 6
fine-grained	1.7	0 to 4
vitrophyric	1.6	0 to 4
KREEP basalt		0 to 3
Breccia		
brown (vitric)	16.7	6 to 30
medium-grade	4.3	1 to 8
with shock melt	0.9	0 to 4
Anorthosite	0.9	0 to 2
plagioclase	10.4	7 to 14
orthopyroxene	0.3	0.3 to 1
clinopyroxene	37.4	30 to 47
olivine	0.3	0.3 to 2
ilmenite	0.4	0.3 to 2
green glass	1.5	0 to 6

Table C: Modal analysis of 15003 (0.02 to 0.2 mm).*from Walker and Papike 1981 (appendix)*

	unit A		unit B				unit C								
Mare component	0.2		0.1	0.4	0.3	0.2	0.6	0.4	0.1	0.3	0.2	0.2	0.2	0.3	0.5
Highland component	0.3	0.40	0.2	0.2	0.5	0.4	0.3	0.4	0.4	0.3	1.9	0.1	1.1	0.3	0.2
Agglutinate	6.7	7.70	6.4	9.2	6.7	8	7.2	5	9.9	7.1	8.2	8.9	9.1	11	10
Minerals															
olivine		0.60	0.4	0.5	0.4	0.7	0.7		0.4	0.1	0.6	0.6	0.7	0.1	0.2
pyroxene	14.8	16.30	11	11.6	16.3	16.5	11.7	16.8	18.5	15.5	13.4	14.3	14.5	12.1	13.9
plagioclase	4.5	2.80	6.3	5.8	6.3	5.1	5.1	5	4.5	4.3	4.9	4.9	5.6	6.2	5.2
ilmenite	0.5	0.50	0.6	0.3	0.5	0.2	0.3	0.3	0.5	0.2	0.5	0.2	0.4	0.7	0.7
Glass															
green	0.9	1.20	1.8	1.2	0.3	0.6	0.9	1.3	0.8	0.7	1.4	1	1.5	0.5	0.7




Figure 3: Chemical composition of A15 deep drill.




Figure 4: Normalized rare-earth-element diagram for three samples of A15 deep drill core (see Table 1).

tables A, B and C. Walker and Papike subdivide 15003 into three units based on different percentages of agglutinates, mare component and highland component (table C).

Chemistry

While there is a lot of other data on this core, the chemical composition as function of depth is sparse, and one needs to look at the composition of the nearby trench samples, 15030 and 15040. Helmke et al. (1973) and Gold et al. (1977) appear to be the only ones who reported complete analyses of the deep drill. Wiesmann and Hubbard (1975) tabulated a few elements from the bottom of each segment and have reported a somewhat complete analysis of the bottom of the drill (table 1). Chou and Pearce (1979) determined the composition of different grain sizes. Reed and Jovanovic (1972) determined Ru, Os, Hg and U.

Gold et al. (1977) noted that there was a distinct difference in chemical composition between 15003 and 15001 (figure 3 and 4). Evensen et al. (1974) tried to




Figure 5: Carbon content determined by Wszolek et al. (1973), maturity from Morris (1976).

characterize the exotic component by analyzing size fractions (see table 2). Wszolek et al. (1973) and Wszolek and Burlingame (1973) determined the carbon chemistry of the deep drill core (figure 5).

Ma et al. (1976) and Lindstrom et al. (1977) analyzed some of the particles extracted from the core (some in table 4, figure 7).

Radiogenic age dating

Pepin et al. (1974) reported the K-Ar age of soils from the core (3 b.y) - but that's not the age of the core!

Cosmogenic isotopes and exposure ages

Russ et al. (1972) and Pepin et al. (1974) showed that cosmic ray produced spallation nuclides are all smooth functions of depth. This indicates that the depositional history of the drill core is coherent and relatively simple.

Hubner et al. (1973), Bogard et al. (1973) and Bogard and Hirsch (1975) found that solar wind components were present throughout the length of the core. Bogard showed that ^4He and $^4\text{He}/^3\text{He}$ were relatively constant.

Russ et al. (1972) and Curtis and Wasserburg (1977) determined the isotopic composition of soils and fresh particles along the length of the core and found substantial variation (figure 8, 9 and 10). They were able to model the neutron fluence that produced these variations and concluded that the bottom of the core "could not have been deposited more than 750 m.y. ago, nor less than 420 my. ago".




Figure 6: Chemical composition as function of depth for Apollo 15 drill core (from Helmke et al. 1972).




Figure 7: Normalized rare-earth-element composition of small rocks found in A15 deep drill (Lindstrom et al. 1977).

Cosmic ray induced profiles of ^{22}Na and ^{26}Al activity (figure 11) were determined by Rancitelli et al. (1975) and Fruchter et al. (1976), ^{53}Mn by Nishiizumi et al. (figure 12) and ^{14}C by Jull et al. (1998).

Other Studies

Lindsay (1973), Heiken et al. (1973, 1976) and Graf (1993) reported detailed grain size data along the length

of the A15 deep drill core. Morris (1976) tabulates the maturity index and Morris (1977) studied the variation with grain size (figure 13).

Smith et al. (1973) and Becker and Clayton (1977) found that nitrogen from the solar wind was considerably lighter in the past using the Apollo 15 deep drill samples.




Figure 8: Variation in the isotopic composition of Gd due to adsorption of neutrons produced by cosmic rays (from Curtis and Wasserburg 1977).




Figure 9: Neutron flux as function of depth of soil sample in Apollo 15 deep drill (Curtis and Wasserburg 1977).

Phakey et al. (1972), Crozaz et al. (1972), Bhandari et al. (1972, 1973), Fleischer et al. (1973, 1974) and Goswami and Lal (1977) determined the density of cosmic ray induced nuclear tracks in mineral grains as a function of depth in the A15 drill core (figure 14). Goswami and Lal (1977) claimed that the data indicated cyclic variation with an apparent periodicity of ~ 200-300 m.y.

Lindsay and Srnka (1975) reported evidence for periodic fluctuation in the micrometeorite bombardment of the moon based on detailed petrography of samples from 120 to 240 cm depth in the A15 drill core. They even proposed that this was because the earth-moon system passed thru dense interstellar clouds at the time of deposition of this core!

These imaginative studies need verification.

Processing

The three deepest segments of the deep drill (15001, 15002 and 15003) would not separate on the Moon and had to be brought back in a bag linked together (Duke and Nagle 1974). *“The ends were plugged on the lunar surface and tapped in the LM. The exteriors of the linked core sections were exposed to atmospheres of the LM and CM cabins and had water spots on them (probably caused by sea water splashing into the cabin through an open door after splashdown). The remaining sections were protected by the nylon bag, but the exteriors were exposed to the air in the cabin.”*




Figure 10: Neutron flux for soil and pebbles in A15 deep drill core (Curtis and Wasserburg 1977).

Early allocations were made from the junctions of the core segments (3.75 g each) and portions of this soil was used for biomedical experiments. Other portions were used for initial allocations. The segments were then mounted in a cradle and split lengthwise with a milling machine (Heiken et al. 1972). Then the segments were transferred to another cabinet, the split core top removed, and the sample dissected into 5 mm intervals along the length down to approximately two-thirds tube diameter. For thermoluminescence studies,




Figure 11: ^{22}Na and ^{26}Al profiles in 15006 showing disturbance at top of Apollo 15 drill (Fruchter et al. 1977).

representative samples for each stratigraphic unit were collected under “red dark-room light”. After dissection, the remaining third of the core was removed to a laminar flow bench where it was impregnated with n-butyl methacrylate to make “peels”. The peels were intended to maintain grain orientation and permanent stratigraphic record.

15003 is the only segment that was encapsulated in epoxy and for which a continuous set of thin sections were prepared.

According to Allton and Waltz (1977) “the Apollo 15 drill core was completely filled and its scale is straightforward and accurately represents in situ lunar conditions.”

Silver (1972) found extensive Pb contamination from the joints in the deep drill.




Figure 12: ^{53}Mn as function of depth at top of Apollo 15 deep drill (Nishiizumi et al. 1976).




Figure 13: Grain size dependence for A15 deep drill samples (Morris 1977).




Figure 14: Apparent cyclic variation in nuclear track density along length of A15 deep drill (Goswami and Lal 1977).

Table 1. Chemical composition of A15 drill-core.

reference weight	Wiesmann and Hubbard 1975						Gold et al. 1977			comparison Korotev87	
	15001	15002	15003	15004	15005	15006	15001	15001	15003	15030	15040
SiO ₂ %							46.4	44.7	47.3	(b) 30 cm	surface
TiO ₂	1.82	1.82	1.75	1.72	1.72	1.73	(a) 1.83	1.77	1.87	(b) 1.7	1.7
Al ₂ O ₃							10.8	10.8	14.5	(b) 14.1	14.2
FeO							18.7	18.1	14.9	(b) 15.9	14.5
MnO							0.23	0.23	0.21	(b) 0.19	0.19
MgO	10.3	10.1	10.1	10.4	10.3	10.1	(a) 11.5	11.9	9.8	(b) 12	11.5
CaO	9.93	10.1	10.5	10.3	10.5	9.93	(a) 8.5	8.3	10.5	(b) 9.6	11.6
Na ₂ O	0.48	0.55	0.51	0.57	0.53	0.48	(a) 0.4	0.34	0.45	(b) 0.41	0.44
K ₂ O	0.3	0.26	0.22	0.21	0.23	0.22	(a) 0.3	0.2	0.25	(b)	
P ₂ O ₅											
S %											
sum											
Sc ppm							31.5		28.9	(b) 31.6	28.5
V							151	147	129	(b) 120	110
Cr							3750	3640	2670	(b) 2840	2710
Co							54.8		41.2	(b) 46.4	46.3
Ni							382		124	(b) 203	252
Cu											
Zn											
Ga											
Ge ppb											
As											
Se											
Rb	8.13	7.17	6.11	5.89	6.39	5.94	(a)				
Sr	136	137	134	132	136	127	(a)			115	150
Y											
Zr	600						(a)			430	370
Nb											
Mo											
Ru											
Rh											
Pd ppb											
Ag ppb											
Cd ppb											
In ppb											
Sn ppb											
Sb ppb											
Te ppb											
Cs ppm										0.27	0.27
Ba	370	334	285	279	297	283	(a) 252		285	(b) 271	259
La	35						(a) 21.6	18.9	24.4	(b) 26.4	26.1
Ce	92.2						(a) 64.5		71.7	(b) 70	68
Pr											
Nd	57.8						(a) 35		48	(b) 41	36
Sm	16.4						(a) 11.2	10.1	13.2	(b) 12.6	12.4
Eu	1.7						(a) 1.13		1.45	(b) 1.36	1.4
Gd											
Tb							1.38		1.8	(b) 2.49	2.5
Dy	22.5						(a) 12		16	(b)	
Ho											
Er	13.5						(a)				
Tm											
Yb	11.8						(a) 7.2		8.4	(b) 8.8	8.3
Lu	1.73						(a) 1.13		1.36	(b) 1.32	1.25
Hf	17.8						(a) 8.2		9.81	(b) 11.8	9.8
Ta							0.97		1.26	(b) 1.24	1.18
W ppb											
Re ppb											
Os ppb											
Ir ppb										4.9	9.3
Pt ppb											
Au ppb										<4	3.9
Th ppm							1.72		3.4	(b) 4.8	4.3
U ppm	1.69	1.52	1.29	1.25	1.33	1.35	(a)			1.1	1.18

technique: (a) IDMS, (b) INAA

Table 2a. Chemical composition of A15 drill-core.

reference	Evensen74												
weight	15001		15002		15003		15004		15005		15006		
depth	241 cm		202		161		121		82		39		
size frac	>74	<16	>74	<16	>74	<16	>74	<16	>74	<16	>74	<16	(a)
Al2O3													
FeO													
MnO													
MgO													
CaO													
Na2O													
K2O	0.29	0.47	0.29	0.27	0.22	0.25	0.22	0.22	0.26	0.26	0.23	0.21	(a)
P2O5													
Se													
Rb	6.63	8.19	7.38	7.23	5.62	6.72		5.88	5.3	8.56	5.86	5.77	(a)
Sr	139	154	120	155	114	153	130	138	119	149	129	135	(a)
Y													
Zr													
Cs ppm													
Ba	377	383	337	355	249	297		268	340	316	273	302	(a)
La													

technique: (a) IDMS

Table 2b. Chemical composition of A15 drill.

reference	Nyquist 1973						
weight	15001	15002	15003	15004	15005	15006	
Se							
Rb	8.01	7.17	6.11	5.89	6.39	5.94	(a)
Sr	136	137	134	132	136	127	(a)
Y							

technique: (a) INAA


Table 3: Composition of A15 Deep Core.

	depth (cm)	K2O %	U ppm	Th ppm	author
15006	5.50	0.254	0.9	4.65	Fruchter 75
15006	9.00	0.24	1.06	4.58	Fruchter 75
15006	14.50	0.195	1.16	4.59	Fruchter 75
15006	21.50	0.22	1.13	4.33	Fruchter 75
15006	33.00	0.24	1.41	5.6	Fruchter 75
15006	39.00		1.27	4.84	Silver 73
15005	51.00	0.2	1.52	6.13	Fruchter 75
15005	54.00		1.38	5.21	Silver 73
15005	79.00	0.093	0.99	4.42	Fruchter 75
15005	79.00		1.37	5.16	Silver 73
15004	95.00	0.144	1.22	5.39	Silver 73
15004	119.00		1.33	5	Silver 73
15003	138.00	0.194	1.18	4.69	Fruchter 75
15003	159.00		1.26	4.85	Silver 73
15002	176.00		1.32	4.96	Silver 73
15002	184.00		1.34	5.24	Silver 73
15002	199.00		1.54	5.8	Silver 73
15001	216.00	0.37	1.89	7.35	Fruchter 75
15001	239.00		1.75	6.58	Silver 73

Table 4. Chemical composition of basaltic rocklets from 15003.

reference weight	Ma 76 mare basalts						Lindstrom 77 mare basalts				KREEP basalts			
SiO2														
TiO2	2	1.6	3.3	1.9	1.6	2.6	1.8	1.6	2	1.9	2.6	2.2	(a)	
Al2O3	10.7	8	9.9	10.7		9.2	9.2	8	10.7	10.7	15.7	15.5	(a)	
FeO	18.4	22.1	21.1	20.5	17.8	22.9	23.3	22.1	18.4	20.4	10.2	10.5	(a)	
MnO	0.268	0.26	0.274	0.262	0.246	0.278	0.27	0.26	0.27	0.26	0.14	0.14	(a)	
MgO	8.9	10	9.4	11.1		11.9	18.9	10	8.9	11.1	9.7	6.4	(a)	
CaO	10.2	8.6	10.5	9.8	10.7	9.7	8	8.6	10.2	9.8	10.2	9.3	(a)	
Na2O	0.25	0.249	0.413	0.298	0.22	0.384	0.241	0.209	0.249	0.295	0.82	0.91	(a)	
K2O	0.047	0.037	0.035	0.039		0.011	0.033	0.037	0.047	0.039	0.55	0.69	(a)	
P2O5														
S %														
sum	<i>note: there are many more analyses in these pubs than can be reentered here.</i>													
Sc ppm	43	38	43	42	35	41	28.5	38.1	42.6	41.9	21.8	23.4	(a)	
V	240	220	200	250		230	210	220	240	250	58	49	(a)	
Cr	4215	5029	3181	4413	4100	4262	5650	5035	4420	4415	1970	1860	(a)	
Co	42	66	45	56	55	54	66.5	66.1	42.1	55.8	19	17.8	(a)	
Ni														
Cu														
Zn														
Ga														
Ge ppb														
As														
Se														
Rb														
Sr														
Y														
Zr											800	1020	(a)	
Nb														
Mo														
Ru														
Rh														
Pd ppb														
Ag ppb														
Cd ppb														
In ppb														
Sn ppb														
Sb ppb														
Te ppb														
Cs ppm														
Ba											690	950	(a)	
La	5.4	4.6	4.9	3.7	2.6		4.02	4.64	5.39	3.67	70.7	84	(a)	
Ce	17.5	15.6	15.7	13.6	6.7		15	15.6	17.5	13.6	173	242	(a)	
Pr														
Nd											113	161	(a)	
Sm	3.6	3.1	3.6	2.9	2		2.82	3.13	3.59	2.9	34	40.7	(a)	
Eu	0.86	0.69	1	0.94	0.67		0.96	0.69	0.86	0.94	2.74	3.08	(a)	
Gd														
Tb	0.98	0.84	0.92	0.74	0.41		0.86	0.84	0.98	0.74	7.33	8.7	(a)	
Dy	5.3	3.9	6.4	3.5	2.5		5	4	5	4	43	47	(a)	
Ho														
Er														
Tm														
Yb	2.8	2	2.5	1.8			1.7	2	2.8	1.8	23	27	(a)	
Lu	0.44	0.33	0.36	0.32	0.26		0.3	0.33	0.44	0.32	3.5	4.2	(a)	
Hf	2.4	2.1	2.4	1.9			1.95	2.06	2.43	1.85	27.5	33.9	(a)	
Ta														
W ppb														
Re ppb														
Os ppb														
Ir ppb														
Pt ppb														
Au ppb														
Th ppm										0.49	0.24	12	15.6	(a)
U ppm														

technique: (a) INAA




15003,655



or ,317 ?



W1



*Note:
The top of 15003
was 121.8 cm
below the lunar
surface (a la Allton
and Waltz 1977).*




epoxy
encapsulated
core









15001-6 References

(note: *There is a vast literature on the lunar drill cores, which can not all be listed at once. Please excuse the complier for his brevity.*)

Allton J.H. and Waltz S.R. (1980) Depth scales for Apollo 15, 16 and 17 drill cores. Proc. 11th Lunar Planet. Sci. Conf. 1463.

Arnold J.R. (1975) A Monte-Carlo model for the gardening of the lunar regolith. *The Moon* 13, 159-172.

Basu A. and Bower J.F. (1976) Petrography of KREEP basalt fragments from Apollo 15 soils. Proc. 7th Lunar Sci. Conf. 659-678.

Basu A. and Bower J.F. (1977) Provenance of Apollo 15 deep drill core sediments. Proc. 8th Lunar Sci. Conf. 2841-2867.

Becker R.H. and Clayton R.N. (1977) Nitrogen isotopes in lunar soils as a measure of cosmic-ray exposure and regolith history. Proc. 8th Lunar Sci. Conf. 3685-3704.

Bhandari N., Goswami J.W. and Lal D. (1972) Apollo 15 regolith: a predominantly aggregation or mixing model? *In* The Apollo 15 Samples, 336-341.

Bhandari N., Goswami Jitendra and Lal D. (1973) Surface irradiation and evolution of the lunar regolith. Proc. 4th Lunar Sci. Conf. 2275-2290.

Bogard D.D. and Nyquist L.E. (1973a) Noble gases in the Apollo 15 drill cores? *In* The Apollo 15 Samples, 342-346.

Bogard D.D. and Nyquist L.E. (1973b) ⁴⁰Ar/³⁹Ar variations in Apollo 15 and 16 regolith. Proc. 4th Lunar Sci. Conf. 1975-1985.

Bogard D.D., Nyquist L.E., Hirsch W.C. and Moore D.R. (1973) Trapped solar and cosmogenic noble gas abundances in Apollo 15 and 16 deep drill samples. *Earth Planet. Sci. Lett.* 21, 52-69.

Bogard D.D. and Hirsch W.C. (1975) Noble gas studies on grain size separates of Apollo 15 and 16 deep drill cores. Proc. 6th Lunar Sci. Conf. 2057-2083.

Carrier D. W., Johnson S.W., Carrasco L. and Schmidt R. (1973) Core sample depth relationships: Apollo 14 and 15. Proc. 3rd Lunar Sci. Conf. 3213-3221.

Carrier David (1974) Apollo drill core depth relationships. *The Moon* 10, 183.

Carr M.H. and Meyer C.E. (1974) The regolith at the Apollo 15 site and its stratigraphic implications. *Geochem. Cosmochim. Acta* 38, 1183-1197.

Chou C. and Pearce G.W. (1979) Apollo 15 deep drill core: Trace element and metallic abundances in size fractions of sample 15002,170. Proc. 10th Lunar Planet. Sci. Conf. 1321-1332.

Clanton U.S., McKay D.S., Taylor R.M. and Heiken G.H. (1973) Relationships of exposure age to size distribution and particle types in the Apollo 15 drill core. *In* The Apollo 15 Samples, 54-56.

Crozaz G., Drozd R., Hohenberg C.M., Hoyt H.P., Ragan D., Walker R.M. and Yuhas D. (1972) Solar flare and galactic cosmic ray studies of Apollo 14 and 15 samples. Proc. 3rd Lunar Sci. Conf. 2917-2931.

Curtis D.B. and Wasserburg G.J. (1977) Stratigraphic processes in the lunar regolith. Proc. 8th Lunar Sci. Conf. 3575-3593.

DesMaris D.J., Hayes J.M. and Meinschein W.G. (1974) The distribution in lunar soil of hydrogen released by pyrolysis. Proc. 5th Lunar Sci. Conf. 1811-1822.

Drake M.J. (1974) Apollo 15 deep-drill-core: Classification, description and inventory of separated particles. JSC curator catalog

Duke M.B. and Nagle J.S. (1976) Lunar Core Catalog (*and many supplements*). JSC 09252. 242 pages.

Evensen N.M., Murthy V.R. and Coscio M.R. (1974) Provenance of KREEP and the exotic component: Elemental and isotopic studies of grain size fractions in lunar soils. Proc. 5th Lunar Sci. Conf. 1401-1417.

Epstein S. and Taylor H.P. (1975) Investigation of the carbon, hydrogen, oxygen and silicon isotope and concentration relationships on the grain surfaces of a variety of lunar soils and in some Apollo 15 and 16 core samples. Proc. 6th Lunar Sci. Conf. 1771-1798.

Fleischer R.L. and Hart H.R. (1973) Particle track record in Apollo 15 deep core from 54 to 80 cm depths. *In* The Apollo 15 Samples, 371-373.

Fleischer R.L. and Hart H.R. (1973) Particle track record in Apollo 15 deep core from 54 to 80 cm depth. *Earth Planet. Sci. Lett.* 18, 420-426.

Fleischer R.L., Hart H.R. and Giard W.R. (1974) Surface history of lunar soil and soil columns. *Geochem. Cosmochim. Acta* 38, 365-380.

- Frick U., Baur H., Funk H., Phinney D., Schafer C., Schultz L. and Signer P. (1973) Diffusion properties of light noble gases in lunar fines. Proc. 4th Lunar Sci. Conf. 1987-2002.
- Fruchter J.S., Rancitelli L.A. and Perkins R.W. (1975) Primordial radionuclide variation in the Apollo 15 and 17 deep core samples and in Apollo 17 igneous rocks and breccias. Proc. 6th Lunar Sci. Conf. 1399-1405.
- Fruchter J.S., Rancitelli L.A. and Perkins R.W. (1976) Recent and long-term mixing of the lunar regolith based on ²²Na and ²⁶Al measurements in Apollo 15, 16 and 17 deep drill stems and drive tubes. Proc. 7th Lunar Sci. Conf. 27-39.
- Fruchter J.S., Rancitelli L.A., Laul J.C. and Perkins R.W. (1977) Lunar regolith dynamics based on analysis of the cosmogenic radionuclides ²²Na, ²⁶Al and ⁵³Mn. Proc. 8th Lunar Sci. Conf. 3595-3605.
- Fryxell R. and Heiken G. (1974) Preservation of lunar core samples: Preparation and interpretation of three-dimensional stratigraphic sections. Proc. 5th Lunar Sci. Conf. 935-966.
- Gold T., Bilson E. and Baron R.L. (1974) Optical properties of the Apollo 15 deep core samples. Proc. 5th Lunar Sci. Conf. 2355-2359.
- Gold T., Bilson E., Baron R.L., Ali M.Z. and Ehmann W.D. (1977) Chemical and optical properties at the Apollo 15 and 16 sites. Proc. 8th Lunar Sci. Conf. 3633-3643.
- Gose W.A., Pearce G.W. and Lindsay J.F. (1975) Magnetic stratigraphy of the Apollo 15 deep drill core. Proc. 6th Lunar Sci. Conf. 3071-3080.
- Goswami J.N. and Lal D. (1977) Particle track correlation studies in lunar soils: Possible long-term periodic fluctuations in ancient meteoritic flux at 1 A.U. Proc. 8th Lunar Sci. Conf. 813-824.
- Graf John (1993) Lunar Soils Grain Size Catalog. NASA Ref. Pub. 1265.
- Heiken G.H., Duke M., Fryxell R., Nagle S., Scott R. and Sellers G.A. (1972) Stratigraphy of the Apollo 15 drill core. NASA TM X-58101.
- Heiken G.H., Duke M., McKay D.S., Clanton U.S., Fryxell R., Nagle S., Scott R. and Sellers G.A. (1973) Preliminary stratigraphy of the Apollo 15 drill core. Proc. 4th Lunar Sci. Conf. 191-213.
- Heiken G.H., Morris R.V., McKay D.S. and Fruland R.M. (1976) Petrographic and ferromagnetic resonance studies of the Apollo 15 deep drill. Proc. 7th Lunar Sci. Conf. 93-111.
- Heiken G.H., Vaniman D.T. and French B.M. (1992) The Lunar Sourcebook. Cambridge Press – see pages 337-342
- Helmke P., Telander K.H., Weiss C.K. and Haskin L.A. (1973) Trace elements in fines from the Apollo 15 deep drill core. Geochim. Cosmochim. Acta 37, 706-708.
- Housley R.M., Cirlin E.H., Goldberg I.B., Crowe H., Weeks R.A. and Perhac R. (1975) Ferromagnetic resonance as a method of studying the micrometeorite bombardment history of the lunar surface. Proc. 6th Lunar Sci. Conf. 3173-3186.
- Hubner W., Heymann D. and Kirsten T. (1973) Inert gas stratigraphy of Apollo 15 drill core sections 15001 and 15003. Proc. 4th Lunar Sci. Conf. 2021-2036.
- Jovanovic S. and Reed G.W. (1979) Regolith layering processes based on studies of low-temperature volatile elements in Apollo core samples. Proc. 10th Lunar Sci. Conf. 1425-1435.
- Jull A.J.T., Cloudt S., Donohue D.J., Sisterson J.M., Reedy R.C. and Masarik J. (1998) ¹⁴C depth profiles in Apollo 15 and 17 cores and lunar rock 68815. Geochim. Cosmochim. Acta 62, 3025-3036.
- Langevin Y. and Maurette M. (1978) Plausible depositional histories for the Apollo 15, 16 and 17 drill core tubes. Proc. 9th Lunar Planet. Sci. Conf. 1765-1786.
- Lindsay John F. (1973) Evolution of lunar soil grain-size and shape parameters. Proc. 4th Lunar Sci. Conf. 215-224.
- Lindsay J. and Srnka L.B. (1975) Galactic dust lanes and lunar soils. Nature **257**, 776-777.
- Lindstrom M.M., Nielsen R.L. and Drake M.J. (1977) Petrology and geochemistry of lithic fragments separated from the Apollo 15 deep-drill core. Proc. 8th Lunar Sci. Conf. 2869-2888.
- McKay D.S., Heiken G., Basu A., Blanford G., Simon S., Reedy R., French B.M. and Papike J.J. (1992) The lunar regolith. In Lunar Source Book: a users guide to the moon, 285-356.
- Ma M-S., Murali A.V. and Schmitt R.A. (1976) Chemical constraints for mare basalt genesis. Proc. 7th Lunar Sci. Conf. 1673-1695.
- Mitchell et al. (1974) Apollo Soil Mechanics Experiment S-200. Space Science Laboratory Series 15, Issue 7, UC Berkeley
- Modzeleski J.E., Modzeleski V.E., Nagy L.A., Nagy B., Hamalton P.B., McEwan W.S. and Urey H.C. (1972) Carbon

- compounds in Apollo 15 lunar samples. *In* The Apollo 15 Samples, 311-315.
- Morris R.V. (1976) Surface exposure indices of lunar soils: A comparative FMR study. Proc. 7th Lunar Sci. Conf. 315-335.
- Morris R.V. (1977) Origin and evolution of the grain-size dependence of the concentration of fine-grained metal in lunar soils: The maturation of lunar soils to a steady stage. Proc. 8th Lunar Sci. Conf. 3719-3747.
- Morris R.V. (1978) In situ reworking (gardening) of the lunar surface: Evidence from the Apollo cores. Proc. 9th Lunar Planet. Sci. Conf. 1801-1811.
- Nagle J.S. and Duke M.D. (1974) Stabilization of lunar core samples. Proc. 5th Lunar Sci. Conf. 967-973.
- Nishiizumi K., Imamura M., Honda M., Russ G.P., Kohl C.P. and Arnold J.R. (1976) ⁵³Mn in the Apollo 15 and 16 drill stems: Evidence for surface mixing. Proc. 7th Lunar Sci. Conf. 41-54.
- Nyquist L.E., Hubbard N.J., Gast P.W., Bansal B.M., Wiesmann H. and Jahn B. (1973) Rb-Sr systematics for chemically defined Apollo 15 and 16 materials. Proc. 4th Lunar Sci. Conf. 1823-1846.
- Papike J.J., Simon S.B. and Laul J.C. (1982) The lunar regolith: Chemistry, mineralogy and petrology. *Rev. Geophys. Space Phys.* 20, 761-826.
- Pepin R.O., Basford J.R., Dragon J.C., Coscio M.R. and Murthy V.R. (1974) Rare gases and trace element in Apollo 15 drill core fines: Depositional chronologies and K-Ar ages and production rates of spallation-produced ³He, ²¹Ne and ³⁸Ar versus depth. Proc. 5th Lunar Sci. Conf. 2149-2184.
- Phakey P.P., Hutcheon I.D., Rajan R.S. and Price P.B. (1972) Radiation effects in soils from five lunar missions. Proc. 3rd Lunar Sci. Conf. 2905-2915.
- Preliminary Science Reports, A15 NASA-SP 289.
- Rancitelli L.A., Fruchter J.S., Felix W.D., Perkins R.W. and Wogman N.A. (1975) Cosmogenic isotopic production in Apollo deep-core samples. Proc. 6th Lunar Sci. Conf. 1891-1899.
- Reed G.W. and Jovanovic S. (1973) Trace element comparisons between mare and Apennine-Front nonmare samples. *In* The Apollo 15 Samples, 247-249.
- Russ C.P., Burnett D.S. and Wasserburg G.J. (1972) Lunar neutron spectroscopy. *Earth Planet. Sci. Lett.* 15, 172-186.
- Silver L.T. (1973) Uranium-Thorium-Lead isotopes and the nature of the Mare surface debris at Hadley-Apennine. *In* The Apollo 15 Samples, 388-390.
- Smith J.W., Kaplan I.R. and Petrowski C. (1973) Carbon, nitrogen, sulfur, helium, hydrogen and metallic iron in Apollo 15 drill stem fines. Proc. 4th Lunar Sci. Conf. 1651-1656.
- Sullivan Tom (1994) Catalog of Apollo Experiment Operations. NASA Ref. 1317
- Swann G.A., Hait M.H., Schaber G.G., Freeman V.L., Ulrich G.E., Wolfe E.W., Reed V.S. and Sutton R.L. (1971) Preliminary descriptions of Apollo 15 sample environments. USGS Interagency Report 36.
- Swann G.A., Bailey N.G., Batson R.M., Freeman V.L., Hait M.H., Head J.W., Holt H.E., Howard K.A., Irwin J.B., Larsson K.B., Muehlerger W.R., Reed V.S., Rennilson J.J., Schaber G.G., Scott D.R., Silver L.T., Sutton R.L., Ulrich G.E., Ulrich G.E., Wilshire H.G. and Wolfe E.W. (1972) Preliminary Geologic Investigation of the Apollo 15 landing site. *In* Preliminary Science Reports, A15 NASA-SP 289.
- Taylor G.R. and Wooley B.C. (1973) Evaluation of lunar samples for the presence of viable organisms. Proc. 4th Lunar Sci. Conf. 2267-2274.
- Walker R.J. and Papike J.J. (1981) The Apollo 15 regolith: Comparative petrology of drive tube 15010/15011 and drill section 15003. Proc. 12th Lunar Planet. Sci. Conf. 485-508.
- Wiesmann H. and Hubbard N. (1975) A compilation of the lunar sample data generated – (unpublished)
- Wszolek P.C. and Burlingame A.L. (1973) Carbon chemistry of the Apollo 15 and 16 deep drill cores. Proc. 4th Lunar Sci. Conf. 1681-1692.
- Wszolek P.C., Jackson R.F. and Burlingame A.L. (1973) Carbon chemistry of the Apollo 15 deep drill stem and a glass-rich sample related to the uniformity of the regolith and lunar surface processes. *In* The Apollo 15 Samples, 324-326.