U.S. DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY

Prepared for the
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

GEOLOGIC INVESTIGATION SERIES I–2650
ATLAS OF MARS: THAUMASIA REGION
SHEET 2 OF 3

PALEOTECTONIC MAP OF THE THAUMASIA REGION, MARS

By
James M. Dohm, Kenneth L. Tanaka, and Trent M. Hare
2001





Contact—Dashed where approximately located or gradational

Fault or graben—Bar and ball on downthrown side of fault; dotted where buried

Scarp—Line marks top of slope; barb points downslope. Forms contact in places

Mare-type (wrinkle) ridge—Symbol on ridge crest; dashed where buried

Subdued mare-type (wrinkle) ridge

Broad (>3 km wide), nearly flat-topped ridge

Narrow (<2 km wide), sharp-crested linear ridge

Narrow (<3 km wide), subdued ridge

Depression or caldera

Crater rim crest

Crater central peak

Crater central pit

Center of figure

INTRODUCTION

The geology of the Thaumasia region (fig. 1) includes a wide array of rock materials, depositional and erosional landforms, and tectonic structures. The region is dominated by the Thaumasia plateau, which includes central high lava plains ringed by highly deformed highlands; the plateau may comprise the ancestral center of Tharsis tectonism (Frey, 1979; Plescia and Saunders, 1982). The extensive structural deformation of the map region, which includes the largest rift system on Mars in both complexity and extent, occurred during the Noachian and Hesperian periods (Tanaka and Davis, 1988; Scott and Dohm, 1990a). The deformation produced small and large extensional and contractional structures (fig. 2) that resulted from stresses related to the formation of Tharsis (Frey, 1979; Wise, 1980; Frey, 1979; Plescia and Saunders, 1982; Banerdt and others, 1982, 1992; Watters and Maxwell, 1986; Tanaka and Davis, 1988; Francis, 1988; Phillips, 1991; Schultz and Tanaka, 1994), from magmatic-driven uplifts, such as at Syria Planum (Tanaka and Davis, 1988; Dohm and others, 1998, Dohm and Tanaka, 1999) and central Valles Marineris (Dohm and others, 1998, Dohm and Tanaka, 1999), and from the Argyre impact (Wilhelms, 1973; Scott and Tanaka, 1986). In addition, volcanic, eolian, and fluvial processes have highly modified older surfaces in the map region. Local volcanic and tectonic activity often accompanied episodes of valley formation. Our mapping depicts and describes the diverse terrains and complex geologic history of this unique and scientifically exciting region.

The geologic sheet 1), paleotectonic (sheet 2), and paleoerosional (sheet 3) maps of the Thaumasia region were compiled on a Viking 1:5,000,000-scale digital photomosaic base. The base is a combination of four quadrangles: the southern portion of Syria Planum (MC-17), most of the southern half of Coprates (MC-18), a large part of Thaumasia (MC-25), and the northwest margin of Argyre (MC-26). The medium-resolution Viking images used for mapping and base preparation also formed the basis of the 1:2,000,000 scale subquadrangle maps. Earlier geologic maps of all or parts of the region include: (1) maps of the Phoenicis Lacus, Coprates, Thaumasia, and Argyre quadrangles at 1:5,000,000 scale based mainly on Mariner 9 images (respectively, Mausoley and others, 1978; McCauley, 1978; Scott and Carr, 1978; Hodges, 1980), (2) the Argyre area at 1:25,000,000 scale (Scott and Carr, 1978) compiled largely from the 1:5,000,000 scale geologic maps, (3) maps showing Sinai, Solis, and Syria Planums on Mars in both geologic and geomorphic units, based on Mariner 9 images (Scott, 1981; Scott and Tanaka, 1981a, b; Scott and others, 1981), (4) the map of the western equatorial region of Mars at 1:15,000,000 scale based on Viking images (Scott and Tanaka, 1986), and (5) the map of the Thaumasia region compiled by Scott and Carr (1978) and series (upper, middle, and lower divisions of systems) defined by Tanaka (1986). To determine the material and modification ages of the Thaumasia geologic units, we compiled crater statistics for 28 units of significant areal extents, which cover 96.6% of the map region. We measured all crater-rim diameters larger than 2 km using the Mars Digital Image Model (at 231 m/pixel; U.S. Geological Survey, 1993a, b) and calculated unit areas from digitally scanned contacts using GIS (tables 1 and 3). The crater statistics consist of total crater populations (including partly buried, degraded, and eroded, pristine impact craters), which may indicate emplacement ages, and superposed craters only, which indicate ages of most recent, major resurfacing (for example, see fig. 5A, *B*). We produced hundreds of crater counts that differ by as much as a factor of 10; we grouped the counts according to local or regional physiographic associations, and all outcrops together for each unit. The crater densities within physiographic provinces or groups vary somewhat; these ranges are typically incorporated in their time-stratigraphic age assignments.

The map units are categorized into channel-system and eolian materials, Valles Marineris interior deposits, western volcanic assemblage (flow and shield materials, Thaumasia Montes Formation, Syria Planum Formation), plateau and high-plains assemblage (plateau sequence, Thaumasia highland rift materials, dissected and subdued cratered terrain materials, subdivided material), circum-Argyre materials, and crater materials. Only materials of impact craters that have ejecta deposits nearly 50 km and greater are mapped; crater materials are assigned relative ages based on age relations with surrounding units and structures. For all units, relative-age assignments represent material emplacement ages; in some cases, erosional and depositional modification of the units somewhat obscures original formation. For example, dissected and etched materials include the eroded outcrops and superposed sediment deposits; therefore, the resulting age became the youngest age assigned to or mapped separately; their relative-age ranges reflect both materials.

Most unit contacts are fairly precise, especially where they are defined by ridge, embayment, and channel and buried relations. Transitional contacts, such as between fractured and dissected materials and adjacent geologic units are also sharp in most places. For example, these contacts commonly form along erosional scarps or partly buried tectonic structure. In other cases, because of a paucity of stratigraphic markers or poor image quality, contacts must be considered gradational and appear dashed on the map.

NOACHIAN SYSTEM

The Noachian System consists of ancient crustal materials formed during the period of early bombardment that produced a high density of impact craters and large impact basins. Cratering, along with water and wind erosion and tectonic deformation, have modified or destroyed many of the primary morphologic features of Noachian units, including a substantial part of the ancient crater population. Thus, Noachian cratered terrains are usually characterized by impact, erosional, and tectonic features that postdate the material. The Noachian materials exhibit a wide variety of impact breccia, including those that are buried under younger units. Eolian and fluvial sediments that mantle and subdue parts of the heavily cratered terrain. Materials of the Noachian System, which include geologic units that straddle the Noachian/Hesperian boundary (fig. 6A; table 4), cover nearly 40% of the Thaumasia region.

The hilly unit (unit Nph) at the base of the plateau sequence forms heavily cratered terrain consisting of high-standing plateaus, irregular topographic highs, prominent ridges, and highly degraded, ancient crater rims mainly in the Thaumasia highland, Daedalia, and Argyre provinces (fig. 2). Locally, tectonic and impact processes (Watters, 1993; Schultz and Tanaka, 1994; Craddock and others, 1990) may have contributed to the prominent relief of the hilly unit. Deep, steep-walled valleys, some as much as 200 km in diameter), and the center of the northeast-trending Thaumasia "bulge" or "rise," which extends for more than 3,000 km across the western equatorial region. This bulge forms a regional slope in the map region that trends southeast from Syria Planum at an elevation greater than 8 km to Argyre Planitia at an elevation less than 4 km (U.S. Geological Survey, 1989). The regional slope is interrupted by rugged highland terrain, which includes rifts, fossae, and fault systems, mesa-tains, and volcanoes. Differential wind and water erosion has eroded rock materials and structures along parts of the south edge of the Thaumasia plateau, resulting in well-defined valley networks, canyons, troughs, badlands, and mesas. The Argyre impact event includes the region, irregular topography produced by the colossal Argyre impact event includes relatively small basins and broad, scarp-bounded mesas.

We distinguish eleven geologic provinces within the Thaumasia region based on their structural, topographic, and geomorphic characteristics (fig. 1; table 2). These characteristics include broad canyons, highlands, large mountain ranges, volcanoes (fig. 4), lava and ridged plains, and surfaces that have been modified by tectonic, volcanic, and erosional processes.

STRATIGRAPHY

Geologic units were identified and mapped on individual, digitally enhanced Viking images and 1:2,000,000-scale photomosaic bases and compiled on the 1:5,000,000-scale digital photomosaic map base. Image resolution of the Thaumasia region ranges from 231 m > 500 m/pixel; those in the range of 50 to 300 m/pixel were most commonly used for mapping. The identification and classification of geologic units into groups and assemblages, where applicable, follows that of Scott and Tanaka (1986) in the region southeast of (1) maps of the Phoenicis Lacus, Coprates, Thaumasia, and Argyre quadrangles at 1:5,000,000 scale based mainly on Mariner 9 images (respectively, Mausorky and others, 1978; McCauley, 1978; Scott and Carr, 1978; Hodges, 1980), (2) the Argyre area at 1:25,000,000 scale (Scott and Carr, 1978) compiled largely from the 1:5,000,000 scale geologic maps, (3) maps showing Sinai, Solis, and Syria Planums on Mars in both geologic and geomorphic units, based on Mariner 9 images (Scott, 1981; Scott and Tanaka, 1981a, b; Scott and others, 1981), (4) the map of the western equatorial region of Mars at 1:15,000,000 scale based on Viking images (Scott and Tanaka, 1986), and (5) the map of the Thaumasia region compiled by Scott and Carr (1978) and series (upper, middle, and lower divisions of systems) defined by Tanaka (1986). To determine the material and modification ages of the Thaumasia geologic units, we compiled crater statistics for 28 units of significant areal extents, which cover 96.6% of the map region. We measured all crater-rim diameters larger than 2 km using the Mars Digital Image Model (at 231 m/pixel; U.S. Geological Survey, 1993a, b) and calculated unit areas from digitally scanned contacts using GIS (tables 1 and 3). The crater statistics consist of total crater populations (including partly buried, degraded, and eroded, pristine impact craters), which may indicate emplacement ages, and superposed craters only, which indicate ages of most recent, major resurfacing (for example, see fig. 5A, *B*). We produced hundreds of crater counts that differ by as much as a factor of 10; we grouped the counts according to local or regional physiographic associations, and all outcrops together for each unit. The crater densities within physiographic provinces or groups vary somewhat; these ranges are typically incorporated in their time-stratigraphic age assignments.

The map units are categorized into channel-system and eolian materials, Valles Marineris interior deposits, western volcanic assemblage (flow and shield materials, Thaumasia Montes Formation, Syria Planum Formation), plateau and high-plains assemblage (plateau sequence, Thaumasia highland rift materials, dissected and subdued cratered terrain materials, subdivided material), circum-Argyre materials, and crater materials. Only materials of impact craters that have ejecta deposits nearly 50 km and greater are mapped; crater materials are assigned relative ages based on age relations with surrounding units and structures. For all units, relative-age assignments represent material emplacement ages; in some cases, erosional and depositional modification of the units somewhat obscures original formation. For example, dissected and etched materials include the eroded outcrops and superposed sediment deposits; therefore, the resulting age became the youngest age assigned to or mapped separately; their relative-age ranges reflect both materials.

Most unit contacts are fairly precise, especially where they are defined by ridge, embayment, and channel and buried relations. Transitional contacts, such as between fractured and dissected materials and adjacent geologic units are also sharp in most places. For example, these contacts commonly form along erosional scarps or partly buried tectonic structure. In other cases, because of a paucity of stratigraphic markers or poor image quality, contacts must be considered gradational and appear dashed on the map.

NOACHIAN SYSTEM

The Noachian System consists of ancient crustal materials formed during the period of early bombardment that produced a high density of impact craters and large impact basins. Cratering, along with water and wind erosion and tectonic deformation, have modified or destroyed many of the primary morphologic features of Noachian units, including a substantial part of the ancient crater population. Thus, Noachian cratered terrains are usually characterized by impact, erosional, and tectonic features that postdate the material. The Noachian materials exhibit a wide variety of impact breccia, including those that are buried under younger units. Eolian and fluvial sediments that mantle and subdue parts of the heavily cratered terrain. Materials of the Noachian System, which include geologic units that straddle the Noachian/Hesperian boundary (fig. 6A; table 4), cover nearly 40% of the Thaumasia region.

The hilly unit (unit Nph) at the base of the plateau sequence forms heavily cratered terrain consisting of high-standing plateaus, irregular topographic highs, prominent ridges, and the far northwest part of the map region and probably includes Noachian basal outcrops (Scott and Tanaka, 1986; Wilhelms and others, 1991).

Resurfaced Noachian units of the plateau sequence include large valley forms, irregular depressions, broad, irregular basins covered by valley networks, furrows, and highly subdued ridges of all shapes and sizes. These highly modified surfaces, which include the materials that partly infill the floors of depressions and assortment of valley forms, are mapped as the dissected and etched unit of the circum-Argyre materials (unit HNdeh). Crater counts (total crater populations and superposed only crater populations; fig. 5*A*, *B*) clearly indicate that the modified material is Noachian and that the material deposited by resurfacing is Early Hesperian or younger; crater statistics along with stratigraphic and crosscutting relations of the Lowell impact crater materials with the surrounding geologic units and structure indicate that valley systems and channels (unit Nvc) were formed by the Lowell impact and Amazonian Lowell impact (Lias and others, 1997). Eolian, mass-wasting, fluvial, and (or) pyroclastic deposits (Baker and others, 1992; Kargel and Strom, 1992) could have formed the dissected and etched terrain.

Proximity to moderately eroded impact craters (mostly >50 km) that have continuous rim and ejecta blankets occur within Noachian and Hesperian materials of the map region. They are mapped as material of fresh to subdued craters (unit ec), which is mainly superposed on the surrounding rock materials.

Areas of Noachian materials and structure (faults and ridges) along the southern margin of the Thaumasia plateau are highly modified by Late Noachian/Early Hesperian sequences of relatively deep and narrow valleys. The dissected outcrops and the younger sediment deposited on their surfaces by the channeling are mapped as interbedded material of dissected (unit HNplh). We subdivided the unit into a high-relief and a low-relief unit based on the regional slope in large rift system (fig. 8*A*, *B*).

Intermediate fractured material (unit HNf) in the Claritas, Thaumasia, and Coracis Fossae and in the Sinai, Thaumasia highland, Daedalia southeast, and Sirenum provinces is cut by fewer and less complex faults than the older fractured material (sheet 2). The faults record tectonism of most of the regional centers that contributed to the structure of the older fractured material, but of lesser intensity. The intermediate fractured material is locally buried by younger plateau sequence materials (unit HNplh) and ridged plains material (unit HN_{rp}) and newer fractured material (unit HN_{fr}). Several units, mapped separately, are assigned to this unit from Scott and Tanaka (1986). The units are assigned to the formal stratigraphic nomenclature of Noachian, Hesperian, and Amazonian age by Scott and Carr (1978) and series (upper, middle, and lower divisions of systems) defined by Tanaka (1986). To determine the material and modification ages of the Thaumasia geologic units, we compiled crater statistics for 28 units of significant areal extents, which cover 96.6% of the map region. We measured all crater-rim diameters larger than 2 km using the Mars Digital Image Model (at 231 m/pixel; U.S. Geological Survey, 1993a, b) and calculated unit areas from digitally scanned contacts using GIS (tables 1 and 3). The crater statistics consist of total crater populations (including partly buried, degraded, and eroded, pristine impact craters), which may indicate emplacement ages, and superposed craters only, which indicate ages of most recent, major resurfacing (for example, see fig. 5*A*, *B*). We produced hundreds of crater counts that differ by as much as a factor of 10; we grouped the counts according to local or regional physiographic associations, and all outcrops together for each unit. The crater densities within physiographic provinces or groups vary somewhat; these ranges are typically incorporated in their time-stratigraphic age assignments.

The map units are categorized into channel-system and eolian materials, Valles Marineris interior deposits, western volcanic assemblage (flow and shield materials, Thaumasia Montes Formation, Syria Planum Formation), plateau and high-plains assemblage (plateau sequence, Thaumasia highland rift materials, dissected and subdued cratered terrain materials, subdivided material), circum-Argyre materials, and crater materials. Only materials of impact craters that have ejecta deposits nearly 50 km and greater are mapped; crater materials are assigned relative ages based on age relations with surrounding units and structures. For all units, relative-age assignments represent material emplacement ages; in some cases, erosional and depositional modification of the units somewhat obscures original formation. For example, dissected and etched materials include the eroded outcrops and superposed sediment deposits; therefore, the resulting age became the youngest age assigned to or mapped separately; their relative-age ranges reflect both materials.

Most unit contacts are fairly precise, especially where they are defined by ridge, embayment, and channel and buried relations. Transitional contacts, such as between fractured and dissected materials and adjacent geologic units are also sharp in most places. For example, these contacts commonly form along erosional scarps or partly buried tectonic structure. In other cases, because of a paucity of stratigraphic markers or poor image quality, contacts must be considered gradational and appear dashed on the map.

NOACHIAN SYSTEM

The Noachian System consists of ancient crustal materials formed during the period of early bombardment that produced a high density of impact craters and large impact basins. Cratering, along with water and wind erosion and tectonic deformation, have modified or destroyed many of the primary morphologic features of Noachian units, including a substantial part of the ancient crater population. Thus, Noachian cratered terrains are usually characterized by impact, erosional, and tectonic features that postdate the material. The Noachian materials exhibit a wide variety of impact breccia, including those that are buried under younger units. Eolian and fluvial sediments that mantle and subdue parts of the heavily cratered terrain. Materials of the Noachian System, which include geologic units that straddle the Noachian/Hesperian boundary (fig. 6A; table 4), cover nearly 40% of the Thaumasia region.

The hilly unit (unit Nph) at the base of the plateau sequence forms heavily cratered terrain consisting of high-standing plateaus, irregular topographic highs, prominent ridges, and highly degraded, ancient crater rims mainly in the Thaumasia highland, Daedalia, and Argyre provinces (fig. 2). Locally, tectonic and impact processes (Watters, 1993; Schultz and Tanaka, 1994; Craddock and others, 1990) may have contributed to the prominent relief of the hilly unit. Deep, steep-walled valleys, some as much as 200 km in diameter), and the center of the northeast-trending Thaumasia "bulge" or "rise," which extends for more than 3,000 km across the western equatorial region. This bulge forms a regional slope in the map region that trends southeast from Syria Planum at an elevation greater than 8 km to Argyre Planitia at an elevation less than 4 km (U.S. Geological Survey, 1989). The regional slope is interrupted by rugged highland terrain, which includes rifts, fossae, and fault systems, mesa-tains, and volcanoes. Differential wind and water erosion has eroded rock materials and structures along parts of the south edge of the Thaumasia plateau, resulting in well-defined valley networks, canyons, troughs, badlands, and mesas. The Argyre impact event includes the region, irregular topography produced by the colossal Argyre impact event includes relatively small basins and broad, scarp-bounded mesas.

We distinguish eleven geologic provinces within the Thaumasia region based on their structural, topographic, and geomorphic characteristics (fig. 1; table 2). These characteristics include broad canyons, highlands, large mountain ranges, volcanoes (fig. 4), lava and ridged plains, and surfaces that have been modified by tectonic, volcanic, and erosional processes.

STRATIGRAPHY

Geologic units were identified and mapped on individual, digitally enhanced Viking images and 1:2,000,000-scale photomosaic bases and compiled on the 1:5,000,000-scale digital photomosaic map base. Image resolution of the Thaumasia region ranges from 231 m > 500 m/pixel; those in the range of 50 to 300 m/pixel were most commonly used for mapping. The identification and classification of geologic units into groups and assemblages, where applicable, follows that of Scott and Tanaka (1986) in the region southeast of

(Schultz and Tanaka, 1994) and has morphologic features (fig. 17*A*) similar to those of the narrower (<50% of Coprates) Wind River Mountain Range of Wyoming (fig. 17*B*). Both mountain ranges exhibit faults, cuestas, hogbacks, and valleys.

Argyre structure (stages 1–2). During stage 1, a large impact event produced the Argyre basin and associated fault-controlled radial valleys and concentric mountain ranges, large fault scarps, and broad ridges. Normal offset of some of the concentric faults may have occurred during stages 1 and 2 because of isostatic adjustment of the basin.

Valles Marineris/Noctis Labyrinthus (stages 2–5). Numerous narrow (<5 km) and broad grabens (>5 km), which are radial to and parallel with Valles Marineris, cut stage 1 and 2 materials in the Sinai, Thaumasia Planum, Coprates, and Thaumasia highland provinces but are buried by stage 3 younger ridged plains material (for example, see fig. 13) of the Sinai province and stage 4 lava flows of the Syria Planum Formation of Syria–Solis province. Faults and grabens cut stage 4 and 5 materials of the Noctis Labyrinthus region outside the map region near Syria Planum. The south-central part of Valles Marineris has been identified as a possible center of stage 2 magmatic-driven tectonic activity (Anderson and others, 1998; Dohm and others, 1998).

VOLCANIC HISTORY

Detailed mapping of volcanic constructs and determination of their stratigraphic relations with surrounding materials and structures suggest that construct-forming volcanism occurred in the region throughout most of the Noachian Period and continued into the Early Hesperian; two distinct periods of construct-forming activity are shown on the geologic map. The older construct-forming activity produced 11 volcanoes in the Coprates, Thaumasia highland, Daedalia, and Sirenum provinces; several smaller domelike structures, which occur in the Coprates and Thaumasia highland provinces, also may represent volcanic activity during this time. Many of the volcanoes formed along extensional fault systems, suggesting a correlation between tectonism (development of deep-seated basement structures) and magmatic intrusions.

Late Noachian and Early Hesperian, younger construct-forming activity produced 3 volcanoes in the Thaumasia highlands province. Also during this time, widespread volcanism resulted in the emplacement of older ridged plains material (unit HN_{rp}; interpreted to consist of lava plains) in the Thaumasia Planum and northwestern part of the Argyre provinces and troughout material (unit HN_{tf}; interpreted to consist mainly of pyroclastic material) in the southern part of Coprates province. Younger rocks probably obscure a more widespread distribution of older ridged plains material; for example, subdued ridges of Daedalia southeast province may represent buried older ridged plains material.

The Hesperian system records younger ridged plains volcanism in the Sinai and Coprates east provinces followed by the emplacement of extensive sheet lava in the Syria–Solis, Daedalia, and Daedalia southeast provinces from the summit areas and flanks of the coronalike feature of Syria Planum and the Tharsis Montes volcanoes. The emplacement of voluminous sheet lava is coeval with waning Syria- and Tharsis-centered tectonism; thus, a significant change from tectonic to volcanic Syria- and Tharsis-dominated activity occurred during the Late Hesperian (stage 4) in the Thaumasia region. Similar to the Tharsis activity, the corona-like feature of Syria Planum may have developed over a long period of time (Middle Noachian to Late Hesperian). Evidence for long-lived volcanotectonic activity at Syria Planum is the wide age range of Syria-centered faults-especially in the Syria-Solis and Thaumasia highland provinces coupled with the large age progression of lava flow material (Scott and Tanaka, 1986; Wilhelms and others, 1991).

Resurfaced Noachian units of the plateau sequence include large valley forms, irregular depressions, broad, irregular basins covered by valley networks, furrows, and highly subdued ridges of all shapes and sizes. These highly modified surfaces, which include the materials that partly infill the floors of depressions and assortment of valley forms, are mapped as the dissected and etched unit of the circum-Argyre materials (unit HNdeh). Crater counts (total crater populations and superposed only crater populations; fig. 5*A*, *B*) clearly indicate that the modified material is Noachian and that the material deposited by resurfacing is Early Hesperian or younger; crater statistics along with stratigraphic and crosscutting relations of the Lowell impact crater materials with the surrounding geologic units and structure indicate that valley systems and channels (unit Nvc) were formed by the Lowell impact and Amazonian Lowell impact (Lias and others, 1997). Eolian, mass-wasting, fluvial, and (or) pyroclastic deposits (Baker and others, 1992; Kargel and Strom, 1992) could have formed the dissected and etched terrain.

Proximity to moderately eroded impact craters (mostly >50 km) that have continuous rim and ejecta blankets occur within Noachian and Hesperian materials of the map region. They are mapped as material of fresh to subdued craters (unit ec), which is mainly superposed on the surrounding rock materials.

Areas of Noachian materials and structure (faults and ridges) along the southern margin of the Thaumasia plateau are highly modified by Late Noachian/Early Hesperian sequences of relatively deep and narrow valleys. The dissected outcrops and the younger sediment deposited on their surfaces by the channeling are mapped as interbedded material of dissected (unit HNplh). We subdivided the unit into a high-relief and a low-relief unit based on the regional slope in large rift system (fig. 8*A*, *B*).

Intermediate fractured material (unit HNf) in the Claritas, Thaumasia, and Coracis Fossae and in the Sinai, Thaumasia highland, Daedalia southeast, and Sirenum provinces is cut by fewer and less complex faults than the older fractured material (sheet 2). The faults record tectonism of most of the regional centers that contributed to the structure of the older fractured material, but of lesser intensity. The intermediate fractured material is locally buried by younger plateau sequence materials (unit HNplh) and ridged plains material (unit HN_{rp}) and newer fractured material (unit HN_{fr}). Several units, mapped separately, are assigned to this unit from Scott and Tanaka (1986). The units are assigned to the formal stratigraphic nomenclature of Noachian, Hesperian, and Amazonian age by Scott and Carr (1978) and series (upper, middle, and lower divisions of systems) defined by Tanaka (1986).

Terraced material (unit HNtr) occurs mostly in the southern and northeastern parts of the Thaumasia Planum and Thaumasia highland provinces, respectively, largely bordering topographic highs. It is marked by a series of terraces with steps and risers, benches, terraces, and subdued wrinkle ridges and faults. The terraced unit is transitional in places with the older ridged plains material and the mountain flank surfaces of the Thaumasia Montes. Fluvial erosion may have formed terraces by exposing older basal materials and (or) by stripping away the upper layers of the older ridged plains material. The older ridged plains material is marked by numerous wrinkle ridges, narrow linear ridges, and faults of varying trends.

The younger construct-forming activity of the map region produced three of the fourteen prominent volcanoes within the Thaumasia highlands province in the Thaumasia province. They are mapped as younger flow and construct material (unit HNto). One example, along a large rift (fig. 8*A*), of the younger flow and construct material is the younger fractured material (unit Hf). These outcrops, which record continued volcanotectonic activity of Apollinaris (Scott and others, 1993) and Hadriaca Patera (Crown and Greeley, 1993) are preserved on a few isolated mountain flanks, as at Cyclops (fig. 4*A*, *B*).

A relatively thin, Late Noachian to Early Hesperian and possibly younger plains-forming material forms the lower flows of the Argyre province and flows that extend toward the basin. The material is mapped as the smooth floor unit of the circum-Argyre materials (unit HNplc), which may include lava flows, impact melt, eolian, fluvial, and (or) possibly glacial origin (Baker and others, 1991; Kargel and Strom, 1992).

Chaotic material (unit HNch) of the channel-system and eolian materials forms several faulted patches of knobby terrain that may be mapped as older fractured material (unit Nf). The older fractured material, which is marked by highly eroded and deformed impact craters, is cut by numerous fault systems consisting of complex fractures and graben-bounding fault that record regional Thaumasia, Syria Planum, and possible Valles Marineris tectonism and local intrusions and uplifts. Where the older fractured material has been highly dissected, it is mapped as fractured and dissected material (unit Nfd); superposition relations with younger material indicate that many faults and valleys within unit Nfd are Noachian (sheets 2 and 3, respectively).

Along the southeast margin of the Thaumasia plateau, a small isolated patch of etched material (unit Nplh) is cut by flat-floored troughs with serrated ridges. In this region and in other places along the plateau's edge, a wide variety of ensemble features occurs, including locally networked trough-and-valley forms, highly eroded impact crater rims, faults, irregular basin floors, and buried craters. These features are probably eolian, fluvial erosion may have highly modified older rock outcrops of the Thaumasia plateau, which include interbedded deposits of impact breccia, lava flows, and eolian and fluvial sediments (Scott and others, 1995; Carr and Chuang, 1997) is recorded during this period. In the region, wind erosion appears to be the dominant resurfacing agent. The Amazonian System materials consist of (1) lava flows of the Syria Planum Formation and (2) eolian deposits. Amazonian materials consist of widespread smooth floor crater material (unit es), Valles Marineris interior deposits (unit Avf), and the younger fractured material (unit Af). Formation of these structures largely ceased during the later part of stage 2. In addition to large scarps, highly resistant lava flows and (or) welded pyroclastic flows in the map region may be the result of extensional or contractional tectonism, erosion, or deposition.

Thaumasia highland rifts (stages 1–3). Rift systems in the Thaumasia highland province are defined by well-developed graben (tens of kilometers wide) and associated normal faults (<5 km wide) and commonly include probable volcanic constructs (for example, see fig. 4). Tectonism from these rifts contributed to the development of the plateau sequence materials (Dohm and Tanaka, 1999) contributed to the development of the rift systems during stages 1 and 2; locally, the rift structures were reactivated during stage 3. The close association of fault systems with volcanoes on Mars (Wise and others, 1979; Scott and Tanaka, 1980; Whitford-Stark, 1982; Scott and Dohm, 1990a; Banerdt and others, 1992), which is similar to strong correlations between volcanic constructs and fault systems of several volcanic fields on Earth (for example, see Scott and Trask, 1971; Shoemaker and others, 1978; Crumpler, 1982; Holm and others, 1991; Dohm, 1995), suggests that magma used deep-seated basement structures as conduits to reach the martian surface during periods of volcanic activity.

Coprates rise (stages 1–2). Contractional tectonism, which may be related to vertical plateau uplift related to magmatism and thermal rise (Dohm and Tanaka, 1999), mainly during stage 2 resulted in the formation of a large (900 km long and 2 to 4 km high) asymmetric ridge (higher on the east); the ridge is the largest of several in the Thaumasia fold belt

2. Late Noachian (stages 1–2)—Decreasing impact rate; widespread aggradational and degradational modification that partly subdued older surfaces; ridged plains volcanism in the Thaumasia Planum province; continued development of Claritas, Thaumasia, Coracis, Melas, and Nectaris Fossae, volcanotectonic centers of Tharsis and Syria Planum, Coprates rise, volcanoes, broad ridges and large scarps, and Thaumasia highland rifts; wrinkle ridge formation and the production of faults and grabens associated with local magmatic-driven centers of tectonic activity near central Valles Marineris and the source region of Warrego Valles; possible reactivation of Argyre impact-related, concentric faults; formation of local valley systems and isolated valleys, which include Warrego Valles.

3. Early Hesperian (stages 2–3)—Extensive ridged plains volcanism in the Sinai and east Coprates provinces; widespread resurfacing of older Noachian rock; ending production of Coracis, Melas, and Nectaris Fossae, wrinkle ridges, broad ridges and large scarps, volcanoes, Thaumasia highland rifts, Coprates rise, and faults and grabens of the Thaumasia Planum province; continued development of Claritas and Thaumasia Fossae, Valles Marineris/Noctis Labyrinthus faults and grabens, volcanotectonic centers of Tharsis and Syria Planum, and local valley systems and isolated valleys.

4. Late Hesperian (stage 4)—Emplacement of extensive sheet flows in the Syria–Solis and Daedalia provinces from the summit areas and flanks of Tharsis Montes volcanoes and Syria Planum; faulting of lava flow materials in the northwest part of the map region; significant waning of tectonic deformation, which includes faults and grabens of Claritas Fossae, Syria Planum, and Valles Marineris/Noctis Labyrinthus, and waning development of valley systems and isolated valleys.

5. Amazonian (stage 5)—Continued emplacement of Tharsis-related lavas in Daedalia province; deposition of smooth crater material, Valles interior deposits, and dune material; minor Tharsis-related faulting in the northwest corner of map region; wind erosion and deposition becomes the dominate resurfacing agent.

REFERENCES CITED

Anderson, R.C., Golombek, M.P., Franklin, B.J., Tanaka, K.L., Dohm, J.M., Lias, J.H., and Peer, B., 1998, Centers of tectonic activity through time for the western hemisphere of Mars [abs.], *in* Abstracts of papers submitted to the Twenty-ninth Lunar and Planetary Science Conference, Houston, March 16–20, 1998: Houston, Lunar and Planetary Institute, no. 1881.

Baker, V.R., 1982, The channels of Mars: Austin, University of Texas Press, 198 p.

Baker, V.R., Strom, R.G., Gulick, V.C., Kargel, J.S., Komatsu, Goro, and Kale, V.S., 1991, Ancient oceans, ice sheets and the hydrological cycle on Mars: Nature, v. 352, p. 589–594.

Banerdt, W.B., Golombek, M.P, and Tanaka, K.L., 1992, Stress and tectonics on Mars, *in* Kieffer, H.H., and Jakosky, B.M., eds., Mars: Tucson, University of Arizona Press, p. 249–297.

Banerdt, W.B., Phillips, R.J., Sleep, N.H., and Saunders, R.S., 1982, Thick-shell tectonics on one-plate planets—Applications to Mars: Journal of Geophysical Research, v. 87, p. 9723–9733.

Brakenridge, G.R., Newsom, H.E., and Baker, V.R., 1985, Ancient hot springs on Mars: Origins and paleoenvironmental significance of small martian valleys: Geology, v. 13, Ibus. 12, p. 859–862.

Cabrol, N.A., Grin, E.A., and Dawidowicz, Gilles, 1997, A model of outflow generation by hydrothermal underpressure drainage in volcano-tectonic environment, Shalbatana Complex, Mars: Icarus, v. 126, p. 455–468.

Carr, M.H., 1989, Recharge of the early atmosphere of Mars by impact-induced release of CO2: Icarus, v. 79, no.2, p. 311–327.

———1995, The martian drainage system and the origin of valley networks and fretted channels: Journal of Geophysical Research, v. 100, no. E4, p. 7479–7507.

Carr, M.H., and Chuang, F.C., 1997, Martian drainage densities: Journal of Geophysical Research, v. 102, no. E4, p. 9145–9152.

Carr, M.H., and Clow, G.D., 1981, Martian channels and valleys—Their characteristics, distributions, and age: Icarus, v. 48, no. 1, p. 91–117.

Clifford, S.M., 1984, The hydrology of Mars—An update: Icarus, v. 50, p. 277–295.

Clifford, S.M., 1993, A model for the hydrologic and climatic behavior of water on Mars: Journal of Geophysical Research, v. 98, p. 10,973–11,016.

Craddock, R.A., Greeley, Ronald, and Christensen, P.R., 1990, Evidence of an ancient impact basin in northern Amazonis Planitia, Mars: Journal of Geophysical Research, v. 95, p. 10,729–10,741.

Dohm, J.M., Barlow, N.G. and Maxwell, T.A., 1993, Geomorphic evolution of the martian highlands through ancient fluvial processes: Journal of Geophysical Research, v. 98, p. 3453–3468.

Crown, D.A., and Greeley, Ronald, 1993, Volcanic geology of Hadriaca Patera and the eastern Hellas region of Mars: Journal of Geophysical Research, v. 98, p. 3431–3451.

Crown, D.A., Price, K.H., and Greeley, Ronald, 1992, Geologic evolution of the east rim of the Hellas basin, Mars: Icarus, v. 100, p. 1–25.

Crumpler, L.S., 1982, Geology of the Mount Taylor region: New Mexico Geological Society Guidebook, 33rd Field Conference, p. 291–298.

Dohm, J.M., 1995, Origin of Stonesnan lake, and volcano-tectonic relations of Mormon and Stonesnan lakes, Arizona: Flagstaff, Arizona, M.S. thesis, Northern Arizona University, Flagstaff, Arizona, 101 p.

Dohm, J.M., Gulick, V.C., and Tanaka, K.L., 1998, Digital structural mapping of Mars: Astronomy & Geophysics, v. 39, p. 3.20–3.22.

Dohm, J.M., and Scott, D.H., 1993, Relation between ages and elevations of martian channels [abs.], *in* Lunar and Planetary Science Conference, 24th, Houston, March 13–17, 1989, Proceedings, p. 515–523.

———1991, Hydrologic activity during Late Noachian and Early Hesperian downwarping of Boreotlas basin, Mars [abs.], *in* Abstracts of papers submitted to the Twenty-second Lunar and Planetary Science Conference, Houston, March 18–22, 1991: Houston, Lunar and Planetary Institute, p. 1377–1378.

Dohm, J.M., and Tanaka, K.L., 1999, Geology of the Thaumasia region, Mars: Planetary and planetocentric, valley origins, and magmatic evolution: Planetary and Space Science, v. 47, p. 411–431.

Dohm, J.M., Tanaka, K.L., and Hare, T.M., 1996, Unraveling the complex geologic history of the Thaumasia region, Mars: Geologic mapping of the Claritas Fossae quadrangle: Lunar and Planetary Science Conference, Houston