Double-β Decay

NODE=S076

OMITTED FROM SUMMARY TABLE

A REVIEW GOES HERE – Check our WWW List of Reviews

NODE=S076

Half-life Measurements and Limits for Double-β Decay

NODE=S076HDB

NODE=S076HDB

In most cases the transitions (Z,A) $\rightarrow$ (Z$-$2,A) + $2e^-$ + (0 or 2) $\overline{\nu}_e$ to the 0^+ ground state of the final nucleus are listed. However, we also list transitions that increase the nuclear charge ($2e^+$, e^+/EC and ECEC) and transitions to excited states of the final nuclei (0^+_i, 2^+, and 2^+_i). In the following Listings, only best or comparable limits or lifetimes for each isotope are reported. For 2ν decay, which is well established, only measured half-lives are reported.

NODE=S076HDB

• • • We do not use the following data for averages, fits, limits, etc. • • •

$t_{1/2}(10^{21}$ yr)	CL%	ISOTOPE	TRANSITION		METHOD	DOCUMENT ID		
>3000	90	^{130}Te	0ν		TeO_2 bolometer	[1] ARNABOLDI	08	
>0.004	90	^{64}Zn	0ν	2K	$ZnWO_4$ scint.	[2] BELLI	08	
>0.22	90	^{64}Zn	0ν		$ZnWO_4$ scint.	[3] BELLI	08	OCCUR=2
>0.001	90	^{108}Cd	0ν	2K	$CdWO_4$ scint.	[4] BELLI	08B	
>0.0013	90	^{114}Cd	2ν	$\beta\beta$	$CdWO_4$ scint.	[5] BELLI	08B	OCCUR=2
>1.1	90	^{114}Cd	0ν	$\beta\beta$	$CdWO_4$ scint.	[6] BELLI	08B	OCCUR=3
>58	90	^{48}Ca	0ν		CaF_2 scint.	[7] UMEHARA	08	
$0.57^{+0.13}_{-0.09}\pm0.08$	68	^{100}Mo	2ν	$0^+\rightarrow0^+_1$	NEMO-3	[8] ARNOLD	07	
>89	90	^{100}Mo	0ν	$0^+\rightarrow0^+_1$	NEMO-3	[9] ARNOLD	07	OCCUR=2
>1.1	90	^{100}Mo	2ν	$0^+\rightarrow2^+$	NEMO-3	[10] ARNOLD	07	OCCUR=3
>160	90	^{100}Mo	0ν	$0^+\rightarrow2^+$	NEMO-3	[11] ARNOLD	07	OCCUR=4
>0.0019	90	^{74}Se	0ν+2ν		γ in Ge det.	[12] BARABASH	07	
>0.0055	90	^{74}Se	0ν+2ν	$0^+\rightarrow2^+_2$	γ in Ge det.	[13] BARABASH	07	OCCUR=2
$>(1.9–6.0)\ 10^{-4}$	90	^{120}Te	0ν		γ in Ge det.	[14] BARABASH	07B	
$>1.9\times10^{-4}$	90	^{120}Te	0ν+2ν		γ in Ge det.	[15] BARABASH	07B	OCCUR=2
$>7.5\times10^{-4}$	90	^{120}Te	0ν+2ν	$0^+\rightarrow2^+$	γ in Ge det.	[16] BARABASH	07B	OCCUR=3
$>1.19\times10^{-4}$	90	^{64}Zn	0ν		CdZnTe calorim.	[17] BLOXHAM	07	
$>1.21\times10^{-4}$	90	^{120}Te	0ν		CdZnTe calorim.	[18] BLOXHAM	07	OCCUR=2
$>2.68\times10^{-6}$	90	^{120}Te	0ν		CdZnTe calorim.	[19] BLOXHAM	07	OCCUR=3
$>9.72\times10^{-6}$	90	^{120}Te	0ν	$0^+\rightarrow2^+_1$	CdZnTe calorim.	[20] BLOXHAM	07	OCCUR=4
22300^{+4400}_{-3100}	68	^{76}Ge	0ν		Enriched HPGe	[21] KLAPDOR-K...	06A	
>1800	90	^{130}Te	0ν		Cryog. det.	[22] ARNABOLDI	05	
>460	90	^{100}Mo	0ν		NEMO-3	[23] ARNOLD	05A	
>100	90	^{82}Se	0ν		NEMO-3	[24] ARNOLD	05A	OCCUR=2
$(7.11\pm0.02\pm0.54)$E-3		^{100}Mo	2ν		NEMO-3	[25] ARNOLD	05A	OCCUR=3
$(9.6\pm0.3\pm1.0)$E-2		^{82}Se	2ν		NEMO-3	[26] ARNOLD	05A	OCCUR=4
>550	90	^{130}Te	0ν		Cryog. det.	[27] ARNABOLDI	04	
>310	90	^{100}Mo	0ν		NEMO-3	[28] ARNOLD	04	
>140	90	^{82}Se	0ν		NEMO-3	[29] ARNOLD	04	OCCUR=2
$(7.68\pm0.02\pm0.54)$E-3		^{100}Mo	2ν		NEMO-3	[30] ARNOLD	04	OCCUR=3
$(10.3\pm0.3\pm0.7)$E-2		^{82}Se	2ν		NEMO-3	[31] ARNOLD	04	OCCUR=4
$0.14^{+0.04}_{-0.02}\pm0.03$	68	^{150}Nd	0ν+2ν	$0^+\rightarrow0^+_1$	γ in Ge det.	[32] BARABASH	04	
11900^{+29900}_{-5000}	99.7	^{76}Ge	0ν		Enriched HPGe	[33] KLAPDOR-K...	04A	
>14	90	^{48}Ca	0ν		CaF_2 scint.	[34] OGAWA	04	
>210	90	^{130}Te	0ν		Cryog. det.	[35] ARNABOLDI	03	
>31	90	^{130}Te	0ν	$0^+\rightarrow2^+$	Cryog. det.	[36] ARNABOLDI	03	OCCUR=2
$0.61\pm0.14^{+0.29}_{-0.35}$	90	^{130}Te	2ν		Cryog. det.	[37] ARNABOLDI	03	OCCUR=3
>110	90	^{128}Te	0ν		Cryog. det.	[38] ARNABOLDI	03	OCCUR=4
$(0.029^{+0.004}_{-0.003})$		^{116}Cd	2ν		$^{116}CdWO_4$ scint.	[39] DANEVICH	03	
>170	90	^{116}Cd	0ν		$^{116}CdWO_4$ scint.	[40] DANEVICH	03	OCCUR=2
>29	90	^{116}Cd	0ν	$0^+\rightarrow2^+$	$^{116}CdWO_4$ scint.	[41] DANEVICH	03	OCCUR=3
>14	90	^{116}Cd	0ν	$0^+\rightarrow0^+_1$	$^{116}CdWO_4$ scint.	[42] DANEVICH	03	OCCUR=4
>6	90	^{116}Cd	0ν	$0^+\rightarrow0^+_2$	$^{116}CdWO_4$ scint.	[43] DANEVICH	03	OCCUR=5
$1.74\pm0.01^{+0.18}_{-0.16}$		^{76}Ge	2ν		Enriched HPGe	[44] DOERR	03	
>15700	90	^{76}Ge	0ν		Enriched HPGe	[45] AALSETH	02B	
>58	90	^{134}Xe	0ν		Liquid Xe Scint.	[46] BERNABEI	02D	

> 1200	90	^{136}Xe	0ν		Liquid Xe Scint.	[47] BERNABEI	02D	OCCUR=2
$15000^{+168000}_{-7500}$		^{76}Ge	0ν		Enriched HPGe	[48] KLAPDOR-K...	02D	
$(7.2 \pm 0.9 \pm 1.8)$E-3		^{100}Mo	2ν		Liq. Ar ioniz.	[49] ASHITKOV	01	
$>$ 4.9	90	^{100}Mo	0ν		Liq. Ar ioniz.	[50] ASHITKOV	01	OCCUR=2
$>$ 1.3	90	^{160}Gd	0ν		Gd_2SiO_5:Ce	[51] DANEVICH	01	
$>$ 1.3	90	^{160}Gd	0ν	$0^+ \rightarrow 2^+$	Gd_2SiO_5:Ce	[52] DANEVICH	01	OCCUR=2
$0.59^{+0.17}_{-0.11} \pm 0.06$		^{100}Mo	$0\nu+2\nu$	$0^+ \rightarrow 0^+_1$	Ge coinc.	[53] DEBRAECKEL...	01	
$>$ 55	90	^{100}Mo	$0\nu,\langle m_\nu \rangle$		ELEGANT V	[54] EJIRI	01	
$>$ 42	90	^{100}Mo	$0\nu,\langle \lambda \rangle$		ELEGANT V	[54] EJIRI	01	OCCUR=2
$>$ 49	90	^{100}Mo	$0\nu,\langle \eta \rangle$		ELEGANT V	[54] EJIRI	01	OCCUR=3
$>$19000	90	^{76}Ge	0ν		Enriched HPGe	[55] KLAPDOR-K...	01	
$1.55 \pm 0.001^{+0.19}_{-0.15}$	90	^{76}Ge	2ν		Enriched HPGe	[56] KLAPDOR-K...	01	OCCUR=2
(9.4 ± 3.2)E-3	90	^{96}Zr	$0\nu+2\nu$		Geochem	[57] WIESER	01	
$0.042^{+0.033}_{-0.013}$		^{48}Ca	2ν		Ge spectrometer	[58] BRUDANIN	00	OCCUR=2
$0.021^{+0.008}_{-0.004} \pm 0.002$		^{96}Zr	2ν		NEMO-2	[59] ARNOLD	99	
> 1.0	90	^{96}Zr	0ν		NEMO-2	[59] ARNOLD	99	OCCUR=2
$(8.3 \pm 1.0 \pm 0.7)$E-2		^{82}Se	2ν		NEMO-2	[60] ARNOLD	98	
$>$ 9.5	90	^{82}Se	0ν		NEMO-2	[61] ARNOLD	98	OCCUR=2
$>$ 2.8	90	^{82}Se	0ν	$0^+ \rightarrow 2^+$	NEMO-2	[62] ARNOLD	98	OCCUR=3
$(7.6^{+2.2}_{-1.4})$E-3		^{100}Mo	2ν		Si(Li)	[63] ALSTON-...	97	OCCUR=2
$(6.82^{+0.38}_{-0.53} \pm 0.68)$E-3		^{100}Mo	2ν		TPC	[64] DESILVA	97	
$(6.75^{+0.37}_{-0.42} \pm 0.68)$E-3		^{150}Nd	2ν		TPC	[65] DESILVA	97	OCCUR=3
$>$ 1.2	90	^{150}Nd	0ν		TPC	[66] DESILVA	97	OCCUR=4
$(3.75 \pm 0.35 \pm 0.21)$E-2		^{116}Cd	2ν	$0^+ \rightarrow 0^+$	NEMO 2	[67] ARNOLD	96	
$0.043^{+0.024}_{-0.011} \pm 0.014$		^{48}Ca	2ν		TPC	[68] BALYSH	96	
0.79 ± 0.10		^{130}Te	$0\nu+2\nu$		Geochem	[69] TAKAOKA	96	
$0.61^{+0.18}_{-0.11}$		^{100}Mo	$0\nu+2\nu$	$0^+ \rightarrow 0^+_1$	γ in HPGe	[70] BARABASH	95	
$(9.5 \pm 0.4 \pm 0.9)$E-3		^{100}Mo	2ν		NEMO 2	DASSIE	95	
$>$ 0.6	90	^{100}Mo	0ν	$0^+ \rightarrow 0^+_1$	NEMO 2	DASSIE	95	OCCUR=4
$0.026^{+0.009}_{-0.005}$		^{116}Cd	2ν	$0^+ \rightarrow 0^+$	ELEGANT IV	EJIRI	95	
$0.017^{+0.010}_{-0.005} \pm 0.0035$		^{150}Nd	2ν	$0^+ \rightarrow 0^+$	TPC	ARTEMEV	93	
0.039 ± 0.009		^{96}Zr	$0\nu+2\nu$		Geochem	KAWASHIMA	93	
2.7 ± 0.1		^{130}Te	$0\nu+2\nu$		Geochem	BERNATOW...	92	
7200 ± 400		^{128}Te	$0\nu+2\nu$		Geochem	[71] BERNATOW...	92	OCCUR=2
$>$ 27	68	^{82}Se	0ν	$0^+ \rightarrow 0^+$	TPC	ELLIOTT	92	
$0.108^{+0.026}_{-0.006}$		^{82}Se	2ν	$0^+ \rightarrow 0^+$	TPC	ELLIOTT	92	OCCUR=2
2.0 ± 0.6		^{238}U	$0\nu+2\nu$		Radiochem	[72] TURKEVICH	91	
$>$ 9.5	76	^{48}Ca	0ν		CaF_2 scint.	YOU	91	
$0.12 \pm 0.01 \pm 0.04$	68	^{82}Se	$0\nu+2\nu$		Geochem.	[73] LIN	88	
$0.75 \pm 0.03 \pm 0.23$	68	^{130}Te	$0\nu+2\nu$		Geochem.	[74] LIN	88	OCCUR=2
1800 ± 700	68	^{128}Te	$0\nu+2\nu$		Geochem.	[75] LIN	88B	
2.60 ± 0.28		^{130}Te	$0\nu+2\nu$		Geochem	[76] KIRSTEN	83	OCCUR=2

[1] ARNABOLDI 08 use high resolution TeO_2 bolometer calorimeter to search for double beta decay of ^{130}Te. Supersedes ARNABOLDI 05. NODE=S076HDB;LINKAGE=AL

[2] BELLI 08 use $ZnWO_4$ scintillation calorimeter to search for neutrinoless double K-shell electron capture decay of ^{64}Zn. Slightly weaker limit is obtained for capture from other shells. The halflife limit for the 2ν mode is 6.2×10^{18} years. NODE=S076HDB;LINKAGE=BE

[3] BELLI 08 use $ZnWO_4$ scintillation calorimeter to search for neutrinoless β^+ plus electron capture decay of ^{64}Zn. The halflife limit for the 2ν mode is 2.1×10^{20} years. NODE=S076HDB;LINKAGE=BI

[4] BELLI 08B use $CdWO_4$ scintillation calorimeter to search for 0ν and 2ν decay with 2K electron captures of ^{108}Cd. The absence of signal at 2×K electron binding energy (48.8 keV) is used to derive the halflife limit. Search for γ at 223 keV gives halflife limit for the 0ν mode of 1.0×10^{18} years. NODE=S076HDB;LINKAGE=EL

[5] BELLI 08B use $CdWO_4$ scintillation calorimeter to search for 2ν $\beta\beta$ decay of ^{114}Cd. NODE=S076HDB;LINKAGE=LL

[6] BELLI 08B use $CdWO_4$ scintillation calorimeter to search for 0ν $\beta\beta$ decay of ^{114}Cd. NODE=S076HDB;LINKAGE=EI

[7] UMEHARA 08 use CaF_2 scintillation calorimeter to search for double beta decay of ^{48}Ca. Limit is significantly more stringent than quoted sensitivity: 18×10^{21} years. NODE=S076HDB;LINKAGE=UM

[8] First exclusive measurement of 2ν-decay to the first excited 0^+_1-state of daughter nucleus. ARNOLD 07 use the NEMO-3 tracking calorimeter to detect all particles emitted in decay. Result agrees with the inclusive (0ν + 2ν) measurement of DEBRAECKELEER 01. NODE=S076HDB;LINKAGE=NA

[9] Limit on 0ν-decay to the first excited 0^+_1-state of daughter nucleus using NEMO-3 tracking calorimeter. Supersedes DASSIE 95. NODE=S076HDB;LINKAGE=NR

[10] Limit on 2ν-decay to the first excited 2^+-state of daughter nucleus using NEMO-3 tracking calorimeter. NODE=S076HDB;LINKAGE=NN

[11] Limit on 0ν-decay to the first excited 2^+-state of daughter nucleus using NEMO-3 tracking calorimeter. NODE=S076HDB;LINKAGE=NO

[12] BARABASH 07 use Ge calorimeter to search for γ-radiation following double electron capture or β^+ plus electron capture decays of ^{74}Sr to the ground state of ^{74}Ge. This limit is based on the search for the 511 keV annihilation radiation. Various other limits, for the capture from different atomic shells and also to the excited states, are reported in the paper. NODE=S076HDB;LINKAGE=BR

[13] BARABASH 07 use Ge calorimeter to search for γ-radiation following double electron capture decay of ^{74}Sr into the second excited 2^+-state of ^{74}Ge. That transition has been considered due to a possible resonance enhancement. The 2ν mode would be suppressed for this decay by its extremely small phase space factor. NODE=S076HDB;LINKAGE=BS

[14] BARABASH 07B use Ge calorimeter to search for γ-radiation following the double electron capture decay of ^{120}Te. This limit is based on the search for the bremsstrahlung radiation. Various limits, for the capture from different atomic shells, are reported that cover the range of halflives shown. NODE=S076HDB;LINKAGE=CA

[15] BARABASH 07B use Ge calorimeter to search for the 511 keV positron annihilation radiation following the β^+ plus electron capture decay of ^{120}Te. NODE=S076HDB;LINKAGE=CB

[16] BARABASH 07B use Ge calorimeter to search for γ-radiation following the double electron capture decay of ^{120}Te into the excited 2^+ state of ^{120}Sn. This limit is based on the search for the γ-radiation from the excited 2^+ state. NODE=S076HDB;LINKAGE=CC

[17] BLOXHAM 07 use CdZnTe solid state detectors to search for the decays of the various double beta detector components, here for double electron capture ^{64}Zn decay. NODE=S076HDB;LINKAGE=BM

[18] BLOXHAM 07 use CdZnTe solid state detectors to search for the decays of the various double beta detector components, here for β^+ plus electron capture ^{120}Te decay. NODE=S076HDB;LINKAGE=BH

[19] BLOXHAM 07 use CdZnTe solid state detectors to search for the decays of the various double beta detector components, here for double electron capture ^{120}Te decay. NODE=S076HDB;LINKAGE=BX

[20] BLOXHAM 07 use CdZnTe solid state detectors to search for the decays of the various double beta detector components, here for double electron capture ^{120}Te decay to the first excited 2^+-state. NODE=S076HDB;LINKAGE=BO

[21] KLAPDOR-KLEINGROTHAUS 06A present re-analysis of data originally published in KLAPDOR-KLEINGROTHAUS 04A. Modified pulse shape analysis leads the authors to claim improved 6σ statistical evidence for observation of 0ν-decay, compared to 4.2σ in KLAPDOR-KLEINGROTHAUS 04A. Analysis of the systematic uncertainty is not presented. Supersedes KLAPDOR-KLEINGROTHAUS 04A. NODE=S076HDB;LINKAGE=KR

[22] Supersedes ARNABOLDI 04. Bolometric TeO_2 detector array CUORICINO is used for high resolution search for $0\nu\beta\beta$ decay. The half-life limit is derived from 3.09 kg yr ^{130}Te exposure. NODE=S076HDB;LINKAGE=AO

[23] NEMO-3 tracking calorimeter containing 6.9 kg of enriched ^{100}Mo is used in ARNOLD 05A. A limit for $0\nu\beta\beta$ half-life of ^{100}Mo is reported. Supersedes ARNOLD 04. NODE=S076HDB;LINKAGE=N1

[24] NEMO-3 tracking calorimeter is used in ARNOLD 05A to place limit on $0\nu\beta\beta$ half-life of ^{82}Se. Detector contains 0.93 kg of enriched ^{82}Se. Supersedes ARNOLD 04. NODE=S076HDB;LINKAGE=N2

[25] ARNOLD 05A use the NEMO-3 tracking calorimeter to determine the $2\nu\beta\beta$ half-life of ^{100}Mo with high statistics and low background (389 days of data taking). Supersedes ARNOLD 04. NODE=S076HDB;LINKAGE=N3

[26] ARNOLD 05A use the NEMO-3 tracking detector to determine the $2\nu\beta\beta$ half-life of ^{82}Se with high statistics and low background (389 days of data taking). Supersedes ARNOLD 04. NODE=S076HDB;LINKAGE=N4

[27] Supersedes ARNABOLDI 03. Bolometric TeO_2 detector array Cuoricino used for high resolution search for $0\nu\beta\beta$ decay. NODE=S076HDB;LINKAGE=AN

[28] ARNOLD 04 use the NEMO-3 tracking detector to determine the limit for $0\nu\beta\beta$ halflife of ^{100}Mo. This represents an improvement, by a factor of ~ 6, when compared with EJIRI 01. NODE=S076HDB;LINKAGE=A1

[29] ARNOLD 04 use the NEMO-3 tracking detector to determine the limit for $0\nu\beta\beta$ halflife of ^{82}Se. This represents an improvement, by a factor of ~ 10, when compared with ELLIOTT 92. It supersedes the limit of ARNOLD 98 for this decay using NEMO-2. NODE=S076HDB;LINKAGE=A2

[30] ARNOLD 04 use the NEMO-3 tracking detector to determine the $2\nu\beta\beta$ halflife of ^{100}Mo with high statistics and low background. The halflife is determined assuming the Single State Dominance. It is in agreement with, and more accurate than, previous determinations. Supersedes DASSIE 95 determination of this quantity with NEMO-2. NODE=S076HDB;LINKAGE=A3

[31] ARNOLD 04 use the NEMO-3 tracking detector to determine the $2\nu\beta\beta$ halflife of ^{82}Se. The halflife is in agreement with ARNOLD 98 with NEMO-2 which it supersedes. NODE=S076HDB;LINKAGE=A4

[32] BARABASH 04 perform an inclusive measurement of the $\beta\beta$ decay of ^{150}Nd into the first excited (0^+_1) state of ^{150}Sm. Gamma radiation emitted in decay of the excited state is detected. NODE=S076HDB;LINKAGE=BA

[33] Supersedes KLAPDOR-KLEINGROTHAUS 02D. Authors present new analysis of event excess seen in Heidelberg-Moscow experiment at $\beta\beta$-decay energy. Enhanced statistics leads to a 4.2 σ evidence for observation of 0ν $\beta\beta$-decay and a finite Majorana neutrino mass. Stated error is purely statistical. No systematic errors are mentioned in the paper. More details can be found in KLAPDOR-KLEINGROTHAUS 04C. NODE=S076HDB;LINKAGE=KL

[34] CaF_2 scintillation calorimeter ELEGANT VI used to set limit on $0\nu\beta\beta$-decay rate of ^{48}Ca. The stated half-life limit benefits from a downward fluctuation on the number of background events. The experimental sensitivity is 5.9×10^{21} yr. at 90 % CL. Replaces YOU 91 as the most stringest experiment using ^{48}Ca. NODE=S076HDB;LINKAGE=OG

[35] Supersedes ALESSANDRELLO 00. Array of TeO_2 crystals in high resolution cryogenic calorimeter. Some enriched in ^{130}Te. Ground state to ground state decay. NODE=S076HDB;LINKAGE=RN

[36] Decay into first excited state of daughter nucleus. NODE=S076HDB;LINKAGE=RP

[37] Two neutrino decay into ground state. Relatively large error mainly due to uncertainties in background determination. Reported value is shorter than the geochemical measurements of KIRSTEN 83 and BERNATOWICZ 92 but in agreement with LIN 88 and TAKAOKA 96. NODE=S076HDB;LINKAGE=RQ

[38] Supersedes ALESSANDRELLO 00. Array of TeO_2 crystals in high resolution cryogenic calorimeter. Some enriched in ^{128}Te. Ground state to ground state decay. NODE=S076HDB;LINKAGE=RO

[39] Calorimetric measurement of 2ν ground state decay of ^{116}Cd using enriched $CdWO_4$ scintillators. Agrees with EJIRI 95 and ARNOLD 96. Supersedes DANEVICH 00. NODE=S076HDB;LINKAGE=VA

[40] Limit on 0ν decay of ^{116}Cd using enriched $CdWO_4$ scintillators. Supersedes DANEVICH 00. NODE=S076HDB;LINKAGE=VB

[41] Limit on 0ν decay of ^{116}Cd into first excited 2^+ state of daughter nucleus using enriched $CdWO_4$ scintillators. Supersedes DANEVICH 00. NODE=S076HDB;LINKAGE=VC

[42] Limit on 0ν decay of ^{116}Cd into first excited 0^+ state of daughter nucleus using enriched $CdWO_4$ scintillators. Supersedes DANEVICH 00. NODE=S076HDB;LINKAGE=VD

[43] Limit on 0ν decay of ^{116}Cd into second excited 0^+ state of daughter nucleus using enriched $CdWO_4$ scintillators. Supersedes DANEVICH 00. NODE=S076HDB;LINKAGE=VE

[44] Results of the Heidelberg-Moscow experiment (KLAPDOR-KLEINGROTHAUS 01 and GUENTHER 97) are reanalyzed using a new simulation of the complete background spectrum. The $\beta\beta2\nu$-decay rate is deduced from a 41.57 kg-y exposure. The result is in agreement and supersedes the above referenced halflives with similar statistical and systematic errors. NODE=S076HDB;LINKAGE=DO

[45] AALSETH 02B limit is based on 117 mol·yr of data using enriched Ge detectors. Background reduction by means of pulse shape analysis is applied to part of the data set. Reported limit is slightly less restrictive than that in KLAPDOR-KLEINGROTHAUS 01 However, it excludes part of the allowed half-life range reported in KLAPDOR-KLEINGROTHAUS 01B for the same nuclide. The analysis has been criticized in KLAPDOR-KLEINGROTHAUS 04B. The criticism was addressed and disputed in AALSETH 04. NODE=S076HDB;LINKAGE=SH

[46] BERNABEI 02D report a limit for the 0ν, $0^+ \rightarrow 0^+$ decay of ^{134}Xe, present in the source at 17%, by considering the maximum number of events for this mode compatible with the fitted smooth background. NODE=S076HDB;LINKAGE=D5

[47] BERNABEI 02D report a limit for the 0ν, $0^+ \rightarrow 0^+$ decay of ^{136}Xe, by considering the maximum number of events for this mode compatible with the fitted smooth background. The quoted sensitivity is 450×10^{21} yr. The Feldman and Cousins method is used to obtain the quoted limit. NODE=S076HDB;LINKAGE=D6

[48] KLAPDOR-KLEINGROTHAUS 02D is an expanded version of KLAPDOR-KLEINGROTHAUS 01B. The authors re-evaluate the data collected by the Heidelberg-Moscow experiment (KLAPDOR-KLEINGROTHAUS 01) and present a more detailed description of their analysis of an excess of counts at the energy expected for neutrinoless double-beta decay. They interpret this excess, which has a significance of 2.2 to 3.1 σ depending on the data analysis, as evidence for the observation of Lepton Number violation and violation of Baryon minus Lepton Number. The analysis has been criticized by AALSETH 02 and others. The criticisms have been addressed in KLAPDOR-KLEINGROTHAUS 02. See also KLAPDOR-KLEINGROTHAUS 02B. GROMOV 06 analysis of the background supports the assignment of the weak γ transitions near 2040 keV to the decay of ^{214}Bi as claimed in KLAPDOR-KLEINGROTHAUS 04A and KLAPDOR-KLEINGROTHAUS 04C, and in earlier works. NODE=S076HDB;LINKAGE=KX

[49] ASHITKOV 01 result for 2ν of ^{100}Mo is in agreement with other determinations of that halflife. NODE=S076HDB;LINKAGE=1S

[50] ASHITKOV 01 result for 0ν of ^{100}Mo is less stringent than EJIRI 01. NODE=S076HDB;LINKAGE=S2

[51] DANEVICH 01 place limit on 0ν decay of ^{160}Gd using Gd_2SiO_5:Ce crystal scintillators. The limit is more stringent than KOBAYASHI 95. NODE=S076HDB;LINKAGE=TH

[52] DANEVICH 01 place limits on 0ν decay of ^{160}Gd into excited 2^+ state of daughter nucleus using Gd_2SiO_5:Ce crystal scintillators. NODE=S076HDB;LINKAGE=TI

[53] DEBRAECKELEER 01 performed an inclusive measurement of the $\beta\beta$ decay into the second excited state of the daughter nucleus. A novel coincidence technique counting the de-excitation photons is employed. The result agrees with BARABASH 95. NODE=S076HDB;LINKAGE=RB

[54] EJIRI 01 uses tracking calorimeter and isotopically enriched passive source. Efficiencies were calculated assuming $\langle m_\nu \rangle$, $\langle \lambda \rangle$, or $\langle \eta \rangle$ driven decay. This is a continuation of EJIRI 96 which it supersedes. NODE=S076HDB;LINKAGE=EJ

[55] KLAPDOR-KLEINGROTHAUS 01 is a continuation of the work published in BAUDIS 99. Isotopically enriched Ge detectors are used in calorimetric measurement. The most stringent bound is derived from the data set in which pulse-shape analysis has been used to reduce background. Exposure time is 35.5 kg y. Supersedes BAUDIS 99 as most stringent result. NODE=S076HDB;LINKAGE=K1

[56] KLAPDOR-KLEINGROTHAUS 01 is a measurement of the $\beta\beta2\nu$-decay rate with higher statistics than GUENTHER 97. The reported value has a larger systematic error than their previous result. NODE=S076HDB;LINKAGE=K2

[57] WIESER 01 reports an inclusive geochemical measurement of ^{96}Zr $\beta\beta$ half life. Their result agrees within 2σ with ARNOLD 99 but only marginally, within 3σ, with KAWASHIMA 93. NODE=S076HDB;LINKAGE=WR

[58] BRUDANIN 00 determine the 2ν halflife of ^{48}Ca. Their value is less accurate than BALYSH 96. NODE=S076HDB;LINKAGE=P2

[59] ARNOLD 99 measure directly the 2ν decay of Zr for the first time, using the NEMO-2 tracking detector and an isotopically enriched source. The lifetime is more accurate than the geochemical result of KAWASHIMA 93. NODE=S076HDB;LINKAGE=LD

[60] ARNOLD 98 measure the 2ν decay of ^{82}Se by comparing the spectra in an enriched and natural selenium source using the NEMO-2 tracking detector. The measured half-life is in agreement, perhaps slightly shorter, than ELLIOTT 92. NODE=S076HDB;LINKAGE=R1

[61] ARNOLD 98 determine the limit for 0ν decay to the ground state of ^{82}Se using the NEMO-2 tracking detector. The half-life limit is in agreement, but less stringent, than ELLIOTT 92. NODE=S076HDB;LINKAGE=R2

[62] ARNOLD 98 determine the limit for 0ν decay to the excited 2^+ state of ^{82}Se using the NEMO-2 tracking detector. NODE=S076HDB;LINKAGE=R3

[63] ALSTON-GARNJOST 97 report evidence for 2ν decay of ^{100}Mo. This decay has been also observed by EJIRI 91, DASSIE 95, and DESILVA 97. NODE=S076HDB;LINKAGE=ZC

[64] DESILVA 97 result for 2ν decay of ^{100}Mo is in agreement with ALSTON-GARNJOST 97 and DASSIE 95. This measurement has the smallest errors. NODE=S076HDB;LINKAGE=DA

[65] DESILVA 97 result for 2ν decay of ^{150}Nd is in marginal agreement with ARTEMEV 93. It has smaller errors. NODE=S076HDB;LINKAGE=DC

[66] DESILVA 97 do not explain whether their efficiency for 0ν decay of ^{150}Nd was calculated under the assumption of a $\langle m_\nu\rangle$, $\langle\lambda\rangle$, or $\langle\eta\rangle$ driven decay. NODE=S076HDB;LINKAGE=DD

[67] ARNOLD 96 measure the 2ν decay of ^{116}Cd. This result is in agreement with EJIRI 95, but has smaller errors. Supersedes ARNOLD 95. NODE=S076HDB;LINKAGE=AR

[68] BALYSH 96 measure the 2ν decay of ^{48}Ca, using a passive source of enriched ^{48}Ca in a TPC. NODE=S076HDB;LINKAGE=Y2

[69] TAKAOKA 96 measure the geochemical half-life of ^{130}Te. Their value is in disagreement with the quoted values of BERNATOWICZ 92 and KIRSTEN 83; but agrees with several other unquoted determinations, e.g., MANUEL 91. NODE=S076HDB;LINKAGE=J3

[70] BARABASH 95 cannot distinguish 0ν and 2ν, but it is inferred indirectly that the 0ν mode accounts for less than 0.026% of their event sample. They also note that their result disagrees with the previous experiment by the NEMO group (BLUM 92). NODE=S076HDB;LINKAGE=Y1

[71] BERNATOWICZ 92 finds ^{128}Te/^{130}Te activity ratio from slope of ^{128}Xe/^{132}Xe vs ^{130}Xe/^{132}Xe ratios during extraction, and normalizes to lead-dated ages for the ^{130}Te lifetime. The authors state that their results imply that "(a) the double beta decay of ^{128}Te has been firmly established and its half-life has been determined ... without any ambiguity due to trapped Xe interferences... (b) Theoretical calculations ... underestimate the [long half-lives of ^{128}Te ^{130}Te] by 1 or 2 orders of magnitude, pointing to a real suppression in the 2ν decay rate of these isotopes. (c) Despite [this], most $\beta\beta$-models predict a *ratio* of 2ν decay widths ... in fair agreement with observation." Further details of the experiment are given in BERNATOWICZ 93. Our listed half-life has been revised downward from the published value by the authors, on the basis of reevaluated cosmic-ray ^{128}Xe production corrections. NODE=S076HDB;LINKAGE=BB

[72] TURKEVICH 91 observes activity in old U sample. The authors compare their results with theoretical calculations. They state "Using the phase-space factors of Boehm and Vogel (BOEHM 87) leads to matrix element values for the ^{238}U transition in the same range as deduced for ^{130}Te and ^{76}Ge. On the other hand, the latest theoretical estimates (STAUDT 90) give an upper limit that is 10 times lower. This large discrepancy implies either a defect in the calculations or the presence of a faster path than the standard two-neutrino mode in this case." See BOEHM 87 and STAUDT 90. NODE=S076HDB;LINKAGE=S

[73] Result agrees with direct determination of ELLIOTT 92. NODE=S076HDB;LINKAGE=LN

[74] Inclusive half life inferred from mass spectroscopic determination of abundance of $\beta\beta$-decay product ^{130}Te in mineral kitkaite (NiTeSe). Systematic uncertainty reflects variations in U-Xe gas-retention-age derived from different uranite samples. Agrees with geochemical determination of TAKAOKA 96 and direct measurement of ARNABOLDI 03. Inconsistent with results of KIRSTEN 83 and BERNATOWICZ 92. NODE=S076HDB;LINKAGE=LO

[75] Ratio of inclusive double beta half lives of ^{128}Te and ^{130}Te determined from minerals melonite ($NiTe_2$) and altaite (PbTe) by means of mass spectroscopic measurement of abundance of $\beta\beta$-decay products. As gas-retention-age could not be determined the authors use half life of ^{130}Te (LIN 88) to infer the half life of ^{128}Te. No estimate of the systematic uncertainty of this method is given. The directly determined half life ratio agrees with BERNATOWICZ 92. However, the inferred ^{128}Te half life disagrees with KIRSTEN 83 and BERNATOWICZ 92. NODE=S076HDB;LINKAGE=LP

[76] KIRSTEN 83 reports "2σ" error. References are given to earlier determinations of the ^{130}Te lifetime. NODE=S076HDB;LINKAGE=D

$\langle m_\nu \rangle$, The Effective Weighted Sum of Majorana Neutrino Masses Contributing to Neutrinoless Double-β Decay

NODE=S076MW

$\langle m_\nu \rangle = |\Sigma\ U^2_{1j} m_{\nu_j}|$, where the sum goes from 1 to n and where n = number of neutrino generations, and ν_j is a Majorana neutrino. Note that U^2_{ej}, not $|U_{ej}|^2$, occurs in the sum. The possibility of cancellations has been stressed. In the following Listings, only best or comparable limits or lifetimes for each isotope are reported.

NODE=S076MW

NODE=S076MW

VALUE (eV)	*CL%*	*ISOTOPE*	*TRANSITION*	*METHOD*	*DOCUMENT ID*		
• • • We do not use the following data for averages, fits, limits, etc. • • •							
< 0.19–0.68	90	^{130}Te	0ν	TeO_2 bolometer	77 ARNABOLDI	08	
< 3.5–22	90	^{48}Ca	0ν	CaF_2 scint.	78 UMEHARA	08	
< 9.3–60	90	^{100}Mo	$0^+ \to 0^+_1$	NEMO-3	79 ARNOLD	07	
< 6500	90	^{100}Mo	$0^+ \to 2^+$	NEMO-3	80 ARNOLD	07	OCCUR=2
0.32±0.03	68	^{76}Ge	0ν	Enriched HPGe	81 KLAPDOR-K...	06A	
< 0.2–1.1	90	^{130}Te		Cryog. det.	82 ARNABOLDI	05	
< 0.7–2.8	90	^{100}Mo	0ν	NEMO-3	83 ARNOLD	05A	
< 1.7–4.9	90	^{82}Se	0ν	NEMO-3	84 ARNOLD	05A	OCCUR=2
< 0.37–1.9	90	^{130}Te		Cryog. det.	85 ARNABOLDI	04	
< 0.8–1.2	90	^{100}Mo	0ν	NEMO-3	86 ARNOLD	04	
< 1.5–3.1	90	^{82}Se	0ν	NEMO-3	86 ARNOLD	04	OCCUR=2
0.1–0.9	99.7	^{76}Ge		Enriched HP Ge	87 KLAPDOR-K...	04A	
< 7.2–44.7	90	^{48}Ca		CaF_2 scint.	88 OGAWA	04	
< 1.1–2.6	90	^{130}Te		Cryog. det.	89 ARNABOLDI	03	
< 1.5–1.7	90	^{116}Cd	0ν	$^{116}CdWO_4$ scint.	90 DANEVICH	03	
< 0.33–1.35	90			Enriched HPGe	91 AALSETH	02B	
<2.9	90	^{136}Xe	0ν	Liquid Xe Scint.	92 BERNABEI	02D	
$0.39^{+0.17}_{-0.28}$		^{76}Ge	0ν	Enriched HPGe	93 KLAPDOR-K...	02D	
< 2.1–4.8	90	^{100}Mo	0ν	ELEGANT V	94 EJIRI	01	
< 0.35	90	^{76}Ge		Enriched HPGe	95 KLAPDOR-K...	01	
<23	90	^{96}Zr		NEMO-2	96 ARNOLD	99	
< 1.1–1.5		^{128}Te		Geochem	97 BERNATOW...	92	
<5	68	^{82}Se		TPC	98 ELLIOTT	92	
<8.3	76	^{48}Ca	0ν	CaF_2 scint.	YOU	91	

77 Limit was obtained using high resolution TeO_2 bolometer calorimeter to search for double beta decay of ^{130}Te. Reported range of limits reflects spread of matrix element calculations used. Supersedes ARNABOLDI 05. NODE=S076MW;LINKAGE=AL

78 Limit was obtained using CaF_2 scintillation calorimeter to search for double beta decay of ^{48}Ca. Reported range of limits reflects spread of QRPA and SM matrix element calculations used. Supersedes OGAWA 04. NODE=S076MW;LINKAGE=UE

79 ARNOLD 07 use NEMO-3 half life limit for 0ν-decay of ^{100}Mo to the first excited 0^+_1-state of daughter nucleus to obtain neutrino mass limit. The spread reflects the choice of two different nuclear matrix elements. This limit is not competitive when compared to the decay to the ground state. NODE=S076MW;LINKAGE=NA

80 ARNOLD 07 use NEMO-3 half life limit for 0ν-decay of ^{100}Mo to the first excited 2^+-state of daughter nucleus to obtain neutrino mass limit. This limit is not competitive when compared to the decay to the ground state. NODE=S076MW;LINKAGE=NR

81 Re-analysis of data originally published in KLAPDOR-KLEINGROTHAUS 04A. Modified pulse shape analysis leads the authors to claim 6σ statistical evidence for observation of 0ν-decay. Authors use matrix element of STAUDT 90. Uncertainty of nuclear matrix element is not reflected in stated error. Supersedes KLAPDOR-KLEINGROTHAUS 04A. NODE=S076MW;LINKAGE=KR

82 Supersedes ARNABOLDI 04. Reported range of limits due to use of different nuclear matrix element calculations. NODE=S076MW;LINKAGE=AO

83 Mass limits reported in ARNOLD 05A are derived from ^{100}Mo data, obtained by the NEMO-3 collaboration. The range reflects the spread of matrix element calculations considered in this work. Supersedes ARNOLD 04. NODE=S076MW;LINKAGE=N1

84 Neutrino mass limits based on ^{82}Se data utilizing the NEMO-3 detector. The range reported in ARNOLD 05A reflects the spread of matrix element calculations considered in this work. Supersedes ARNOLD 04. NODE=S076MW;LINKAGE=N2

85 Supersedes ARNABOLDI 03. Reported range of limits due to use of different nuclear matrix element calculations. NODE=S076MW;LINKAGE=AN

86 ARNOLD 04 limit is based on the nuclear matrix elements of SIMKOVIC 99, STOICA 01 and CIVITARESE 03. NODE=S076MW;LINKAGE=A1

87 Supersedes KLAPDOR-KLEINGROTHAUS 02D. Event excess at $\beta\beta$-decay energy is used to derive Majorana neutrino mass using the nuclear matrix elements of STAUDT 90. The mass range shown is based on the authors evaluation of the uncertainties of the STAUDT 90 matrix element calculation. If this uncertainty is neglected, and only statistical errors are considered, the range in $\langle m \rangle$ becomes (0.2–0.6) eV at the 3 σ level. NODE=S076MW;LINKAGE=KL

[88] Calorimetric CaF_2 scintillator. Range of limits reflects authors' estimate of the uncertainty of the nuclear matrix elements. Replaces YOU 91 as the most stringest limit based on ^{48}Ca. NODE=S076MW;LINKAGE=OG

[89] Supersedes ALESSANDRELLO 00. Cryogenic calorimeter search. Reported a range reflecting uncertainty in nuclear matrix element calculations. NODE=S076MW;LINKAGE=RR

[90] Limit for $\langle m_\nu \rangle$ is based on the nuclear matrix elements of STAUDT 90 and ARNOLD 96. Supersedes DANEVICH 00. NODE=S076MW;LINKAGE=VF

[91] AALSETH 02B reported range of limits on $\langle m_\nu \rangle$ reflects the spread of theoretical nuclear matrix elements. Excludes part of allowed mass range reported in KLAPDOR-KLEINGROTHAUS 01B. NODE=S076MW;LINKAGE=SH

[92] BERNABEI 02D limit is based on the matrix elements of SIMKOVIC 02. The range of neutrino masses based on a variety of matrix elements is 1.1–2.9 eV. NODE=S076MW;LINKAGE=D5

[93] KLAPDOR-KLEINGROTHAUS 02D is a detailed description of the analysis of the data collected by the Heidelberg-Moscow experiment, previously presented in KLAPDOR-KLEINGROTHAUS 01B. Matrix elements in STAUDT 90 have been used. See the footnote in the preceding table for further details. See also KLAPDOR-KLEINGROTHAUS 02B. NODE=S076MW;LINKAGE=KX

[94] The range of the reported $\langle m_\nu \rangle$ values reflects the spread of the nuclear matrix elements. On axis value assuming $\langle \lambda \rangle = \langle \eta \rangle = 0$. NODE=S076MW;LINKAGE=EJ

[95] KLAPDOR-KLEINGROTHAUS 01 uses the calculation by STAUDT 90. Using several other models in the literature could worsen the limit up to 1.2 eV. This is the most stringent experimental bound on m_ν. It supersedes BAUDIS 99B. NODE=S076MW;LINKAGE=K1

[96] ARNOLD 99 limit based on the nuclear matrix elements of STAUDT 90. NODE=S076MW;LINKAGE=LD

[97] BERNATOWICZ 92 finds these majorana neutrino mass limits assuming that the measured geochemical decay width is a limit on the 0ν decay width. The range is the range found using matrix elements from HAXTON 84, TOMODA 87, and SUHONEN 91. Further details of the experiment are given in BERNATOWICZ 93. NODE=S076MW;LINKAGE=BB

[98] ELLIOTT 92 uses the matrix elements of HAXTON 84. NODE=S076MW;LINKAGE=L

Limits on Lepton-Number Violating ($V+A$) Current Admixture

NODE=S076ETA

NODE=S076ETA

For reasons given in the discussion at the beginning of this section, we list only results from 1989 and later. $\langle \lambda \rangle = \lambda \sum U_{ej} V_{ej}$ and $\langle \eta \rangle = \eta \sum U_{ej} V_{ej}$, where the sum is over the number of neutrino generations. This sum vanishes for massless or unmixed neutrinos. In the following Listings, only best or comparable limits or lifetimes for each isotope are reported.

NODE=S076ETA

$\langle \lambda \rangle$ (10^{-6})	CL%	$\langle \eta \rangle$ (10^{-8})	CL%	ISOTOPE	METHOD	DOCUMENT ID		
• • • We do not use the following data for averages, fits, limits, etc. • • •								
<120	90			^{100}Mo	$0^+ \to 2^+$	[99] ARNOLD	07	
$0.692^{+0.058}_{-0.056}$	68	$0.305^{+0.026}_{-0.025}$	68	^{76}Ge	Enriched HPGe	[100] KLAPDOR-K...	06A	
< 2.5	90			^{100}Mo	0ν, NEMO-3	[101] ARNOLD	05A	
< 3.8	90			^{82}Se	0ν, NEMO-3	[102] ARNOLD	05A	OCCUR=2
< 1.5–2.0	90			^{100}Mo	0ν, NEMO-3	[103] ARNOLD	04	
< 3.2–3.8	90			^{82}Se	0ν, NEMO-3	[104] ARNOLD	04	OCCUR=2
< 1.6–2.4	90	< 0.9–5.3	90	^{130}Te	Cryog. det.	[105] ARNABOLDI	03	
< 2.2	90	<2.5	90	^{116}Cd	$^{116}CdWO_4$ scint.	[106] DANEVICH	03	
< 3.2–4.7	90	< 2.4–2.7	90	^{100}Mo	ELEGANT V	[107] EJIRI	01	
< 1.1	90	<0.64	90	^{76}Ge	Enriched HPGe	[108] GUENTHER	97	
< 4.4	90	<2.3	90	^{136}Xe	TPC	[109] VUILLEUMIER	93	
		<5.3		^{128}Te	Geochem	[110] BERNATOW...	92	

[99] ARNOLD 07 use NEMO-3 half life limit for 0ν-decay of ^{100}Mo to the first excited 2^+-state of daughter nucleus to limit the right-right handed admixture of weak currents $\langle \lambda \rangle$. This limit is not competitive when compared to the decay to the ground state. NODE=S076ETA;LINKAGE=NA

[100] Re-analysis of data originally published in KLAPDOR-KLEINGROTHAUS 04A. Modified pulse shape analysis leads the authors to claim 6σ statistical evidence for observation of 0ν-decay. Authors use matrix element of MUTO 89 to determine $\langle \lambda \rangle$ and $\langle \eta \rangle$. Uncertainty of nuclear matrix element is not reflected in stated errors. NODE=S076ETA;LINKAGE=KR

[101] ARNOLD 05A derive limit for $\langle \lambda \rangle$ based on ^{100}Mo data collected with NEMO-3 detector. No limit for $\langle \eta \rangle$ is given. Supersedes ARNOLD 04. NODE=S076ETA;LINKAGE=N1

[102] ARNOLD 05A derive limit for $\langle \lambda \rangle$ based on ^{82}Se data collected with NEMO-3 detector. No limit for $\langle \eta \rangle$ is given. Supersedes ARNOLD 04. NODE=S076ETA;LINKAGE=N2

[103] ARNOLD 04 use the matrix elements of SUHONEN 94 to obtain a limit for $\langle \lambda \rangle$, no limit for $\langle \eta \rangle$ is given. This limit is more stringent than the limit in EJIRI 01 for the same nucleus. NODE=S076ETA;LINKAGE=A1

[104] ARNOLD 04 use the matrix elements of TOMODA 91 and SUHONEN 91 to obtain a limit for $\langle \lambda \rangle$, no limit for $\langle \eta \rangle$ is given. NODE=S076ETA;LINKAGE=A2

[105] Supersedes ALESSANDRELLO 00. Cryogenic calorimeter search. Reported a range reflecting uncertainty in nuclear matrix element calculations. NODE=S076ETA;LINKAGE=RR

[106] Limits for $\langle \lambda \rangle$ and $\langle \eta \rangle$ are based on nuclear matrix elements of STAUDT 90. Supersedes DANEVICH 00. NODE=S076ETA;LINKAGE=VG

107 The range of the reported $\langle \lambda \rangle$ and $\langle \eta \rangle$ values reflects the spread of the nuclear matrix elements. On axis value assuming $\langle m_{\nu} \rangle$=0 and $\langle \lambda \rangle$=$\langle \eta \rangle$=0, respectively. NODE=S076ETA;LINKAGE=EJ

108 GUENTHER 97 limits use the matrix elements of STAUDT 90. Supersedes BALYSH 95 and BALYSH 92. NODE=S076ETA;LINKAGE=J1

109 VUILLEUMIER 93 uses the matrix elements of MUTO 89. Based on a half-life limit 2.6×10^{23} y at 90%CL. NODE=S076ETA;LINKAGE=V

110 BERNATOWICZ 92 takes the measured geochemical decay width as a limit on the 0ν width, and uses the SUHONEN 91 coefficients to obtain the least restrictive limit on η. Further details of the experiment are given in BERNATOWICZ 93. NODE=S076ETA;LINKAGE=BB

Double-β Decay REFERENCES

NODE=S076

ARNABOLDI	08	PR C78 035502	C. Arnaboldi *et al.*		REFID=52519
BELLI	08	PL B658 193	P. Belli *et al.*	(INFN Gran Sasso)	REFID=52084
BELLI	08B	EPJ A36 167	P. Belli *et al.*		REFID=52497
UMEHARA	08	PR C78 058501	S. Umehara *et al.*		REFID=52522
ARNOLD	07	NP A781 209	R. Arnold *et al.*	(NEMO-3 Collab.)	REFID=51598
BARABASH	07	NP A785 371	A.S. Barabash *et al.*		REFID=51770
BARABASH	07B	JPG 34 1721	A.S. Barabash *et al.*		REFID=51964
BLOXHAM	07	PR C76 025501	T. Bloxham *et al.*	(COBRA Collab.)	REFID=51848
GROMOV	06	PPNL 3 157	K. Gromov *et al.*		REFID=51382
KLAPDOR-K...	06A	MPL A21 1547	H.V. Klapdor-Kleingrothaus, I.V. Krivosheina		REFID=51401
ARNABOLDI	05	PRL 95 142501	C. Arnaboldi *et al.*	(CUORICINO Collab.)	REFID=50930
ARNOLD	05A	PRL 95 182302	R. Arnold *et al.*	(NEMO-3 Collab.)	REFID=50934
AALSETH	04	PR D70 078302	C.E. Aalseth *et al.*		REFID=50235
ARNABOLDI	04	PL B584 260	C. Arnaboldi *et al.*		REFID=49824
ARNOLD	04	JETPL 80 377	R. Arnold *et al.*	(NEMO3 Detector Collab.)	REFID=50411
		Translated from ZETFP 80 429.			
BARABASH	04	JETPL 79 10	A.S. Barabash *et al.*		REFID=50551
KLAPDOR-K...	04A	PL B586 198	H.V. Klapdor-Kleingrothaus *et al.*		REFID=49889
KLAPDOR-K...	04B	PR D70 078301	H.V. Klapdor-Kleingrothaus, A. Dietz, I.V. Krivosheina		REFID=50234
KLAPDOR-K...	04C	NIM A522 371	H.V. Klapdor-Kleingrothaus *et al.*		REFID=50554
OGAWA	04	NP A730 215	I. Ogawa *et al.*		REFID=49796
ARNABOLDI	03	PL B557 167	C. Arnaboldi *et al.*		REFID=49258
CIVITARESE	03	NP A729 867	O. Civitarese, J. Suhonen		REFID=49794
DANEVICH	03	PR C68 035501	F.A. Danevich *et al.*		REFID=49600
DOERR	03	NIM A513 596	C. Doerr, H.V. Klapdor-Kleingrothaus		REFID=50553
AALSETH	02	MPL A17 1475	C.E. Aalseth *et al.*		REFID=48629
AALSETH	02B	PR D65 092007	C.E. Aalseth *et al.*	(IGEX Collab.)	REFID=48793
BERNABEI	02D	PL B546 23	R. Bernabei *et al.*	(DAMA Collab.)	REFID=48990
KLAPDOR-K...	02	hep-ph/0205228	H.V. Klapdor-Kleingrothaus		REFID=48688
KLAPDOR-K...	02B	PPNL 110 57	H.V. Klapdor-Kleingrothaus, A. Dietz, I.V. Krivosheina		REFID=49114
KLAPDOR-K...	02D	FP 32 1181	H.V. Klapdor-Kleingrothaus, A. Dietz, I.V. Krivosheina		REFID=49193
SIMKOVIC	02	hep-ph/0204278	F. Simkovic, P. Domin, A. Faessler		REFID=49198
ASHITKOV	01	JETPL 74 529	V.D. Ashitkov *et al.*		REFID=48730
		Translated from ZETFP 74 601.			
DANEVICH	01	NP A694 375	F.A. Danevich *et al.*		REFID=48113
DEBRAECKEL...	01	PRL 86 3510	L. De Braeckeleer *et al.*		REFID=48467
EJIRI	01	PR C63 065501	H. Ejiri *et al.*		REFID=48163
KLAPDOR-K...	01	EPJ A12 147	H.V. Klapdor-Kleingrothaus *et al.*		REFID=48460
KLAPDOR-K...	01B	MPL A16 2409	H.V. Klapdor-Kleingrothaus *et al.*		REFID=48628
STOICA	01	NP A694 269	S. Stoica, H.V. Klapdor-Kleingrothous		REFID=50526
WIESER	01	PR C64 024308	M.E. Wieser, J.R. De Laeter		REFID=48224
ALESSAND...	00	PL B486 13	A. Alessandrello *et al.*		REFID=47726
BRUDANIN	00	PL B495 63	V.B. Brudanin *et al.*		REFID=47917
DANEVICH	00	PR C62 045501	F.A. Danevich *et al.*		REFID=47810
ARNOLD	99	NP A658 299	R. Arnold *et al.*	(NEMO Collab.)	REFID=47325
BAUDIS	99	PR D59 022001	L. Baudis *et al.*	(Heidelberg-Moscow Collab.)	REFID=46702
BAUDIS	99B	PRL 83 41	L. Baudis *et al.*	(Heidelberg-Moscow Collab.)	REFID=47031
SIMKOVIC	99	PR C60 055502	F. Simkovic *et al.*		REFID=50525
ARNOLD	98	NP A636 209	R. Arnold *et al.*	(NEMO-2 Collab.)	REFID=49250
ALSTON-...	97	PR C55 474	M. Alston-Garnjost *et al.*	(LBL, MTHO+)	REFID=45230
DESILVA	97	PR C56 2451	A. de Silva *et al.*	(UCI)	REFID=45865
GUENTHER	97	PR D55 54	M. Gunther *et al.*	(Heidelberg-Moscow Collab.)	REFID=45265
ARNOLD	96	ZPHY C72 239	R. Arnold *et al.*	(BCEN, CAEN, JINR+)	REFID=45277
BALYSH	96	PRL 77 5186	A. Balysh *et al.*	(KIAE, UCI, CIT)	REFID=45405
EJIRI	96	NP A611 85	H. Ejiri *et al.*	(OSAK)	REFID=45334
TAKAOKA	96	PR C53 1557	N. Takaoka, Y. Motomura, K. Nagao	(KYUSH, OKAY)	REFID=45407
ARNOLD	95	JETPL 61 170	R.G. Arnold *et al.*	(NEMO Collab.)	REFID=44332
		Translated from ZETFP 61 168.			
BALYSH	95	PL B356 450	A. Balysh *et al.*	(Heidelberg-Moscow Collab.)	REFID=44401
BARABASH	95	PL B345 408	A.S. Barabash *et al.*	(ITEP, SCUC, PNL+)	REFID=44334
DASSIE	95	PR D51 2090	D. Dassie *et al.*	(NEMO Collab.)	REFID=44148
EJIRI	95	JPSJ 64 339	H. Ejiri *et al.*	(OSAK, KIEV)	REFID=44333
KOBAYASHI	95	NP A586 457	M. Kobayashi, M. Kobayashi	(KEK, SAGA)	REFID=44175
SUHONEN	94	PR C49 3055	J. Suhonen, O. Civitarese		REFID=50527
ARTEMEV	93	JETPL 58 262	V.A. Artemiev *et al.*	(ITEP, INRM)	REFID=44331
		Translated from ZETFP 58 256.			
BERNATOW...	93	PR C47 806	T. Bernatowicz *et al.*	(WUSL, TATA)	REFID=43219
KAWASHIMA	93	PR C47 R2452	A. Kawashima, K. Takahashi, A. Masuda	(TOKYC+)	REFID=44329
VUILLEUMIER	93	PR D48 1009	J.C. Vuilleumier *et al.*	(NEUC, CIT, VILL)	REFID=43446
BALYSH	92	PL B283 32	A. Balysh *et al.*	(MPIH, KIAE, SASSO)	REFID=42083
BERNATOW...	92	PRL 69 2341	T. Bernatowicz *et al.*	(WUSL, TATA)	REFID=42181
BLUM	92	PL B275 506	D. Blum *et al.*	(NEMO Collab.)	REFID=41979
ELLIOTT	92	PR C46 1535	S.R. Elliott *et al.*	(UCI)	REFID=42190
EJIRI	91	PL B258 17	H. Ejiri *et al.*	(OSAK)	REFID=41483
MANUEL	91	JPG 17 S221	O.K. Manuel	(MISSR)	REFID=45408
SUHONEN	91	NP A535 509	J. Suhonen, S.B. Khadkikar, A. Faessler	(JYV+)	REFID=41945
TOMODA	91	RPP 54 53	T. Tomoda		REFID=45867
TURKEVICH	91	PRL 67 3211	A. Turkevich, T.E. Economou, G.A. Cowan	(CHIC+)	REFID=41832
YOU	91	PL B265 53	K. You *et al.*	(BHEP, CAST+)	REFID=41598
STAUDT	90	EPL 13 31	A. Staudt, K. Muto, H.V. Klapdor-Kleingrothaus		REFID=41834
MUTO	89	ZPHY A334 187	K. Muto, E. Bender, H.V. Klapdor	(TINT, MPIH)	REFID=43301
LIN	88	NP A481 477	W.J. Lin *et al.*		REFID=49526
LIN	88B	NP A481 484	W.J. Lin *et al.*		REFID=49527
BOEHM	87	Massive Neutrinos	F. Bohm, P. Vogel	(CIT)	REFID=41833
Cambridge Univ. Press, Cambridge					
TOMODA	87	PL B199 475	T. Tomoda, A. Faessler	(TUBIN)	REFID=43303
HAXTON	84	PPNP 12 409	W.C. Haxton, G.J. Stevenson		REFID=43302
KIRSTEN	83	PRL 50 474	T. Kirsten, H. Richter, E. Jessberger	(MPIH)	REFID=10425