



Labeling Format Information:			
Fonts: Universe Cond regular, bold and bold italic.			
Drug Facts:	8.5 pt	Leading:	6.5 pt
Header:	8 pt	Bullets:	5 pt
Subheader:	6 pt	Barlines:	2.5 pt
Body Text:	6 pt	Hairlines:	0.5 pt
Drug Facts (continued):	8 pt		
Horizontal Scale: 90%		Average Kerning: 0	

**BLISTER CARTON PEPCID MAX STRENGTH FCT 5 (VERSION E)**  
**3004095 A-21**  
**JULY 17, 2003**

00000000



**Drug Facts**

**Active ingredient (in each tablet)**  
Famotidine 20 mg.....Acid reducer

**Uses**  
• relieves heartburn associated with acid indigestion and sour stomach  
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**  
**Do not use**  
• if you have trouble swallowing  
• with other acid reducers  
• if you have kidney disease, except under the advice and supervision of a doctor



**Stop use and ask a doctor if**  
• stomach pain continues • you need to take this product for more than 14 days

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
• adults and children 12 years and over:  
• to **prevent** symptoms, swallow 1 tablet with a glass of water  
• to **relieve** symptoms, swallow 1 tablet with a glass of water from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn  
• do not use more than 2 tablets in 24 hours  
• children under 12 years: ask a doctor

**TABLETS**  
**Pepcid AC**  
**MAXIMUM STRENGTH**

**Do not use if the individual blister unit is open or torn.**  
• 1 tablet relieves heartburn due to acid indigestion (read Package insert before use).  
• PEPCID AC prevents heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.  
*The makers of Pepcid AC do not manufacture store brands.*



**Pepcid AC** **MAXIMUM STRENGTH**

25 TABLETS

00000000

**NEW! MAXIMUM STRENGTH**

NDC 16837-855-25

**Pepcid AC**  
Famotidine Tablets 20mg Acid Reducer

**Just One Tablet**

**Prevents & Relieves Heartburn Due to Acid Indigestion**

25 TABLETS

**MAXIMUM STRENGTH**

**Pepcid AC**

*The makers of Pepcid AC do not manufacture store brands*

Distributed by:  
Johnson & Johnson • MERCK  
CONSUMER PHARMACEUTICALS CO.  
FORT WASHINGTON, PA 19094 USA  
Generic and trademark of Merck & Co., Inc.  
Please visit our web site at:  
www.pepcidac.com

7 16837 85525 5

The Pepcid AC Brand name is a guarantee of our commitment to you. We have worked hard to earn your trust and we'll work even harder to maintain it.

Johnson & Johnson • MERCK • **Pepcid AC** • Allentown, PA • Committed



**Drug Facts (continued)**

**Other information**  
• read the directions and warnings before use  
• keep the carton and package insert. They contain important information.  
• store at 20° - 30° C (68° - 86° F)  
• protect from moisture

**Inactive ingredients**  
carnauba wax, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

**Questions or comments? 1-800-755-4008**

**Labeling Format Information:**

<b>Fonts:</b> Univers Condensed regular, bold and bold oblique.	
<b>Drug Facts:</b> 10 pt	<b>Leading:</b> 6.5 pt
<b>Header:</b> 8 pt	<b>Bullets:</b> 5 pt
<b>Subheader:</b> 6 pt	<b>Barlines:</b> 2.5 pt
<b>Body Text:</b> 6 pt	<b>Hairlines:</b> 0.5 pt
<b>Drug Facts (continued):</b> 8 pt	

Horizontal Scale: 95%      Average Kerning: 0

Do not use if printed foil seal under bottle cap is open or torn.

**Active ingredient (in each tablet)** ..... **Purpose**  
Famotidine 20 mg ..... Acid reducer

**Uses** • relieves heartburn associated with acid indigestion and sour stomach  
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings** **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers.

**Do not use**  
• if you have trouble swallowing • with other acid reducers • if you have kidney disease, except under the advice and supervision of a doctor

**Stop use and ask a doctor if • stomach pain continues**

LOT  
EXP.

000000

**MAXIMUM STRENGTH**

NDC 16837-855-50

**Pepcid<sup>®</sup> AC**

Famotidine Tablets 20mg Acid Reducer

**Just One Tablet**

**Prevents & Relieves Heartburn Due to Acid Indigestion**

50 TABLETS

• you need to take this product for more than 14 days  
**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water • to prevent symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes** before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor. **Other information** • read the directions, warnings and accompanying product information before use • keep the carton and package insert. They contain important information. • store at 20° - 30°C (68° - 86°F) • protect from moisture

Dist. by: **Johnson & Johnson • MERCK**  
CONSUMER PHARMACEUTICALS CO. ©Registered trademark  
FORT WASHINGTON, PA 19034 USA of Merck & Co., Inc.

150% OF SIZE

Do not use if printed foil seal under bottle cap is open or torn.

**Active ingredient (in each tablet)** ..... **Purpose**  
Famotidine 20 mg ..... Acid reducer

**Uses** • relieves heartburn associated with acid indigestion and sour stomach  
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings** **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers.

**Do not use**  
• if you have trouble swallowing • with other acid reducers • if you have kidney disease, except under the advice and supervision of a doctor

**Stop use and ask a doctor if • stomach pain continues**

LOT  
EXP.

000000

**MAXIMUM STRENGTH**

NDC 16837-855-50

**Pepcid<sup>®</sup> AC**

Famotidine Tablets 20mg Acid Reducer

**Just One Tablet**

**Prevents & Relieves Heartburn Due to Acid Indigestion**

50 TABLETS

• you need to take this product for more than 14 days  
**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water • to prevent symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes** before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor. **Other information** • read the directions, warnings and accompanying product information before use • keep the carton and package insert. They contain important information. • store at 20° - 30°C (68° - 86°F) • protect from moisture

Dist. by: **Johnson & Johnson • MERCK**  
CONSUMER PHARMACEUTICALS CO. ©Registered trademark  
FORT WASHINGTON, PA 19034 USA of Merck & Co., Inc.

**Labeling Format Information:**


Fonts: Universe Cond regular, bold and bold italic.

<b>Drug Facts:</b>	NA	<b>Leading:</b>	4.5 pt
<b>Header:</b>	NA	<b>Bullets:</b>	4.5 pt
<b>Subheader:</b>	4.5 pt	<b>Barlines:</b>	NA
<b>Body Text:</b>	4.5 pt	<b>Hairlines:</b>	NA
<b>Drug Facts (continued):</b>	NA		


Horizontal Scale: 77%

Average Kerning: -15

**LABEL PEPCID MAX STRENGTH FCT 50 (VERSION E)**  
**3004095 E-14**  
**JULY 18, 2003**



130% OF SIZE




Labeling Format Information:			
Fonts: Universe Cond regular, bold and bold italic.			
Drug Facts:	NA	Leading:	4.5 pt
Header:	4.5 pt	Bullets:	4.5 pt
Subheader:	4.5 pt	Barlines:	NA
Body Text:	4.5 pt	Hairlines:	NA
Drug Facts (continued):	NA		
Horizontal Scale: 76%		Average Kerning: -4	

**POUCH PEPCID MAX STRENGTH FCT CR 1 (VERSION E)**  
**3004095 G-5**  
**JULY 17, 2003**

FRONT

BACK




PEPCID AC MAX STRENGTH TRADE POUCH  
2/3/03

<b>Labeling Format Information:</b>	
<b>Fonts:</b> Helvetica Neue condensed and bold condensed Zapf Humanist 601BT bold and bold italic Humanist 777BT bold condensed	
<b>Drug Facts:</b>	<b>Leading:</b>
<b>Header:</b>	<b>Bullets:</b>
<b>Subheader:</b>	<b>Barlines:</b>
<b>Body Text:</b>	<b>Hairlines:</b>
<b>Drug Facts (continued):</b>	
<b>Avg Horizontal Scale:</b>	<b>Avg Kerning:</b>

Labeling Format Information:	
<b>Fonts:</b> Century Bold, Helvetica narrow, compressed, condensed bold, Zapf Humanist 601BT bold Humanist 777BT black italic	
<b>Drug Facts:</b>	<b>Leading:</b> 7 pt
<b>Header:</b>	<b>Bullets:</b>
<b>Subheader:</b>	<b>Barlines:</b>
<b>Body Text:</b> 6.5 pt	<b>Hairlines:</b>
<b>Drug Facts (continued):</b>	
<b>Avg Horizontal Scale:</b> 100%	<b>Avg Kerning:</b> 0

FILE SCALE: 100%



**Drug Facts**

**Active ingredient (in each tablet)**  
Famotidine 20 mg..... Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble swallowing
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

**Stop use and ask a doctor if**

- stomach pain continues
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years and over:
- to relieve symptoms, swallow 1 tablet with a glass of water
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**MAXIMUM STRENGTH**

# Pepcid<sup>®</sup> AC

75 TABLETS



8726570

**NEW! MAXIMUM STRENGTH**

NDC 16837-855-75

# Pepcid<sup>®</sup> AC

Famotidine Tablets 20mg Acid Reducer

## Just One Tablet

**Prevents & Relieves Heartburn Due to Acid Indigestion**

75 TABLETS



**MAXIMUM STRENGTH**

# Pepcid<sup>®</sup> AC

The makers of Pepcid AC do not manufacture store brands

Distributed by:  
Johnson & Johnson • MERCK  
CONSUMER PHARMACEUTICALS CO.  
PO BOX 1084  
KILGORE, TX 75132 USA

© 2003 Johnson & Johnson  
Please visit our web site at:  
[www.pepcidac.com](http://www.pepcidac.com)



7 16837 85575 0

**Drug Facts (continued)**

**Other information**

- read the directions and warnings before use
- keep the carton and package insert. They contain important information.
- store at 20° - 30°C (68° - 86°F)
- protect from moisture

**Inactive ingredients** carnauba wax, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

**Questions or comments?** 1-800-755-4008

**Do not use if carton is open or printed foil seal under bottle cap is open or torn.**

- 1 tablet relieves heartburn due to acid indigestion (Read Package insert before use).
- PEPCID AC prevents heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

**The makers of Pepcid AC do not manufacture store brands.**



**Labeling Format Information:**

<b>Fonts:</b> Universe Cond regular, bold and bold italic.	
<b>Drug Facts:</b> 8.5 pt	<b>Leading:</b> 6.5 pt
<b>Header:</b> 8 pt	<b>Bullets:</b> 5 pt
<b>Subheader:</b> 6 pt	<b>Barlines:</b> 2.5 pt
<b>Body Text:</b> 6 pt	<b>Hairlines:</b> 0.5 pt
<b>Drug Facts (continued):</b> 8 pt	
Horizontal Scale: 100%	Average Kerning: 0

**CTN PEPCID MAX STR 75ct CLUB CODE  
3004404 A-1  
JULY 30, 2003**

**Drug Facts**  
**Active ingredient (in each tablet)** Famotidine 20 mg.  
**Purpose** Acid reducer

**Uses**  
 • relieves heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages  
 • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**  
**Warning alert:** Do not use if you are allergic to famotidine or other acid reducers  
**Do not use**  
 • if you have trouble swallowing  
 • with other acid reducers  
 • if you have kidney disease, except under the advice and supervision of a doctor  
**Stop use and ask a doctor if**  
 • you need to take this product for more than 14 days  
**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 • adults and children 12 years and over:  
 • to relieve symptoms, swallow 1 tablet with a glass of water  
 • to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn  
 • do not use more than 2 tablets in 24 hours  
 • children under 12 years: ask a doctor

**Peppcid<sup>®</sup> AC** **MAXIMUM STRENGTH**



**50 TABLETS**

**NEW! MAXIMUM STRENGTH**

NDC 16837-855-50

**Peppcid<sup>®</sup> AC**

Famotidine Tablets 20mg Acid Reducer

**Just One Tablet**

**Prevents & Relieves Heartburn Due to Acid Indigestion**



**50 TABLETS**

**MAXIMUM STRENGTH**

**Peppcid<sup>®</sup> AC**

The makers of Peppcid AC do not manufacture store brands

Distributed by:  
 Johnson & Johnson • MERCK  
 CONSUMER PHARMACEUTICALS CO.  
 FORT WASHINGTON, PA 19004 USA  
 ©Registered trademark of Merck & Co., Inc.  
 Please visit our web site at:  
 www.peppcidac.com



7 16837 85550 7

**Drug Facts (continued)**

**Other information**  
 • read the directions and warnings before use  
 • keep the carton and package insert. They contain important information.  
 • store at 20° - 30° C (68° - 86° F)  
 • protect from moisture


**Inactive ingredients** carnauba wax, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

**Questions or comments?** 1-800-755-4008


**Do not use if carton is open or printed foil seal under bottle cap is open or torn.**

• 1 tablet relieves heartburn due to acid indigestion (Read Package Insert before use).  
 • PEPCID AC prevents heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.  
**The makers of Peppcid AC do not manufacture store brands.**

00000000



00000000



00000000

**Labeling Format Information:**

<b>Fonts:</b> Universe Cond regular, bold and bold italic.	
<b>Drug Facts:</b> 8.5 pt	<b>Leading:</b> 6.5 pt
<b>Header:</b> 8 pt	<b>Bullets:</b> 5 pt
<b>Subheader:</b> 6 pt	<b>Barlines:</b> 2.5 pt
<b>Body Text:</b> 6 pt	<b>Hairlines:</b> 0.5 pt
<b>Drug Facts (continued):</b> 8 pt	
Horizontal Scale: 100%	Average Kerning: 0

**BOTTLE CARTON PEPCID MAX STRENGTH FCT 50 (VERSION E)**  
**3004095 C-15**  
**JULY 18, 2003**



Do not use if printed foil seal under bottle cap is open or torn.

**Active ingredient (in each tablet)** ..... **Purpose**  
 Famotidine 20 mg ..... Acid reducer

**Uses** • relieves heartburn associated with acid indigestion and sour stomach  
 • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings** **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers.

**Do not use**  
 • if you have trouble swallowing • with other acid reducers • if you have kidney disease, except under the advice and supervision of a doctor

**Stop use and ask a doctor if** • stomach pain continues

LOT  
 EXP.

**7836340**

**MAXIMUM STRENGTH**

NDC 16837-855-75

**Pepcid<sup>®</sup>**  
 Famotidine Tablets 20mg Acid Reducer  
**Just One Tablet**  
**AC**

**Prevents & Relieves Heartburn Due to Acid Indigestion**

**75 TABLETS**

• you need to take this product for more than 14 days  
**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** • adults and children 12 years and over: • to **relieve** symptoms, swallow 1 tablet with a glass of water  
 • to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor. **Other information** • read the directions, warnings and accompanying product information before use • keep the carton and package insert. They contain important information. • store at 20° - 30°C (68° - 86°F) • protect from moisture

Dist. by: **Johnson & Johnson** • MERCK  
 CONSUMER PHARMACEUTICALS CO. ©Registered trademark of Merck & Co., Inc.  
 FORT WASHINGTON, PA 19034 USA

150% OF SIZE

Do not use if printed foil seal under bottle cap is open or torn.

**Active ingredient (in each tablet)** ..... **Purpose**  
 Famotidine 20 mg ..... Acid reducer

**Uses** • relieves heartburn associated with acid indigestion and sour stomach  
 • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings** **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers.

**Do not use**  
 • if you have trouble swallowing • with other acid reducers • if you have kidney disease, except under the advice and supervision of a doctor

**Stop use and ask a doctor if** • stomach pain continues

LOT  
 EXP.

**7836340**

**MAXIMUM STRENGTH**

NDC 16837-855-75

**Pepcid<sup>®</sup>**  
 Famotidine Tablets 20mg Acid Reducer  
**Just One Tablet**  
**AC**

**Prevents & Relieves Heartburn Due to Acid Indigestion**

**75 TABLETS**

• you need to take this product for more than 14 days  
**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** • adults and children 12 years and over: • to **relieve** symptoms, swallow 1 tablet with a glass of water  
 • to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor. **Other information** • read the directions, warnings and accompanying product information before use • keep the carton and package insert. They contain important information. • store at 20° - 30°C (68° - 86°F) • protect from moisture

Dist. by: **Johnson & Johnson** • MERCK  
 CONSUMER PHARMACEUTICALS CO. ©Registered trademark of Merck & Co., Inc.  
 FORT WASHINGTON, PA 19034 USA

**Labeling Format Information:**

**Fonts:** Universe Cond regular, bold and bold italic.

<b>Drug Facts:</b>	NA	<b>Leading:</b>	4.5 pt
<b>Header:</b>	NA	<b>Bullets:</b>	4.5 pt
<b>Subheader:</b>	4.5 pt	<b>Barlines:</b>	NA
<b>Body Text:</b>	4.5 pt	<b>Hairlines:</b>	NA
<b>Drug Facts (continued):</b>	NA		

Horizontal Scale: 77%

Average Kerning: -15

**LBL PEPCID MAX STR 75ct CLUB CODE  
 3004404 B-1  
 JULY 30, 2003**

Do not use if the individual blister unit is open or torn.

WHAT YOU SHOULD KNOW ABOUT:



### A non-prescription stomach medicine

MAXIMUM STRENGTH PEPCID® AC contains a medicine, different from antacids, that doctors have prescribed for years to treat acid-related problems in millions of people.

- Product Benefits:**
- 1 tablet relieves heartburn due to acid indigestion.
  - MAXIMUM STRENGTH PEPCID® AC prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods or beverages.
  - It contains famotidine, a prescription-proven medicine.




Shown actual size

### How to use MAXIMUM STRENGTH PEPCID® AC

Use MAXIMUM STRENGTH PEPCID® AC to relieve or prevent these symptoms:

- Heartburn
- Acid indigestion
- Sour stomach



#### TO RELIEVE SYMPTOMS

Swallow 1 tablet with a glass of water (do not chew).

#### TO PREVENT SYMPTOMS

Swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn.

MAXIMUM STRENGTH PEPCID® AC can be used up to twice daily (up to 2 tablets in 24 hours).


- Do not take more than 1 tablet at a time.
- Do not take more than 2 tablets in 24 hours.

### Excess acid: a burning problem

It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the wrong place (the esophagus), or too much acid, can cause burning pain and discomfort that interfere with everyday activities.

**Heartburn – Caused by acid in the esophagus**

A valve-like muscle called the lower esophageal sphincter (LES) is relaxed in an open position



#### Tips for Managing Heartburn

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

INSERT PEPCID MAX STRENGTH FCT 5 (VERSION 3) 3004316 B-5  
JULY 18, 2003

Just ONE  
and  
heartburn's done!



This coupon good only on purchase of product indicated. Any other use constitutes fraud. COUPON CANNOT BE BOUGHT, TRANSFERRED OR SOLD. LIMIT — ONE COUPON PER PURCHASE. VOID IF TAXED, RESTRICTED OR PROHIBITED BY LAW. To the retailer, Johnson & Johnson • Merck Consumer Pharmaceuticals Co. will reimburse you for the face value of this coupon plus 8¢ if submitted in compliance with Johnson & Johnson • Merck Consumer Pharmaceuticals Company Coupon Redemption Policy. Cash value 1/20th of one cent. Send coupons to Johnson • Merck Consumer Pharmaceuticals, Dept. 00045, 1 Fawcett Drive, Del Rio, TX 78840. ©JUMCP 2003.

Save \$2.00  
on your next purchase of  
any Pepcid® AC strength or Pepcid® Complete  
25 count or larger

MANUFACTURER COUPON NO EXPIRATION DATE  
00000000  
Registered trademark of Merck & Co., Inc.  
00000000  
Distributed by:  
Johnson & Johnson • MERCK  
CONSUMER PHARMACEUTICALS CO  
FORT WASHINGTON, PA 19034 USA

Please call 1-800-755-4008 for more information  
and a guide of how to help your heartburn.  
www.pepcidac.com

### Questions or comments?


• You need to take this product for more than 14 days.  
• You have trouble swallowing or stomach pain continues. You may have a serious condition that may need different treatment.  
• You are allergic to famotidine or other acid reducers. Do not use with other acid reducers. Do not use if you have kidney disease, except under the advice and supervision of a doctor. This product should not be given to children under 12 years old, unless directed by a doctor. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Heartburn and acid indigestion are common, but you should stop taking MAXIMUM STRENGTH PEPCID® AC and see your doctor promptly if:

### Know when to see your doctor

The ingredient in MAXIMUM STRENGTH PEPCID® AC, famotidine, has been prescribed by doctors for years to treat millions of patients safely and effectively. The active ingredient in MAXIMUM STRENGTH PEPCID® AC has been taken safely with many frequently prescribed medications.

### Excellent safety record



In clinical studies, MAXIMUM STRENGTH PEPCID® AC film-coated tablets were significantly better than placebo tablets (tablets without the medicine) in relieving and preventing heartburn.

### Proven effective in clinical studies

Do not use if the individual pouch is open or torn.

WHAT YOU SHOULD KNOW ABOUT:



### A non-prescription stomach medicine

MAXIMUM STRENGTH PEPCID® AC contains a medicine, different from antacids, that doctors have prescribed for years to treat acid-related problems in millions of people.

- Product Benefits:**
- 1 tablet relieves heartburn due to acid indigestion.
  - MAXIMUM STRENGTH PEPCID® AC prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods or beverages.
  - It contains famotidine, a prescription-proven medicine.



Shown actual size


### How to use MAXIMUM STRENGTH PEPCID® AC

Use MAXIMUM STRENGTH PEPCID® AC to relieve or prevent these symptoms:

- Heartburn
- Acid indigestion
- Sour stomach

MAXIMUM STRENGTH PEPCID® AC can be used up to twice daily (up to 2 tablets in 24 hours).

- Do not take more than 1 tablet at a time.
- Do not take more than 2 tablets in 24 hours.



#### TO RELIEVE SYMPTOMS

Swallow 1 tablet with a glass of water (do not chew).

#### TO PREVENT SYMPTOMS


Swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn.

### Excess acid: a burning problem

It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the wrong place (the esophagus), or too much acid, can cause burning pain and discomfort that interfere with everyday activities.

**Heartburn – Caused by acid in the esophagus**

A valve-like muscle called the lower esophageal sphincter (LES) is relaxed in an open position



Burning pain/discomfort

Excess acid moves up into esophagus

#### Tips for Managing Heartburn

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.



Name (please print) \_\_\_\_\_  
 Address \_\_\_\_\_  
 City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_ Apt. \_\_\_\_\_  
 Age:  under 35  35-49  50-54  55-64  65 yrs+  
 Please circle one:  Dr.  Mr.  Mrs.  Miss.

Yes, I'd like to receive future communications from McNeil-PPC, Inc., and Johnson & Johnson • Merck Consumer Pharmaceuticals Co.  
 @ Johnson & Johnson • Merck 2003  
 Douglas, AZ 85655-6607  
 P.O. Box 6607  
**SEND TO: Pepcid AC / Pepcid Complete \$3 Offer 780**  
 Form may not be mechanically reproduced. LIMIT TWO OFFERS PER FAMILY AND/OR ADDRESS, PER YEAR. No group or organization requests will be honored. Your offer may not be transferred or assigned. Offer void where prohibited or taxed. You should expect to receive your coupon in 8 weeks. Cannot be combined with any other offers.

**SAVE \$3.00 on Pepcid AC any strength or Pepcid Complete**  
 Here's how you can get \$3.00 off your next purchase: BUY: Pepcid AC (any strength) or Pepcid Complete Package 25 count or larger. SEND: This completed certificate, UPC from the Pepcid AC or Pepcid Complete package, plus dated cash register receipt with purchase price circled. RECEIVE: \$3.00 Coupon by mail.

**VALUABLE MAIL-IN OFFER**  
 and a guide of how to help your heartburn.  
 Distributed by: **Johansen-Johnson • MERCK**  
 www.pepcid.com  
 Registered trademark of Merck & Co., Inc.  
 CONSUMER PHARMACEUTICALS CO.  
 FORT WASHINGTON, PA 19034 USA


### Questions or comments?

Heartburn and acid indigestion are common, but you should stop taking MAXIMUM STRENGTH PEPCID® AC and see your doctor promptly if:  
 • You have trouble swallowing or stomach pain continues. You may have a serious condition that may need different treatment.  
 • You need to take this product for more than 14 days.  
**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers. Do not use with other acid reducers. Do not use if you have kidney disease, except under the advice and supervision of a doctor. This product should not be given to children under 12 years old, unless directed by a doctor. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

### Know when to see your doctor

The ingredient in MAXIMUM STRENGTH PEPCID® AC, famotidine, has been prescribed by doctors for years to treat millions of patients safely and effectively. The active ingredient in MAXIMUM STRENGTH PEPCID® AC has been taken safely with many frequently prescribed medications.

### Excellent safety record



Do not use if printed foil seal under bottle cap is open or torn.



WHAT YOU SHOULD KNOW ABOUT:



**A non-prescription stomach medicine**

MAXIMUM STRENGTH PEPCID<sup>®</sup> AC contains a medicine, different from antacids, that doctors have prescribed for years to treat acid-related problems in millions of people.

- Product Benefits:**
- 1 tablet relieves heartburn due to acid indigestion.
  - MAXIMUM STRENGTH PEPCID<sup>®</sup> AC prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods or beverages.
  - It contains famotidine, a prescription-proven medicine.



**How to use MAXIMUM STRENGTH PEPCID<sup>®</sup> AC**

Use MAXIMUM STRENGTH PEPCID<sup>®</sup> AC to relieve or prevent these symptoms:

- Heartburn
- Acid indigestion
- Sour stomach

MAXIMUM STRENGTH PEPCID<sup>®</sup> AC can be used up to twice daily (up to 2 tablets in 24 hours).

- Do not take more than 1 tablet at a time.
- Do not take more than 2 tablets in 24 hours.



**TO RELIEVE SYMPTOMS**

Swallow 1 tablet with a glass of water (do not chew).

**TO PREVENT SYMPTOMS**


Swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn.

**Excess acid: a burning problem**

It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the wrong place (the esophagus), or too much acid, can cause burning pain and discomfort that interfere with everyday activities.

**Heartburn – Caused by acid in the esophagus**

A valve-like muscle called the lower esophageal sphincter (LES) is relaxed in an open position



**Tips for Managing Heartburn**

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

Age:  under 35  35-49  50-54  55-64  65 yrs+  
E-mail: \_\_\_\_\_  
City: \_\_\_\_\_  
Address: \_\_\_\_\_  
Apt. \_\_\_\_\_  
Zip: \_\_\_\_\_  
State: \_\_\_\_\_



Yes, I'd like to receive future communications from McNeil-PPC, Inc., Splenda, Inc. and Johnson & Johnson • Merck Consumer Pharmaceuticals Co.  
Please circle one: Dr. Mr. Mrs. Miss.  
Name (please print) \_\_\_\_\_  
P.O. Box 6607  
Douglas, AZ 85655-6607  
© Johnson & Johnson • Merck 2003  
00000000  
NOTE: Offer good only in U.S.A. and APFD address. This request form may not be mechanically reproduced. LIMIT TWO OFFERS PER FAMILY AND/OR ADDRESS. PER YEAR. No group or organization requests will be honored. Your offer rights may not be transferred or assigned. Offer valid where prohibited or taxed. You should expect to receive your coupon in 8-9 weeks. Cannot be combined with any other offers.

**Save \$3.00 on Pepcid<sup>®</sup> AC any strength or Pepcid<sup>®</sup> Complete**  
**VALUABLE MAIL-IN OFFER**  
RECEIVE: \$3.00 Coupon by mail.  
SEND TO: Pepcid AC / Pepcid Complete \$3 Offer 780  
SEND: This completed certificate, UPC from the Pepcid AC or Pepcid Complete Package 25 count or Larger, cash register receipt with purchase price circled.  
BUY: Pepcid AC (any strength) or Pepcid Complete Package 25 count or Larger.  
Here's how you can get \$3.00 off your next purchase: product with purchase of 25 count or larger

Registered trademark of Merck & Co., Inc.  
FORT WASHINGTON, PA 19034 USA  
00000000  
CONSUMER PHARMACEUTICALS CO  
Distributors: Johnson & Johnson • MERCK  
www.pepcid.com  
Please call 1-800-755-4008 for more information and a guide of how to help your heartburn.


**Questions or comments?**

seek professional assistance or contact a Poison Control Center immediately.  
As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. Keep this and all drugs out of the reach of children, in case of accidental overdose, other doctor. This product should not be given to children under 12 years old, unless directed by a doctor.  
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use with other acid reducers. Do not use if you have kidney disease, except under the advice and supervision of a doctor.  
• You need different treatment.  
• You have trouble swallowing or stomach pain continues. You may have a serious condition that may need different treatment.

**Know when to see your doctor**

PEPCID<sup>®</sup> AC and acid indigestion are common, but you should stop taking MAXIMUM STRENGTH PEPCID<sup>®</sup> AC and see your doctor promptly if:  
• You have trouble swallowing or stomach pain continues. You may have a serious condition that may need different treatment.

**Excellent safety record**



In clinical studies, MAXIMUM STRENGTH PEPCID<sup>®</sup> AC film-coated tablets were significantly better than placebo tablets (tablets without the medicine) in relieving and preventing heartburn.  
Percent of heartburn episodes completely relieved within 3 hours

**Proven effective in clinical studies**