

Multiple scattering and p_T -broadening at RHIC energies

Gábor Papp^{a,b} Gergely G. Barnaföldi^c George Fai^b
Péter Lévai^c Yi Zhang^b

^a*HAS Research Group for Theoretical Physics, Eötvös University, Budapest,
Pázmány P. s. 1/A, Budapest, 1117, Hungary*

^b*Center for Nuclear Research, Department of Physics, Kent State University,
Kent, Ohio 44242, USA*

^c*Research Institute for Particle and Nuclear Physics, P.O.B. 49., 1525 Budapest
114., Hungary*

Presented by: Gábor Papp

Abstract

In ultrarelativistic heavy-ion collisions, in the $2 \text{ GeV} < p_T < 6 \text{ GeV}$ transverse-momentum region, the soft and semi-hard multiple scattering of the incoming nucleons results in the broadening of the expected hadronic (e.g. pion) p_T spectra. Thus, higher transverse-momentum regions are populated than in proton-proton (pp) collisions. In a perturbative QCD based calculation we include the intrinsic transverse momentum (k_T) of the partons in the nucleon (determined from pp collisions), augmented by the extra broadening obtained via a systematic analysis of pA collisions in the energy range $17 < s^{1/2} < 39 \text{ A GeV}$. The original polynomial spectra are modified, and an exponential spectrum appears in the region $2 \text{ GeV} < p_T < 4 \text{ GeV}$. At present RHIC energies ($\sqrt{s} = 56, 130 \text{ A GeV}$), the slope of the spectra coincides with that of fluid-dynamical descriptions, but it lacks any thermal origin. We determine and discuss the size of the modifications originating in multiple scattering, which lead to this state of affairs.
