



United States  
Department of  
Agriculture




NRCS

Natural  
Resources  
Conservation  
Service

In cooperation with  
the Oklahoma Agricultural  
Experiment Station  
and the  
Oklahoma Conservation  
Committee

# Supplement to the Soil Survey of Cherokee County, Oklahoma





# How To Use This Soil Survey Supplement

---

This document, in conjunction with the Web Soil Survey, supplements the Soil Survey of Cherokee County, Oklahoma, published in 1970. Find a map of your area of interest on the Web Soil Survey at <http://websoilsurvey.nrcs.usda.gov>. Note the map unit symbols in the area. Turn to the **Contents** in this supplement. The **Contents** lists the map units by symbol and name and shows the page where each map unit is described. Also see the **Contents** for sections of this publication that may address your specific needs.

Advancements in technology and increases in the intensity and variety of land uses have produced a need for updated soils information. In preparation for this publication, the correlation for the Soil Survey of Cherokee County was amended in April 1997 and February 2001. This publication and the Web Soil Survey include the recorrelated map unit legend and updated information regarding major soil properties and the use and management of the soils. In some cases, the name of the map unit and the name of the soil series have changed from the first publication. The map unit symbols and map delineations have not changed.

## Web Soil Survey

The latest detailed soil maps and updated tabular data, including soil properties and interpretations, are available on the Web Soil Survey at <http://websoilsurvey.nrcs.usda.gov>.

## Archived Soil Survey

Comprehensive descriptions of the detailed soil map units and additional information about the soils in the survey area are archived in the original Soil Survey of Cherokee County, Oklahoma. Archived soil surveys are available from many libraries, from the local office of the Natural Resources Conservation Service, and from the Cherokee County Conservation District in Tahlequah, Oklahoma.

---

This document is a publication of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (formerly the Soil Conservation Service) has leadership for the Federal part of the National Cooperative Soil Survey.

Major fieldwork for the Soil Survey of Cherokee County, Oklahoma, was completed in 1966. Soil names and descriptions were approved in 1967 and were amended in April 1997 and February 2001. Unless otherwise indicated, statements in this publication refer to conditions in the survey area in 1997. The maps for the survey were recompiled utilizing photographic imagery at 1:24,000 and rectified to 1995 digital orthophotography for SSURGO digitizing. The survey was made cooperatively by the Natural Resources Conservation Service, the Oklahoma Agricultural Experiment Station, and the Oklahoma Conservation Commission. It is part of the technical assistance furnished to the Cherokee County Conservation District.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

# Contents

---

<b>How To Use This Soil Survey Supplement</b> .....	i
<b>Contents</b> .....	iii
<b>Foreword</b> .....	v
How Soil Surveys Are Made .....	2
General Nature of the Survey Area .....	3
Relief and Drainage .....	3
Settlement and Development .....	3
Natural Resources .....	4
Transportation and Industry .....	4
Agriculture .....	5
Climate .....	5
<i>Table 1.--Temperature and Precipitation</i> .....	6
<i>Table 2.--Freeze Dates in Spring and Fall</i> .....	7
<i>Table 3.--Growing Season</i> .....	7
<b>Detailed Soil Map Units</b> .....	9
<i>Table 4.--Acreage and Proportionate Extent of the Soils</i> .....	11
CaB—Captina silt loam, 1 to 3 percent slopes .....	12
CkD—Clarksville very gravelly silt loam, 1 to 8 percent slopes .....	12
CIE—Clarksville stony silt loam, 5 to 20 percent slopes .....	13
CIF—Clarksville stony silt loam, 20 to 50 percent slopes .....	13
CoC—Coweta fine sandy loam, 3 to 5 percent slopes .....	14
DAM—Large dam .....	15
DnB—Dennis silt loam, 1 to 3 percent slopes .....	15
EdC—Eldorado silt loam, 3 to 5 percent slopes .....	16
EID—Eldorado stony silt loam, 3 to 12 percent slopes .....	16
Es—Elsah very gravelly loam, 0 to 3 percent slopes, frequently flooded .....	17
HcC—Hector fine sandy loam, 3 to 5 percent slopes .....	18
HIE—Hector-Linker association, 8 to 30 percent slopes .....	18
JaA—Jay silt loam, 1 to 3 percent slopes .....	19
LkC—Linker fine sandy loam, 3 to 5 percent slopes .....	20
LoB—Tonti gravelly silt loam, 1 to 3 percent slopes .....	20
M-W—Miscellaneous water .....	21
NaB—Newtonia silt loam, 1 to 3 percent slopes .....	21
NaC—Newtonia silt loam, 3 to 5 percent slopes .....	22
NaC2—Newtonia silt loam, 3 to 5 percent slopes, eroded .....	23
NixB—Nixa gravelly silt loam, 0 to 3 percent slopes .....	23
OkA—Okemah silty clay loam, 0 to 1 percent slopes .....	24
OkB—Okemah silty clay loam, 1 to 3 percent slopes .....	25
OkC—Okemah silty clay loam, 3 to 5 percent slopes .....	25
Os—Osage clay, 0 to 1 percent slopes, occasionally flooded .....	26
PIT—Pits .....	27
Rs—Rock outcrop-Hector complex, 40 to 100 percent slopes .....	27
SaA—Britwater silt loam, 0 to 1 percent slopes .....	28
SaB—Britwater silt loam, 1 to 3 percent slopes .....	28

---

SgB—Britwater gravelly silt loam, 1 to 3 percent slopes .....	29
SgD—Britwater gravelly silt loam, 3 to 8 percent slopes .....	30
Sm—Healing silt loam, 0 to 1 percent slopes, occasionally flooded .....	30
Sn—Razort gravelly loam, 0 to 3 percent slopes, occasionally flooded .....	31
SrA—Stigler silt loam, 0 to 1 percent slopes .....	32
SuC2—Apperson silty clay loam, 3 to 5 percent slopes, eroded .....	32
TkA—Taloka silt loam, 0 to 1 percent slopes .....	33
TrD—Shidler-Rock outcrop complex, 2 to 8 percent slopes .....	34
TrF—Shidler-Rock outcrop complex, 15 to 50 percent slopes .....	34
Vd—Verdigris silt loam, 0 to 1 percent slopes, occasionally flooded .....	35
VrdA—Verdigris silt loam, 0 to 1 percent slopes, frequently flooded .....	36
W—Water .....	36
<b>Use and Management of the Soils</b> .....	39
Cropland .....	39
Rangeland .....	40
Similarity Index .....	42
Rangeland Management .....	43
Ecological Site Descriptions .....	44
<b>Formation and Classification of the Soils</b> .....	51
Formation of the Soils .....	51
Classification of the Soils .....	51
<i>Table 5.--Classification of the Soils</i> .....	53
<b>References</b> .....	55
<b>Glossary</b> .....	57

Issued December 2006

# Foreword

---

This soil survey supplement contains information that can be used in conjunction with the previously published soil survey and with online resources. It provides valuable information for land-planning programs in Oklahoma. It contains predictions of soil behavior for selected land uses. This supplement also highlights limitations and hazards inherent in the soil, improvements needed to overcome the limitations, and the impact of selected land uses on the environment.

Soil surveys are designed for many different users. Farmers, ranchers, foresters, and agronomists can use a survey to evaluate the potential of the soil and the management needed for maximum food and fiber production. Planners, community officials, engineers, developers, builders, and home buyers can use a survey to plan land use, select sites for construction, and identify special practices needed to ensure proper performance. Conservationists, teachers, students, and specialists in recreation, wildlife management, waste disposal, and pollution control can use a survey to help them understand, protect, and enhance the environment.

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are shallow to bedrock. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are not suited for use as septic tank absorption fields. A high water table makes a soil very limited for basements or underground installations.

These and many other soil properties that affect land use are described in soil surveys. Each soil in the survey area is described. Information on specific uses is given for each soil. Help in using this publication and additional information are available at the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.



Ronald L. Hilliard  
State Conservationist  
Natural Resources Conservation Service





# Supplement to the Soil Survey of Cherokee County, Oklahoma

---

By Everett L. Cole, Natural Resources Conservation Service  
(formerly the Soil Conservation Service)

Fieldwork by Peter Warth and Dock J. Polone, Natural Resources  
Conservation Service (formerly the Soil Conservation Service)

United States Department of Agriculture,  
Natural Resources Conservation Service,  
in cooperation with  
the Oklahoma Agricultural Experiment Station and  
the Oklahoma Conservation Commission

This supplement provides information to update the original Soil Survey of Cherokee County, Oklahoma, which was issued December 1970. The original tables and maps are not included in this document. Updated tables and new digital maps on updated photography are available from the Web Soil Survey at <http://websoilsurvey.nrcs.usda.gov>. The tables were generated from the NRCS National Soil Information System (NASIS) and are also available from the NRCS Soil Data Mart at <http://soildatamart.nrcs.usda.gov>.

CHEROKEE COUNTY is in the northeastern part of Oklahoma (fig. 1). It is bordered by Delaware and Mayes Counties on the north, by Wagoner and Muskogee Counties on the west, by Sequoyah County on the south, and by Adair County on the east.

Cherokee County has an area of about 497,235 acres, or 777 square miles. Tahlequah, the county seat, is in the central part of the county. The main source of income in the county is the sale of livestock and livestock products. Small grains, corn, grain sorghum, and improved grasses and legumes for hay are grown as feed




Figure 1.—Location of Cherokee County in Oklahoma.

for livestock. Pasture grasses and native grasses in woodland are grazed by livestock. The sale of timber supplements farm income. Woodland occupies about 61 percent of Cherokee County.

## **How Soil Surveys Are Made**

Soil surveys are made to provide information about the soils and miscellaneous areas in a survey area. The information includes a description of the soils and miscellaneous areas and their location and a discussion of their suitability, limitations, and management for specified uses. Soil scientists observe the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They dig many holes to study the soil profile, which is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

The soils and miscellaneous areas in a survey area are in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind or segment of the landscape. By observing the soils and miscellaneous areas in a survey area and relating their position to specific segments of the landscape, soil scientists develop a concept, or model, of how the soils were formed. Thus, during mapping, this model enables the soil scientists to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Individual soils on the landscape commonly merge into one another as their characteristics gradually change. To construct an accurate map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists record the characteristics of the soil profiles that they study. They note color, texture, size, and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in a survey area and determining their properties, the soil scientists assign the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil scientists classify and name the soils in a survey area, they compare the individual soils with similar soils in the same taxonomic class in other areas so that they can confirm data and assemble additional data based on experience and research.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop

yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists locate and identify the significant natural bodies of soil in a survey area, they draw the boundaries of these bodies on aerial photographs and identify each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

The descriptions, names, and delineations of the soils in a survey area do not always fully agree with those of the soils in adjacent survey areas. Differences are the result of a better knowledge of soils, modifications in series concepts, or variations in the intensity of mapping or in the extent of the soils in the survey areas.

## **General Nature of the Survey Area**

This section gives general information about this survey area. It describes relief and drainage, settlement and development, natural resources, transportation and industry, agriculture, and climate.

### **Relief and Drainage**

Cherokee County is mostly hilly, but there are small, more gently sloping areas. The general slope is from the northeast to the southwest. The Neosho (Grand) and Illinois Rivers drain most of the county. Fourteen Mile, Clear, Double Spring, and Pecan Creeks are the main streams draining into the Neosho (Grand) River. Elk, Barren Fork, and Caney Creeks drain into the Illinois River. These rivers and creeks have entrenched about 200 to 400 feet and have imparted a hilly relief to most of the survey area.

Landforms in the county include prairies and areas of cherty limestone and sandstone. The prairies are mostly nearly level to sloping. The areas of cherty limestone and sandstone are hilly. The divides between drainageways are nearly level to sloping and range from  $\frac{1}{4}$  to 4 miles in width, but the slopes that extend into the drains are strongly sloping to very steep. The larger areas that are drained consist of nearly level to very gently sloping flood plains and nearly level to sloping benches. The drainageways range from 200 feet wide along the smallest streams to about 2 miles wide along some of the rivers.

### **Settlement and Development**

The early settlement of Cherokee County was by the Cherokee, Delaware, and Seneca Indians. When Oklahoma became a state, the county was named for the Cherokee Indians. Land was allotted to the Indians on basis of its cash value, the largest allotments in size being in the hilly, timbered areas. Land could be leased for farming subject to the supervision of the Indian Agency. The sale or lease of land brought non-Indian settlers into the area.

Most of the early settlers farmed. In the hilly, timbered areas, the land had to be cleared before it could be cultivated. Some settlers quit farming and sold out because they found that the areas they had bought were not suitable for cultivation. Other farmers acquired these areas, and farmed units increased in size.

Two large reservoirs were built on streams in the survey area for municipal and industrial purposes. Some industry was established.

Tahlequah, the old Cherokee Capital, is the county seat of Cherokee County. Hulbert, which is smaller, is the only other sizable town in the county.

## **Natural Resources**

The natural resources of the survey area are mainly water, timber, gravel, limestone, wild game, fish, and scenic beauty.

The water supply for towns comes mainly from wells, perennial streams, and reservoirs. Hydroelectric and flood-control reservoirs furnish cheap electricity and irrigation water. Wells, springs, and farm ponds supply water for livestock. The water in the lakes and streams is crystal clear and of high quality.

The income from timber is less than it was in past years. Most of the timber has been cut over, and the trees that were left to propagate the stands are of poor quality. This timber is now used mainly for the production of tool handles, gunstocks, and special furniture parts.

In the survey area, deposits of gravel are abundant along streams, but the gravel is little used except as road material and in cement mixtures. The gravel is used for roads mainly on gently sloping soils because most of the soils in hilly areas contain chert, sandstone, or shale material.

Limestone is the most common mineral in the area, and some of it is mined for commercial and agricultural purposes. Sample specimens of lead and zinc also occur.

Clear running streams and the lakes in the hilly, timbered areas provide scenic beauty that attracts thousands of visitors each year. Visitors are most numerous when the redbud and dogwood trees bloom in the spring and when the hardwood trees turn from green to brilliant colors in the fall.

Wildlife and fish are abundant in the survey area. Big game has been restocked in the Cookson Hills and Cherokee Wildlife Management Areas. These management areas are open to deer hunting for short periods. Quail, dove, and duck are hunted during season in all areas. These and other kinds of game are stocked by commercial and private game farms. Fishing is a year-round sport on the streams and lakes.

## **Transportation and Industry**

Although there are no railroads within the survey area, bus service and freight trucking are available in Tahlequah and other communities. Federal and State highways form a network of paved roads in the county. State Routes 10 and 82 extend north and south across the survey area. State Route 51 and U.S. Highway 62 are the main east and west roads. State Routes 80 and 100 provide access to other parts of the county.

In farm areas, roads of graded chert, sandstone, gravel, shale, and dirt provide access to hard-surfaced roads in nearly all weather. Some areas in the hilliest part, however, have few roads.

Grain elevators are located in adjoining counties, and most dairy farmers market milk in nearby large cities.

Livestock in the survey area is marketed at auctions near Tahlequah. Poultry and poultry products are sold to egg-processing and broiler-processing markets outside of the survey area.

Small sawmills are scattered throughout the survey area. Timber products are produced and used locally. Vegetables and craft products are also important to the local economy.

## Agriculture

In Cherokee County, the first settlers, who were Indians, found forests of pine and oak and numerous grassy savannahs and open prairie flats. Peaches and tobacco were among the first crops grown on the relatively few acres of cultivated land, but wheat and corn soon became the principal crops. By the mid-1930s, strawberries, tomatoes, blackberries, and green beans were important crops. Since early in the 1940s, the acreage of cultivated crops in the county has declined steadily, and the acreage used for pasture and livestock enterprises has increased.

For many years the trend has been for the number of farms to decrease and the average size of farms to increase. The number of farms in the county decreased from 1,798 in 1954 to 1,221 in 2002. Over the same period, the average size of a farm increased from 129 to 181 acres.

Livestock and poultry production are now the principle farm products in the survey area.

## Climate

The Natural Resources Conservation Service National Water and Climate Center, Portland, Oregon, helped prepare this section.

Cherokee County has a mild, temperate climate. It occupies a portion of the Ozark Highland in the northeastern part of Oklahoma (USDA, 1970). This is an area of wooded hills, small prairies, pasturelands, many streams, and portions of two large lakes. Elevations range from 600 to 1,500 feet above sea level. Rainfall in the area is ample for farming and the encouragement of natural beauty and tourism. The major changes in the weather and the heaviest rains occur when warm, moisture-laden air from the Gulf of Mexico meets cool, drier air from the Pacific Ocean and Arctic areas.

Because of the continental effects, there are pronounced daily and seasonal changes in temperature and variations in seasonal and annual rainfall. Seasonal changes are gradual, but each season is distinct. Winters are mild and short. They have brief cold periods and occasional snows. Severe local storms and the heaviest rains are most numerous in spring and early summer. Summers are long and hot. The heat is moderated by occasional showers or thunderstorms and moderate winds. A secondary period of heavy precipitation occurs early in fall and is followed by many pleasant, sunny days and cool nights.

The climate tables were created using data from the climate station at Tahlequah, Oklahoma. Thunderstorm days, relative humidity, percent sunshine, and wind information were estimated using data from the first order station at Fort Smith, Arkansas.

The table "Temperature and Precipitation" gives data for the survey area as recorded at Tahlequah in the period 1971 to 2000. The table "Freeze Dates in Spring and Fall" shows probable dates of the first freeze in fall and the last freeze in spring. The table "Growing Season" provides data on the length of the growing season.

In winter, the average temperature is 40.0 degrees F and the average daily minimum temperature is 28.9 degrees. The lowest temperature on record, which occurred at Tahlequah on December 23, 1989, is -14 degrees. In summer, the average temperature is 78.6 degrees and the average daily maximum temperature is 90.2 degrees. The highest temperature, which occurred at Tahlequah on July 13, 1954, is 114 degrees.

Growing degree days are shown in the table "Temperature and Precipitation." They are equivalent to "heat units." During the month, growing degree days accumulate by the amount that the average temperature each day exceeds a base temperature (50 degrees F). The normal monthly accumulation is used to schedule single or

Supplement to the Soil Survey of Cherokee County, Oklahoma

successive plantings of a crop between the last freeze in spring and the first freeze in fall.

The average annual total precipitation is about 47.78 inches. Of this, about 31.07 inches, or 65 percent, usually falls in April through October. The growing season for most crops falls within this period. The heaviest 1-day rainfall during the period of record was 6.6 inches at Tahlequah on October 9, 1970. Thunderstorms occur on about 57 days each year. Most occur in May, which averages 8.5.

The average seasonal snowfall is 6.3 inches. The greatest snow depth at any one time during the period of record was 14 inches recorded on March 6, 1989. On an average, 4 days per year have at least 1 inch of snow on the ground. The heaviest 1-day snowfall on record was 14 inches recorded on January 9, 1977.

The average relative humidity in mid-afternoon is about 55 percent. Humidity is higher at night, and the average at dawn is about 85 percent. The sun shines 72 percent of the time possible in summer and 52 percent in winter. The prevailing wind is from the northeast. Average wind speed is highest, 9.4 miles per hour, in March.

Table 1.--Temperature and Precipitation

[Recorded in the period 1971-2000 at Tahlequah, Oklahoma]

Month	Temperature						Precipitation				
	Average daily maximum	Average daily minimum	Average	2 years in 10 will have--		Average number of growing degree days*	Avg.	2 years in 10 will have--		Average number of days with 0.10 inch	Average snowfall
				Max. temp. higher than	Min. temp. lower than			Less than--	More than--		
°F	°F	°F	°F	°F	Units	In	In	In		In	
January----	48.3	26.5	37.4	72	-2	17	2.41	0.82	3.89	4	2.6
February---	54.1	30.6	42.4	78	1	44	2.35	1.03	3.66	4	2.0
March-----	63.8	39.6	51.7	85	14	155	4.15	2.22	5.95	6	0.3
April-----	72.6	47.8	60.2	88	25	318	40.2	1.95	6.14	6	0.0
May-----	79.2	56.9	68.1	91	36	552	5.61	3.50	7.54	7	0.0
June-----	86.3	64.5	75.4	97	47	755	5.15	2.22	7.71	6	0.0
July-----	92.0	68.7	80.3	104	53	934	3.55	1.16	5.87	4	0.0
August-----	92.3	67.7	80.0	103	53	917	3.19	1.30	4.99	4	0.0
September--	84.3	60.8	72.5	100	36	668	5.27	2.53	7.97	6	0.0
October----	73.9	50.1	62.0	90	27	383	4.28	1.62	6.53	5	0.0
November---	60.7	39.3	50.0	81	16	119	4.51	2.37	6.61	5	0.6
December---	50.7	29.7	40.2	75	3	29	3.30	1.22	5.17	4	0.9
Yearly:											
Average--	71.5	48.5	60.0	---	---	---	---	---	---	---	---
Extreme--	110	-14	---	105	-6	---	---	---	---	---	---
Total----	---	---	---	---	---	4,893	47.78	36.48	62.69	61	6.3

\* A growing degree day is a unit of heat available for plant growth. It can be calculated by adding the maximum and minimum daily temperatures, dividing the sum by 2, and subtracting the temperature below which growth is minimal for the principal crops in the area (50 degrees F).

Supplement to the Soil Survey of Cherokee County, Oklahoma

Table 2.--Freeze Dates in Spring and Fall

[Recorded in the period 1971-2000 at Tahlequah, Oklahoma]

Probability	Temperature		
	24 °F or lower	28 °F or lower	32 °F or lower
Last freezing temperature in spring:			
1 year in 10 later than--	Apr. 6	Apr. 14	Apr. 20
2 years in 10 later than--	Mar. 29	Apr. 9	Apr. 16
5 years in 10 later than--	Mar. 15	Mar. 29	Apr. 8
First freezing temperature in fall:			
1 year in 10 earlier than--	Oct. 31	Oct. 16	Oct. 7
2 years in 10 earlier than--	Nov. 6	Oct. 22	Oct. 13
5 years in 10 earlier than--	Nov. 17	Nov. 3	Oct. 25

Table 3.--Growing Season

[Recorded in the period 1971-2000 at Tahlequah, Oklahoma]

Probability	Daily Minimum Temperature During growing season		
	Higher than 24 °F	Higher than 28 °F	Higher than 32 °F
	Days	Days	Days
9 years in 10	216	198	179
8 years in 10	226	205	186
5 years in 10	246	219	199
2 years in 10	265	232	212
1 year in 10	276	240	219





# Detailed Soil Map Units

---

Detailed soil maps for the survey area are available online at <http://websoilsurvey.nrcs.usda.gov>. The map units on the detailed soil maps represent the soils and miscellaneous areas in the survey area. The map unit descriptions in this section, along with the maps, can be used to determine the suitability and potential of a unit for specific uses. They also can be used to plan the management needed for those uses.

A map unit delineation on the detailed soil maps represents an area on the landscape and consists of one or more soils or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils or miscellaneous areas. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils and miscellaneous areas are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, are mapped without areas of minor components of other taxonomic classes. Consequently, map units are made up of the soils or miscellaneous areas for which they are named and some areas of included soils that belong to other taxonomic classes.

Most included soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting or similar inclusions. They may or may not be mentioned in the map unit description. Other included soils and miscellaneous areas, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, inclusions. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. The included areas of contrasting soils or miscellaneous areas are mentioned in the map unit descriptions. A few included areas may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of included areas in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into segments that have similar use and management requirements. The delineation of such landscape segments on the map provides sufficient information for the development of resource plans, but if intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit.

Soils that have profiles that are almost alike make up a soil series. Except for differences in texture of the surface layer or of the underlying layers, all the soils of

## Supplement to the Soil Survey of Cherokee County, Oklahoma

a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer or of the underlying layers. They also can differ in slope, stoniness, salinity, wetness, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into soil phases. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Verdigris silt loam, 0 to 1 percent slopes, occasionally flooded, is a phase of the Verdigris series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Shidler-Rock outcrop complex, 2 to 8 percent slopes, is an example.

This survey includes *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

The table "Acreage and Proportionate Extent of the Soils" gives the acreage and proportionate extent of each map unit. The Glossary defines many of the terms used in describing the soils or miscellaneous areas.

Supplement to the Soil Survey of Cherokee County, Oklahoma

Table 4.--Acreage and Proportionate Extent of the Soils

Map symbol	Soil name	Acres	Percent
CaB	Captina silt loam, 1 to 3 percent slopes-----	8,502	1.7
CkD	Clarksville very gravelly silt loam, 1 to 8 percent slopes-----	52,323	10.5
ClE	Clarksville stony silt loam, 5 to 20 percent slopes-----	24,962	5.0
ClF	Clarksville stony silt loam, 20 to 50 percent slopes-----	96,997	19.5
CoC	Coweta fine sandy loam, 3 to 5 percent slopes-----	1,680	0.3
DAM	Large dam-----	10	*
DnB	Dennis silt loam, 1 to 3 percent slopes-----	945	0.2
EdC	Eldorado silt loam, 3 to 5 percent slopes-----	4,059	0.8
ELD	Eldorado stony silt loam, 3 to 12 percent slopes-----	1,662	0.3
Es	Elsah very gravelly loam, 0 to 3 percent slopes, frequently flooded-----	19,953	4.0
HcC	Hector fine sandy loam, 3 to 5 percent slopes-----	21,858	4.4
HlE	Hector-Linker association, 8 to 30 percent slopes-----	89,570	18.0
JaA	Jay silt loam, 1 to 3 percent slopes-----	5,588	1.1
LkC	Linker fine sandy loam, 3 to 5 percent slopes-----	6,045	1.2
LoB	Tonti gravelly silt loam, 1 to 3 percent slopes-----	22,199	4.5
M-W	Miscellaneous water-----	104	*
NaB	Newtonia silt loam, 1 to 3 percent slopes-----	8,309	1.7
NaC	Newtonia silt loam, 3 to 5 percent slopes-----	2,310	0.5
NaC2	Newtonia silt loam, 3 to 5 percent slopes, eroded-----	601	0.1
NixB	Nixa gravelly silt loam, 0 to 3 percent slopes-----	124	*
OkA	Okemah silty clay loam, 0 to 1 percent slopes-----	3,351	0.7
OkB	Okemah silty clay loam, 1 to 3 percent slopes-----	7,606	1.5
OkC	Okemah silty clay loam, 3 to 5 percent slopes-----	2,265	0.5
Os	Osage clay, 0 to 1 percent slopes, occasionally flooded-----	1,875	0.4
PIT	Pits-----	281	*
Rs	Rock outcrop-Hector complex, 40 to 100 percent slopes-----	11,021	2.2
SaA	Britwater silt loam, 0 to 1 percent slopes-----	1,027	0.2
SaB	Britwater silt loam, 1 to 3 percent slopes-----	6,005	1.2
SgB	Britwater gravelly silt loam, 1 to 3 percent slopes-----	10,326	2.1
SgD	Britwater gravelly silt loam, 3 to 8 percent slopes-----	19,203	3.9
Sm	Healing silt loam, 0 to 1 percent slopes, occasionally flooded-----	4,191	0.8
Sn	Razort gravelly loam, 0 to 3 percent slopes, occasionally flooded-----	4,811	1.0
SrA	Stigler silt loam, 0 to 1 percent slopes-----	3,920	0.8
SuC2	Apperson silty clay loam, 3 to 5 percent slopes, eroded-----	2,485	0.5
TkA	Taloka silt loam, 0 to 1 percent slopes-----	1,355	0.3
TrD	Shidler-Rock outcrop complex, 2 to 8 percent slopes-----	9,182	1.8
TrF	Shidler-Rock outcrop complex, 15 to 50 percent slopes-----	21,543	4.3
Vd	Verdigris silt loam, 0 to 1 percent slopes, occasionally flooded-----	787	0.2
VrdA	Verdigris silt loam, 0 to 1 percent slopes, frequently flooded-----	126	*
W	Water-----	18,074	3.6
	Total-----	497,235	100.0

\* Less than 0.1 percent.

## **CaB—Captina silt loam, 1 to 3 percent slopes**

### **Map Unit Setting**

*Major land resource area:* 116A  
*Elevation:* 700 to 1,400 feet  
*Mean annual precipitation:* 42 to 46 inches  
*Mean annual air temperature:* 54 to 57 degrees F  
*Frost-free period:* 160 to 195 days

### **Major Component Description**

#### **Captina**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Paleoterraces on uplands  
*Position on the landform:* Treads  
*Parent material:* Loess over colluvium and/or residuum weathered from cherty limestone  
*Slope:* 1 to 3 percent  
*Runoff:* Very high  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Slow  
*Slowest permeability class within a depth of 80 inches:* Slow  
*Drainage class:* Moderately well drained  
*Available water capacity:* About 7.3 inches  
*Depth to the top of the seasonal high water table:* 1.0 to 2.0 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 3e  
*Ecological site number and name:* R116AY092OK, Smooth Chert Savannah PE 72–80  
*Typical profile:*  
A—0 to 7 inches; silt loam  
BA—7 to 23 inches; silty clay loam  
Bt—23 to 38 inches; silty clay loam  
Bx—38 to 62 inches; gravelly silty clay loam

## **CkD—Clarksville very gravelly silt loam, 1 to 8 percent slopes**

### **Map Unit Setting**

*Major land resource area:* 116A  
*Elevation:* 700 to 1,300 feet  
*Mean annual precipitation:* 40 to 46 inches  
*Mean annual air temperature:* 54 to 57 degrees F  
*Frost-free period:* 160 to 200 days

### **Major Component Description**

#### **Clarksville**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Position on hillslope:* Backslopes  
*Parent material:* Loamy colluvium over residuum weathered from cherty limestone  
*Slope:* 1 to 8 percent  
*Runoff:* Low  
*Soil depth:* Greater than 60 inches

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer: Moderate*

*Slowest permeability class within a depth of 80 inches: Moderate*

*Drainage class: Somewhat excessively drained*

*Available water capacity: About 4.7 inches*

*Depth to the top of the seasonal high water table: Greater than 6 feet*

*Flooding: None*

*Ponding: None*

*Land capability classification, nonirrigated: 4e*

*Typical profile:*

A—0 to 10 inches; very gravelly silt loam

Bt—10 to 40 inches; very gravelly silty clay loam

BC—40 to 60 inches; very gravelly silty clay

## ***CIE—Clarksville stony silt loam, 5 to 20 percent slopes***

### ***Map Unit Setting***

*Major land resource area: 116A*

*Elevation: 700 to 1,200 feet*

*Mean annual precipitation: 40 to 46 inches*

*Mean annual air temperature: 54 to 57 degrees F*

*Frost-free period: 160 to 200 days*

### ***Major Component Description***

#### ***Clarksville***

*Extent: 100 percent of the map unit*

*Geomorphic setting: Hillslopes on hills on uplands*

*Position on hillslope: Backslopes*

*Parent material: Loamy colluvium over residuum weathered from cherty limestone*

*Slope: 5 to 20 percent*

*Runoff: Low*

*Soil depth: Greater than 60 inches*

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer: Moderate*

*Slowest permeability class within a depth of 80 inches: Moderate*

*Drainage class: Somewhat excessively drained*

*Available water capacity: About 4.7 inches*

*Depth to the top of the seasonal high water table: Greater than 6 feet*

*Flooding: None*

*Ponding: None*

*Land capability classification, nonirrigated: 6e*

*Typical profile:*

A—0 to 10 inches; stony silt loam

Bt—10 to 40 inches; very gravelly silty clay loam

BC—40 to 60 inches; very gravelly silty clay

## ***CIF—Clarksville stony silt loam, 20 to 50 percent slopes***

### ***Map Unit Setting***

*Major land resource area: 116A*

*Elevation: 700 to 1,200 feet*

*Mean annual precipitation: 40 to 46 inches*

*Mean annual air temperature:* 54 to 57 degrees F

*Frost-free period:* 160 to 200 days

### **Major Component Description**

#### **Clarksville**

*Extent:* 100 percent of the map unit

*Geomorphic setting:* Hillslopes on hills on uplands

*Position on hillslope:* Backslopes

*Parent material:* Loamy colluvium over residuum weathered from cherty limestone

*Slope:* 20 to 50 percent

*Runoff:* Medium

*Soil depth:* Greater than 60 inches

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate

*Slowest permeability class within a depth of 80 inches:* Moderate

*Drainage class:* Somewhat excessively drained

*Available water capacity:* About 4.7 inches

*Depth to the top of the seasonal high water table:* Greater than 6 feet

*Flooding:* None

*Ponding:* None

*Land capability classification, nonirrigated:* 8e

*Typical profile:*

A—0 to 10 inches; stony silt loam

Bt—10 to 40 inches; very gravelly silty clay loam

BC—40 to 60 inches; very gravelly silty clay

## **CoC—Coweta fine sandy loam, 3 to 5 percent slopes**

### **Map Unit Setting**

*Major land resource area:* 112

*Elevation:* 700 to 1,100 feet

*Mean annual precipitation:* 37 to 45 inches

*Mean annual air temperature:* 57 to 62 degrees F

*Frost-free period:* 200 to 220 days

### **Major Component Description**

#### **Coweta**

*Extent:* 100 percent of the map unit

*Geomorphic setting:* Hillslopes on hills on uplands

*Position on hillslope:* Backslopes

*Parent material:* Loamy residuum weathered from sandstone and shale

*Slope:* 3 to 5 percent

*Runoff:* Medium

*Soil depth:* 10 to 20 inches to paralithic bedrock

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate

*Slowest permeability class within a depth of 80 inches:* Moderately slow

*Drainage class:* Well drained

*Available water capacity:* About 1.7 inches

*Depth to the top of the seasonal high water table:* Greater than 6 feet

*Flooding:* None

*Ponding:* None

*Land capability classification, nonirrigated:* 4s

*Ecological site number and name:* R112XY086OK, Shallow Prairie (Eastern) PE  
62–80

*Typical profile:*

- A—0 to 6 inches; fine sandy loam
- Bw—6 to 13 inches; gravelly fine sandy loam
- Cr—13 to 20 inches; bedrock

## **DAM—Large dam**

### **Map Unit Setting**

*Major land resource area:* 116A  
*Elevation:* 700 to 2,000 feet  
*Mean annual precipitation:* 30 to 35 inches  
*Mean annual air temperature:* 60 to 61 degrees F  
*Frost-free period:* 200 to 210 days

### **Major Component Description**

#### **Large dam**

*Extent:* 100 percent of the map unit  
*Definition:* This map unit consists of dams constructed to impound water for ponds and lakes.  
*Geomorphic setting:* Artificial levees  
*Parent material:* Mine spoil or earthy fill  
*Slope:* 3 to 45 percent  
*Runoff:* Very high  
*Land capability classification, nonirrigated:* 8

## **DnB—Dennis silt loam, 1 to 3 percent slopes**

### **Map Unit Setting**

*Major land resource area:* 112  
*Elevation:* 500 to 1,200 feet  
*Mean annual precipitation:* 37 to 46 inches  
*Mean annual air temperature:* 57 to 62 degrees F  
*Frost-free period:* 190 to 220 days

### **Major Component Description**

#### **Dennis**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Position on hillslope:* Shoulders  
*Parent material:* Clayey residuum weathered from shale  
*Slope:* 1 to 3 percent  
*Runoff:* Very high  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Slow  
*Slowest permeability class within a depth of 80 inches:* Slow  
*Drainage class:* Somewhat poorly drained  
*Available water capacity:* About 10.5 inches  
*Depth to the top of the seasonal high water table:* 1.0 to 2.5 feet  
*Flooding:* None  
*Ponding:* None

*Land capability classification, nonirrigated:* 2e

*Ecological site number and name:* R112XY059OK, Loamy Prairie (Northeast) PE  
62–80

*Typical profile:*

A—0 to 13 inches; silt loam

BA—13 to 17 inches; silty clay loam

Bt—17 to 44 inches; silty clay loam

BC—44 to 66 inches; silty clay loam

## ***EdC—Eldorado silt loam, 3 to 5 percent slopes***

### ***Map Unit Setting***

*Major land resource area:* 112

*Elevation:* 500 to 1,000 feet

*Mean annual precipitation:* 37 to 43 inches

*Mean annual air temperature:* 58 to 62 degrees F

*Frost-free period:* 200 to 220 days

### ***Major Component Description***

#### ***Eldorado***

*Extent:* 100 percent of the map unit

*Geomorphic setting:* Hillslopes on hills on uplands

*Position on hillslope:* Footslopes

*Parent material:* Gravelly, loamy residuum weathered from cherty limestone

*Slope:* 3 to 5 percent

*Runoff:* Low

*Soil depth:* Greater than 60 inches

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate

*Slowest permeability class within a depth of 80 inches:* Moderate

*Drainage class:* Well drained

*Available water capacity:* About 5.6 inches

*Depth to the top of the seasonal high water table:* Greater than 6 feet

*Flooding:* None

*Ponding:* None

*Land capability classification, nonirrigated:* 3e

*Ecological site number and name:* R112XY059OK, Loamy Prairie (Northeast) PE  
62–80

*Typical profile:*

A—0 to 10 inches; silt loam

BA—10 to 18 inches; very gravelly silt loam

Bt1—18 to 30 inches; very gravelly silty clay loam

Bt2—30 to 63 inches; very gravelly silty clay loam

## ***EID—Eldorado stony silt loam, 3 to 12 percent slopes***

### ***Map Unit Setting***

*Major land resource area:* 112

*Elevation:* 500 to 1,000 feet

*Mean annual precipitation:* 37 to 43 inches

*Mean annual air temperature:* 58 to 62 degrees F

*Frost-free period:* 200 to 220 days



### **Major Component Description**

#### **Eldorado**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Position on hillslope:* Backslopes  
*Parent material:* Gravelly, loamy residuum weathered from cherty limestone  
*Slope:* 3 to 12 percent  
*Runoff:* Medium  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate  
*Slowest permeability class within a depth of 80 inches:* Moderate  
*Drainage class:* Well drained  
*Available water capacity:* About 4.7 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 6e  
*Ecological site number and name:* R112XY059OK, Loamy Prairie (Northeast) PE 62–80  
*Typical profile:*  
A—0 to 11 inches; stony silt loam  
BA—11 to 22 inches; very gravelly silt loam  
Bt—22 to 63 inches; extremely gravelly silty clay loam

### **Es—Elsah very gravelly loam, 0 to 3 percent slopes, frequently flooded**

#### **Map Unit Setting**

*Major land resource area:* 116A  
*Elevation:* 340 to 1,500 feet  
*Mean annual precipitation:* 36 to 48 inches  
*Mean annual air temperature:* 54 to 57 degrees F  
*Frost-free period:* 180 to 200 days

### **Major Component Description**

#### **Elsah**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Flood plains in drainageways in valleys  
*Parent material:* Gravelly, loamy alluvium  
*Slope:* 0 to 3 percent  
*Runoff:* Low  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate  
*Slowest permeability class within a depth of 80 inches:* Moderate  
*Drainage class:* Somewhat excessively drained  
*Available water capacity:* About 5.5 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* Frequent  
*Ponding:* None  
*Land capability classification, nonirrigated:* 5w  
*Typical profile:*  
A—0 to 15 inches; very gravelly loam  
Bw1—15 to 22 inches; very gravelly loam  
Bw2—22 to 62 inches; very gravelly loam

## **HcC—Hector fine sandy loam, 3 to 5 percent slopes**

### **Map Unit Setting**

*Major land resource area:* 117  
*Elevation:* 500 to 2,400 feet  
*Mean annual precipitation:* 46 to 50 inches  
*Mean annual air temperature:* 61 to 64 degrees F  
*Frost-free period:* 175 to 210 days

### **Major Component Description**

#### **Hector**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Mountain slopes on mountains  
*Position on the landform:* Mountainflanks  
*Parent material:* Loamy residuum weathered from sandstone  
*Slope:* 3 to 5 percent  
*Runoff:* Very high  
*Soil depth:* 10 to 20 inches to lithic bedrock  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderately rapid  
*Slowest permeability class within a depth of 80 inches:* Impermeable  
*Drainage class:* Well drained  
*Available water capacity:* About 1.8 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 4s  
*Ecological site number and name:* R117XY088OK, Shallow Savannah PE 72–76  
*Typical profile:*  
    A—0 to 7 inches; fine sandy loam  
    Bw—7 to 15 inches; gravelly fine sandy loam  
    R—15 to 30 inches; bedrock

## **HIE—Hector-Linker association, 8 to 30 percent slopes**

### **Map Unit Setting**

*Major land resource area:* 117  
*Elevation:* 500 to 2,800 feet  
*Mean annual precipitation:* 43 to 50 inches  
*Mean annual air temperature:* 57 to 64 degrees F  
*Frost-free period:* 175 to 210 days

### **Major Component Description**

#### **Hector**

*Extent:* 56 percent of the map unit  
*Geomorphic setting:* Mountain slopes on mountains  
*Position on the landform:* Mountainflanks  
*Parent material:* Loamy residuum weathered from sandstone  
*Slope:* 8 to 30 percent  
*Runoff:* Very high  
*Soil depth:* 10 to 20 inches to lithic bedrock  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderately rapid  
*Slowest permeability class within a depth of 80 inches:* Impermeable  
*Drainage class:* Well drained

*Available water capacity:* About 1.7 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 7e  
*Ecological site number and name:* R117XY088OK, Shallow Savannah PE 72–76  
*Typical profile:*  
A—0 to 7 inches; stony fine sandy loam  
Bw—7 to 15 inches; gravelly fine sandy loam  
R—15 to 30 inches; bedrock

**Linker**

*Extent:* 44 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Position on hillslope:* Backslopes  
*Parent material:* Loamy residuum weathered from sandstone  
*Slope:* 8 to 30 percent  
*Runoff:* High  
*Soil depth:* 20 to 40 inches to lithic bedrock  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate  
*Slowest permeability class within a depth of 80 inches:* Impermeable  
*Drainage class:* Well drained  
*Available water capacity:* About 4.0 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 7e  
*Ecological site number and name:* R117XY075OK, Sandy Savannah PE 72–76  
*Typical profile:*  
A—0 to 6 inches; stony fine sandy loam  
BA—6 to 11 inches; gravelly loam  
Bt—11 to 32 inches; gravelly sandy clay loam  
R—32 to 35 inches; bedrock

**JaA—Jay silt loam, 1 to 3 percent slopes**

**Map Unit Setting**

*Major land resource area:* 116A  
*Elevation:* 1,200 to 1,500 feet  
*Mean annual precipitation:* 42 to 46 inches  
*Mean annual air temperature:* 54 to 57 degrees F  
*Frost-free period:* 160 to 195 days

**Major Component Description**

**Jay**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Position on hillslope:* Shoulders  
*Parent material:* Loamy residuum weathered from siltstone  
*Slope:* 1 to 3 percent  
*Runoff:* Medium  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderately slow

*Slowest permeability class within a depth of 80 inches:* Slow  
*Drainage class:* Moderately well drained  
*Available water capacity:* About 9.2 inches  
*Depth to the top of the seasonal high water table:* 2.0 to 3.0 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 2e  
*Ecological site number and name:* R116AY059OK, Loamy Prairie (Northeast) PE 72–80  
*Typical profile:*  
    Ap—0 to 13 inches; silt loam  
    Bt—13 to 26 inches; silty clay loam  
    B'x—26 to 70 inches; silty clay loam

### **LkC—Linker fine sandy loam, 3 to 5 percent slopes**

#### **Map Unit Setting**

*Major land resource area:* 117  
*Elevation:* 500 to 2,800 feet  
*Mean annual precipitation:* 43 to 50 inches  
*Mean annual air temperature:* 61 to 64 degrees F  
*Frost-free period:* 175 to 210 days

#### **Major Component Description**

##### **Linker**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Position on hillslope:* Backslopes  
*Parent material:* Loamy residuum weathered from sandstone  
*Slope:* 3 to 5 percent  
*Runoff:* High  
*Soil depth:* 20 to 40 inches to lithic bedrock  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate  
*Slowest permeability class within a depth of 80 inches:* Impermeable  
*Drainage class:* Well drained  
*Available water capacity:* About 5.2 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 3e  
*Ecological site number and name:* R117XY075OK, Sandy Savannah PE 72–76  
*Typical profile:*  
    Ap—0 to 6 inches; fine sandy loam  
    BA—6 to 11 inches; gravelly loam  
    Bt—11 to 39 inches; gravelly clay loam  
    R—39 to 42 inches; bedrock

### **LoB—Tonti gravelly silt loam, 1 to 3 percent slopes**

#### **Map Unit Setting**

*Major land resource area:* 116A  
*Elevation:* 900 to 1,400 feet

*Mean annual precipitation:* 42 to 46 inches  
*Mean annual air temperature:* 54 to 57 degrees F  
*Frost-free period:* 160 to 200 days

**Major Component Description**

**Tonti**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Position on hillslope:* Shoulders  
*Parent material:* Loamy residuum weathered from cherty limestone  
*Slope:* 1 to 3 percent  
*Runoff:* Very high  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Slow  
*Slowest permeability class within a depth of 80 inches:* Slow  
*Drainage class:* Moderately well drained  
*Available water capacity:* About 6.3 inches  
*Depth to the top of the seasonal high water table:* 1.5 to 2.5 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 3s  
*Typical profile:*  
    Ap—0 to 10 inches; gravelly silt loam  
    Bt—10 to 22 inches; gravelly silty clay loam  
    Btx—22 to 30 inches; very gravelly silty clay loam  
    BC—30 to 66 inches; very gravelly silty clay

**M-W—Miscellaneous water**

**Map Unit Setting**

*Major land resource area:* 116A  
*Elevation:* 850 to 1,200 feet  
*Mean annual precipitation:* 32 to 36 inches  
*Mean annual air temperature:* 60 to 61 degrees F  
*Frost-free period:* 200 to 210 days

**Major Component Description**

**Miscellaneous water**

*Extent:* 100 percent of the map unit  
*Definition:* This unit consists of areas of waste water, such as sewage lagoons and industrial waste water.  
*Geomorphic setting:* Sewage lagoons  
*Land capability classification, nonirrigated:* 8w

**NaB—Newtonia silt loam, 1 to 3 percent slopes**

**Map Unit Setting**

*Major land resource area:* 112  
*Elevation:* 500 to 1,200 feet  
*Mean annual precipitation:* 37 to 43 inches  
*Mean annual air temperature:* 57 to 63 degrees F  
*Frost-free period:* 200 to 220 days

**Major Component Description**

**Newtonia**

*Extent:* 100 percent of the map unit

*Geomorphic setting:* Hillslopes on hills on uplands

*Position on hillslope:* Shoulders

*Parent material:* Loess and/or silty residuum

*Slope:* 1 to 3 percent

*Runoff:* Low

*Soil depth:* Greater than 60 inches

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate

*Slowest permeability class within a depth of 80 inches:* Moderate

*Drainage class:* Well drained

*Available water capacity:* About 11.1 inches

*Depth to the top of the seasonal high water table:* Greater than 6 feet

*Flooding:* None

*Ponding:* None

*Land capability classification, nonirrigated:* 2e

*Ecological site number and name:* R112XY059OK, Loamy Prairie (Northeast) PE 62–80

*Typical profile:*

A—0 to 10 inches; silt loam

AB—10 to 16 inches; silt loam

Bt1—16 to 26 inches; silty clay loam

Bt2—26 to 54 inches; silty clay loam

BC—54 to 62 inches; silty clay loam

**NaC—Newtonia silt loam, 3 to 5 percent slopes**

**Map Unit Setting**

*Major land resource area:* 112

*Elevation:* 500 to 1,200 feet

*Mean annual precipitation:* 37 to 43 inches

*Mean annual air temperature:* 57 to 63 degrees F

*Frost-free period:* 200 to 220 days

**Major Component Description**

**Newtonia**

*Extent:* 100 percent of the map unit

*Geomorphic setting:* Hillslopes on hills on uplands

*Position on hillslope:* Backslopes

*Parent material:* Loess and/or silty residuum

*Slope:* 3 to 5 percent

*Runoff:* Low

*Soil depth:* Greater than 60 inches

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate

*Slowest permeability class within a depth of 80 inches:* Moderate

*Drainage class:* Well drained

*Available water capacity:* About 11.1 inches

*Depth to the top of the seasonal high water table:* Greater than 6 feet

*Flooding:* None

*Ponding:* None

*Land capability classification, nonirrigated:* 3e

*Ecological site number and name:* R112XY059OK, Loamy Prairie (Northeast) PE  
62–80

*Typical profile:*

A—0 to 10 inches; silt loam  
AB—10 to 16 inches; silt loam  
Bt1—16 to 26 inches; silty clay loam  
Bt2—26 to 54 inches; silty clay loam  
BC—54 to 62 inches; silty clay loam

***NaC2—Newtonia silt loam, 3 to 5 percent slopes, eroded***

***Map Unit Setting***

*Major land resource area:* 112  
*Elevation:* 500 to 1,200 feet  
*Mean annual precipitation:* 37 to 43 inches  
*Mean annual air temperature:* 57 to 63 degrees F  
*Frost-free period:* 200 to 220 days

***Major Component Description***

***Newtonia***

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Position on hillslope:* Backslopes  
*Parent material:* Loess and/or silty residuum  
*Slope:* 3 to 5 percent  
*Runoff:* Low  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate  
*Slowest permeability class within a depth of 80 inches:* Moderate  
*Drainage class:* Well drained  
*Available water capacity:* About 10.1 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 3e  
*Ecological site number and name:* R112XY856OK, Eroded Loamy Prairie PE  
62–80

*Typical profile:*

A—0 to 6 inches; silt loam  
AB—6 to 12 inches; silt loam  
Bt1—12 to 22 inches; silty clay loam  
Bt2—22 to 54 inches; silty clay loam  
BC—54 to 62 inches; silty clay loam

***NixB—Nixa gravelly silt loam, 0 to 3 percent slopes***

***Map Unit Setting***

*Major land resource area:* 116A  
*Elevation:* 800 to 1,500 feet  
*Mean annual precipitation:* 42 to 46 inches  
*Mean annual air temperature:* 54 to 57 degrees F  
*Frost-free period:* 160 to 195 days

### **Major Component Description**

#### **Nixa**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Position on hillslope:* Shoulders  
*Parent material:* Loamy residuum weathered from cherty limestone  
*Slope:* 0 to 3 percent  
*Runoff:* High  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Very slow  
*Slowest permeability class within a depth of 80 inches:* Very slow  
*Drainage class:* Moderately well drained  
*Available water capacity:* About 6.2 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 3s  
*Ecological site number and name:* R116AY092OK, Smooth Chert Savannah PE 72–80  
*Typical profile:*  
A—0 to 4 inches; gravelly silt loam  
E—4 to 12 inches; gravelly silt loam  
Bw—12 to 19 inches; very gravelly silty clay loam  
E'—19 to 23 inches; very gravelly silt loam  
B'x—23 to 42 inches; very gravelly silt loam  
C—42 to 80 inches; very gravelly clay loam

### **OkA—Okemah silty clay loam, 0 to 1 percent slopes**

#### **Map Unit Setting**

*Major land resource area:* 112  
*Elevation:* 500 to 1,000 feet  
*Mean annual precipitation:* 37 to 46 inches  
*Mean annual air temperature:* 57 to 64 degrees F  
*Frost-free period:* 190 to 220 days

### **Major Component Description**

#### **Okemah**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Terraces on uplands  
*Position on the landform:* Treads  
*Parent material:* Clayey and loamy colluvium or alluvium over clayey residuum weathered from shale  
*Slope:* 0 to 1 percent  
*Runoff:* Very high  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Slow  
*Slowest permeability class within a depth of 80 inches:* Slow  
*Drainage class:* Somewhat poorly drained  
*Available water capacity:* About 10.6 inches  
*Depth to the top of the seasonal high water table:* 1.0 to 2.5 feet  
*Flooding:* None  
*Ponding:* None



*Land capability classification, nonirrigated:* 1

*Ecological site number and name:* R112XY059OK, Loamy Prairie (Northeast) PE  
62–80

*Typical profile:*

- A—0 to 11 inches; silty clay loam
- BA—11 to 20 inches; silty clay loam
- Bt—20 to 40 inches; silty clay
- BC—40 to 62 inches; silty clay

### ***OkB—Okemah silty clay loam, 1 to 3 percent slopes***

#### ***Map Unit Setting***

*Major land resource area:* 112

*Elevation:* 500 to 1,000 feet

*Mean annual precipitation:* 37 to 46 inches

*Mean annual air temperature:* 57 to 64 degrees F

*Frost-free period:* 190 to 220 days

#### ***Major Component Description***

##### ***Okemah***

*Extent:* 100 percent of the map unit

*Geomorphic setting:* Terraces on uplands

*Position on the landform:* Treads

*Parent material:* Clayey and loamy colluvium or alluvium over clayey residuum  
weathered from shale

*Slope:* 1 to 3 percent

*Runoff:* Very high

*Soil depth:* Greater than 60 inches

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive  
layer:* Slow

*Slowest permeability class within a depth of 80 inches:* Slow

*Drainage class:* Somewhat poorly drained

*Available water capacity:* About 10.6 inches

*Depth to the top of the seasonal high water table:* 1.0 to 2.5 feet

*Flooding:* None

*Ponding:* None

*Land capability classification, nonirrigated:* 2e

*Ecological site number and name:* R112XY059OK, Loamy Prairie (Northeast) PE  
62–80

*Typical profile:*

- A—0 to 11 inches; silty clay loam
- BA—11 to 20 inches; silty clay loam
- Bt—20 to 40 inches; silty clay
- BC—40 to 62 inches; silty clay

### ***OkC—Okemah silty clay loam, 3 to 5 percent slopes***

#### ***Map Unit Setting***

*Major land resource area:* 112

*Elevation:* 500 to 1,000 feet

*Mean annual precipitation:* 37 to 46 inches

*Mean annual air temperature:* 57 to 64 degrees F

*Frost-free period:* 190 to 220 days

### **Major Component Description**

#### **Okemah**

*Extent:* 100 percent of the map unit

*Geomorphic setting:* Terraces on uplands

*Position on the landform:* Risers

*Parent material:* Clayey and loamy colluvium or alluvium over clayey residuum weathered from shale

*Slope:* 3 to 5 percent

*Runoff:* Very high

*Soil depth:* Greater than 60 inches

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Slow

*Slowest permeability class within a depth of 80 inches:* Slow

*Drainage class:* Somewhat poorly drained

*Available water capacity:* About 10.6 inches

*Depth to the top of the seasonal high water table:* 1.0 to 2.5 feet

*Flooding:* None

*Ponding:* None

*Land capability classification, nonirrigated:* 3e

*Ecological site number and name:* R112XY059OK, Loamy Prairie (Northeast) PE 62–80

*Typical profile:*

A—0 to 11 inches; silty clay loam

BA—11 to 20 inches; silty clay loam

Bt—20 to 40 inches; silty clay

BC—40 to 62 inches; silty clay

### **Os—Osage clay, 0 to 1 percent slopes, occasionally flooded**

#### **Map Unit Setting**

*Major land resource area:* 112

*Elevation:* 740 to 800 feet

*Mean annual precipitation:* 37 to 42 inches

*Mean annual air temperature:* 55 to 61 degrees F

*Frost-free period:* 190 to 200 days

### **Major Component Description**

#### **Osage**

*Extent:* 100 percent of the map unit

*Geomorphic setting:* Flood plains in river valleys

*Parent material:* Clayey alluvium

*Slope:* 0 to 1 percent

*Runoff:* Very high

*Soil depth:* Greater than 60 inches

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Very slow

*Slowest permeability class within a depth of 80 inches:* Very slow

*Drainage class:* Poorly drained

*Available water capacity:* About 6.2 inches

*Depth to the top of the seasonal high water table:* 0.0 to 1.0 foot

*Flooding:* Occasional

*Ponding:* None

*Land capability classification, nonirrigated:* 5w

*Ecological site number and name:* R112XY045OK, Heavy Bottomland PE 62–80

*Typical profile:*

A1—0 to 11 inches; clay

A2—11 to 60 inches; clay

## ***PIT—Pits***

### ***Map Unit Setting***

*Major land resource area:* 116A

*Elevation:* 500 to 2,200 feet

*Mean annual precipitation:* 22 to 48 inches

*Mean annual air temperature:* 57 to 64 degrees F

*Frost-free period:* 190 to 240 days

### ***Major Component Description***

#### ***Pits***

*Extent:* 100 percent of the map unit

*Geomorphic setting:* Borrow pits

*Parent material:* Mine spoil or earthy fill

*Slope:* 0 to 4 percent

*Runoff:* High

*Land capability classification, nonirrigated:* 8s

## ***Rs—Rock outcrop-Hector complex, 40 to 100 percent slopes***

### ***Map Unit Setting***

*Major land resource area:* 117

*Elevation:* 500 to 2,400 feet

*Mean annual precipitation:* 22 to 50 inches

*Mean annual air temperature:* 57 to 64 degrees F

*Frost-free period:* 175 to 240 days

### ***Major Component Description***

#### ***Rock outcrop***

*Extent:* 50 percent of the map unit

*Geomorphic setting:* Hillslopes on hills on uplands

*Position on hillslope:* Shoulder summits

*Slope:* 40 to 100 percent

*Runoff:* Very high

*Land capability classification, nonirrigated:* 8s

#### ***Hector***

*Extent:* 45 percent of the map unit

*Geomorphic setting:* Mountain slopes on mountains

*Position on the landform:* Mountainflanks

*Parent material:* Loamy residuum weathered from sandstone

*Slope:* 40 to 60 percent

*Runoff:* Very high

*Soil depth:* 10 to 20 inches to lithic bedrock

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderately rapid

*Slowest permeability class within a depth of 80 inches:* Impermeable

*Drainage class:* Well drained

*Available water capacity:* About 1.7 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 8e  
*Ecological site number and name:* R117XY079OK, Savannah Breaks PE 72–76  
*Typical profile:*  
A—0 to 7 inches; stony fine sandy loam  
Bw—7 to 15 inches; gravelly fine sandy loam  
R—15 to 30 inches; bedrock

### ***Additional Components***

*Linker soils:* 5 percent of the map unit

## ***SaA—Britwater silt loam, 0 to 1 percent slopes***

### ***Map Unit Setting***

*Major land resource area:* 116A  
*Elevation:* 1,000 to 1,400 feet  
*Mean annual precipitation:* 38 to 48 inches  
*Mean annual air temperature:* 54 to 58 degrees F  
*Frost-free period:* 160 to 195 days

### ***Major Component Description***

#### ***Britwater***

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Paleoterraces on uplands  
*Position on the landform:* Treads  
*Parent material:* Loamy alluvium derived from cherty limestone  
*Slope:* 0 to 1 percent  
*Runoff:* Low  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate  
*Slowest permeability class within a depth of 80 inches:* Moderate  
*Drainage class:* Well drained  
*Available water capacity:* About 8.4 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 1  
*Ecological site number and name:* R116AY092OK, Smooth Chert Savannah PE 72–80  
*Typical profile:*  
Ap—0 to 18 inches; silt loam  
Bt—18 to 32 inches; gravelly silty clay loam  
2Bt—32 to 63 inches; very gravelly silty clay loam

## ***SaB—Britwater silt loam, 1 to 3 percent slopes***

### ***Map Unit Setting***

*Major land resource area:* 116A  
*Elevation:* 1,000 to 1,400 feet

*Mean annual precipitation:* 38 to 48 inches  
*Mean annual air temperature:* 54 to 58 degrees F  
*Frost-free period:* 160 to 195 days

**Major Component Description**

**Britwater**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Paleoterraces on uplands  
*Position on the landform:* Treads  
*Parent material:* Loamy alluvium derived from cherty limestone  
*Slope:* 1 to 3 percent  
*Runoff:* Low  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate  
*Slowest permeability class within a depth of 80 inches:* Moderate  
*Drainage class:* Well drained  
*Available water capacity:* About 8.4 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 2e  
*Ecological site number and name:* R116AY092OK, Smooth Chert Savannah PE 72–80  
*Typical profile:*  
    Ap—0 to 18 inches; silt loam  
    Bt—18 to 32 inches; gravelly silty clay loam  
    2Bt—32 to 63 inches; very gravelly silty clay loam

**SgB—Britwater gravelly silt loam, 1 to 3 percent slopes**

**Map Unit Setting**

*Major land resource area:* 116A  
*Elevation:* 1,000 to 1,400 feet  
*Mean annual precipitation:* 38 to 48 inches  
*Mean annual air temperature:* 54 to 58 degrees F  
*Frost-free period:* 160 to 195 days

**Major Component Description**

**Britwater**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Paleoterraces on uplands  
*Position on the landform:* Treads  
*Parent material:* Loamy alluvium derived from cherty limestone  
*Slope:* 1 to 3 percent  
*Runoff:* Low  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate  
*Slowest permeability class within a depth of 80 inches:* Moderate  
*Drainage class:* Well drained  
*Available water capacity:* About 7.9 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None

*Ponding:* None

*Land capability classification, nonirrigated:* 3e

*Ecological site number and name:* R116AY092OK, Smooth Chert Savannah PE  
72–80

*Typical profile:*

Ap—0 to 18 inches; gravelly silt loam

Bt—18 to 32 inches; gravelly silty clay loam

2Bt—32 to 63 inches; very gravelly silty clay loam

## ***SgD—Britwater gravelly silt loam, 3 to 8 percent slopes***

### ***Map Unit Setting***

*Major land resource area:* 116A

*Elevation:* 1,000 to 1,400 feet

*Mean annual precipitation:* 38 to 48 inches

*Mean annual air temperature:* 54 to 58 degrees F

*Frost-free period:* 160 to 195 days

### ***Major Component Description***

#### ***Britwater***

*Extent:* 100 percent of the map unit

*Geomorphic setting:* Paleoterraces on uplands

*Position on the landform:* Risers

*Parent material:* Loamy alluvium derived from cherty limestone

*Slope:* 3 to 8 percent

*Runoff:* Medium

*Soil depth:* Greater than 60 inches

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate

*Slowest permeability class within a depth of 80 inches:* Moderate

*Drainage class:* Well drained

*Available water capacity:* About 7.9 inches

*Depth to the top of the seasonal high water table:* Greater than 6 feet

*Flooding:* None

*Ponding:* None

*Land capability classification, nonirrigated:* 4e

*Ecological site number and name:* R116AY092OK, Smooth Chert Savannah PE  
72–80

*Typical profile:*

Ap—0 to 18 inches; gravelly silt loam

Bt—18 to 32 inches; gravelly silty clay loam

2Bt—32 to 63 inches; very gravelly silty clay loam

## ***Sm—Healing silt loam, 0 to 1 percent slopes, occasionally flooded***

### ***Map Unit Setting***

*Major land resource area:* 116A

*Elevation:* 500 to 1,000 feet

*Mean annual precipitation:* 42 to 46 inches

*Mean annual air temperature:* 54 to 57 degrees F

*Frost-free period:* 160 to 195 days

### **Major Component Description**

#### **Healing**

*Extent:* 100 percent of the map unit

*Geomorphic setting:* Flood plains in valleys

*Parent material:* Silty alluvium

*Slope:* 0 to 1 percent

*Runoff:* Low

*Soil depth:* Greater than 60 inches

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate

*Slowest permeability class within a depth of 80 inches:* Moderate

*Drainage class:* Well drained

*Available water capacity:* About 11.2 inches

*Depth to the top of the seasonal high water table:* Greater than 6 feet

*Flooding:* Occasional

*Ponding:* None

*Land capability classification, nonirrigated:* 2w

*Ecological site number and name:* R116AY050OK, Loamy Bottomland PE 72–80

*Typical profile:*

A—0 to 12 inches; silt loam

Bt1—12 to 43 inches; silt loam

Bt2—43 to 60 inches; gravelly silt loam

### **Sn—Razort gravelly loam, 0 to 3 percent slopes, occasionally flooded**

#### **Map Unit Setting**

*Major land resource area:* 116A

*Elevation:* 500 to 1,000 feet

*Mean annual precipitation:* 42 to 46 inches

*Mean annual air temperature:* 52 to 57 degrees F

*Frost-free period:* 160 to 195 days

### **Major Component Description**

#### **Razort**

*Extent:* 100 percent of the map unit

*Geomorphic setting:* Flood plains in valleys

*Parent material:* Silty alluvium

*Slope:* 0 to 3 percent

*Runoff:* Low

*Soil depth:* Greater than 60 inches

*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate

*Slowest permeability class within a depth of 80 inches:* Moderate

*Drainage class:* Well drained

*Available water capacity:* About 8.4 inches

*Depth to the top of the seasonal high water table:* Greater than 6 feet

*Flooding:* Occasional

*Ponding:* None

*Land capability classification, nonirrigated:* 3e

*Ecological site number and name:* R116AY050OK, Loamy Bottomland PE 72–80

*Typical profile:*

A—0 to 12 inches; gravelly loam

Bw—12 to 43 inches; gravelly loam

C—43 to 60 inches; very gravelly loam

## **SrA—Stigler silt loam, 0 to 1 percent slopes**

### **Map Unit Setting**

*Major land resource area:* 116A  
*Elevation:* 500 to 1,100 feet  
*Mean annual precipitation:* 36 to 54 inches  
*Mean annual air temperature:* 60 to 64 degrees F  
*Frost-free period:* 190 to 220 days

### **Major Component Description**

#### **Stigler**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Paleoterraces on uplands  
*Position on the landform:* Treads  
*Parent material:* Loamy and clayey alluvium and/or colluvium over sandstone and shale  
*Slope:* 0 to 1 percent  
*Runoff:* Negligible  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Very slow  
*Slowest permeability class within a depth of 80 inches:* Very slow  
*Drainage class:* Somewhat poorly drained  
*Available water capacity:* About 10.2 inches  
*Depth to the top of the seasonal high water table:* 1.0 to 2.0 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 3s  
*Ecological site number and name:* R116AY092OK, Smooth Chert Savannah PE 72–80  
*Typical profile:*  
Ap—0 to 6 inches; silt loam  
E—6 to 12 inches; silt loam  
BE—12 to 18 inches; silty clay loam  
Bt—18 to 37 inches; silty clay loam  
Btg—37 to 65 inches; silty clay loam

## **SuC2—Apperson silty clay loam, 3 to 5 percent slopes, eroded**

### **Map Unit Setting**

*Major land resource area:* 112  
*Elevation:* 800 to 1,200 feet  
*Mean annual precipitation:* 36 to 44 inches  
*Mean annual air temperature:* 57 to 64 degrees F  
*Frost-free period:* 190 to 220 days

### **Major Component Description**

#### **Apperson**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Position on hillslope:* Backslopes  
*Parent material:* Calcareous clayey residuum weathered from limestone  
*Slope:* 3 to 5 percent



*Runoff:* Very high  
*Soil depth:* 40 to 60 inches to lithic bedrock  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Very slow  
*Slowest permeability class within a depth of 80 inches:* Impermeable  
*Drainage class:* Somewhat poorly drained  
*Available water capacity:* About 8.0 inches  
*Depth to the top of the seasonal high water table:* 1.5 to 2.0 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 3e  
*Ecological site number and name:* R112XY810OK, Eroded Claypan Prairie PE 62–80  
*Typical profile:*  
A—0 to 3 inches; silty clay loam  
BA—3 to 17 inches; silty clay loam  
Bt—17 to 48 inches; silty clay  
R—48 to 60 inches; bedrock

### **TkA—Taloka silt loam, 0 to 1 percent slopes**

#### **Map Unit Setting**

*Major land resource area:* 112  
*Elevation:* 500 to 1,200 feet  
*Mean annual precipitation:* 37 to 45 inches  
*Mean annual air temperature:* 57 to 63 degrees F  
*Frost-free period:* 190 to 220 days

#### **Major Component Description**

##### **Taloka**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Paleoterraces on uplands  
*Position on the landform:* Treads  
*Parent material:* Loamy and clayey alluvium and/or colluvium derived from sandstone and shale  
*Slope:* 0 to 1 percent  
*Runoff:* Very high  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Very slow  
*Slowest permeability class within a depth of 80 inches:* Very slow  
*Drainage class:* Somewhat poorly drained  
*Available water capacity:* About 11.2 inches  
*Depth to the top of the seasonal high water table:* 0.5 to 2.0 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 3s  
*Ecological site number and name:* R112XY059OK, Loamy Prairie (Northeast) PE 62–80  
*Typical profile:*  
Ap—0 to 12 inches; silt loam  
E—12 to 22 inches; silt loam  
Bt1—22 to 38 inches; silt loam  
Bt2—38 to 49 inches; silty clay  
BC—49 to 62 inches; silty clay loam

## **TrD—Shidler-Rock outcrop complex, 2 to 8 percent slopes**

### **Map Unit Setting**

*Major land resource area:* 112  
*Elevation:* 500 to 2,200 feet  
*Mean annual precipitation:* 22 to 48 inches  
*Mean annual air temperature:* 57 to 64 degrees F  
*Frost-free period:* 190 to 240 days

### **Major Component Description**

#### **Shidler**

*Extent:* 62 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Position on hillslope:* Shoulders; backslopes  
*Parent material:* Loamy residuum weathered from cherty limestone  
*Slope:* 2 to 8 percent  
*Runoff:* Very high  
*Soil depth:* 4 to 20 inches to lithic bedrock  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate  
*Slowest permeability class within a depth of 80 inches:* Impermeable  
*Drainage class:* Well drained  
*Available water capacity:* About 1.8 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 7s  
*Ecological site number and name:* R112XY098OK, Very Shallow (Eastern) PE 62–80  
*Typical profile:*  
    A—0 to 9 inches; silty clay loam  
    R—9 to 20 inches; bedrock

#### **Rock outcrop**

*Extent:* 38 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Slope:* 2 to 8 percent  
*Runoff:* Very high  
*Land capability classification, nonirrigated:* 8s

## **TrF—Shidler-Rock outcrop complex, 15 to 50 percent slopes**

### **Map Unit Setting**

*Major land resource area:* 112  
*Elevation:* 500 to 2,200 feet  
*Mean annual precipitation:* 22 to 48 inches  
*Mean annual air temperature:* 57 to 64 degrees F  
*Frost-free period:* 190 to 240 days

### **Major Component Description**

#### **Shidler**

*Extent:* 62 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Position on hillslope:* Backslopes

*Parent material:* Loamy residuum weathered from cherty limestone  
*Slope:* 15 to 25 percent  
*Runoff:* Very high  
*Soil depth:* 4 to 20 inches to lithic bedrock  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate  
*Slowest permeability class within a depth of 80 inches:* Impermeable  
*Drainage class:* Well drained  
*Available water capacity:* About 1.8 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* None  
*Ponding:* None  
*Land capability classification, nonirrigated:* 7e  
*Ecological site number and name:* R112XY098OK, Very Shallow (Eastern) PE 62–80  
*Typical profile:*  
    A—0 to 9 inches; silty clay loam  
    R—9 to 20 inches; bedrock

**Rock outcrop**

*Extent:* 38 percent of the map unit  
*Geomorphic setting:* Hillslopes on hills on uplands  
*Slope:* 15 to 50 percent  
*Runoff:* Very high  
*Land capability classification, nonirrigated:* 8s

**Vd—Verdigris silt loam, 0 to 1 percent slopes,  
occasionally flooded**

**Map Unit Setting**

*Major land resource area:* 112  
*Elevation:* 500 to 1,000 feet  
*Mean annual precipitation:* 38 to 47 inches  
*Mean annual air temperature:* 57 to 64 degrees F  
*Frost-free period:* 190 to 230 days

**Major Component Description**

**Verdigris**

*Extent:* 100 percent of the map unit  
*Geomorphic setting:* Flood plains in river valleys  
*Parent material:* Silty alluvium  
*Slope:* 0 to 1 percent  
*Runoff:* Low  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate  
*Slowest permeability class within a depth of 80 inches:* Moderate  
*Drainage class:* Well drained  
*Available water capacity:* About 11.8 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* Occasional  
*Ponding:* None  
*Land capability classification, nonirrigated:* 2w  
*Ecological site number and name:* R112XY050OK, Loamy Bottomland PE 62–80

*Typical profile:*

A—0 to 24 inches; silt loam  
Bw—24 to 60 inches; silty clay loam

***VrdA—Verdigris silt loam, 0 to 1 percent slopes,  
frequently flooded***

***Map Unit Setting***

*Major land resource area:* 112  
*Elevation:* 500 to 1,000 feet  
*Mean annual precipitation:* 35 to 56 inches  
*Mean annual air temperature:* 55 to 65 degrees F  
*Frost-free period:* 190 to 240 days

***Major Component Description***

***Verdigris***

*Extent:* 85 percent of the map unit  
*Geomorphic setting:* Flood plains in river valleys  
*Parent material:* Silty alluvium  
*Slope:* 0 to 1 percent  
*Runoff:* Low  
*Soil depth:* Greater than 60 inches  
*Slowest permeability class of the soil to a depth of 60 inches or above a restrictive layer:* Moderate  
*Slowest permeability class within a depth of 80 inches:* Moderate  
*Drainage class:* Well drained  
*Available water capacity:* About 11.8 inches  
*Depth to the top of the seasonal high water table:* Greater than 6 feet  
*Flooding:* Frequent  
*Ponding:* None  
*Land capability classification, nonirrigated:* 5w  
*Ecological site number and name:* R112XY050OK, Loamy Bottomland PE 62–80  
*Typical profile:*  
A—0 to 31 inches; silt loam  
Bw—31 to 60 inches; silty clay loam

***Additional Components***

*Osage soils:* 4 percent of the map unit  
*Roebuck soils:* 4 percent of the map unit  
*Cupco soils:* 3 percent of the map unit  
*Barge soils:* 2 percent of the map unit  
*Lightning soils:* 2 percent of the map unit

***W—Water***

***Map Unit Setting***

*Major land resource area:* 116A  
*Elevation:* 250 to 4,000 feet  
*Mean annual precipitation:* 22 to 48 inches  
*Mean annual air temperature:* 57 to 64 degrees F  
*Frost-free period:* 190 to 240 days

***Major Component Description***

**Water**

*Extent:* 100 percent of the map unit

*Definition:* This unit consists of areas of water, including ponds, lakes, rivers, and other bodies of fresh water.

*Land capability classification, nonirrigated:* 8w



# Use and Management of the Soils

---

For general and detailed information regarding the use and management of the map units in this survey, see the soil reports and report descriptions on the Web Soil Survey at <http://websoilsurvey.nrcs.usda.gov>. A soil survey is an inventory and evaluation of the soils in the survey area. It can be used to adjust land uses to the limitations and potentials of natural resources and the environment. Also, it can help to prevent soil-related failures in land uses.

In preparing a soil survey, soil scientists, conservationists, engineers, and others collect extensive field data about the nature and behavioral characteristics of the soils. They collect data on erosion, droughtiness, flooding, and other factors that affect various soil uses and management. Field experience and collected data on soil properties and performance are used as a basis in predicting soil behavior.

Information developed during a soil survey can be used to plan the use and management of soils for crops and pasture; as rangeland and woodland; as sites for buildings, sanitary facilities, highways and other transportation systems, and parks and other recreational facilities; and for wildlife habitat. It can be used to identify the potentials and limitations of each soil for specific land uses and to help prevent construction failures caused by unfavorable soil properties.

Interpretive ratings help engineers, planners, and others understand how soil properties influence important nonagricultural uses, such as building site development and construction materials. The ratings indicate the most restrictive soil features affecting the suitability of the soils for these uses.

Soils are rated in their natural state. Only normal practices for the rated use are considered. Unusual modifications to the site or soil material are not considered in the ratings. Where soils have limitations, engineers and others may be able to modify soil features or adjust the plans for a structure to compensate for most of the limitations. Most of these modifications, however, are costly. The final decision in selecting a site for a particular use generally involves weighing the costs of site preparation and maintenance.

Planners and others using soil survey information can evaluate the effect of specific land uses on productivity and on the environment in all or part of the survey area. The survey can help planners to maintain or create a land use pattern in harmony with the natural soil.

Contractors can use a survey to locate sources of sand, gravel, roadfill, and topsoil. They can use it to identify areas where bedrock, wetness, or very firm soil layers can cause difficulty in excavation.

Health officials, highway officials, engineers, and others may also find a soil survey useful. The survey can help them plan the safe disposal of wastes and locate sites for pavements, sidewalks, campgrounds, playgrounds, lawns, and trees and shrubs.

## Cropland

The main management concerns affecting nonirrigated cropland are conserving moisture, controlling soil blowing and water erosion, and maintaining soil fertility and tilth.

Conserving moisture primarily involves reducing the rates of evaporation and runoff and increasing the rate of water infiltration. Applying conservation tillage and conservation cropping systems, farming on the contour, stripcropping, establishing field windbreaks, and leaving crop residue on the surface conserve moisture.

Generally, a combination of several practices is needed to control soil blowing and water erosion. Conservation tillage, stripcropping, field windbreaks, tall-grass barriers, contour farming, conservation cropping systems, crop residue management, diversions, and grassed waterways help to prevent excessive soil loss.

Soil fertility can be effectively maintained by applying inorganic and organic fertilizers, including manure; incorporating crop residue or green manure crops into the soil; and using proper crop rotations. Controlling erosion helps to prevent the loss of organic matter and plant nutrients and thus helps to maintain productivity, although the level of fertility can be reduced even in areas where erosion is controlled. All of the soils that are used for nonirrigated crops in the survey area respond well to applications of fertilizer.

On irrigated soils, the main management concerns are efficient water use, nutrient management, erosion control, soil tilth, pest control, weed control, and timely planting and harvesting. An irrigation system that provides optimum control and distribution of water at minimum cost is needed. Overirrigation wastes water, leaches plant nutrients, and causes erosion. Also, it can create drainage problems, raise the water table, and increase soil salinity.

Planners of management systems for individual fields or farms should consider obtaining specific information from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

## Rangeland

Mark Moseley, range conservationist, Natural Resources Conservation Service, helped prepare this section.

Range, grazed forestland, and native pasture provide forage for livestock in the survey area.

*Range* is defined as land on which the native vegetation (the climax, or natural potential, plant community) is predominantly grasses, grasslike plants, forbs, and shrubs suitable for grazing and browsing. Range includes natural grasslands, savannas, many wetlands, some deserts, tundra, and certain shrub and forb communities. Range receives no regular or frequent cultural treatment. The composition and production of the plant community are determined by soil, climate, topography, overstory canopy, and grazing management.

*Grazed forestland* is defined as land on which the understory includes, as an integral part of the forest plant community, plants that can be grazed without significant impairment of other forest values.

*Native pasture* is defined as land on which the potential (climax) vegetation is forest but which is used and managed primarily for the production of native forage plants. Native pasture includes cutover forestland and forestland that has been cleared and is managed for native or naturalized forage plants.

Most of the rangeland in Cherokee County is in areas that are not economical to cultivate. These areas are on gently sloping to steep side slopes and a few narrow, very gently sloping to sloping summits. A few native-grass meadows are managed for hay production in the central and north-central parts of the county. Three distinct rangeland types are present. In the southeastern part of the county, most of the soils are loamy and are moderately deep or shallow over sandstone. These soils support an oak savannah rangeland that has low productivity due to the shallow rooting depth and low available water capacity. In the northeastern and western parts of the county, the soils are loamy and are dominantly moderately deep. There are also some



shallow and deep soils over shale or over shale interbedded with sandstone. The soils in the northeastern and western parts of the county support mid and tall grasses, and productivity is moderate. In the north-central, central, and south-central parts of the county, the soils are loamy and are dominantly moderately deep. There are also some shallow and deep soils over sandstone or over sandstone interbedded with shale. The soils in the north-central, central, and south-central parts of the county support tall and mid grasses that are moderately productive.

Approximately 75 percent of the annual production on rangeland grows in April, May, and June, coinciding with spring rains and moderate temperatures. A secondary growth period generally occurs in September and October, coinciding with fall rains and cooling temperatures.

Most of the local ranches and livestock farms are cow-calf operations. There are also some pure stocker enterprises and some ranches that diversify their cow-calf operation with stockers to provide greater flexibility.

Several livestock operations supplement the grazing of native rangeland with introduced grasses, such as bermudagrass and "plains" bluestem. Forage crops are also used. Protein, hay, and small-grain crops are used to supplement livestock through winter.

Droughts of varying lengths occur, and short-term summer droughts are common. Longer periods of drought, some lasting several months, are also frequent.

The pre-settlement vegetation evolved with periodic natural fires, droughts, migratory grazing by bison, and impact from many other wildlife species. The bison would heavily impact an area and then move to other grazing range.

Early settlement brought continuous grazing and eliminated much of the high-quality vegetation on some ecological sites. Areas that were once open-savannah ecological sites with a mixture of grasses, forbs, and scattered trees are now covered with oaks, a few tall and mid grasses, and low successional grasses and forbs. Some prairie sites are now growing low successional grasses and forbs instead of tall grasses. The amount of forage presently produced may be less than half of that originally produced. On some sites, the number of eastern redcedar has increased significantly due to a lack of prairie fires. Remnants of the original plant species are still found on most rangeland, however, and progressive grazing management can allow these high quality plants to re-establish without reseeding.

The table "Rangeland Productivity and Characteristic Plant Communities" is available on the Soil Data Mart at <http://soildatamart.nrcs.usda.gov>. This table shows, for each soil, the ecological site; the total annual production of vegetation in favorable, normal, and unfavorable years; the characteristic vegetation; and the average percentage of each species. Only those soils that are used as rangeland or are suited to use as rangeland are listed. Explanation of the column headings in this table follows.

An *ecological site* for rangeland is a distinctive kind of land and vegetation with specific physical characteristics that makes it different from other kinds of land in its ability to produce a distinctive kind and amount of vegetation.

Many different ecological sites are in the survey area. Over time, the combination of plants best suited to a particular soil and climate has become dominant. If the soil is not excessively disturbed, this group of plants is the natural plant community for the site. Natural plant communities are not static but vary slightly from year to year and place to place.

The relationship between soils and vegetation was ascertained during this survey; thus, ecological sites generally can be determined directly from the soil map. Soil properties that affect moisture supply and plant nutrients have the greatest influence on the productivity of range plants. Soil reaction, salt content, and a seasonal high water table are also important. The "Electronic Field Office Technical Guide," which is available at the local office of the Natural Resources Conservation Service or online

at <http://www.nrcs.usda.gov/technical/efotg/>, can provide specific information about ecological sites.

*Total production* is the amount of vegetation that can be expected to grow annually on well managed rangeland. It includes all vegetation, whether or not it is palatable to grazing animals. It includes the current year's growth of leaves, twigs, and fruit of woody plants. It does not include the increase in stem diameter of trees and shrubs. It is expressed in pounds per acre of air-dry vegetation for favorable, normal, and unfavorable years. In a favorable year, the amount and distribution of precipitation and the temperatures make growing conditions substantially better than average. In a normal year, growing conditions are near the historical monthly average. In an unfavorable year, growing conditions are well below average, generally because of low available soil moisture. Figure 2 shows a typical growth curve that represents the percentage of total growth that occurs each month for native vegetation and other forage. Dry weight is the total annual yield per acre of air-dry vegetation. Yields are adjusted to a percent of air-dry moisture content. The relationship of green weight to air-dry weight varies according to such factors as stage of maturity, exposure, amount of shade, recent rains, and unseasonable dry periods.

*Characteristic vegetation* consists of the grasses, forbs, and shrubs that make up most of the potential natural plant community on each soil. The plants are listed by common name. Under composition, the anticipated percentage of the total annual production is given for each species making up the characteristic vegetation. The amount that can be used as forage depends on the kinds of grazing animals and on the grazing season.

### Similarity Index

The similarity index indicates on a percentage basis the extent to which the present plant community resembles a specified vegetative state on an ecological site. The Natural Resources Conservation Service uses similarity index two ways.

	Jan.	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sept.	Oct.	Nov.	Dec.
IMPROVED BERMUDAGRASS				5	25	35	20	10	5			
WEeping LOVEGRASS			3	20	25	20	15	6	11			
INTRODUCED BLUESTEM				3	15	26	22	18	10	1		
SMALL GRAIN GRAZEOUT	3	9	29	27	18				1	4	6	3
FORAGE SORGHUM						14	33	33	20			
NATIVE GRASS	1	1	2	10	20	27	16	8	5	2	2	1

Figure 2.—Typical growth curves for various kinds of forage in the survey area. The growth curve for each kind of forage indicates the percentage of the total annual growth that occurs each month.

The first use compares the present vegetation on an ecological site to the presumed historic vegetation for that site. A similarity index of 70 would suggest that the present plant community contains 70 percent of the presumed historic plant community for that site. This comparison provides a basis for examining the extent and direction of changes between current vegetation and historic vegetation.

The second use measures how near the current plant community is to the landowners goal for the land. The management goal for rangeland is not necessarily a similarity index of 100 as compared to the historic plant community. Therefore, the similarity index can represent the percentage of the plant community that resembles a desired plant community.

Abnormal disturbances that change the natural plant community include repeated overuse by livestock, excessive burning, erosion, and cultivation. Grazing animals select the most palatable plants. These plants eventually die if they are continually grazed at a severity that does not allow for recovery. A very severe disturbance can completely destroy the natural community. Under such conditions, the less desirable plants, such as annuals and weed-like plants, can increase in abundance. If the plant community and the soils have not deteriorated significantly and proper range management is applied, the site can eventually return to predominantly natural plants.

Knowledge of the ecological site is necessary as a basis for planning and applying the management needed to maintain or improve the desired plant community for selected uses. Such information is needed to support management objectives, planned grazing systems, stocking rates, and wildlife management practices; to improve the potential of an area for recreational uses; and to improve the condition of watersheds.

## **Rangeland Management**

Rangeland management requires a knowledge of the kinds of soil in an area and of the potential natural plant community. It also requires an evaluation of the similarity index.

Effective range management conserves rainfall, enhances water quality, reduces the hazard of downstream flooding, improves yields, provides forage for livestock and wildlife, enhances recreational opportunities, and protects the soil. The main management concern is recognizing important changes in the plant cover or the range trend. These changes occur gradually and can be overlooked.

Each range manager should evaluate the type of plant community that best supports the ranch and then apply management and ecological principles to achieve the goals. The desired plant community should be within the capabilities of the land.

The range management practices used in Cherokee County include prescribed grazing, stock-water developments, and fences. If undesirable plants become dominant, range seeding, brush management, or prescribed burning are commonly used.

Range management includes four major considerations:

- *Grazing distribution*, which is achieved by managing livestock to graze all parts of the grazing unit equally.
- *Selective grazing*, which occurs because animals graze preferred plants to balance their diets. If selective grazing occurs repeatedly, the preferred plants are damaged.
- *Proper stocking rates*, which are achieved by balancing animal numbers with forage production.
- *Rest periods*, during which grazed plants are given enough rest to recover and to maintain growth.

An important principle of range management is that forage production is controlled by rainfall while composition is determined by grazing management.

The setting of stocking rates is not an exact science because of influences from grazing management, season of use, mix of livestock, and seasonal production of forage. Some general rules, however, can be helpful. To maintain a nutritional cover of plants, about 50 percent of the annual growth of the most important grazing plants should remain at the end of the grazing season. Plants can be removed not only through grazing by livestock but also through grazing by rodents, insects, and wildlife and through the deterioration caused by climatic variations. Because of these factors, a safe initial stocking rate for livestock should be calculated on the basis of 25 percent of the total annual growth, by weight, of the vegetation.

For example, production could be 3,500 pounds per acre of air-dry grasses, forbs, and limited woody species for an average season on a Loamy Prairie ecological site with a similarity index of 70 to the historic plant community. Twenty-five percent of this production would be 875 pounds per acre.

A 1,000-pound cow and her calf is equivalent to one animal unit (AU) and consume about 2.6 percent of her body weight (26 pounds) of forage per day. Therefore, in 1 month an animal unit could consume 790 pounds of native vegetation, depending on the quality and stage of growth of the plants (26 pounds per day times 365 days per year divided by 12 months per year).

Dividing 875 pounds (the forage allocation) by 26 pounds (the forage required per day for 1 animal unit) suggests that 1 acre of Loamy Prairie ecological site with a similarity index of 70 would feed 1 cow and calf for 33.6 days. To convert forage available from 1 acre to animal unit months (AUM), the available forage (875 pounds) is divided by the amount required to feed 1 animal unit for 1 month (790 pounds). Therefore, 1 acre would provide 1.1 AUM of grazing and 10.9 acres would feed 1 cow and calf for 12 months.

Another approach is to calculate the annual forage needs of an animal unit (790 pounds per month times 12 months equals 9,490 pounds). Dividing the 875 pounds of usable forage per acre into the 9,490 pounds needed by the cow and calf reveals that approximately 10.9 acres is needed for 1 cow annually. Stocking rate calculations should be adjusted for animal size, grazing system, and grazing season.

More information about planning a grazing program is available from the local office of the Natural Resources Conservation Service.

## Ecological Site Descriptions

Thirteen ecological sites are recognized in Cherokee County. The ecological site identifier has eleven characters. The "R" indicates an ecological site. The next four characters identify the major land resource area. The sixth character identifies the major land resource unit subdivision. The next three characters identify the individual ecological site number. The final two characters identify the state. The identifier is followed by the proper name for the ecological site. The following descriptions include a list of the plants that are characteristic of the sites. Detailed ecological site descriptions are available at the local office of the Natural Resources Conservation Service.

**R112XY010OK, Claypan Prairie PE 62-80.**—This site is in nearly level to moderately sloping areas on uplands. Elevations range from 500 feet on the east side of the major land resource area to 1,100 feet on the west side. The plant community is composed of about 90 to 95 percent grasses, 1 to 3 percent legumes, 1 to 3 percent forbs, and 1 to 3 percent woody species. The main grasses are big bluestem (25 to 35 percent), Indiangrass (25 to 35 percent), little bluestem (10 to 20 percent), switchgrass (5 to 10 percent), Canada wildrye (2 to 5 percent), and other grasses (0 to 5 percent). The other grasses include dropseed species, sideoats grama, Scribner's panicum, hairy grama, buffalograss, red lovegrass, tridens, and knotroot bristlegass. The main legumes are leadplant (trace to 2 percent), Illinois

## Supplement to the Soil Survey of Cherokee County, Oklahoma

bundleflower (trace to 2 percent), scurfpea (trace to 2 percent), and sessile-leaved tickclover (trace to 2 percent). Other legumes include wild indigo, slimflower scurfpea, and yellow neptune. The main forbs are black sampson (trace to 2 percent) and dotted gayfeather (trace to 2 percent). Other forbs include ashy sunflower, milkweeds, sagewort, goldenrods, and stinging nettle. The main woody species are poison ivy (trace to 2 percent) and wild grape (trace to 2 percent). Other woody species include winged elm and sumacs. When conditions are favorable, production is about 4,500 pounds per acre. Normal production is about 2,800 pounds per acre. When conditions are unfavorable, production is about 2,000 pounds per acre.

**R112XY045OK, Heavy Bottomland PE 62-80.**—This site is in nearly level to concave areas on flood plains that are occasionally or frequently flooded. The soils are deep, clayey, and have very slow permeability. This site is usually droughty but is often excessively wet during periods of high rainfall. The plant community is composed of 75 to 85 percent grasses, 10 to 20 percent forbs, and 10 to 15 percent woody species. The main grasses are prairie cordgrass (25 to 35 percent), big bluestem (15 to 25 percent), switchgrass (5 to 15 percent), eastern gamagrass (5 to 15 percent), and Indiangrass (5 to 15 percent). Other grasses include Canada wildrye, Florida paspalum, green muhly, tall dropseed, scouringrush, Virginia wildrye, and sedges. The main forbs are giant ragweed (0 to 5 percent), yarrow species (0 to 5 percent), coralberry (0 to 5 percent), American licorice (0 to 5 percent), Illinois bundleflower (0 to 5 percent), Maximilian sunflower (0 to 5 percent), sawtooth sunflower (0 to 5 percent), tall goldenrod (0 to 5 percent), wholeleaf rosinweed (0 to 5 percent), Baldwin's ironweed (0 to 5 percent), dogbane (0 to 5 percent), Pennsylvania smartweed (0 to 5 percent), swamp smartweed (0 to 5 percent), and white crownbeard (0 to 5 percent). The main woody species are pecan (5 to 10 percent), elderberry (0 to 5 percent), black willow (0 to 5 percent), buckbrush (0 to 5 percent), common buttonbush (0 to 5 percent), eastern cottonwood (0 to 5 percent), green ash (0 to 5 percent), amorpha (0 to 5 percent), pin oak (0 to 5 percent), and sycamore (0 to 5 percent). When conditions are favorable, production is about 7,500 to 10,000 pounds per acre. Normal production is about 5,000 to 7,500 pounds per acre. When conditions are unfavorable, production is about 4,000 to 5,000 pounds per acre.

**R112XY050OK, Loamy Bottomland PE 62-80.**—This site is in areas of highly productive, deep, loamy soils on bottomlands. Because of their position and depth, the soils receive and store more moisture than is normal for the area. This site is dominated by tall grasses. Prairie cordgrass is dominant in sloughs, and big bluestem and little bluestem are dominant on second bottoms. The plant community is composed of 70 to 80 percent grasses, 10 to 20 percent forbs, and 5 to 10 percent woody species. The main grasses are big bluestem (10 to 25 percent), switchgrass (10 to 25 percent), prairie cordgrass (10 to 25 percent), Indiangrass (10 to 25 percent), little bluestem (5 to 15 percent), eastern gamagrass (trace to 5 percent), Florida paspalum (trace to 5 percent), broadleaf uniola (trace to 5 percent), tall dropseed (trace to 5 percent), Scribner's panicum (trace to 5 percent), switchcane (trace to 5 percent), sedges (0 to 2 percent), rushes (0 to 2 percent), and wildryes (0 to 2 percent). The main forbs are perennial sunflowers (0 to 2 percent), heath aster (0 to 2 percent), rough coneflower (0 to 2 percent), wholeleaf rosinweed (0 to 2 percent), goldenrods (0 to 2 percent), compassplant (trace to 5 percent), and dotted gayfeather (trace to 5 percent). The main woody species are pecan (5 to 10 percent), walnut (trace to 5 percent), poison ivy (0 to 2 percent), passionvine (0 to 2 percent), winged elm (0 to 2 percent), sumacs (0 to 2 percent), and indigobush (0 to 2 percent). When conditions are favorable, production is about 10,000 pound per acre. When conditions are unfavorable, production is about 6,000 pounds per acre.

**R112XY056OK, Loamy Prairie PE 62-80.**—This site is in areas of nearly level to rolling uplands. Slopes commonly range up to 20 percent but in rare cases can approach 30 percent. The plant community is a tall-grass prairie composed of about

90 percent grasses and grasslike species, 10 percent forbs, and a trace of shrubs. The main grasses are big bluestem (40 percent), little bluestem (20 percent), Indiangrass (10 percent), eastern gamagrass (10 percent), and other grasses (trace to 5 percent). The other grasses include blue grama, buffalograss, Canada wildrye, porcupinegrass, prairie Junegrass, purple lovegrass, Scribner's panicum, sideoats grama, sedges, tall dropseed, and Virginia wildrye. This site has many forbs in trace amounts. The more abundant of these forbs include button snakeroot, catclaw sensitivebriar, compassplant, dotted gayfeather, fringe-leafed ruellia, hairy sunflower, echinacea, pitchersage, purple prairie clover, roundhead lespedeza, plains evening primrose, slender lespedeza, stiff sunflower, tall gayfeather, Virginia tephrosia, and white prairie clover. Many other species of forbs may be present. The main woody species are New Jersey tea and leadplant. Buckbrush, osage orange, and eastern redcedar are common invaders. When conditions are favorable, production ranges from 5,000 to 6,500 pounds per acre. Normal production ranges from about 3,500 to 5,000 pounds per acre. When conditions are unfavorable, production ranges from 2,500 to 3,500 pounds.

**R112XY059OK, Loamy Prairie (Northeast) PE 62-80.**—This site is in areas of nearly level to moderately steep uplands. It is on convex slopes of low ridges and on the side slopes of moderately steep ridges in broad valleys. Slopes range from 0 to 20 percent. Areas of this site are commonly referred to as the “bluestem ranges,” “bluestem meadows,” or “true prairies.” Elevations range from 450 to 700 feet. The plant community is composed of 85 to 95 percent grasses or grasslike species, 5 to 15 percent forbs, 1 to 5 percent legumes, and 1 to 2 percent woody species. The main grasses are big bluestem (5 to 35 percent), little bluestem (5 to 25 percent), Indiangrass (5 to 10 percent), switchgrass (trace to 5 percent), eastern gamagrass (trace to 5 percent), Florida paspalum (trace to 5 percent), tall dropseed (trace to 2 percent), purpletop (trace to 3 percent), jointtail grass (trace to 2 percent), and Scribner's panicum (trace to 1 percent). The main forbs are compassplant (trace to 2 percent), wholeleaf rosinweed (trace to 2 percent), black sampson (trace to 2 percent), dotted gayfeather (trace to 2 percent), ashy sunflower (trace to 2 percent), heath aster (trace to 1 percent), milkweeds (trace to 2 percent), goldenrods (trace to 1 percent), and sagewort (trace to 2 percent). The main legumes are leadplant (trace to 5 percent), sessile-leaved tickclover (trace to 3 percent), Illinois bundleflower (trace to 2 percent), scurfpea (trace to 2 percent), wild indigos (trace to 2 percent), and wild alfalfa (trace to 2 percent). The main woody species are American elm (trace to 5 percent), winged elm (trace to 4 percent), sumac (trace to 5 percent), and indigobush (trace to 5 percent). When conditions are favorable, production is about 7,000 pounds. When conditions are unfavorable, production is about 3,500 pounds per acre.

**R112XY086OK, Shallow Prairie (Eastern) PE 62-80.**—This site is in areas of rocky sandstone and ridges on rather steep slopes developed from sandstone and shale. Some very shallow ledge rock and shale is associated with this site. Elevations range from 700 to 1,200 feet. The main grasses, which make up 75 to 90 percent of the plant community, are big bluestem (10 to 25 percent), switchgrass (10 to 25 percent), little bluestem (5 to 20 percent), Indiangrass (5 to 25 percent), and wildryes (5 to 10 percent). Other grasses include Florida paspalum (trace to 5 percent), dropseed species (trace to 5 percent), sideoats grama (trace to 5 percent), Scribner's panicum (trace to 5 percent), jointtail grass (0 to 2 percent), longspike tridens (0 to 2 percent), purpletop (0 to 2 percent), silver bluestem (0 to 2 percent), and buffalograss (0 to 2 percent). The main legumes are tephrosia (trace to 5 percent), catclaw sensitivebriar (trace to 5 percent), sessile-leaved tickclover (0 to 2 percent), scurfpea (0 to 2 percent), wild indigo (0 to 2 percent), slimflower scurfpea (0 to 2 percent), and leadplant (0 to 2 percent). The main forbs are black sampson (trace to 5 percent), tall gayfeather (trace to 5 percent), perennial sunflowers (0 to 2 percent), heath aster

(0 to 2 percent), milkweeds (0 to 2 percent), goldenrods (0 to 2 percent), and sagewort (0 to 2 percent). Woody species grow in minor amounts. The main woody species are skunkbush (0 to 2 percent), wild grape (0 to 2 percent), poison ivy (0 to 2 percent), coralberry (0 to 2 percent), hackberry (0 to 2 percent), winged elm (0 to 2 percent), and indigobush (0 to 2 percent). These species increase in abundance if the site is abused. When conditions are favorable, production is about 4,200 pounds per acre. When conditions are unfavorable, production is about 2,800 pounds per acre.

**R112XY098OK, Very Shallow (Eastern) PE 62-80.**—This site is in nearly level to gently sloping areas of very shallow soils. The surface layer is typically 6 to 10 inches deep over limestone. Bedrock severely restricts available water capacity and limits root development. Limestone rock fragments are typically on the surface. The plant community is composed of about 75 to 80 percent grasses and grasslike species, 10 percent legumes, 10 percent forbs, and 5 percent woody species. The main grasses are sideoats grama (25 percent), blue grama (15 percent), silver bluestem (10 percent), sedges (5 percent), dropseed species (5 percent), Scribner's panicum (5 percent), hairy grama (5 percent), fall witchgrass (5 percent), and buffalograss (5 percent). Tall grasses are in the deeper pockets of soil and make up about 10 percent of the community. The main tall grasses are big bluestem (trace to 5 percent), little bluestem (trace to 10 percent), Indiangrass (trace to 5 percent), switchgrass (trace to 5 percent), Canada wildrye (trace to 5 percent), and Virginia wildrye (trace to 5 percent). The main legumes are catclaw sensitivebriar (trace to 2 percent), tephrosia (trace to 2 percent), willowleaf sunflower (trace to 2 percent), pitcher sage (trace to 2 percent), deathcamas (trace to 2 percent), greenthread (trace to 2 percent), heath aster (trace to 2 percent), skullcap (trace to 2 percent), milkweeds (trace to 2 percent), dotted gayfeather (trace to 2 percent), sagewort (trace to 2 percent), pricklypear cactus (trace to 2 percent), and goldenrods (trace to 2 percent). The main woody species are poison ivy (0 to 2 percent), Jersey tea (0 to 2 percent), elm (0 to 2 percent), sumacs (0 to 2 percent), blackberry (0 to 2 percent), and coralberry (0 to 2 percent). When conditions are favorable, production is about 2,500 pounds per acre. When conditions are unfavorable, production is about 500 pounds per acre.

**R112XY810OK, Eroded Claypan Prairie PE 62-80.**—This site is in areas of R112XY010OK, Claypan Prairie PE 62-80, that have been subject to erosion. Part or all of the A horizon has been removed from the soil. The soil integrity has been disturbed. Because of the past erosion and the probability of further erosion, the plant community can only be determined by onsite inspection. Production levels are unknown. See R112XY010OK, Claypan Prairie PE 62-80, for a description of the plant community on the parent site.

**R112XY856OK, Eroded Loamy Prairie PE 62-80.**—This site is in areas of R112XY056OK, Loamy Prairie PE 62-80, that have been subject to erosion. Part or all of the A horizon has been removed from the soil. The soil integrity has been disturbed. Because of the past erosion and the probability of further erosion, the plant community can only be determined by onsite inspection. Production levels are unknown. See R112XY056OK, Loamy Prairie PE 62-80, for a description of the plant community on the parent site.

**R116AY050OK, Loamy Bottomland PE 72-80.**—This site is in areas of highly productive, deep, loamy soils on bottomlands. Because of the position and depth of the soils, this site receives and stores more moisture than is normal for the area. This site is dominated by tall grasses. Prairie cordgrass is dominant in sloughs, and big bluestem and little bluestem are dominant on second bottoms. The plant community is composed of 70 to 80 percent grasses, 10 to 20 percent forbs, and 5 to 10 percent woody species. The main grasses are big bluestem (10 to 25 percent), switchgrass (10 to 25 percent), prairie cordgrass (10 to 25 percent), Indiangrass (10 to 25 percent), little bluestem (5 to 15 percent), eastern gamagrass (trace to 5 percent),

Florida paspalum (trace to 5 percent), broadleaf uniola (trace to 5 percent), tall dropseed (trace to 5 percent), Scribner's panicum (trace to 5 percent), switchcane (trace to 5 percent), sedges (0 to 2 percent), rushes (0 to 2 percent), and wildryes (0 to 2 percent). The main forbs are perennial sunflowers (0 to 2 percent), heath aster (0 to 2 percent), rough coneflower (0 to 2 percent), wholeleaf rosinweed (0 to 2 percent), goldenrods (0 to 2 percent), compassplant (trace to 5 percent), and dotted gayfeather (trace to 5 percent). The main woody species are pecan (5 to 10 percent), walnut (trace to 5 percent), poison ivy (0 to 2 percent), passionvine (0 to 2 percent), winged elm (0 to 2 percent), sumacs (0 to 2 percent), and indigobush (0 to 2 percent). When conditions are favorable, production is about 10,000 pound per acre. When conditions are unfavorable, production is about 6,000 pounds per acre.

**R116AY059OK, Loamy Prairie (Northeast) PE 72-80.**—This site is in level to gently rolling areas of deep, moderately permeable soils at elevations ranging from 500 to 700 feet. The main grasses, which make up about 70 to 80 percent of the plant community, are big bluestem, little bluestem, Indiangrass, switchgrass, and prairie dropseed. Jointtail grass, purpletop, tall dropseed, meadow dropseed, and similar species make up about 20 to 30 percent of the plant community. Broomsedge bluestem, splitbeard bluestem, and buffalograss are typically present if the site has deteriorated. They make up trace amounts to 5 percent of the plant community. The main forbs are sumpweed, western ragweed, Baldwin's ironweed, common broomweed, and sumpweed. When conditions are favorable, production ranges from about 5,250 to 7,000 pounds per acre. Normal production ranges from about 3,500 to 5,250 pounds per acre. When conditions are unfavorable, production ranges from about 500 to 3,500 pounds per acre.

**R116AY092OK, Smooth Chert Savannah PE 72-80.**—This site is in nearly level to moderately steep areas in the Ozark Highlands on any slope aspect. Elevations range from 600 to 1,500 feet. The position of the soils on the landscape ranges from ridge crests to footslopes. Available water capacity and fertility are very low due to a high content of chert in the soils. The plant community is composed of 65 to 75 percent grasses and grasslike species, 5 to 10 percent forbs, and 25 to 35 percent trees and shrubs. The main grasses and grasslike species are little bluestem (20 to 30 percent), big bluestem (15 to 25 percent), Indiangrass (10 to 20 percent), switchgrass (10 to 20 percent), uniola (5 to 10 percent), wildryes (5 to 10 percent), and others (35 to 45 percent). The main forbs are black sampson (2 to 5 percent), perennial sunflower (2 to 5 percent), tephrosia (2 to 5 percent), catclaw sensitivebriar (2 to 5 percent), sessile-leaved tickclover (2 to 5 percent), prairie clover (0 to 5 percent), leadplant (0 to 5 percent), and others (0 to 5 percent). The main woody species are post oak (10 to 20 percent), blackjack oak (10 to 20 percent), red oak (5 to 10 percent), white oak (5 to 10 percent), and shortleaf pine (5 to 10). The main shrubs are coralberry (0 to 5 percent), sumacs (0 to 5 percent), Jersey tea (0 to 5 percent), huckleberry (0 to 5 percent), and others (0 to 5 percent). When conditions are favorable, production is about 4,200 pounds per acre. Normal production is about 3,200 pounds per acre. When conditions are unfavorable, production is about 2,250 pounds per acre.

**R117XY075OK, Sandy Savannah PE 72-76.**—This site is in nearly level to steeply sloping areas of sandy soils on uplands. It supports a cover of tall grasses and mid grasses mixed with oak, hickory, and scrub pine. The main grasses, which make up about 50 percent of the plant community, are big bluestem, little bluestem, Indiangrass, and switchgrass. Other grasses include low panicums and carex species. The main cool-season plants are Canada wildrye, Virginia wildrye, and carex species. Woody species make up about 30 percent of the plant community. The main woody species are post oak, blackjack oak, southern red oak, hickory, persimmon, and sassafras. When conditions are favorable, production ranges from about 3,250 to 4,000 pounds per acre. Normal production ranges from about 2,500 to 3,250 pounds



per acre. When conditions are unfavorable, production ranges from about 500 to 2,500 pounds per acre.

**R117XY079OK, Savannah Breaks PE 72-76.**—This site is in steep or very steep (30 percent or greater) areas of rocky soils that have large fragments of sandstone on or near the surface. Most of the rock ledges are in horizontal beds and are commonly on hillsides banded with shale. Large amounts of bare rock on the surface restrict forage production. Surface runoff is rapid. This site supports a cover of tall grasses mixed with low-grade hardwoods and some scrub pine. The plant community includes about 50 to 60 percent big bluestem, Indiangrass, little bluestem, switchgrass, Canada wildrye, Virginia wildrye, low panicum, and carex species. The grasses are in about that order of importance. Forbs make up about 10 percent of the plant community. The main forbs are huckleberry, Jersey tea, asters, and legumes. Trees and shrubs make up about 30 percent of the vegetation. The main woody species are post oak, blackjack oak, southern red oak, hickory, and sassafras. When conditions are favorable, production is about 4,000 pounds per acre. When conditions are unfavorable, production is about 2,500 pounds per acre.

**R117XY088OK, Shallow Savannah PE 72-76.**—This site is in areas of rugged topography on low mountainous ridges, typically oriented in an easterly to westerly direction. The ridges have horizontal beds of sandstone, shale, and some slate. The horizontal bedding of sandstone and the presence of shale limit the available water capacity, thereby reducing the total available moisture to below the average for the area. Slopes range from 6 to 30 percent. Elevations range from 400 to 2,600 feet. The characteristic vegetation consists of scrubby oaks interspersed with tall grasses. The plant community is composed of about 15 to 20 percent oak species and 70 percent big bluestem, little bluestem, and Indiangrass. The main grasses are big bluestem (10 to 30 percent), Indiangrass (10 to 25 percent), little bluestem (5 to 20 percent), dropseed species (5 to 10 percent), jointtail grass (trace to 5 percent), Scribner's panicum (trace to 5 percent), purpletop (0 to 2 percent), sand lovegrass (0 to 2 percent), and purple lovegrass (0 to 2 percent). The main legumes are perennial lespedezas (trace to 5 percent), Illinois bundleflower (trace to 5 percent), sessile-leaved tickclover (0 to 2 percent), scurfpea (0 to 2 percent), wild indigos (0 to 2 percent), and slimflower scurfpea (0 to 2 percent). The main forbs are black sampson (trace to 5 percent), dotted gayfeather (trace to 5 percent), perennial sunflowers (trace to 5 percent), heath aster (0 to 2 percent), and goldenrods (0 to 2 percent). The main woody species are sumacs (trace to 5 percent), post oak (trace to 5 percent), blackjack oak (trace to 5 percent), plum (0 to 2 percent), poison ivy (0 to 2 percent), cedar elm (0 to 2 percent), and winged elm (0 to 2 percent). Shortleaf pine is on the dry edge of the humid zone. When conditions are favorable, production is about 3,500. When conditions are unfavorable, production is about 1,750 pounds per acre.



# Formation and Classification of the Soils

---

This section summarizes the major factors of soil formation and describes the system of soil classification. The classification of each soil in the survey area is shown in the table “Classification of the Soils” at the end of this section. The Official Soil Series Descriptions, including the range of important characteristics of the soils, for the series in this survey area are online at <http://soils.usda.gov/technical/classification/osd/>. Characteristics of the soil and the material in which it formed are identified for each soil series. A pedon, a small three-dimensional area of soil, which is typical of the series is described. The detailed description of each soil horizon follows standards in the “Soil Survey Manual” (Soil Survey Division Staff, 1993). Many of the technical terms used in the descriptions are defined in “Soil Taxonomy” (Soil Survey Staff, 1999).

## Formation of the Soils

Soil is produced by the action of soil-forming processes on materials deposited or accumulated by geologic agencies. The characteristics of the soil at any given point are determined by the physical and mineralogical composition of the parent materials; the climate under which the soil material has accumulated and existed since accumulation; the plant and animal life on and in the soil; the relief, or lay of the land; and the length of time the forces of soil development have acted on the soil material.

Climate and vegetation are the active factors of soil formation. They act on parent material that has accumulated through the weathering of rocks and slowly change it into a natural body that has genetically related horizons. The effects of climate and vegetation are conditioned by relief. Parent material also affects the kind of profile that can be formed and, in extreme cases, determines it almost entirely. Finally, time is needed for the changing of the parent material into a soil profile. The time may be long or short, but some time is always required for differentiation of horizons. Generally, a long time is required for the development of distinct horizons.

The factors of soil formation are so closely interrelated in their effects that few generalizations can be made regarding the effects of any one unless conditions are specified for the other four.

## Classification of the Soils

The system of soil classification used by the National Cooperative Soil Survey has six categories (Soil Survey Staff, 1999 and 2003). Beginning with the broadest, these categories are the order, suborder, great group, subgroup, family, and series. Classification is based on soil properties observed in the field or inferred from those observations or from laboratory measurements. The categories are described in the following paragraphs.

**ORDER.** Twelve soil orders are recognized. The differences among orders reflect the dominant soil-forming processes and the degree of soil formation. Each order is identified by a word ending in *sol*. An example is Mollisol.

**SUBORDER.** Each order is divided into suborders primarily on the basis of properties that influence soil genesis and are important to plant growth or properties that reflect the most important variables within the orders. The last syllable in the name of a suborder indicates the order. An example is Ustoll (*Ust*, meaning dry, plus *oll*, from Mollisol).

**GREAT GROUP.** Each suborder is divided into great groups on the basis of close similarities in kind, arrangement, and degree of development of pedogenic horizons; soil moisture and temperature regimes; and base status. Each great group is identified by the name of a suborder and by a prefix that indicates a property of the soil. An example is Argiustolls (*Argi*, meaning argillic horizonation, plus *ustoll*, the suborder of the Mollisols that has an ustic moisture regime).

**SUBGROUP.** Each great group has a typic subgroup. Other subgroups are intergrades or extragrades. The typic is the central concept of the great group; it is not necessarily the most extensive. Intergrades are transitions to other orders, suborders, or great groups. Extragrades have some properties that are not representative of the great group but do not indicate transitions to any other known kind of soil. Each subgroup is identified by one or more adjectives preceding the name of the great group. The adjective *Typic* identifies the subgroup that typifies the great group. An example is Typic Argiustolls.

**FAMILY.** Families are established within a subgroup on the basis of physical and chemical properties and other characteristics that affect management. Generally, the properties are those of horizons below plow depth where there is much biological activity. Among the properties and characteristics considered are particle-size class, mineral content, temperature regime, thickness of the root zone, consistence, moisture equivalent, slope, and permanent cracks. A family name consists of the name of a subgroup preceded by terms that indicate soil properties. An example is fine-loamy, mixed, superactive, thermic Typic Argiustolls.

**SERIES.** The series consists of soils that have similar horizons in their profile. The horizons are similar in color, texture, structure, reaction, consistence, mineral and chemical composition, and arrangement in the profile. The texture of the surface layer or of the substratum can differ within a series.

Supplement to the Soil Survey of Cherokee County, Oklahoma

Table 5.--Classification of the Soils

Soil name	Family or higher taxonomic class
Apperson-----	Fine, smectitic, thermic Aquic Hapluderts
Barge-----	Fine-silty, mixed, active, nonacid, thermic Mollic Udarents
Britwater-----	Fine-loamy, mixed, active, mesic Typic Paleudalfs
Captina-----	Fine-silty, siliceous, active, mesic Typic Fragiudults
Clarksville-----	Loamy-skeletal, siliceous, semiactive, mesic Typic Paleudults
Coweta-----	Loamy, siliceous, superactive, thermic, shallow Typic Hapludolls
Cupco-----	Fine-silty, siliceous, active, thermic Typic Epiaqualfs
Dennis-----	Fine, mixed, active, thermic Aquic Argiudolls
Eldorado-----	Loamy-skeletal, mixed, active, thermic Typic Paleudolls
Elsah-----	Loamy-skeletal, mixed, superactive, nonacid, mesic Typic Udifluvents
Healing-----	Fine-silty, mixed, superactive, mesic Typic Argiudolls
Hector-----	Loamy, siliceous, subactive, thermic Lithic Dystrudepts
Jay-----	Fine-silty, mixed, active, thermic Oxyaquic Fragiudalfs
Lightning-----	Fine, mixed, active, thermic Chromic Vertic Epiaqualfs
Linker-----	Fine-loamy, siliceous, semiactive, thermic Typic Hapludults
Newtonia-----	Fine-silty, mixed, superactive, thermic Typic Paleudolls
Nixa-----	Loamy-skeletal, siliceous, active, mesic Glossic Fragiudults
Okemah-----	Fine, mixed, active, thermic Aquic Paleudolls
Osage-----	Fine, smectitic, thermic Typic Epiaquerts
Razort-----	Fine-loamy, mixed, active, mesic Mollic Hapludalfs
Roebuck-----	Fine, smectitic, thermic Aeric Epiaquerts
Shidler-----	Loamy, mixed, active, thermic Lithic Haplustolls
Stigler-----	Fine, mixed, active, thermic Aquic Paleudalfs
Taloka-----	Fine, mixed, active, thermic Mollic Albaqualfs
Tonti-----	Fine-loamy, mixed, active, mesic Typic Fragiudults
Verdigris-----	Fine-silty, mixed, superactive, thermic Cumulic Hapludolls



# References

---

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. <http://soils.usda.gov/technical/>

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service. U.S. Department of Agriculture Handbook 436. <http://soils.usda.gov/technical/>

Soil Survey Staff. 2003. Keys to soil taxonomy. 9th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. <http://soils.usda.gov/technical/>

United States Department of Agriculture. 1970. Soil survey of Cherokee and Delaware Counties, Oklahoma.





# Glossary

---

Many of the terms relating to landforms, geology, and geomorphology are defined in more detail in the "National Soil Survey Handbook" (available at local offices of the Natural Resources Conservation Service or on the Internet).

**ABC soil.** A soil having an A, a B, and a C horizon.

**AC soil.** A soil having only an A and a C horizon. Commonly, such soil formed in recent alluvium or on steep, rocky slopes.

**Aeration, soil.** The exchange of air in soil with air from the atmosphere. The air in a well aerated soil is similar to that in the atmosphere; the air in a poorly aerated soil is considerably higher in carbon dioxide and lower in oxygen.

**Aggregate, soil.** Many fine particles held in a single mass or cluster. Natural soil aggregates, such as granules, blocks, or prisms, are called peds. Clods are aggregates produced by tillage or logging.

**Alkali (sodic) soil.** A soil having so high a degree of alkalinity (pH 8.5 or higher) or so high a percentage of exchangeable sodium (15 percent or more of the total exchangeable bases), or both, that plant growth is restricted.

**Alluvial fan.** A low, outspread mass of loose materials and/or rock material, commonly with gentle slopes. It is shaped like an open fan or a segment of a cone. The material was deposited by a stream at the place where it issues from a narrow mountain valley or upland valley or where a tributary stream is near or at its junction with the main stream. The fan is steepest near its apex, which points upstream, and slopes gently and convexly outward (downstream) with a gradual decrease in gradient.

**Alluvium.** Unconsolidated material, such as gravel, sand, silt, clay, and various mixtures of these, deposited on land by running water.

**Alpha,alpha-dipyridyl.** A compound that when dissolved in ammonium acetate is used to detect the presence of reduced iron (Fe II) in the soil. A positive reaction implies reducing conditions and the likely presence of redoximorphic features.

**Animal unit month (AUM).** The amount of forage required by one mature cow of approximately 1,000 pounds weight, with or without a calf, for 1 month.

**Aquic conditions.** Current soil wetness characterized by saturation, reduction, and redoximorphic features.

**Argillic horizon.** A subsoil horizon characterized by an accumulation of illuvial clay.

**Aspect.** The direction toward which a slope faces. Also called slope aspect.

**Association, soil.** A group of soils or miscellaneous areas geographically associated in a characteristic repeating pattern and defined and delineated as a single map unit.

**Available water capacity (available moisture capacity).** The capacity of soils to hold water available for use by most plants. It is commonly defined as the difference between the amount of soil water at field moisture capacity and the amount at wilting point. It is commonly expressed as inches of water per inch of soil. The capacity, in inches, in a 60-inch profile or to a limiting layer is expressed as:

Supplement to the Soil Survey of Cherokee County, Oklahoma

Very low .....	0 to 3
Low .....	3 to 6
Moderate .....	6 to 9
High .....	9 to 12
Very high .....	more than 12

- Backslope.** The position that forms the steepest and generally linear, middle portion of a hillslope. In profile, backslopes are commonly bounded by a convex shoulder above and a concave footslope below.
- Backswamp.** A flood-plain landform. Extensive, marshy or swampy, depressed areas of flood plains between natural levees and valley sides or terraces.
- Badland.** A landscape that is intricately dissected and characterized by a very fine drainage network with high drainage densities and short, steep slopes and narrow interfluves. Badlands develop on surfaces that have little or no vegetative cover overlying unconsolidated or poorly cemented materials (clays, silts, or sandstones) with, in some cases, soluble minerals, such as gypsum or halite.
- Basal area.** The area of a cross section of a tree, generally referring to the section at breast height and measured outside the bark. It is a measure of stand density, commonly expressed in square feet.
- Base saturation.** The degree to which material having cation-exchange properties is saturated with exchangeable bases (sum of Ca, Mg, Na, and K), expressed as a percentage of the total cation-exchange capacity.
- Base slope** (geomorphology). A geomorphic component of hills consisting of the concave to linear (perpendicular to the contour) slope that, regardless of the lateral shape, forms an apron or wedge at the bottom of a hillside dominated by colluvium and slope-wash sediments (for example, slope alluvium).
- Bedding plane.** A planar or nearly planar bedding surface that visibly separates each successive layer of stratified sediment or rock (of the same or different lithology) from the preceding or following layer; a plane of deposition. It commonly marks a change in the circumstances of deposition and may show a parting, a color difference, a change in particle size, or various combinations of these. The term is commonly applied to any bedding surface, even one that is conspicuously bent or deformed by folding.
- Bedrock.** The solid rock that underlies the soil and other unconsolidated material or that is exposed at the surface.
- Bedrock-controlled topography.** A landscape where the configuration and relief of the landforms are determined or strongly influenced by the underlying bedrock.
- Bisequum.** Two sequences of soil horizons, each of which consists of an illuvial horizon and the overlying eluvial horizons.
- Blowout.** A saucer-, cup-, or trough-shaped depression formed by wind erosion on a preexisting dune or other sand deposit, especially in an area of shifting sand or loose soil or where protective vegetation is disturbed or destroyed; the adjoining accumulation of sand derived from the depression, where recognizable, is commonly included. Blowouts are commonly small.
- Bottom land.** An informal term loosely applied to various portions of a flood plain.
- Boulders.** Rock fragments larger than 2 feet (60 centimeters) in diameter.
- Breaks.** A landscape or tract of steep, rough or broken land dissected by ravines and gullies and marking a sudden change in topography.
- Breast height.** An average height of 4.5 feet above the ground surface; the point on a tree where diameter measurements are ordinarily taken.
- Brush management.** Use of mechanical, chemical, or biological methods to make conditions favorable for reseeding or to reduce or eliminate competition from woody vegetation and thus allow understory grasses and forbs to recover. Brush management increases forage production and thus reduces the hazard of erosion. It can improve the habitat for some species of wildlife.

- Butte.** An isolated, generally flat-topped hill or mountain with relatively steep slopes and talus or precipitous cliffs and characterized by summit width that is less than the height of bounding escarpments; commonly topped by a caprock of resistant material and representing an erosion remnant carved from flat-lying rocks.
- Calcareous soil.** A soil containing enough calcium carbonate (commonly combined with magnesium carbonate) to effervesce visibly when treated with cold, dilute hydrochloric acid.
- Caliche.** A general term for a prominent zone of secondary carbonate accumulation in surficial materials in warm, subhumid to arid areas. Caliche is formed by both geologic and pedologic processes. Finely crystalline calcium carbonate forms a nearly continuous surface-coating and void-filling medium in geologic (parent) materials. Cementation ranges from weak in nonindurated forms to very strong in indurated forms. Other minerals (e.g., carbonates, silicate, and sulfate) may occur as accessory cements. Most petrocalcic horizons and some calcic horizons are caliche.
- Canopy.** The leafy crown of trees or shrubs. (See Crown.)
- Canyon.** A long, deep, narrow valley with high, precipitous walls in an area of high local relief.
- Capillary water.** Water held as a film around soil particles and in tiny spaces between particles. Surface tension is the adhesive force that holds capillary water in the soil.
- Catena.** A sequence, or “chain,” of soils on a landscape that formed in similar kinds of parent material and under similar climatic conditions but that have different characteristics as a result of differences in relief and drainage.
- Cation.** An ion carrying a positive charge of electricity. The common soil cations are calcium, potassium, magnesium, sodium, and hydrogen.
- Cation-exchange capacity.** The total amount of exchangeable cations that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. The term, as applied to soils, is synonymous with base-exchange capacity but is more precise in meaning.
- Catsteps.** See Terracettes.
- Channery soil material.** Soil material that has, by volume, 15 to 35 percent thin, flat fragments of sandstone, shale, slate, limestone, or schist as much as 6 inches (15 centimeters) along the longest axis. A single piece is called a chanter.
- Chemical treatment.** Control of unwanted vegetation through the use of chemicals.
- Chiseling.** Tillage with an implement having one or more soil-penetrating points that shatter or loosen hard, compacted layers to a depth below normal plow depth.
- Clay.** As a soil separate, the mineral soil particles less than 0.002 millimeter in diameter. As a soil textural class, soil material that is 40 percent or more clay, less than 45 percent sand, and less than 40 percent silt.
- Clay depletions.** See Redoximorphic features.
- Clay film.** A thin coating of oriented clay on the surface of a soil aggregate or lining pores or root channels. Synonyms: clay coating, clay skin.
- Clay beds.** Old, buried, alluvial clay sediments.
- Claypan.** A dense, compact, slowly permeable subsoil layer that contains much more clay than the overlying materials, from which it is separated by a sharply defined boundary. A claypan is commonly hard when dry and plastic and sticky when wet.
- Climax plant community.** The stabilized plant community on a particular site. The plant cover reproduces itself and does not change so long as the environment remains the same.
- Coarse fragments.** Mineral or rock particles larger than 2 millimeters in diameter.
- Coarse textured soil.** Sand or loamy sand.
- Cobble (or cobblestone).** A rounded or partly rounded fragment of rock 3 to 10 inches (7.6 to 25 centimeters) in diameter.

- Cobbly soil material.** Material that has 15 to 35 percent, by volume, rounded or partially rounded rock fragments 3 to 10 inches (7.6 to 25 centimeters) in diameter. Very cobbly soil material has 35 to 60 percent of these rock fragments, and extremely cobbly soil material has more than 60 percent.
- COLE (coefficient of linear extensibility).** See Linear extensibility.
- Colluvium.** Unconsolidated, unsorted earth material being transported or deposited on side slopes and/or at the base of slopes by mass movement (e.g., direct gravitational action) and by local, unconcentrated runoff.
- Complex slope.** Irregular or variable slope. Planning or establishing terraces, diversions, and other water-control structures on a complex slope is difficult.
- Complex, soil.** A map unit of two or more kinds of soil or miscellaneous areas in such an intricate pattern or so small in area that it is not practical to map them separately at the selected scale of mapping. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas.
- Concretions.** See Redoximorphic features.
- Conglomerate.** A coarse grained, clastic sedimentary rock composed of rounded or subangular rock fragments more than 2 millimeters in diameter. It commonly has a matrix of sand and finer textured material. Conglomerate is the consolidated equivalent of gravel.
- Conservation cropping system.** Growing crops in combination with needed cultural and management practices. In a good conservation cropping system, the soil-improving crops and practices more than offset the effects of the soil-depleting crops and practices. Cropping systems are needed on all tilled soils. Soil-improving practices in a conservation cropping system include the use of rotations that contain grasses and legumes and the return of crop residue to the soil. Other practices include the use of green manure crops of grasses and legumes, proper tillage, adequate fertilization, and weed and pest control.
- Conservation tillage.** A tillage system that does not invert the soil and that leaves a protective amount of crop residue on the surface throughout the year.
- Consistence, soil.** Refers to the degree of cohesion and adhesion of soil material and its resistance to deformation when ruptured. Consistence includes resistance of soil material to rupture and to penetration; plasticity, toughness, and stickiness of puddled soil material; and the manner in which the soil material behaves when subject to compression. Terms describing consistence are defined in the "Soil Survey Manual."
- Contour stripcropping.** Growing crops in strips that follow the contour. Strips of grass or close-growing crops are alternated with strips of clean-tilled crops or summer fallow.
- Control section.** The part of the soil on which classification is based. The thickness varies among different kinds of soil, but for many it is that part of the soil profile between depths of 10 inches and 40 or 80 inches.
- Coprogenous earth (sedimentary peat).** A type of limnic layer composed predominantly of fecal material derived from aquatic animals.
- Corrosion (geomorphology).** A process of erosion whereby rocks and soil are removed or worn away by natural chemical processes, especially by the solvent action of running water, but also by other reactions, such as hydrolysis, hydration, carbonation, and oxidation.
- Corrosion (soil survey interpretations).** Soil-induced electrochemical or chemical action that dissolves or weakens concrete or uncoated steel.
- Cover crop.** A close-growing crop grown primarily to improve and protect the soil between periods of regular crop production, or a crop grown between trees and vines in orchards and vineyards.
- Crop residue management.** Returning crop residue to the soil, which helps to maintain soil structure, organic matter content, and fertility and helps to control erosion.

- Cropping system.** Growing crops according to a planned system of rotation and management practices.
- Cross-slope farming.** Deliberately conducting farming operations on sloping farmland in such a way that tillage is across the general slope.
- Crown.** The upper part of a tree or shrub, including the living branches and their foliage.
- Cuesta.** An asymmetric ridge capped by resistant rock layers of slight or moderate dip (commonly less than 15 percent slopes); a type of homocline produced by differential erosion of interbedded resistant and weak rocks. A cuesta has a long, gentle slope on one side (dip slope) that roughly parallels the inclined beds; on the other side, it has a relatively short and steep or clifflike slope (scarp) that cuts through the tilted rocks.
- Decreasers.** The most heavily grazed climax range plants. Because they are the most palatable, they are the first to be destroyed by overgrazing.
- Deferred grazing.** Postponing grazing or resting grazing land for a prescribed period.
- Deflocculation.** Dispersion, or breaking up, of soil aggregates into individual particles. Sodium salts, for example, deflocculate, or disperse, granulated particles of clay to form a clay that runs together, or puddles.
- Depth, soil.** Generally, the thickness of the soil over bedrock. Very deep soils are more than 60 inches deep over bedrock; deep soils, 40 to 60 inches; moderately deep, 20 to 40 inches; shallow, 10 to 20 inches; and very shallow, less than 10 inches.
- Diatomaceous earth.** A geologic deposit of fine, grayish siliceous material composed chiefly or entirely of the remains of diatoms.
- Dip slope.** A slope of the land surface, roughly determined by and approximately conforming to the dip of the underlying bedrock.
- Diversion (or diversion terrace).** A ridge of earth, generally a terrace, built to protect downslope areas by diverting runoff from its natural course.
- Drainage class (natural).** Refers to the frequency and duration of wet periods under conditions similar to those under which the soil formed. Alterations of the water regime by human activities, either through drainage or irrigation, are not a consideration unless they have significantly changed the morphology of the soil. Seven classes of natural soil drainage are recognized—*excessively drained, somewhat excessively drained, well drained, moderately well drained, somewhat poorly drained, poorly drained, and very poorly drained*. These classes are defined in the “Soil Survey Manual.”
- Drainage, surface.** Runoff, or surface flow of water, from an area.
- Drainageway.** A general term for a course or channel along which water moves in draining an area. A term restricted to relatively small, linear depressions that at some time move concentrated water and either do not have a defined channel or have only a small defined channel.
- Draw.** A small stream valley that generally is shallower and more open than a ravine or gulch and that has a broader bottom. The present stream channel may appear inadequate to have cut the drainageway that it occupies.
- Duff.** A generally firm organic layer on the surface of mineral soils. It consists of fallen plant material that is in the process of decomposition and includes everything from the litter on the surface to underlying pure humus.
- Dune.** A low mound, ridge, bank, or hill of loose, windblown granular material (generally sand), either barren and capable of movement from place to place or covered and stabilized with vegetation but retaining its characteristic shape.
- Earthy fill.** See Mine spoil.
- Ecological site.** An area where climate, soil, and relief are sufficiently uniform to produce a distinct natural plant community. An ecological site is the product of all the environmental factors responsible for its development. It is typified by an

association of species that differ from those on other ecological sites in kind and/or proportion of species or in total production.

- Eluviation.** The movement of material in true solution or colloidal suspension from one place to another within the soil. Soil horizons that have lost material through eluviation are eluvial; those that have received material are illuvial.
- Endosaturation.** A type of saturation of the soil in which all horizons between the upper boundary of saturation and a depth of 2 meters are saturated.
- Eolian deposit.** Sand-, silt-, or clay-sized clastic material transported and deposited primarily by wind, commonly in the form of a dune or a sheet of sand or loess.
- Ephemeral stream.** A stream, or reach of a stream, that flows only in direct response to precipitation. It receives no long-continued supply from melting snow or other source, and its channel is above the water table at all times.
- Episaturation.** A type of saturation indicating a perched water table in a soil in which saturated layers are underlain by one or more unsaturated layers within 2 meters of the surface.
- Erosion.** The wearing away of the land surface by water, wind, ice, or other geologic agents and by such processes as gravitational creep.
- Erosion (geologic).* Erosion caused by geologic processes acting over long geologic periods and resulting in the wearing away of mountains and the building up of such landscape features as flood plains and coastal plains.  
Synonym: natural erosion.
- Erosion (accelerated).* Erosion much more rapid than geologic erosion, mainly as a result of human or animal activities or of a catastrophe in nature, such as a fire, that exposes the surface.
- Erosion surface.** A land surface shaped by the action of erosion, especially by running water.
- Escarpment.** A relatively continuous and steep slope or cliff breaking the general continuity of more gently sloping land surfaces and resulting from erosion or faulting. Most commonly applied to cliffs produced by differential erosion.  
Synonym: scarp.
- Fallow.** Cropland left idle in order to restore productivity through accumulation of moisture. Summer fallow is common in regions of limited rainfall where cereal grain is grown. The soil is tilled for at least one growing season for weed control and decomposition of plant residue.
- Fan remnant.** A general term for landforms that are the remaining parts of older fan landforms, such as alluvial fans, that have been either dissected or partially buried.
- Fertility, soil.** The quality that enables a soil to provide plant nutrients, in adequate amounts and in proper balance, for the growth of specified plants when light, moisture, temperature, tilth, and other growth factors are favorable.
- Field moisture capacity.** The moisture content of a soil, expressed as a percentage of the oven-dry weight, after the gravitational, or free, water has drained away; the field moisture content 2 or 3 days after a soaking rain; also called *normal field capacity*, *normal moisture capacity*, or *capillary capacity*.
- Fill slope.** A sloping surface consisting of excavated soil material from a road cut. It commonly is on the downhill side of the road.
- Fine textured soil.** Sandy clay, silty clay, or clay.
- Firebreak.** An area cleared of flammable material to stop or help control creeping or running fires. It also serves as a line from which to work and to facilitate the movement of firefighters and equipment. Designated roads also serve as firebreaks.
- First bottom.** An obsolete, informal term loosely applied to the lowest flood-plain steps that are subject to regular flooding.

- Flaggy soil material.** Material that has, by volume, 15 to 35 percent flagstones. Very flaggy soil material has 35 to 60 percent flagstones, and extremely flaggy soil material has more than 60 percent flagstones.
- Flagstone.** A thin fragment of sandstone, limestone, slate, shale, or (rarely) schist 6 to 15 inches (15 to 38 centimeters) long.
- Flood plain.** The nearly level plain that borders a stream and is subject to flooding unless protected artificially.
- Flood-plain landforms.** A variety of constructional and erosional features produced by stream channel migration and flooding. Examples include backswamps, flood-plain splays, meanders, meander belts, meander scrolls, oxbow lakes, and natural levees.
- Flood-plain splay.** A fan-shaped deposit or other outspread deposit formed where an overloaded stream breaks through a levee (natural or artificial) and deposits its material (commonly coarse grained) on the flood plain.
- Flood-plain step.** An essentially flat, terrace-like alluvial surface within a valley that is frequently covered by floodwater from the present stream; any approximately horizontal surface still actively modified by fluvial scour and/or deposition. May occur individually or as a series of steps.
- Fluvial.** Of or pertaining to rivers or streams; produced by stream or river action.
- Foothills.** A region of steeply sloping hills that fringes a mountain range or high-plateau escarpment. The hills have relief of as much as 1,000 feet (300 meters).
- Footslope.** The concave surface at the base of a hillslope. A footslope is a transition zone between upslope sites of erosion and transport (shoulders and backslopes) and downslope sites of deposition (toeslopes).
- Forb.** Any herbaceous plant not a grass or a sedge.
- Forest cover.** All trees and other woody plants (underbrush) covering the ground in a forest.
- Forest type.** A stand of trees similar in composition and development because of given physical and biological factors by which it may be differentiated from other stands.
- Fragipan.** A loamy, brittle subsurface horizon low in porosity and content of organic matter and low or moderate in clay but high in silt or very fine sand. A fragipan appears cemented and restricts roots. When dry, it is hard or very hard and has a higher bulk density than the horizon or horizons above. When moist, it tends to rupture suddenly under pressure rather than to deform slowly.
- Genesis, soil.** The mode of origin of the soil. Refers especially to the processes or soil-forming factors responsible for the formation of the solum, or true soil, from the unconsolidated parent material.
- Gilgai.** Commonly, a succession of microbasins and microknolls in nearly level areas or of microvalleys and microridges parallel with the slope. Typically, the microrelief of clayey soils that shrink and swell considerably with changes in moisture content.
- Gleyed soil.** Soil that formed under poor drainage, resulting in the reduction of iron and other elements in the profile and in gray colors.
- Grassed waterway.** A natural or constructed waterway, typically broad and shallow, seeded to grass as protection against erosion. Conducts surface water away from cropland.
- Gravel.** Rounded or angular fragments of rock as much as 3 inches (7.6 centimeters) in diameter. An individual piece is a pebble.
- Gravelly soil material.** Material that has 15 to 35 percent, by volume, rounded or angular rock fragments, not prominently flattened, as much as 3 inches (7.6 centimeters) in diameter.
- Green manure crop (agronomy).** A soil-improving crop grown to be plowed under in an early stage of maturity or soon after maturity.

- Ground water.** Water filling all the unblocked pores of the material below the water table.
- Gully.** A small channel with steep sides caused by erosion and cut in unconsolidated materials by concentrated but intermittent flow of water. The distinction between a gully and a rill is one of depth. A gully generally is an obstacle to farm machinery and is too deep to be obliterated by ordinary tillage; a rill is of lesser depth and can be smoothed over by ordinary tillage.
- Hard bedrock.** Bedrock that cannot be excavated except by blasting or by the use of special equipment that is not commonly used in construction.
- Hardpan.** A hardened or cemented soil horizon, or layer. The soil material is sandy, loamy, or clayey and is cemented by iron oxide, silica, calcium carbonate, or other substance.
- Head slope** (geomorphology). A geomorphic component of hills consisting of a laterally concave area of a hillside, especially at the head of a drainageway. The overland waterflow is converging.
- High-residue crops.** Such crops as small grain and corn used for grain. If properly managed, residue from these crops can be used to control erosion until the next crop in the rotation is established. These crops return large amounts of organic matter to the soil.
- Hill.** A generic term for an elevated area of the land surface, rising as much as 1,000 feet above surrounding lowlands, commonly of limited summit area and having a well defined outline. Slopes are generally more than 15 percent. The distinction between a hill and a mountain is arbitrary and may depend on local usage.
- Hillslope.** A generic term for the steeper part of a hill between its summit and the drainage line, valley flat, or depression floor at the base of a hill.
- Horizon, soil.** A layer of soil, approximately parallel to the surface, having distinct characteristics produced by soil-forming processes. In the identification of soil horizons, an uppercase letter represents the major horizons. Numbers or lowercase letters that follow represent subdivisions of the major horizons. An explanation of the subdivisions is given in the "Soil Survey Manual." The major horizons of mineral soil are as follows:
- O horizon.*—An organic layer of fresh and decaying plant residue.
- A horizon.*—The mineral horizon at or near the surface in which an accumulation of humified organic matter is mixed with the mineral material. Also, a plowed surface horizon, most of which was originally part of a B horizon.
- E horizon.*—The mineral horizon in which the main feature is loss of silicate clay, iron, aluminum, or some combination of these.
- B horizon.*—The mineral horizon below an A horizon. The B horizon is in part a layer of transition from the overlying A to the underlying C horizon. The B horizon also has distinctive characteristics, such as (1) accumulation of clay, sesquioxides, humus, or a combination of these; (2) prismatic or blocky structure; (3) redder or browner colors than those in the A horizon; or (4) a combination of these.
- C horizon.*—The mineral horizon or layer, excluding indurated bedrock, that is little affected by soil-forming processes and does not have the properties typical of the overlying soil material. The material of a C horizon may be either like or unlike that in which the solum formed. If the material is known to differ from that in the solum, an Arabic numeral, commonly a 2, precedes the letter C.
- Cr horizon.*—Soft, consolidated bedrock beneath the soil.
- R layer.*—Consolidated bedrock beneath the soil. The bedrock commonly underlies a C horizon, but it can be directly below an A or a B horizon.
- Humus.** The well decomposed, more or less stable part of the organic matter in mineral soils.



- Hydrologic soil groups.** Refers to soils grouped according to their runoff potential. The soil properties that influence this potential are those that affect the minimum rate of water infiltration on a bare soil during periods after prolonged wetting when the soil is not frozen. These properties are depth to a seasonal high water table, the infiltration rate and permeability after prolonged wetting, and depth to a very slowly permeable layer. The slope and the kind of plant cover are not considered but are separate factors in predicting runoff.
- Igneous rock.** Rock that was formed by cooling and solidification of magma and that has not been changed appreciably by weathering since its formation. Major varieties include plutonic and volcanic rock (e.g., andesite, basalt, and granite).
- Illuviation.** The movement of soil material from one horizon to another in the soil profile. Generally, material is removed from an upper horizon and deposited in a lower horizon.
- Impervious soil.** A soil through which water, air, or roots penetrate slowly or not at all. No soil is absolutely impervious to air and water all the time.
- Increasers.** Species in the climax vegetation that increase in amount as the more desirable plants are reduced by close grazing. Increasers commonly are the shorter plants and the less palatable to livestock.
- Infiltration.** The downward entry of water into the immediate surface of soil or other material, as contrasted with percolation, which is movement of water through soil layers or material.
- Infiltration capacity.** The maximum rate at which water can infiltrate into a soil under a given set of conditions.
- Infiltration rate.** The rate at which water penetrates the surface of the soil at any given instant, usually expressed in inches per hour. The rate can be limited by the infiltration capacity of the soil or the rate at which water is applied at the surface.
- Intake rate.** The average rate of water entering the soil under irrigation. Most soils have a fast initial rate; the rate decreases with application time. Therefore, intake rate for design purposes is not a constant but is a variable depending on the net irrigation application. The rate of water intake, in inches per hour, is expressed as follows:
- |                     |                 |
|---------------------|-----------------|
| Less than 0.2 ..... | very low        |
| 0.2 to 0.4 .....    | low             |
| 0.4 to 0.75 .....   | moderately low  |
| 0.75 to 1.25 .....  | moderate        |
| 1.25 to 1.75 .....  | moderately high |
| 1.75 to 2.5 .....   | high            |
| More than 2.5 ..... | very high       |
- Interfluve.** A landform composed of the relatively undissected upland or ridge between two adjacent valleys containing streams flowing in the same general direction. An elevated area between two drainageways that sheds water to those drainageways.
- Interfluve** (geomorphology). A geomorphic component of hills consisting of the uppermost, comparatively level or gently sloping area of a hill; shoulders of backwearing hillslopes can narrow the upland or can merge, resulting in a strongly convex shape.
- Intermittent stream.** A stream, or reach of a stream, that does not flow year-round but that is commonly dry for 3 or more months out of 12 and whose channel is generally below the local water table. It flows only during wet periods or when it receives ground-water discharge or long, continued contributions from melting snow or other surface and shallow subsurface sources.
- Internal soil drainage.** The downward movement of water through the soil profile. The rate of movement is determined by the texture, structure, and other

characteristics of the soil profile and underlying layers and by the height of the water table, either permanent or perched. Relative terms for expressing internal drainage are *none*, *very slow*, *slow*, *medium*, *rapid*, and *very rapid*.

**Invaders.** On range, plants that encroach into an area and grow after the climax vegetation has been reduced by grazing. Generally, plants invade following disturbance of the surface.

**Iron depletions.** See Redoximorphic features.

**Irrigation.** Application of water to soils to assist in production of crops. Methods of irrigation are:

*Basin.*—Water is applied rapidly to nearly level plains surrounded by levees or dikes.

*Border.*—Water is applied at the upper end of a strip in which the lateral flow of water is controlled by small earth ridges called border dikes, or borders.

*Controlled flooding.*—Water is released at intervals from closely spaced field ditches and distributed uniformly over the field.

*Drip (or trickle).*—Water is applied slowly and under low pressure to the surface of the soil or into the soil through such applicators as emitters, porous tubing, or perforated pipe.

*Furrow.*—Water is applied in small ditches made by cultivation implements.

Furrows are used for tree and row crops.

*Sprinkler.*—Water is sprayed over the soil surface through pipes or nozzles from a pressure system.

*Subirrigation.*—Water is applied in open ditches or tile lines until the water table is raised enough to wet the soil.

**Karst** (topography). A kind of topography that formed in limestone, gypsum, or other soluble rocks by dissolution and that is characterized by closed depressions, sinkholes, caves, and underground drainage.

**Knoll.** A small, low, rounded hill rising above adjacent landforms.

**$K_{sat}$ .** Saturated hydraulic conductivity. (See Permeability.)

**Lacustrine deposit.** Material deposited in lake water and exposed when the water level is lowered or the elevation of the land is raised.

**Lake plain.** A nearly level surface marking the floor of an extinct lake filled by well sorted, generally fine textured, stratified deposits, commonly containing varves.

**Lamellae.** Thin bands of fibers of translocated clay that constitute illuvial, and in many cases argillic, horizons in sandy upland soils.

**Landslide.** A general, encompassing term for most types of mass movement landforms and processes involving the downslope transport and outward deposition of soil and rock materials caused by gravitational forces; the movement may or may not involve saturated materials. The speed and distance of movement, as well as the amount of soil and rock material, vary greatly.

**Leaching.** The removal of soluble material from soil or other material by percolating water.

**Linear extensibility.** Refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. Linear extensibility is used to determine the shrink-swell potential of soils. It is an expression of the volume change between the water content of the clod at  $1/3$ - or  $1/10$ -bar tension (33kPa or 10kPa tension) and oven dryness. Volume change is influenced by the amount and type of clay minerals in the soil. The volume change is the percent change for the whole soil. If it is expressed as a fraction, the resulting value is COLE, coefficient of linear extensibility.

**Liquid limit.** The moisture content at which the soil passes from a plastic to a liquid state.

**Loam.** Soil material that is 7 to 27 percent clay particles, 28 to 50 percent silt particles, and less than 52 percent sand particles.

- Loess.** Material transported and deposited by wind and consisting dominantly of silt-sized particles.
- Low strength.** The soil is not strong enough to support loads.
- Low-residue crops.** Such crops as corn used for silage, peas, beans, and potatoes. Residue from these crops is not adequate to control erosion until the next crop in the rotation is established. These crops return little organic matter to the soil.
- Map unit.** Any soil, miscellaneous land type, soil complex, or undifferentiated soil group shown on a detailed soil map and identified by a symbol.
- Marl.** An earthy, unconsolidated deposit consisting chiefly of calcium carbonate mixed with clay in approximately equal proportions; formed primarily under freshwater lacustrine conditions but also formed in more saline environments.
- Mass movement.** A generic term for the dislodgment and downslope transport of soil and rock material as a unit under direct gravitational stress.
- Masses.** See Redoximorphic features.
- Meander belt.** The zone within which migration of a meandering channel occurs; the flood-plain area included between two imaginary lines drawn tangential to the outer bends of active channel loops.
- Mechanical treatment.** Use of mechanical equipment for seeding, brush management, and other management practices.
- Medium textured soil.** Very fine sandy loam, loam, silt loam, or silt.
- Mesa.** A broad, nearly flat topped and commonly isolated landmass bounded by steep slopes or precipitous cliffs and capped by layers of resistant, nearly horizontal rocky material. The summit width is characteristically greater than the height of the bounding escarpments.
- Metamorphic rock.** Rock of any origin altered in mineralogical composition, chemical composition, or structure by heat, pressure, and movement at depth in the earth's crust. Nearly all such rocks are crystalline.
- Mine spoil.** An accumulation of displaced earthy material, rock, or other waste material removed during mining or excavation. Also called earthy fill.
- Mineral soil.** Soil that is mainly mineral material and low in organic material. Its bulk density is more than that of organic soil.
- Minimum tillage.** Only the tillage essential to crop production and prevention of soil damage.
- Miscellaneous area.** A kind of map unit that has little or no natural soil and supports little or no vegetation.
- Miscellaneous land type.** A map unit consisting of areas of land that have little or no natural soil, that are too nearly inaccessible for orderly examination, or that occur where, for other reasons, it is not feasible to classify the soil.
- Moderately coarse textured soil.** Coarse sandy loam, sandy loam, or fine sandy loam.
- Moderately fine textured soil.** Clay loam, sandy clay loam, or silty clay loam.
- Mollic epipedon.** A thick, dark, humus-rich surface horizon (or horizons) that has high base saturation and pedogenic soil structure. It may include the upper part of the subsoil.
- Morphology, soil.** The physical makeup of the soil, including the texture, structure, porosity, consistence, color, and other physical, mineral, and biological properties of the various horizons, and the thickness and arrangement of those horizons in the soil profile.
- Mottling, soil.** Irregular spots of different colors that vary in number and size. Descriptive terms are as follows: abundance—*few*, *common*, and *many*; size—*fine*, *medium*, and *coarse*; and contrast—*faint*, *distinct*, and *prominent*. The size measurements are of the diameter along the greatest dimension. *Fine* indicates less than 5 millimeters (about 0.2 inch); *medium*, from 5 to 15 millimeters (about 0.2 to 0.6 inch); and *coarse*, more than 15 millimeters (about 0.6 inch).

**Mountain.** A generic term for an elevated area of the land surface, rising more than 1,000 feet (300 meters) above surrounding lowlands, commonly of restricted summit area (relative to a plateau) and generally having steep sides. A mountain can occur as a single, isolated mass or in a group forming a chain or range. Mountains are formed primarily by tectonic activity and/or volcanic action but can also be formed by differential erosion.

**Mudstone.** A blocky or massive, fine grained sedimentary rock in which the proportions of clay and silt are approximately equal. Also, a general term for such material as clay, silt, claystone, siltstone, shale, and argillite and that should be used only when the amounts of clay and silt are not known or cannot be precisely identified.

**Munsell notation.** A designation of color by degrees of three simple variables—hue, value, and chroma. For example, a notation of 10YR 6/4 is a color with hue of 10YR, value of 6, and chroma of 4.

**Natric horizon.** A special kind of argillic horizon that contains enough exchangeable sodium to have an adverse effect on the physical condition of the subsoil.

**Natural drainage.** Refers to conditions that existed during the development of a soil, as opposed to altered drainage, which is commonly the result of artificial drainage or irrigation but may be caused by the sudden deepening of channels or the blocking of drainage outlets. Seven different classes of natural drainage are recognized.

**Neutral soil.** A soil having a pH value of 6.6 to 7.3. (See Reaction, soil.)

**Nodules.** See Redoximorphic features.

**Nose slope** (geomorphology). A geomorphic component of hills consisting of the projecting end (laterally convex area) of a hillside. The overland waterflow is predominantly divergent. Nose slopes consist dominantly of colluvium and slope-wash sediments (for example, slope alluvium).

**Nutrient, plant.** Any element taken in by a plant essential to its growth. Plant nutrients are mainly nitrogen, phosphorus, potassium, calcium, magnesium, sulfur, iron, manganese, copper, boron, and zinc obtained from the soil and carbon, hydrogen, and oxygen obtained from the air and water.

**Open space.** A relatively undeveloped green or wooded area provided mainly within an urban area to minimize feelings of congested living.

**Organic matter.** Plant and animal residue in the soil in various stages of decomposition. The content of organic matter in the surface layer is described as follows:

Very low .....	less than 0.5 percent
Low .....	0.5 to 1.0 percent
Moderately low .....	1.0 to 2.0 percent
Moderate .....	2.0 to 4.0 percent
High .....	4.0 to 8.0 percent
Very high .....	more than 8.0 percent

**Paleoterrace.** An erosional remnant of a terrace that retains the surface form and alluvial deposits of its origin but was not emplaced by, and commonly does not grade to, a present-day stream or drainage network.

**Pan.** A compact, dense layer in a soil that impedes the movement of water and the growth of roots. For example, *hardpan*, *fragipan*, *claypan*, *plowpan*, and *traffic pan*.

**Parent material.** The unconsolidated organic and mineral material in which soil forms.

**Ped.** An individual natural soil aggregate, such as a granule, a prism, or a block.

**Pedisediment.** A layer of sediment, eroded from the shoulder and backslope of an erosional slope, that lies on and is being (or was) transported across a gently sloping erosional surface at the foot of a receding hill or mountain slope.

**Pedon.** The smallest volume that can be called “a soil.” A pedon is three dimensional and large enough to permit study of all horizons. Its area ranges from about 10 to 100 square feet (1 square meter to 10 square meters), depending on the variability of the soil.

**Percolation.** The movement of water through the soil.

**Permeability.** The quality of the soil that enables water or air to move downward through the profile. The rate at which a saturated soil transmits water is accepted as a measure of this quality. In soil physics, the rate is referred to as “saturated hydraulic conductivity,” which is defined in the “Soil Survey Manual.” In line with conventional usage in the engineering profession and with traditional usage in published soil surveys, this rate of flow continues to be expressed as “permeability.” Terms describing permeability, measured in inches per hour, are as follows:

Impermeable .....	less than 0.0015 inch
Very slow .....	0.0015 to 0.06 inch
Slow .....	0.06 to 0.2 inch
Moderately slow .....	0.2 to 0.6 inch
Moderate .....	0.6 inch to 2.0 inches
Moderately rapid .....	2.0 to 6.0 inches
Rapid .....	6.0 to 20 inches
Very rapid .....	more than 20 inches

**pH value.** A numerical designation of acidity and alkalinity in soil. (See Reaction, soil.)

**Phase, soil.** A subdivision of a soil series based on features that affect its use and management, such as slope, stoniness, and flooding.

**Plastic limit.** The moisture content at which a soil changes from semisolid to plastic.

**Plasticity index.** The numerical difference between the liquid limit and the plastic limit; the range of moisture content within which the soil remains plastic.

**Plateau** (geomorphology). A comparatively flat area of great extent and elevation; specifically, an extensive land region that is considerably elevated (more than 100 meters) above the adjacent lower lying terrain, is commonly limited on at least one side by an abrupt descent, and has a flat or nearly level surface. A comparatively large part of a plateau surface is near summit level.

**Playa.** The generally dry and nearly level lake plain that occupies the lowest parts of closed depressions, such as those on intermontane basin floors. Temporary flooding occurs primarily in response to precipitation and runoff. Playa deposits are fine grained and may or may not have a high water table and saline conditions.

**Plinthite.** The sesquioxide-rich, humus-poor, highly weathered mixture of clay with quartz and other diluents. It commonly appears as red mottles, usually in platy, polygonal, or reticulate patterns. Plinthite changes irreversibly to an ironstone hardpan or to irregular aggregates on repeated wetting and drying, especially if it is exposed also to heat from the sun. In a moist soil, plinthite can be cut with a spade. It is a form of laterite.

**Plowpan.** A compacted layer formed in the soil directly below the plowed layer.

**Ponding.** Standing water on soils in closed depressions. Unless the soils are artificially drained, the water can be removed only by percolation or evapotranspiration.

**Poorly graded.** Refers to a coarse grained soil or soil material consisting mainly of particles of nearly the same size. Because there is little difference in size of the particles, density can be increased only slightly by compaction.

**Pore linings.** See Redoximorphic features.

**Potential native plant community.** See Climax plant community.

- Potential rooting depth (effective rooting depth).** Depth to which roots could penetrate if the content of moisture in the soil were adequate. The soil has no properties restricting the penetration of roots to this depth.
- Prescribed burning.** Deliberately burning an area for specific management purposes, under the appropriate conditions of weather and soil moisture and at the proper time of day.
- Productivity, soil.** The capability of a soil for producing a specified plant or sequence of plants under specific management.
- Profile, soil.** A vertical section of the soil extending through all its horizons and into the parent material.
- Proper grazing use.** Grazing at an intensity that maintains enough cover to protect the soil and maintain or improve the quantity and quality of the desirable vegetation. This practice increases the vigor and reproduction capacity of the key plants and promotes the accumulation of litter and mulch necessary to conserve soil and water.
- Rangeland.** Land on which the potential natural vegetation is predominantly grasses, grasslike plants, forbs, or shrubs suitable for grazing or browsing. It includes natural grasslands, savannas, many wetlands, some deserts, tundras, and areas that support certain forb and shrub communities.
- Reaction, soil.** A measure of acidity or alkalinity of a soil, expressed as pH values. A soil that tests to pH 7.0 is described as precisely neutral in reaction because it is neither acid nor alkaline. The degrees of acidity or alkalinity, expressed as pH values, are:

Ultra acid .....	less than 3.5
Extremely acid .....	3.5 to 4.4
Very strongly acid .....	4.5 to 5.0
Strongly acid .....	5.1 to 5.5
Moderately acid .....	5.6 to 6.0
Slightly acid .....	6.1 to 6.5
Neutral .....	6.6 to 7.3
Slightly alkaline .....	7.4 to 7.8
Moderately alkaline .....	7.9 to 8.4
Strongly alkaline .....	8.5 to 9.0
Very strongly alkaline .....	9.1 and higher

- Red beds.** Sedimentary strata that are mainly red and are made up largely of sandstone and shale.
- Redoximorphic concentrations.** See Redoximorphic features.
- Redoximorphic depletions.** See Redoximorphic features.
- Redoximorphic features.** Redoximorphic features are associated with wetness and result from alternating periods of reduction and oxidation of iron and manganese compounds in the soil. Reduction occurs during saturation with water, and oxidation occurs when the soil is not saturated. Characteristic color patterns are created by these processes. The reduced iron and manganese ions may be removed from a soil if vertical or lateral fluxes of water occur, in which case there is no iron or manganese precipitation in that soil. Wherever the iron and manganese are oxidized and precipitated, they form either soft masses or hard concretions or nodules. Movement of iron and manganese as a result of redoximorphic processes in a soil may result in redoximorphic features that are defined as follows:
1. Redoximorphic concentrations.—These are zones of apparent accumulation of iron-manganese oxides, including:
    - A. Nodules and concretions, which are cemented bodies that can be removed from the soil intact. Concretions are distinguished from nodules on

the basis of internal organization. A concretion typically has concentric layers that are visible to the naked eye. Nodules do not have visible organized internal structure; *and*

B. Masses, which are noncemented concentrations of substances within the soil matrix; *and*

C. Pore linings, i.e., zones of accumulation along pores that may be either coatings on pore surfaces or impregnations from the matrix adjacent to the pores.

2. Redoximorphic depletions.—These are zones of low chroma (chromas less than those in the matrix) where either iron-manganese oxides alone or both iron-manganese oxides and clay have been stripped out, including:

A. Iron depletions, i.e., zones that contain low amounts of iron and manganese oxides but have a clay content similar to that of the adjacent matrix; *and*

B. Clay depletions, i.e., zones that contain low amounts of iron, manganese, and clay (often referred to as silt coatings or skeletons).

3. Reduced matrix.—This is a soil matrix that has low chroma *in situ* but undergoes a change in hue or chroma within 30 minutes after the soil material has been exposed to air.

**Reduced matrix.** See Redoximorphic features.

**Regolith.** All unconsolidated earth materials above the solid bedrock. It includes material weathered in place from all kinds of bedrock and alluvial, glacial, eolian, lacustrine, and pyroclastic deposits.

**Relief.** The relative difference in elevation between the upland summits and the lowlands or valleys of a given region.

**Residuum (residual soil material).** Unconsolidated, weathered or partly weathered mineral material that accumulated as bedrock disintegrated in place.

**Rill.** A very small, steep-sided channel resulting from erosion and cut in unconsolidated materials by concentrated but intermittent flow of water. A rill generally is not an obstacle to wheeled vehicles and is shallow enough to be smoothed over by ordinary tillage.

**Rippable.** Rippable bedrock or hardpan can be excavated using a single-tooth ripping attachment mounted on a tractor with a 200–300 drawbar horsepower rating.

**Riser.** The vertical or steep side slope (e.g., escarpment) of terraces, flood-plain steps, or other stepped landforms; commonly a recurring part of a series of natural, steplike landforms, such as successive stream terraces.

**Road cut.** A sloping surface produced by mechanical means during road construction. It is commonly on the uphill side of the road.

**Rock fragments.** Rock or mineral fragments having a diameter of 2 millimeters or more; for example, pebbles, cobbles, stones, and boulders.

**Root zone.** The part of the soil that can be penetrated by plant roots.

**Runoff.** The precipitation discharged into stream channels from an area. The water that flows off the surface of the land without sinking into the soil is called surface runoff. Water that enters the soil before reaching surface streams is called ground-water runoff or seepage flow from ground water.

**Saline soil.** A soil containing soluble salts in an amount that impairs growth of plants. A saline soil does not contain excess exchangeable sodium.

**Sand.** As a soil separate, individual rock or mineral fragments from 0.05 millimeter to 2.0 millimeters in diameter. Most sand grains consist of quartz. As a soil textural class, a soil that is 85 percent or more sand and not more than 10 percent clay.

**Sandstone.** Sedimentary rock containing dominantly sand-sized particles.

**Saturated hydraulic conductivity ( $K_{sat}$ ).** See Permeability.

- Saturation.** Wetness characterized by zero or positive pressure of the soil water. Under conditions of saturation, the water will flow from the soil matrix into an unlined auger hole.
- Scarification.** The act of abrading, scratching, loosening, crushing, or modifying the surface to increase water absorption or to provide a more tillable soil.
- Sedimentary rock.** A consolidated deposit of clastic particles, chemical precipitates, or organic remains accumulated at or near the surface of the earth under normal low temperature and pressure conditions. Sedimentary rocks include consolidated equivalents of alluvium, colluvium, drift, and eolian, lacustrine, and marine deposits. Examples are sandstone, siltstone, mudstone, claystone, shale, conglomerate, limestone, dolomite, and coal.
- Sequum.** A sequence consisting of an illuvial horizon and the overlying eluvial horizon. (See Eluviation.)
- Series, soil.** A group of soils that have profiles that are almost alike, except for differences in texture of the surface layer. All the soils of a series have horizons that are similar in composition, thickness, and arrangement.
- Shale.** Sedimentary rock that formed by the hardening of a deposit of clay, silty clay, or silty clay loam and that has a tendency to split into thin layers.
- Sheet erosion.** The removal of a fairly uniform layer of soil material from the land surface by the action of rainfall and surface runoff.
- Shoulder.** The convex, erosional surface near the top of a hillslope. A shoulder is a transition from summit to backslope.
- Shrub-coppice dune.** A small, streamlined dune that forms around brush and clump vegetation.
- Side slope (geomorphology).** A geomorphic component of hills consisting of a laterally planar area of a hillside. The overland waterflow is predominantly parallel. Side slopes are dominantly colluvium and slope-wash sediments.
- Silica.** A combination of silicon and oxygen. The mineral form is called quartz.
- Silica-sesquioxide ratio.** The ratio of the number of molecules of silica to the number of molecules of alumina and iron oxide. The more highly weathered soils or their clay fractions in warm-temperate, humid regions, and especially those in the tropics, generally have a low ratio.
- Silt.** As a soil separate, individual mineral particles that range in diameter from the upper limit of clay (0.002 millimeter) to the lower limit of very fine sand (0.05 millimeter). As a soil textural class, soil that is 80 percent or more silt and less than 12 percent clay.
- Siltstone.** An indurated silt having the texture and composition of shale but lacking its fine lamination or fissility; a massive mudstone in which silt predominates over clay.
- Similar soils.** Soils that share limits of diagnostic criteria, behave and perform in a similar manner, and have similar conservation needs or management requirements for the major land uses in the survey area.
- Sinkhole.** A closed, circular or elliptical depression, commonly funnel shaped, characterized by subsurface drainage and formed either by dissolution of the surface of underlying bedrock (e.g., limestone, gypsum, or salt) or by collapse of underlying caves within bedrock. Complexes of sinkholes in carbonate-rock terrain are the main components of karst topography.
- Similarity index.** The present composition of the plant community on an ecological site in relation to the potential natural plant community for that site. Similarity index is expressed as *excellent*, *good*, *fair*, or *poor*, on the basis of how much the present plant community has departed from the potential.
- Site index.** A designation of the quality of a forest site based on the height of the dominant stand at an arbitrarily chosen age. For example, if the average height attained by dominant and codominant trees in a fully stocked stand at the age of 50 years is 75 feet, the site index is 75.



**Slickensides** (pedogenic). Grooved, striated, and/or glossy (shiny) slip faces on structural peds, such as wedges; produced by shrink-swell processes, most commonly in soils that have a high content of expansive clays.

**Slope.** The inclination of the land surface from the horizontal. Percentage of slope is the vertical distance divided by horizontal distance, then multiplied by 100. Thus, a slope of 20 percent is a drop of 20 feet in 100 feet of horizontal distance. In this survey, classes for simple slopes are as follows:

Nearly level .....	0 to 1 percent
Very gently sloping .....	1 to 3 percent
Gently sloping .....	3 to 5 percent
Moderately sloping .....	5 to 8 percent
Strongly sloping .....	8 to 12 percent
Moderately steep .....	12 to 20 percent
Steep .....	20 to 45 percent
Very steep .....	45 percent and higher

Classes for complex slopes are as follows:

Nearly level .....	0 to 3 percent
Gently undulating .....	1 to 5 percent
Undulating .....	1 to 8 percent
Gently rolling .....	5 to 12 percent
Rolling .....	5 to 15 percent
Hilly .....	8 to 30 percent
Steep .....	20 to 45 percent
Very steep .....	45 percent and higher

**Slope alluvium.** Sediment gradually transported down the slopes of mountains or hills primarily by nonchannel alluvial processes (i.e., slope-wash processes) and characterized by particle sorting. Lateral particle sorting is evident on long slopes. In a profile sequence, sediments may be distinguished by differences in size and/or specific gravity of rock fragments and may be separated by stone lines. Burnished peds and sorting of rounded or subrounded pebbles or cobbles distinguish these materials from unsorted colluvial deposits.

**Sodic (alkali) soil.** A soil having so high a degree of alkalinity (pH 8.5 or higher) or so high a percentage of exchangeable sodium (15 percent or more of the total exchangeable bases), or both, that plant growth is restricted.

**Sodicity.** The degree to which a soil is affected by exchangeable sodium. Sodicity is expressed as a sodium adsorption ratio (SAR) of a saturation extract, or the ratio of  $\text{Na}^+$  to  $\text{Ca}^{++} + \text{Mg}^{++}$ . The degrees of sodicity and their respective ratios are:

Slight .....	less than 13:1
Moderate .....	13-30:1
Strong .....	more than 30:1

**Sodium adsorption ratio (SAR).** A measure of the amount of sodium (Na) relative to calcium (Ca) and magnesium (Mg) in the water extract from saturated soil paste. It is the ratio of the Na concentration divided by the square root of one-half of the Ca + Mg concentration.

**Soft bedrock.** Bedrock that can be excavated with trenching machines, backhoes, small rippers, and other equipment commonly used in construction.

**Soil.** A natural, three-dimensional body at the earth's surface. It is capable of supporting plants and has properties resulting from the integrated effect of climate and living matter acting on earthy parent material, as conditioned by relief and by the passage of time.

**Soil separates.** Mineral particles less than 2 millimeters in equivalent diameter and ranging between specified size limits. The names and sizes, in millimeters, of separates recognized in the United States are as follows:

Very coarse sand .....	2.0 to 1.0
Coarse sand .....	1.0 to 0.5
Medium sand .....	0.5 to 0.25
Fine sand .....	0.25 to 0.10
Very fine sand .....	0.10 to 0.05
Silt .....	0.05 to 0.002
Clay .....	less than 0.002

**Solum.** The upper part of a soil profile, above the C horizon, in which the processes of soil formation are active. The solum in soil consists of the A, E, and B horizons. Generally, the characteristics of the material in these horizons are unlike those of the material below the solum. The living roots and plant and animal activities are largely confined to the solum.

**Stone line.** In a vertical cross section, a line formed by scattered fragments or a discrete layer of angular and subangular rock fragments (commonly a gravel- or cobble-sized lag concentration) that formerly was draped across a topographic surface and was later buried by additional sediments. A stone line generally caps material that was subject to weathering, soil formation, and erosion before burial. Many stone lines seem to be buried erosion pavements, originally formed by sheet and rill erosion across the land surface.

**Stones.** Rock fragments 10 to 24 inches (25 to 60 centimeters) in diameter if rounded or 15 to 24 inches (38 to 60 centimeters) in length if flat.

**Stony.** Refers to a soil containing stones in numbers that interfere with or prevent tillage.

**Strath terrace.** A type of stream terrace; formed as an erosional surface cut on bedrock and thinly mantled with stream deposits (alluvium).

**Stream terrace.** One of a series of platforms in a stream valley, flanking and more or less parallel to the stream channel, originally formed near the level of the stream; represents the remnants of an abandoned flood plain, stream bed, or valley floor produced during a former state of fluvial erosion or deposition.

**Stripcropping.** Growing crops in a systematic arrangement of strips or bands that provide vegetative barriers to wind erosion and water erosion.

**Structure, soil.** The arrangement of primary soil particles into compound particles or aggregates. The principal forms of soil structure are—*platy* (laminated), *prismatic* (vertical axis of aggregates longer than horizontal), *columnar* (prisms with rounded tops), *blocky* (angular or subangular), and *granular*. *Structureless* soils are either *single grained* (each grain by itself, as in dune sand) or *massive* (the particles adhering without any regular cleavage, as in many hardpans).

**Stubble mulch.** Stubble or other crop residue left on the soil or partly worked into the soil. It protects the soil from wind erosion and water erosion after harvest, during preparation of a seedbed for the next crop, and during the early growing period of the new crop.

**Subsoil.** Technically, the B horizon; roughly, the part of the solum below plow depth.

**Subsoiling.** Tilling a soil below normal plow depth, ordinarily to shatter a hardpan or claypan.

**Substratum.** The part of the soil below the solum.

**Subsurface layer.** Any surface soil horizon (A, E, AB, or EB) below the surface layer.

**Summer fallow.** The tillage of uncropped land during the summer to control weeds and allow storage of moisture in the soil for the growth of a later crop. A practice common in semiarid regions, where annual precipitation is not enough to produce

a crop every year. Summer fallow is frequently practiced before planting winter grain.

- Summit.** The topographically highest position of a hillslope. It has a nearly level (planar or only slightly convex) surface.
- Surface layer.** The soil ordinarily moved in tillage, or its equivalent in uncultivated soil, ranging in depth from 4 to 10 inches (10 to 25 centimeters). Frequently designated as the "plow layer," or the "Ap horizon."
- Surface soil.** The A, E, AB, and EB horizons, considered collectively. It includes all subdivisions of these horizons.
- Talus.** Rock fragments of any size or shape (commonly coarse and angular) derived from and lying at the base of a cliff or very steep rock slope. The accumulated mass of such loose broken rock formed chiefly by falling, rolling, or sliding.
- Taxadjuncts.** Soils that cannot be classified in a series recognized in the classification system. Such soils are named for a series they strongly resemble and are designated as taxadjuncts to that series because they differ in ways too small to be of consequence in interpreting their use and behavior. Soils are recognized as taxadjuncts only when one or more of their characteristics are slightly outside the range defined for the family of the series for which the soils are named.
- Terrace (conservation).** An embankment, or ridge, constructed across sloping soils on the contour or at a slight angle to the contour. The terrace intercepts surface runoff so that water soaks into the soil or flows slowly to a prepared outlet. A terrace in a field generally is built so that the field can be farmed. A terrace intended mainly for drainage has a deep channel that is maintained in permanent sod.
- Terrace (geomorphology).** A steplike surface, bordering a valley floor or shoreline, that represents the former position of a flood plain, lake, or seashore. The term is usually applied both to the relatively flat summit surface (tread) that was cut or built by stream or wave action and to the steeper descending slope (scarp or riser) that has graded to a lower base level of erosion.
- Terracettes.** Small, irregular steplike forms on steep hillslopes, especially in pasture, formed by creep or erosion of surficial materials that may be induced or enhanced by trampling of livestock, such as sheep or cattle.
- Texture, soil.** The relative proportions of sand, silt, and clay particles in a mass of soil. The basic textural classes, in order of increasing proportion of fine particles, are *sand, loamy sand, sandy loam, loam, silt loam, silt, sandy clay loam, clay loam, silty clay loam, sandy clay, silty clay, and clay*. The sand, loamy sand, and sandy loam classes may be further divided by specifying "coarse," "fine," or "very fine."
- Tilth, soil.** The physical condition of the soil as related to tillage, seedbed preparation, seedling emergence, and root penetration.
- Toeslope.** The gently inclined surface at the base of a hillslope. Toeslopes in profile are commonly gentle and linear and are constructional surfaces forming the lower part of a hillslope continuum that grades to valley or closed-depression floors.
- Topsoil.** The upper part of the soil, which is the most favorable material for plant growth. It is ordinarily rich in organic matter and is used to topdress roadbanks, lawns, and land affected by mining.
- Trace elements.** Chemical elements, for example, zinc, cobalt, manganese, copper, and iron, in soils in extremely small amounts. They are essential to plant growth.
- Tread.** The flat to gently sloping, topmost, laterally extensive slope of terraces, flood-plain steps, or other stepped landforms; commonly a recurring part of a series of natural steplike landforms, such as successive stream terraces.
- Upland.** An informal, general term for the higher ground of a region, in contrast with a low-lying adjacent area, such as a valley or plain, or for land at a higher elevation

than the flood plain or low stream terrace; land above the footslope zone of the hillslope continuum.

**Valley fill.** The unconsolidated sediment deposited by any agent (water, wind, ice, or mass wasting) so as to fill or partly fill a valley.

**Variants, soil.** A soil having properties sufficiently different from those of other known soils to justify a new series name, but occurring in such a limited geographic area that creation of a new series is not justified.

**Variation.** Refers to patterns of contrasting colors assumed to be inherited from the parent material rather than to be the result of poor drainage.

**Water-holding capacity.** See Available water capacity.

**Weathering.** All physical disintegration, chemical decomposition, and biologically induced changes in rocks or other deposits at or near the earth's surface by atmospheric or biologic agents or by circulating surface waters but involving essentially no transport of the altered material.

**Well graded.** Refers to soil material consisting of coarse grained particles that are well distributed over a wide range in size or diameter. Such soil normally can be easily increased in density and bearing properties by compaction. Contrasts with poorly graded soil.

**Wilting point (or permanent wilting point).** The moisture content of soil, on an oven-dry basis, at which a plant (specifically a sunflower) wilts so much that it does not recover when placed in a humid, dark chamber.

**Windthrow.** The uprooting and tipping over of trees by the wind.

**Winnowing.** The removal of clay and silt particles from the soil by strong winds. The coarser particles remain, and the soil becomes sandier and more highly erodible as the process continues.

# **NRCS Accessibility Statement**

---

The Natural Resources Conservation Service (NRCS) is committed to making its information accessible to all of its customers and employees. If you are experiencing accessibility issues and need assistance, please contact our Helpdesk by phone at 1-800-457-3642 or by e-mail at [ServiceDesk-FTC@ftc.usda.gov](mailto:ServiceDesk-FTC@ftc.usda.gov). For assistance with publications that include maps, graphs, or similar forms of information, you may also wish to contact our State or local office. You can locate the correct office and phone number at <http://offices.sc.egov.usda.gov/locator/app>.