14240

Trench Soil (vacuum container) 184 grams




Figure 1: Soil Mechanics Trench at Apollo 14. It was only about 30 cm deep. AS14-64-9161.

CC: Okay. And Al, one question, did you get the SESC sample out of the bottom of the trench?

CDR: Well, I told you the trench was kind of a miserable thing, because the walls kept falling down. And I could get a sample from the bottom, but it wouldn't be from the bottom, I'm afraid.

CC: I guess we'd still like the SESC sample form the bottom of the trench, even though it probably isn't the bottom.

CDR: Well, I'll tell you. I'll go back and whack at it a little bit. See what I can do.

CDR: We're digging the bottom of the trench for you, Fredo. I'm redigging the trench. I can't believe it!

LMP: What's the matter. Al?

CDR: Oh, that (vacuum) seal came off that thing.

Introduction

14240 was collected from the bottom of the Soil Mechanics Trench (figure 1) and brought back in a special environmental sample container (SESC) which was opened by Burlingame in the UC Berkeley Space Science Laboratory (figure 5). However, the lid was not even finger tight and the sample probably not sealed (although the vacuum in the can may have held the lid on). The gas in the SESC was mainly N_2 so the lid may have leaked in the LRL glovebox, rather than in the LM, CM or during transit.

Petrography

The Soil Mechanics Trench was about 25 - 36 cm deep (Mitchell et al. 1971). 14240 was taken from the bottom (although it may contain material slumped from the walls). The trench samples were taken during the 2nd EVA from the rim of a ~6 meter crater at station G. It was planned to dig this trench to be 60 cm deep, but was cut instead to ~30 cm because the walls were caving in. According to Mitchell et al., there was noticeable color change in the wall of the trench. "The upper 3 to 5 cm were dark brown and fine grained. Next, a very thin layer (0.5 cm thick or less) of black, glassy particles were encountered. Beneath this layer was a much lighter colored and coarser grained material."

Subsurface samples 14240 (and 14149 from the same trench) are both immature soils (Is/FeO = 46 and 53; Morris 1978). The soil collected from the top of the trench (14148) was more mature Is/FeO = 74.

Various small particles found in the SESC were set aside (figures 7-9).

Chemistry

Rose et al. (1972) and Laul et al. (1972) analyzed 14240 (table 1). Cadogan et al. (1972) and Moore et al. (1972) reported the carbon content (figure 4).

Processing

The samples from the bottom of this trench (14149 and 14240) may have been contaminated by surface material that fell into the trench during digging. As you can see form the transcript, 14240 was the second attempt to sample the bottom of the trench. The slightly lower maturity seems to indicate that it was less contaminated by slumped surface material.

This sample was opened in a dry He glove box at Berkeley (figure 6), and thus may be free of contamination at the LRL (or at least have a different source(s) of contamination). It has been kept in a He "bean pot" at JSC, separate from other lunar samples, all these years.

The official LRL weight was 168 grams based on the difference between the weight of hardware and weight of returned sample, but the actual weight of sample was found to be 184 grams when the sample was opened in the UCB SSL Berkeley (Burlingame 1971).




Figure 2: Normalized rare-earth-element diagram for 14240 compared with 14149 (data from Laul et al. 1972).




Figure 3: Chemical composition of 14240 compared with other lunar soils.




Figure 4: Carbon content and matruity of lunar soils. Data for 14240 from compilations by Cadogen et al. (1972), desMarais et al. (1976) and Morris et al. (1978).

Table 1. Chemical composition of 14240.

reference weight SiO2 % TiO2 Al2O3 FeO MnO MgO CaO Na2O K2O P2O5	1.8 19.4 10.9 0.125 12 0.717 0.57	2 18.5 0.13 12 0.745 0.6	(a) (a) (a) (a) (a)	Rose72 47.77 1.67 17.99 10.02 0.14 9.47 11.25 0.7 0.54 0.55	(b) (b) (b) (b) (b) (b) (b) (b) (b)	14149 Laul82 for comparis 1.6 17.9 10 0.13 9.9 11 0.78 0.58	son
S % sum Sc ppm	22		(a)	28	(b)	21	
V Cr Co Ni Cu Zn Ga Ge ppb As	55 1430 36	40	(a) (a)	52 1574 33 320 16 23 7.5	(b) (b) (b) (b) (b) (b)	40 1232 27.5 320	1
Se Rb Sr Y Zr Nb Mo Ru Rh Pd ppb Ag ppb Cd ppb In ppb Sn ppb Sb ppb Te ppb	790		(a)	13 390 250 930 42	(b) (b) (b) (b) (b)	170 680	
Cs ppm Ba La Ce Pr	790 70 214		(a) (a) (a)	67	(b)	870 60.5 170	

(a) (a)

(a) 23

(a)

(a)

(a)

(b)

110

26.5

2.2

5.4

37 8.2

3.3 20.2

2.8

21.9 3

13.3

3.6




Figure 5: Drawing of SESC container (Allton 1989).



Figure 6: Opening the SESC in the SESC holder at the Berkeley Lab (Burlingame et al. 1971).

technique: (a) INAA, (b) "microchemical"

Nd

 Sm

Eu

Gd

Tb Dy

Ho Er Tm

Yb Lu

Hf

Ta

W ppb Re ppb Os ppb


Ir ppb
Pt ppb
Au ppb
Th ppm
U ppm

31 2.65

24

3.2

20



References for 14240

Allton J.H. (1989) Catalog of Apollo Lunar Surface Geological Sampling Tools and Containers. JSC 23454.

Baur H., Frick U., Funk H., Schultz L. and Signer P. (1972) Thermal release of He, Ne, and Ar from lunar fines and minerals. Proc. 3rd Lunar Sci. Conf. 1947-1966.

Burlingame A.L., Holland P., McFadden W.H., Simoneit B.R., Wilder J.T. and Wszolek P.C. (1971) UCB Space Sciences Laboratory Simulation 3 sand transfer and transfer of Apollo 14 SESC Lunar Material. June 17, 1971

Cadogen P.H., Eglinton G., Maxwell J.R. and Pillinger C.T. (1971) Carbon chemistry of lunar surface. Nature 231, 29.

Cadogen P.H., Eglinton G., Firth J.N.M., Maxwell J.R., Mays B.J. and Pillinger C.T. (1972) Survey of lunar carbon compounds. Proc. 3rd Lunar Sci. Conf. 2069-2090.

DesMarais D.J., Hayes J.M. and Meinshein W.G. (1973b) The distribution in lunar soil of carbon released by pryrolysis. Proc. 4th Lunar Sci. Conf. 1543-1558.

DesMarais D.J., Basu A., Hayes J.M. and Meinshein W.G. (1975) Evolution of carbon isotopes, agglutinates and the lunar regolith. Proc. 6th Lunar Sci. Conf. 2353-2374.

Epstein S. and Taylor H.P. (1972) O18/O16, Si30/Si28, C13/C12 and D/H studies of Apollo 14 and 15 samples. Proc. 3rd Lunar Sci. Conf. 1429-1455.

Flory D.A., Wikstrom S., Gupta S., Gibert J.M. and Oro J. (1972) Organogenic compounds in Apollo 11, 12 and 14 lunar samples. Proc. 3rd Lunar Sci. Conf. 2091-2108.

Gehrke C.W., Zumwalt R.W., Kuo K., Aue W.A., Stalling D.L., Kvenvolden K.A. and Ponnamperuma C. (1972) Amino acid analysis of Apollo 14 fines. Proc. 3rd Lunar Sci. Conf. 2119-2129.

Graf J.C. (1993) Lunar Soils Grain Size Catalog. NASA Pub. 1265

Holland P.T., Simoneit B.R., Wszolek P.C. and Burlingame A.L. (1972) Compounds of carbon and other volatile elements in Apollo 14 and 15 samples. Proc. 3rd Lunar Sci. Conf. 2131-2149.

Laul J.C., Wakita H., Showalter D.L., Boynton W.V. and Schmitt R.A. (1972) Bulk, rare earth, and other trace elements in Apollo 14 and 15 and Luna 16 samples. Proc. 3rd Lunar Sci. Conf. 1181-1201

Modzeleski V.E., et al. (1973) Carbon compounds in pyrolysates and amino-acid in extracts of Apollo 14 lunar samples. Nature 242, 50.

Moore C.B., Lewis C.F., Cripe J.D., Delles F.M., Kelly W.R. and Gibson E.K. (1972) Total carbon, nitrogen and sulfur in Apollo 14 lunar samples. Proc. 3rd Lunar Sci. Conf. 2051-2058.

Morris R.V. (1978) The surface exposure (maturity) of lunar soils: Some concepts and Is/FeO compilation. Proc. 9th Lunar Planet . Sci. Conf. 2287-2298.

Morris R.V., Score R., Dardano C. and Heiken G. (1983) Handbook of Lunar Soils. JSC 19069

Rose H.J., Cuttitta F., Annell C.S., Carron M.K., Christian R.P., Dwornik E.J., Greenland L.P. and Ligon D.T. (1972) Compositional data for twenty-one Fra Mauro lunar materials. Proc. 3rd Lunar Sci. Conf. 1215-1231.

Swann G.A., Bailey N.G., Batson R.M., Eggleton R.E., Hait M.H., Holt H.E., Larson K.B., McEwen M.C., Mitchell E.D., Schaber G.G., Schafer J.P., Shepard A.B., Sutton R.L., Trask N.J., Ulrich G.E., Wilshire H.G. and Wolf E.W. (1971)



Figure 7: 14240,22 is the first rock found in 14240 and weighs 2.49 g. It appears to be a dark matrix breccia, but has never been studied. It is about 1.2 cm across. (from Burlingame et al. 1971).




Figure 8: One of three small rocks in 14240,23. It appears to be light matrix breccia, but has never been studied. It is about 1 cm across.



Figure 9: Additional small rocks found in 14240, now collectively numbered 14240,23 (weight 5.06 g). Never studied.

Preliminary geologic investigations of the Apollo 14 landing site. In Apollo 14; Preliminary Science Report. NASA SP-272, 39-85.

Swann G.A., Bailey N.G., Batson R.M., Eggleton R.E., Hait M.H., Holt H.E., Larson K.B., Reed V.S., Schaber G.G., Sutton R.L., Trask N.J., Ulrich G.E. and Wilshire H.G. (1977)

Geology of the Apollo 14 landing site in the Fra Mauro highlands. U.S. Geological Survey Professional Paper 880

Warner J. (1970) Apollo 14 Lunar Sample Information Catalog. NASA MSC