Superconducting Clusters and Colossal Effects in Underdoped Cuprates

Gonzalo Alvarez Computer Science and Mathematics Division Oak Ridge National Laboratory

Matthias Mayr Mas-Plank-Institut für Festkörperforschung

Adrian Moreo and Elbio Dagotto University of Tennessee

Phenomenological models for the antiferromagnetic vs. d-wave superconductivity competition in cuprates are studied [1] using conventional Monet Carlo techniques. The analysis suggests that cuprates may show a variety of different behaviors in the very underdoped regime: local coexistence, stripes, or, if disorder is present, states with Nanoscale superconducting clusters. The transition from an antiferromagnetic to a super-conducting State does not seem universal. Under suitable conditions, non-superconducting Cu-oxides could rapidly [2] become superconducting by the influence of weak perturbations that align the randomly oriented phase of the superconducting clusters in the mixed state. Consequences of these ideas for angle resolved photoemission and scanning tunneling microscopy experiments [3] will also be discussed.

- [1] Alvarez et al., cond-mat/0401474, to appear in PRB.
- [2] I. Bozovic et al., PRL 93, 157002, (2004).
- [3] A. Ino et al., PRB 62, 4127 (2000); K. Lang et al., Nature 415, 412 (2002).