

32


**NASA TECHNICAL
MEMORANDUM**

REPORT NO. 53905

ON THE REAL SINGULARITIES OF THE N-BODY PROBLEM

By Hans J. Sperling
Aero-Astroynamics Laboratory

September 12, 1969



NASA

*George C. Marshall Space Flight Center
Marshall Space Flight Center, Alabama*

N70-37562
(ACCESSION NUMBER)
102
(PAGES)
TMX-53905
(NASA CR OR TMX OR AD NUMBER)

(THRU)

(CODE)
130
(CATEGORY)

NASA - GEORGE C. MARSHALL SPACE FLIGHT CENTER

IN-AERO-68-6
was changed to TM X-53905 on September 12, 1969

October 15, 1968

ON THE REAL SINGULARITIES OF THE N-BODY PROBLEM

By

Hans J. Sperling

TECHNICAL AND SCIENTIFIC STAFF
AERO-ASTRODYNAMICS LABORATORY
RESEARCH AND DEVELOPMENT OPERATIONS

Abstract

A collision singularity in the N-body problem has the property that all position vectors have finite limits and at least one mutual distance approaches zero as time approaches the instant of the singularity. General properties of a collision singularity are investigated and it is found that they are analogous to those of a collision of all bodies at the same point and time. A more detailed study of a simple type of such a singularity, namely that of several simultaneous binary collisions, is presented.

Introduction

Little is known about the singularities of the equations of motion of the N-body problem of celestial mechanics, particularly if complex values of the independent and dependent variables are admitted. Only the three simplest cases have received considerable attention, as manifested by the bibliography: the binary collision in the three-body problem (e.g., Sundman [4]) and in the N-body problem (e.g., Wintner [2]), the triple collision in the three-body problem (e.g., Sundman [1], Siegel [1]), and the collision of all N bodies at the same point and instant (e.g., Block [2], Wintner [2]); all of the above studies are restricted to real values of all variables. For a study of the binary collision in the N-body problem for complex values of the variables, see Sperling [3].

It is presently unknown which types of real singularities (in the sense of complex analysis as well as with regard to their dynamical interpretation) are possible in the N-body problem; the only general result is a remarkable statement by von Zeipel [2] (cf. 205), asserting that a singularity is a collision singularity (cf. 203) if the mutual distances remain bounded as time approaches the instant of the singularity. "Collision singularities" are such that the N bodies separate into distinct "clusters" with all bodies in each cluster colliding at the instant of the singularity at a well-defined point, while the clusters remain apart from each other. Since von Zeipel's statement seems to be virtually unknown and particularly in view of Wintner's [2, p. 431] remark: "A note of H. von Zeipel indicates a consideration to the effect that, if U becomes infinite when t tends to a finite value, then J must tend to infinity, unless all bodies tend to definite limiting positions. But it seems to be hard to fill in the

gaps ", we will subsequently present a detailed proof, following essentially von Zeipel's ideas. Chazy's [6] attempt of a proof is only a brief sketch and must be considered at least vague. The introduction of " cluster coordinates " (cf. 103-104) enables us to prove in a simple manner that the existence of finite limits of the mutual distances implies the existence of the limits of the positions and relative positions as t approaches a finite value (cf. 216); Wintner [2, p. 327] mentions this as an undecided problem (cf. von Zeipel [2, p. 4]).

It will be shown that the known important properties of a collision of all bodies at the same instant hold, with obvious modifications, also for a collision singularity, the main result probably being that the bodies of each cluster are positioned close to a central configuration when the time is close to the instant of the singularity.

A detailed study of the case of several simultaneous binary collisions concludes the paper. This case has already been considered by Lahaye [3]; his treatment, however, is at least incomplete: his assumption, that the asymptotic behavior of a perturbed two-body problem equals that of the two-body problem as the instant of a binary collision is approached, is not justified without proof, and he does not show that his solutions are the only ones. The main result of the investigation is that this singularity gives rise to an algebraic branch point of the second order on the real time axis, the same as for one binary collision; in particular, the motion can be analytically continued in the real beyond this singularity.

While writing this paper, I was informed (an abstract in the Notices Amer. Math. Soc. 15 (1968) 631 and private communication) that H. Pollard has worked in the same field. Details of his work have not been available to me at the time of the completion of this paper.

On the notation

Let $t^* \neq \pm \infty$ (later $t^* = 0$) be the instant of the (collision) singularity;
all limits are taken as $t \rightarrow t^*$ ($t < t^*$).

c, \underline{c} denote positive constants

b, \underline{b} denote functions of t or $s = t^{\frac{1}{3}}$, bounded on an interval
 $[t_0, t^*[$ with suitable $t_0 < t^*$;

e, \underline{e} denote functions of t or s , with $e \rightarrow 0, \underline{e} \rightarrow 0$ as $t \rightarrow t^*$.

The above symbols have fixed meaning, at least within the given context, if they are indexed; without indexes they are unspecified, i. e., they may denote different constants or functions from one occurrence to the next. For example, in

$$\int_0^t b(\tau) d\tau = b(t) \quad \text{and in} \quad f(t) = b(t) + t b(t) \quad \text{the } b\text{'s}$$

occurring in the same equation may denote different functions.

The absolute value of a vector \underline{z} is $z (= |\underline{z}| = +(\underline{z}\underline{z})^{\frac{1}{2}})$.

\sim denotes asymptotic behavior as $t \rightarrow t^* (= 0)$ or $s \rightarrow 0$: $f \sim g$ is equivalent to $f = g + o(g)$.

Differentiation with respect to time t is denoted by $\dot{}$: $df/dt = \dot{f}$.

Differentiation with respect to s is denoted by $'$: $df/ds = f'$.

The gradient of a scalar f with respect to a vector \underline{z} is denoted

$$\text{by } \text{grad}_{\underline{z}} f = \frac{\partial f}{\partial \underline{z}} \quad .$$

Subscript following semicolon: 0: some fixed value; 1, 2, 3: the three components of a vector.

100 Equations and integrals of motion

For reference we list here the equations and integrals of motion and some related concepts and equations, in the two coordinate systems used later. The geometric basis is the 3-dimensional euclidean space.

101 (Cartesian) inertial coordinates, the center of mass resting at the origin. N particles (=bodies) with masses $m_k > 0$ are moving according to Newton's law of gravitation. Let γ be the gravitational constant, \underline{z}_k the position vector from the origin to the body m_k , and

$$\underline{z}_{jk} = \underline{z}_k - \underline{z}_j \quad (101.1)$$

the relative position vector from m_j to m_k .

Set

$$M = \sum_{k=1}^N m_k \quad (101.2)$$

102 Equations of motion

$$\ddot{\underline{z}}_k = -\gamma \sum_{\substack{j=1 \\ j \neq k}}^N m_j \frac{\underline{z}_k - \underline{z}_j}{|\underline{z}_k - \underline{z}_j|^3} = -\gamma \sum_{\substack{j=1 \\ j \neq k}}^N m_j \frac{\underline{z}_{jk}}{z_{jk}^3} = -\frac{1}{m_k} \frac{\partial U}{\partial \underline{z}_k} \quad k = 1 \dots N, \quad (102.1)$$

potential energy

$$U = -\gamma \sum_{1 \leq j < k \leq N} \sum_{1 \leq j < k \leq N} \frac{m_j m_k}{|\underline{z}_k - \underline{z}_j|} = -\gamma \sum_{1 \leq j < k \leq N} \sum_{1 \leq j < k \leq N} \frac{m_j m_k}{z_{jk}}, \quad (102.2)$$

kinetic energy

$$V = \frac{1}{2} \sum_{j=1}^N m_j \dot{\underline{z}}_j \dot{\underline{z}}_j = \frac{1}{2} \sum_{j=1}^N m_j |\dot{\underline{z}}_j|^2, \quad (102.3a)$$

$$V = \frac{1}{4M} \sum_{j=1}^N \sum_{k=1}^N m_j m_k z_{jk} z'_{jk} \quad , \quad (102.3b)$$

polar inertia momentum

$$J = \sum_{j=1}^N m_j z_j z'_j = \sum_{j=1}^N m_j z_j^2 \quad (102.4a)$$

$$= \frac{1}{2M} \sum_{j=1}^N \sum_{k=1}^N m_j m_k z_{jk} z'_{jk} = \frac{1}{M} \sum_{1 < j < k \leq N} m_j m_k z_{jk}^2 \quad , \quad (102.4b)$$

angular momentum

$$\underline{K} = \sum_{j=1}^N m_j z_j \times z'_j = \frac{1}{2M} \sum_{j=1}^N \sum_{k=1}^N m_j m_k z_{jk} \times z'_{jk} \quad , \quad (102.5)$$

conservation of energy (energy theorem)

$$V + U = h \quad (= \text{constant of energy}) \quad , \quad (102.6)$$

conservation of angular momentum

$$\underline{K} = \underline{\text{constant}} \quad , \quad (102.7)$$

center of mass rests at the origin

$$\sum_{j=1}^N m_j z_j = \underline{0} \quad , \quad (102.8)$$

Lagrange-Jacobi equation

$$J'' = -2U + 4h = 2V + 2h \quad ; \quad (102.9)$$

the positions z_k are expressed in terms of the relative positions

$$\text{by } M z_k = \sum_{j=1}^N m_j z_{jk} \quad . \quad (102.10)$$


103 "Cluster coordinates". Partition the set of all N bodies into L nonempty disjoint subsets, called "clusters" in the following. Let N_ℓ , $1 \leq \ell \leq L$, be the number of bodies in the ℓ -th cluster. Obviously, $N_1 + \dots + N_L = N$. (Observe that subscripts are separated by commas in cluster coordinates and that $\ell = 1 \dots L$, $\kappa = 1 \dots N_\ell$). Notation:

$m_{\ell, k}$ = mass of the k -th body in the ℓ -th cluster,

$\underline{z}_{\ell, k}$ = position vector from the origin to $m_{\ell, k}$,

\underline{Z}_ℓ = position vector from the origin to the center of mass of the ℓ -th cluster,

$\underline{\zeta}_{\ell, k}$ = position vector from the center of mass of the ℓ -th cluster to $m_{\ell, k}$.



Obviously,

$$\underline{z}_{\ell, k} = \underline{Z}_\ell + \underline{\zeta}_{\ell, k} \quad ; \quad (103.1)$$

define M_ℓ and \underline{Z}_ℓ by

$$M_\ell = \sum_{k=1}^{N_\ell} m_{\ell, k} \quad \text{and} \quad M_\ell \underline{Z}_\ell = \sum_{k=1}^{N_\ell} m_{\ell, k} \underline{z}_{\ell, k} \quad ; \quad (103.2)$$

then

$$\sum_{k=1}^{N_\ell} m_{\ell, k} \zeta_{\ell, k} = 0 \quad (103.3)$$

are

$$M_{\ell, k}^{\zeta} = \sum_{j=1}^{N_\ell} m_{\ell, j} (\zeta_{\ell, k} - \zeta_{\ell, j}) \quad (103.4)$$

104 Equations of motion

$$\ddot{z}_{\ell, k} = -\gamma \sum_{\lambda=1}^L \sum_{\substack{n=1 \\ \lambda \neq \ell}}^{N_\lambda} m_{\lambda, n} \frac{z_{\ell, k} - z_{\lambda, n}}{|z_{\ell, k} - z_{\lambda, n}|^3}, \quad (104.1)$$

$$M_{\ell, k}^{\ddot{z}} = -\gamma \sum_{\lambda=1}^L \sum_{\substack{n=1 \\ \lambda \neq \ell}}^{N_\lambda} \sum_{k=1}^{N_\ell} m_{\ell, k} m_{\lambda, n} \frac{z_{\ell, k} - z_{\lambda, n}}{|z_{\ell, k} - z_{\lambda, n}|^3} \quad (104.2)$$

$$z_{\ell, k} - z_{\lambda, n} = z_{\ell} - z_{\lambda} + \zeta_{\ell, k} - \zeta_{\lambda, n},$$

$$\ddot{\zeta}_{\ell, k} = -\gamma \sum_{\substack{i=1 \\ j \neq k}}^{N_\ell} m_{\ell, j} \frac{\zeta_{\ell, k} - \zeta_{\ell, j}}{|\zeta_{\ell, k} - \zeta_{\ell, j}|^3} -$$

(104.3a)

$$\gamma \sum_{\substack{\lambda=1 \\ \lambda \neq \ell}}^L \sum_{n=1}^{N_\lambda} m_{\lambda, n} \left[\frac{z_{\ell} - z_{\lambda} + \zeta_{\ell, k} - \zeta_{\lambda, n}}{|z_{\ell} - z_{\lambda} + \zeta_{\ell, k} - \zeta_{\lambda, n}|^3} - \frac{1}{M_{\ell, k}} \sum_{\mu=1}^{N_\ell} m_{\ell, \mu} \frac{z_{\ell} - z_{\lambda} + \zeta_{\ell, \mu} - \zeta_{\lambda, n}}{|z_{\ell} - z_{\lambda} + \zeta_{\ell, \mu} - \zeta_{\lambda, n}|^3} \right]$$

or

$$\ddot{u}_{l,k} = - \frac{1}{m_{l,k}} \frac{\partial U_l}{\partial \underline{u}_{l,k}} + \dots, k \quad (104.3b)$$

Define the potential energy of the l -th cluster by

$$U_l = -\gamma \sum_{k=1}^{N_l} \sum_{\substack{j=1 \\ k < j}}^{N_l} \frac{m_{l,j} m_{l,k}}{|\underline{r}_{l,j} - \underline{r}_{l,k}|} \quad (104.4)$$

its kinetic energy by

$$V_l = \frac{1}{2} \sum_{k=1}^{N_l} m_{l,k} \dot{\underline{r}}_{l,k} \dot{\underline{r}}_{l,k} \quad (104.5)$$

its (total) energy by

$$h_l = V_l + U_l \quad (104.6)$$

(observe that h_l is not a constant in general) ,

its polar inertia momentum by

$$J_l = \sum_{k=1}^{N_l} m_{l,k} \underline{r}_{l,k} \underline{r}_{l,k} \quad (104.7a)$$

$$= \frac{1}{2M_l} \sum_{j=1}^{N_l} \sum_{k=1}^{N_l} m_{l,j} m_{l,k} (\underline{r}_{l,k} - \underline{r}_{l,j}) (\underline{r}_{l,k} - \underline{r}_{l,j}), \quad (104.7b)$$

and its angular momentum by

$$K_{\omega l} = \sum_{k=1}^{N_l} m_{l,k} \zeta_{\omega l,k} \times \zeta_{\omega l,k} \quad ; \quad (104.8)$$

define further the kinetic energy of the centers of mass of all clusters by

$$V^0 = \frac{1}{2} \sum_{l=1}^L M_l \dot{Z}_{\omega l} \dot{Z}_{\omega l} \quad , \quad (104.9)$$

their polar inertia momentum by

$$J^0 = \sum_{l=1}^L M_l Z_{\omega l} Z_{\omega l} \quad , \quad (104.10)$$

and their angular momentum by

$$K^0 = \sum_{l=1}^L M_l Z_{\omega l} \times Z_{\omega l} \quad ; \quad (104.11)$$

write

$$U = \sum_{l=1}^L U_l + U^+ = U^* + U^+ \quad , \quad (104.12)$$

observing that U^+ consists exactly of those terms of U which contain the difference of coordinates of masses in different clusters.

Using (103.3), it is easy to show that

$$V = \sum_{l=1}^L V_l + V^0 = V^* + V^0 \quad , \quad (104.13)$$

$$J = \sum_{l=1}^L J_l + J^0 = J^* + J^0 \quad , \quad (104.14)$$

$$\underline{K} = \sum_{\ell=1}^L \underline{K}_{\ell} + \underline{K}^0 = \underline{K}^* + \underline{K}^0 \quad , \quad (104.15)$$

and

$$h = \sum_{\ell=1}^L h_{\ell} + V^0 + U^+ = h^* + V^0 + U^+ \quad . \quad (104.16)$$

200 Classification and properties of real singularities

As mentioned before, we restrict ourselves (necessarily) to real values of the independent variable time t and the dependent variables, the positions and velocities; consider only real analytic continuation of the solution of the equations of motion and exclude the two cases that $t = (\pm)\infty$, or that the finite t is an accumulation point of singularities from both sides: then, if there is a singularity at t^* , this point is the right or left endpoint of an open interval on which the solution is holomorphic. Without restriction of generality, we will assume that t^* is the right endpoint of such an interval.

201 For definiteness, we refer to the equations of motion (102.1), but analogous statements obviously hold for the equations in the form (104.2), (104.3). Let $t_0, z_{k;0}, z'_{k;0}$ be finite and real such that $|z_{k;0} - z_{j;0}| \neq 0$ for $j \neq k$; by the existence theorem for analytic differential equations there exists an interval $|t - t_0| < \delta$ and on it a unique holomorphic solution $z_k(t), z'_k(t)$, $k = 1 \dots N$, of (102.1) satisfying $z_k(t_0) = z_{k;0}, z'_k(t_0) = z'_{k;0}$. It is noteworthy (and soon to be used) that an estimate of δ can be made involving only $\min |z_{k;0} - z_{j;0}|$ and $\max |z'_{k;0}|$, but not $z_{k;0}$ (cf. Siegel [3], pp. 23-25). Continue this solution holomorphically from t_0 for increasing t ; the finite point t^* is a singularity, if the solution is holomorphic on $[t_0, t^*[$, but not on $[t_0, t^*]$.

202 Theorem a): t^* is a singularity iff $\liminf(\min z_{jk}) = 0$ as $t \rightarrow t^*$.
 b): t^* is a singularity iff $\lim(\min z_{jk}) = 0$ as $t \rightarrow t^*$.

Proof:

1. If t^* is a singularity then $\liminf(\min z_{jk}) = 0$: assume contrariwise that $\liminf(\min z_{jk}) \geq d_0 > 0$.

Then there exist numbers t_1 and d_1 , $t_0 \leq t_1 < t^*$ and $0 < d_1 \leq d_0$, such that $z_{jk} \geq d_1$ on $[t_1, t^*[$; we infer that $|z_{jk} z_{jk}^{-3}| = z_{jk}^{-2} \leq d_1^{-2}$

and, using the equations of motion (102.1), $|z''_{jk}| \leq \gamma M d_1^{-2}$. From

$$z_k = z_k(t_1) + (t - t_1) z'_k(t_1) + \int_{t_1}^t (t - \tau) z''_k(\tau) d\tau$$

it follows that z_k and z'_k have finite limits z_k^* and z'^*_k with $z^*_{jk} = |z^*_k - z^*_j| \geq d_1$ for $j \neq k$ as $t \rightarrow t^*$. The existence theorem

(cf. 201), applied to initial values very close to t^* , z_k^* , z_k^{\setminus} , states that the solution remains holomorphic at t^* , contrary to the assumption.

2. If $\liminf(\min z_{jk}) = 0$, then t^* is a singularity: if at least one z_k^{\setminus} is unbounded as $t \rightarrow t^*$, then there is a singularity at t^* . Assume now that all z_k^{\setminus} remain bounded; then, by integration, all z_k^{\setminus} are bounded. In

$$U^{\setminus} = \sum_{j=1}^3 \sum_{k=1}^N \frac{\partial U}{\partial z_{k;j}} z_{k;j}^{\setminus} \quad \text{the } \frac{\partial U}{\partial z_{k;j}} \text{ are bounded by (102.1) and the}$$

assumption; thus, U^{\setminus} and therefore U are bounded. This is inconsistent with the fact that $\liminf(\min z_{jk}) = 0$ implies $\limsup U = -\infty$.

3. If t^* is a singularity, then $\lim(\min z_{jk}) = 0$: assume contrariwise that $\limsup(\min z_{jk}) \geq d_0 > 0$. There exists a sequence $t_\nu \rightarrow t^*$ such that $z_{jk}(t_\nu) \geq d_1 > 0$, $-U \leq d_2$, and by the energy theorem $V \leq d_3$, implying $|z_k^{\setminus}(t_\nu)| \leq d_4$. Apply the existence theorem (cf. 201) to t_ν , $z_k(t_\nu)$, $z_k^{\setminus}(t_\nu)$ as initial values; if t_ν is sufficiently close to t^* , t^* is contained in the interval $|t - t_\nu| < \delta$ where the solution remains holomorphic, yielding a contradiction (observe that δ is a function of d_1 and d_4 only, but not of t_ν).

4. If $\lim(\min z_{jk}) = 0$, then t^* is a singularity: this is implied by 2.

Observe that a) does not require the energy theorem (102.6) for its proof.

203 Definition: A singularity t^* is called a collision singularity if all position vectors \underline{z}_k have finite limits as $t \rightarrow t^*$.

Hence, by 202 theorem, at least one of the mutual distances z_{jk} converges to zero as $t \rightarrow t^*$ for this type of singularity.

According to the original definition, the partition of the bodies into clusters (cf. 103) is quite arbitrary; in the following this partition is always used in conjunction with a collision singularity and assumed to be consistent with it, i. e.: two bodies m_j and m_k are in the same cluster iff z_{jk} is very small as $t \rightarrow t^*$ = instant of the collision singularity, or in terms of cluster coordinates:

$$\begin{aligned} \text{all } |\xi_{m_l, k} - \xi_{m_l, j}| & \text{ are very small compared with} \\ & (203.1) \\ |\underline{z}_{m_l} - \underline{z}_{m_\lambda}|, \quad l \neq \lambda, & \text{ on the considered time interval.} \end{aligned}$$

204 Lemma: Let the $\underline{z}_k = \underline{z}_k(t)$ be holomorphic on $t_0 \leq t < t^*$.
Then $\lim J \leq \infty$ exists as $t \rightarrow t^*$.

Proof (cf. Wintner [2] p. 327):

Let t^* be a singularity, otherwise the statement is trivial; then $\lim(\min z_{jk}) = 0$, hence $\lim(-U) = +\infty$. From $J'' = 4h - 2U$ it follows that J'' is ultimately positive; hence, J' is ultimately increasing and therefore ultimately does not change sign, and we conclude that J is ultimately monotonic (increasing or decreasing), implying the existence of the limit of J . Since $J \geq 0$, $\lim J$ is ∞ or a nonnegative number.

205 Theorem (von Zeipel): If there is a singularity at t^* and $\lim J < \infty$ as $t \rightarrow t^*$, then t^* is a collision singularity.

Proof: Assume that the solution of the N-body problem is holomorphic on $[t_0, t^*[$, $t_0 < t^*$.

206 Let first be $\lim J = 0$: then $z_k \rightarrow 0$ and $z_{jk} \rightarrow 0$ by (102.4), implying that $\omega_k \rightarrow 0$ and $\omega_{jk} \rightarrow 0$; all positions z_k and all relative positions z_{jk} have the finite limit zero and we have a collision singularity. All bodies collide at time t^* at the center of mass.

207 Now let $0 < \lim J = J(t^*) < \infty$; we partition the set of all mutual distances z_{jk} into four disjoint subsets as follows:

- first type: $\lim z_{jk} > 0$ exists;
- second type: $\lim z_{jk} = 0$ exists;
- third type: $\lim z_{jk}$ does not exist and $\liminf z_{jk} > 0$;
- fourth type: $\lim z_{jk}$ does not exist and $\liminf z_{jk} = 0$,
 $\limsup z_{jk} > 0$.

We show in the following that $0 < \lim J < \infty$ implies the existence and finiteness of all $\lim z_{jk}$. In a first case (cf. 208), we assume that there are no mutual distances of the fourth type and prove our assertion directly; in a second case (cf. 209-215), we assume that there is at least one distance of the fourth type and lead this assumption to a contradiction. 216 lemma then concludes the proof.

208 First case: there are no mutual distances of the fourth type. There exist a t_{00} , $t_0 \leq t_{00} < t^*$, and two constants a_1 and a_2 , $0 < a_1 < a_2$, such that on $[t_{00}, t^*[$

- for all mutual distances of the first and third type: $z_{jk} > a_2$,
- and for all mutual distances of the second type: $z_{jk} < a_1$.

We consider now the solution on $[t_{00}, t^*[$. Introduce cluster coordinates such that two bodies are in the same cluster iff for their mutual distance $z_{jk} < a_1$ holds.

Note: Because the mutual distances z_{jk} are continuous on $[t_{00}, t^*[$ and either $z_{jk} < a_1 < a_2$ or $z_{jk} > a_2 > a_1$, the separation of the bodies into clusters is unique on this interval; i. e., each body stays in one and the same cluster.

All mutual distances of bodies in the same cluster, say the l -th, are by the definition of the cluster of the second type, i. e., $|\zeta_{\omega l, k} - \zeta_{\omega l, j}| \rightarrow 0$ as $t \rightarrow t^*$. By (103.4) this implies $\zeta_{l, k} \rightarrow 0$, hence $\zeta_{\omega l, k} \rightarrow 0$, i. e., all $\lim_{\omega l, k} \zeta_{\omega l, k}$ exist (and are zero).

Consider the equation of motion (104.2) for the center of mass $Z_{\omega l}$ of the l -th cluster; on the right of this equation only relative positions and mutual distances of bodies in different clusters occur, i. e., the mutual distances are of the first or third type; hence $|z_{\omega l, k} - z_{\omega \lambda, \kappa}| > a_2$, $|(z_{\omega l, k} - z_{\omega \lambda, \kappa})| z_{\omega l, k} - z_{\omega \lambda, \kappa}|^{-3} < a_2^{-2}$, and $|Z_{\omega l}''| < \gamma M a_2^{-2}$. This together with

$$Z_{\omega l} = Z_{\omega l}(t_{00}) + (t - t_{00}) Z_{\omega l}'(t_{00}) + \int_{t_{00}}^t (t - \tau) Z_{\omega l}''(\tau) d\tau$$

implies that $\lim_{\omega l} Z_{\omega l}$ exists and is finite as $t \rightarrow t^*$. We conclude that $\lim_{\omega l, k} z_{\omega l, k} = \lim_{\omega l} Z_{\omega l} + \lim_{\omega l, k} \zeta_{\omega l, k}$, $\lim_{\omega l, k} z_{\omega l, k}$, $\lim_{\omega l, k} z_{jk}$, and $\lim_{\omega l, k} z_k$ all exist and are finite; i. e., in this first case all mutual distances have finite limits.

209 Second case: There is at least one mutual distance of the fourth type. The existence of $\lim J = J(t^*) > 0$ implies that for every $\epsilon > 0$ there exists a t_{0*} , $t_0 \leq t_{0*} < t^*$, such that

$$|J(t) - J(t^*)| < \epsilon \quad \text{on } [t_{0*}, t^*[\quad . \quad (209.1)$$

Assume without restriction that $\epsilon < \frac{1}{2} J(t^*)$; then $J_{\min} > \frac{1}{2} J(t^*)$ on $[t_{0*}, t^*[$, and with (102.4) it follows that

$$\max_{jk} z_{jk} \cdot \left(\frac{J(t^*)}{M} \right)^{\frac{1}{2}} = a_3 \quad \text{on } [t_{0*}, t^*[\quad (209.2)$$

and

$$z_{jk} < \left(\frac{3 M J(t^*)}{2 m_{\min}^2} \right)^{\frac{1}{2}} = a_4 \quad \text{on } [t_{0*}, t^*[\quad , \quad (209.3)$$

and by (102.10)

$$z_k < a_4 \quad , \quad k = 1 \dots N, \quad \text{on } [t_{0*}, t^*[\quad . \quad (209.4)$$

210 Define $n(\tau, d)$ as the number of mutual distances z_{jk} for which $z_{jk}(\tau) \geq d$ holds; consider $n(\tau, d)$ on the set $\{ \tau, d : t_{0*} \leq \tau < t^*, 0 < d \leq a_3 \}$; $\frac{1}{2}N(N-1)$ being the total number of mutual distances, $n(\tau, d)$ satisfies

$$1 \leq n(\tau, d) \leq \frac{1}{2}N(N-1) \quad \text{on this set ;} \quad (210.1)$$

the first inequality holding because of (209.2).

Define

$$\mathfrak{N}(t, d) = \text{g. l. b. } \mathfrak{S}(t, d) \quad (210.2)$$

$$\mathfrak{S}(t, d) = \{ n(\tau, d) : \tau \in [t, t^*[, t_{0*} \leq t < t^*, 0 < d \leq a_3 , \\ t \text{ fixed, } d \text{ fixed} \} ;$$

since each $n(\tau, d)$ is a natural number satisfying (210.1), their greatest lower bound exists and is the minimum of all competing $n(\tau, d)$. $\mathfrak{N}(t, d)$ satisfies

$$1 \leq \mathfrak{N}(t, d) \leq \frac{1}{2}N(N-1) - 1 \quad \text{on} \quad \{ t, d : t_{0*} \leq t < t^* , 0 < d \leq a_3 \} , \quad (210.3)$$

the second inequality holding because of $\lim(\min z_{jk}) = 0$. $\mathfrak{N}(t, d)$ is thus a number such that at every point of the interval $[t, t^*[$ at least $\mathfrak{N}(t, d)$ mutual distances are $\geq d$. Furthermore,

$$\mathfrak{N}(t_\sigma, d_\sigma) \geq \mathfrak{N}(t, d) \quad \text{for} \quad t \leq t_\sigma < t^* , 0 < d_\sigma \leq d : \quad (210.4)$$

for, $\mathfrak{S}(t_\sigma, d_\sigma)$ is a subset of $\mathfrak{S}(t, d)$ and hence $\text{g.l.b.} \mathfrak{S}(t_\sigma, d_\sigma) \geq \text{g.l.b.} \mathfrak{S}(t, d)$.

Define

$$\mathfrak{N}^* = \text{l.u.b.} \{ \mathfrak{N}(t, d) : t_{0*} \leq t < t^* , 0 < d \leq a_3 \} ; \quad (210.5a)$$

since each $\mathfrak{N}(t, d)$ is a natural number satisfying (210.3), their least upper bound exists and is the maximum of all competing $\mathfrak{N}(t, d)$.

Let t_m and d_m , $t_{0*} \leq t_m < t^*$, $0 < d_m \leq a_3$, be such that

$$\mathfrak{N}^* = \mathfrak{N}(t_m, d_m) ; \quad (210.5b)$$

by (210.4) we have

$$\mathfrak{N}^* = \mathfrak{N}(t_m, d_m) = \mathfrak{N}(t, d) \quad \text{for} \quad t_m \leq t < t^* , 0 < d \leq d_m . \quad (210.6)$$

Note: There are always at least $\mathfrak{N}(t, d)$ mutual distances with $z_{jk} \geq d$ on $[t, t^*[$, but the individual mutual distances satisfying this may change from one instant to another.

211 We define now the constants used in the subsequent analysis and further restrict in this context the interval on which we consider the solution. First, choose t_{0**} , $t_{0*} \leq t_{0**} < t^*$, and $a_5 > 0$ such that for all mutual distances of the first and of the third type (i.e., those with $\liminf z_{jk} > 0$)

$$z_{jk} > a_5 \quad \text{on} \quad [t_{0**}, t^*[\quad , \quad (211.1)$$

and that for all mutual distances of the fourth type

$$\limsup z_{jk} > a_5 \quad \text{on} \quad [t_{0**}, t^*[\quad . \quad (211.2)$$

By (210.6) d_m can be replaced by a smaller positive number without changing \mathfrak{N}^* ; replace d_m by d_* (subsequently defined). Then

$$\text{at every point of } [t_{0**}, t^*[\quad \text{at least} \quad (211.3)$$

$$\mathfrak{N}^* \text{ mutual distances are } \geq d_* .$$

Set

$$d_* = \left(10^{-Q} \frac{2 M J(t^*)}{m_{\min}^2} \right)^{\frac{1}{2}} \quad , \quad (211.4)$$

$$\epsilon_* = \left(10^{-Q-10} \frac{m_{\min}^2}{M^2} \right)^{\frac{1}{2}} d_* \quad , \quad (211.5)$$

and determine the natural number Q such that

$$d_* < \min(d_m, a_5) \quad \text{and} \quad \epsilon_* < 10^{-1} d_* \quad ; \quad (211.6)$$

set further

$$a_6 = \left(\gamma M^2 M^{-1} \frac{a_4}{\ell \min} d_*^{-2} \right)^{\frac{1}{2}} \quad , \quad (211.7)$$

$$a_7 = a_6 + |t_{0**} - t^*| \gamma M d_*^{-2} \quad , \quad (211.8)$$

$$a_5 = 2M \left(a_7^2 + \gamma M a_4 d_*^{-2} \right) , \quad (211.9)$$

$$\epsilon = \frac{M}{2} \epsilon_*^2 = 10^{-2Q-10} J(t^*) . \quad (211.10)$$

Now a t_* , $t_* < t^*$, can be determined such that

- (1) $t_* \geq \max(t_{0**}, t_m, t^* - \epsilon a_8^{-1})$,
- (2) all mutual distances of the second type satisfy $z_{jk} < \epsilon_*$, and (211.11)
- (3) $|J - J(t^*)| < \epsilon$ with ϵ from (211.10) .

We show: there exist $t_{1*}, t_2, t_3, t_* < t_{1*} < t_2 < t_3 < t^*$, such that

- (a) every mutual distance of the fourth type is $> a_5$ at least once on $[t_{1*}, t_2]$;
- (b) exactly $\frac{1}{2}N(N-1) - \mathfrak{N}^*$ mutual distances are $< \epsilon_*$ on $[t_2, t_3]$.

(a): Select a t_{1*} on $]t_*, t^*[$ and consider the mutual distances z_{pq} of the fourth type for t increasing from t_{1*} ; since they satisfy $\limsup z_{pq} > a_5$ (cf. (211.2)), for each of them there is an instant t_{pq} on $[t_{1*}, t^*[$ such that $z_{pq}(t_{pq}) > a_5$. Let $t_{2*}, t_{1*} \leq t_{2*} < t^*$, be such that on $[t_{1*}, t_{2*}]$ for each of them there is such an instant t_{pq} where $z_{pq}(t_{pq}) > a_5$ ($t_{2*} = \max(t_{pq})$ will do) .

(b): Consider all mutual distances on $[t_{2*}, t^*[$; assume that contrary to the assertion at every point of this interval less than $\frac{1}{2}N(N-1) - \mathfrak{N}^*$ mutual distances are $< \epsilon_*$; then always more than \mathfrak{N}_2^* mutual distances are $\geq \epsilon_*$, and we have $\mathfrak{N}(t_{2*}, \epsilon_*) > \mathfrak{N}^*$, a contradiction to the maximum property of \mathfrak{N}^* (cf. (210.6), (211.6), (211.11)).

Correction to para. 4
at bottom of p. 20.

(b): Consider all mutual distances on $[t_{2*}, t^*[$. Assume that at every point of this interval more than \mathfrak{N}^* mutual distances are $\geq \epsilon_*$, i. e. that $n(\tau, \epsilon_*) > \mathfrak{N}^*$, $\tau \in [t_{2*}, t^*[$; then $\mathfrak{N}(t_{2*}, \epsilon_*) > \mathfrak{N}^*$, a contradiction to the maximum property of \mathfrak{N}^* (cf. (210.6), (211.6), (211.11)). Hence $n(\tau_\nu, \epsilon_*) \leq \mathfrak{N}^*$ for infinitely many τ_ν in $[t_{2*}, t^*[$, and since $\mathfrak{N}(t_{2*}, d_*) = \mathfrak{N}^*$ by the choice of t_* , ϵ_* , and d_* , we have also $n(\tau, \epsilon_*) \geq n(\tau, d_*) \geq \mathfrak{N}^*$ for $\tau \in [t_{2*}, t^*[$. It follows that $n(\tau_\nu, \epsilon_*) = \mathfrak{N}^*$, i. e. exactly $\frac{1}{2}N(N-1) - \mathfrak{N}^*$ mutual distances are $< \epsilon_*$ at τ_ν ; take one of these τ_ν for t_2 , then there is a nearby $t_3 > t_2$ such that by their continuity the number of mutual distances being $< \epsilon_*$ on $[t_2, t_3]$ still equals $\frac{1}{2}N(N-1) - \mathfrak{N}^*$.

212 We have shown that $\frac{1}{2}N(N-1) - \eta^*$ mutual distances are $< \epsilon_*$ on $[t_2, t_3]$, while all other η^* mutual distances are $\geq d_*$ on this interval. Set

$$\mathfrak{S}_{\epsilon_*} = \{ z_{jk} : z_{jk} < \epsilon_* \text{ on } [t_2, t_3] \} , \quad (212.1)$$

$$\mathfrak{S}_{d_*} = \{ z_{jk} : z_{jk} \geq d_* \text{ on } [t_2, t_3] \} ;$$

because the mutual distances are continuous functions of time, there is no exchange of elements between $\mathfrak{S}_{\epsilon_*}$ and \mathfrak{S}_{d_*} . We know that all mutual distances of the first and of the third type are $> d_*$ on $[t_2, t^*[$ and therefore in \mathfrak{S}_{d_*} (cf. (211.1)), and all of the second type $< \epsilon_*$ on $[t_2, t^*[$ and therefore in $\mathfrak{S}_{\epsilon_*}$ (cf. (211.11)). We prove that at least one mutual distance of the fourth type must be in $\mathfrak{S}_{\epsilon_*}$, i. e., $< \epsilon_*$ on $[t_2, t_3]$: assume contrariwise that all (and at least one) mutual distances of the fourth type are in \mathfrak{S}_{d_*} . None of the mutual distances in $\mathfrak{S}_{\epsilon_*}$ (all of them are of the second type) can increase beyond ϵ_* on $[t_2, t^*[$, i. e., no mutual distance can change from $\mathfrak{S}_{\epsilon_*}$ to \mathfrak{S}_{d_*} , and therefore none of the η^* mutual distances in \mathfrak{S}_{d_*} can decrease below d_* , because then less than η^* mutual distances would be $\geq d_*$: while by the definition of η^* always at least η^* mutual distances are $\geq d_*$ on $[t_2, t^*[$. This last result is contrary to the foregoing assumption that there exists a mutual distance of the fourth type in \mathfrak{S}_{d_*} , because this must necessarily decrease below d_* on $[t_2, t^*[$. Thus we have shown that there is a mutual distance z_{pq} of the fourth type in $\mathfrak{S}_{\epsilon_*}$.

It follows that there is a t_{3*} , $t_3 < t_{3*} < t^*$, such that $z_{pq}(t_{3*}) = d_*$.

Consider the mutual distances in $\mathfrak{S}_{\epsilon_*}$ for t increasing from t_3 ; it follows now that

there is a t_4 , $t_3 < t_4 < t^*$, such that

- (1) at least one mutual distance of $\mathfrak{S}_{\epsilon_*}$ equals d_* at t_4
 and (212.2)
 (2) all mutual distances of $\mathfrak{S}_{\epsilon_*}$ are $< d_*$ on $[t_2, t_4[$.

Considering the mutual distances in $\mathfrak{S}_{\epsilon_*}$ for t decreasing from t_2 ,

we find analogously (cf. 211 (a)) that

there is a t_1 , $t_* < t_1 < t_2$, such that

- (1) at least one mutual distance of $\mathfrak{S}_{\epsilon_*}$ equals d_* at t_1
 and (212.3)
 (2) all mutual distances of $\mathfrak{S}_{\epsilon_*}$ are $< d_*$ on $]t_1, t_3]$.

Observe that

all mutual distances of \mathfrak{S}_{d_*} remain $\geq d_*$ on $[t_1, t_4]$. (212.4)

Introduce cluster coordinates such that two bodies are in the same cluster iff their mutual distance is $< \epsilon_*$, i.e., it is in $\mathfrak{S}_{\epsilon_*}$ (, and in different clusters iff their mutual distance is $\geq d_*$, i.e., it is in \mathfrak{S}_{d_*}) on $[t_2, t_3]$.

By (104.7) and (104.14) we have

$$J^* < \frac{M}{2} \epsilon_*^2 \quad \text{on } [t_2, t_3] \quad (212.5)$$

and

$$J^* > \frac{m_{\min}^2}{2M} d_*^2 \quad \text{at } t_1 \quad \text{and at } t_4 ; \quad (212.6)$$

it follows that J^* has a minimum on $]t_1, t_4[$, and (cf. (211.5))

$$J^*_{\max} - J^*_{\min} > \frac{m_{\min}^2}{2M} d_*^2 (1 - 10^{-Q-10}) \quad \text{on } [t_1, t_4] . \quad (212.7)$$

213 J^0 has a maximum on $]t_1, t_4[$: with $J^0 = J - J^*$ we find

$$J^0 > J(t^*) - \epsilon - \frac{M}{2} \epsilon_*^2 = J(t^*) (1 - 2 \cdot 10^{-2Q-10}) \quad \text{on } [t_2, t_3] \quad (213.1)$$

and

$$J^0 < J(t^*) + \epsilon - \frac{m_{\min}^2}{2M} d_*^2 = J(t^*) (1 - 10^{-Q} + 10^{-2Q-10}) \quad \text{at } t_1 \quad \text{and at } t_4 ; \quad (213.2)$$

the right side of (213.1) is greater than the right side of (213.2) (recall that $Q \geq 1$, natural).

214 Let a maximum of J^0 , existing by 213, be at t_c , $t_1 < t_c < t_4$;

$$\text{then } J^{0\prime}(t_c) = 0 \quad , \quad J^{0\prime\prime}(t_c) \leq 0 \quad . \quad (214.1)$$

We estimate $J^{0\prime\prime}$ on $[t_1, t_4]$: by (104.10)

$$J^{0\prime\prime} = 2 \sum_{\ell=1}^L M_{\ell} \underline{z}_{\ell}^{\prime} \underline{z}_{\ell}^{\prime\prime} + 2 \sum_{\ell=1}^L M_{\ell} \underline{z}_{\ell} \underline{z}_{\ell}^{\prime\prime} \quad ; \quad (214.2)$$

all z_k are bounded (cf. (209.4)), therefore, all Z_{ℓ} are bounded by (103.2) :

$$Z_{\ell} < a_4 \quad \text{on } [t_1, t_4] \quad . \quad (214.3)$$

Estimate of $\underline{z}_{\ell}^{\prime\prime}$ (cf. (212.4)) :

$$|(\underline{z}_{\ell, k} - \underline{z}_{\lambda, \kappa})| \underline{z}_{\ell, k} - \underline{z}_{\lambda, \kappa} |^{-3} \leq d_*^{-2} \quad \text{for } \ell \neq \lambda \quad (214.4)$$

and from the equation of motion (104.2) for \underline{z}_{ℓ} we find

$$|\underline{z}_{\ell}^{\prime\prime}| < \gamma M d_*^{-2} \quad . \quad (214.5)$$

Estimate of $\underline{z}_{\ell}^{\prime}$:

at t_c , (214.1) and (214.2) imply that

$$M_{\underline{z}} \underline{z} \underline{z} \leq \sum_{\ell=1}^L M_{\underline{z}} \underline{z} \underline{z} \leq \left| \sum_{\ell=1}^L M_{\underline{z}} \underline{z} \underline{z} \right| < M a_4 \gamma M d_*^{-2} ,$$

hence (cf. (211.7))

$$| \underline{z} \underline{z} (t_c) | < \left(\gamma M^2 M_{\underline{z}}^{-1} \min a_4 d_*^{-2} \right)^{\frac{1}{2}} = a_6 , \quad (214.6)$$

and from

$$\underline{z} \underline{z} = \underline{z} \underline{z} (t_c) + \int_{t_c}^t \underline{z} \underline{z} (\tau) d\tau$$

it follows that (cf. (211.8))

$$\begin{aligned} | \underline{z} \underline{z} | &\leq | \underline{z} \underline{z} (t_c) | + | t_1 - t_4 | \gamma M d_*^{-2} \\ &< a_6 + | t_{0^{**}} - t^* | \gamma M d_*^{-2} = a_7 \quad \text{on } [t_1, t_4] ; \end{aligned} \quad (214.7)$$

thus finally (cf. (211.9))

$$| J^{00} | < 2M \left(a_7^2 + \gamma M a_4 d_*^{-2} \right) = a_8 \quad \text{on } [t_1, t_4] . \quad (214.8)$$

215 Write (cf. (209.1))

$$J = J(t^*) + \mathcal{E}_1 , \quad | \mathcal{E}_1 | < \epsilon , \quad (215.1)$$

and

$$J^0 = J^0(t_c) + \mathcal{E}_2 , \quad \mathcal{E}_2 = \int_{t_c}^t (t - \tau) J^{00}(\tau) d\tau , \quad (215.2)$$

$$| \mathcal{E}_2 | < | t_1 - t_4 | a_8 < | t_* - t^* | a_8 \leq \epsilon \quad (\text{cf. (211.11)}) .$$

Hence,

$$J^* = J - J^0 = J(t^*) - J^0(t_c) + \mathcal{E}_1 - \mathcal{E}_2 , \quad (215.3)$$

and

$$J_{\max}^* - J_{\min}^* < 4\epsilon = 2M\epsilon_*^2 = 2 \cdot 10^{-Q-10} \frac{m_{\min}^2}{M} d_*^2, \quad (215.4)$$

this being inconsistent with the inequality (212.7).

Thus, the assumption that there is a mutual distance of the fourth type is false, and taking 208 into account, we deduce that all mutual distances have finite limits. The proof is concluded by the following

216 Lemma: Let the N-body motion be holomorphic on $[t_0, t^*[$, $t_0 < t^*$, and let all mutual distances z_{jk} have finite limits as $t \rightarrow t^*$. Then all relative positions \underline{z}_{jk} and all positions \underline{z}_k have finite limits as $t \rightarrow t^*$.

I. e., at t^* the motion remains holomorphic, or there occurs a collision singularity. The proof is analogous to that in 208, observing that the set of mutual distances of the third type is empty.

300 Properties of a collision singularity

301 This part generalizes results which have been established for a collision of all bodies ("simultaneous collision"), to each cluster of the configuration developing prior to a collision singularity. The principal result is that, for t close to t^* , the configuration of each cluster is close to a central configuration. Our proofs are similar to those of Wintner, with some extensions and modifications.

302 Observe that a collision singularity at t^* is characterized by: at least one mutual distance converges to zero, and $J \geq 0$ remains bounded and has a limit as $t \rightarrow t^*$. Without restriction of generality we may assume that $t^* = 0$ and that the approach is from $t < 0$. Also, without mentioning it always, we assume that $t < 0$ is very close to the instant of collision singularity $t = 0$, such that for the cluster coordinates holds:

$$\begin{aligned} \underline{c}_{l,k} &\rightarrow 0 \text{ is very small on } [t, 0[, \text{ and} \\ |Z_{\underline{m}l} - Z_{\underline{m}\lambda}| &\geq \text{a certain positive constant on } [t, 0[, \quad l \neq \lambda . \end{aligned}$$

It follows also that $Z_{\underline{m}l}''$ is bounded (cf. (104.2)), consequently $\lim_{t \rightarrow 0} Z_{\underline{m}l}$ and $\lim_{t \rightarrow 0} Z_{\underline{m}l}'$, and finally $\lim_{t \rightarrow 0} Z_{\underline{m}l}''$ exist as $t \rightarrow 0$, as seen ~~by integrating~~ ~~from~~ (104.2).

303 Differential equation and ultimate behavior of J^* :

Using (102.9): $J'' = 4h - 2U$, (104.12): $U = U^* + U^+$, and (104.14):

$J = J^* + J^0$, we find

$$J^{*\prime\prime} = -2U^* + 4h - 2U^+ - J^{0\prime\prime} ; \quad (303.1)$$

both U^+ and $J^{0\prime\prime}$ (cf. 214) are bounded on the considered interval, and we write

$$J^{*\prime\prime} = -2U^* + b . \quad (303.2)$$

From $J^{*\prime\prime} = -2U^* + b$ and $-U^* \rightarrow \infty$ as $t \rightarrow 0$ we conclude that:

$$J^{*\prime\prime} \rightarrow +\infty,$$

$J^{*\prime}$ is monotonically increasing and ultimately does not change sign, and (303.3)

J^* is ultimately monotonically decreasing.

The last statement follows from the facts that J^* is ultimately monotonic (since $J^{*\prime}$ does not change sign) and $J^* > C$, $J^* \rightarrow 0$.

304 Estimate of $K^{*\prime}$:

The differential equation (104.3) for $\zeta_{l,k}$ and the definition (104.8) of $K^{*\prime}$ yield

$$K^{*\prime} = \sum_{k=1}^{N_l} m_{l,k} \zeta_{l,k} \times \zeta_{l,k}^{\prime\prime} = \sum_{k=1}^{N_l} m_{l,k} \zeta_{l,k} \times P_{l,k}. \quad (304.1)$$

Write, expanding in a binomial series,

$$|Z_{l,\lambda} - Z_{l,\lambda} + \zeta_{l,r} - \zeta_{l,s}|^{-3} = |Z_{l,\lambda} - Z_{l,\lambda}|^{-3} (1 + W_{l\lambda rs}), \quad (304.2)$$

where $W_{l\lambda rs}$ is an absolutely convergent series, each term of which contains a $\zeta_{p,q}$ or a component of a $\zeta_{p,q}$ as a factor; it is easily seen that the terms in $P_{l,k}$ not containing such a factor cancel out, i. e., $P_{l,k} = 0$ if all $\zeta_{p,q} = 0$, and we conclude that each term of $P_{l,k}$ contains a factor of the type mentioned above.

From the definition (104.7) of J^* and (104.14)

$$\zeta_{p,q} < cJ^{*\frac{1}{2}}, \quad \zeta_{p,q} \zeta_{r,s} < cJ^*, \quad (304.3)$$

and the same inequalities for the components of $\zeta_{p,q}$; hence,

$$|K^{*\prime}| < cJ^* \quad \text{and} \quad K^{*\prime} = J^* b. \quad (304.4)$$

Integrating from t to $t = 0$ and observing that (cf. (303.3))

$$J^* = J^*(t) = \max J^*(\tau), \quad \tau \in [t, 0[\quad (304.5)$$

we find

$$\underline{K}_l = \underline{K}_{l;0} + tJ^* \underline{b} \quad (304.6)$$

and

$$\underline{K}^* = \underline{K}_0^* + tJ^* \underline{b} \quad , \quad K_0^{*2} = K_0^{*2} + tJ^* \underline{b} \quad . \quad (304.7)$$

$$305 \quad \frac{1}{4} J^{*2} + K_0^{*2} \leq J^* (J^{**} + d) :$$

Using on

$$\frac{1}{2} J^{*2} = \sum_{\ell=1}^L \sum_{k=1}^{N_\ell} m_{\ell,k} \zeta_{\ell,k} \zeta'_{\ell,k}$$

the inequality

$$(\sum a_\nu b_\nu)^2 \leq (\sum a_\nu^2)(\sum b_\nu^2)$$

with

$$a_\nu = m_{\ell,k}^{\frac{1}{2}} \zeta_{\ell,k} \quad , \quad b_\nu = m_{\ell,k}^{\frac{1}{2}} \zeta'_{\ell,k} \quad ,$$

we find

$$\frac{1}{4} J^{*2} \leq J^* \sum \sum m_{\ell,k} (\zeta_{\ell,k} \zeta'_{\ell,k})^2 \zeta_{\ell,k}^{-2}$$

and similarly

$$K_0^{*2} \leq J^* \sum \sum m_{\ell,k} (\zeta_{\ell,k} \times \zeta'_{\ell,k})^2 \zeta_{\ell,k}^{-2} \quad .$$

Using $(ab)^2 + (a \times b)^2 = a^2 b^2$, addition yields

$$\frac{1}{4} J^{*2} + K_0^{*2} \leq J^* \sum \sum m_{\ell,k} \zeta_{\ell,k} \zeta'_{\ell,k} = 2 J^* V^* \quad (305.1)$$

(cf. (104.5), (104.13)) .

Substitute for K_0^{*2} from (304.7) and from (104.13): $V = V^* + V^0$,

from (104.14): $J = J^* + J^0$, and (102.9): $J^{**} = 2h + 2V$; then

$$\frac{1}{4} J^{*2} + K_0^{*2} \leq J^* (J^{**} - 2h + J^0 - 2V^0 - t \underline{b}) \quad ,$$

and since $-2h + J^{0''} - 2V^0 - tb$ is bounded on $[t, 0[$, we finally get

$$\frac{1}{4} J^{*''2} + K_0^{*2} \leq J^* (J^{*''} + d) \quad (305.2)$$

306 The inequality (305.2) implies that $K_0^* = 0$ and that $\lim J^{*''} J^{*''2} \geq 0$ exists:

define
$$Q = d J^{*''\frac{1}{2}} + \left(\frac{1}{4} J^{*''2} + K_0^{*2} \right) / J^{*''\frac{1}{2}} \quad ; \quad (306.1)$$

then

$$Q' = \frac{1}{2} J^{*''\frac{1}{2}} J^{*''} \left\{ d + J^{*''} - \left(\frac{1}{4} J^{*''2} + K_0^{*2} \right) / J^{*''} \right\} \quad ,$$

and since $\{..\} \geq 0$, we see that $J^{*''}$ and Q' are of the same sign, implying that Q ultimately cannot increase, since $J^{*''}$ ultimately decreases; hence, Q has a limit Q_0 as $t \rightarrow 0$, $0 \leq Q_0 < \infty$, and this implies because of $J^{*''} \rightarrow 0$ that

$$K_0^{*2} = 0 \quad \text{and} \quad (0 \leq) \lim J^{*''\frac{1}{2}} J^{*''2} (< \infty) \quad \text{exists} \quad . \quad (306.2)$$

§07 $J^{*''\frac{1}{2}} J^{*''} \geq c > 0$:

$J^{*''} = -2U^* + b$ implies

$$J^{*''\frac{1}{2}} J^{*''} = -2J^{*''\frac{1}{2}} U^* + J^{*''\frac{1}{2}} b \quad ; \quad (307.1)$$

now

$$J^{*''} > c \min \zeta_{l,k}^2 \quad , \quad J^{*''\frac{1}{2}} > c \min \zeta_{l,k} \quad , \quad -U^* > c (\min \zeta_{l,k})^{-1} \quad ,$$

therefore $J^{*''\frac{1}{2}} (-U^*) > c > 0$. The second term in (307.1) goes to zero, and the desired inequality

$$J^{*''\frac{1}{2}} J^{*''} \geq c > 0 \quad (307.2)$$

follows.

308 $J^{*\frac{1}{2}} J^{*\prime\prime} \geq c > 0$ and $\frac{1}{4} J^{*\prime\prime 2} \leq J^* (J^{*\prime\prime} + d)$ imply $\lim J^{*\frac{1}{2}} J^{*\prime\prime 2} > 0$:

define (cf. (306.1))

$$Q = d J^{*\frac{1}{2}} + \frac{1}{4} J^{*\frac{1}{2}} J^{*\prime\prime 2} ;$$

then

$$\begin{aligned} Q J^{*\frac{1}{2}} &= d J^* + \frac{1}{4} J^{*\prime\prime 2} \\ &= d J^* + \frac{1}{2} \int_0^t J^{*\prime\prime} J^{*\prime\prime} dt \quad , \quad \text{since } J^* \rightarrow 0 \text{ by (306.2)} . \end{aligned}$$

Substitute from (307.2), then

$$Q J^{*\frac{1}{2}} \geq d J^* + \frac{c}{2} \int_0^t J^{*\frac{1}{2}} J^{*\prime\prime} dt = d J^* + c J^{*\frac{1}{2}} ,$$

and this implies

$$Q \geq d J^{*\frac{1}{2}} + c ,$$

i. e.,

$$\lim Q = \frac{1}{4} \lim J^{*\frac{1}{2}} J^{*\prime\prime 2} = \mu^* > 0 \quad . \quad (308.1)$$

309 From (308.1) asymptotic expressions for J^* and $J^{*\prime\prime}$ as $t \rightarrow 0$ are easily derived:

(308.1) is equivalent to

$$J^{*\frac{1}{2}} J^{*\prime\prime 2} = 4\mu^* + e \quad , \quad e = e(t) \rightarrow 0 ; \quad (309.1)$$

separating and integrating we find

$$J^* = \left(\frac{3}{2} \mu^{*\frac{1}{2}} \right)^{\frac{4}{3}} t^{\frac{4}{3}} + t^{\frac{4}{3}} e$$

or

$$J^* \sim \left(\frac{9}{4} \mu^* \right)^{\frac{2}{3}} t^{\frac{4}{3}} \quad . \quad (309.2)$$

Substitution into (309.1) yields

$$J^{*\lambda} = (12\mu^{*2})^{\frac{1}{3}} t^{\frac{1}{3}} + t^{\frac{1}{3}} e$$

or

$$J^{*\lambda} \sim (12\mu^{*2})^{\frac{1}{3}} t^{\frac{1}{3}} .$$

(309.3)

310 $|J^{*\lambda}| < cJ^{*\lambda^{\frac{1}{2}}}$ ultimately:

differentiation of (303.1) yields

$$J^{*\lambda} = -2U^{*\lambda} - 2U^{*\lambda} - J^{*\lambda} ; \quad (310.1)$$

by (104.10) $J^{0\lambda}$ is a linear combination of terms $Z_{\lambda}^{\lambda} Z_{\lambda}^{\lambda}$ and $Z_{\lambda} Z_{\lambda}^{\lambda}$, Z_{λ}^{λ} , Z_{λ}^{λ} , and Z_{λ}^{λ} are bounded, and by the equation of motion (104.2) for Z_{λ}^{λ} , Z_{λ}^{λ} is a linear combination of the components of the $\zeta_{p,q}^{\lambda}$ with bounded coefficients, plus a bounded function. Since $|\zeta_{p,q}^{\lambda}| < cV^{*\frac{1}{2}}$ and this inequality holds also for the components of $\zeta_{p,q}^{\lambda}$, we get

$$|J^{0\lambda}| < cV^{*\frac{1}{2}} . \quad (310.2)$$

Similarly, U^{+} is a linear combination of the components of the $\zeta_{p,q}^{\lambda}$ with bounded coefficients, plus a bounded function, hence

$$|U^{+}| < cV^{*\frac{1}{2}} . \quad (310.3)$$

$U^{*\lambda}$ is a linear combination of terms $(\zeta_{p,q}^{\lambda} - \zeta_{p,s}^{\lambda})(\zeta_{p,q}^{\lambda} - \zeta_{p,s}^{\lambda})|\zeta_{p,q}^{\lambda} - \zeta_{p,s}^{\lambda}|^{-3}$,

this term's absolute value being $|\zeta_{p,q}^{\lambda} - \zeta_{p,s}^{\lambda}| |\zeta_{p,q}^{\lambda} - \zeta_{p,s}^{\lambda}|^{-2}$; since

$|\zeta_{p,q}^{\lambda} - \zeta_{p,s}^{\lambda}|^{-2} < cU^{*2}$, we have

$$|U^{*\lambda}| < cV^{*\frac{1}{2}} U^{*2} . \quad (310.4)$$

By means of the energy equation $U^{*} + V^{*} = h - U^{+} - V^{0}$ and the Lagrange-Jacobi equation (303.2): $J^{*\lambda} = -2U^{*} + b$, both V^{*} and U^{*}

can be replaced by $J^{*\nu}$, and we finally get the desired inequality

$$|J^{*\nu}| < c J^{*\nu \frac{3}{2}} \quad . \quad (310.5)$$

311 Asymptotic expression for $J^{*\nu}$. We prove $J^{*\frac{1}{2}} J^{*\nu} \sim \mu^*$:

$\frac{1}{4} J^{*\nu^2} \leq J^*(J^{*\nu} + d)$ (cf. (305.2), (306.2)) implies

$$\liminf J^{*\frac{1}{2}} J^{*\nu} \geq \frac{1}{4} \lim J^{*-\frac{1}{2}} J^{*\nu^2} = \mu^* \quad ; \quad (311.1)$$

it remains to be shown that

$$\limsup J^{*\frac{1}{2}} J^{*\nu} \leq \mu^* \quad . \quad (311.2)$$

Introduce

$$F = J^{*\nu^3} \quad ;$$

then (311.3)

$$F' = 3J^{*\nu^2} J^{*\nu} \quad , \quad F'' = 6J^{*\nu} J^{*\nu^2} + 3J^{*\nu^2} J^{*\nu} \quad ,$$

and as consequence of (309.3) we have

$$F \sim 12\mu^{*2} t \quad . \quad (311.4)$$

Consider now F' (≥ 0 obviously) instead of $J^{*\frac{1}{2}} J^{*\nu}$: we get

$$\begin{aligned} \liminf F' &= 3 \liminf (J^{*-\frac{1}{2}} J^{*\nu^2}) (J^{*\frac{1}{2}} J^{*\nu}) \\ &= 12\mu^* \liminf J^{*\frac{1}{2}} J^{*\nu} \geq 12\mu^{*2} \quad , \end{aligned} \quad (311.5)$$

and it remains to be shown that

$$\limsup F' \leq 12\mu^{*2} \quad . \quad (311.6)$$

Case 1: Assume that $\liminf F' = \limsup F' = \lim F' = d < \infty$; integration of this relation $F' \sim d$ to $F \sim dt$ yields by comparison with (311.4) $d = 12\mu^{*2}$, and the proof is complete.

Case 2: Assume that $\lim F' = +\infty$ and let $d > 0$ be arbitrarily large ; then ultimately $F' > d$ and by integration $F > dt$, contradicting (311.4).

Case 3: Assume finally that $\liminf F' < \limsup F'$, implying that $\overbrace{(\liminf F' < +\infty)}$.

There exist two positive numbers d_1 and d_2 and an infinite sequence of intervals $t_\alpha < t < t_\beta$, both t_α and t_β going to zero, such that

$$0 < 12L^{*2} < d_1 = F'(t_\alpha) < F'(t) < F'(t_\beta) = d_2, \quad t \in]t_\alpha, t_\beta[. \quad (311.7)$$

Write

$$F'' = \frac{1}{F} \left(\frac{2}{3} F'^2 + 3^{-\frac{3}{2}} F'^{\frac{5}{2}} \frac{J^{*''}}{J^{*'' \frac{5}{2}}} \right)$$

and use (310.5) and (311.4); then

$$|F''| < \frac{c}{t} \left(F'^2 + F'^{\frac{5}{2}} \right)$$

and on $]t_\alpha, t_\beta[$, since $F' \leq d_2$:

$$|F''| < \frac{c}{t}.$$

Integration over $[t_\alpha, t_\beta]$ yields

$$\begin{aligned} 0 < d_2 - d_1 = F'(t_\beta) - F'(t_\alpha) &= \int_{t_\alpha}^{t_\beta} F''(t) dt \leq \int_{t_\alpha}^{t_\beta} |F''(t)| dt \\ &< c \left| \log \frac{t_\beta}{t_\alpha} \right| = c \log \frac{-t_\alpha}{-t_\beta} \end{aligned}$$

(observe that $t_\alpha < t_\beta < 0$); this implies

$$\log \frac{-t_\alpha}{-t_\beta} > \frac{d_2 - d_1}{c}, \quad \text{i. e., ultimately } \frac{-t_\alpha}{-t_\beta} \geq T > 1. \quad (311.8)$$

From (311.7) it follows that

$$F(t_\beta) - F(t_\alpha) = \int_{t_\alpha}^{t_\beta} F'(t) dt > d_1 (t_\beta - t_\alpha),$$

implying

$$\frac{F(t_\beta)}{-t_\beta} + d_1 > \left(\frac{F(t_\alpha)}{-t_\alpha} + d_1 \right) \frac{-t_\alpha}{-t_\beta} ;$$

this last inequality is ultimately inconsistent, since

by (311.4) $\frac{F(t_\beta)}{-t_\beta}$ and $\frac{F(t_\alpha)}{-t_\alpha}$ both $\rightarrow -12\mu^{*2}$,

by (311.7) $12\mu^{*2} \neq d_1$, and

by (311.8) ultimately $\frac{-t_\alpha}{-t_\beta} > 1$.

Thus, $J^{*\frac{1}{2}} J^{*N} \sim \mu^*$ is true, and it follows that

$$J^{*N} \sim \left(\frac{2}{3} \mu^* \right)^{\frac{2}{3}} t^{-\frac{2}{3}} . \quad (311.9)$$

312 h_ℓ remains bounded as $t \rightarrow 0$: the energy h_ℓ of the ℓ -th cluster is by (104.5) and (104.6)

$$h_\ell = \frac{1}{2} \sum_{k=1}^{N_\ell} m_{\ell, k} \zeta_{\ell, k}^2 + U_\ell , \quad (312.1)$$

and by differentiation we get

$$h'_\ell = \sum_{k=1}^{N_\ell} m_{\ell, k} \zeta_{\ell, k} \zeta_{\ell, k}'' + U'_\ell ;$$

substitute from the equation of motion (104.3): then

$$h'_\ell = - \sum_{k=1}^{N_\ell} (\partial U_\ell / \partial \zeta_{\ell, k}) \zeta_{\ell, k}' + \sum_{k=1}^{N_\ell} m_{\ell, k} \zeta_{\ell, k} P_{\ell, k} + U'_\ell ,$$

the first and last term on the right cancel each other, thus

$$h'_l = \sum_{k=1}^{N_l} m_{l,k} \zeta'_{l,k} P_{l,k}$$

and by integration

$$h_l = \text{const.} + \sum_{k=1}^{N_l} m_{l,k} \int \zeta'_{l,k} P_{l,k} dt \quad . \quad (312.2)$$

Now $J'' \sim \left(\frac{2}{3} \right)^{\frac{2}{3}} t^{-\frac{2}{3}}$ implies $V^* < ct^{-\frac{2}{3}}$ and

$$|\zeta'_{l,k}| < c(-t)^{-\frac{1}{3}} \quad , \quad (312.3)$$

and since $P_{l,k}$ is bounded on $[t, 0[$, we get

$$\left| \int_t^0 \zeta'_{l,k} P_{l,k} dt \right| < c \left| \int_t^0 (-t)^{-\frac{1}{3}} dt \right| = ct^{\frac{2}{3}} \quad , \quad (312.4)$$

together with (312.2) implying the boundedness of h_l .

313 The Lagrange-Jacobi equation for J_l :

from the definition (104.7) we get

$$J''_l = 2 \sum_{k=1}^{N_l} m_{l,k} \zeta'_{l,k} \zeta'_{l,k} + 2 \sum_{k=1}^{N_l} m_{l,k} \zeta_{l,k} \zeta''_{l,k} \quad ,$$

and substituting V_l from (104.5) and for $\zeta''_{l,k}$ from the equation of motion (104.3),

$$J''_l = 4V_l - 2 \sum_{k=1}^{N_l} \zeta_{l,k} (\partial U_l / \partial \zeta_{l,k}) + 2 \sum_{k=1}^{N_l} m_{l,k} \zeta_{l,k} P_{l,k} \quad ;$$

on the right, the third term is bounded, and the second term equals $+2U_l$ (by Euler's theorem about homogeneous functions), and using $h_l = V_l + U_l$, where h_l is bounded (cf. 312), we find

$$J_l'' = -2U_l + b \quad . \quad (313.1)$$

As in (303.3) we conclude (observing that $J_l \rightarrow 0$ by the assumption of a collision singularity):

$$\begin{aligned} J_l'' &= -2U_l + b \quad \text{and} \quad U_l \rightarrow -\infty \quad \text{as} \quad t \rightarrow 0 \quad \text{imply:} \\ J_l'' &\rightarrow +\infty \\ J_l' &\text{ ultimately does not change sign (and is negative)} \\ J_l &\text{ is ultimately monotonically decreasing to zero.} \end{aligned} \quad (313.2)$$

314 Estimate of K_l (cf. 304) :

(304.1), together with

$$\zeta_{l,k} < c J_l^{\frac{1}{2}} \quad , \quad (314.1)$$

yields

$$|K_l'| < c J_l^{\frac{1}{2}} \quad \text{or equivalently} \quad K_l' = J_l^{\frac{1}{2}} b \quad , \quad (314.2)$$

and integrating from t to $t=0$, observing that by (313.2)

$J_l = J_l(t) = \max J_l(\tau)$, $\tau \in [t, 0[$, we have

$$K_l = K_{l;0} + J_l^{\frac{1}{2}} t b \quad \text{and} \quad K_l^2 = K_{l;0}^2 + J_l^{\frac{1}{2}} t b \quad . \quad (314.3)$$

315 $\frac{1}{4} J_l'^2 + K_{l;0}^2 \leq J_l (J_l'' + d)$:

analogous to the process used in 305 leading to (305.1), we arrive at

$$\frac{1}{4} J_l'^2 + K_l^2 \leq 2J_l V_l \quad ; \quad (315.1)$$

by (313.1): $2V_l = 2h_l - 2U_l = J_l'' + 2h_l - b$; using (314.3),

and observing that h_l is bounded, we get

$$\frac{1}{4} J_l'^2 + K_{l;0}^2 \leq J_l (J_l'' + d_1) + J_l^{\frac{1}{2}} (-t) d_2 \quad . \quad (315.2)$$

As in 307 for J^* , it can be shown that (313.1): $J_l'' = -2U_l + b$
 implies $J_l^{\frac{1}{2}} J_l'' \geq c > 0$, hence $J_l^{\frac{1}{2}} \leq c J_l J_l''$; substitute this into the
 last term of (315.2):

$$\frac{1}{4} J_l'^2 + K_{l;0}^2 \leq J_l (J_l'' + d_1 + c J_l'' \cdot (-t)) \quad (315.3)$$

Finally, with $J_l'' \leq J_l'' + c$, $J_l'' \leq c t^{-\frac{2}{3}}$ (cf. (311.9)):

$(-t) J_l'' \leq c (-t)^{\frac{1}{3}}$, and we get

$$\frac{1}{4} J_l'^2 + K_{l;0}^2 \leq J_l (J_l'' + d) \quad (315.4)$$

316 In full analogy to the development in 306 - 311, the following results
 can be derived:

$$K_{l;0} = 0 \quad \text{or} \quad K_{l;0} = 0 \quad ; \quad (316.1)$$

$$\lim J_l^{-\frac{1}{2}} J_l'^2 = 4\mu_l > 0 \quad \text{exists} \quad ; \quad (316.2)$$

$$J_l \sim \left(\frac{9}{4} \mu_l \right)^{\frac{2}{3}} t^{\frac{4}{3}}, \quad (316.3)$$

$$J_l' \sim (12\mu_l^2)^{\frac{1}{3}} t^{\frac{1}{3}}, \quad (316.4)$$

$$J_l'' \sim \left(\frac{2}{3} \mu_l \right)^{\frac{2}{3}} t^{-\frac{2}{3}}. \quad (316.5)$$

317 Direct proof of $K_{l;0} = 0$:

we had seen that (cf. (312.3))

$$|\zeta_{l,k}'| < c(-t)^{-\frac{1}{3}} \quad \text{or equivalently} \quad \zeta_{l,k}' = t^{-\frac{1}{3}} b, \quad (317.1)$$

and by integration from t to $t = 0$

$$\underline{r}_{\underline{m}l, k} = t^{\frac{1}{2}} \underline{b}_{\underline{m}} \quad (317.2)$$

results. Therefore

$$\underline{K}_{\underline{m}l} = \sum_{k=1}^{N_l} m_{l, k} \underline{r}_{\underline{m}l, k} \times \underline{r}_{\underline{m}l, k} = t^{\frac{1}{2}} \underline{b}_{\underline{m}} \quad , \quad (317.3)$$

i. e., $\underline{K}_{\underline{m}l} \rightarrow 0$ as $t \rightarrow 0$. (317.4)

(317.4) together with (314.3) implies

$$\underline{K}_{\underline{m}l} = J_l^{\frac{1}{2}} t \underline{b}_{\underline{m}} \quad \text{and} \quad \underline{K}_{\underline{m}l}^2 = J_l t^2 \underline{b}_{\underline{m}} \quad ,$$

leading to a simple derivation of the inequality (315.4).

318 Corollary. The estimate of $\underline{K}_{\underline{m}l}$ (cf. 314) can be developed in full analogy with that of $\underline{K}_{\underline{m}l}^*$ (cf. 304) : indeed,

$$\underline{P}_{\underline{m}l, k} = 0 \quad \text{if} \quad \underline{\zeta}_{\underline{m}l, \mu} = 0 \quad , \quad \mu = 1 \dots N_l \quad , \quad (318.1)$$

and writing

$$\underline{Z}_{\underline{m}l} - \underline{Z}_{\underline{m}l} = \underline{\zeta}_{\underline{m}l, \mu} + \underline{\zeta}_{\underline{m}l, \nu} = \underline{w}_{\underline{m}l, \mu} + \underline{\zeta}_{\underline{m}l, \nu}$$

and

$$|\underline{w}_{\underline{m}l, \mu} + \underline{\zeta}_{\underline{m}l, \nu}|^{-3} = \underline{w}_{\underline{m}l, \mu}^{-3} (1 + \underline{\zeta}_{\underline{m}l, \nu} \underline{R}_{\underline{m}l, \mu \nu}) \quad ,$$

where $\underline{R}_{\underline{m}l, \mu \nu}$ is bounded on $[t_0, t^*[$, t_0 sufficiently close to t^* ,

(318.1) shows that all terms in $\underline{P}_{\underline{m}l, k}$ not containing a factor $\underline{\zeta}_{\underline{m}l, \nu}$

or $\underline{\zeta}_{\underline{m}l, \nu}$ cancel out. Since

$$\underline{\zeta}_{\underline{m}l, \nu} < c J_l^{\frac{1}{2}} \quad ,$$

it follows that

$$|\underline{K}_{\underline{m}l}^*| < c J_l \quad \text{and} \quad \underline{K}_{\underline{m}l}^* = J_l \underline{b}_{\underline{m}} \quad ;$$

for the further steps see (304.4) ff. .

319 In 319 - 323 we will show that each cluster's configuration is close to a central configuration if $t < 0$ is close to the instant $t = 0$ of a collision singularity; more precisely: the N_ℓ bodies of the ℓ -th cluster are close to a central configuration in the N_ℓ -body problem.

We consider throughout the following an arbitrary, but fixed cluster, say the ℓ -th.

Let ψ be one of the components of $\zeta_{\ell, k}$; the conditions that the ℓ -th cluster form a central configuration are

$$J_\ell \frac{\partial U_\ell}{\partial \psi} = -\frac{1}{2} U_\ell \frac{\partial J_\ell}{\partial \psi} \quad \text{for all } \psi . \quad (319.1)$$

320 We introduce new dependent variables $\tilde{\zeta}_k$ by

$$\tilde{\zeta}_k = t^{-\frac{2}{3}} \zeta_{\ell, k} , \quad \tilde{\psi} = t^{-\frac{2}{3}} \psi ; \quad (320.1)$$

then

$$\tilde{\zeta}_k = t^{-\frac{2}{3}} \zeta_{l,k} \quad , \quad (320.2)$$

$$\tilde{J} = t^{-\frac{4}{3}} J_l \quad , \quad (320.3)$$

and

$$\tilde{U} = t^{\frac{2}{3}} U_l \quad . \quad (320.4)$$

The conditions (319.1) for a central configuration transform into

$$\tilde{J} \frac{\partial \tilde{U}}{\partial \tilde{\psi}} = -\frac{1}{2} \tilde{U} \frac{\partial \tilde{J}}{\partial \tilde{\psi}} \quad \text{for all } \tilde{\psi} \quad , \quad (320.5)$$

the Lagrange-Jacobi equation $J_l'' = -2U_l + b_l$ into

$$t^2 \tilde{J}'' + \frac{8}{3} t \tilde{J}' + \frac{4}{9} \tilde{J} = -2\tilde{U} + t^{\frac{2}{3}} b \quad (320.6)$$

and the equation of motion (104.3b) into

$$m_{l,k} \left(t^2 \tilde{\zeta}_{k}'' + \frac{4}{3} t \tilde{\zeta}_{k}' - \frac{2}{9} \tilde{\zeta}_{k} \right) = - \frac{\partial \tilde{U}}{\partial \tilde{\zeta}_{k}} + t^{\frac{4}{3}} m_{l,k} P_{l,k} ; \quad (320.7)$$

differentiating this last equation we get

$$\begin{aligned} m_{l,k} \left(t^3 \tilde{\zeta}_{k}''' + \frac{10}{3} t^2 \tilde{\zeta}_{k}'' + \frac{10}{9} t \tilde{\zeta}_{k}' \right) \\ = -t \frac{d}{dt} \frac{\partial \tilde{U}}{\partial \tilde{\zeta}_{k}} + t^{\frac{7}{3}} m_{l,k} \frac{d}{dt} P_{l,k} + \frac{4}{3} t^{\frac{4}{3}} m_{l,k} P_{l,k} . \end{aligned} \quad (320.8)$$

321 From (316.3) - (316.5) it follows that

$$t \tilde{J}' = - \frac{4}{3} t^{-\frac{4}{3}} J_l + t^{-\frac{1}{3}} J_l' \sim 0 \quad , \quad (321.1)$$

and similarly

$$t^2 \tilde{J}'' \sim 0 \quad ; \quad (321.2)$$

these relations together with (320.6) imply that

$$\frac{2}{9} J^{\sim} \sim -U^{\sim} \quad . \quad (321.3)$$

Substitute $\zeta_{\omega l, k}^{\sim} = t^{\frac{2}{3}} \zeta_{\omega k}^{\sim} + \frac{2}{3} t^{-\frac{1}{3}} \zeta_{\omega k}^{\sim}$ into the energy equation (312.1); we get

$$\frac{1}{2} \sum m_{l, k} (t \zeta_{\omega k}^{\sim}) (t \zeta_{\omega k}^{\sim}) + \frac{1}{3} t J^{\sim} + \frac{2}{9} J^{\sim} = t^{\frac{2}{3}} h_l - U^{\sim} \quad , \quad (321.4)$$

implying by (321.2) and (321.3) that

$$t \zeta_k^{\sim} \sim 0 \quad . \quad (321.5)$$

By (316.5) or (316.3) and (321.3) U^{\sim} is bounded, therefore

$\liminf \zeta_k^{\sim} > 0$; hence also $\frac{\partial U^{\sim}}{\partial \zeta_{\omega k}^{\sim}}$ is bounded, and from (320.7) it

follows that

$$t^2 \zeta_{\omega k}^{\sim} \text{ is bounded as } t \rightarrow 0 \quad . \quad (321.6)$$

In (320.8) the first and second term on the right are bounded since $t \zeta_{\omega k}^{\sim} \rightarrow 0$ by (321.5), and the third term is obviously bounded; therefore

$$t^5 \zeta_{\omega k}^{\sim} \text{ is bounded as } t \rightarrow 0 \quad . \quad (321.7)$$

322 (Tauberian) Lemma: If $tf^{\sim}(t)$ has a finite limit \mathfrak{Q} and $t^3 f^{\sim}(t)$ remains bounded as $t \rightarrow 0$, then $\lim t^2 f^{\sim}(t) = -\mathfrak{Q}$ as $t \rightarrow 0$. (cf. Wintner [2] pp. 279 - 280)

323 Applying 322 to (321.5), (321.6), and (321.7), we conclude that

$$t^2 \zeta_{\omega k}^{\sim} \sim 0 \quad ; \quad (323.1)$$

using this, together with (321.5) in the equation of motion (320.7), the asymptotic relation

$$-\frac{2}{9} m_{\ell, k} \zeta_{\omega k}^{\sim} \sim -\frac{\partial U^{\sim}}{\partial \zeta_{\omega k}^{\sim}} \quad \text{or} \quad -\frac{2}{9} m_{\ell, k} \psi^{\sim} \sim -\frac{\partial U^{\sim}}{\partial \psi^{\sim}} \quad (323.2)$$

results. The left side is $-\frac{1}{9} \frac{\partial J^{\sim}}{\partial \psi^{\sim}}$ and, multiplying both sides by $-J^{\sim} \neq 0$ and using (321.3), the desired result (cf. (320.5))

$$J^{\sim} \frac{\partial U^{\sim}}{\partial \psi^{\sim}} \sim -\frac{1}{2} U^{\sim} \frac{\partial J^{\sim}}{\partial \psi^{\sim}} \quad (323.3)$$

follows.

400 Simultaneous binary collisions

We shall consider the situation that there are L_1 pairs of bodies such that the two bodies in each pair collide at $t = 0$, while the centers of mass of the pairs and the remaining L_2 bodies approach distinct positions; thus

$$L_1 + L_2 = L = \text{number of clusters,}$$

$$2L_1 + L_2 = N = \text{number of bodies.}$$

401 Change to a new coordinate system such that the motion of the second body in each cluster is referred to its first body. Introduce the new coordinate


$$\xi_{ml} = \zeta_{ml,2} - \zeta_{ml,1} \quad ; \quad (401.1)$$

using $m_{l,1}\zeta_{ml,1} + m_{l,2}\zeta_{ml,2} = 0$ (cf. (103.3)), we find

$$m_{l,2}\xi_{ml} = -(m_{l,1} + m_{l,2})\zeta_{ml,1}$$

and

$$m_{l,1}\xi_{ml} = +(m_{l,1} + m_{l,2})\zeta_{ml,2} \quad . \quad (401.2)$$



402 The transformed equations of motion (104.3) read

$$\ddot{\xi}_{ml} = -\Gamma_l \frac{\xi_{ml}}{\xi_{ml}^3} + Q_{ml} \quad , \quad \Gamma_l = \gamma(m_{l,1} + m_{l,2}) \quad , \quad Q_{ml} = -\frac{m_{l,1} + m_{l,2}}{m_{l,2}} P_{ml} \quad (402.1)$$

the energy equation (104.6) changes into

$$J_l = V_l + \dots = \frac{m_{l,1} m_{l,2}}{m_{l,1} + m_{l,2}} \xi_l^2 - \gamma \frac{m_{l,1} m_{l,2}}{\xi_l} ; \quad (402.2)$$

observing that

$$J_l = \frac{m_{l,1} m_{l,2}}{m_{l,1} + m_{l,2}} \xi_l^2 , \quad (402.3)$$

the Lagrange-Jacobi equation reads

$$\left(\xi_l^2 \right)'' = \frac{2\Gamma_l}{\xi_l} + b \quad (402.4)$$

403 From the Lagrange-Jacobi equation (402.4) we derive now the asymptotic expression

$$\xi_l = \left(\frac{9}{2} \dots \right)^{\frac{1}{3}} t^{\frac{2}{3}} (1 + \dots b) \quad \text{as } t \rightarrow 0 : \quad (403.1)$$

Abbreviate temporarily

$$\xi_l^2 = \Xi_l ;$$

(313.2) implies that Ξ_l ultimately decreases monotonically to zero,

and (316.3) that

$$\Xi_l = bt^{\frac{2}{3}} . \quad (403.2)$$

From the energy expression (402.2) follows $\xi_l^2 \xi_l^{\prime 2} \xi_l^{\prime \prime 2} \rightarrow 0$, and since

$K_l \rightarrow 0$, we find $\Xi_l^{\prime} = \frac{1}{2} \xi_l \xi_l^{\prime} \rightarrow 0$ by means of the identity

$$K_l K_l + (\xi_l \xi_l^{\prime}) (\xi_l^{\prime} \xi_l^{\prime \prime}) = \xi_l^2 \xi_l^{\prime 2} \xi_l^{\prime \prime 2} ;$$

Ξ'_l is not identically zero since Ξ_l is not constant.

Multiply the Lagrange-Jacobi equation

$$\Xi'_l \ddot{\Xi}_l = \frac{2\Gamma_l}{\Xi_l^{3/2}} + b$$

by Ξ'_l and integrate from $t=0$ to t , $t < 0$; we get

$$\Xi_l'^2 = 8\Gamma_l \Xi_l^{-1/2} + \int_0^t \Xi_l'(\tau) b(\tau) d\tau \quad (403.3)$$

Taking into account that by (402.4) Ξ_l'' is ultimately positive, implying that Ξ_l' ultimately increases, i.e., does not change sign,

~~$\Xi_l'(\tau) = \Xi_l'(t) + \dots$~~

the last integral can be written as $b\Xi_l$; hence, by the first mean value theorem

$$\Xi_l'^2 = 8\Gamma_l \Xi_l^{-1/2} (1 + b\Xi_l^{1/2})$$

and substituting into the last term from (403.2), we get

$$\Xi_l'^2 = 8\Gamma_l \Xi_l^{-1/2} (1 + bt^{2/3})$$

yielding (403.1) by integration.

404 In order to get an asymptotic expression for ξ_{ml} as $t \rightarrow 0$, we first introduce a new independent variable s by

$$s = t^{\frac{1}{3}} \quad (404.1)$$

into the equations of motion (402.1); denoting differentiation with respect to s by a prime ' and substituting for ξ_l from (403.1), we get

$$s^2 \xi_{ml}'' - 2s \xi_{ml}' + 2\xi_{ml} = s^2 b \xi_{ml} + s^6 Q_{ml} \quad (404.2)$$

Introduce further new dependent variables u_{ml} by

$$\xi_{ml} = s u_{ml} \quad ; \quad (404.3)$$

it follows that $\xi_l = s u_l$, and since $\xi_l = s^2 b$ by (403.1), both u_l and \underline{u}_l remain bounded and \underline{u}_l can be written as $\underline{u}_l = s \underline{b}$.

The application of (404.3) transform (404.2) into

$$u_{ml}'' = b u_{ml} + s^3 Q_{ml} \quad ; \quad (404.4)$$

substitute $\underline{u}_l = s \underline{b}$ and integrate $u_{ml}'' = s \underline{b} + s^3 Q_{ml}$ twice; then

$$u_{ml} = s a_{ml} + s^3 b_{ml} \quad , \quad (404.5)$$

$$\xi_{ml} = s^2 a_{ml} + s^4 b_{ml} \quad , \quad (404.6)$$

and by comparison with (403.1)

$$a_{ml}^2 = \left(\frac{9}{2} \Gamma_l \right)^{\frac{2}{3}} \neq 0 \quad . \quad (404.7)$$

405 Without restriction of generality, we can assume that all (finitely many) components of the \underline{a}_l are different from zero; this can always be achieved by a constant rotation of the coordinates. We now transform the equations of motion (104.2) and (402.1) into a standard form of a system of differential equations of the first order, using again $s = t^{\frac{1}{3}}$ as the independent variable and introducing new dependent variables \underline{w}_l , $\underline{\omega}_l$, \underline{W}_l , and $\underline{\Omega}_l$ by

$$\underline{Z}_l = \underline{Z}_{l;0} + s^2 \underline{W}_l, \quad \underline{\Omega}_l = s \underline{W}'_l, \quad (405.1)$$

$$\xi_{l,j} = s^2 a_{l;j} (1 + s w_{l;j}), \quad \underline{\omega}_l = s \underline{w}'_l \quad j=1,2,3. \quad (405.2)$$

Since $\underline{Z}_l = \underline{Z}_{l;0} + s^3 \underline{Z}'_{l;0} + s^6 \underline{b}_l$, we have $\underline{w}_l \rightarrow 0$ as $s \rightarrow 0$, and (404.6) implies that $\underline{\omega}_l \rightarrow 0$.

Write for the further discussion (104.2) as

$$Z_{\underline{m}l}^{\prime\prime} = B_{\underline{m}l} \quad (405.3)$$

and (402.1) as

$$\xi_{\underline{m}l}^{\prime\prime} = -\Gamma_l \frac{\xi_{\underline{m}l}}{\xi_l^3} + b_{\underline{m}l} \quad , \quad (405.4)$$

where $B_{\underline{m}l}$ and $b_{\underline{m}l}$ are bounded as $s, t \rightarrow 0$ and can be expanded into power series of s , $W_{l;j}$, and $w_{l;j}$, which converge for sufficiently small absolute values of those quantities.

406 Transformation of (405.3) by (405.1) gives

$$\begin{aligned} s \Omega_{\underline{m}l}^{\prime} &= -\Omega_{\underline{m}l} + 2W_{\underline{m}l} + 9s^4 B_{\underline{m}l} \\ s W_{\underline{m}l}^{\prime} &= \Omega_{\underline{m}l} \quad . \end{aligned} \quad (406.1)$$

407 From the first equation of (405.2) we find

$$\xi_l^2 = s^4 \left(a_l^2 + 2s \chi_l + s^2 b \right) \quad , \quad (407.1)$$

where

$$\chi_l = a_{l;1}^2 w_{l;1} + a_{l;2}^2 w_{l;2} + a_{l;3}^2 w_{l;3} \quad ,$$

hence

$$\xi_l^{-3} = s^{-6} \frac{2}{9\Gamma_l} \left(1 - 3a_l^{-2} \chi_l + s^2 b \right) \quad . \quad (407.2)$$

Introduction of s and $w_{\underline{m}l}$ into (405.4) and substitution from (407.2) lead to

$$s^2 w_{l;j}^{\prime\prime} + 4s w_{l;j}^{\prime} = -2w_{l;j} + 6a_l^{-2} \chi_l + s b_{l;j}^* \quad , \quad (407.3)$$

where \underline{b}_l^* has the same properties as \underline{b}_l in 405. With \underline{w}_l (cf. (405.2)) we finally get the following system:

$$\begin{aligned} s \omega'_{l;j} &= -3 \omega_{l;j} - 2 w_{l;j} + 6 a_l^{-2} \chi_l + s b_{l;j}^* \\ s w'_{l;j} &= w_{l;j} \end{aligned} \quad j=1, 2, 3 \quad (407.4)$$

408 The full system of the transformed equations of motion (406.1) and (407.4) reads

$$\begin{aligned} s \Omega'_{l;j} &= -\Omega_{l;j} + 2 W_{l;j} + s B_{l;j}^* \\ s W'_{l;j} &= \Omega_{l;j} \end{aligned} \quad \begin{array}{l} j=1, 2, 3 \\ l=1 \text{ -- } L_1 + L_2 \end{array} \quad (408.1a)$$

$$\begin{aligned} s \omega'_{l;j} &= -3 \omega_{l;j} - 2 w_{l;j} + 6 a_l^{-2} \chi_l + s b_{l;j}^* \\ s w'_{l;j} &= w_{l;j} \end{aligned} \quad \begin{array}{l} j=1, 2, 3 \\ l=1 \text{ -- } L_1 \end{array} \quad (408.1b)$$

the nonlinear parts $s B_{l;j}^*$ and $s b_{l;j}^*$ are power series in all variables, converging if their absolute values are sufficiently small.

It is apparent that the matrix of the coefficients of the linear part on the right of (408.1) is in "block form"; consequently, the set of its elementary divisors is the union of the sets of the elementary divisors of the individual blocks (a block is the 6×6 matrix for a particular l in (408.1a) or (408.1b)).

The eigenvalues of a block in (408.1a) are $+1, +1, +1, -2, -2, -2$, and all elementary divisors are linear; similarly, the eigenvalues of

a block in (408.1b) are $+1, -1, -1, -2, -2, -4$, and again all elementary divisors are linear.

By a well-known theorem about systems of differential equations of the type (408.1) it follows that the solution with the property that all $W_{\mu l}$, $\Omega_{\mu l}$, $w_{\mu l}$, and $\omega_{\mu l}$ converge to zero as $s \rightarrow 0$ can be expanded into power series of s , which converge for sufficiently small $|s|$; the solution depends on $4L_1 + 3L_2$ parameters.

409 The theorem used in 408 has been proved by J. Horn: Ueber die Reihenentwicklung der Integrale eines Systems von Differentialgleichungen in der Umgebung gewisser singulärer Stellen. J. Reine Angew. Math. 116 (1896) 265 - 306 ; 117 (1897) 104 - 128 ; 254 - 266. For details and a survey of this and related problems see H. Dulac: Points singuliers des équations différentielles. Gauthier-Villars, Paris 1934. A modern simple proof (of a special case) can be found in Siegel [1] pp. 156 - 165.

410 Alternate proof of the boundedness of h_l . The principal tool for the study of the multiple binary collisions is the Lagrange-Jacobi equation (cf. (402.4)), enabling us to derive the asymptotic expression (403.1) for ξ_l ; its applicability, in turn, is based on the boundedness of the energy h_l as t approaches the instant of collision. We deduced this boundedness from general results about collisions (cf. 312). In the following we shall give a direct proof of the boundedness of h_l , independent of the previous general results; for the method see Arenstorf [3].

According to 318, Q_l in equation (402.1) can be written as $\xi_l b$, provided that ξ_l is sufficiently small, i. e., that t is sufficiently close to the instant of collision $t=0$. From

$$h_l' = \frac{m_{l,1} m_{l,2}}{m_{l,1} + m_{l,2}} \xi_l' Q_l \quad (\text{cf. 312})$$

$$(410.1)$$

$$= \xi_l \xi_l' b$$

we get

$$h_l'^2 \leq \xi_l^2 b^2 \xi_l' \xi_l' \quad (b^2, \text{ to indicate } \underline{\text{positive}} \text{ bounded function})$$

and substituting for $\xi_l' \xi_l'$ from (402.2),

$$h_l'^2 \leq \xi_l b^2 (\xi_l h_l + \gamma m_{l,1} m_{l,2})$$

and because of $\xi_l \rightarrow 0$:

$$h_l'^2 \leq 1 + |h_l| \quad (\text{for sufficiently small } \xi_l).$$

Denote by $H_\ell(t_1)$ the maximum of $|h_\ell|$ on $[t_0, t_1]$; then by integration from t_0 to t and then to t_1 , $t_0 < t \leq t_1$:

$$|h_\ell(t) - h_\ell(t_0)| \leq \left(1 + H_\ell(t_1)\right)^{\frac{1}{2}}(t - t_0) \leq \left(1 + H_\ell(t_1)\right)^{\frac{1}{2}}(t_1 - t_0),$$

hence

$$|h_\ell(t)| \leq |h_\ell(t_0)| + \left(1 + H_\ell(t_1)\right)^{\frac{1}{2}}(t_1 - t_0) \quad \text{on } [t_0, t_1]$$

and

$$H_\ell(t_1) = \max_{t \in [t_0, t_1]} |h_\ell(t)| \leq |h_\ell(t_0)| + \left(1 + H_\ell(t_1)\right)^{\frac{1}{2}}(t_1 - t_0).$$

For nonnegative a, b, x the inequality $x \leq a + b(1 + x)^{\frac{1}{2}}$ implies $x \leq a + \frac{1}{2}b^2 + b(1 + a + \frac{1}{4}b^2)^{\frac{1}{2}} \leq a + b + ba^{\frac{1}{2}} + b^2$; applying this, we find

$$H_\ell(t_1) \leq |h_\ell(t_0)| + (t_1 - t_0) \left(1 + |h_\ell(t_0)|^{\frac{1}{2}} + (t_1 - t_0)\right),$$

and it is obvious that the right side remains bounded as $t_1 \rightarrow 0$.

411 We finally formulate the following

Theorem: Let all variables be real and let the solution of the N-body problem be holomorphic on $[t_0, 0[$, $t_0 < 0$; let there be at $t = 0$ a collision singularity such that several binary collisions (and only these) take place at this instant. Then the position vectors, distances, and the products $t^{\frac{1}{3}} \cdot (\text{velocity vector})$ can be expanded into power series of $t^{\frac{1}{3}}$. Therefore, the solution can be analytically continued in the real beyond $t = 0$ into $t > 0$.

Bibliography on the singularities
of the equations of motion of celestial mechanics

This bibliography is intended to be a complete listing of papers on the title subject; the major part of the papers deals with the classical N-body problem (with newtonian gravitational forces acting on particles), but publications on the following topics have been included: regularization of the equations of motion, non-newtonian forces, and central configurations and homographic solutions, these last ones in view of their importance for the collision problem.

The only publications known to me as containing more than just a few references are (Sokolov [30],) Szebehely [13], Vernić [3], and Wintner [2]. Being a first attempt at collecting the pertinent literature, this bibliography will have its share of errors and omissions; particularly, no systematic search could be made for the Russian literature. Also, since to many of the included papers I do not have access, there will be listings which are only marginally or not at all connected with the topics intended to be covered. No attempt has been made to include all publications on collision and regularization in (1) the two-body problem and (2) in Euler's problem of two fixed centers; likewise, papers only applying regularizing transformations to other problems have not been listed systematically.

An effort has been made to provide references to review journals; the publications and the abbreviations used are:

- AJ Astronomischer Jahresbericht [1899 -]
- JFM Jahrbuch über die Fortschritte der Mathematik [1868 - 1944]
- MR Mathematical Reviews [1940 -]
- Z' Zentralblatt für Mathematik und ihre Grenzgebiete [1931 -];

- A 2 -

other review publications (and most of JFM) are not available to me at present. The abbreviations of names of journals have been chosen as much as possible as in MR 32 (1966).

An asterisk * preceding the page number of a reference to a review publication indicates that the paper is listed only, but not reviewed. A tilde ~ indicates that the name of the journal is the same as in the foregoing reference.

I shall be grateful for any comments as to errors, omissions, etc., that will lead to an improvement of this bibliography. In particular, I would appreciate receiving reprints of new papers dealing with the singularity problem and/or related topics.

Agostinelli, Cataldo

- [1] Sul problema dei tre corpi.
Rend. Sem. Mat. Fis. Milano 21 (1950) 165-195 (1951)
AJ51 *101 MR13 698 ZM44 388
- [2] Il problema dei tre corpi dall'epoca di Lagrange ai giorni nostri.
Atti Convegno Lagrangiano (Torino, 1963). (Atti Accad. Sci. Torino 98 (1963/64), suppl.) 304-326 Accad. Sci., Torino 1964
MR32 110 #621

Altavista, C. A.

- [1] Some considerations on the homographic solutions of the three body problem.
As. Argentina Astr. Bol. Nr. 4 93 pp 52-57.
Actas de la Cuarta Reunión Científica de la Asociación, realizada en el Observatorio Astronómico de Córdoba, los días 20, 21 y 22 de Octubre de 1961 (ed. C. Jaschek, C. Hernandez, L. Milone).
AJ62 *204, *82(title)
- [2] Homographic solutions in the n body problem with an arbitrary law of attraction.
As. Argentina Astr. Bol. Nr. 6 117 pp 94.
Actas de la sexta reunión científica de la Asociación Argentina de Astronomía, realizada en la Universidad de Tucumán (Horoco Molle) los días 22, 23, 24 y 25 de Mayo de 1963.
AJ64 *197

d'Ambly, Carl-Cünter

- [1] Kollineare Zentralkonfigurationen des n-Körperproblems.
Astronom. Nachr. 288 (1964) 81-85
AJ64 197/198 MR30 852 #4574

Amoroso, L.

- [1] La soluzione del problema dei tre corpi secondo Sundman.
Boll. della Matheſis 6 (1914) 14-22
AJ16 *96 JFM45 1174; *1182

Andoyer, H.

- [1] Sur l'équilibre relatif de n corps.
Bull. Astronom. 23 (1906) 50-59
AJ8 169
- [2] Sur les solutions périodiques voisines des positions d'équilibre relatif, dans le problème des n corps.
Bull. Astronom. 23 (1906) 129-146.

Arenstorff Richard F. = Schulz-Arenstorff, Richard F.

- [1] [Schulz-Arenstorff]
On the representation of the solutions of the restricted problem of three bodies by power series.
Army Ballistic Missile Agency, Redstone Arsenal, Alabama.
Report DC-TR-1-59 March 30, 1959 29pp
- [2] New Regularization of the Restricted Problem of Three Bodies.
Astronom. J. 68 (1963) 548-555
AJ63 181/182
- [3] Regularization theory for the elliptic restricted three body problem.
George C. Marshall Space Flight Center, Huntsville, Alabama.
R-COMP-IN-6-68 July 1, 1968 57 pp (to be published in J. Differential Equations)

Armellini, G.

- [1] Sul moto di un punto attratto da più centri fissi.
Atti Accad. Naz. Lincei. Rend. Cl. Sci. Fis. Mat. Natur.
(5) 22₁ (1913) 672-678
AJ15 162/163 JFM44 827/828
- [2] Sopra la soluzione delle equazioni differenziali dei moti di un punto attratto da più centri fissi posti in linea retta.
~ (5) 23₁ (1914) 770-776
AJ16 97
- [3] Estensione della soluzione del Sundman dal caso di corpi ideali al caso di sferrette elastiche omogenee.
~ (5) 24₁ (1915) 184-190
AJ17 62/63 JFM45 1176
- [4] Sur la solution analytique du problème restreint des trois corps.
C. R. Acad. Sci. Paris 158 (1914) 253-255
AJ15 163/163 AJ16 97 JFM45 1175
- [5] Un théorème général sur le problème des n corps.
~ 158 (1914) 680-683
AJ16 96/97 JFM45 1175
- [6] Ricerche sopra la previsione dell'urto nel problema dei tre corpi.
Atti Accad. Naz. Lincei. Rend. Cl. Sci. Fis. Mat. Natur.
(5) 27₁ (1919) 87-91
AJ21 *120 JFM46 1375

Banachiewicz, Thadee

- [1] Sur un cas particulier du problème des n corps.
C. R. Acad. Sci. Paris 142 (1906) 510-512
AJ8 170 JFM37 739

Bardon, Jose Robla

- [1] Regularización del problema plano de los tres cuerpos.
Rev. Ci. (Lima) 38 (1937) No. 420 99-133
AJ40 *73 JFM63 1295 ZM18 280

Bartky, Walter

- [1] = MacMillan, W. U. [2]

Bashenow, G. M. = Baženov, G. M.

- [1] Einige Spezialfälle des n-Körperproblems.
Charkow Buli. Nr. 6 (1946) 3-5
AJ46 56

Belorizky, David

- [1] Sur la convergence des séries dans la solution du problème des trois corps donnée par M. Sundman.
C. R. Acad. Sci. Paris 193 (1931) 314-317
AJ33 *56 ZM2 212
- [2] Sur la solution du problème des trois corps, donnée par M. Sundman.
~ 193 (1931) 766-768
AJ34 *42 ZM3 134
- [3] Sur l'application des méthodes de M. Sundman aux problèmes de la mécanique céleste.
~ 193 (1931) 1321-1323
AJ34 42 ZM3 234
- [4] Application pratique des méthodes de M. Sundman a un cas particulier des trois corps.
Bull. Astronom. (2) I 6 (1931) 417-434
AJ33 *56
- [5] Sur la nature des chocs dans le problème des trois corps à trois degrés de liberté.
C. R. Acad. Sci. Paris 194 (1932) 769-771
AJ34 *42 ZM4 373
- [6] Sur le rayon de convergence des séries dans le problème de deux corps, traité par la méthode de Levi-Civita.
~ 194 (1932) 1449-1452
AJ34 *42 ZM5 31
- [7] Sur l'intégration du problème des deux corps.
Bull. Sci. Math. (2) 57 (1933) 45-49
JFM59 812 ZM7 35
- [8] Recherches sur l'application pratique des solutions générales du problème des trois corps.
These Paris 1933 Gauthier-Villars 73 pp
AJ35 49

- [9] Recherches sur l'application pratique des solutions générales du problème des trois corps.
J. des Observateurs 16 (1933) 109-132; 149-172; 189-211
AJ35 49
- [10] Sur la stabilité dans le problème des trois corps.
Bull. Sci. Math (2) 62 (1938) 271-288
ZM22 414
- [11] Les points singuliers dans le problème restreint des trois corps.
C. R. Acad. Sci. Paris 207 (1938) 1382-1384
AJ40 *73 ZM20 *175
- [12] Chocs triples imaginaires dans le problème plan des trois corps.
~ 208 (1939) 558-560
ZM21 165
- [13] Les chocs dans le problème des trois corps et dans l'espace.
~ 208 (1939) 966-969
ZM21 165
- [14] Chocs d'une nouvelle espèce dans le problème des trois corps.
~ 213 (1941) 558-560
AJ43 *51 MR5 79 ZM26 76
- [15] Les chocs triples imaginaires et la régularisation des intégrales dans le problème des trois corps.
J. des Observateurs (Marseille) 25 (1942) 81-127
AJ46 56/57
- [16] Sur deux cas particuliers du problème des trois corps.
C. R. Acad. Sci. Paris 223 (1946) 193-196
AJ46 *57 MR8 291
- [17] Sur la régularisation du problème des trois corps.
~ 231 (1950) 1428-1429
AJ50 91 MR19 550 ZM40 395
- [18] Sur la régularisation du problème des trois corps par la variable de Sundman.
Bull. Astronom. 16 (1952) 327-331
AJ52 88 MR14 802
- [19] Sur la régularisation du mouvement dans le problème restreint.
Notes Travaux Section Astr. Inst. Math. Beograd 2 (1958) 7-10
AJ58 127
- [20] La régularisation du mouvement dans le problème général des trois corps par la variable de Sundman.
C. R. Acad. Sci. Paris Ser. A 262 (1966) 1133-1136

Benedikt, Elliot T.

- [1] Collision trajectories in the three-body problem.
J. Astronaut. Sci. 6 (1959) 17-24, 30.
AJ59 75; *145

Bilimowitsch, Anton D. = Bilimovitch = Bilimović

- [1] Einige partikuläre Lösungen des Problems der n Körper.
Astronom. Nachr. 189 (1911) c181-c186
AJ13 142
- [2] O nekim specijalnim slučajevima problema n tela.
(Über einige Spezialfälle des Problems der n Körper)
Glas Srpske kr. akademije nauka (Berichte der Serbischen Kgl.
Akademie der Wissenschaften. Belgrad) 59 Erste Abt. (1927) 17-43
AJ31 46 JFM54 832
- [3] [Bilimovitch, A.; Petronievitch, B.]
Beitrag zur elementaren Lösung der zwei speziellen Fälle des
Dreikörperproblems.
Astronom. Nachr. 258 (1936) 209
AJ38 73/74 ZM13 325
- [4] Über die Rolle der Archimedischen gleichseitigen Polyeder im
 n -Körperprobleme.
Bull. Acad. Serbe Sci. Arts Cl. Sci. Math. Natur. Sci. Math.
A. 7 (1941) 197-212 (15 plates)
AJ50 *91 MR10 746
- [5] Über einen Sonderfall des Vierkörperproblems.
~ A. 7 (1941) 213-227
AJ50 *91 MR10 746

Birkhoff, George David

- [1] The restricted problem of three bodies.
Rend. Circ. Mat. Palermo 39₁ (1915) 265-334
= Collected Works 1 682-751
- [2] Recent advances in dynamics.
Science N. S. 51 (1920) 51-55 = Collected Works 2 106-110
- [3] Dynamical Systems.
American Math. Soc. 1927 (revised ed. 1966)
- [4] Sur le problème restreint des trois corps. I, II.
Ann. Scuola Norm. Sup. Pisa I: (2)4 (1935) 267-306
II: (2)5 (1936) 1-42 (ZM: 9-50) = Collected Works 2 466-
505; 668-709
I: JFM61 1480-1483 ZM12 128/129 II: ZM14 84/85

Bisconcini, Giulio

- [1] Sul problema dei tre corpi. Condizioni d'urto di due di essi.
Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.
(5) 122 (1903) 552-557
AJ6 181 JFM34 769
- [2] Sur le problème des trois corps. Trajectoires le long desquelles
deux au moins des trois corps se choquent. Conditions qui
entraînent un choc.
Acta Math. 30 (1905/06) 49-92

Block, Henrik

- [1] Sur les chocs dans le problème des trois corps.
Arkiv for Matematik, Astronomi och Fysik vol 5 no 9 (1909) 32 pp
= Lunds Medd. Nr. 42
AJ10 194
- [2] Sur une classe de singularités dans le problème de n corps.
Lunds Meddel. Sér. II, Nr. 6, Lund 1909 57 pp
AJ11 266

Boas, Ralph Philip, Jr.

- [1] A Tauberian Theorem Connected with the Problem of Three Bodies.
Amer. J. Math. 61 (1939) 161-164
ZM20 17

Bohlin, K.

- [1] Note sur le problème des deux corps et sur une intégration nouvelle dans le problème des trois corps.
Bull. Astronom. 28 (1911) 113-119
AJ13 141/142

Bourget, H.

- [1] Le problème des trois corps.
L'Astronomie. Bulletin de la société astronomique de France.
Paris 28 (1914) 65-69
AJ16 96

Breglia, Emilia

- [1] Su alcuni casi particolari del problema dei tre corpi.
Giorn. Mat. Battaglini 54 (1916) 151-173
AJ18 169

Brendel, M.

- [1] Neue Methode zur Berechnung der Störungen.
Astronom. Nachr. 224 (1925) c229-c236
AJ27 55
- [2] Über die Lagrangeschen strengen Lösungen des Dreikörperproblems.
Monatsh. Math. Phys. 43 (1936) 298-304
AJ38 74 ZM14 233

Brocard, H.

- [1] Pour la bibliographie du problème des trois corps.
Interméd. des math. 27 (1919) 99
JFM47 835

Broucke, Roger A.

- [1] Recherches d'orbites périodique dans le problème restreint plan (système Terre-Lune).
Dr. Sci. dissertation, Univ. of Louvain 1962
- [2] = Deprit, André [2]
- [3] Regularizations of the Plane Restricted Three-Body Problem.
Jet Propulsion Laboratory, California Institute of Technology
Pasadena, Calif., February 28, 1964 14pp
- [4] Regularizations of the plane restricted three-body problem.
Icarus 4 (1965) 8-18
AJ65 196 MR30 852 #4575
- [5] Symmetrization of the Two-Body Problem.
Jet Propulsion Laboratory, Space Programs Summary, No. 37-33,
Vol. IV. June 30, 1965
April 1 - May 31, 1965 23-28,32
- [6] Regularized Special Perturbation Techniques Using Levi-Civita
Variables.
American Institute of Aeronautics and Astronautics
Aerospace Sciences Meeting, 3rd, New York, N. Y. Jan. 24-26,
1966. ALAA Paper No. 66-8 11pp
- [7] On changes of Independent Variable in Dynamical Systems and
Applications to Regularization.
Icarus 7 (1967) 221-232

Brumberg, W. A. (V. A.)

- [1] Constant configurations in the problem of four bodies and their
stability. (Russian, English sum.)
Astronom. Zh. 34 (1957) 55-74
AJ57 125 MR19 1141
- [2] Trajectories of double collision in the restricted problem of
three bodies. (Russian; English sum.)
Bull. des Instituts für theoret. Astron., Leningrad 7(1960)
833-843
AJ60 188

Bruns, H.

- [1] Bemerkungen zur Theorie der allgemeinen Störungen.
Astronom. Nachr. 109 (1884) c215-c222
JFM16 1100

Buchanan, D.

- [1] A new isosceles triangle solution of the three-body problem.
Trans. Amer. Math. Soc. 16 (1915) 259-274
AJ18 168/169 JFM45 1177
- [2] Asymptotic isosceles triangle solutions for unequal masses.
Rend. Circ. Mat. Palermo 62 (1938/39) 239-248
AJ43 50 MR1 124

Burdet, Claude Alain

- [1] = Stiefel, Eduard [6]
- [2] Regularization of the Two Body Problem.
Z. Angew. Math. Phys. (ZAMP) 18 (1967) 434-438
- [3] Theory of Kepler Motion: The General Perturbed Two Body Problem.
Dissertation No. 4010 Swiss Federal Institute of Technology
Zurich 1968
= Z. Angew. Math. Phys. (ZAMP) 19 (1968) 345-368

Burrau, Carl

- [1] Recherches numériques concernant des solutions périodiques d'un cas spécial du problème des trois corps. I, II.
Astronom. Nachr. I: 135 (1894) No. 3230 c233-c240
II: 136 (1894) No. 3251 c161-c174
JFM25 1872/1873
- [2] Über einige in Aussicht genommene Berechnungen, betreffend einen Specialfall des Dreikörperproblems.
Vierteljschr. Astronom. Ges. 41 (1906) 261-266
AJ8 170 JFM37 968

Carathéodory, Constantin

- [1] Über die strengen Lösungen des Dreikörperproblems.
Bayer. Akad. Wiss. Math.-Natur. Kl. S.-B. 257-267 (1933)
= Mathematische Schriften 2 387-396
JFM 592 1614 ZM8 134

Cesco, R. P.

- [1] Sobre las soluciones homográficas del problema de los tres cuerpos. (English summary)
Observ. astron. Univ. nac. La Plata, Ser. astron. 25, Nr. 2, 11p (1959)
AJ60 188 MR28 193 #988 ZM113 392
- [2] On the homographic solutions of the three-body problem.
As. Argentina Astr. Bol. Nr. 2 23
Actas de la Segunda reunión científica de la Asociación Argentina de Astronomía, realizada en el Observatorio Astronómico de La Plata, los días 23, 24 y 25 de Setiembre de 1959
AJ60 *188, *77(title)
- [3] Some theorems and results in the three-body problem.
Proc. Intern. Meeting Problems Astrometry and Celestial Mech., La Plata 1960. Astron. Obs., Natl. Univ. of La Plata, Argentina 1961. 81-92

- [4] On the exact solutions of the problem of three bodies.
Univ. Nac. Tucumán Rev. Ser. A 14 (1962) 253-261
= Obs. Astr. La Plata Separata Astr. Nr. 31
AJ62 *205 MR31 825 #4589 7M132 218
- [5] On the general solution of the problem of three bodies.
Information Bull. Southern Hemisph. Nr. 2 27-28 (review of a
presentation given at Stockholm in Aug. 1962)
AJ62 *205
- [6] Sobre la solución general del problema de los tres cuerpos.
Actas de la quinta reunión científica de la Ascciación, realizada
en el Observatorio Astronómico de San Juan los días 11, 12, y 13
de julio 1962. As. Argentina Astr. Bol. Nr. 5, 75pp 13
AJ63 *182, *79(title)
- [7] Sobre la solución general del problema de los tres cuerpos.
Obs. Astr. Univ. Nacional La Plata (Astr.) 25 (1963) Nr. 4 5-30
(5-6 English sum.)
AJ63 *182 ZM118 403

Chazy, Jean

- [1] Sur certaines trajectoires du problème des n corps.
C. R. Acad. Sci. Paris 157 (1913) 688-691
AJ15 164 JFM44 1075/1076
- [2] Sur les points singuliers de l'intégrale générale du problème
des n corps.
~ 157 (1913) 1398-1400
AJ15 164 JFM44 1076
- [3] Sur certaines trajectoires du problème des n corps.
Bull. Astronom. 35 (1918) 321-389
AJ20 93
- [4] Remarque sur le problème de deux corps et de trois corps.
C. R. Acad. Sci. Paris 168 (1919) 81-83
AJ21 117 JFM47 845
- [5] Sur les solutions du problème des trois corps où les trois
corps forment un triangle isoscèle.
~ 169 (1919) 526-529
AJ21 117 JFM47 845
- [6] Sur les singularités impossibles du problème des n corps.
~ 170 (1920) 575-577
AJ22 101
- [7] Sur l'allure du mouvement dans le problème des trois corps
quand le temps croît indefiniment.
~ 170 (1920) 1560-1563
AJ22 101 JFM47 845
- [8] Sur la stabilité à la Poisson dans le problème des trois corps.
~ 173 (1921) 517-519
- [9] Sur la stabilité dans le problème des trois corps.
~ 173 (1921) 576-578

- [10] Sur l'allure du mouvement dans le problème des trois corps quand le temps croît indéfiniment.
Ann. Sci. École Norm. Sup. (3) 39 (1922) 29-130
- [11] Sur l'arrivée dans le système solaire d'un astre étranger.
Proceedings of the International Mathematical Congress
Toronto 1924. Vol. II (1928) 19-21
JFM54 1022
- [12] Sur le problème rectiligne des trois corps.
Bull. Soc. Math. France 55 (1927) 222-268
JFM53 892
- [13] Sur l'allure finale du mouvement dans le problème des trois corps.
J. Math. Pures Appl. (9) 8(1929) 353-380
JFM55 567
- [14] Sur la stabilité à la Poisson dans le problème des trois corps.
Atti del Congresso Internazionale dei Matematici Bologna
1928 (VI) Vol 5 (1931) 15-16
JFM57₁ 1023
- [15] Sur les multiplicités singulières du problème des trois corps.
Bull. Sci. Math. (2) 56 (1932) 79-104
ZM4 165
- [16] Sur l'allure finale du mouvement dans le problème des trois corps.
Bull. Astronom. (Mém. Var.) (2) 8 (1932) 403-436
AJ36 51 JFM58 1267
- [17] Sur les solutions d'Euler et de Lagrange du problème des trois corps.
C. R. Acad. Sci. Paris 224 (1947) 1677-1680
AJ47 *41 MR9 161 ZM29 79
- [18] L'intégration du problème des trois corps par Sundman et ses conséquences.
Bull. Astronom. 16 (1952) 175-190
AJ52 *89 MR14 *95

Chilmi, H. F. = Hil'mi, G. F.

Choudry, Ram Kishore

- [1] Existence of Analytical Continuation and Its Parametric Representation for $\mu \neq 0$ Corresponding to the Circular Orbits for $\mu = 0$ in the Restricted Problem of Three Bodies in Three Dimensional Coordinate System.
J. Indian Math. Soc. new series 28 (1964) 119-138
- [2] Birkhoff's Rings in the Restricted Problem of Three Bodies in a Three-Dimensional Co-ordinate System.
Proc. Indian Acad. Sci. Sect. A 62 (1965) 252-270

Clavo Rivera, Carlos A.

- [1] Sobre la regularizacion del problema piano de los tres cuerpos.
(Spanish)
Rev. Ci.(Lima) 49 (1947) 3-69
AJ47 *41 AJ48 76/77 AJ49 *89 MR9 161 ZM30 223

Crenna, Mario

- [1] Esposizione elementare di un caso notevole del problema ristretto dei tre corpi.
Period. Mat. (4) 23 (1943) 28-33
AJ45 69 MR8 *98

Czuchry, Andrew J.

- [& Pitkin, Edward T.]
[1] Regularization in Optimal Trajectory Problems.
AIAA J. 6 (1968) 1209-1210

Delie, André

- [1] = Deprit, A. [1]
- [2] = Deprit, A. [2]
- [3] = Deprit, A. [4]

Deprit, André

- [1] [& Delie, A.]
Systèmes matériels de trois masses ponctuelles.
Mouvement plan en coordonnées symétriques.
Ann. Soc. Sci. Bruxelles Ser. I 75 (1961) 5-44
MR23B 521 #B2991 ZM101 420
- [2] [& Delie, A.]
Courbure de la métrique riemannienne du problème plan des trois corps.
Acad. Roy. Belg. Bull. Cl. Sci (5) 48 (1962) 652-658
AJ62 *206 MR26 1125 #5962
- [3] [& Broucke, R.]
Régularisation du Problème Restreint Plan des Trois Corps par Représentations Conformes.
Icarus 2 (1963) 207-218
MR28 193 #987 AJ63 *183 ZM134 440
- [4] [& Delie, A.]
Régularisation du problème des trois corps.
Arch. Rational Mech. Anal. 12 (1963) 325-353
AJ63 *183 MR26 1125/1126 #5963 ZM113 393

Dramba, Constantin

- [1] Sur les singularités du problème restreint des trois corps.
C. R. Acad. Sci. Paris 202 (1936) 1736-1738
AJ38 *75 ZM14 333

- [2] Sur la régularisation du problème restreint des trois corps pour les valeurs de l'excentricité, voisines de l'unité.
(2. Congr. Interbalkan. des Math., Bucarest, 12. IX. 1937)
Bull. Math. Soc. Roumaine Sci. 40 (1938) no. 112 277-280
AJ40 *74 ZM20 175
- [3] Sur les singularités réelles et imaginaires dans le problème des trois corps.
Bull. Math. Soc. Roumaine Sci. Vol 41 no. 2 (1939) 21-75
JFM65 908 ZM20 407/408
- [4] Singularités imaginaires du problème isoscèle plan des trois corps.
C. R. Acad. Sci. Paris 210 (1940) 131-133
AJ42 *51 MR1 183/184 ZM23 273
- [5] Singularités imaginaires du problème isoscèle des trois corps.
~ 210 (1940) 393-395
AJ42 *51 MR2 24/25 ZM23 407
- [6] Le problème plan de Lagrange et les chocs triples imaginaires.
Bull. Math. Soc. Roumaine Sci. 43 (1941) 3-6
MR7 90/91
- [7] Les chocs triples imaginaires dans le problème général des trois corps.
Mathematica, Timișoara 22 (1946) 74-80
AJ46 *57 MR8 291
- [8] Les chocs triples imaginaires dans le problème rectiligne des trois corps.
Acad. Roum. Bull. Sect. 29 (1946) 79-82
AJ46 *58 MR10 156 ZM61 *418
- Dziobek, Otto
[1] Über einen merkwürdigen Fall des Vielkörperproblems.
Astronom. Nachr. 152 (1900) c33-c46
AJ2 197
- Ebert, W.
[1] Sur un système d'équations différentielles qui équivant au problème des n corps, mais admet une intégral de plus.
C. R. Acad. Sci. Paris 131 (1900) 251-253
JFM31 890
- Egerváry, E.
[1] On a generalization of the Lagrangian solution of the problem of three bodies.
C. R. Acad. Sci. USSR (NS) 55 (1947) 793-795 (AJ48: 805-807)
AJ47 *42 AJ48 77 MR9 211 ZM29 234/235

Euler, Leonhard

- [1] Theoria motuum planetarum et cometarum. (Berlin 1744)
Opera omnia ser. 2 vol 28 105-251
- [2] De motu rectilineo trium corporum se mutuo attrahentium.
Novi commentarii academiae scientiarum Petropolitanae
11 (1765) 17-20, 144-151
= Opera Omnia ser. 2, 25 281-289

Fieber, H.

- [1] Zur Frage der Lagrangeschen Lösungen beim n-Körperproblem.
Z. Angew. Math. Mech. 39 (1959) 398-399
ZM99 398/399
- [2] Zu den Lagrangeschen Lösungen des n-Körperproblems.
Monatsh. Math. 64 (1960) 1-14
AJ60 *610 MR22 1564 #9260 ZM99 398
- [3] Zu den Lagrangeschen Lösungen des n-Körperproblems. II.
~ 67 (1963) 7-11
AJ63 *84 MR26 866 #4523 ZM113 393

Finlay-Freundlich, E. = Freundlich, Erwin

Frادlin, B. N.

- [1] = Sokolov [30]

Fransén, A. E.

- [1] Ett specialfall af tre-Kroppars-problemet.
Ofversigt af Kongl. Vetenskaps-Akademiens Förhanlingar 52
(1895) 783-805
JFM26 1098

Freundlich, Erwin

- [1] Über die singulären Stellen der Lösungen des n-Körper-Problems.
(Preussische) Akademie der Wissenschaften, Berlin.
Sitzungsberichte 168-188 (1918)
AJ20 93
- [2] Über ein Paradoxon in den Jacobischen Vorlesungen über Dynamik.
Astronom. Nachr. 208 (1918/19) c209-c212
- [3] [Finlay-Freundlich, E.]
Celestial Mechanics.
Pergamon Press. New York London 1958 150pp

García, Godofredo

- [1] [& Rosenblatt, A.]
Sobre el problema de los tres cuerpos en el plano.
Rev. Ci. (Lima) 38 (1937) no. 419 57-94
AJ39 *63 JFM63 1295 ZM17 137
- [2] [& Rosenblatt, A.]
Sur la régularisation du problème plan des trois corps.
C. R. Acad. Sci. Paris 204 (1937) 1711
AJ39 *63 ZM16 180

- [3] On the regularisation of the plane problem of three bodies.
Actas Ac. Ci. Lima 5 (1942) 101-113
= Rev. Ci. (Lima) 44 (1942) 471-483
= Rev. Ci. (Lima) 45 (1943) 53; 91
MR4 174
- [4] On a new form of the inequalities of K. Sundman in the problem
of three bodies. (Spanish)
Actas Acad. Ci. Lima 5 (1942) 117-121
MR4 174
- [5] Generalization of the formula of Lagrange. Generalization of
Sundman's inequality, deduced from the homogeneity of the
potential and Birkhoff's auxiliary function for the problems
of n bodies and of infinitely many bodies (gravitational or
nebulous gas). (Spanish)
~ 5 (1942) 132-144
MR4 174
- [6] Generalization of Sundman's fundamental equality to the case of
more than three bodies.
Proc. Nat. Acad. Sci. U.S.A. 28 (1942) 425-427
MR4 57
- [7] Generalization of the inequality of Sundman to the case of more
than three bodies and to the case of a gravitational gas.
~ 28 (1942) 428-432
MR4 57
- [8] The three-body problem. (Spanish)
Rev. Ci. (Lima) 44 (1942) 159-186
MR4 57
- [9] On the problem of n bodies and the motion of a gravitational
gas (or a nebulous one). (Spanish)
~ 44 (1942) 457-469
MR4 174
- [10] On the regularization of the two dimensional three-body problem.
(Spanish)
~ 44 (1942) 471-483
MR5 17
- [11] On a generalization of a new inequality derived from K. Sundman's
auxiliary function in the problem of three bodies generalized
by George D. Birkhoff. (Spanish)
Actas Acad. Ci. Lima 6 (1943) 3-11
MR4 174
- [12] On the regularization of two dimensional three-body problem.
(Spanish)
Publ. Inst. Math. Univ. Nac. Litoral 5 (1943) 221-230
MR6 75; 334(correction) MR7 620

- [13] Generalization of the formula of Lagrange. Generalization of Sundman's inequality, deduced from the homogeneity of the potential and Birkhoff's auxiliary function for the problems of n bodies and of infinitely many bodies (gravitational or nebulous gas). (Spanish)
Rev. Ci. (Lima) 45 (1943) 3-15
MR5 17
- [14] On the regularization of the two dimensional three-body problem I. (Spanish)
~ 45 (1943) 53-90
MR5 17
- [15] On the regularization of the two dimensional three-body problem. II. Expansion into a series of the regularizing coordinates and of the time as a function of the auxiliary variable of K. Sundman and T. Levi-Civita. (Spanish)
~ 45 (1943) 91-~~93~~ 133
MR5 17
- [16] On the regularization of the two dimensional three-body problem. III. Conditions for collision in the three-body problem. (Spanish)
~ 45 (1943) 9pp
MR5 17
- [17] On the regularization of the two dimensional three-body problem. IV. Analytic continuation beyond the collision of two of the bodies. Orbits in the neighborhood of a collision of two of the bodies. (Spanish)
~ 45 (1943) 13pp
MR5 17
- [18] Generalization of the Lagrange-Birkhoff inequality for a dissipative system in an approximately Newtonian field. (Spanish)
Actas Acad. Ci. Lima 7 (1944) 431-434
MR7 92
- [19] Celestial mechanics and its evolution. (Spanish)
Rev. Ci. (Lima) 45 (1944) 145-164
MR6 *75
- [20] Generalization of the theory of the virial. (Spanish)
~ 45 (1944) 281-292
= Actas Acad. Ci. Lima 7 (1944) 225-236
MR6 190 MR7 620
- [21] The generalization of the equality of Lagrange and the inequality of Sundman in the case of more than three bodies. (Spanish)
Facultad de Ingeniería Montevideo. Publ. Inst. Math. Estadística 1, 129-136 = Bol. Fac. Ingen. Montevideo 3 (Año 10) (1946) 177-184
AJ46 *58 MR8 495 ZM60 *465

Giacaglia, Giorgio E. O.

- [1] = Szebehely, Victor [6]
- [2] Periodic Orbits of Collision in the Restricted Problem of Three Bodies.
Astronom. J. 72 (1967) 386-391
- [3] Regularization of the Restricted Problem of Four Bodies.
~ 72 (1967) 674-678

Gylden

- [1] Über eine Methode die Störungen eines Kometen vermittelst rasch konvergierender Ausdrücke darzustellen.
Bull. de l'Acad. Imp. des Sciences de S.-Petersburg 14 (1869)
195-231
JFM2 852/853

Hadamard, Jacques Salomon

- [1] Sur un mémoire de M. Sundman.
Bull. Sci. Math. (2) 39 (1915) 249-264
AJ18 *170 JFM45 1171-1172

Hamy, Maurice

- [1] Sur le calcul approché des inégalités d'ordre élevé.
C. R. Acad. Sci. Paris 147 (1908) 1251-1256
JFM39 998
- [2] Sur une propriété de l'équation obtenue en égalant à zéro la distance de deux planètes P, P_1 qui ne se rencontrent pas en des points réels.
~ 193 (1931) 1357-1360
ZM3 380
- [3] Sur une propriété de l'équation obtenue en égalant à zéro la distance de deux planètes P, P_1 qui ne se rencontrent pas en des points réels.
~ 194 (1932) 146-149
ZM3 380
- [4] Sur l'équation obtenue en égalant à zéro la distance de deux planètes qui ne se rencontrent pas en des points réels.
~ 194 (1932) 329-333
ZM3 380

Happel, H.

- [1] Das Dreikörperproblem.
K. F. Koehler Verlag. Leipzig 1941 526pp
MR15 652/653 ZM25 23

Haviland, E. K.

- [1] On the singularities and the qualitative behavior of the solutions of the problem of central forces.
Amer. J. Math. 65 (1943) 477-491
MR5 16

Heinrich, Wladimír Wáclaw

- [1] Sur une transformation des théories classiques des perturbations, permettant d'aborder l'étude du choc planétaire. Mem. Soc. roy. des lettres et sciences, classe des sciences 1946 (XII) 30pp
= Publ. Inst. Astr. Univ. Charles de Prague (2) Nr. 21
AJ46 59
- [2] Sur la régularisation du choc planétaire. Calcul des séries. ~ (XIII) 20pp
= ~ (2) Nr. 22
AJ46 59

Herrick, Samuel

- [1] Universal Variables.
Astronom. J. 70 (1965) 309-315
AJ65 205

Hil'mi, G. F. = Chilmi, H. F. = Khilmi, G. F.

- [1] The virial theorem in celestial mechanics. (Russian)
Doklady Akad. Nauk SSSR (N.S.) 70 (1950) 393-396.
MR11 466
- [2] Qualitative methods in the n-body problem. (Russian)
Moscow 1958
- [3] [Chilmi, H. F.]
Qualitative Methoden beim n-Körperproblem der Himmelsmechanik.
Akademie-Verlag Berlin 1961
- [4] [Khilmi, G. F.]
Qualitative Methods in the Many Body Problem.
Gordon and Breach New York 1961

Hinrichsen, J. J. L.

- [1] On the problem of n bodies. (abstract)
Bull. Amer. Math. Soc. 36 (1930) 205
JFM56₁ *786
- [2] On the problem of n bodies.
Trans. Amer. Math. Soc. 36 (1934) 306-314
AJ36 *52 ZM9 234
- [3] Bounds for the libration points in a restricted problem of n-bodies. (abstract)
Bull. Amer. Math. Soc. 48 (1942) 45-46
- [4] The libration points in an n-body problem.
Amer. Math. Monthly 50 (1943) 231-237
MR4 227

Hintze, K. H.

- [1] Zur Regularisierung des doppelten Zweierstosses.
Monatsb. Deutsch. Akad. Wiss. Berlin 7 (1965) 179-181
= Mitt. Sternw. Babelsberg Nr. 26
AJ65 *199

Holm, S.

- [1] Om den liksidiga-triangel-lösningen av trekroppars problemet.
Populär Astronomisk Tidsskrift (Stockholm) 8 (1928) 15-21
AJ30 53

Holtom, Carl

- [1] Permanent configurations in the n-body problem.
Trans Amer. Math. Soc. 54 (1943) 520-543
MR5 79

Hopfner, F.

- [1] Über eine Verallgemeinerung der relativen kanonischen
Koordinaten von Jacobi.
Astronom. Nachr. 195 (1913) c257-c262
AJ15 162 JFM44 1073

Hoppe, R.

- [1] Erweiterung der bekannten Speciallösung des Dreikörperproblems.
Archiv der Mathematik und Physik 64 (1879) 218-233
JFM11 650

Iglesias Garrido, Tomás

- [1] Eine Verallgemeinerung der Lagrangeschen partikularen Lösungen
des Dreikörperproblems. (Spanish)
Revista Mat. Hisp.-Amer. (4) 11 (1951) 266-276
AJ51 *102 MR14 324 ZM44 388

Ingelgem, R. van

- [1] Le problème des trois corps.
Ciel et Terre 42 (1927) 128-138
= Louvain, Publ. du Laboratoire d'Astronomie 3 (1927) 267-279
AJ29 54

Izsak, J.

- [1] Zur Regularisierung des Einzentrumproblems.
Mitt. Sternw. Ungar. Akad. Wiss. Budapest-Szabadsaghegy
Nr. 39 3-5 (1956)
AJ56 122

Jablonski, E.

- [1] Contribution à l'étude du cas le plus général du choc dans un
système de points matériel soumis à la loi de Newton.
C. R. Acad. Sci. Paris 164 (1917) 995-997
AJ19 113

Kaplan, Wilfred

- [1] Topology of the Two-Body problem.
Amer. Math. Monthly 49 (1942) 316-323
MR3 280

Karamata, Jovan

- [1] Über eine Tauberschen Satz im Dreikörperproblem.
Amer. J. Math. 61 (1939) 769-770
MR1 11

Kiveliiovitch, (Michel?)

- [1] Sur les conditions d'un choc binaire dans le problème des trois corps.
C. R. Acad. Sci. Paris 182 (1926) 116-118
AJ28 45
- [2] Le calcul des conditions du choc binaire dans le problème des trois corps.
~ 182 (1926) 263-264
- [3] Sur les orbites périodiques du problème des trois corps avec chocs de deux corps.
~ 184 (1927) 1533-1535
AJ29 54
- [4] Sur le problème des trois corps avec chocs successifs d'un corps avec deux autres.
~ 186 (1928) 361-362
AJ30 *55 JFM54 1022
- [5] Sur les chocs dans le problème d'un corps attiré par deux centres fixes.
~ 193 (1931) 513-516
- [6] Sur les vitesses nulles dans le problème des trois corps.
~ 194 (1932) 1317-1318
AJ34 43 ZM4 331
- [7] Sur quelques cas particuliers du problème des trois corps avec chocs.
~ 195 (1932) 763-765
AJ34 43 ZM6 228
- [8] Sur les points singuliers du problème des trois corps.
Bull. Astronom. (Mém. Var.) (2) 7 (1932) 75-127
AJ34 43 JFM58 1392
- [9] Étude sur les chocs des corps qui s'attirent proportionnellement à la distance.
~ (2) 7 (1932) 415-417
AJ35 50 JFM58 1393
- [10] Régularisation des chocs binaires pour des forces proportionnelles à l'inverse d'une puissance quelconque de distance.
Atti. Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (6)
16 (1932) 227-230
ZM6 228

- [11] Sur les orbites du problème des trois corps.
Bull. Astronom. (Mém. Var.) (2) 8 (1933) 71-75
AJ35 *50
- [12] Le nombre des chocs dans le problème des n corps qui s'attirent
inversement à une puissance quelconque de la distance.
Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (6)
17 (1933) 140; 224-227
AJ35 *50 JFM59 815 ZM7 181

Klemperer, W. B.

- [1] Rosette configurations of gravitating bodies in homographic
equilibrium.(abstract)
Astronom. J. 65 (1960) 492
AJ60 *191
- [2] Some Properties of Rosette Configurations of Gravitating
Bodies in Homographic Equilibrium.
~ 67 (1962) 162-167
AJ62 207

Kobold, H.

- [1] Eine Bemerkung zur Störungstheorie.
Astronom. Nachr. 272 (1942) 212-213
MR5 17 ZM60 461

Kurth, Rudolf

- [1] On Lagrange's triangular solution of the problem of three
bodies.
Arch. Math. 8 (1957) 381-392
= Astr. Contr. Univ. Manchester (3) No. 51
AJ57 127 MR19 1141 ZM81 178
- [2] Introduction to the Mechanics of the Solar System.
Pergamon Press. New York London 1959 177pp

Kusnetzoff, W. = Kuznecov, V.

- [1] Sulla regolarizzazione del problema generale dei tre corpi.
Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.
(6) 11 (1930) 568-574
AJ32 47 JFM56₁ 779
- [2] Sur une équation du problème des trois corps analogue à l'équa-
tion de Képler du problème des deux corps.
Stud. Cerc. Astronom. 9 (1964) 171-174
AJ64 *201

Kustaanheimo, Paul

- [1] Über die Bestimmung der oskulierenden Bahn eines künstlichen
Erdsatelliten.
Comment Phys.-Math. Soc. Fennica 26, No. 1 22pp (1961)
ZM102 401

- [2] Spinor regularization of the Kepler motion.
Ann. Univ. Turku. Ser. AI 73 (1964) 7pp
= Publication No. 102 of the Astronomical Observatory,
Helsinki, Finland
AJ64 *201 MR30 852 #4571
- [3] Relativistic spinor regularization of the Kepler motion.
Comment. Phys.-Math. Soc. Fennica 30 No. 9 4 pp (1964)
= Publ. Astr. Obs. Helsinki No. 103
AJ64 *201
- [4] [& Stiefel, E.]
Perturbation theory of Kepler motion based on spinor
regularization.
J. Reine Angew. Math. 218 (1965) 204-219
= Publ. Astr. Obs. Helsinki No. 110
AJ65 *200 MR31 824 #4584

Kuznecov, V. = Kusnetzoff, W.

Lagrange, J. L.

- [1] Essai sur le problème des trois corps.(1772)
Oeuvres 6 229-324

Lahaye, Edmond

- [1] Sur la résolution du problème des trois corps par des itérations
intégrales convergentes réelles.
Acad. Roy. Belg. Cl. Sci. (Mém. Coll. in-4°) (2) 11 (1937)
No. 6 67pp
JFM63 732
- [2] Le choc quadruple dans le problème des n corps.
Acad. Roy. Belg. Bull. Cl. Sci. (5) 29 (1943) 576-590
MR7 224
- [3] Les chocs binaires simultanés réels dans le problème des N corps.
~ (5) 32 (1946) 678-693
AJ47 *42 MR9 211
- [4] Les chocs triples réels dans le problème des N corps.
~ (5) 33 (1947) 89-104
AJ47 *42 MR9 211
- [5] La régularisation du mouvement dans le problème des N corps
et les itérations intégrales convergentes.
~ (5) 33 (1947) 652-671
MR9 626

Lampariello, G.

- [1] Intorno alle soluzioni stazionarie del problema dei tre corpi.
Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.
(6) 29 (1939) 298-303
ZM22 33

Lehmann-Filhés, R.

- [1] Ueber zwei Fälle des Vielkörperproblems.
Astronom. Nachr. 127 (1891) c137-c144
JFM23 1222

Leimanis, Eugène

- [1] [& Minorsky, N.]
Dynamics and Nonlinear Mechanics.
Wiley, New York 1958
AJ58 *130 MR20 476 #2877 ZM81 179/180
- [2] Qualitative methods in general dynamics and celestial mechanics.
Appl. Mech. Rev. 12 (1959) 665-670
AJ60 *610 MR22 *1523 #7847

Lemaître, G.

- [1] Coordonnées symétriques dans le problème des trois corps.
Acad. Roy. Belg. Bull. Cl. Sci. (5) 38 (1952) 582-592; 1218-1234
AJ52 91 MR14 910
(I) ZM48 177 (II) ZM49 247
- [2] Régularisation dans le problème des trois corps.
~ (5) 40 (1954) 759-767
AJ54 104/105 MR16 964/965 ZM57 161
- [3] Regularization of the Three-Body Problem.
Vistas in Astronomy Vol 1 207-215
Pergamon Press 1955
- [4] The Three Body Problem.
NASA CR-110, October 1964, 40 pp

Lense, Josef

- [1] Über algebroiden und algebromorphe Funktionen.
Bayer. Akad. Wiss. Math.-Natur. Kl. S -B. (1956) 237-242
MR19 257

Levi-Civita, T.

- [1] Sur les trajectoires singulières du problème restreint des trois corps.
C. R. Acad. Sci. Paris 136 (1903) 82-84
= Opere Matematiche II, 271-274
JFM34 768
- [2] Condition du choc dans le problème restreint des trois cc.ps.
~ 136 (1903) 221-223
= Opere Matematiche II, 275-277
AJ5 204 JFM34 768/769
- [3] Traiettorie singolari ed urti nel problema ristretto dei tre corpi.
Ann. Mat. Pura Appl. (3) 9 (1903) 1-32
= Opere Matematiche II, 279-308
AJ5 204 JFM34 769

- [4] Sur la résolution qualitative du problème restreint des trois corps.
Verhandl. des III Internat. Math.-Kongresses Heidelberg, 1904
402-408
= Opere Matematiche II, 411-417
- [5] Sur la résolution qualitative du problème restreint des trois corps.
Acta Math. 30 (1906) 305-327
= Opere Matematiche II, 419-439
JFM37 *739
- [6] Sulla regolarizzazione del problema piano dei tre corpi.
Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.
(5^a) 24₂(1915) 61-75; 235-248
= Opere Matematiche III, 477-493
AJ17 65 JFM45 1172
- [7] Sul problema piano dei tre corpi. I-III.
Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.
(5^a) 24₂ (1915) 421-434; 485-501; 553-569
= Opere Matematiche III, 511-525; 526-544; 545-564
AJ17 65/66 JFM45 1172-1174
- [8] Sopra due trasformazioni canoniche desunte dal moto parabolico.
Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.
(5^a) 25₁ (1916) 446-458
= Opere Matematiche III, 573-587
JFM46 1373
- [9] Sur la régularisation du problème des trois corps.
C. R. Acad. Sci. Paris 162 (1916) 625-628
= Opere Matematiche III, 589-593
AJ18 170/171 JFM46 1373
- [10] Sur la régularisation du problème des trois corps.
Acta Math. 42 (1920) 99-144
= Opere Matematiche IV, 217-264
- [11] Die Regularisierung des Dreikörperproblems und ihre Tragweite.
in: Fragen der klassischen und relativistischen Mechanik
1. Vortrag (1921) 1-26 1924 Springer-Verlag
- [12] Sur les chocs dans le problème des trois corps.
Comptes rendus du 2^{me} Congrès International de Mécanique
Appliquée (Zürich 1926) Zürich, Orell Füssli 1927, 96-106
= Opere Matematiche IV, 417-432

Lévy, Jacques

- [1] Les approches dans le problème des trois corps.
Thèse Paris 1943 Gauthier-Villars, Paris 1943 59pp
AJ46 *60 MR8 98

Lewis, D. C.

- [1] Rejection to Infinity in the Problem of Three Bodies When the Total Energy is Negative.
in: First Compilation of Papers on Trajectory Analysis and Guidance Theory NASA SP-141 (1967) 73-96.
- [2] Comments on the Sundman Inequality
in: Second Compilation of Papers on Trajectory Analysis and Guidance Theory, NASA PM-67-21 (1968) 129-139

Lindemann, F.

- [1] Über die Bewegung von Massenpunkten, die dem Newtonschen Anziehungsgesetze unterworfen sind (Problem der n Körper).
Bayer. Akad. Wiss. Matn.-Natur. Kl. Abh. Neue Folge Heft 28 (1935) 1-31
AJ37 *61 JFM12 1484/1485 ZM11 373

Lindow, M.

- [1] Ein Nomogramm zur Thieleschen Transformation.
Astronom. Nachr. 214 (1921) c263-c264
AJ23 *111
- [2] Eine Transformation für das Problem der $n+1$ Körper.
~ 219 (1923) c141-c154
AJ25 51/52
- [3] Der Kreisfall in Problem der $3+1$ Körper.
~ 220 (1924) c369-c380 + 2 plates
AJ26 54
- [4] Der Kreisfall in Problem der $n+1$ Körper.
~ 228 (1926) c233-c248
AJ28 46

Longley, W. R.

- [1] Some particular solutions in the problem of n bodies.
Bull. Amer. Math. Soc. 13 (1906/07) 324-335

Lovett, E. O.

- [1] Special Periodic Solutions of the Problem of n Bodies.
Science N.S. 17 (1903) 107
AJ5 203

MacMillan, William Duncar

- [1] A proof of Wilczynski's Theorem.(abstract)
Bull. Amer. Math. Soc. (2) 19 (1912/13) 300-301
AJ18 178 JFM44 *841
- [2] Permanent configurations in the problem of four bodies.
Trans. Amer. Math. Soc. 34 (1932) 838-875
AJ35 *50 ZM5 418
- [3] Dynamics of rigid bodies.
McGraw-Hill 1936. Dover Publ., New York, reprint 1960.

Mangad, Moshe

- [1] Regularized Solutions of the Two-Body Problem and the Restricted Problem.
Astronom. J. 72 (1967) 467-471

Maikoff, André, Jr. = Markov, A. jr.

- [1] La loi des dix tiers et la classification des chocs dans le problème des trois corps.
Byulleten' Astronomicheskogo Instituta no. 14 (1927) 159-160
AJ28 45/46
- [2] The law of ten thirds and the classification of collisions in the general problem of three bodies. (Russian; English sum.)
J. Soc. Phys. Leningrad 22 (1929) 81-97
JFM 57 1550 *im*
- [3] Über einige Bewegungsfälle ~~im~~ Dreikörperproblem.
Bulletin de l'Institut astronomique (Leningrad) Nr. 21,
2 (1928) 5-10
AJ30 54

Marcolongo, R.

- [1] Le recenti ricerche di K. Sundman sul problema matematico dei tre corpi.
Giorn. Mat. Battaglini 52 (1914) 171-186
AJ16 *95 JFM45 *1174
- [2] Il problema dei tre corpi da Newton (1686) ai nostri giorni.
Atti Accad. Sci. Fis. Mat. Napoli (2) 16 no. 6
= Manuali Hoepli 1919 166pp
= Nuovo Cimento (6) 9 (1915) 309-367; 10 (191?) 89-130
AJ17 64/65 AJ18 *170; *178 JFM45 1174 JFM47 834

Marković, Ž.

- [1] Duh novijih istraživanja u problemu triju tijela.
Glasnik Hrv. Prir. Društva 31 (1919) 1-24

Matukuma, T.

- [1] Singular points in the problem of two bodies.
Tokyo Phys. Math. Soc. Proc. Ser. 2, 9 (1917) 116-128
AJ19 112

Meffroy, Jean

- [1] Sur un nouveau cas particulier du problème des trois corps.
C. R. Acad. Sci. Paris 254 (1962) 818-820
AJ62 208 MR24B 182 #B1200 ZM102 400/401
- [2] Sur un cas particulier du mouvement rectiligne du problème des n corps.
~ 255 (1962) 477-479
AJ62 208 MR26 671 #3505 ZM114 152

Mendes, M.

- [1] Recherches sur le problème des n corps à masses variables.
Ann. Fac. Sc. Univ. Toulouse III.s. 27 1-170 (1935)
AJ38 *77 JFM612 1485 ZM15 183/184

Merman, G. A.

- [1] On Chazy's investigations in the problem of three bodies.
(Russian)
Byull. Inst. Theoret. Astronom. 5 (1954) 594-605
MR17 905
- [2] Zu Arbeiten von R. Vernic über die Regularisierung und die periodischen Lösungen des Dreikörperproblems. (Russian, German sum.)
~ 6, no. 6 (1956) 408-415
(English translation: NASA N67 13573 December 5, 1966)
MR19 227
- [3] Recherches qualitatives dans le problème de trois corps.
(Russian, French sum.)
~ 6 no. 10 (1958) 687-712
AJ58 130 MR20 940 #5684 MR21 1440 #7786
- [4] Sur la représentation de la solution générale du problème des trois corps par des séries convergentes. (Russian, French sum.)
~ 6 no. 10 (1958) 713-732
AJ58 130 MR20 940 #5685 MR21 1440/1441 #7787

Mihajlović, Dobrivoje = Mihailović, D. = Michailovitch, D.

- [1] Les intégrales premières dans le problème d'entrechoc de trois corps. (Serbian; Russian, French sum.)
Vesnik Društva mat. fiz. Srbije 1 no. 1 (1949) 45-61
AJ49 *88 MR11 465, 872 (corr) ZM39 *402
- [2] Sur les directions des mouvements instantanés relatifs dans le problème d'entre-choc de trois corps de la mécanique céleste. (Serbian; Russian, French sum.)
~ 1 no. 2 (1949) 11-16
AJ49 *88 MR11 465, 872(corr.) ZM35 114
- [3] Réduction du système des équations différentielles du mouvement dans un cas spécial du problème des trois corps. (Serbo-Croatian; French sum.)
~ 2 no. 1-2 (1950) 81-97
MR12 550 ZM39 402
- [4] Interprétation de quelques résultats dans un cas spécial du problème des trois corps. (Serbo-Croatian; French sum.)
~ 2 no. 3-4 (1950) 27-34
AJ52 *93 MR14 101 ZM54 76

- [5] Contribution a l'étude d'un problème particulier de n corps.
(Serbian; 33-34 French sum.)
Vesnik Društva mat. fiz. Srbije 3 no. 1-2 (1951) 11-34
MR14 101 ZM43 390
- [6] Généralisation de certains résultats dans le problème du choc
de trois corps. (Serbo-Croatian; French sum.)
Univ. Beogradu. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.
no. 12 (1957) 32 pp
AJ58 *131 MR20 258 #1576 ZM77 374

Milanković, M. = Milankowitha, M. = Milankovitsch, M.

- [1] Nebaska mehanika. (Celestial mechanics) (Serbian)
Beograd 1935 333pp
AJ38 *77 JFM612 1480 ZM14 233

Mittag-Leffler, G.

- [1] see: Weierstrass, Karl

Montille, H.

- [1] Bibliographie du problème des trois corps.
Interméd. des Math. (2) 1 (1922) 39
JFM48 1083

Moulton, Forest Ray

- [1] On a class of particular solutions of the problem of four
bodies.
Trans. Amer. Math. Soc. 1 (1900) 17-29
JFM31 889
- [2] On a class of particular solutions of the problem of four
bodies.
Popular Astronomy 8 (1900) 224
AJ2 180
- [3] On straight line solutions of the problem of n bodies. (abstract)
Bull. Amer. Math. Soc. 7 (1900/01) 249-250
- [4] The straight line solutions of the problem of N bodies.
Ann. of Math. (2) 12 (1910) 1-17
AJ16 98 JFM41 794
- [5] On orbits of ejection and collision in the problem of three
bodies. (abstract)
Bull. Amer. Math. Soc. (2) 19 (1912/13) 450
JFM44 *841
- [6] Closed orbits of ejection and related periodic orbits.
Painlevé's theorem.
Proc. London Math. Soc. (2) 11 (1913) 367-397
AJ16 99 JFM43 1083

- [7] The problem of three bodies.
Popular Astronomy 22 (1914) 197-207
AJ16 96
- [8] The Straight-line Solutions of the Problem of n Bodies.
chapter VIII p. 285-298 of: Moulton, F. R. (et al.) Periodic
Orbits Carnegie Institution of Washington, Washington 1920
Publication No. 161. Reprint: Johnson Reprint Corp. 1963

Müntz, Ch. H.

- [1] Die Ähnlichkeitsbewegungen beim allgemeinen n-Körperproblem.
Math. Z. 15 (1922) 169-187

Nikliborc, Władysław

- [1] Über das allgemeine Dreikörperproblem. I. Diskussion einer
Ungleichung; II. Grundlegende Formeln für die Geschwindigkeits-
komponenten.
Studia Math. 8 (1939) 28-67; 92-128
AJ41 *64 ZM20 318/319

Painlevé, Paul

- [1] Sur les singularités des équations de la Dynamique.
C. R. Acad. Sci. Paris 123 (1896) 636-639
JFM27 607/608
- [2] Sur les singularités des équations de la Dynamique et sur
le problème des trois corps.
~ 123 (1896) 871-873
JFM27 608
- [3] Sur les cas du problème des trois corps (et de n corps) où
deux des corps se choquent au bout d'un temps fini.
~ 125 (1897) 1078-1081
JFM28 652
- [4] Leçons sur la théorie analytique des équations différentielles,
professées à Stockholm 1895.
A. Hermann. Paris 1897
JFM28 262/266

Paul, E. W.

- [1] Function-Theoretic Analysis of Kepler's Equation.
Westinghouse Electric Corp. AFOSR Document Number 1123
(for Air Force Office of Scientific Research) 1961

Pedersen, Peder

- [1] Die Erweiterung der Lagrangeschen Dreieckslösungen im
allgemeinen Dreikörperproblem.
Astronom. Nachr. 211 (1920) c33-c42
AJ22 102

Perron, Oskar

- [1] Über eine Schar periodischer Lösungen des ebenen Dreikörperproblems (Mondbahnen).
Bayer. Akad. Wiss. Math.-Natur. Kl. S.-B. H.1 (1936) 157-176
ZM14 333

Petronievitch, B.

- [1] = Bilimovitch, A. [3]

Picard, Émile

- [1] Le problème des trois corps. À propos des recherches récentes de M. Sundman.
Bull. Sci. Math. (2) 37 (1913) 313-320
= Rev. gén. Sci. 24 (1913) 722-725
= Traité d'Analyse T. III (1928) Note 1 p. 601-608
AJ16 96 JFM44 824

Pierce, David A.

- [1] = Szebehely, Victor [5]
- [2] A Solution of the Regularized Equations of Motion of the Restricted Problem.
Astronom. J. 71 (1966) 545-561
- [3] Applications of a Solution of the Regularized Equations of Motion of the Restricted Problem.
~ 71 (1966) 562-565
- [4] = Szebehely, V. [9]
- [5] [& Standish, E. Myles Jr.]
Family of Consecutive Collision Earth-to-Moon Trajectories.
AIAA J. 4 (1966) 709-711
- [6] = Szebehely, Victor [12]

Pitkin, Edward T.

- [1] = Czuchry, A. J. [1]

Pizzetti, Paolo

- [1] Sopra alcune equazioni fondamentali nel problema degli n corpi.
Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur. 38 (1903) 954-961
JFM34 770/771
- [2] Casi particolari del problema dei tre corpi.
Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (5) 13₁ (1904) 17-26
AJ6 181 JFM35 724/725
- [3] Generalizzazione di alcune formule di Sundman.
Giorn. Mat. Battaglini 53 (1915) 276-283
AJ18 *170 JFM45 *1182

Plummer, H. C.

- [1] Note on the use of conjugate functions in some dynamical systems.
Proc. Roy. Irish Acad. Sect. A 32 (1914) 7-14
- [2] An Introductory Treatise on Dynamical Astronomy.
Cambridge U. P. 1918. Dover Publications reprint 1960

Pollard, Harry

- [1] Mathematical introduction to celestial mechanics.
Prentice-Hall. Englewood Cliffs, N. J. 1966
ZM141 238/239
- [2] Some nonlinear Tauberian theorems.
Proc. Amer. Math. Soc. 18 (1967) 399-401

Popoff, Kyrille

- [1] Sur la convergence des séries de la Balistique extérieure.
C. R. Acad. Sci. Paris 183 (1926) 266-267
- [2] Sur la convergence des séries de la Mécanique céleste.
~ 183 (1926) 472-474
- [3] Sur le problème des trois corps.
Atti del Congresso Internazionale dei Matematici Bologna
3-10 settembre 1928 (VI) tomo V 49-53
- [4] Sur le problème des trois corps.
Math. Ann. 101 (1929) 579-590
AJ31 46 JFM55₂ 1101

Putiata, T. V.

- [1] = Sokolov [30]

Pylarinos, O.

- [1] Zum Dreikörperproblem.
Math. Ann. 114 (1937) 150-160
AJ39 *65 ZM16 237
- [2] Über die Lagrangeschen Fälle im verallgemeinerten
Dreikörperproblem.
Math. Z. 47 (1942) 357-372
AJ43 *52 MR3 281 ZM26 23/24
- [3] Über das Dreikörperproblem.
Acta Math. 81 (1949) 257-263
AJ49 *89 MR11 545 ZM34 406

Rauch, L. M.

- [1] [& Riddell, W. C.]
The iterative solutions of the analytical N-body problem.
J. Soc. Indust. Appl. Math. 8 (1960) 568-581
MR22 1825 #10744 ZM101 233
- [2] The Algebraic Structure of the N-Body problem.
NASA CR-111 October 1964 32 pp

- [3] The Regularized Explicit Solution of the Analytic N-Body Problem.
NASA CR-123 October 1964 28pp

Richards, Paul B.

- [1] Lectures on regularization.
in: Space Mathematics, part 1. Lectures in Applied Mathematics,
vol. 5. pp 100-118
American Mathematical Society, Providence, Rhode Island 1966

Riddell, W. C.

- [1] = Rauch, L. M. [1]

Rivera Clavo = Clavo Rivera

Rosenblatt, Alfredo

- [1] Sur le cas de la collision générale dans le problème des trois corps.
Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (6)
3 (1926) 69-75
- [2] Sur la régularisation du problème plan des trois.
~ (6) 3 (1926) 271-274
- [3] Sur la régularisation du problème plan des trois corps.
Bull. Sci. Math. (2) 52 (1928) 195-202
JFM54 836
- [4] = García, Godofredo [2]
- [5] On the movement of a cosmic cloud.
Amer. J. Math. 66 (1944) 268-280
MR6 75 ZM60 *465

Rosenthal, Jenny E.

- [1] The equation of stability of periodic orbits of the restricted problem of three bodies in Thiele's regularizing coordinates.
Amer. J. Math. 53 (1931) 626-630
ZM2 211

Rössler, Max

- [1] Numerische Erfahrungen bei der Störungsrechnung in der Himmelsmechanik mit der Methode der Spinor-Regularisierung.
Z. Angew. Math. Phys. (ZAMP) 16 (1965) 395-402
AJ65 203
- [2] Analytische Störungstheorie im Dreikörperproblem mit Hilfe von regularisierenden Variablen.
Dissertation Nr. 3772 (1966) 71pp
Eidgenössische Technische Hochschule Zürich
- [3] = Stiefel, E. [6]

Samoilowa-Jachontowa, N.

- [1] Die singulären Punkte der Differentialgleichungen des Zwei-Körperproblems.
Byulleten' Astronomicheskogo Instituta no. 15 (1927) 169-176
AJ29 54/55
- [2] Über die Anwendung einer regularisierenden Veränderlichen zur Berechnung der Störungen. (German; Russian sum.)
~ no. 24 (1929) 35-44
AJ31 50
- [3] Sur le calcul des perturbations planétaires au moyen d'une nouvelle variable indépendante.
C. R. Acad. Sci. Paris 190 (1930) 1544-1546
AJ32 49 JFM562 1349
- [4] Die Störungen der kleinen Planeten als Funktionen der regularisierenden Veränderlichen.
Byulleten' Astronomicheskogo Instituta no. 27 (1931) 75-86
AJ33 57
- [5] Über die Anwendung einer regularisierenden Veränderlichen zur gruppenweisen Berechnung der Planeten-Störungen. (Russian, German sum.)
~ no. 41 (1936) 4-10
AJ39 67

Samter, H.

- [1] Das Zweizentren-Problem in der Störungstheorie.
Astronom. Nachr. 217 (1922) c129-c152
AJ24 67/68
- [2] Spezielle Störungen der Planeten und Kometen, numerische Behandlung besonderer Fälle des Dreikörperproblems.
Encykl. Math. Wiss. VI 2B 19 (1922)

Schanzle, Allan F.

- [1] Regularization and power series solution for the third order representation of the restricted 3 body problem.
Frank J. Seiler Research Laboratory SRL 65-0009 Dec. 1965
DDC: AD627 653

Schubart, Joachim

- [1] Zur Regularisierung des Zweierstosses im Dreikörperproblem.
Astronom. Nachr. 282 (1955) 262-264
AJ55 133 MR17 906

Schulze-Arenstorff, Richard F. = Arenstorff, Richard F.

Sconzo, Pasquale

- [1] A heuristic approach to the introduction of a universal variable.
Mem. Soc. Astronom. Ital. 38 (1967) 21 pp

- [2] An Algorithm to Obtain Series Expansions in the Three-Body Problem.
in: First Compilation of Papers on Trajectory Analysis and Guidance Theory NASA SP 141 (1967) 97-120

Selder, Helmut

- [1] Über singuläre Stellen in den Differentialgleichungen des Dreikörperproblems.
Bayer. Akad. Wiss. Math.-Natur. Kl. S.-B. 1964, Abt. II, (1965) 13-48
AJ65 *203 MR32 1346 #7852

Seeliger, H.

- [1] Über den neuen Stern im Andromedanebel (?).
Astronom. Nachr. 113 (1885) c358
[2] Über einen Fehler in den Jacobischen Vorlesungen über Dynamik.
Astronom. Nachr. 212 (1920/21) c1-c4
AJ22 103

Sémirot, Pierre

- [1] Application de la transformation de Sundman au problème d'Euler.
C. R. Acad. Sci. Paris 212 (1941) 848-850
AJ43 *52 MR3 215 ZM25 113
[2] Chocs imaginaires dans le problème des trois corps.
~ 212 (1941) 974-977
AJ43 *52 MR5 79 ZM25 270
[3] Chocs et solutions périodiques dans le problème des trois corps.
These Paris 1943 101p
= Annales de l'Observatoire de Bordeaux 18 fasc. 1, 101pp, 9 Fig. 1943
AJ45 63 MR8 176/177
[4] Conditions de choc dans le problème des trois corps.
C. R. Acad. Sci. Paris 218 (1944) 389-391
MR7 224 ZM61 *418
[5] Chocs triples imaginaires dans le problème des trois corps.
Mathematica, Timișoara 23 (1948) 85-87
AJ48 *78 MR10 333 ZM31 40

Siegel, Carl Ludwig

- [1] Der Dreierstoss.
Ann. of Math. (2) 42 (1941) 127-168
= Gesammelte Abhandlungen II, 169-210
AJ43 *52 JFM67 785 MR2 263 ZM24 364/365
[2] On the modern development of celestial mechanics.
Amer. Math. Monthly 48 (1941) 430-435
= Gesammelte Abhandlungen II, 211-216
MR3 215

- [3] Vorlesungen über Himmelsmechanik.
Springer-Verlag Berlin 1956
AJ56 *124 AJ57 *129 MR18 178 ZM70 454/455

Skljanskiĭ, A. L. = Skljanskij = Sklianskiĭ = Sklianski

- [1] On trajectories of a double collision in the problem of three bodies interacting with forces proportional to the logarithms of the mutual distances. (Russian)
Ukrain. Mat. Ž 6 (1954) 348-369
AJ55 *133 MR17 198 ZM57 161/162
- [2] On the theory of pairwise collision in the generalized three-body problem. (Russian)
~ 7 (1955) 160-166
AJ55 *133 MR17 542 ZM65 395
- [3] Sur la question de la classification des cointonations dans le problème généralisé des trois corps. (Russian; 80-81 French sum.)
~ 9 (1957) 66-81
MR19 368/369
- [4] Sur les trajectoires singulières du problème général des trois corps avec le choc binaire. (Russian; French sum.)
~ 9 (1957) 163-175
MR19 784
- [5] On the application of Sundman's and Chazy's methods to the generalized three-body problem. (Russian; English sum.)
~ 11 (1959) 380-392
MR24B 55/56 #D359

Sludsky, Th. = Sludskiĭ = Sloudsky

- [1] Sur le problème de plusieurs corps. (Russian)
Mat. Sbornik 9 (1878 (79?)) 536-545
JFM11 *646

Sokolov, Ju. (Yu.) = Sokoloff, George

- [1] Sur la limite inférieure des rayons de convergence des développements des coordonnées dans le problème des trois corps.
Kiev Bull. Ac. Sc. = Zapyski 14 (1925) 12-14
JFM51 779
- [2] Conditions d'une collision générale des trois corps qui s'attirent mutuellement suivant la loi de Newton.
Académie des Sciences de l'Ukraine. Mémoires de la classe des Sciences Physiques et Mathématiques tome IX, livr. 1 (1928) 1-64
JFM67 785

- [3] Sur la collision générale dans le cas symétrique du problème des trois corps. (Ukrainian; French sum.)
J. Inst. Math. Acad. Sci. Ukraine 1 (1934/35) 27-33
AJ37 *63 JFM612 1483 ZM12 38/39
- [4] Sur le choc dans le problème restreint des trois corps qui s'attirent (ou se repoussent) mutuellement proportionnellement à leurs masses et à une fonction de la distance. (Ukrainian; French sum.)
154-155 French sum.)
~ Nr. 3/4 (1934/35) 133-155
AJ37 *63 JFM612 1484 ZM12 128
- [5] Sur certaines singularités dans le problème restreint des trois corps. (Ukrainian; French sum.)
~ (1935/36) 107-118
JFM612 1484
- [6] Sur le choc dans le problème des trois corps qui s'attirent proportionnellement à leurs masses et à une fonction de la distance.
Acad. Roy. Belg. Bull. Cl. Sci. (5) 22 (1936) 295-305
AJ38 *79 ZM14 85
- [7] Sur les trajectoires singulières dans le problème des trois corps qui s'attirent mutuellement proportionnellement à leurs masses et à une fonction de la distance.
~ (5) 22 (1936) 540-551
AJ38 *79 ZM14 233
- [8] Sur un théorème de Weierstrass. (Ukrainian; 51 French sum.)
J. Inst. Math. Acad. Sci. Ukraine Nr. 2 47-51
ZM14 332
- [9] Sur une généralisation d'un théorème de M. J. Chazy. (Ukrainian; 7 French sum.)
~ (1937) Nr. 1 3-7
ZM18 *94
- [10] Remarques sur le problème des n corps généralisé. (Ukrainian; 100 French sum.)
~ (1937) Nr. 3 95-100
AJ39 *66 ZM18 96
- [11] Sur un mémoire de Kiveliovitch. (Ukrainian)
~ 4 (1938) 69-72
ZM18 280
- [12] Sur les points singuliers des intégrales du problème rectiligne des trois corps, qui s'attirent proportionnellement à l'inverse d'une puissance quelconque de la distance. (Ukrainian; 15-16 French sum.)
Acad. Sci. RSS Ukraine. Rec. Trav. (Zbirnik Prace) Inst. Mat. 1938, no. 1, 3-16
AJ40 *76 ZM21 164/165

- [13] Sur les points singuliers des intégrales dans le cas symétrique du problème des trois corps généralisé. (Ukrainian; 19-20 French sum.)
Acad Sci. RSS Ukraine Rec. Trav. (Zbirnik Prace) Inst. Mat. 1940, no. 3, 3-20
ZM25 113
- [14] Sur la collision générale des trois corps qui s'attirent mutuellement proportionnellement à l'inverse d'une puissance quelconque de la distance. (Ukrainian; 45 French sum.)
~ 1940, no. 4, 7-45 (1940)
MR2 207 ZM23 408
- [15] Sur le mouvement symétrique d'un système des points matériels qui agissent avec des forces dépendants des distances mutuelles. (Ukrainian; 114-115 French sum.)
~ 1940, no. 5, 99-115
MR2 207 ZM25 113/114
- [16] Sur les pôles des coordonnées dans le mouvement symétrique d'un système de points matériels qui s'agissent avec des forces dépendantes des distances mutuelles. (Ukrainian; 49-50 French sum.)
~ 1941, no. 6, 33-50
MR2 326 ZM24 363
- [17] Sur les trajectoires de collision simultanée de trois points matériels dont les forces attractives ou répulsives dépendent de leurs distances mutuelles.
C. R. (Doklady) Acad. Sci. URSS (N.S.) 33 (1941) 112-115
MR5 190/191
- [18] Sur un nouveau cas d'intégrabilité dans le problème rectiligne des trois corps.
~ 46 (1945) 95-98
MR7 224
- [19] On a spatial homographic motion of a system of three material points. (Russian)
Doklady Akad. Nauk. SSSR (N.S.) 58 (1947) 369-371
AJ47 *44 AJ49 *89 MR9 310 ZM30 83
- [20] On the trajectories of the rejection to infinity of three material points moving under the influence of their mutual interaction. (Russian)
~ 58 (1947) 539-542
AJ49 *89 MR9 625 ZM30 83
- [21] On infinite trajectories of a system of mass points under the influence of their mutual actions. (Ukrainian; Russian sum.)
Akad. Nauk. Ukrain. RSR. (Zbirnik Prace) Inst. Mat. 1948, No. 10, 142-164 (1948)
MR12 136

- [22] On the motion of a system of three material points on a straight line. (Russian)
Ukrain. Mat. Ž. 1 (1949) no. 3, 3-40
MR13 996 ZM49 *406 ZM59 170/171
- [23] On a general case of symmetric motion of a system of three material points. (Russian)
~ 2 (1950) no. 3, 7-44
MR13 996 ZM54 76
- [24] On rectilinear motion with a common collision of a system of three material points mutually attracting according to an exponential law. (Russian)
~ 2 (1950) no. 4, 18-24
MR13 996 ZM54 *76 ZM59 171
- [25] On the motion on a straight line of a system of three material points, each acting with forces proportional to the logarithms of their mutual distances. (Russian)
~ 2 (1950) no. 4, 25-36
MR13 996 ZM54 *76 ZM59 171
- [26] Osobyie traektorii systemy svobodnyh materialnyh toček. (Singular trajectories of a system of free material points) (Russian)
Monografii Instituta Matematiki, vyp. I. Akad. Nauk. Ukrainskoi SSR Kiev, 1951 126pp
MR14 910/912
- [27] On some general characteristics of the behavior of a material system in the neighborhood of a singular instant of time. (Ukrainian; Russian sum.)
Dopovidi Akad. Nauk Ukrain. RSR 1951, 227-231 (1951)
MR15 748
- [28] On a case of integrability of the equations of symmetric motion of a system of three material points. (Russian)
Ukrain. Mat. Ž. 3 (1951) 347-380
MR15 168 ZM45 263/264
- [29] Recherches sur la théorie des trajectoires singulières d'un système des points matériels libres. (Russian, French sum.)
~ 11 (1959) 3-15
MR22 1041 #6109
- [30] [& Putiata, T. V., & Fradlin, B. N.]
Works of Ukrainian Scientists on Analytical and Celestial Mechanics.
Prikladnaia Mekhanika 3 (1967) 89-99
(rev. in: International Aerospace Abstracts A68-13238)

Speiser, Andreas

- [1] Topologische Fragen aus der Himmelsmechanik.
Vierteljschr. Naturforsch. Ges. Zürich 85 (1940),
Beiblatt 32 (Rudolf Fueter Festschrift) 204-213
AJ42 *51 MR2 326 ZM23 273

Sperling, Hans J.

- [1] Computation of Keplerian Conic Sections.
ARS J. 31 (1961) 660-661
- [2] The binary collision in the N-body problem of celestial mechanics.
George C. Marshall Space Flight Center, Huntsville, Alabama.
R-AERO-IN-11-66 July 27, 1966 43pp
- [3] The Binary Collision in the N-Body Problem.
Icarus 9 (1968) 305-314
- [4] On the real singularities of the N-body problem.

Standish, E. Myles Jr.

- [1] = Szebehely, V. [5]
- [2] = Pierce, D. A. [5]

Steffensen, J. F.

- [1] Über die Integration des Dreikörperproblems in der Ebene.
Astronom. Nachr. 176 (1907) c221-c236
AJ9 167
(cf. von Zeipel [1])

Stiefel, Eduard

- [1] Many-Body Problems and Interplanetary Flight.
in: Groves, G. V. (Ed.): Dynamics of Rockets and Satellites.
p. 71-112
North-Holland Publ. Co. Amsterdam 1965
- [2] Koordinatenstörungen in regularisierten Variablen.
in: (Stiefel, E.: Ed.) Mathematische Methoden der
Himmelsmechanik und Astronautik 87-95
Bibliographisches Institut, Mannheim 1966 350pp
Mathematisches Forschungsinstitut Oberwolfach. Berichte 1
(Tagung 15.-21. März 1964)
MR34 723 #3979
- [3] [& Waldvogel, J.]
Problème restreint des trois corps. Généralisation de la
régularisation de Birkhoff pour le mouvement du mobile dans
l'espace à trois dimensions.
C. R. Acad. Sci. Paris 260 (1965) 805
AJ65 203 MR30 852 #4572 ZM123 *182
- [4] Neuere numerische Methoden in der Himmelsmechanik.
Nordisk Tidskr. Informations-Behandling 5 (1965) 51-60
AJ65 *203
- [5] = Kustaanheimo, P. [4]

- [6] [& Rössler, M. & Waldvogel, J. & Burdet, C. A.]
Report to NASA on: Methods of regularization for computing
orbits in celestial mechanics. 1966 124 pp
Institute for Applied Mathematics of the Swiss Federal
Institute of Technology. = NASA CR-769

Strömgren, E.

- [1] A new class of periodic solutions in the general problem of
three bodies.
Monthly Notices of the R.A.S. 80 (1919/20) 12-22
- [2] Zusatz zu: Pedersen, P. (Astronom. Nachr. 211 (1920)
c33-c42).
Astronom. Nachr. 211 (1920) c41-c44
- [3] Librationen und Ejektionsbahnen im allgemeinen Dreikörper-
problem, nebst einigen Beispiele periodischer Bahnen im
Vierkörperproblem.
Astronom. Nachr. 214 (1921) c26-c28
AJ23 109/110

Stumpff, Karl

- [1] Eine einfache symmetrische Ableitung der Lagrangeschen
partikulären Lösungen des Dreikörperproblems.
Astronom. Nachr. 280 (1951) 91-93
ZM44 204
- [2] Himmelsmechanik. I
II
VEB Deutscher Verlag der Wissenschaften Berlin I 1959;
II 1965
I: ZM92 239/240
- [3] A new solution of Hill's lunar problem.
Proc. Intern. Astronom. Union, Symp. 25, Thessaloniki, 1964
Academic Press, New York 1966 261-265
- [4] On the application of spinors to the problems of celestial
mechanics.
NASA TN D-4447 May 1968 41pp.

Subbotin, M. F.

- [1] Sur quelques propriétés du mouvement dans le problème des
n corps.
C. R. Acad. Sci. SSSR = Doklady, N.S. 27 (1940) 440-442
AJ42 *52 JFM66 1011 MR2 263 ZM25 114

Sundman, Karl Frithiof

- [1] Recherches sur le problème des trois corps.
Acta Societatis Scientiarum Fennicae 34 (1907) no. 6
2 + 43 pp
AJ9 166

- [2] Nouvelles recherches sur le problème des trois corps.
Acta Societatis Scientiarum Fennicae 35 (1909) no. 9
1-27
AJ14 *168
- [3] Sur les singularités réelles dans le problème des trois corps.
Compte rendu Congr. des Mathém. à Stockholm (22-25 sept.) 1909
Leipzig B. G. Teubner 1910 62-75
AJ12 191 JFM41 1019
- [4] Mémoire sur le problème des trois corps.
Acta Math. 36 (1912) 105-179
AJ14 160 v43 826/827

Szebehely, Victor G.

- [1] On the problem of three bodies in a plane.
Math. Mag. 26 (1952/53) 59-66
MR14 589/590 ZM48 177
- [2] The generalized inverse problem of orbit computation.
in: Space Research II (Ed: van de Hulst, H.C.) 318-338
Proceedings of the Second International Space Science
Symposium. Florence, April 10-14, 1961
North-Holland Publ. Co., Amsterdam 1961
- [3] The Restricted Problem of Three Bodies. (Lectures)
Summer Institute in Dynamical Astronomy August 1961 119pp
- [4] Application of the restricted problem of three bodies to
space mechanics.
Space Sci. Rev. 2 (1963) 219-249
AJ63 190
- [5] [& Pierce, D. A. & Standish, E. M.]
A group of earth-to-moon trajectories with consecutive
collisions.
in: Progress in Astronautics and Aeronautics
Vol. 14 Celestial Mechanics and Astrodynamics (Szebehely, V.,
Ed.) 35-51
Academic Press 1964
- [6] [& Giacaglia, Giorgio E.O.]
On the Elliptic Restricted Problem of Three Bodies.
Astronom. J. 69 (1964) 230-235
AJ64 203
- [7] Perturbation of the regularized equations of the restricted
problem of three bodies.
~ 69 (1964) 309-315
AJ64 203
- [8] Regularization in celestial mechanics.
Proc. Conf. Senior Navy Mathematicians, Monterey, Calif. 1965
Office of Naval Research, Washington 1967
- [9] [& Pierce, D. A.]
Regularizing Transformations of the Jacobian Integral.
Astronom. J. 71 (1966) 886-890

- [10] Solution of the Restricted Problem of Three Bodies by Power Series.
Astronom. J. 71 (1966) 968-975
- [11] Selection of Regularizing Functions.
~ 72 (1967) 370-373
- [12] [& Pierce, D. A.]
Advantages of Regularization in Space Dynamics.
AIAA J. 5 (1967) 1520-1522
- [13] Theory of Orbits. The Restricted Problem of Three Bodies.
Academic Press, New York 1967 668pp

Thiele, T. N.

- [1] Recherches numériques concernant des solutions périodiques d'un cas spécial du problème des trois corps (troisième mémoire).
Astronom. Nachr. 138 (1895) No. 3289 c1-c10
JFM26 1098

Tscherny, S. D.

- [1] Geometrische Lösung zweier spezieller Fälle des Problems der drei Körper.
Astronom. Nachr. 171 (1906) c129-c136; c349-c350
AJ8 170/171 JFM37 967/968
- [2] Geometrische Lösung des Problems der drei Körper in zwei speziellen Fällen. (Russian)
Kiew 1907 15pp
AJ10 194

Uno, Tosio

- [1] Sur les singularités des équations différentielles dans le problème des trois corps.
Ann. Mat. Pura Appl. (4) 14 (1935) 111-137
= Sendai Astronomiaj Raportoj 1 Nr. 10 (1935)
AJ37 *64 AJ38 80 JFM61 861/862 ZM12 128

Veiga de Oliveira, Fernando Vasco Alves da

- [1] Soluções homográficas no problema generalizado dos n corpos.
Faculdade de Ciências, Lisboa, 1956 82pp
MR19 993
- [2] Solutions homographiques du problème généralisé des n corps.
(Portuguese, French sum.)
Univ. Lisboa. Revista Fac. Ci. A (2) 6 (1957/58) 5-82
AJ59 *151 MR21 477/478 ZM82 173

Vernić, Radovan

- [1] Periodische Lösungen im Dreikörperproblem.
Glasnik Mat.-Fiz. Astronom. Ser. II Društvo Mat. Fiz.
Hrvatske 8 (1953) 247-266
AJ53 99 MR16 181 ZM53 438/439

- [2] Die Stossbedingungen im Dreikörperproblem.
Glasnik Mat.-Fiz. Astronom. Ser. II Društvo Mat. Fiz.
Hrvatske 9 (1954) 3-13
AJ54 107 MR16 868 ZM56 173/174
- [3] Diskussion der Sundmanschen Lösung des Dreikörperproblems.
Südslavische Akademie der Wissenschaften und Künste.
Zagreb 1954 146pp
AJ54 107/108 MR16 867/868 ZM57 61 ZM59 172
- [4] Numerische Auflösung des allgemeinen Dreikörperproblems.
(Croatian, German sum.)
Rad. Jugoslav. Akad. Znanosti Umjet nosti Zagreb 302 (1955)
45-76
AJ55 134 MR17 94 ZM65 394/395
- [5] Kritische Betrachtungen über die Zusammenstösse im
Mehrkörperproblem.(Serbo-Croatian, German sum.)
Rad. Jugoslav. Akad. Znan. Umjet. 314. Odjel Mat. Fiz.
Tehn. Nauke 7 (1957) 5-85
AJ57 *129 MR21 113 #622a MR22 1825 #10746 ZM125 123/124
- [6] Die Lösung des Mehrkörperproblems.(Serbo-Croatian, German sum.)
Rad. Jugoslav. Akad. Znan. Umjet. 314. Odjel Mat. Fiz.
Tehn. Nauke 7 (1957) 111-186 (1 insert)
AJ57 *129 MR21 113 #622b MR22 1825 #10745 ZM125 123/124

Vietoris, L.

- [1] Zur Geometrie der ebenen Massenanziehungsprobleme.
Math. Z. 19 (1924) 130-135
= Publ. og mindre Medd. fra Københavns Obs. No. 42
AJ26 55

von Brunn, A.

- [1] Bemerkungen zum 3-Körperproblem.
Schriften der Naturforschenden Ges. in Danzig NF 153
48 pp (1920)
AJ22 101 AJ24 66/67

von Zeipel, H.

- [1] Bemerkungen zu dem Aufsatz "Über die Integration des
Dreikörperproblems in der Ebene." A. N. 4214.
Astronom. Nachr. 176 (1907) c313-c314
(AJ9 167)
- [2] Sur les singularités du problème des n corps.
Arkiv för Matematik Astronomi och Fysik 4 (1908) no. 32 4 pp
AJ10 195 JFM39 *792
- [3] Sur le problème des trois corps.
Vierteljschr. Astronom. Ges. 52 (1917) 56-79
AJ19 111/112

Waldvogel, Jörg

- [1] Regularisierung und konforme Abbildung.
in: (Stiefel, E.: Ed.) Mathematische Methoden der
Himmelsmechanik und Astronautik 33-45
Bibliographisches Institut, Mannheim 1966 350pp
Mathematisches Forschungsinstitut Oberwolfach. Berichte. 1
(Tagung 15.-21. März 1964)
MR34 723 #35:9
- [2] = Stiefel, E. [3]
- [3] = Stiefel, E. [6]
- [4] Die Verallgemeinerung der Birkhoff-Regularisierung für das
räumliche Dreikörperproblem.
Dissertation Nr. 3816 (1966) 82pp
Eidgenössische Technische Hochschule Zürich
- [5] Die Verallgemeinerung der Birkhoff-Regularisierung für das
räumliche Dreikörperproblem.
Bull. Astronom. (3) 2 (1967) 295-341

Weierstrass, Karl [Ed.: Mittag-Leffler, G.]

- [1] Zur Biographie von Weierstrass.
Acta Math. 35 (1912) 29-65

Wilczynski, E. J.

- [1] Ricerche geometriche intorno al problema dei tre corpi.
Ann. Mat. Pura Appl. (3) 21 (1913) 1-31
AJ18 *178 JFM44 824/825

Williams, Carol

- [1] Solution of Birkhoff's Equations of the Regularized Restricted
Problem.
Astronom. J. 71 (1966) 976-982

Williams, W. L.

- [1] Permanent configurations in the problem of five bodies.
Trans. Amer. Math. Soc. 44 (1938) 563-579
AJ40 *78 ZM20 319

Wintner, Aurel

- [1] Zur effektiven Stabilität des Mondes.
Math. Z. 32 (1930) 675-698
AJ33 57 JFM56 1 779
- [2] The Analytical Foundations of Celestial Mechanics.
Princeton University Press 1941

Woronez, P.

- [1] Einige besondere Fälle der Bewegung eines Systems materieller Punkte unter Einwirkung gegenseitiger Kräfte.
Kiew 20pp. (Russian) (1904?)
JFM35 727/728

Yarov-Yarovoi, M. S.

- [1] Integration of regularized equations of the two-body problem.
Differential Equations 1 (1967) 746-760
transl. from: Differentsial'nye Uravneniya 1 (1965) 961-976
- [2] Solution of regularized equations in perturbation theory.
~ 1 (1967) 943-964
~ 1 (1965) 1204-1230

Yeomans, Donald K.

- [1] Exposition of Sundman's regularization of the three body problem.
NASA TM X-55636 Dec. 1966 105pp (N67 14919)

Zagar, F.

- [1] Sopra il problema dei tre corpi nel caso in cui le costanti delle aree siano tutte nulle.
Memoire della Società Astronomica Italiana 3 (1925) 130-136
AJ27 *51

Zeipel, H. von = von Zeipel, H.

Zuser, Karl

- [1] Die Riemannsche Fläche des Zweikörperproblems.
Bayer. Akad. Wiss. Math.-Natur. Kl. S.-B. (1960) 61-93
MR26 746 #3898 ZM131 213

ADDENDUM TO THE BIBLIOGRAPHY

Deprit, André

- [5] Concerning Kustaanheimo-Stiefel's regularization.
Z. Angew. Math. Phys. (ZAMP) 19 (1968) 369-372

Ferroli, D.

- [1] On the regularization of the problem of three bodies.
J. Indian Math. Soc. 18 (1929) 158-167
JFM55₂ 1102

Jacobi, Carl Gustav Jakob

- [1] Vorlesungen über Dynamik (2nd ed.).
Gesammelte Werke, Supplementband. Berlin 1884

Pollard, Harry

- [3] [& Saari, D. G.]
Singularities of the n-body problem. I. (abstract)
Notices Amer. Math. Soc. 15 (1968) 631

Saari, D. G.

- [1] = Pollard, H. [3]

Schaub, Werner

- [1] Über einen speziellen Fall des Fünfkörperproblems.
Astronom. Nachr. 236 (1929) c33-c54
JFM 55₂ 1102

Sludsky, Th.

- [2] Note sur quelques cas particuliers du problème de plusieurs corps.
Soc. imp. natur. Moscou 1892 437-440
JFM 24 880/881

Stiefel, Eduard

- [0] Die Renaissance der Himmelsmechanik.
Elem. Math. 19 (1964) 97-106
ZM 128 413

Szebehely, Victor G.

- [11] MR 36 482 #2380