



United States Department of the Interior

FISH AND WILDLIFE SERVICE

New Mexico Ecological Services Field Office
2105 Osuna NE
Albuquerque, New Mexico 87113
Phone: (505) 346-2525 Fax: (505) 346-2542

September 12, 2007

Cons. # 22420-2007-F-0056

Memorandum

To: Area Manager, Albuquerque Area Office, U.S. Bureau of Reclamation,
Albuquerque, New Mexico

From: Field Supervisor, U.S. Fish and Wildlife Service, New Mexico Ecological Services
Field Office, Albuquerque, New Mexico

Subject: U.S. Fish and Wildlife Service's Biological Opinion on the Effects of Actions
Associated with the Biological Assessment for the Corrales Siphon River
Maintenance Project

This document transmits the U.S. Fish and Wildlife Service's (Service) biological opinion on the effects of the proposed Corrales Siphon River Maintenance Project (Corrales Siphon Project). The Corrales Siphon river maintenance priority site is located at River Mile 199.7, on the west side of the Rio Grande, approximately 700 feet downstream of the Arroyo de la Barranca confluence in Bernalillo County, New Mexico. The Corrales Siphon conveys water under the Rio Grande to the Corrales Main Canal for the Middle Rio Grande Conservancy District. The Bureau of Reclamation (Reclamation) is proposing to protect the inverted siphon and associated infrastructure from damage caused by potential westward migration of the Rio Grande. Reclamation proposes to protect the bank by moving the river eastward using a bioengineering technique designed to improve and create habitat for the Rio Grande silvery minnow (*Hybognathus amarus*) (silvery minnow) and the southwestern willow flycatcher (*Empidonax traillii extimus*) (flycatcher). This biological opinion concerns the effects of the proposed action on the endangered silvery minnow, flycatcher and their designated critical habitat. Your request for formal consultation, in accordance with section 7 of the Endangered Species Act (ESA) of 1973, as amended (16 U.S.C. 531 *et seq.*) was received on February 1, 2007.

This biological opinion is based on information submitted in the January 2007 Middle Rio Grande Corrales Siphon Biological Assessment (Biological Assessment) and other sources of information available to the Service. A complete administrative record of this consultation is on file at the Service's New Mexico Ecological Services Field Office (NMESFO).

This biological opinion does not rely on the regulatory definition of "destruction or adverse modification" of critical habitat at 50 CFR 402.02. Instead, we have relied upon the statute and the August 6, 2004, Ninth Circuit Court of Appeals decision in *Gifford Pinchot Task Force v.*

USDI Fish and Wildlife Service (CIV No. 03-35279) to complete the following analysis with respect to designated critical habitat. This consultation analyzes the effects of the action and its relationship to the function and conservation role of silvery minnow and flycatcher designated critical habitat to determine whether the current proposal destroys or adversely modifies designated critical habitat. This document represents our biological opinion for the silvery minnow and its designated critical habitat, and the flycatcher and its designated critical habitat, in accordance with section 7 of the Act.

You have determined that the proposed project “may affect, but is not likely to adversely affect,” the flycatcher, its designated critical habitat. We concur with these determinations for the following reasons:

The flycatcher is a migrant through this portion of the Rio Grande and may be present between April and June, and again in August. Suitable nesting habitat does not currently exist within the project area and the nearest known occupied habitats are located within the Isleta and San Juan pueblos, approximately 30 miles south and 70 miles north of the project area, respectively. Although a small amount of riparian vegetation within the project area may be disturbed by the project, the proposal includes the planting of riparian and wetland plants in newly created wetland areas that could eventually mature and create potentially suitable flycatcher habitat. Noise from construction activities could affect individual birds moving in the area; however, construction will not occur during the flycatcher breeding or migration season to avoid noise impacts.

Designated flycatcher critical habitat is not within the project area. Designated critical habitat exists upstream on the Rio Grande and would not be affected by the Corrales Siphon Project. Effects to individual flycatchers are not likely to occur since no nesting occurs in the vicinity of the project area, and opportunities for nesting habitat in current and future breeding habitats are not significantly affected. The number of flycatcher territories in the Middle Rio Grande Management Unit has exceeded recovery goals (100 territories) for the past three years (Ahlers and Doster 2007). Additionally, flycatcher habitat restoration projects (see Environmental Baseline, below) are occurring throughout the basin, improving conditions for the flycatcher. Given the relatively small riparian vegetation disturbance created by this project, the effects of this project on flycatcher are discountable.

The remainder of this biological opinion will deal with the effects of implementation of the proposed action on the silvery minnow and its designated critical habitat.

Consultation History

The proposed action is addressed in the Reclamation and U.S. Army Corp of Engineers’ 2003 Programmatic Biological Assessment and the Service’s *Biological Opinion and Conference Report entitled, Bureau of Reclamation’s Water and River Maintenance Operations, Army Corps of Engineers’ Flood Control Operation, and Related Non-Federal actions on the Middle Rio Grande, New Mexico* (2003 Middle Rio Grande Biological Opinion). The proposed Corrales Siphon Project tiers off this 2003 section 7 consultation, which emphasized the use of

bioengineering in conjunction with river maintenance projects whenever possible.

BIOLOGICAL OPINION

DESCRIPTION OF PROPOSED ACTION

The Biological Assessment provided for this project by Reclamation, contains the comprehensive description of the purpose and need for the proposed action and the project area, details on construction of the bioengineering project, revegetation, adaptive management, a description of environmental commitments, and effects determination for listed species and critical habitat. The material contained in the Biological Assessment is herein incorporated by reference (U.S. Bureau of Reclamation 2007). The following description of the proposed action is a summary of the material in the biological assessment and should not be considered as the complete description.

Purpose and Objective

The Corrales Siphon Project is proposed by Reclamation as one of their priority river maintenance sites. The Corrales Siphon priority site is located at River Mile 199.7, on the west side of the Rio Grande, approximately 700 feet downstream of the Arroyo de la Barranca confluence in Bernalillo County, New Mexico. It conveys water under the Rio Grande to the Corrales Main Canal for the Middle Rio Grande Conservancy District. Reclamation is proposing to protect the inverted siphon and associated infrastructure from damage caused by potential westward migration of the Rio Grande by protecting the bank and moving it eastward with a bioengineering project. In addition to protecting irrigation infrastructure and public health and safety, this project is designed to improve and create habitat for the silvery minnow and the flycatcher (U.S. Bureau of Reclamation 2007).

Project Description

Reclamation proposes to install a bio-engineered bankline on the west side of the Rio Grande to protect the existing Corrales Siphon. The main objective of this bankline is to provide long term stability through the establishment of deep rooted native vegetation on the bankline. Coir (a biodegradable erosion control product) fabric will be used to provide temporary stability (approximately three years) until the trees and shrubs become established. However, while trees and shrubs may protect the bankline, vegetation alone generally does not prevent toe scour. Therefore, the design includes the installation of stone below the bio-engineered bankline to prevent toe scour.

Access to the Corrales Siphon priority site will be by existing dirt roads along the Corrales Main Canal that are maintained by the Middle Rio Grande Conservancy District. The bio-engineered bankline will be installed by placing fill in the existing channel eastward from the existing bankline. Once the bankline is built out to the desired location, fill will be placed at a 2:1 (H:V [horizontal:vertical]) slope. Stone will then be placed directly on the channel bottom, without excavating. The toe will be thickened, to resist toe scour, and will have an approximate width of 9 feet and height of 1.5 feet. Stone will extend another 3.5 feet up the new bankline, at an

approximate width of 7 feet. This stone will be 12-inch nominal size, at a 2:1 (H:V) slope to the approximate 4,000-cfs water surface elevation. The thickness of the stone layer will be approximately 3.1 feet, measured perpendicular to the bankline.

The bio-engineered bankline will be constructed above the stone. The bankline will have three layers, each offset to the west by two feet. Each layer consists of a coir fabric block (16" × 9" × 120" [H × W × L]) with both a bottom and top coir fabric mat, and two lifts of soil. The bottom mat extends 6.25 feet, and the top mat 4 feet, from the edge of the coir fiber block. The 2-foot width directly behind the coir blocks will be backfilled with topsoil and loosely compacted. This fill material will then be planted with native riparian vegetation (see "Revegetation" below) to help ensure bank stability and improve habitat.

Behind the 2-foot area containing plantings, soil will be placed in two lifts. Each lift will be placed at 10 inches thick, and then will be wetted and compacted to an 8-inch thickness. The total compacted thickness of each layer will be approximately 16 inches. The bottom lift will be placed directly on top of the bottom coir fabric mat. The top coir fabric mat will then be placed on top of the two lifts. Stakes of either wood or metal will be used to hold both the top and bottom coir fabric mats in place, based on the pattern recommended by the manufacturer. The new bankline is designed so that flows greater than approximately 4,000 cfs should inundate only the stone bankline protection, while a flow of 10,000 cfs should inundate the second lift of coir fabric/block. After the third lift, the ground will be graded at a 4:1 (H:V) slope to reach the ground elevation as defined by the newly placed fill. This final sloped area will be seeded and protected against erosion caused by runoff associated with rainfall.

Fill material for the bio-engineered bankline will be obtained from the adjacent Pueblo of Sandia habitat restoration project funded by the Fish and Wildlife Service's Management of Exotics for the Recovery of Endangered Species Program (MERES) (see Pueblo of Sandia MERES Project, below). This project includes excavation of a side channel in the bar east of the Corrales Siphon project site. Sediment from the Pueblo of Sandia MERES project will be hauled across the Rio Grande and used as fill between the bio-engineered bankline and the existing bankline. The sediment will be hauled using articulated dump trucks or scrapers. The location at which the river will be crossed will be selected at the time of construction, based on hydrologic and morphologic conditions; selection will emphasize finding a crossing point with a shallow depth and sufficiently solid bed to allow safe equipment access. Reclamation estimates that 500-1000 round trips across the river will be needed over a one month period. Crews are expected to work a total of 18 days (following a 5-4 sequence of 9 hour days resulting in 2 or 3 day weekends). Equipment will create a 30 foot wide swath of disturbance and the crossing will be approximately 220 feet across the channel. Imported fill material may also be used if necessary.

Replacing the steep bankline with approximately one acre of bio-engineered bankline is designed to create a gradient of benthic habitat at various water levels. Establishment of riparian vegetation on the stabilized bankline would provide nursery habitat for silvery minnow when inundated during the spawning season. Although the wetted area at base flows will be reduced, the habitat value of the remaining wetted area will be improved by replacing one acre of deep higher water velocity river channel with a bio-engineered bankline having a gradient of water

velocities and depths.

Construction would occur during fall 2007, starting in September, when river flows are minimal. Construction equipment utilized would include the following: bulldozers, excavators (land-track and amphibious), water trucks, scrapers, dump trucks, loaders, and motor graders. The total amount of disturbance created during construction is estimated to be approximately one acre, most of which is within the existing floodway.

The Biological Assessment states that Reclamation would use Best Management Practices (BMPs) for placement of rock and other fill for constructing the stabilized bankline and would continue to coordinate with the Service on construction techniques to avoid harm to silvery minnow in the work area (U.S. Bureau of Reclamation 2007). According to a Reclamation biologist associated with the Corrales Siphon Project, the BMPs that are currently used during river maintenance projects involving in-channel work are (M. Porter, Reclamation, *in litt.* 2007):

1. Use the excavator to gently place riprap (or sediment) in the proper location rather than dropping it into the water. This gives fish the opportunity to swim away from the rock being placed, reducing the probability of physical harm.
2. Push sediment fill from the bankline using a bulldozer rather than dumping it directly from an excavator or truck into the water. This provides fish the opportunity to swim away from the sediment being moved, reducing the probability of physical harm.
3. Maintain an open water area for fish to retreat to during placement of riprap and sediment fill. This will be accomplished by maintaining a connection to the river from the construction area to prevent the formation of isolated pools or channels, where fish could become trapped. This BMP allows fish the opportunity to swim away from potential harm in the work area.
4. When crossing the wetted river channel with heavy equipment, the path will be placed in a shallow riffle hardened with gravel. This is preferable to the shortest distance across the river, which would have deeper water and a faster current. This BMP is intended to reduce the chances of a spill, allow the trucks to traverse the river quicker, and reduce turbidity.

Revegetation

Approximately 150 poles (cottonwood [*Populus deltoides*] or black willow [*Salix gooddingii*]) will be planted throughout disturbed areas of the project. Approximately 200 containerized New Mexico olives (or other native shrubs, at the discretion of Reclamation biologists) will be planted in the disturbed area of the project. Coyote willows (*Salix exigua*) will be planted as part of the bio-engineered bankline. Additional coyote willows may be planted elsewhere in the project area. The disturbed area will be planted with a grass seed mix to be specified by Reclamation biologists (U.S. Bureau of Reclamation 2007).

Pueblo of Sandia MERES Project

The Pueblo of Sandia is in the planning and design stages of a project to be funded under the

U.S. Fish and Wildlife Service's MERES Project habitat restoration program. The site for this project is the large bar on the opposite side of the Rio Grande from the Corrales Siphon priority site. Designs for this project are being developed by SWCA Environmental Consultants, a firm employed by the Pueblo. Reclamation has committed to coordinating with SWCA to ensure compatibility between the two projects. The MERES project involves construction of one or more secondary channels through the bar. Earth material from the secondary channel excavation may be hauled across the river by Reclamation, to be used as fill behind the bio-engineered bankline. According to the Biological Assessment, the MERES project is an essential component for protecting the siphon. Splitting flow into a second channel with the MERES project would spread the volume of water out over a greater surface area and increase the width-to-depth ratio of the river. The resulting decrease in depth and velocity would benefit juvenile and adult silvery minnow by increasing the total amount of preferred habitat conditions (<40 cm deep and <10 cm/s water velocity). The increased width-to-depth ratio would also reduce water velocities and allow sediment deposition in the shallower channels, which would improve habitat conditions for silvery minnow (U.S. Bureau of Reclamation 2007). The MERES project is considered in a separate consultation.

The Biological Assessment states that if the MERES project is canceled or otherwise fails to result in construction of secondary channels through the bar across the river from the priority site, Reclamation may construct a side channel through the bar and that the maximum volume of earth material to be excavated by Reclamation for this secondary channel is 10,000 cubic yards. This Biological Opinion does not consider the effects of this action. Therefore, if this element is added to the Corrales Siphon project, Reclamation must reinitiate consultation with the Service.

Environmental Commitments

- Construction of the stabilized bankline would be implemented during low flows (200-1000 cfs) to minimize the area of disturbance at the construction site.
- All construction debris and waste would be disposed of at an approved landfill facility.
- Best Management Practices would be implemented and utilized to prevent stormwater runoff and water pollution from entering the Rio Grande during construction activities.

Adaptive Management

Reclamation is proposing to use adaptive management principles to ensure that the project purposes are fulfilled and that undesirable conditions do not develop. The Biological Assessment states that based on post-project monitoring, additional work may be done at the project site to address perceived deficiencies and unforeseeable developments. For this site, likely adaptive management activities include the following:

- Replanting vegetation if initial survival rates are too low.
- Herbicide spraying or other measures to control weeds.
- Performing additional earthwork, rock placement, and coir revetment placement to repair any damage that occurs to the bio-engineered bankline. Earthwork in the channels through the bar opposite the project site may also occur. The maximum earthwork quantity for this adaptive management activity is 10,000 cubic yards, the maximum toe rock quantity is 600 cubic yards, and the maximum additional quantity of coir fabric is 1,500 square yards

Access for adaptive management activities will be the same as for initial construction, except that work on the channels through the bar on the east side of the river may require access through the Pueblo of Sandia. All adaptive management activities will occur during a 6-year period, beginning at the conclusion of project construction. Owing to the inherent uncertainty in adaptive management activities, the Biological Assessment states that the actual work performed will very likely require material quantities that are much less than the maximum quantities specified (U.S. Bureau of Reclamation 2007).

As currently proposed, earthwork in the channels through the bar opposite the proposed bioengineered bankline will be considered during the consultation process for the Pueblo of Sandia's MERES project. However, if any additional earthwork by Reclamation is required, reinitiation of consultation will be necessary because the effects of such work on listed species and their designated critical habitat were not thoroughly described in the Biological Assessment, and thus, have not been thoroughly considered in this biological opinion. If Reclamation determines that any of the other activities described above for "adaptive management" are required, Reclamation should communicate with the Service to determine if reinitiation of consultation is required.

Action Area

The action area is defined as the area from the Angostura Diversion Dam to the Isleta Diversion Dam and the entire width of the 100 year Rio Grande floodplain within that reach.

STATUS OF THE SPECIES

RIO GRANDE SILVERY MINNOW

Description

The silvery minnow currently occupies a 170-mile reach of the Middle Rio Grande, New Mexico, from Cochiti Dam, Sandoval County, to the headwaters of Elephant Butte Reservoir, Socorro County (U.S. Fish and Wildlife Service 1994). The silvery minnow is a stout minnow, with moderately small eyes, a small, sub-terminal mouth, and a pointed snout that projects beyond the upper lip (Sublette *et al.* 1990). The back and upper sides of the silvery minnow are silvery to olive, the broad mid-dorsal stripe is greenish, and the lower sides and abdomen are silver. Maximum length attained is about 3.5 inches (in). The only readily apparent sexual dimorphism is the expanded body cavity of ripe females during spawning (Bestgen and Propst 1994).

In the past, the silvery minnow was included with other species of the genus *Hybognathus* due to morphological similarities. Phenetic and phylogenetic analyses corroborate the hypothesis that it is a valid taxon, distinctive from other species of *Hybognathus* (Cook *et al.* 1992, Bestgen and Propst 1994). It is now recognized as one of seven species in the genus *Hybognathus* in the United States and was formerly one of the most widespread and abundant minnow species in the Rio Grande basin of New Mexico, Texas, and Mexico (Pflieger 1980, Bestgen and Platania

1991). Currently, *Hybognathus amarus* is the only remaining endemic pelagic spawning minnow in the Middle Rio Grande. The speckled chub (*Extrarius aestivalus*), Rio Grande shiner (*Notropis jemezianus*), phantom shiner (*Notropis orca*), and bluntnose shiner (*Notropis simus*) are either extinct or have been extirpated from the Middle Rio Grande (Bestgen and Platania 1991).

Legal Status

The silvery minnow was federally listed as endangered under the ESA on July 20, 1994 (U.S. Fish and Wildlife Service 1994). The species is also listed as an endangered species by the state of New Mexico. Primary reasons for listing the silvery minnow are described below in the Reasons for Listing section.

Designated critical habitat for the silvery minnow was designated on February 19, 2003 (68 FR 8088). The critical habitat designation extends approximately 157 from Cochiti Dam, Sandoval County, New Mexico downstream to the utility line crossing the Rio Grande, a permanent identified landmark in Socorro County, New Mexico. The critical habitat designation defines the lateral extent (width) as those areas bounded by existing levees or, in areas without levees, 300 feet (ft) or riparian zone adjacent to each side of the bank full stage of the Middle Rio Grande. Some developed lands within the 300 ft lateral extent are not considered critical habitat because they do not contain the primary constituent elements of designated critical habitat and are not essential to the conservation of the silvery minnow. Lands located within the lateral boundaries of the critical habitat designation, but not considered critical habitat include: developed flood control facilities, existing paved roads, bridges, parking lots, dikes, levees, diversion structures, railroad tracks, railroad trestles, water diversion and irrigation canals outside of natural stream channels, the Low Flow Conveyance Channel, active gravel pits, cultivated agricultural land, and residential, commercial, and industrial developments. The Pueblo lands of Santo Domingo, Santa Ana, Sandia, and Isleta within this area are not included in the critical habitat designation. Except for these Pueblo lands, the remaining portion of the silvery minnow's occupied range in the Middle Rio Grande in New Mexico is designated as critical habitat (68 FR 8088).

Habitat

The silvery minnow travels in schools and tolerates a wide range of habitats (Sublette *et al.* 1990); yet, generally prefers low velocity (<0.33 ft per second) areas over silt or sand substrate that are associated with shallow [< 15.8 inch (in)] braided runs, backwaters or pools (Dudley and Platania 1997). Habitat for the silvery minnow includes stream margins, side channels, and off-channel pools where water velocities are low or reduced from main-channel velocities. Stream reaches dominated by straight, narrow, incised channels with rapid flows are not typically occupied by silvery minnow (Sublette *et al.* 1990, Bestgen and Platania 1991).

Adult silvery minnow are most commonly found in backwaters, pools, and habitats associated with debris piles; whereas, young of year (YOY) occupy shallow, low velocity backwaters with silt substrates (Dudley and Platania 1997). A study conducted between 1994 and 1996 characterized habitat availability and use at two sites in the Middle Rio Grande at Rio Rancho and Socorro. From this study Dudley and Platania (1997) reported that the silvery minnow was

most commonly found in habitats with depths less than 19.7 in. Over 85 percent were collected from low-velocity habitats (<0.33 ft/sec) (Dudley and Platania 1997, Watts *et al.* 2002).

Designated Critical Habitat

The Service has determined the primary constituent elements (PCEs) of silvery minnow designated critical habitat based on studies on silvery minnow habitat and population biology (68 FR 8088). They include:

1. A hydrologic regime that provides sufficient flowing water with low to moderate currents capable of forming and maintaining a diversity of aquatic habitats, such as, but not limited to the following: backwaters (a body of water connected to the main channel, but with no appreciable flow), shallow side channels, pools (that portion of the river that is deep with relatively little velocity compared to the rest of the channel), and runs (flowing water in the river channel without obstructions) of varying depth and velocity – all of which are necessary for each of the particular silvery minnow life-history stages in appropriate seasons (e.g., the silvery minnow requires habitat with sufficient flows from early spring (March) to early summer (June) to trigger spawning, flows in the summer (June) and fall (October) that do not increase prolonged periods of low- or no flow, and relatively constant winter flow (November through February));
2. The presence of eddies created by debris piles, pools, or backwaters, or other refuge habitat within unimpounded stretches of flowing water of sufficient length (i.e., river miles) that provide a variation of habitats with a wide range of depth and velocities;
3. Substrates of predominantly sand or silt; and
4. Water of sufficient quality to maintain natural, daily, and seasonally variable water temperatures in the approximate range of greater than 1°C (35°F) and less than 30°C (85°F) and reduce degraded conditions (e.g., decreased dissolved oxygen, increased pH).

These PCEs provide for the physiological, behavioral, and ecological requirements essential to the conservation of the silvery minnow.

Life History

The species is a pelagic spawner that produces 3,000 to 6,000 semi-buoyant, non-adhesive eggs during a spawning event (Platania 1995, Platania and Altenbach 1998). The majority of adults spawn in about a one-month period in late spring to early summer (May to June) in association with spring runoff. Platania and Dudley (2000, 2001) found that the highest collections of silvery minnow eggs occurred in mid- to late May. In 1997, Smith (1999) collected the highest number of eggs in mid-May, with lower frequency of eggs being collected in late May and June. These data suggest multiple silvery minnow spawning events during the spring and summer, perhaps concurrent with flow spikes. Artificial spikes have apparently induced silvery minnow

to spawn (Platania and Hoagstrom 1996). It is unknown if individual silvery minnow spawn more than once a year or if some spawn earlier and some later in the year.

Platania (2000) found that development and hatching of eggs are correlated with water temperature. Eggs of the silvery minnow raised in 30°C water hatched in approximately 24 hours while eggs reared in 20-24°C water hatched within 50 hours. Eggs were 0.06 in size upon fertilization, but quickly swelled to 0.12 in. Recently hatched larval fish are about 0.15 in standard length and grow about 0.005 in size per day during the larval stages. Eggs and larvae have been estimated to remain in the drift for 3-5 days, and could be transported from 134 to 223 mi downstream depending on river flows (Platania 2000). Approximately three days after hatching the larvae move to low velocity habitats where food (mainly phytoplankton and zooplankton) is abundant and predators are scarce. YOY attain lengths of 1.5 to 1.6 in by late autumn (U.S. Fish and Wildlife Service 1999). Age-1 fish are 1.8 to 1.9 in by the start of the spawning season. Most growth occurs between June (post spawning) and October, but there is some growth in the winter months. In the wild, maximum longevity is about 25 months, but very few survive more than 13 months (U.S. Fish and Wildlife Service 1999). Captive fish have lived up to four years (C. Altenbach, City of Albuquerque, *pers. comm.* 2003).

Platania (1995) suggested that historically the downstream transport of eggs and larvae of the silvery minnow over long distances was likely beneficial to the survival of their populations. This behavior may have promoted recolonization of reaches impacted during periods of natural drought (Platania 1995). The spawning strategy of releasing floating eggs allows the silvery minnow to replenish populations downstream, but the current presence of diversion dams (Angostura, Isleta, and San Acacia Diversion Dams) prevents recolonization of upstream habitats (Platania 1995). As populations are depleted upstream and diversion structures prevent upstream movements, isolated extirpations of the species through fragmentation may occur (U.S. Fish and Wildlife Service 1999). Adults, eggs and larvae are also transported downstream to Elephant Butte Reservoir. It is believed that none of these fish survive because of poor habitat and predation from reservoir fishes (U.S. Fish and Wildlife Service 1999).

The silvery minnow is herbivorous (feeding primarily on algae); this is indicated indirectly by the elongated and coiled gastrointestinal tract (Sublette *et al.* 1990). Additionally, detritus, including sand and silt, is filtered from the bottom (Sublette *et al.* 1990, U.S. Fish and Wildlife Service 1999).

Population Dynamics

Generally, a population of silvery minnow consists of only two age classes: YOY and Age-1 (U.S. Fish and Wildlife Service 1999). The majority of spawning silvery minnow is one year old. Two year old fish comprise less than 10 percent of the spawning population. High silvery minnow mortality occurs during or subsequent to spawning, consequently very few adults are found in late summer. By December, the majority (greater than 98 percent) of individuals are YOY (Age 0). This population ratio does not change appreciably between January and June, as Age 1 fish usually constitute over 95 percent of the population just prior to spawning.

Platania (1995) found that a single female in captivity could broadcast 3,000 eggs in eight hours. Females produce 3 to 18 clutches of eggs in a 12-hour period. The mean number of eggs in a clutch is approximately 270 (Platania and Altenbach 1998). In captivity, silvery minnow have been induced to spawn as many as four times in a year (C. Altenbach, City of Albuquerque, *pers. comm.* 2000). It is not known if they spawn multiple times in the wild. The high reproductive potential of this fish appears to be one of the primary reasons that it has not been extirpated from the Middle Rio Grande. However, the short life span of the silvery minnow increases the population instability. When two below-average flow years occur consecutively, a short-lived species such as the silvery minnow can be impacted, if not completely eliminated from dry reaches of the river (U.S. Fish and Wildlife Service 1999).

Distribution and Abundance

Historically, the silvery minnow occurred in 2,465 mi of rivers in New Mexico and Texas. They were known to have occurred from Española upstream from Cochiti Lake; in the downstream portions of the Chama and Jemez Rivers; throughout the Middle and Lower Rio Grande to the Gulf of Mexico; and in the Pecos River from Sumner Reservoir downstream to the confluence with the Rio Grande (Sublette *et al.* 1990, Bestgen and Platania 1991). The current distribution of the silvery minnow is limited to the Rio Grande between Cochiti Dam and Elephant Butte Reservoir, which amounts to approximately 5 percent of its historic range.

The construction of mainstem dams, such as Cochiti Dam and irrigation diversion dams have contributed to the decline of the silvery minnow. The construction of Cochiti Dam in particular has affected the silvery minnow by reducing the magnitude and frequency of flooding events that help to create and maintain habitat for the species. In addition, the construction of Cochiti Dam has resulted in degradation of silvery minnow habitat within the Cochiti Reach. Flow in the river at Cochiti Dam is now generally clear, cool, and free of sediment. There is relatively little channel braiding, and areas with reduced velocity and sand or silt substrates are uncommon. Substrate immediately downstream of the dam is often armored cobble (rounded rock fragments generally 3 to 12 in in diameter). Further downstream the riverbed is gravel with some sand material. Ephemeral tributaries including Galisteo Creek and Tonque Arroyo introduce sediment to the lower sections of this reach, and some of this is transported downstream with higher flows (U.S. Fish and Wildlife Service 1999, 2001). The Rio Grande below Angostura Dam becomes a predominately sand bed river with low, sandy banks in the downstream portion of the reach. The construction of Cochiti Dam also created a barrier between silvery minnow populations (U.S. Fish and Wildlife Service 1999). As recently as 1978, the silvery minnow was collected upstream of Cochiti Lake; however surveys since 1983 suggest that the fish is now extirpated from this area (U.S. Fish and Wildlife Service 1999).

Silvery minnow catch rates declined two to three orders of magnitude between 1993 and 2004. Additionally, relative abundance of silvery minnow declined from approximately 50 percent of the total fish community in 1995 to about 5 percent in 2004. However, in 2004, the October density of silvery minnow was significantly higher ($p < 0.05$) than in 2003 and autumnal catch rates increased by over an order of magnitude between those years. Silvery minnow catch rates in 2004 were comparable to those in 2001. Catch rates in 2005 were even higher. October catch

rates in 2005 (3,899) increased nearly 50 times over catch rates for 2004 (78) (Dudley *et al.* 2005).

Augmentation, throughout this period, likely sustained the silvery minnow population. Approximately, 1,000,000 silvery minnow have been released (primarily in the Angostura Reach) since 2000 (see Environmental Baseline). Captively propagated and released fish supplemented the native adult population and most likely also took advantage of the good spawning conditions of 2004 and 2005.

The silvery minnow was the most abundant taxon in October 2005 captures; it comprised about 72 percent of the total catch (Dudley *et al.* 2005). The species was nearly twice as abundant as the next most-abundant taxon (western mosquitofish). The increase in abundance of silvery minnow in 2005 has been comparable to previous years with above average precipitation (e.g., mid-1990s) (Dudley *et al.* 2005). These monitoring results from 2005 indicate that the status of the species has improved markedly compared to Fall of 2004.

Increased discharge in the Rio Grande during 2004 contrasted with the extended low-flow conditions observed throughout the Middle Rio Grande during 2003 and 2002. The timing of the 2004 runoff flow was typical of a flow increase that would normally occur at the onset of the spring runoff period. Elevated and extended flows during 2004 likely resulted in more favorable conditions for the growth and survivorship of newly hatched silvery minnow larvae. It is possible that even low numbers of eggs and larvae could have resulted in greatly increased recruitment success because of the inundation of shoreline habitats, abandoned side channels, and backwaters. Low velocity and shallow areas provide the warm and productive habitats required by larval fishes to successfully complete their early life history.

Spring runoff in 2005 was also above average, leading to a peak of over 6,000 cfs at Albuquerque and sustained high flows (> 3,000 cfs) for more than two months. These flows improved conditions for both spawning and recruitment. October monitoring indicated a significant increase in silvery minnow in the Middle Rio Grande, increasing to 3,899 total silvery minnow captured from 2 and 78 in 2003 and 2004, respectively.

In 2006, however, spring runoff was extremely low and although there were several peaks in the natural hydrograph in June, July, August, and September, only a small number of silvery minnow eggs were documented in June and July. October samples yielded only 166 silvery minnow. None of the silvery minnow collected were YOY, indicating poor recruitment, likely due to channel drying in June and July, after the late and minimal spawn (Dudley *et al.* 2006a).

Middle Rio Grande Distribution

Since the early 1990s, the density of silvery minnow generally increased from upstream (Angostura Reach) to downstream (San Acacia Reach). During surveys in 1999, over 98 percent of the silvery minnow captured were downstream of San Acacia Diversion Dam (Dudley and Platania 2002). This distributional pattern has been observed since 1994 (Dudley and Platania 2002) and is attributed to downstream drift of eggs and larvae and the inability of adults to repopulate upstream reaches because of diversion dams.

However, in 2004 and 2005, Dudley *et al.* (2005 and 2006a) found that this pattern reversed. Catch rates were highest in the Angostura Reach and approximately equal in the Isleta and San Acacia reaches. The Angostura Reach yielded the most silvery minnow (n=2,226) in 2004, followed by the Isleta Reach (n=442), and San Acacia Reach (n=371). Routine augmentation of silvery minnow in the Angostura Reach (nearly 900,000 since 2000), and the transplanting of silvery minnow rescued from drying reaches (approximately 770,000 since 2003) explains this change in pattern. Additionally, good spawning conditions (i.e., high and sustained spring runoff) throughout the Middle Rio Grande during April and May followed by wide-scale drying in the Isleta and San Acacia reaches from June-September exacerbated the skew. High spring runoff and perennial flow in the Angostura Reach appeared to result in relatively high survival and recruitment of larval and juvenile silvery minnow compared to previous drought years (2002-2003). In contrast, large portions of the Rio Grande south of Isleta Diversion Dam were dewatered in 2004 and young silvery minnow in these areas were either subjected to poor recruitment conditions (i.e., lack of nursery habitats during low-flows) or they were trapped in drying pools where they perished.

Sampling in early 2006 indicates populations are again higher downstream. Of the 6,143 silvery minnow caught in March 2006, 33 were found in the Angostura Reach, 2,445 were found in the Isleta Reach, and 3,665 were caught in the San Acacia Reach.

Reasons for Listing/Threats to Survival

The silvery minnow was federally listed as endangered for the following reasons:

1. Regulation of stream waters, which has led to severe flow reductions, often to the point of dewatering extended lengths of stream channel;
2. Alteration of the natural hydrograph, which impacts the species by disrupting the environmental cues the fish receives for a variety of life functions, including spawning;
3. Both the stream flow reductions and other alterations of the natural hydrograph throughout the year can severely impact habitat availability and quality, including the temporal availability of habitats;
4. Actions such as channelization, bank stabilization, levee construction, and dredging result in both direct and indirect impacts to the silvery minnow and its habitat by severely disrupting natural fluvial processes throughout the floodplain;
5. Construction of diversion dams fragment the habitat and prevent upstream migration;
6. Introduction of nonnative fishes that directly compete with, and can totally replace the silvery minnow, as was the case in the Pecos River, where the species was totally replaced in a time frame of 10 years by its congener the plains minnow (*Hybognathus placitus*); and

7. Discharge of contaminants into the stream system from industrial, municipal, and agricultural sources also impact the species (U.S. Fish and Wildlife Service 1993b, 1994).

These reasons for listing continue to threaten the species throughout its currently occupied range in the Middle Rio Grande.

Recovery Efforts

The final recovery plan for the silvery minnow was released in July 1999 (U.S. Fish and Wildlife Service 1999). The Recovery Plan has been updated and revised and a draft revised Recovery Plan (U.S. Fish and Wildlife Service 2007) was released for public comment on January 18, 2007 (72 FR 2301).

The draft revised Recovery Plan describes recovery goals for the silvery minnow and actions to complete these (U.S. Fish and Wildlife Service 2007). The three goals identified for the recovery and delisting of the silvery minnow are:

1. Prevent the extinction of the silvery minnow in the middle Rio Grande of New Mexico.
2. Recover the silvery minnow to an extent sufficient to change its status on the List of Endangered and Threatened Wildlife from endangered to threatened (downlisting).
3. Recover the silvery minnow to an extent sufficient to remove it from the List of Endangered and Threatened Wildlife (delisting).

Downlisting (Goal 2) for the silvery minnow may be considered when three populations (including at least two that are self-sustaining) of the species have been established within the historic range of the species and have been maintained for at least five years.

Delisting (Goal 3) of the species may be considered when three self-sustaining populations have been established within the historic range of the species and they have been maintained for at least ten years (U.S. Fish and Wildlife Service 2007).

ENVIRONMENTAL BASELINE

Under section 7(a)(2) of the ESA, when considering the effects of the action on federally listed species, we are required to take into consideration the environmental baseline. Regulations implementing the ESA (50 FR 402.02) define the environmental baseline as the past and present impacts of all Federal, State, or private actions and other human activities in the action area; the anticipated impacts of all proposed Federal actions in the action area that have undergone formal or early section 7 consultation; and the impacts of State and private actions that are contemporaneous with the consultation in progress. The environmental baseline defines the

current status of the species and its habitat in the action area to provide a platform to assess the effects of the action now under consultation.

Drought, as an overriding condition of the last decade in the southwest, is an important factor in the environmental baseline. However, stream conditions in 2004 and 2005 improved over previous years. The United States Geological Survey (USGS) in Albuquerque, New Mexico reported that stream flow conditions in 2005 were well above average to significantly above average statewide leading to a peak of over 6,000 cfs at Albuquerque and sustained high flows (> 3,000 cfs) for more than two months. These flows improved conditions for both spawning and recruitment.

The 2006 spring runoff was well below average because of lower than normal snowpack. In May 2006, year to date precipitation was well below average with the snow pack at 20 percent of average in the Rio Grande Basin. Fortunately, a strong monsoon season led to the wettest period of record in July and August. Consequently, only 26.5 mi of river dried in the summer of 2006, the lowest amount since 2001. Despite this monsoonal precipitation, reservoir levels continue to be below average across the state. It will take a least another year or two of well above average precipitation to reach pre-drought reservoir conditions.

The 2007 runoff was above average. Additionally, a one time deviation in Cochiti operations (Corps 2007) allowed managed releases of native flow during the spawn. Flows below Cochiti exceeded 3,000 cfs for ten days in May.

Since 1996, Reclamation has relied heavily on leases of SJC water to provide supplemental water by the Middle Rio Grande Endangered Species Act Collaborative Program to implement the 2003 Middle Rio Grande Water Operations Biological Opinion. Supplemental water has been used to create spawning pulses and recruitment flows for the silvery minnow and to meet minimum flow requirements for silvery minnow and flycatchers. From 1996-2003, Reclamation leased an average of 46,318 afy of SJC water from willing leasers to provide the average (1996-2001) of 63,109 afy of supplemental water needed for these purposes.

Status of the Species within the Action Area

The population of silvery minnows in the Action Area and throughout the Middle Rio Grande is highly variable over time (see Status of the Species). The most recent fall sample of silvery minnow reported silvery minnow in the action area at an estimated density of 1.33 per 100 meters squared (m^2) (Dudley et al. 2006).

Past actions have eliminated and severely altered habitat conditions for the silvery minnow. These actions can be broadly categorized as changes to the natural hydrology of the Rio Grande and changes to the morphology of the channel and floodplain. Other factors that influence the environmental baseline are water quality, the release of captively propagated silvery minnow; silvery minnow rescue efforts, on-going research efforts, and past projects in the Middle Rio Grande. Also of importance is the current drought, the expected weather pattern for the near future, and how it may affect flow in the Rio Grande. Each of these topics is discussed below.

Changes in Hydrology

There have been two primary changes in hydrology as a result of the construction of dams on the Rio Chama and Rio Grande that affect the silvery minnow: Loss of water and changes to the magnitude and duration of peak flows.

Loss of Water

Prior to measurable human influence on the system, up to the fourteenth century, the Rio Grande was a perennially flowing, aggrading river with a shifting sand substrate (Biella and Chapman 1977). There is now strong evidence that the Middle Rio Grande first began drying up periodically after the development of Colorado's San Luis Valley in the mid to late 1800s (Scurlock 1998). After humans began exerting more influence on the river, there are two documented occasions when the river became intermittent; during prolonged, severe droughts in 1752 and 1861 (Scurlock 1998). The silvery minnow historically survived low-flow periods because such events were infrequent and of lesser magnitude than they are today. There were also no diversion dams to block repopulation of upstream areas, the fish had a much greater geographical distribution, and there were oxbow lakes, cienegas, and sloughs associated with the Rio Grande that supported fish until the river became connected again.

Water management and use has resulted in a large reduction of suitable habitat for the silvery minnow. Agriculture accounts for 90 percent of surface water consumption in the Middle Rio Grande (Bullard and Wells 1992). The average annual diversion of water in the Middle Rio Grande by the Middle Rio Grande Conservation District (MRGCD) was 535,280 af for the period from 1975 to 1989 (U.S. bureau of Reclamation 1993). In 1990, total water withdrawal (groundwater and surface water) from the Rio Grande Basin in New Mexico was 1,830,628 af, significantly exceeding a sustainable rate (Schmandt 1993). Water withdrawals have not only reduced overall flow quantities, but also caused the river to become locally intermittent and/or dry for extended reaches. Irrigation diversions and drains significantly reduce water volumes in the river. However, the total water use (surface and groundwater) in the Middle Rio Grande by the MRGCD may range from 28 – 37 percent (S.S. Papadopulos & Associates, Inc. 2000; U.S. Geological Survey 2002). A portion of the water diverted by the MRGCD returns to the river and may be re-diverted (in some cases more than once) (Bullard and Wells 1992; MRGCD, *in litt.* 2003).

River reaches particularly susceptible to drying are immediately downstream of the Isleta Diversion Dam (river mile 169), a 5-mi reach near Tome (river miles 150-155), a 5-mi reach near the U.S. Highway 60 Bridge (river miles 127-132), and an extended 36-mi reach from near Brown's Arroyo (downstream of Socorro) to Elephant Butte Reservoir. Extensive fish kills, including tens of thousands of silvery minnow, have occurred in these lower reaches when the river has dried (C. Shroeder, Service, *pers. comm.* 2002). Since 1996, an average of 32 mi of the Rio Grande has dried, mostly in the San Acacia Reach. The most extensive drying occurred in 2003 and 2004 when 60 and 68.7 mi, respectively, were dewatered. Most documented drying events lasted an average of two weeks, before flows returned.

Predatory birds have been observed hunting and consuming fish from isolated pools during river intermittence (J. Smith, NMESFO, *pers. comm.* 2003). Although the number of fish present in any pool is unknown, it must be assumed that many of the fish preyed upon in these pools are silvery minnow. Thus, while some dead silvery minnow were collected during the shorter drying events, it is assumed that many more mortalities occurred than were documented.

Changes to Size and Duration of Peak Flows

Water management has also resulted in a loss of peak flows that historically initiated spawning. The reproductive cycle of the silvery minnow is tied to the natural river hydrograph. A reduction in peak flows and/or altered timing of flows may inhibit reproduction. Since completion of Elephant Butte Dam in 1916, four additional dams have been constructed on the Middle Rio Grande, and two have been constructed on one of its major tributaries, the Rio Chama (Scurlock 1998). Construction and operation of these dams, which are either irrigation diversion dams (Angostura, Isleta, San Acacia) or flood control and water storage dams (Elephant Butte, Cochiti, Abiquiu, El Vado), have modified the natural flow of the river. Mainstem dams store spring runoff and summer inflow, which would normally cause flooding, and release this water back into the river channel over a prolonged period of time. These releases are often made during the winter months, when low-flows would normally occur. The releases depart significantly from natural conditions, and can substantially alter the habitat. In spring and summer, artificially low-flows may limit the amount of habitat available to the species and may also limit dispersal of the species (U.S. Fish and Wildlife Service 1999).

Mainstem dams and the altered flows they create can affect habitat by preventing overbank flooding, trapping nutrients, altering sediment transport regimes, prolonging summer base flows, modifying or eliminating native riparian vegetation, and creating reservoirs that favor non-native fish species. These changes may affect the silvery minnow by reducing its food supply; altering its preferred habitat, preventing dispersal, and providing a continual supply of non-native fish that may compete with or prey upon them. Altered flow regimes may also result in improved conditions for other native fish species that occupy the same habitat, causing those populations to expand at the expense of the silvery minnow (U.S. Fish and Wildlife Service 1999).

In addition to providing a cue for spawning, flood flows also maintain a channel morphology to which the silvery minnow is adapted. The changes in channel morphology that have occurred from the loss of flood flows are discussed below.

Changes in Channel Morphology

Historically, the Rio Grande was sinuous, braided, and freely migrated across the floodplain. Changes in natural flow regimes, narrowing and deepening of the channel, and restraints to lateral channel migration (i.e., jetty jacks) adversely affected the silvery minnow. These effects result directly from constraints placed on channel capacity by structures built in the floodplain. These anthropogenic changes have and continue to degrade and eliminate spawning, nursery, feeding, resting, and refugia areas required for species' survival and recovery (U.S. Fish and Wildlife Service 1993a).

The active river channel within occupied habitat is being narrowed by the encroachment of vegetation, resulting from continued low-flows and the lack of overbank flooding. The lack of flood flows has allowed non-native riparian vegetation such as salt cedar and Russian olive to encroach on the river channel (U. S. Bureau of Reclamation 2001). These non-native plants are very resistant to erosion, resulting in narrowing of the channel. When water is confined to a narrower cross-section, its velocity increases and the ability to carry sediments is enhanced. Fine sediments such as silt and sand are carried away leaving coarser bed materials such as gravel and cobble. Habitat studies during the winter of 1995 and 1996 (Dudley and Platania 1996), demonstrated that a wide, braided river channel with low velocities resulted in higher catch rates of silvery minnow, and narrower channels resulted in fewer fish captured. The availability of wide, shallow habitats that are important to the silvery minnow is decreasing. Narrow channels have few backwater habitats with low velocities that are important for silvery minnow fry and YOY.

Within the current range of the silvery minnow, human development and use of the floodplain have greatly restricted the width available to the active river channel. A comparison of river area between 1935 and 1989 shows a 52 percent reduction, from 26,598 acres (10,764 ha) to 13,901 acres (5,626 ha) (Crawford *et al.* 1993). These data refer to the Rio Grande from Cochiti Dam downstream to the “Narrows” in Elephant Butte Reservoir. Within the same stretch, 234.6 mi of levees occur, including levees on both sides of the river. Analysis of aerial photography taken by Reclamation in February 1992, for the same river reach, shows that of the 180 mi of river, only 1 mi, or 0.6 percent of the floodplain has remained undeveloped.

Development in the floodplain, makes it difficult, if not impossible, to send large quantities of water downstream that would create low velocity side channels that the silvery minnow prefers. As a result, reduced releases have decreased available habitat for the silvery minnow and allowed encroachment of non-native species into the floodplain.

Water Quality

Both point (pollution discharges from a pipe) and non-point (diffuse sources of pollution) sources affect the Middle Rio Grande. Major point sources are WWTPs and feedlots. Major non-point sources include agricultural activities (e.g., fertilizer and pesticide application, livestock grazing), storm water run off, and mining activities.

Effluents from WWTPs contain contaminants that may affect the water quality of the river. For that reason the water quality of the effluent is extremely important. In the project area, the largest WWTP discharges are from two Rio Rancho facilities and Bernalillo (mean annual discharge flows are 80.4, 2.5, 0.9, and 0.7 cfs, respectively) (Bartolino and Cole 2002). Since 1998, total residual chlorine (chlorine) and ammonia, as nitrogen (ammonia), have been discharged unintentionally at concentrations that exceed protective levels for the silvery minnow. Chlorine concentrations of 0.013 mg/L can be harmful to the silvery minnow. Records also show that the monthly maximum concentration of ammonia during July 2001 was 14 mg/L. At pH 8 and water temperature of 25 °C, ammonia concentrations as low of 3.1 mg/L can be harmful to larval fathead minnow (U.S. Environmental Protection Agency 1999). The fathead minnow has

been suggested as a surrogate to evaluate the effects of various chemicals on the silvery minnow (Buhl 2002).

Although we do not have complete records for the Rio Rancho and Bernalillo WWTPs, in the summer of 2000, the Rio Rancho WWTP released approximately one million gallons of raw sewage into the Rio Grande. Chlorine treatment was maximized in an attempt to reduce the public health risk. Ammonia was reported at 37 mg/L on July 13, 2000, and at 17.1 mg/L on July 27, 2000 (City of Rio Rancho, *in litt.* 2000). Nonetheless, no violations of chlorine or ammonia effluent limits were recorded. This suggests that averaging measurements and/or the frequency of water quality measurements is insufficient to detect water quality situations that would be toxic to silvery minnow. The Rio Rancho WWTP now uses ultraviolet disinfection (Dee Fuerst, City of Rio Rancho, *pers. comm.* 2003). However, high concentrations of ammonia could still be discharged during an upset. Spills from the Rio Rancho City sewage system are treated with chlorine, which may lead to chlorine being flushed to the Rio Grande.

In addition to chlorine and ammonia, WWTP effluents may also include cyanide, chloroform, organophosphate pesticides, semi-volatile compounds, volatile compounds, heavy metals, and pharmaceuticals and their derivatives, which can pose a health risk to silvery minnow when discharged in concentrations that exceed the protective water quality criteria (J. Lusk, Service, *in litt.* 2003). Even if the concentration of a single element or compound is not harmful by itself, chemical mixtures may be more than additive in their toxicity to silvery minnow (Buhl 2002). The long-term effects and overall impacts of chemicals on the silvery minnow are not known.

Large precipitation events wash sediments and pollutants into the river from surrounding lands through storm drains and intermittent tributaries. Contaminants of concern to the silvery minnow that are frequently found in storm water include the metals aluminum, cadmium, lead, mercury, and zinc, organics such as oils, the industrial solvents trichloroethene and tetrachloroethene (TCE), and the gasoline additive methyl tert-butyl ether (U.S. Geological Survey 2001).

Harwood (1995) studied the North Floodway Channel (Floodway) of Albuquerque, which drains an urban area of about 90 square miles and crosses Pueblo of Sandia lands. He found that storm water contributions of dissolved lead, zinc, and aluminum were significant and posed a threat to the water quality of the Rio Grande. Because the Floodway crosses lands of the Pueblo of Sandia and enters their portion of the Rio Grande, they requested that the Environmental Protection Agency conduct toxicity tests on water in the Rio Grande collected below the Floodway. Aquatic crustaceans exposed to this water were found to have significant reproductive impairment and mortality when compared with controls. Additionally, larval fish also experienced significant mortality and/or narcosis when exposed to water and bed sediment collected from this same area on April 22, 2002 (http://oaspub.epa.gov/enviro/pcs_det_reports_detail_report?npdesid=NM0022250). This study indicates that storm water runoff can impact the water quality of the Rio Grande and the aquatic organisms that live in the river.

In a cooperative study, the New Mexico Environment Department (NMED) detected elevated polychlorinated biphenyl (PCBs) contamination of the San Jose Drain (NMED DOE Oversight Bureau Correspondence and Transmittal Letter, signed S. Yanicak, to G. Turner, DOE, Subject: 2002 – 2003 Cooperative Polychlorinated Biphenyl (PCB) Study Data, Dated June 6, 2006). The San Jose Drain empties into an area near the confluence of the Tijeras Arroyo (and SDC) with the Rio Grande. The PCB pollution was detected in sediment and storm water runoff and in fish tissue collected downstream. Concentrations of PCBs in fish tissues were elevated above the threshold by which fish consumption advisories would recommend that no fish be eaten by people (R. Ford-Schmid, NMED, electronic communication, June 24, 2004). The San Jose Drain empties into the Rio Grande in close proximity of the SDC Island Site increasing the probability that sediment forming the island may be contaminated with PCBs as well.

Sediment is the sand, silt, organic matter, and clay portion of the river bed, or the same material suspended in the water column. Ong *et al.* (1991) recorded the concentrations of trace elements and organochlorine pesticides in suspended sediment and bed sediment samples collected from the Middle Rio Grande between 1978 and 1988. These data were compared to numerical sediment quality criteria (Probable Effects Criteria [PEC]) proposed by MacDonald *et al.* (2000). According to MacDonald *et al.* (2000) most of the PEC provide an accurate basis for predicting sediment toxicity to aquatic life and a reliable basis for assessing sediment quality in freshwater ecosystems. Although the PEC were developed to assess bed (bottom) sediments, they also provide some indication of the potential adverse effects to organisms consuming these same sediments when suspended in the water column.

Semi-volatile organic compounds are a large group of environmentally important organic compounds. Three groups of compounds, polycyclic aromatic hydrocarbons (PAHs), phenols, and phthalate esters, were included in the analysis of bed sediment collected by the USGS (Levings *et al.* 1998). These compounds were abundant in the environment, are toxic and often carcinogenic to organisms, and could represent a long-term source of contamination. The analysis of the PAH data by Levings *et al.* (1998) show one or more PAH compounds were detected at 14 sites along the Rio Grande with the highest concentrations found below Albuquerque and Santa Fe. Polycyclic aromatic hydrocarbons and other semi-volatile compounds affect the sediment quality of the Rio Grande and may affect silvery minnow behavior, habitat, feeding, and health.

Pesticide contamination occurs from agricultural activities, as well as from the cumulative impact of residential and commercial landscaping activities. The presence of pesticides in surface water depends on the amount applied, timing, location, and method of application. Water quality standards have not been set for many pesticides, and existing standards do not consider cumulative effects of several pesticides in the water at the same time. Roy *et al.* (1992) reported that DDE, a degradation product of DDT, was detected most frequently in whole body fish collected throughout the Rio Grande. He suggested that fish in the lower Rio Grande may be accumulating DDE in concentrations that may be harmful to fish and their predators.

In addition to the compounds discussed above, several other constituents are present and affect the water quality of the Rio Grande. These include nutrients such as nitrates and phosphorus, total dissolved solids (salinity), and radionuclides. Each of these also has the potential to affect the aquatic ecosystem and health of the silvery minnow. As the river dries, pollutants will be concentrated in the isolated pools. Even though these pollutants do not cause the immediate death of silvery minnow, the evidence suggests that the amount and variety of pollutants present in the Rio Grande, could compromise their health and fitness (Rand and Petrocelli 1985).

Silvery Minnow Propagation and Augmentation

In 2000, the Service identified captive propagation as an appropriate strategy to assist in the recovery of the silvery minnow. Captive propagation is conducted in a manner that will, to the maximum extent possible, preserve the genetic and ecological distinctiveness of the silvery minnow and minimize risks to existing wild populations.

Silvery minnow are currently housed at four facilities in New Mexico including: the Dexter Fish Hatchery; New Mexico State University Coop Unit (Las Cruces); the Service's New Mexico Fishery Resources Office (NMFRO), and the City of Albuquerque's propagation facilities. These facilities are actively propagating and rearing silvery minnow. Silvery minnow are also held in South Dakota at the U.S. Geological Survey, Biological Resources Division Lab, but there is no active spawning program at this facility.

Since 2000 approximately 1,000,000 silvery minnow have been propagated using both adult wild silvery minnow and wild caught eggs and then released into the wild. Wild gravid adults are successfully spawned in captivity at the City's propagation facilities. Eggs are raised and released as larval fish. Marked fish have been released by the NMFRO since 2002 under a formal augmentation effort funded by the Collaborative Program. Silvery minnow are released into the Angostura Reach of the river near Alameda Bridge to ensure downstream repopulation. Eggs left in the wild have a very low survivorship and this ensures that an adequate number of spawning adults are present to repopulate the river each year. While hatcheries continue to successfully spawn silvery minnow, wild eggs are collected to ensure genetic diversity within the remaining population.

Genetic Diversity

Genetic data have been collected for the Rio Grande silvery minnow. The data set includes information from eight generations: one generation that preceded the precipitous decline that occurred in the last decade (1987), three generations that preceded the augmentation program (1999, 2000, 2001; Alò & Turner, 2005), and four generations that were supplemented with captively spawned and/or captively reared stocks (2002-2005; Turner et al. 2005). The following information was derived from studies of this data set.

Overall, mitochondrial (mt) DNA gene diversity declined nearly 18 percent between 1987 and 2005. In addition, researchers have identified other changes:

- There have been two sharp declines in mitochondrial haplotype diversity in the "wild"

Rio Grande silvery minnow population. The first occurred in 1999, the second in 2001. Each loss of diversity followed a sharp decline in abundance of Rio Grande silvery minnow: between 1995 and 1997, and again between 1999 and 2000, catch rates declined by an order of magnitude (Dudley et al. 2004). These declines in diversity coincided with extensive river drying in the San Acacia Reach of the Rio Grande.

- Microsatellite allelic diversity was less in 1999, but detected diversity was greater from 1999 to 2002. Although numerical abundance of the wild population continued to decline drastically after 2001, reaching extremely low levels in 2003, there was no substantial loss of allelic diversity over that time period.
- Declines in heterozygosity were recorded for the Rio Grande silvery minnow from 1987 to 1999 and between 2000 and 2002. However, heterozygosity increased between 2002 and 2005. Supplemental stocking with captive-reared wild caught-eggs between 2001 and 2003 may have temporarily alleviated loss of alleles and heterozygosity in the wild (Turner et al. 2004).

Permitted and/or Authorized Take

Take is authorized by section 10, and incidental take is permitted under section 7. These permits and/or authorizations are issued by the Service. Applicants for section 10 permits must also acquire a permit from the State to “take” or collect silvery minnow. Many of the permits issued under section 10 allow take for the purpose of collection and salvage of silvery minnow and eggs for captive propagation. Eggs, larvae, and adults are also collected for scientific studies to further our knowledge about the species and how best to conserve the silvery minnow. Because of the population decline from 2002-2004, the Service has reduced the amount of take permitted for voucher specimens in the wild.

Incidental take of silvery minnow is authorized through section 7 consultation associated with the 2003 BO, the City of Albuquerque Drinking Water Project (U.S. Fish and Wildlife Service 2004), the Isleta Island Removal Project, the Tiffany Plug Removal Project, and the Interstate Stream Commission’s (ISC) Habitat Restoration Project. In 2005 the Service revised the ITS for the 2003 BO using a formula that incorporates October monitoring data, habitat conditions during the spawn (spring runoff), and augmentation. Annual estimated take for the 2003 BO now fluctuates relative to the total number of RGSM found in October across 20 population monitoring locations.

Factors Affecting Species Environment within the Action Area

On the Middle Rio Grande, the following past and present federal, state, private, and other human activities, in addition to those discussed above, have affected the silvery minnow and its designated critical habitat:

1. Release of Carryover Storage from Abiquiu Reservoir to Elephant Butte Reservoir: The Army Corps of Engineers (Corps) consulted with the Service on the release of water during the winter of 1995. Ninety-eight thousand af of water was released from November 1, 1995 to March 31, 1996, at a rate of 325 cfs. This discharge is above the

historic winter flow rate. Substantial changes in the flow regime that do not mimic the historic hydrograph can be detrimental to the silvery minnow.

2. Corrales, Albuquerque, and Belen Levees: These levees contribute to floodplain constriction and habitat degradation for the silvery minnow. Levees at these sites result in a reduction in the amount and quality of suitable habitat for the silvery minnow.
3. Santa Ana River Restoration Project: In August 1999, Reclamation consulted with the Service on a restoration project located on Santa Ana Pueblo in an area where the river channel was incising and eroding into the levee system. This project included a Gradient Restoration Facility (GRF), channel re-alignment, bioengineering, riverside terrace lowering, and erodible bank lines. The primary component of the Santa Ana Restoration Project is the GRF, which should control river hydraulics upstream of its location and also river bed control. The GRF was designed to: (1) store more sand sediments at a stable slope for the current sediment supply; (2) decrease the velocities and depths and increase the width in the river channel upstream; (3) be hydraulically submerged at higher flows while simultaneously increasing the frequency and duration of overbank flows upstream; (4) provide velocities and depths suitable for passage of the silvery minnow through the structure; and (5) halt or limit further channel degradation upstream of its location. The channel re-alignment involved moving the river away from the levee system and over the grade control structure, and involves excavation of a new river channel and floodplain. Another significant component of the Santa Ana Restoration project is riverside terrace lowering for the creation of a wider floodplain. The bioengineering and deformable bank lines also assist in establishing the new channel bank and regenerating native species vegetation in the floodplain.
4. Creation of a Conservation Pool for Storage of Native Water in Abiquiu and Jemez Canyon Reservoirs and Release of a Spike Flow: The City created space (100,000 af) in Abiquiu Reservoir and the Corps created space in Jemez Canyon Reservoir to store Rio Grande Compact credit water for use in 2001, 2002, and 2003 for the benefit of listed species. The conservation pool was created with the understanding that the management of this water would be decided in later settlement meetings or during water operations conference calls. In addition, a supplemental release (spike) occurred in May 2001 to accommodate movement of sediment as a part of habitat restoration and construction on the Rio Grande and Jemez River on the Santa Ana Pueblo.
5. Programmatic Biological Opinions on the Effects of Actions Associated with the U.S. Bureau of Reclamation's, U.S. Army Corps of Engineers', and non-federal Entities' Discretionary Actions Related to Water Management on the Middle Rio Grande: In 2001 and 2003, the Service issued jeopardy biological opinions on the effects of water operations and management activities in the Middle Rio Grande on the silvery minnow and flycatcher. In 2002, the Service issued a jeopardy biological opinion for the silvery minnow. The current opinion, issued on March 17, 2003, contains one RPA with multiple elements. These elements set forth a flow regime in the Middle Rio Grande and describe habitat improvements necessary to alleviate jeopardy to both the silvery minnow

and flycatcher. For example, the elements require augmentation in the Rio Grande of an additional million silvery minnow over the life of the project and 484 acres of habitat restoration.

6. Albuquerque Drinking Water Project: The Drinking Water Project, involves the construction and operation of: (1) A new surface diversion dam north of Paseo del Norte Bridge, (2), conveyance of raw water from the point of diversion to the new water treatment plant, (3) a new water treatment plant on Chappell Road NE, (4) transmission of treated (potable) water to residential and commercial customers throughout the Albuquerque metropolitan area, and (5) aquifer storage and recovery. This consultation covers through 2060. During typical operations, the project will divert a total of 94,000 afy of raw water from the Rio Grande (47,000 afy of City SJC water and 47,000 afy of Rio Grande native water) at a near constant rate of about 130 cfs. Diversions of native water would be reduced if flows above the new diversion site were less than 260 cfs and all diversions would cease at levels below 195 cfs. Peak diversion operations will consist of up to 103,000 afy being diverted at a rate of up to 142 cfs. Consultation on this project was completed in 2004. Construction is currently underway with operations likely to begin in 2010.
7. Silvery minnow salvage and relocation: During river drying, the Service's silvery minnow salvage crew captures and relocates silvery minnow. Since 1996, approximately 770,000 silvery minnow have been rescued and relocated to wet reaches, the majority of which were released in the Angostura Reach. Studies are being conducted to determine survival rates for salvaged fish.
8. Habitat Restoration Projects: Several habitat restoration projects have been completed in the Albuquerque reach through the MRGESACP. These projects include two woody debris installation projects to encourage the development of pools and wintering habitat, and a river bar modification project south of the I-40 Bridge designed to create side and backwater channels on an existing bar as well as modify the top surface of the bar to create habitat over a range of flows. Additionally, in 2005, the ISC started a multi-year habitat restoration program that implements several island, bar, and bank line modification techniques throughout the Albuquerque Reach. Approximately 24 acres of habitat were restored in the Phase I. Phase II is scheduled to begin in winter 2007.
9. Bernalillo and Sandia Priority Site Projects: The Bernalillo and Sandia Priority Site Projects proposed by Bureau of Reclamation (Reclamation) are necessary to protect the integrity of the east levee and canal system along the Albuquerque Reach of the Middle Rio Grande between the U.S. Highway 550 bridge and the northern boundary of the Pueblo of Sandia. The banks of the river were close to the east levee and posed a potentially serious threat to project facilities and public health and safety. The Project created a secondary high flow channel, realigned the main river channel, and installed bendway weirs to reduce bank erosion threatening the levee.

10. Middle Rio Grande Conservation District: Improvements to physical and operational components of the irrigation system since 2001 have contributed to a reduction in the total diversion of water from the Rio Grande by the MRGCD. Prior to 2001, average diversions were 630,000 afy and now average 370,000 afy. The change was possible because of the considerable efforts of MRGCD to install new gages, automated gates at diversions, and scheduling and rotation of diversions among water users. The new operations reduce the amount of water diverted; however, this also reduces return flows that previously supported flow in the river. The river below Isleta Diversion Dam may be drier than in the past, but small inflows may contribute to maintaining flows.
11. Pilot Water Leasing Project: The City of Albuquerque and Albuquerque Bernalillo County Water Utility Authority with six conservation groups established a fund in February 2007 that will provide the opportunity to lease water from Rio Grande farmers and have that water remain in the river channel to support the silvery minnow. This program supports the need for reliable sources of water to support conservation programs as identified by the MRGESACP (2004).

Summary

The remaining population of the silvery minnow is restricted to approximately 5 percent of its historic range. Every year since 1996, there has been at least one drying event in the river that has negatively affected the silvery minnow population. The population is unable to expand its distribution because poor habitat quality and Cochiti Dam prevent upstream movement and Elephant Butte Reservoir blocks downstream movement (U.S. Fish and Wildlife Service 1999). Augmentation of silvery minnow with captive-reared fish will continue; however, continued monitoring and evaluation of these fish is necessary to obtain information regarding the survival and movement of individuals.

Water withdrawals from the river and water releases from dams severely limit the survival of silvery minnow. The consumption of shallow groundwater and surface water for municipal, industrial, and irrigation uses continues to reduce the amount of flow in the Rio Grande and eliminate habitat for the silvery minnow (U.S. Bureau of Reclamation 2003). However, under state law, the municipal and industrial users are required to offset the effects of groundwater pumping on the surface water system. The City of Albuquerque, for example, has been offsetting their surface water depletions with 60,000 afy returning to the river from the WWTP (U.S. Bureau of Reclamation 2003). The effect of water withdrawals means that discharge from WWTPs and irrigation return flows will have greater importance to the silvery minnow and a greater impact on water quality. Lethal levels of chlorine and ammonia have been released from the WWTPs in the last several years. In addition, a variety of organic chemicals, heavy metals, nutrients, and pesticides have been documented in storm water channels feeding into the river and contribute to the overall degradation of water quality.

Various conservation efforts have been undertaken in the past and others are currently being carried out in the middle Rio Grande. Silvery minnow abundance has increased over 2002-2003 population levels. However, the threat of extinction for the silvery minnow continues because of increased reliance on captive propagation, the fragmented and isolated nature of currently

occupied habitat, and the absence of silvery minnow in other parts of the historic range. The increased abundance of silvery minnow from 2004-2005 is a positive sign. Nevertheless, the threats that endanger this species are still present as evidenced by the sharp population decline of 2006.

EFFECTS OF THE ACTION

Effects of the action refer to the direct and indirect effects of an action on the species or designated critical habitat, together with the effects of other activities that are interrelated and interdependent with that action, which will be added to the environmental baseline. Indirect effects are those that are caused by the proposed action and are later in time, but are still reasonably certain to occur.

Silvery minnow are extremely likely to be present in the project area, as evidenced by data from one of the long-term silvery minnow monitoring sites, which is located approximately 0.3 miles downstream of the project area. Monitoring from October 2006 indicates that the approximate density of silvery minnow at the Rio Rancho monitoring site (River Mile 200) is 1.33 silvery minnow per 100 meters square (m^2) (Dudley *et al.* 2006).

Direct Adverse Effects

There are potential direct adverse effects to silvery minnow that may occur in the immediate project area during construction. Silvery minnow may be present when construction equipment is working along the bankline or crossing the river channel and/or when fill material is placed in the wetted channel.

The proposed action is likely to have direct short-term adverse effects on silvery minnow during construction of the proposed project. Heavy equipment will need to work at the river's edge during bankline construction and cross the active channel to haul fill across the river from the Pueblo of Sandia's MERES project site to the bioengineered bankline project area. It is expected that because as the heavy equipment displaces water, any fish in the area would flee. Fleeing from the disturbance represents an expenditure of energy that the fish would not have without the project. This form of harassment will be short in duration.

Equipment working on the bankline of the river, placement of fill material in the wetted channel, and equipment crossing wetted portions of the river channel may also potentially affect water quality. During construction, localized increases in turbidity and suspended sediments will likely occur. Direct effects from excess suspended sediments to a variety of fish include: alarm reaction, abandonment of cover, avoidance response, reduction in feeding rates, increase in coughing rate, increased respiration, physiological stress, poor condition, reduced growth, delayed hatching, and mortality (Newcombe and Jensen 1996).

Indirect Adverse Effects

The indirect effects of sediment mobilization due to the use of heavy equipment along the bankline, placement of fill material into the wetted channel, and crossing the river include the potential smothering and mortality of algae and aquatic invertebrates, depressed rates of growth,

reproduction, and recruitment or reduced physiological function of invertebrates. Decreases in primary production are associated with increases in sedimentation and turbidity and produce negative cascading effects through depleted food availability to zooplankton, insects, mollusks, and fish.

Designated Critical Habitat

This biological opinion does not rely on the regulatory definition of “destruction or adverse modification” of designated critical habitat at 50 CFR 402.02. Instead, we have relied upon the statutory provisions of the ESA to complete the following analysis with respect to designated critical habitat for the silvery minnow.

In the project area, deeper habitats along the bankline, as well as main channel areas used for equipment crossings, will be unusable during the construction period. These effects are expected to be short in duration (one month) and will be replaced by lower velocity habitats that may be inundated at high flows. Additionally, the proposed action, in concert with the Pueblo of Sandia’s MERES project, would create new habitat for the silvery minnow in the project area by increasing the channel complexity. However, the creation of secondary channels will be considered in the consultation on the MERES project.

As part of the Corrales Siphon Project, replacing the steep bankline with approximately one acre of bio-engineered bankline would create a gradient of benthic habitat at various water levels. Establishment of riparian vegetation on the stabilized bankline would provide nursery habitat when inundated during the spawning season. This habitat is necessary for development and hatching of eggs and the survival of the species from larvae to adult. Low-velocity habitat provides food, shelter, and sites for reproduction, which are essential for the survival and reproduction of Rio Grande silvery minnow. The wetted area at base flows will be reduced, but the habitat value of the remaining wetted area is likely to be improved. Replacing one acre of deep higher water velocity river channel with a bio-engineered bankline having a gradient of water velocities and depths is expected to increase in the value of designated critical habitat over the long term.

Therefore, we find that the effects to the function and conservation role of critical habitat relative to the entire designation are not significant because the impacts will be temporary and occur in a very small area relative to the overall critical habitat designation. Therefore, we conclude that the primary constituent elements of silvery minnow critical habitat will serve the intended conservation role for species with implementation of the proposed action.

Cumulative Effects

Cumulative effects include the effects of future state, tribal, local or private actions that are reasonably certain to occur in the action area considered in this biological opinion. Future federal actions that are unrelated to the proposed action are not considered in this section because they require separate consultation pursuant to section 7 of the Act. Cumulative effects include:

- Increases in development and urbanization in the historic floodplain that result in reduced peak flows because of the flooding threat. Development in the floodplain makes it more difficult, if not impossible, to transport large quantities of water that

- would overbank and create low velocity habitats that silvery minnow prefer.
- Increased urban use of water, including municipal and private uses. Further use of surface water from the Rio Grande will reduce river flow and decrease available habitat for the silvery minnow.
 - Contamination of the water (i.e., sewage treatment plants, runoff from small feed lots and dairies, and residential, industrial, and commercial development). A decrease in water quality and gradual changes in floodplain vegetation from native riparian species to non-native species (i.e., saltcedar) could adversely affect the silvery minnow and its habitat. Silvery minnow larvae require shallow, low velocity habitats for development. Therefore, encroachment of non-native species results in less habitat available for the silvery minnow.
 - Human activities that may adversely impact the silvery minnow by decreasing the amount and suitability of habitat include dewatering the river for irrigation; increased water pollution from non-point sources; habitat disturbance from recreational use, suburban development, and removal of large woody debris.
 - Wildfires and wildfire suppression in the riparian areas along the Rio Grande may have an adverse affect on silvery minnow. Wildfires are a fairly common occurrence in the bosque (riparian area) along the Rio Grande. Although fire retardant, which is toxic to aquatic species, is generally not used in close proximity to the Rio Grande, other fire suppression techniques, such as scooping water from the Rio Grande in large buckets, may harm silvery minnow. Silvery minnow could potentially be scooped up along with the water and dropped onto burning areas.

The Service anticipates that these types of activities will continue to threaten the survival and recovery of the silvery minnow by reducing the quantity and quality of habitat through continuation and expansion of habitat degrading actions.

CONCLUSION

After reviewing the current status of the silvery minnow and its designated critical habitat, the environmental baseline for the action area, the effects of the proposed action, and cumulative effects, it is the Service's biological opinion that the Corrales Siphon Project, as proposed in the January 2007, Biological Assessment is not likely to jeopardize the continued existence of the silvery minnow or result in adverse modification of designated critical habitat. The Corrales Siphon Project will create adverse effects to silvery minnow and its prey base, which are assumed to be present in the main channel construction zone, through the use of heavy equipment along the bankline and crossing the active channel, and placement of fill material in the wetted channel of the Rio Grande.

Although the proposed action has the potential to cause temporary adverse effects to water quality prey base for the silvery minnow during the construction phase, it is anticipated that these

impacts will not affect the function or intended conservation role of designated critical habitat relative to the conservation of the silvery minnow and to the overall critical habitat designation. The implementation of the proposed action is not expected to impede the survival or recovery of the silvery minnow within Middle Rio Grande or range-wide. In fact, the proposed project is designed to increase the quality of the habitat for silvery minnow in the long-term.

INCIDENTAL TAKE STATEMENT

Section 9 of the Act and federal regulation pursuant to section 4(d) of the Act prohibit the take of endangered and threatened species, respectively, without special exemption. Take is defined as to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture or collect, or to attempt to engage in any such conduct. Harm is further defined by the Service to include significant habitat modification or degradation that results in death or injury to listed species by significantly impairing essential behavioral patterns, including breeding, feeding, or sheltering. Harass is defined by the Service as intentional or negligent actions that create the likelihood of injury to listed species to such an extent as to significantly disrupt normal behavior patterns which include, but are not limited to, breeding, feeding or sheltering. Incidental take is defined as take that is incidental to, and not the purpose of, the carrying out of an otherwise lawful activity. Under the terms of section 7(b)(4) and section 7(o)(2), taking that is incidental to and not intended as part of the agency action is not considered to be prohibited taking under the Act provided that such taking is in compliance with the terms and conditions of this incidental take statement.

The measures described below are non-discretionary, and must be undertaken by Reclamation so that they become binding conditions of any grant or permit issued, as appropriate, for the exemption in section 7(o)(2) to apply. The action agency has a continuing duty to regulate the activity covered by this incidental take statement. If Reclamation (1) fails to assume and implement the terms and conditions or (2) fails to require adherence to the terms and conditions of the incidental take statement through enforceable terms that are added to the permit or grant document, the protective coverage of section 7(o)(2) may lapse. In order to monitor the impact of incidental take, Reclamation must report the progress of the action and its impact on the species to the Service as specified in the incidental take statement. [50 CFR §402.14(i)(3)]

Amount or Extent of Take Anticipated

The Service has developed the following incidental take statement based on the premise that the Corrales Siphon Project will be implemented as proposed. Take is expected in the form of harm and harass during: (1) the use of heavy equipment along the bankline and crossing the active channel of the Rio Grande; and (2) placement of fill materials directly into the wetted channel of the Rio Grande.

The Service anticipates that take in the form of harassment may affect up to 244 silvery minnow during project construction. We base this figure on the following assumptions. According to the BA, the area of disturbance along the bankline is approximately one acre. In addition, the

distance across the river from the MERES project site to the Corrales Siphon project site is approximately 220 ft and the swath of disturbance is 50 ft wide (30 ft for equipment and 10 ft on either side that silvery minnows would avoid). We estimate that this area would be newly disturbed at the start of each work day for up to 18 days. Any minnows present would be harassed and likely to stay away until activity ceases at the end of the day. This would add an additional 4.6 acres of disturbed area due to river crossings. Consequently a total of 6.6 ac of wetted habitat will be disturbed. Currently, the average density of silvery minnow in the project area is 1.33/100 m², therefore, approximately 244 silvery minnow will be harassed by construction, fill placement in the river, and movement of equipment through the water.

The Service notes that this number is only a best estimate of the amount of take that is likely under the proposed action. Thus, estimated incidental take may be modified from the above number should population monitoring information, data from silvery minnow rescue operations, or other research indicate substantial deviations from estimated values. In this case, further consultation, may be necessary. In addition, any take that may be associated with the Pueblo of Sandia's MERES project, which will be carried out in conjunction with the Corrales Siphon project, will be estimated under a separate section 7 consultation process.

Effect of the Take

The Service has determined that this level of anticipated take is not likely to result in jeopardy to the silvery minnow or destruction or adverse modification of designated critical habitat.

Reasonable and Prudent Measures

The Service believes the following Reasonable and Prudent Measures (RPMs) are necessary and appropriate to minimize impacts of incidental take of the silvery minnow due to activities associated with the proposed project.

1. Minimize take of silvery minnow due to construction and river crossing activities.
2. Manage for the protection of water quality from activities associated with the project.

Terms and Conditions

Compliance with the following terms and conditions must be achieved in order to be exempt from the prohibitions of section 9 of the ESA. These terms and conditions implement the Corrales Siphon Project described above and outline required reporting/monitoring requirements. These terms and conditions are non-discretionary.

To implement RPM 1, Reclamation shall:

- 1.1 Monitor for the presence/absence of silvery minnow in the project area prior to, at least four times during, and after construction.
- 1.2 Report survey results and any findings of injured or dead silvery minnow to the Service.
- 1.3 Use information collected from Terms and Conditions 1.1 – 1.2 to develop new or modify existing BMPs to minimize the adverse effects of this project and future

river maintenance projects. Any changes made to the project must be made in coordination with the Service to determine if reinitiation of consultation is required.

To implement RPM 2, Reclamation shall:

- 2.1 Transport fill materials with heavy equipment across the Rio Grande as few times as possible to minimize disturbance of sediments.
- 2.2 Avoid, to the extent practicable, crossing the wetted channel of the river with heavy equipment during flows greater than 900 cfs.
- 2.3 Monitor water quality, including turbidity and dissolved oxygen before, during, and after equipment operates in the river channel.
- 2.4 Use information collected from Term and Condition 2.3 to develop new or modify existing BMPs to minimize the adverse effects of this project and future river maintenance projects. Any changes made to the project must be made in coordination with the Service to determine if reinitiation of consultation is required.

CONSERVATION RECOMMENDATIONS

Section 7(a)(1) of the ESA directs Federal agencies to utilize their authorities to further the purposes of the ESA by carrying out conservation programs for the benefit of endangered and threatened species. Conservation recommendations are discretionary agency activities to minimize or avoid adverse effects of a proposed action on listed species or critical habitat, to help implement recovery plans, or to develop information. The Service recommends the following conservation activities:

1. Encourage adaptive management of flows and conservation of water to benefit listed species.
2. Work to further conduct habitat/ecosystem restoration projects in the Middle Rio Grande to benefit the silvery minnow.
3. Continue to use bio-engineering methods whenever possible in conjunction with river maintenance projects.

RE-INITIATION NOTICE

This concludes formal consultation on the action(s) described in the January 7, 2007 biological assessment. As provided in 50 CFR § 402.16, re-initiation of formal consultation is required where discretionary federal agency involvement or control over the action has been retained (or is authorized by law) and if: (1) The amount or extent of incidental take is exceeded; (2) new information reveals effects of the agency action that may affect listed species or designated critical habitat in a manner or to an extent not considered in this biological opinion; (3) the agency action is subsequently modified in a manner that causes an effect to the listed species or designated critical habitat not considered in this biological opinion, including if the Pueblo of Sandia's MERES project is not implemented in conjunction with the Corrales Siphon project, as

described; (3) adaptive management that includes additional earth work is needed to repair or maintain the project after the initial construction phase; or (4) a new species is listed or critical habitat designated that may be affected by the action. In instances where the amount or extent of incidental take is exceeded, any operations causing such take must cease pending re-initiation.

In future correspondence on this project, please refer to consultation number 22420-2007-F-0056. If you have any questions or would like to discuss any part of this biological opinion, please contact Jennifer Parody of my staff at (505) 761-4710.

Wally Murphy

cc:

Assistant Regional Director, Region 2 (ES), U.S. Fish and Wildlife Service, Albuquerque, NM
Regional Section 7 Coordinator, Region 2 (ES), U.S. Fish and Wildlife Service, Albuquerque, NM

LITERATURE CITED

- Ahlers, D. and R. Doster. Southwestern Willow Flycatcher Surveys and Nest Monitoring. Presentation and abstract given at the Middle Rio Grande Endangered Species Act Collaborative Program Second Annual Spring Symposium, April 16-17, 2007.
- Alò, D., and T.F. Turner. 2005. Effects of habitat fragmentation on effective population size in the endangered Rio Grande silvery minnow. *Conservation Biology* 19: 1138 – 1148.
- Bartolino, J.R. and J.C. Cole, 2002. Ground-Water Resources of the Middle Rio Grande Basin, New Mexico. U.S. Geological Survey Circular 1222. 132 pp.
- Bestgen, K. and S.P. Platania. 1991. Status and Conservation of the Rio Grande Silvery Minnow, *Hybognathus amarus*. *Southwestern Naturalist* 26(2):225–232.
- Bestgen, K. and D.R. Propst. 1994. Redescription, Geographic Variation, and Taxonomic Status of the Rio Grande Silvery Minnow, *Hybognathus amarus* (Girard, 1856). Contribution 69. Larval Fish Laboratory, Colorado State University.
- Biella, J., and R. Chapman (eds.). 1977. Archeological Investigations in Cochiti Reservoir, New Mexico. Vol. 1: A Survey of Regional Variability. Report submitted to the National Park Service, Santa Fe, for the U.S. Army Corps of Engineers, Albuquerque, New Mexico.
- Buhl, K. J. 2002. The Relative Toxicity of Waterborne Inorganic Contaminants to the Rio Grande Silvery Minnow (*Hybognathus amarus*) and Fathead Minnow (*Pimephales promelas*) in a Water Quality Simulating that in the Rio Range, New Mexico. Final Report to the U.S. Fish and Wildlife Service, Study No. 2F33 9620003. U.S. Geological Survey, Columbia Environmental Research Center, Yankton Field Research Station, Yankton SD.
- Bullard, T.F., and S.G. Wells . 1992. Hydrology of the Middle Rio Grande from Velarde to Elephant Butte Reservoir, New Mexico. U.S. Department of the Interior, U.S. Fish and Wildlife Service Research Publication 179.
- Caldwell, C. 2002. Hybridization Potential and Spawning Behavior of Rio Grande silvery Minnow (*Hybognathus amarus*) and Plains Minnow (*Hybognathus palcitus*). Interim report submitted to U.S. Fish and Wildlife Service Ecological Services Field Office, September 2002. 18 pp.
- CH2MHill. 2003. Hydrologic Effects of the Proposed City of Albuquerque Drinking Water Project on the Rio Grande and Rio Chama Systems: Updated for New Conservation and Curtailment Conditions. Technical Report. Prepared for City of Albuquerque Public Works Department Water Resources Strategy Implementation.
- Crawford, C., A. Cully, R. Leutheuser, M. Sifuentes, L. White, and J. Wilber. 1993. Middle Rio Grande Ecosystem; Bosque Biological Management Plan. Middle Rio Grande

- Dudley, R.K., and S.P. Platania. 1996. Rio Grande Silvery Minnow Winter Population Habitat Use Monitoring Project, April 1996. Summary of four trips (December 1995–March 1996). Report to the U.S. Army Corps of Engineers, Albuquerque. 12 pp.
- Dudley, R.K., and S.P. Platania. 1997. Habitat Use of the Rio Grande Silvery Minnow Report to U.S. Bureau of Reclamation, Albuquerque, New Mexico. 88 pp.
- Dudley, R.K., and S.P. Platania. 2002. Summary of Population Monitoring of Rio Grande Silvery Minnow (1994–2002). Report to New Mexico Ecological Services Field Office, September 10, 2002, Albuquerque, New Mexico. 14pp.
- Dudley, R.K., S.P. Platania, and S.J. Gottlieb. 2005. Summary of the Rio Grande Silvery Minnow Population Monitoring Program Results from December 2005. American Southwest Ichthyological Research Foundation, Albuquerque, New Mexico.
- Dudley, R.K., S.P. Platania, and S.J. Gottlieb. 2006. Summary of the Rio Grande Silvery Minnow Population Monitoring Program Results from October 2006. November 22, 2006. American Southwest Ichthyological Research Foundation, Albuquerque, New Mexico.
- Franklin, I. R. 1980. Evolutionary Change in Small Populations. Pages 135 – 148 in M. E. Soulé and B. A. Wilcox (editors) Conservation Biology: an evolutionary-ecological perspective. Sinauer Associates, Inc. Sunderland, Massachusetts.
- Harwood, A.K. 1995. The Urban Stormwater Contribution of Dissolved Trace Metal from the North Floodway Channel, Albuquerque, NM, to the Rio Grande. University of New Mexico, Water Resources Program, Professional Project Report.
- Levings, G.W., D.F. Healy, S.F. Richey, and L.F. Carter. 1998. Water Quality in the Rio Grande Valley, Colorado, New Mexico, and Texas, 1992-95. U.S. Geological Survey Circular 1162. <http://water.usgs.gov/pubs/circ1162> (viewed on May 18, 1998) .
- MacDonald, D.D., C.G. Ingersoll, and T.A. Berger. 2000. Development and Evaluation of Consensus-based Sediment Quality Guidelines for Freshwater Ecosystems. Archives of Environmental Toxicology and Chemistry 39:20–31.
- Middle Rio Grande Endangered Species Act Collaborative Program. 2004. Program Water Acquisition Final Position Paper. Water Acquisition and Management Subcommittee. July 12, 2004.
- Newcombe, C. P. and J. O. T. Jensen. 1996. Channel Suspended Sediment and Fisheries: A Synthesis for Quantitative Assessment of Risk and Impact. *North American Journal of Fisheries Management* 16:693-727
- Ong, K., T.F. O'Brien, and M.D. Rucker. 1991. Reconnaissance Investigation of Water Quality, Bottom Sediment, and Biota Associated with Irrigation Drainage in the Middle Rio Grande

- and Bosque del Apache National Wildlife Refuge in New Mexico 1988–89: U.S. Geological Survey Water-Resources Investigations Report 91-4036, Albuquerque, New Mexico.
- Pflieger, W. 1980. *Hybognathus nuchalis* Agassiz. In D. Lee, C. Gilbert, C. Hucutt, R. Jenkins, McCallister, and J. Stauffer, eds., Atlas of North American Freshwater Fishes. North Carolina State Museum of Natural History, Raleigh, North Carolina. 177 pp.
- Platania, S.P. 1995. Reproductive Biology and Early Life-history of Rio Grande Silvery Minnow, *Hybognathus amarus*. U.S. Army Corps of Engineers, Albuquerque, New Mexico. 23 pp.
- Platania, S. P. 2000. Effects of Four Water Temperatures Treatments On Survival, Growth, and Developmental Rates of Rio Grande Silvery Minnows, *Hybognathus amarus*, Eggs and Larvae. Report to U.S. Fish and Wildlife Service, Albuquerque, New Mexico.
- Platania, S.P., and C. Altenbach. 1998. Reproductive Strategies and Egg Types of Seven Rio Grande Basin Cyprinids. *Copeia* 1998(3): 559–569.
- Platania, S.P., and R. Dudley. 2001. Summary of Population Monitoring of Rio Grande Silvery Minnow (21–27 February 2001). Report to the Bureau of Reclamation and Corps of Engineers. Albuquerque. 7pp.
- Platania, S.P., and R. Dudley. 2005. 2004 Summary of Population Monitoring of Rio Grande Silvery Minnow. Report to the Bureau of Reclamation and Corps of Engineers. Albuquerque. 193pp.
- Platania, S.P., and C.W. Hoagstrom. 1996. Response of Rio Grande Fish Community to and Artificial Flow Spike: Monitoring Report Rio Grande Silvery Minnow Spawning Peak Flow. New Mexico Ecological Services State Office, Albuquerque, New Mexico.
- Platania, S.P., Michael A. Farrington, W. Howard Brandenburg, Sara J. Gottlieb, and Robert K. Dudley 2003. Movement Patterns Of Rio Grande Silvery Minnow *Hybognathus amarus*, in the San Acacia Reach of the Rio Grande During 2002 Final Report. 38 pp.
- Rand, G.M., and Petrocelli, S.R. 1985. Fundamentals of Aquatic Toxicology B Methods and Applications. Hemisphere Publishing Corporation, New York. 666 pp.
- Roy, Richard, T.F. O'Brien, and M. Rusk-Maghini. 1992. Organochlorine and Trace Element Contaminant Investigation of the Rio Grande, New Mexico. U.S. Fish and Wildlife Service, New Mexico Ecological Services Office, Albuquerque, NM. 39 pp.
- S.S. Papadopulos & Associates, Inc. 2000. Middle Rio Grande Water Supply Study. Boulder, CO: August 4, 2000.
- Schmandt, J. 1993. Water and Development in the Rio Grande/Río Bravo Basin. University of

Texas Press, Austin, Texas.

Scurlock, D. 1998. From the Rio to the Serria: An Environmental History of the Middle Rio Grande Basin. USDA Rocky Mountain Research Station, Fort Collins, Colorado, 80526.

Smith, J.R. 1999. Summary of Easy Egg Catching in the LFCC in the 9 Mile Study Reach during Spring 1998 Operation. U.S. Fish and Wildlife Service Report Submitted to the U.S. Bureau of Reclamation, Albuquerque, New Mexico on April 28, 1999.

Sublette, J., M. Hatch, and M. Sublette. 1990. The Fishes of New Mexico. Univ. New Mexico Press, Albuquerque, New Mexico. 393 pp.

Turner T.F., and M. J. Osborne. 2004. Genetic Consequences of Supportive Breeding in the Endangered Rio Grande Silvery Minnow (*Hybognathus amarus*): Genetic Evaluation of Wild and Captively Reared and Propagated Stocks, 1999-2004. An annual report submitted to the Middle Rio Grande ESA Collaborative Program, 31pp

U.S. Army Corps of Engineers. 2007. Environmental Assessment for a Temporary Deviation in the Operation of Cochiti Dam, Sandoval County, New Mexico. 69 pp.

U.S. Bureau of Reclamation. 1993. Final Supplement to the Final Environmental Impact Statement-River Maintenance program for the Rio Grande-Velarde to Caballo Dam-Rio Grande and Middle Rio Grande projects, New Mexico. 140 pp.

U.S. Bureau of Reclamation. 2003. Biological Assessment of Bureau of Reclamation's Water and River Maintenance Operations, Army Corps of Engineers' Flood Control Operations, and Related Non-Federal Actions on the Middle Rio Grande, New Mexico, March 1, 2003 – February 28, 2013. February 19, 2003.

U.S. Bureau of Reclamation. 2006. 2006 Middle Rio Grande Annual Operating Plan Report. Albuquerque Area Office, Water Management Division. June 2006. Albuquerque, New Mexico.

U.S. Bureau of Reclamation. 2007. Middle Rio Grande Project Corrales Siphon Biological Assessment. January 2007. Submitted to the U.S. Fish and Wildlife Service by the U.S. Bureau of Reclamation, Albuquerque Area Office, Albuquerque, New Mexico.

U.S. Fish and Wildlife Service. 1993a. Proposal Rule to List the Rio Grande Silvery Minnow as Endangered, with Critical Habitat. 58 Federal Register 11821-11828.

U.S. Fish and Wildlife Service. 1993b. Notice of 12-month Petition Finding/Proposal to List *Empidonax traillii extimus* as an Endangered Species, and to Designate Critical Habitat. Federal Register 58:39495–39522..

- U.S. Fish and Wildlife Service. 1994. Endangered and Threatened Wildlife and Plants; Final Rule to list the Rio Grande Silvery Minnow as an Endangered Species. Federal Register 59:36988–37001.
- U.S. Fish and Wildlife Service. 1999. Rio Grande Silvery Minnow (*Hybognathus amarus*) Recovery Plan. Region 2, U.S. Fish and Wildlife Service, Albuquerque, New Mexico. 138 pp.
- U.S. Fish and Wildlife Service. 2001. Programmatic Biological Opinion on the Effects of Actions Associated with the U.S. Bureau of Reclamation's, U.S. Army Corps of Engineers', and non-Federal Entities' Discretionary Actions Related to Water Management on the Middle Rio Grande, New Mexico, June 29, 2001.
- U.S. Fish and Wildlife Service. 2003. Biological and Conference Opinions on the Effects of Actions Associated with the Programmatic Biological Assessment of Bureau of Reclamation's Water and River Maintenance Operations, Army Corps of Engineers' Flood Control Operation, and Related Non-Federal Actions on the Middle Rio Grande, New Mexico. March 17, 2003. Southwest Region, Albuquerque, New Mexico.
- U.S. Fish and Wildlife Service. 2004. Biological Opinion on the Effects of Actions Associated with the Programmatic Biological Assessment (BA) for the City of Albuquerque Drinking Water Project. February 13, 2004. New Mexico Ecological Services Field Office. Albuquerque, New Mexico.
- U.S. Fish and Wildlife Service. 2005. Endangered and Threatened Wildlife and Plants; Designation of Critical Habitat for the Southwestern Willow Flycatcher (*Empidonax traillii extimus*). Federal Register 70:60935–60984.
- U.S. Fish and Wildlife Service. 2007. Rio Grande Silvery Minnow (*Hybognathus amarus*) Draft Revised Recovery Plan. January 2007. Southwest Region, Albuquerque, New Mexico.
- U.S. Geologic Survey. 2002. Ground-water resources of the Middle Rio Grande basin, New Mexico. Circular 1222.
- U.S. Geologic Survey. 2001. Selected Findings and Current Perspectives on Urban and Agricultural Water Quality by the National Water-Quality Assessment Program, FS-047-<http://water.usgs.gov/pubs/FS/fs-047-01/pdf/fs047-01.pdf>
- Watts, H.E., C.W. Hoagstrom, and J.R. Smith. 2002. Observations on Habitat Associated with Rio Grande Silvery Minnow, *Hybognathus amarus* (Girard). Submitted to U.S. Army Corps of Engineers, Albuquerque District and City of Albuquerque Water Resources Division, June 28, 2002.

