LIST OF TABLES AND FIGURES

Chapter 2: Historical Background, Terminology, Evolution of Recommendations, and Measurement

Table 2-1. Glossary of terms

Table 2-2. Selected physical activity recommendations in the United States (1965–1996)

Table 2-3. Assessment procedures and their potential use in epidemiologic research

Table 2-4. Classification of physical activity intensity, based on physical activity lasting up to 60 minutes

Table 2.5. Correlation of two survey instruments with physiologic measures of caloric exchange

Chapter 3: Physiologic Responses and Long-Term Adaptations to Exercise

Table 3-1. A summary of hormonal changes during an episode of exercise

Table 3-2. A hypothetical example of alterations in selected physiological variables consequent to a 6-month endurance training program in a previously sedentary man compared with those of a typical elite endurance runner

Figure 3-1. Changes in cardiac output (A), heart rate (B), and stroke volume (C) with increasing rates of work on the cycle ergometer

Figure 3-2. Changes in arterial and mixed venous oxygen content with increasing rates of work on the cycle ergometer

Figure 3-3. Changes in oxygen uptake and blood lactate concentrations with increasing rates of work on the cycle ergometer

Figure 3-4. Changes in $\dot{V}O_2$ max with increasing age from 6 to 18 years of age in boys and girls

Figure 3-5. Changes in $\dot{V}O_2$ max with aging, comparing an active population and sedentary population (the figure also illustrates the expected increase in $\dot{V}O_2$ max when a previously sedentary person begins an exercise program)

Chapter 4: The Effects of Physical Activity on Health and Disease

Table 4-1. Population-based studies of association of physical activity or cardiorespiratory fitness with total cardiovascular diseases

Table 4-2. Population-based studies of association of physical activity or cardiorespiratory fitness with coronary heart disease

Table 4-3. Population-based studies of association of physical activity with stroke (CVA)

Table 4-4. Population-based cohort studies of association of physical activity with hypertension

Table 4-5. Epidemiologic studies of leisure-time or leisure-time plus occupational physical activity and colon cancer

Table 4-6. Epidemiologic studies of leisure-time or leisure-time plus occupational physical activity and hormone-dependent cancers in women

Table 4-7. Epidemiologic studies of leisure-time or total physical activity or cardiorespiratory fitness and prostate cancer

Table 4-8. Cohort studies of association of physical activity with non–insulin-dependent diabetes mellitus (NIDDM)

Table 4-9. Longitudinal population-based studies of physical activity as related to depressive symptoms

Table 4-10. Duration of various activities to expend 150 kilocalories for an average 70 kg adult

Chapter 5: Patterns and Trends in Physical Activity

Table 5-1. Sources of national and state-based data on physical activity

Table 5-2. Percentage of adults aged 18+ years reporting no participation in leisure-time physical activity, by various demographic characteristics, National Health Interview Survey (NHIS), Third National Health and Nutrition Examination Survey (NHANES III), and Behavioral Risk Factor Surveillance System (BRFSS), United States

Table 5-3. Percentage of adults aged 18+ years reporting participation in no activity; regular, sustained activity; and regular, vigorous activity, by state,* Behavioral Risk Factor Surveillance System (BRFSS), 1994, United States

Table 5-4. Percentage of adults aged 18+ years reporting participation in regular, sustained physical activity (5+ times per week for 30+ minutes per occasion), by various demographic characteristics, National Health Interview Survey (NHIS) and Behavioral Risk Factor Surveillance System (BRFSS), United States

Table 5-5. Percentage of adults aged 18+ years participating in regular, vigorous physical activity (3+ times per week for 20+ minutes per occasion at 50+ percent of estimated age- and sex-specific maximum cardiorespiratory capacity) by various demographic characteristics, National Health Interview Survey (NHIS) and Behavioral Risk Factor Surveillance System (BRFSS), United States

Table 5-6. Percentage of adults aged 18+ years reporting participation in selected common physical activities in the prior 2 weeks, by sex and age, National Health Interview Survey (NHIS), United States, 1991

Table 5-7. Percentage of adults aged 18+ years reporting participation in any strengthening activities or stretching exercises in the prior 2 weeks, by various demographic characteristics, National Health Interview Survey (NHIS), United States, 1991

Table 5-8. Trends in the percentage of adults aged 18+ years reporting participation in no activity; regular, sustained activity; and regular, vigorous activity, by sex, National Health Interview Survey (NHIS) and Behavioral Risk Factor Surveillance System (BRFSS), from 1985–1994

Table 5-9. Percentage of young people reporting no participation in vigorous or moderate physical activity during any of the 7 days preceding the survey, by demographic group, 1992 National Health Interview Survey-Youth Risk Behavior Survey (NHIS-YRBS) and 1995 Youth Risk Behavior Survey (YRBS), United States

Table 5-10. Percentage of young people reporting participation in vigorous physical activity during 3 or more of the 7 days preceding the survey, by demographic group, 1992 National Health Interview Survey-Youth Risk Behavior Survey (NHIS-YRBS) and 1995 Youth Risk Behavior Survey (YRBS), United States

Table 5-11. Percentage of young people reporting participation in strengthening or toning activities during 3 or more of the 7 days preceding the survey, by demographic group, 1992 National Health Interview Survey-Youth Risk Behavior Survey (NHIS-YRBS) and 1995 Youth Risk Behavior Survey (YRBS), United States

Table 5-12. Percentage of young people reporting participation in stretching activities during 3 or more of the 7 days preceding the survey, by demographic group, 1992 National Health Interview Survey-Youth Risk Behavior Survey (NHIS-YRBS) and 1995 Youth Risk Behavior Survey (YRBS), United States

Table 5-13. Percentage of young people reporting participation in walking or bicycling for 30 minutes or more during 5 or more of the 7 days preceding the survey, by demographic group, 1992 National Health Interview Survey-Youth Risk Behavior Survey (NHIS-YRBS) and 1995 Youth Risk Behavior Survey (YRBS), United States

Table 5-14. Percentage of young people reporting participation in selected physical activities during 1 or more of the 7 days preceding the survey, by demographic group, 1992 National Health Interview Survey-Youth Risk Behavior Survey (NHIS-YRBS), United States

Table 5-15. Percentage of students in grades 9–12 reporting enrollment in physical education class, daily attendance in physical education class, and participation in exercise or sports for at least 20 minutes during an average physical education class, by demographic group, 1995 Youth Risk Behavior Survey (YRBS), United States

Table 5-16. Percentage of students in grades 9–12 reporting participation on at least one sport team run by a school or by other organizations during the year preceding the survey, by demographic group, 1995 Youth Risk Behavior Survey (YRBS), United States

Figure 5-1. Percentage of adults aged 18+ years reporting no participation in leisure-time physical activity by sex and age

Figure 5-2. Percentage of adults aged 18+ years reporting no participation in leisure-time physical activity by month

Figure 5-3. Trends in leisure-time physical activity of adults aged 18+ years, NHIS

Figure 5-4. Trends in the percentage of adults aged 18+ years participating in no leisure-time activity, BRFSS

Chapter 6: Understanding and Promoting Physical Activity

Table 6-1. Summary of theories and models used in physical activity research

Table 6-2. Studies of interventions to increase physical activity among adults

Table 6-3. Summary of progress toward *Healthy People 2000* objective 1.10

Table 6-4. Percentage of all physical education courses in which more than one class period was devoted to each activity, by activity, School Health Policies and Programs Study, 1994

Table 6-5. Studies of interventions to increase physical activity among children and adolescents

Figure 6-1. Results of the Minnesota Heart Health Program on physical activity. Graph compares the percentage of respondents reporting regular physical activity in intervention cities and the secular trend estimated from control cities

Figure 6-2. Moderate-to-vigorous and vigorous physical activity observed during Child and Adolescent Trial for Cardiovascular Health (CATCH) physical education classes

INDEX

A

Abdominal fat, 135
Absolute intensity scale, 66
Acid-base equilibrium, 62
Adenosine diphosphate (ADP), 65
Adenosine triphosphate (ATP), 65, 66
Adipocytes, 128
Adipose tissue, 128, 133, 134
Adolescents and physical activity
  bicycling activities, 200, 205
  bone mass development, 131, 132
  breast cancer incidence, 117
  cardiovascular disease risk factors, 91, 102
  high school physical education, 204, 205
  injuries, 142
  no regular activity, 189
  obesity, 43, 47, 133
  recommendations, 28–29
  regular, vigorous activity, 191, 196–197, 205
  school-based interventions, 6, 236–243
  sports team participation 200
  stretching activities, 194, 200–201
  surveys, 175, 205
  trends, 8
  walking activities, 200, 205
Adults and physical activity
  assessment procedures, 30
  cardiovascular response, 75
  communications strategies, 229–230, 231
  community approaches, 227–229, 234
  determinants, modifiable, 215, 234
  exercise enjoyment, 215, 216
  health care settings, 226–227, 242
  individual approaches, 217, 226, 234
  promotion, 217, 234
  pulmonary ventilation rates in untrained, 64
  recommendations, 24–27, 43
  self-efficacy, 214–215, 217, 248
  self-monitoring, 217, 226, 234
  social support, 216, 226
  surveys, 175
  trends, 8
  worksites, 229–231, 234, 229
Aerobic exercise, 66
Aerobics Center Longitudinal Study, 86
Affective disorders, 135
African Americans
  adult activity interventions, 232, 236, 232
  bicycling activities by, 203
  Fitness Through Churches Project, 245
  high school physical education enrollment, 205
  no regular physical activity, 177, 192, 195
  physical activity trends in, 8
  regular, sustained physical activity, 183
  regular, vigorous physical activity, 185, 187, 196–197
  sports team participation, 200
  strengthening activities by, 189, 191, 198, 193
  stretching activities by, 191, 194, 201
  walking activities by, 203
Age factors, 74–76
  cardiorespiratory capacity and, 187
  exercise intensity and, 31–33
  hypertension and, 103
  no regular activity and, 177, 179, 192, 195
  physical activity and mortality rates, 86
  regular, sustained activity and, 183
  regular, vigorous activity and, 185, 187
  weight gain prevention and, 133, 232–233
Agility. *See* Psychomotor performance
Alabama
  Physical Activity for Risk Reduction project, 232

Amenorrhea, 131, 143
American Academy of Pediatrics, 28
American Alliance for Health, Physical Education, Recreation and Dance, 3, 244
American Association of Cardiovascular and Pulmonary Rehabilitation, 23
American Association of Health, Physical Education, and Recreation (AAHPER), 18
  Health Related Physical Fitness Test, 18
  Youth Fitness Test, 18
American Cancer Society (ACS), 112
American College of Sports Medicine (ACSM), 5
  cardiorespiratory fitness (endurance) and, 4, 20
  consultation for report by, 3
  Guidelines for Exercise Testing and Prescription, 23
  National Coalition for Promoting Physical Activity, 244
  physical activity recommendations, 33, 148
American Diabetes Association, 127
American Heart Association (AHA), 244
  cardiorespiratory endurance (fitness) and, 4
  consultation for report by, 3
  National Coalition for Promoting Physical Activity, 244
  physical activity counseling recommendation, 244
American Hospital Association, 23
American Medical Association (AMA)
  exercise and physical fitness, 17
  Guidelines for Adolescent Preventive Services (GAPS), 28, 242
  Health and Fitness Program, 17
  physical activity counseling recommendation, 244
Anemia, 143
Angina pectoris, 45, 110–112, 143
Anovulation, 143
Anxiety, 8, 137, 150
Anxiety disorders, 135, 136
Aortic aneurysms, 103
Aortic valve stenosis, 45
Appetite, 135
Arrhythmias, 110, 112, 143
Arterial baroreflex, resetting of, 63
Arterial-mixed venous oxygen (A-$\bar{v}O_2$), 62, 64, 70, 75–77
Arterial vasodilatation, 111
Arthritis, 7, 129–130, 142
Arthritis, rheumatoid, 129
Asthma, 143
Atherosclerosis, 5, 102, 103, 110–111, 128
Australian Heart Week (1990) campaign, 233

B

B-lymphocytes. *See* Lymphocytes
Balance. *See* Equilibrium
Bank of America physical activities program, 230
Basal metabolic rate, 66
Bed rest, prolonged, 71, 72
Behavioral sciences, adult
  ecological perspective, 214–215, 244
  health-belief model, 213, 217
  intervention studies. *See under* Intervention studies (adolescent; adult; children)
  learning theories, 211, 214, 226, 228, 230
  planned behavior theory, 213–214, 226
  reasoned action theory, 213–214, 217
  relapse prevention model, 213, 226, 228
  social learning (cognitive) theory, 214, 217, 226, 228, 230, 235, 236
  social support, 220, 214, 227, 228
  transtheoretical model, 213, 235
Behavioral sciences, children and adolescents, 234–243
  planned behavior theory, 213
  reasoned action theory, 213
  social learning (cognitive) theory, 214
Biogenic monoamines, 141
Blood flow, 63, 64, 65, 71, 111, 112, 128

Blood pressure, 16, 71, 90, 110, 111, 145
adaptation to exercise, 73
diastolic, 63, 70, 72, 102, 110
end-diastolic volume, 71, 72
mean arterial, 63
response to resistance exercise, 65
systolic, 63, 64, 70, 102, 110
See related Hypertension; Hypotension
Blood volume, 70, 71, 72
Body composition, 21, 22, 35, 54, 134
Body fat, 7, 35, 102, 128–129, 134, 135, 145
Body mass index, 35, 90, 102, 126–127, 133, 134
Body surface area to mass ratio, 73–74
Body temperature, 62, 64, 141
Body weight, loss practices, 50, 44
Bone density, 69, 72, 73, 75, 130, 131, 132
Bone marrow, 67
Breast cancer, 7, 117–119, 123
British Association for the Advancement of Science, 19
Building Your Fitness Futures program, 232

C

Calcium, balance, 72
Caloric expenditure, 147
Calorimetry, 21, 29, 32
indirect, 21, 32
Cancer, 43, 67, 149
breast, 7, 117–119
colon, 4, 5, 7, 114, 144, 145, 149
endometrial, 7, 120–121, 149
hormone-dependent in women, 117–121
mortality incidence, 113
ovarian, 7, 120, 149
physical activity and, 7
prostate, 7, 121–122, 124–125
testicular, 7, 124, 149
rectal, 7, 113, 116
uterine, 117, 120–121
Capillaries, 71
density of, 63, 65, 73
endurance training and, 69
Carbohydrate, adenosine triphosphate production, 66
Carbon dioxide (CO_2), 18, 32, 61–62, 66
Cardiac output ($\dot{Q}$), 62,71
contribution to mean arterial blood pressure, 63
and heart rate, 62
maximal ($\dot{Q}$ max), 62, 70, 71
rest vs. exercise, 63
stroke volume, 62, 63, 65
Cardiomyopathy, hypertrophic, 45
Cardiorespiratory endurance (fitness), 4, 6, 17
age and, 187
cholesterol, total, and, 102
epidemiologic studies of, 131, 137, 143, 147, 182, 187, 201–205
interventions, 244
measurement of, 19–20, 32–37
multiple sclerosis and, 233
population-based studies, 85-90
prostate cancer and, 121–125
regular, intermittent exercise, 5
sex factors and, 187
See also Physical fitness
Cardiovascular diseases, 87
of adolescents, interventions for, 236–243
community-based intervention programs for, 227–229
of children, interventions for, 236–243
diabetes risk factors for, 127
health care interventions for, 242
myocardial infarction, 5, 43, 45, 112, 143
physical activity and, 43–48, 43–45, 47
Cardiovascular system adaptations, 5, 7, 19
to exercise, 7, 21, 61–62, 65, 70, 71, 87
myocardial wall stress in, 63–64, 71
Cartilage, 130, 143
Catecholamine, 64, 66, 74

Causality, 144–145
Centers for Disease Control and Prevention (CDC), 5
  Behavioral Risk Factor Surveillance System (BRFSS), 175, 205
  exercise recommendations, 23, 28, 33, 148
  Guidelines for School and Community Health Programs to Promote Physical Activity Among Youths, 237, 244
  School Health Policies and Programs Study, 236–237
  Youth Risk Behavior Survey, 175, 176, 189, 193–199, 205
Cerebrovascular disorders, 7, 47, 45, 102–103, 104–107, 110
Child and Adolescent Trial for Cardiovascular Health (CATCH) study, 239, 244
Children and physical activity, 75
  assessment procedures, 29
  behavioral research, 234–243
  bone mass development, 131, 132
  cardiovascular factors, 75, 91, 102
  environmental factors, 73–74
  goals, 28, 43
  hypertension, 87
  injuries, 142
  minimum health standard, 17–18, 19
  no regular activity, 4
  obesity, 43, 47, 133, 134
  obesity interventions, 248
  regular, vigorous activity, 191
  school-based interventions, 6
  school program interventions, 236–243
Cholesterol, 19, 23, 47, 91, 102, 110–111
Church programs, 245
Cognition, 135, 141, 142
College Alumni Study, 36
Colon cancer, 4, 5, 7, 113–117, 146, 149
Colorectal cancer, 113
Communications interventions, 230–231
Community-based programs, 6, 227–229, 245–246
Community behavioral approaches, 227–229
Coronary artery
  perfusion pressure increase, 64
  vasodilation of, 64
Coronary artery bypass, 45
Coronary circulation, 63–64
Coronary disease, 16, 23, 28, 35, 37, 49, 47, 133, 140, 144–147, 149
  inverse association with physical activity, 91
  physical activity and, 4, 5, 7, 87, 90–91
  population-based studies, 92–101
Coronary plaque, 110, 111, 112
Crime, 246–247

D

Dehydration, 75, 143
Dementia, 136
Depression, 8, 135, 136, 140, 150
Detraining, 21, 61, 72
Diabetes mellitus, 4–6, 28, 35, 37, 43, 90, 125, 133, 144–149, 232
Diabetic retinopathy, 128
Diastolic blood pressure. *See under* Blood pressure
Diet, 5, 12–13, 116, 127, 128, 134, 232–233
Dietary Guidelines for Americans, 5, 28, 244
Disability. *See* Physically handicapped
Disuse atrophy, 72–73
Dyslipoproteinemia, 133

E

Eating disorders, 136
Edema, 133
Educational factors
  no physical activity and, 177, 178, 196
  regular sustained physical activity, 183
  regular, vigorous physical activity, 183, 187
  strengthening activity, 191
  stretching activity, 191

Eisenhower, President Dwight D., 17, 18
Elderly persons
behavioral intervention programs for, 233
cardiovascular response to exercise, 75–76
cold stress in, 74
community-based programs for, 245–246
falling, 7, 132
health-related quality of life, 142
maximal oxygen uptake in, 32, 70
osteoporosis, 130–133
physical activity interventions for, 233
physical assessment procedures for, 30
psychomotor performance of, 35
resistance training and, 44
Electrocardiographic changes, 111
Emotional functioning, 141
Emphysema, 140
End-diastolic volume. *See under* Blood pressure
Endocrine factors, 66, 67, 70
Endocrine glands, 5, 7
hormonal responses to exercise, 66
See related Hormones
Endometrial cancer, 7, 114, 120, 149
Endorphins, 141
Endurance training, 4, 18, 19, 21, 61, 63, 65
capillary increase by, 69
health benefits of, 7, 43
health-related quality of life, 142
insulin sensitivity and, 127
ischemia and, 112
lactate threshold and, 67
metabolic adaptations, 69–70
muscle fibers in, 67
obesity and, 135
osteoporosis and, 150
Enkephalins, 141
Environmental exposure, 19, 73–74
air pollution, 74
cold climate disorders, 74
hot and humid conditions, 63, 73, 143
Epicondylitis, 143
Equilibrium, 35, 44
Ergometer tests. *See under* Exercise tests
Erythropoietin, 68, 74
Estradiol-progesterone, 68
Estrogen, 130, 131
Estrogen replacement therapy, 132
Exercise, 20, 21, 140. *See also* Physical activity; Physical activity, specific; Physical fitness; Physical fitness programs
Exercise physiology
research, 18–20
responses to, 61
textbooks on, 61
Exercise tests
accelerometers, 32
bicycle ergometry, 62–63, 64, 66, 74–75
maximal, 86, 87, 90
motion sensors, 31–32
for muscle fitness, 34–35
pedometers, 31
stabilometers, 32
submaximal, 86, 87, 89
treadmill, 32, 34, 122
Exercise training
American College of Sports Medicine recommendations, 22
benefits, 7
bone adaptations, 67, 69
cardiovascular diseases and, 45
definition of, 20
diabetes mellitus and, 128
frequency of, 61
interval vs. continuous, 19
and lipoprotein, HDL, 43
muscle, skeletal, adaptations, 67, 69
triglycerides and, 111
ventricular fibrillation, 112

F

Falls, 143, 150
Fatty acids, 111
Fibrinogen, 43
Fibrinolysis, 43, 112
Fitness Through Churches Project, 245
Foot injuries, 128, 143
Fractures, 130–132, 143

G

Gastrointestinal system
  problems, 130
  transit time, 122
Genetic factors
  diabetes mellitus, 126–127
  maximal oxygen uptake and, 66, 70
  training and, 65
Glossary, 21
Glucagon, 69
Glucose intolerance, 72, 123
Glucose tolerance, 127
Glucose-6-phosphate, 132
Glycogen, muscle storage of, 69
Glycolysis
  energy system for, 65–67
  muscle fiber capacity, 66
Go For Health (GFH), 239
Governor's Councils on Physical Fitness and Sports, 245
Guidelines for Adolescent Preventive Services (GAPS), 28
Guidelines for Exercise Testing and Prescription (ACSM), 23, 28
Guidelines for School and Community Health Programs to Promote Physical Activity Among Youths, 237, 244

H

Handicapped. *See* Physically handicapped
Harvard University, 16
  alumni study, 86
  Fatigue Laboratory, 19
Health, 16–18, 22, 141
  American College of Sports Medicine 1990 recommendations, 22–23
  exercise physiology research and, 18–20
  physical activity recommendations, 28–30
  World Health Organization definition of, 141
Health and Human Services (HHS),
  Department of, 245
  Office of Public Health and Science, 3
Health and Religion Project (HARP), 245
Health-related fitness, 20, 22
Health-related quality of life, 141–142, 150
Healthy People 2000, 23
  cardiorespiratory fitness, 244
  daily, moderate physical activity, 181, 200
  exercise intensity in, 33
  leisure-time activity target, 177
  muscle strength, endurance, 187, 189, 192, 199, 200
  objectives, 5, 175, 237, 245
  physical education, 205
  regular, vigorous activity, 182
  worksite programs, 229–231
Heart, adaptation to exercise, 71
Heart defects, congenital, 45
Heart disease, 142
Heart failure, congestive, 45, 103
Heart rate (HR), 31–32, 62, 72, 73
  coronary circulation and, 66
  maximal, 21
  mean daily, 31
  oxygen consumption and, 63
  resting, 19, 31
  training response and, 71
Heat stress disorders, 74
Heat exhaustion, 74

Heat stroke, 74
Hematocrit, 43
Hematuria, 143
Hemoconcentration, 74
Hemoglobinuria, 143
High blood pressure *See* Hypertension
Hill's causality criteria, 144–145
Hip fracture, 130, 132
Hispanics
  bicycling activities by, 203
  child behavioral intervention program, 232
  diabetes mellitus and occupational physical activity, 126
  high school physical education enrollment and, 205
  no regular activity by, 177, 195
  physical activity trends in, 8
  regular, sustained activity by, 183
  regular, vigorous activity by, 183, 187, 183, 196–197
  sports team participation by, 200
  strengthening activities by, 191, 193, 198,
  stretching activities by, 191, 194, 201
  walking activities by, 203
Home care programs, 46
Hyaline cartilage, 130
Hydrogen ions (H+) concentration, 64
Hygiene, 11–18
Hyperglycemia, 127
Hyperinsulinism, 72
Hyperplasia, 69
Hypertension, 4–5, 7, 23, 47, 43, 63, 66, 71, 126–127, 133
  obesity and, 133
  physical activity and, 103, 110, 144–145, 149
  population studies of, 108–109
Hyperthermia, 143
Hypertrophy, 69, 71, 76, 103
Hypoglycemia, 127–128, 143
Hypotension, 63, 74

I

Immobilization, 71–72, 130
Immune system, responses to exercise, 7, 67
Immunoglobins, 67
Immunosuppression, 143
Inactivity. *See* Physical inactivity
Indian Health Service, 232
Industrywide Network for Social, Urban, and Rural Efforts (INSURE) project, 227
Infection control and exercise, 67
Injuries, 5
  exercise-related, 8, 44, 69, 150, 248
  joint, 129
  musculoskeletal, 142–143
  sports-related, 7
Insulin, 44, 67, 68, 72, 125–129
Intermodal Surface Transportation Efficiency Act, 247
International Consensus Conference
  of Physical Activity Guidelines for Adolescents (1993), 28
  on Physical Activity, Physical Fitness, and Health (1988), 22
Interpersonal relationships, 46
  behavioral sciences theories, 213,
  social support role in activity, 214, 243
Intervention studies, adolescent, 8, 236
  accessibility, 236
  church programs for, 245
  determinants, modifiable, 243
  factors influencing, 243
  health care settings, 242
  outdoor activities, 243
  parental involvement, 243
  school-community programs, 242, 245
  school programs, 236–243, 236–243
  self-efficacy, 242, 248
  societal barriers, 246–247
  societal resources, 247

Intervention studies, adult, 8, 217
communications, 229–230
community approaches to, 227–229, 234, 227–229, 234, 245
church programs for, 245
environmental approach to, 244–245
factors influencing, 215–217
health care settings, 226–227
individual approaches, 217, 226, 234
mental disorders, 136
policy approaches, 244–245
promotion of, 226, 234
societal barriers, 246–247
societal resources, 247
worksites, 229–231, 229–231, 236
Intervention studies, children, 8, 236
accessibility, 243
church programs for, 246
determinants, modifiable, 243
factors influencing, 243
health care settings, 242
outdoor activities, 243
parental involvement, 243
school-community programs, 242, 245
school programs, 236–243
self-efficacy, 242, 248
societal barriers, 246–247
societal resources, 247
Intervertebral disc displacement, 142
Intra-abdominal fat distribution, 128–129
Ischemia, 110, 111–112

J

Johnson, President Lyndon B., 18
Johnson & Johnson Live for Life program, 230

K

Kennedy, President John F., 18
Kilocalorie (kcal), 21, 29, 140, 143, 146–147, 148
Kilojoule (kjoule), 21, 29
Know Your Body (KYB) program, 238

L

Lacerations, 143
Lactate threshold (LT), 66, 67, 69, 70
Lactates, 66, 67, 70, 74
Leukocytes, 128
Ligaments, 69
Lipoproteins, 110
HDL, 43, 91, 102
LDL, 91, 102
lipase activity, 111
profile, 111, 145
Low back pain, 233–234
Lung diseases, obstructive, 233–234
Lymphocytes, 67

M

Magnetic resonance imaging, 35
Marfan syndrome, 45
Mass spectrometer, 32
Maximal oxygen uptake ($\dot{V}O_2$ max), 21, 23, 32–34, 62–63, 66, 67, 69–70, 72, 75–77, 110
Media. *See* Communications interventions
Men and physical activity
no regular activity, 4, 8, 177, 178, 188, 189
regular, sustained activity, 183, 188, 205
regular vigorous activity, 185, 187, 188, 205
selected physical activities, 188
strengthening activities, 191
trends, 8
Menopause, 130, 131
Mental disorders, 135
Mental health, 4, 8, 135–141, 150

Mental retardation, 73
Metabolic equivalent (MET), 21, 29, 32, 33, 66, 148, 204
Metabolic rate, 66
Metabolism
  aerobic, 20, 21
  bed rest and disturbances of, 72
  benefits, 7
  carbohydrate, 128
  energy expenditure, 134
  glucose, 65, 128
  fat, 68
  muscle, skeletal, 65–67, 71–72
  protein, 66
  response to exercise, 18–19, 121, 64, 69–70
Metropolitan Life Insurance Company weight tables, 133
Minnesota Heart Health Program (MHHP), 227–228, 232
Minnesota Leisure-Time Physical Activity Questionnaire, 31, 36
Missouri "Bootheel" behavioral sciences study, 229, 232
Mitochondria, 66
Monocyte-macrophage system, 67
Mortality, 85–87, 149
  all-cause, 133
  diabetes mellitus, 125
  heart disease, 87
  lowering, 7
  premature, 4, 16
  traffic fatalities, 246
Multiple sclerosis, 73, 233–234
Muscle contractions, 34
Muscle fatigue, 65
Muscle fibers, 21
  fast- and slow-twitch, 65, 67, 69, 73
Muscles, skeletal, 5
  adaptations to exercise, 7, 44, 65, 67, 69–70
  atrophy of, 69, 72–73
  capillaries in trained, 71
  energy metabolism of, 65–67
  fibers in, 65, 67, 69, 73
  immobilization and, 72–73
  insulin and, 125, 130
  metabolic adaptations of, 69–70
  multiple sclerosis and, 233–234
  soreness in, 69
  structural damage to, 69
Muscular endurance (fitness), 21, 34–35
Muscular strength, 34, 44
Myocardial contraction, 65, 72
Myocardial infarction, 5, 44, 45, 112, 143
Myocardium, 111
Myosin ATP, 65

N

National Association for Sport and Physical Education, National Physical Education Standards, 244
National Coalition for Promoting Physical Activity, 244
National Institutes of Health (NIH), 5
  Consensus Development Conference Statement, Physical Activity and Cardiovascular Health, 5, 23, 28, 48, 148, 245
  intervention campaigns, 245
  National Physical Education Standards, 244
Native Americans, 12
  adult physical activity interventions, 232
  behavioral intervention program,232
Neoplasms. *See* Cancer
Neural factors, 67
Neuromas, 143
Nitric oxide, 132
Nitrogen, balance, 72

O

Obesity, 7, 43, 133–135, 150, 248
abdominal, 35, 128–129
adult physical activity interventions, 232–234
behavioral intervention programs, 232–233
in adolescents, 102
in children, 102
childhood intervention, 244
trends in, 47, 46
Occupational medicine, 15
Occupational physical activity, 113, 116, 175, 189. *See also* Worksite physical fitness programs

Olympic Games, 12, 15
Osteoarthritis, 7, 129–130, 133, 149–150
Osteoporosis, 7, 23, 43, 69, 130–133, 150. *See related* Bone density
Otitis externa, 143
Ovarian cancer, 7, 114, 116–118, 149
Overtraining, 21, 140
Overweight, 133. *See also* Obesity
Oxidative capacity, of muscle fibers, 65, 67
Oxidative energy system, 65, 66
Oxygen
arterial-mixed venous, 62, 63, 70
ATP production within mitochondria, 66
body's use of, 61
delivery, 74
extraction, blood flow, 63
myocardial demand/use, 63, 64
Oxygen consumption ($\dot{V}O_2$), 18, 31, 32, 34, 66, 70, 74, 110
Oxygen uptake. *See* Maximal oxygen uptake

P

Pain threshold, 130
Paleolithic rhythm, 11
Pawtucket Heart Health Program (PHHP), 229
Pediatricians, physical activity counseling by, 244
Peripheral vascular disease, 45
Personality disorders, 136
Phosphocreatine (PCr), 65
Physical activity, 21
of adolescents. *See* Adolescents and physical activity
of adults. *See* Adults and physical activity
adverse effects of, 142–144
approaches to, 46–47
of children. *See* Children and physical activity
definition of, 20
dosage, 146–148
duration of, 44, 147, 148
evolution of recommendations, 22–28
frequency of, 44
intensity of, 29–33, 35–36, 44
measures of, 211–215
no regular, 15–16, 23, 50, 46–48, 177–189, 188, 195, 248
regular, intermittent, 11, 148
regular, sustained, 4, 6, 23, 37, 49, 43, 110, 146–147, 182–183, 244
regular, vigorous, 4, 6, 11, 23, 37, 50, 110, 127–128, 146–147, 182–187, 188, 244
research considerations, 47, 150
social environmental approaches to, 244–245
surveys of, 175, 177
Physical activity, specific
aerobics, 200, 205
baseball, 129, 143, 200, 205
basketball, 143, 200, 205
bicycling, 4, 143, 144, 187, 200,203
boxing, 143
carpentry, 140
dancing, 14, 143, 144, 148, 187, 200, 205,
football, 129, 143, 200, 205
Frisbee, 197–198, 205
gardening, 8, 140, 144, 147, 187
golfing, 140
hockey, 140, 148
horseback riding, 14

housecleaning, 147, 148, 200, 205
jogging, 140, 142, 144, 148, 187, 198, 205,
kickball, 12
lacrosse, 12
racquetball, 143, 197–198, 205
running, 4, 12, 65, 66, 70, 129, 140, 142, 143, 144, 148, 187, 198, 205
skating, 197–198, 205
skateboarding, 197–198, 205
skiing, 197–198, 205
soccer, 129, 143, 197–198, 205
softball, 144, 197–198, 205,
stair climbing, 127, 147, 187
squash, 197–198, 205
swimming, 140, 143, 144, 197–198, 205,
tennis, 14, 140, 197–198, 205
volleyball, 4, 148,
walking, 4, 8, 14, 33, 127, 140, 144, 148, 187, 197–198, 203, 204, 205, 233
weight lifting, 129, 143
yard work, 4, 8, 147, 187, 202, 200,205, 205
Physical Activity for Risk Reduction (PARR) project, 232
Physical education, 8, 16–18
enrollment in, 4
in high school, 205, 205
school program interventions, 243, 246–249
Physical examination, 6, 45, 47
Physical fitness, 16–18, 21
assessment procedures, direct monitoring, 31–36
assessment procedures, self-reporting, 29–31
definition of, 20
level of, 61
maintenance of, 71–72
maximal oxygen uptake and. *See* Maximal oxygen uptake
measurement of, 33–35
physical activity relationship, 43
worksite programs, 48, 46, 48
See also Cardiorespiratory endurance (fitness)
Physical fitness programs
detraining and, 61, 72
endurance training, 61, 63, 65, 67, 69–70
exercise training, 61, 67, 69
resistance training, 61, 65, 69, 70
Physical functioning, 141
Physical inactivity, 5, 6, 72, 73, 145–146, 148
bone loss and muscle atrophy in, 69
diseases of, 15–16
exercise programs and, 37
health burden of lifestyle, 42–43
mortality and, 86
percentage of, 4
physical activity recommendations for, 29
physiological alterations after endurance training, 70
societal inducements for, 254–247
working toward recommendations, 44
Physically handicapped, 73
behavioral intervention programs, 233–234
childhood interventions, 233, 244
no regular activity and, 189
regular, moderate activity and, 189
regular, vigorous activity and, 189
Physician-based Assessment and Counseling for Exercise (PACE), 227
Plasma insulin concentration, 128
Plasma lipid/lipoprotein, 110, 111
Plasma volume, 71, 72, 74
Platelet function, 43
Population attributable risk (PAR), 145–146
Postmenopause, 131
Postpoliomyelitis syndrome, 73
Power, 21, 85
President's Citizens Advisory Committee on the Fitness of American Youth, 18
President's Conference on Fitness of American Youth, 18
President's Council on Physical Fitness, 18, 23
President's Council on Physical Fitness and Sports (PCPFS), 3, 5, 18, 227, 245

President's Conference on Youth Fitness, 18
Presidential Physical Fitness Award, 18
Preventive medicine, 11–18
Prostaglandin, 68, 113–117, 124
Prostate cancer, 7, 121–122, 124–125
Proteoglycan synthesis, 130
Psychological assessments, 136–137
Psychomotor performance, 17, 19–20, 35, 44
Pulmonary ventilation, 64, 71
Pulse rate, 16

Q

Quadriplegia, 75
Quality of life, 8, 141–142, 150
Quetelet's index, 133

R

Rating of perceived exertion (RPE) scale, 33
Reaction time, 21
Receptors, sensory, 141
Rectal cancer, 7, 113, 116, 122, 149
Relative perceived exertion (RPE), 21
Renin-angiotensin system, 68
Resistance training, 4, 19, 21, 29, 37, 44, 61, 65, 69
  adolescent, 196, 198–200
  elderly persons and, 7, 132–133
  glucose-insulin dynamics, 128
  muscle, skeletal effects of, 69
  obesity and, 135
  osteoporosis and, 150
  risk factors, 67
  strengthening activities for, 187, 189, 191–192, 193, 199–200
  sex factors, 70
Respiration rate (RR), 70, 71
Respiratory system, 5
  adaptation to exercise, 71
  physiological alterations after endurance training, 70
  resistance exercise and, 65
  response to exercise, 61–62, 64
Retraining, 21
Rhabdomyolysis, 143
Rhode Island
  Department of Parks and Recreation, 229
  Health and Religion Project, 245
  Pawtucket Heart Health Program, 229
Roosevelt, President Franklin D., 17

S

Schizophrenia, 136
School program interventions, 6, 236–243, 246–248
Sedentary persons. *See* Physical inactivity
Self concept, 130, 141, 142
Self-help, 13–15, 29–31
Senior citizens. *See* Elderly persons
Sex factors
  in exercise training, 70, 76–77
  hormonal responses to exercise, 67
  hormone-dependent cancers in men, 121–125
  hormone-dependent cancers in women, 114–121
  physical inactivity and, 177–178
  specific physical activities and, 187–189
  strengthening, stretching activities by, 191
  weight gain prevention and, 232
Shoulder dislocation, 142
Skeletal muscles. *See* Muscles, skeletal
Skin, receiving cardiac output at rest vs. exercise, 63
Skinfold measures, 133, 134
Sleep disorders, 136

Social environment
barriers, 246–247
behavioral influences, 215,
resources, 247
Social functioning, 141
Social sciences. *See* Behavioral sciences, adult; children and adolescents
Socioeconomic factors
bicycling activities and, 203
physical inactivity and, 177, 178, 196
resistance training and, 193
stretching activities and, 194
walking activities and, 200
Speed, 21, 203–204
Splanchnic circulation, 63
Sports, Play, and Active Recreation for Kids (SPARK) study, 239
Stanford Adolescent Heart Health Program, 238
Stanford Five-City Project (SFCP), 229
Strength, 21, 22
Strength testing, 16, 34
Strength training. *See* Resistance training
Stretching activities, 187, 191–192
Stroke volume (SV), 62, 71–73
Substance use disorders, 136
Suicide, 135, 140
Systolic blood pressure (SBP), 63, 64, 70

T

T-lymphocytes. *See* Lymphocytes
Tai chi chuan, 12, 113
Taoism, 12
Tecumseh questionnaire, 31
Temperature. *See* Body temperature
Tendinitis, 143
Tendons, 69
Testicular cancer, 7, 124, 147
Thrombosis, 102, 110, 112
Thymus gland, 67
Tidal volume (TV), 70, 71
Training heart rate (THR), 21
Transient constriction, 111
Triglycerides, 111
Trust for Public Land, The, 245

U

Ulnar nerve palsies, 143
United Kingdom Testicular Cancer Study Group, 122
United States regions
East, 12
Midwest, 229
North central, 177, 187
Northeast, 177, 187
South, 177, 189, 187
Southwest, 12
West, 177, 182, 187, 189, 187
U.S. Preventive Services Task Force, 28
Urogenital system, 143
Uterine cancer, 114, 120–121

V

Vasoconstriction, 76
Ventilatory volume ($\dot{V}_E$), 70, 74
Ventricular dysfunction, left, 103
Ventricular end-diastolic volume, 71, 72
Ventricular fibrillation, 112
Vertebral fractures, 130
Virginia
Smyth County program, 245

W

Water-electrolyte imbalance, 143
Weather factors
seasonality, 184, 196, 196, 200, 204
summer months, 184, 187
walking, bicycling activities and, 204
Weight Gain Prevention Program, 232

West Virginia
elderly mall walking campaign in, 233
Whites
behavior intervention program for, 232
bicycling activities by, 203
high school physical education enrollment, 205
no regular activity, 177, 188, 195
physical activity trends in, 8
regular, sustained activity, 183
regular, vigorous activity, 183, 187, 188, 196–197
sports team participation, 200
strengthening activities by, 189, 191, 193, 199
stretching activities by, 191, 194, 201
walking activities by, 203
Wise Weighs program, 232
Women and physical activity
no regular activity, 177, 178, 188, 205
regular, sustained activity, 183, 188, 205
regular, vigorous activity, 185, 187, 188, 189
selected physical activities, 188
strengthening activities, 191
trends, 8
Work rate, 64, 71
energy metabolism and, 65
energy systems and, 65–66
increasing, 62–63, 64
Worksite physical fitness programs, 48, 46, 48, 184
World Health Organization, definition of health, 141
World War I, 16
World War II, 17, 18, 19

Y

YMCA, 23
Yoga, 12

Z

Zuni Diabetes Project, 232