

# Postharvest Pathology

Peter L. Sholberg<sup>1</sup> and William S. Conway<sup>2</sup>

<sup>1</sup>Agriculture and Agri-Food Canada, Pacific Agri-Food Research Centre  
Summerland, BC, Canada

<sup>2</sup>Produce Quality and Safety Laboratory, USDA/ARS  
Henry A. Wallace Beltsville Agricultural Research Center Beltsville, MD

**Introduction:** Losses caused by postharvest diseases are greater than generally realized because the value of fresh fruits and vegetables increases several-fold while passing from the field to the consumer (Eckert and Sommer, 1967). Postharvest losses are estimated to range from 10 to 30% per year despite the use of modern storage facilities and techniques (Harvey, 1978).

Postharvest diseases affect a wide variety of crops particularly in developing countries which lack sophisticated postharvest storage facilities (Jeffries and Jeger, 1990). Infection by fungi and bacteria may occur during the growing season, at harvest time, during handling, storage, transport and marketing, or even after purchase by the consumer (Dennis, 1983). The reduction of losses in perishable food crops because of postharvest diseases has become a major objective of international organizations (Kelman, 1989). The reality is that there is a portending food crisis that will require the concerted efforts of all who are involved in food production to double their efforts. In fact, to adequately feed the world's expected 10 billion people within the next 40 to 50 years, food production efficiency and distribution will need to be improved immensely (Campbell, 1998).

Specific causes of postharvest losses of fruits and vegetables may be classed as parasitic, nonparasitic, or physical (Cappellini and Ceponis, 1984). This chapter deals with the parasitic causes that are of microbiological origin that begin as latent infections before harvest or occur at or after harvest during storage. Fungi are more commonly found attacking fruit and bacteria are more common as postharvest pathogens of vegetables. This chapter will provide a general overview of the subject touching on noteworthy research where it can be used to illustrate postharvest pathology. The reader is encouraged to consult the references for specific information on the topics that are covered.

**Preharvest Factors that Influence Postharvest Pathology:** Postharvest losses vary each year. Prevailing weather while the crop is growing and at harvest contribute greatly to the possibility of decay. Certain cultivars are more prone to decay than are others to specific pathogens. In a recent study, it was found that resistance of major apple cultivars to the fungi that cause blue mold, gray mold, bull's-eye rot, and Mucor rot was dependent on cultivar (Spotts et al., 1999). Condition of the crop, as determined by fertilizer and soil factors, are very important in susceptibility of the crop to disease. Maturity of the crop at harvest, handling and type of storage have a great deal of influence on how long the crop can be stored without decay. Examples are given below that demonstrate how these preharvest factors lead to disease in specific crops.

*Weather:* Weather affects many factors related to plant diseases, from the amount of inoculum that overwinters successfully to the amount of pesticide residue that remains on the crop at harvest (Conway, 1984). Abundant inoculum and favorable conditions for infection during the season often result in heavy infection by the time the produce is harvested. For example, conidia of the fungus that causes bull's-eye rot are rain dispersed from cankers and infected bark to fruit especially if rainfall is prolonged near harvest time, causing rotten fruit in cold storage several months later (Spotts, 1990).

Pinpoint or storage scab of apple caused by the same fungus that causes apple scab, and gray mold caused by the fungus *Botrytis cinerea*, are also very much influenced by the weather. Storage scab only occurs in years with unusually wet Summers and early Falls, when the fruit remain wet for a day or more.

These late season infections may not become visible until the apples are in storage (Pierson et al., 1971). Flowers and fruit are infected by *Botrytis cinerea* most effectively when it is wet. For example, in grapes infection occurs at 15 to 20 °C (59 to 68 °F) in the presence of free water after approximately 15 h (Bulit and Dubos, 1988). In wet seasons strawberries and raspberry crops may be harvested in apparently sound condition, only to decay during transit and marketing (Snowdon, 1990).

Postharvest decay involves further development of pre-harvest infections together with new infections arising from germination of spores on the fruit surface. From these examples it is apparent that decay often has a weather component making thorough weather records an important source of information for predicting possible decay in storage.

*Physiological Condition:* Condition of produce at harvest determines how long the crop can be safely stored. For example, apples are picked slightly immature to ensure that they can be stored safely for several months. The onset of ripening and senescence in various fruit and vegetables renders them more susceptible to infection by pathogens (Kader, 1985). On the other hand, fruit and vegetables can be made less prone to decay by management of crop nutrition. For example, calcium has been more closely related to disease resistance than any other cation associated with the cell wall (Sams, 1994).

In a study on the effect of increased flesh calcium content of apples on storage decay fruit were treated with solutions of CaCl<sub>2</sub> by dipping, vacuum, or pressure infiltration. Both vacuum and pressure infiltration increased calcium content of the fruit sufficiently to significantly reduce decay (Conway, 1982). Increased calcium contents in potatoes and peaches have also been documented with reduced postharvest decay (Conway, 1989). In general, produce containing adequate levels of calcium do not develop physiological disorders and can be stored longer before they breakdown or decay. Conversely, high nitrogen content in fruit predisposes them to decay (Conway, 1984). In pears, it has been found that management of trees for low nitrogen and high calcium in the fruit reduced severity of postharvest fungal decay (Sugar et al., 1992). Apple cultivars can be selected for resistance to certain postharvest diseases (Spotts et al., 1999). For example, 'Royal Gala' is extremely resistant to wound pathogens, 'Granny Smith' to skin punctures, and 'Braeburn' to infiltration of fungal spores into the core.

*Fungicide Sprays:* Certain pre-harvest sprays are known to reduce decay in storage. Several studies have been done on the effectiveness of pre-harvest ziram fungicide application on pome fruit and show an average reduction in decay of about 25 to 50% with a single spray (Sugar and Spotts, 1995). Iprodione has been used for several years as a pre-harvest spray 1 day before harvest to prevent infection of stone fruit by *Monilinia* spp. In combination with wax and/or oil its decay control spectrum is increased and it will also control postharvest fungi such as *Rhizopus*, and *Alternaria* (Ogawa, et al. 1992). Several new fungicides that are being developed, or have recently been registered promise to protect produce from a number of diseases after harvest. For example, cyprodinil prevented gray mold infection in apple 3 mo after it was applied (Sholberg and Bedford, 1999). The new class of strobilurin fungicides promise to provide postharvest control of several diseases in fruit and vegetables. They are especially effective against fruit scab on apples and should reduce the presence of pin point scab in storage.

### **Postharvest Factors that Influence Decay:**

*Packing Sanitation:* It is important to maintain sanitary conditions in all areas where produce is packed. Organic matter (culls, extraneous plant parts, soil) can act as substrates for decay-causing pathogens. For example, in apple and pear packinghouses, the flumes and dump tank accumulate spores (Blanpied and Purnasiri, 1968) and may act as sources of contamination if steps are not taken to destroy or remove them.

Chlorine readily kills microorganisms suspended in dump tanks and flumes if the amount of available chlorine is adequate. A level of 50 to 100 ppm of active chlorine provides excellent fungicidal activity (Spotts and Peters, 1980). Chlorine measured as hypochlorous acid can be obtained by adding chlorine gas, sodium hypochlorite, or dry calcium hypochlorite. Although chlorine effectively kills spores in water it

does not protect wounded tissue against subsequent infection from spores lodged in wounds. Organic matter in the water inactivates chlorine, and levels of chlorine must be constantly monitored. The use of a sand filter in association with chlorination improves its efficiency probably because it removes organic matter (Sholberg and Owen, 1990). Chlorine is sensitive to pH (Dychdala 1983); hypochlorite solutions with higher pH values (7.5 to 8.5) are more stable but less fungicidal, whereas at lower pH values (5.5 to 6.5) the solutions are less stable but more fungicidal.

Recently, chlorine dioxide has replaced chlorine in some sanitizing processes, because several disadvantages limit the use of chlorine, including its unpleasant odor. Chlorine dioxide is not corrosive and is effective over a wide pH range (Spotts and Peters, 1980). Recently, in precisely controlled tests in water or as a foam, chlorine dioxide was found to be effective against common postharvest decay fungi on fruit packinghouse surfaces (Roberts and Reymond, 1994). Peracetic acid is another material that could be used (Mari et al., 1999). It has greater stability and faster biocidal properties than chlorine dioxide, but is more corrosive.

The search goes on for effective and economical sanitizing agents. New and old products alike, are continually being evaluated under present day packing operations. Interest in ozone has been rekindled with development of more efficient ozone generators. Acetic acid in the form of a gas is being evaluated for possible use as a sanitizing agent on several crops (Sholberg, 1998). It was as effective as SO<sub>2</sub> in preventing Gray Mold decay in table grapes stored for 2 mo (Sholberg et al., 1996).

### **Postharvest Treatments:**

Products used for postharvest decay control should only be used after the following critical points are considered (Ogawa and Manji, 1984):

- Type of pathogen involved in the decay.
- Location of the pathogen in the produce.
- Best time for application of the treatment.
- Maturity of the host.

Environment during storage, transportation and marketing of produce. Specific materials are selected based on these conditions and fall into either chemical or biological categories listed below.

*Fungicide treatments:* Several fungicides are presently used as postharvest treatments for control of a wide spectrum of decay-causing microorganisms. However, when compared to preharvest pest control products the number is very small. Many former products that were used after harvest are no longer permitted because of concerns with residues and possible toxic effects, the most notable being products that contained benomyl. Other products have been lost as effective controls due to development of resistance by the target pathogen. For example, intensive and continuous use of fungicides for control of blue and green mold on citrus has led to resistance by the causal pathogens of these diseases (Eckert, 1988). Resistance has been reported in many other crops to several different fungicides with different modes of action (Delp, 1988). Resistance development continues to be an important problem. It has led to the "Fungicide Resistance Action Committee" (FRAC, 1998), a cooperative effort between the various producers of fungicides to delay resistance by recommending specific management guidelines.

Examples of postharvest chemical treatments that are presently used are thiabendazole, dichloran, and imazalil. However, resistance to thiabendazole and imazalil is widespread (Holmes and Eckert, 1999) and their use as effective materials is declining. Preservatives or antimicrobial food additives are not generally thought of as postharvest treatments but they do control decay, and in some cases are the only means of control. These products include sodium benzoate, the parabens, sorbic acid, propionic acid, SO<sub>2</sub>, acetic acid, nitrites and nitrates, and antibiotics such as nisin (Chichester and Tanner, 1972). For example, in California gray mold of stored table grapes is prevented by fumigation with SO<sub>2</sub> (Luvisi et al., 1992). The demand for new postharvest fungicide treatments is strong, especially since the loss of iprodione in 1996.

Fludioxinil was granted an emergency registration in 1998 to curb potential losses in nectarines,

peaches, and plums that would have resulted (Forster and Adaskaveg, 1999). Not all postharvest pathogens are presently controlled by materials that are available. For example, *Mucor piriformis*, a major postharvest pathogen of apples and winter pears in the Pacific Northwest is not controlled by any registered fungicide (Spotts and Dobson, 1989). There is a dire need for new fungicide treatments that could in part be alleviated by using biological control agents (Wisniewski and Wilson, 1992; Utkhede and Sholberg, 1993).

*Biological Control of Postharvest Pathogens:* Postharvest biological control is a relatively new approach and offers several advantages over conventional biological control (Wilson and Pusey, 1985; Pusey, 1996):

Exact environmental conditions can be established and maintained.

The biocontrol agent can be targeted much more efficiently.

Expensive control procedures are cost-effective on harvested food.

Several biological control agents have been developed in recent years, and a few have actually been registered for use on fruit crops. The first biological control agent developed for postharvest use was a strain of *Bacillus subtilis* (Pusey and Wilson, 1984). It controlled peach brown rot, but when a commercial formulation of the bacterium was made, adequate disease control was not obtained (Pusey, 1989). More recently, a strain of *Pseudomonas syringae* van Hall was found that controlled both Blue and Gray Mold of pome fruit (Janisiewicz and Marchi, 1992). It was subsequently registered, and is now sold commercially for postharvest disease control (Janisiewicz and Jeffers, 1997).

Other bacterial microorganisms are being developed for postharvest disease control. For example, strains of *Bacillus pumilus* and *Pseudomonas fluorescens* have been identified that exhibit successful control of *B. cinerea* in field trials of strawberry (Swalding and Jeffries, 1998). Yeasts such as *Pichia guilliermondii* (Wisniewski et al., 1991) and *Cryptococcus laurentii*, a yeast that occurs naturally on apple leaves, buds, and fruit (Roberts, 1990) were the first to be applied for control of postharvest decay on fruit.

The yeast, *Candida oleophilia* has been registered for control of postharvest decay on fruit crops. The yeasts, *Cryptococcus infirmo-minutus* and *Candida sake* successfully control brown rot and blue mold on sweet cherry (Spotts et al., 1998), and three diseases of apple (Vinas et al., 1998), respectively, and may be developed commercially.

Although there is no doubt that biocontrols are effective, they do not always give consistent results. This could be because biocontrol efficacy is so directly affected by the amount of pathogen inoculum present (Roberts, 1994). Compatibility with chemicals used during handling is also important. Indications are that biological control agents must be combined with other disease control strategies if they are to provide acceptable control.

*Irradiation for Postharvest Decay Control:* Although ultraviolet light has a lethal effect on bacteria and fungi that are exposed to the direct rays, there is no evidence that it reduces decay of packaged fruits and vegetables (Hardenburg et al., 1986). More recently, low doses of ultraviolet light irradiation (254 nm UV-C) reduced postharvest brown rot of peaches (Stevens et al., 1998). In this case, the low dose ultraviolet light treatments had two effects on brown rot development; reduction in the inoculum of the pathogen and induced resistance in the host. However, it has not become a practical postharvest treatment as yet and requires more research.

Gamma radiation has been studied for controlling decay, disinfestation, and extending the storage and shelf-life of fresh fruits and vegetables. Dosages of 1.5 to 2 kilogray (kGy), and some cases 3.0 kGy (300 krad), have been effective in controlling decay in several products (Hardenburg et al., 1986). A dose of 250 Gy has an adverse effect on grapefruits increasing skin pitting, scald, and decay. Low doses of 150 for fruit flies and 250 gray (Gy) for codling moth are acceptable quarantine procedures (Meheriuk and Gaunce, 1994). Commercial application of gamma radiation is limited due to the cost and size of equipment needed for the treatment and to uncertainty about the acceptability of irradiated foods to the consumer (Hardenburg et al., 1986). Gamma irradiation may be used more in the future once methyl bromide is no longer available to control insect infestation in stored products. All uses of methyl bromide are being phased out to avoid

any further damage to the protective layer of ozone surrounding the earth.

*Effect of Storage Environment on Postharvest Decay:* Commercial producers and handlers modify temperature, RH, and atmospheric composition during prestorage, storage, and transit to control decay (Spotts, 1984). For optimum decay control, two or more factors often are modified simultaneously.

*Temperature and RH:* Proper management of temperature is so critical to postharvest disease control that all other treatments can be considered as supplements to refrigeration (Sommer, 1989). Fruit rot fungi generally grow optimally at 20 to 25 °C (68 to 77 °F) and can be conveniently divided into those with a growth minimum of 5 to 10 °C (41 to 50 °F), or -6 to 0 °C (21.2 to 32 °F). Fungi with a minimum growth temperature below -2 °C (28.4 °F) cannot be completely stopped by refrigeration without freezing fruit. However, temperatures as low as possible are desirable because they significantly slow growth and thus reduce decay.

High temperature may be used to control postharvest decay on crops that are injured by low temperatures such as mango, papaya, pepper, and tomato (Spotts, 1984). Although hot water generally is more effective, hot air has been used to control decay in crops that are injured by hot water. Heating of pears at temperatures from 21 to 38 °C (69.8 to 100.4 °F) for 1 to 7 days reduced postharvest decay (Spotts and Chen, 1987). Decay in 'Golden Delicious' apples was reduced by exposure to 38 °C (100.4 °F) for 4 days (Sams et al., 1993) and virtually eliminated when treated after inoculation (Fallik et al., 1995; Klien et al., 1997). Heat treatment eliminates incipient infections and improves coverage by fungicides (Couey, 1989). The primary obstacle to the widespread use of heat to control postharvest fruit diseases or insect infestation is the sensitivity of many fruit to the temperatures required for effective treatment.

Both low and high RH have been related to postharvest decay control. Perforated polyethylene bags for fruit and vegetable storage create RH about 5 to 10% above that in storage rooms. Although shrivel and weight loss are reduced, decay may be increased (Spotts, 1984). Crops such as apples and pears with well-developed cuticle and epidermis, tolerate lower RH levels that help prevent storage decay. Often fungal spore germination is inhibited at low RH, and small differences in RH can have significant effects in relation to the degree of postharvest decay (Spotts and Peters, 1981).

*Modified or Controlled Atmospheres:* Alterations in O<sub>2</sub> and CO<sub>2</sub> concentrations are sometimes provided around fruit and vegetables (Spotts, 1984). With close control of these gases, the synthetic atmosphere is commonly called a controlled atmosphere; the term modified atmosphere is used when there is little possibility of adjusting gas composition during storage or transportation (Sommer, 1989). Because the pathogen respire as does produce, lowering the O<sub>2</sub> or raising the CO<sub>2</sub> above 5% can suppress pathogenic growth in the host. In crops such as stone fruits, a direct suppression occurs when fungal respiration and growth are reduced by the high CO<sub>2</sub> of the modified atmosphere. For example, CO<sub>2</sub> added to air has been widely utilized in the transport of 'Bing' cherries, primarily to suppress Gray Mold and Brown Rot. Low O<sub>2</sub> does not appreciably suppresses fungal growth until the concentration is below 2%. Important growth reductions result if the O<sub>2</sub> is lowered to 1% or lower although there is a danger that the crop will start respiring anaerobically and develop off-flavor. Other technologies that have been tested for lowering postharvest decay with limited success have been storage and transport under low O<sub>2</sub> and the use of carbon monoxide (Spotts, 1984; Sommer, 1989).

**Postharvest Diseases of Fruits:** Fruit crops are attacked by a wide range of microorganisms in the postharvest phase (Snowdon, 1990; Ogawa and English, 1991). Actual disease only occurs when the attacking pathogen starts to actively grow in the host. Diseases are loosely classified according to their signs and symptoms. Signs are visible growths of the causal agents, and symptoms the discernible responses produced by the host. In many diseases there is local discoloration and disruption of tissue, with the formation of obvious lesions. Postharvest diseases are caused primarily by microscopic bacteria and fungi, with fungi the most important causal agent in fruit crops. Fungi are further subdivided into classes

and are described as lower fungi, characterized by the production of sporangia which give rise to numerous sporangiospores, or higher fungi, described as ascomycetes, deuteromycetes, and basidiomycetes. Ascomycetes are exemplified by fruiting bodies that release sexual spores when mature. Deuteromycetes, a form of ascomycete, only release asexual spores. They are more common than the sexual ascomycete stage in postharvest crops. Deuteromycetes are further subdivided into hyphomycetes and coelomycetes based on spore and structural characteristics. The anamycetes contain important soil pathogens that form survival structures known as sclerotia that allow them to survive in the absence of the host. These fungi and the rust and smut fungi are examples of basidiomycetes. Table 1 lists many important diseases of fruit crops according to host and causal agents.

**Postharvest Diseases of Vegetables:** Postharvest diseases of vegetables are caused by microscopic fungi and bacteria (Snowdon, 1992; Howard et al., 1994). Bacteria are more common as pathogens of vegetables than fruit because in general vegetables are less acidic than fruit. They are visible under the light microscope as mostly single-celled rods. Bacteria are capable of very rapid multiplication under the right conditions of pH, temperature, and nutrition. They are classified according to their size, shape, reaction to certain stains, and behavior on various growth media (Krieg and Holt, 1984). The term “vegetable” encompasses a range of plant parts, and the common definition is a culinary one, denoting consumption as a savory rather than as a dessert food (Snowdon, 1992). Many vegetables are fruits in the botanical sense, with notable examples being tomatoes, peppers, squashes, and cucumbers. Table 2 lists many of the important diseases of vegetable crops according to host and causal agent.

**New Directions for Postharvest Plant Pathology:** Postharvest plant pathology has changed its emphasis in recent years. Food safety has emerged as a key element in decay control programs. Continued failure to effectively control certain postharvest diseases and the need for more environmentally friendly crop control materials is driving a new approach to disease control. Integrated postharvest decay control is the concept that offers the most promise for the future. Society can no longer rely on one or two control strategies but must enlist the entire spectrum of strategies to reduce postharvest losses.

**Food Safety Issues:** The two most important causes of unsafe food are microbial toxins (Hsieh and Gruenwedel, 1990) and contamination of horticultural products by fecal coliforms (Gould, 1973). The microbial toxins can be subdivided into bacterial toxins and toxins produced by fungi or mycotoxins. An example of a microbial toxin that is extremely toxic are the botulinum toxins produced by the anaerobic bacterium, *Clostridium botulinum*. Interest in toxins produced by fungi was stimulated by the death of 100,000 turkey poults in England in 1960. Aflatoxins produced by fungi in the peanut meal used to feed the birds was the cause. Studies have since shown aflatoxins to be potent carcinogens that may occur in nuts and grain (Phillips, 1984; Ellis et al., 1991). Other toxins have been identified that are produced by the same fungi that cause postharvest decay. For example, patulin produced by *Penicillium* and *Aspergillus* spp. can be found in apple and pear products. Patulin is toxic to many biological systems but its role in causing animal and human disease is unclear (Hsieh and Gruenwedel, 1990). Studies on contamination of horticultural products by fecal coliforms has increased dramatically because of documented incidences of food poisoning from apple juice and seed sprouts. Definite interactions have been shown between plant pathogens and foodborne human pathogens such as *Salmonella* and *Listeria*. A study, involving more than 400 samples each of healthy and soft rotted commodities collected in retail markets, indicated that the presence of *Salmonella* on produce affected by bacterial soft rot was twice that of healthy samples (Wells and Butterfield, 1997). Controlled experiments with potato, carrot and pepper tissues inoculated with a strain of *Salmonella* confirmed that bacterial soft rot infection increased multiplication of *Salmonella* by at least three- to ten-fold compared to multiplication on uninfected tissues. Similarly, populations of *Listeria monocytogenes*, inoculated into decayed apple tissue, continually increased on fruit decayed by *Glomerella cingulata*, but did not survive after 5 days on fruit decayed by *Penicillium expansum* (Conway et al., 2000). The pH of the decayed area declined from pH 4.7 to 3.7 in the case of *P. expansum*, but in the case of *G.*

*cingulata* pH increased from pH 4.7 to 7.0. This pH modification may be responsible for affecting growth of the foodborne pathogen. Contamination of produce with human pathogens is an important issue that must be addressed while at the same time limiting decay caused by postharvest pathogens and maintaining product quality.

**Integrated Control of Postharvest Diseases:** Effective and consistent control of storage diseases is dependent upon integration of the following practices:

- Select disease resistant cultivars where possible.
- Maintain correct crop nutrition by use of leaf and soil analysis.
- Irrigate based on crop requirements and avoid overhead irrigation.
- Apply pre-harvest treatments to control insects and diseases.
- Harvest the crop at the correct maturity for storage.
- Apply postharvest treatments to disinfest and control diseases and disorders on produce.
- Maintain good sanitation in packing areas and keep dump water free of contamination.
- Store produce under conditions least conducive to growth of pathogens.

Integration of cultural methods and biological treatments with yeast biocontrols has been studied on pears (Sugar et al., 1994). It was found that early harvest, low fruit nitrogen, high fruit calcium, yeast or yeast plus fungicide treatment, and controlled atmosphere storage all reduced severity of blue mold and side rot. These results demonstrated that unrelated cultural and biological methods that influenced pear decay susceptibility can be combined into an integrated program to substantially reduce decay.

In another example of an integrated strategy 'Gala' apples were heat treated at 38 °C (100.4 °F) for 4 days, followed by calcium infiltration with 2% CaCl<sub>2</sub>, and then treated with the microbial antagonist, *Pseudomonas syringae* (Conway et al., 1999). The combined strategy was much more effective than any single strategy for the following reasons. Heat treatment reduced the pathogen population on the fruit surface but did not provide any residual protection. The residual protection was provided by calcium, and the biocontrol agent added to the control provided by the heat treatment.

As a general rule, alternatives to chemical control are often less effective than many fungicides. It is highly unlikely that any one alternative method alone will give the same level of control as fungicides. Therefore, it will generally be necessary to combine several alternative methods to develop an integrated strategy to successfully reduce postharvest decay.

#### **References:**

- Blanpied, G.D. and A. Purnasiri. 1968. *Penicillium* and *Botrytis* rot of McIntosh apples handled in water. Plant Dis. Rep. 52:865-867.
- Bulit, J. and B. Dubos. 1988. Botrytis bunch rot and blight. In: R.C. Pearson and A.C. Goheen (eds) Compendium of Grape Diseases, APS Press, St. Paul MN, pp. 13-15.
- Campbell, C.L. 1998. Food security and plant pathologists. Phytopathol. News 32:70.
- Cappellini, R.A. and M.J. Ceponis. 1984. Postharvest losses in fresh fruits and vegetables. In: H.E. Moline (ed) Postharvest Pathology of Fruits and Vegetables: Postharvest Losses in Perishable Crops, Univ. Calif., Agric. Exp. Sta., Bull. No. 1914 (Pub NE-87), pp. 24-30.
- Chichester, D.F. and F.W. Tanner. 1972. Antimicrobial food additives. In: T.E. Furia (ed) CRC Handbook of Food Additives, Vol. 1, CRC Press, Boca Raton FL, pp. 115-184.
- Conway, W.S. 1982. Effect of postharvest calcium treatment on decay of 'Delicious' apples. Plant Dis. 66:402-403.
- Conway, W.S. 1984. Preharvest factors affecting postharvest losses from disease In: H.E. Moline (ed) Postharvest Pathology of Fruits and Vegetables: Postharvest Losses in Perishable Crops, Univ. Calif., Agric. Exp. Sta., Bull. No. 1914 (Pub NE-87), pp. 11-16.
- Conway, W.S. 1989. Altering nutritional factors after harvest to enhance resistance to postharvest disease.

Phytopathology 79:1384-1387.

- Conway, W.S., W.J. Janisiewicz, J.D. Klein and C.E. Sams. 1999. Strategy for combining heat treatment, calcium infiltration, and biological control to reduce postharvest decay of 'Gala' apples. *HortScience* 34:700-704.
- Conway, W.S., B. Leverentz, R.A. Saftner, W.J. Janisiewicz, C.E. Sams and E. Leblanc. 2000. Survival and growth of *Listeria monocytogenes* on fresh-cut apple slices and its interaction with *Glomerella cingulata* and *Penicillium expansum*. *Plant Dis.* 84:177-181.
- Couey, H.M. 1989. Heat treatment for control of postharvest diseases and insect pests of fruits. *HortScience* 24:198-202.
- Delp, C.J. 1988. Fungicide resistance in North America. APS Press, St. Paul MN, pp. 11-16.
- Dennis, C. 1983. Postharvest Pathology of Fruits and Vegetables. Acad. Press, London.
- Dychdala, G.R. 1983. Chlorine and chlorine compounds. In: SS. Block (ed) Disinfection, Sterilization, and Preservation. 3<sup>rd</sup> ed., Lea & Febiger Co., Philadelphia, pp. 157-182.
- Eckert, J.W. and N.F. Sommer. 1967. Control of diseases of fruits and vegetables by postharvest treatment. *Ann. Rev. Plant Pathol.* 5:391-432.
- Eckert, J.W. 1988. Dynamics of benzimidazole-resistant penicillia in the development of postharvest decay of citrus and pome fruits. In: C.J. Delp (ed) Fungicide Resistance in North America, APS Press, St. Paul MN, pp. 31-35.
- Ellis, W.O., J.P. Smith, B.K. Simpson and J.H. Oldham. 1991. Aflatoxins in food: occurrence, biosynthesis, effects on organisms, detection, and methods of control. *Crit. Rev. Food Sci. Nut.* 30:403-439.
- Fallik, E., S. Grinberg, M. Gambourg, J.D. Klien and S. Lurie. 1995. Prestorage heat treatment reduces pathogenicity of *Penicillium expansum* in apple fruit. *Plant Pathol.* 45:92-97.
- Forster, H. and J.E. Adaskaveg. 1999. Fludioxinil, a new reduced risk postharvest fungicide for management of fungal decays of stonefruit. *Phytopathology* 89:S26.
- FRAC, Fungicide Resistance Action Committee. 1998. 1998 status report and recommended fungicide resistance management guidelines. Global Crop Prot. Fed., Brussels, Belgium.
- Gould, W.A. 1973. Micro-contamination of horticultural products. *HortScience* 8:116-119.
- Hardenburg, R.E., A.E. Watada and C.Y. Wang. 1986. The commercial storage of fruits and vegetables, and florist and nursery stocks. USDA Agric. Handbook No. 66.
- Harvey, J.M. 1978. Reduction of losses in fresh fruits and vegetables. *Ann. Rev. Phytopath.* 16:321-341.
- Holmes, G.J. and J.W. Eckert. 1999. Sensitivity of *Penicillium digitatum* and *P. italicum* to postharvest citrus fungicides in California. *Phytopathology* 89:716-721.
- Howard, R.J., J.A. Garland and W.L. Seaman. 1994. Diseases and Pests of Vegetable Crops in Canada: An Illustrated Compendium. Co-published by the Entomological Society of Canada and The Canadian Phytopathological Society, Ottawa, Ontario, Canada.
- Hsieh, D.P. and S.H. Gruenwedel. 1990. Microbial toxins. In: C. Winter, J. Seiber and C. Nuckton (eds) Chemicals in the Human Food Chain, Van Nostrand Reinhold, pp. 239-267.
- Janisiewicz, W.J. and Marchi, A. 1992. Control of storage rots on various pear cultivars with a saprophytic strain of *Pseudomonas syringae*. *Plant Dis.* 76:555-560.
- Janisiewicz, W.J. and S.N. Jeffers. 1997. Efficacy of commercial formulation of two biofungicides for control of blue mold and gray mold of apples in cold storage. *Crop Prot.* 16:629-633.
- Jeffries, P. and M.J. Jeger. 1990. The biological control of postharvest diseases of fruit. *Biocontrol News Info.* 11:333-336.
- Kader, A.A. 1985. Biochemical and physiological basis for effects of controlled and modified atmospheres on fruits and vegetables. *Food Technol.* 40:99-104.
- Kelman, A. 1989. Introduction: the importance of research on the control of postharvest diseases of perishable food crops. *Phytopathology* 79:1374.
- Klien, J.D., W.S. Conway, B.D. Whitaker and C.E. Sams. 1997. *Botrytis cinerea* decay in apples is inhibited by postharvest heat and calcium treatment. *J. Amer. Soc. Hort. Sci.* 122:91-94.



- Krieg, N.R. and J.G. Holt. 1984. Bergey's manual of systemic bacteriology. Vol 1. Williams and Wilkins, Baltimore MD.
- Luvisi, D.A., H.H. Shorey, J.L. Smilanick, J.F. Thompson, B.H. Gump and J. Knutson. 1992. Sulfur dioxide fumigation of table grapes. Univ. Calif., Div. Agric. Nat. Res., Bull. No. 1932.
- Mari, M., T. Cembali, E. Baraldi and L. Casalini. 1999. Peracetic acid and chlorine dioxide for postharvest control of *Monilinia laxa* in stone fruits. Plant Dis. 83:773-776.
- Meheriuk, M. and A.P. Gaunce. 1994. Temperate fruits. In: R.E. Paull and J.W. Armstrong (eds) Insect Pests and Fresh Horticultural Products: Treatments and Responses, CAB Intl, Wallingford UK, pp. 291-307.
- Ogawa, J.M. and B.T. Manji. 1984. Control of postharvest diseases by chemical and physical means. In: H. Moline (ed) Postharvest Pathology of Fruits and Vegetables: Postharvest Losses in Perishable Crops. Univ. Calif., Agric. Exp. Sta., U.C. Bull. 1914 (Pub. NE-87), pp. 55-66.
- Ogawa, J.M. and H. English. 1991. Diseases of Temperate Zone Tree Fruit and Nut Crops. Univ. Calif., Div. Agric. Nat. Res., Pub. 3345.
- Ogawa, J.M., J.E. Adaskaveg and K.E. Conn. 1992. Efficacy of iprodione wax/oil mixtures for control of postharvest decay of fruit caused by *Rhizopus* and *Alternaria* spp. Phytopathology 82:1064.
- Phillips, D.J. 1984. Mycotoxins as a postharvest problem. In: H.E. Moline (ed) Postharvest Pathology of Fruits and Vegetables: Postharvest Losses in Perishable Crops. Univ. Calif., Agric. Exp. Sta., Bull No. 1914 (Pub. NE-87), pp. 50-54.
- Pierson, C.F., M.J. Ceponis, and L.P. McColloch. 1971. Market Diseases of Apples, Pears, and Quinces. USDA, ARS, Agric. Handbook No. 376.
- Pusey, P.L. and C.L. Wilson. 1984. Postharvest biological control of stone fruit brown rot by *Bacillus subtilis*. Plant Dis. 68:753-756.
- Pusey, P.L. 1989. Use of *Bacillus subtilis* and related organisms as biofungicides. Pesticide Sci. 27:133-140.
- Pusey, P.L. 1996. Micro-organisms as agents in plant disease control. In: L.G. Copping (ed) Crop Protection Agents from Nature: Natural Products and Analogues, Royal Soc. Chem., Cambridge UK, pp. 426-443.
- Roberts, R.G. 1990. Biological control of gray mold of apple by *Cryptococcus laurentii*. Phytopathology 80:526-530.
- Roberts, R. 1994. Integrating biological control into postharvest disease management strategies. HortScience 29:758-762.
- Roberts, R.G. and S.T. Reymand. 1994. Chlorine dioxide for reduction of postharvest pathogen inoculum during handling of tree fruits. Appl. Environ. Micro. 60:2864-2868.
- Sams, C.E., W.S. Conway, J.A. Abbott, R.J. Lewis and N. Ben-Shalom. 1993. Firmness and decay of apple following postharvest pressure infiltration of calcium and heat treatment. J. Amer. Soc. Hort. Sci. 118:623-627.
- Sams, C.E. 1994. Management of postharvest disease resistance in horticultural crops: introduction to the colloquium. HortScience 29:746.
- Sholberg, P.L. and G.R. Owen. 1990. Populations of propagules of *Penicillium* spp. during immersion dumping of apples. Can. Plant Dis. Surv. 70:11-14.
- Sholberg, P.L., A.G. Reynolds and A.P. Gaunce. 1996. Fumigation of table grapes with acetic acid to prevent postharvest decay. Plant Dis. 80:1425-1428.
- Sholberg, P.L. 1998. Fumigation of fruit with short-chain organic acids to reduce the potential of postharvest decay. Plant Dis. 82:689-693.
- Sholberg, P.L. and K.E. Bedford. 1999. Use of cyprodinil for control of *Botrytis cinerea* on apple. Phytopathology 89:S72.
- Snowdon, A.L. 1990. Color Atlas of Post-Harvest Diseases and Disorders of Fruits and Vegetables, Vol. 1, General Introduction and Fruits. CRC Press, Boca Raton FL.
- Snowdon, A.L. 1992. Color Atlas of Post-Harvest Diseases and Disorders of Fruits and Vegetables, Vol. 2,

- Vegetables. CRC Press, Boca Raton FL.
- Sommer, N.F. 1989b. Suppressing postharvest disease with handling practices and controlled environments. In: J.H. LaRue and R.S. Johnson (ed) Peaches, Plums, and Nectarines Growing and Handling for Fresh Market. Univ. Calif., DANR Pub. No. 3331, pp. 179-190.
- Spotts, R.A. and B.B. Peters. 1980. Chlorine and chlorine dioxide for control of 'd' Anjou' pear decay. *Plant Dis.* 64:1095-1097.
- Spotts, R.A. and B.B. Peters. 1981. The effect of relative humidity on spore germination of pear decay fungi and 'd' Anjou' pear decay. *Acta Hort.* 124:75-78.
- Spotts, R.A. 1984. Environmental modification for control of postharvest decay. In: H.E. Moline (ed) *Postharvest Pathology of Fruits and Vegetables: Postharvest Losses in Perishable Crops*, Univ. of Calif., Agric. Exp. Station, Bull. No. 1914 (Pub. NE-87), pp. 67-72.
- Spotts, R.A. and P.M. Chen. 1987. Prestorage heat treatment for control of decay of pear fruit. *Phytopathology* 77:1578-1582.
- Spotts, R.A. and R.L. Dobson. 1989. Effects of the experimental fungicide RH 886 on *Mucor piriformis*. *Pesticide Sci.* 25:391-399.
- Spotts, R.A. 1990. Bull's-eye rot. In: A.L. Jones and H.S. Aldwinckle (eds) *Compendium of Apple and Pear Diseases*, APS Press, St. Paul MN, pp 56.
- Spotts, R.A., L.A. Cervantes, T.J. Facticeau and T. Chand-Goyal. 1998. Control of brown rot and blue mold of sweet cherry with preharvest iprodione, postharvest *Cryptococcus infirmo-miniatius*, and modified packaging. *Plant Dis.* 82:1158-1160.
- Spotts, R.A., L.A. Cervantes and E.A. Mielke. 1999. Variability in postharvest decay among apple cultivars. *Plant Dis.* 83:1051-1054.
- Stevens, C., V.A. Khan, J.Y. Lu, C.L. Wilson, P.L. Pusey, M.K. Kabwe, E.C.K. Igwegbe, E. Chalutz and S. Droby. 1998. The germicidal and hormetic effects of UV-C light on reducing brown rot disease and yeast microflora of peaches. *Crop Protection* 17:75-84.
- Sugar, D., T.L. Righetti, E.E. Sanchez and H. Khemira. 1992. Management of nitrogen and calcium in pear tree for enhancement of fruit resistance to postharvest decay. *HortTechnol.* 2:382-387.
- Sugar, D. R.G. Roberts, R.J. Hilton, T.L. Righetti and E.E. Sanchez. 1994. Integration of cultural methods with yeast treatment for control of postharvest fruit decay in pear. *Plant Dis.* 78:791-795.
- Sugar, D. and R. Spotts. 1995. Preharvest strategies to reduce postharvest decay. 1995 Wash. Tree Fruit Postharv. Conf. Proc., Wash. State Hort. Assoc., Wenatchee WA.
- Swalding, I.R. and P.J. Jeffries. 1998. Antagonistic properties of two bacterial biocontrol agents of grey mould disease. *Biocontrol Sci. Technol.* 8:439-448.
- Utkhede, R.S. and P.L. Sholberg. 1993. Biological control of diseases of temperate fruit trees. *Curr. Top. Bot. Res.* 1:313-334.
- Vinas, I., J. Usall, N. Texido and V. Sanchis. 1998. Biological control of major postharvest pathogens in apple with *Candida sake*. *Int. J. Food Micro.* 40:9-16.
- Wells, J. M. and J. E. Butterfield. 1997. *Salmonella* contamination associated with bacterial soft rot of fresh fruits and vegetables in the marketplace. *Plant Disease* 81:867-872.
- Wilson, C.L. and P.L. Pusey. 1985. Potential for biological control of postharvest plant diseases. *Plant Disease* 69:375-378.
- Wisniewski, M., C. Biles, S. Droby, R. McLaughlin, C. Wilson, and E. Chalutz. 1991. Mode of action of the postharvest biocontrol yeast *Pichia guilliermondii*. 1. Characterization of attachment to *Botrytis cinerea*. *Physiol. Molec. Plant Pathol.* 39:245-258.
- Wisniewski, M.E. and C.L. Wilson. 1992. Biological control of postharvest diseases of fruits and vegetables: recent advances. *HortScience* 27:94-98.

### Web Sites

A quickly changing resource on plant pathology: <http://www.sci.soc.org>

Postharvest information network: <http://postharvest.tfrec.wsu.edu/>  
 Postharvest technology research and information center: <http://postharvest.ucdavis.edu/>  
 Sidney postharvest lab with links to other sites: <http://www.postharvest.com.au/Default.html>

Table 1. Important postharvest diseases of fruit.

<b>Fruit</b>	<b>Disease</b>	<b>Causal Agent</b>	<b>Fungal class/Type</b>	
Avocado	Anthracnose	<i>Glomerella cingulata</i>	Pyrenomycete	
	Cercospora spot	<i>Pseudocercospora purpurea</i>	Hyphomycete	
	Dothiorella rot	<i>Botryosphaeria ribis</i>	Loculoascomycete	
	Scab	<i>Sphaceloma persae</i>	Coleomycete	
	Stem-end rots	<i>B. theobromae</i> , <i>Phomopsis perseae</i> , <i>Thyronectria pseudotrichia</i>	Deuteromycetes	
Banana	Anthracnose	<i>Colletotrichum musae</i>	Coelomycete	
	Cigar-end rot	<i>Trachysphaera fructigena</i> , <i>Verticillium theobromae</i>	Deuteromycetes	
	Crown rot	<i>C. musae</i> , <i>Fusarium pallidoroseum</i> , <i>V. theobromae</i>	Deuteromycetes	
	Finger rot	<i>B. theobromae</i>	Coelomycete	
	Pitting disease	<i>Pyricularia grisea</i>	Hyphomycete	
	Sigatoka disease	<i>Mycosphaerella</i> spp.	Loculoascomycete	
	Gray mold	<i>Botrytis cinerea</i>	Hyphomycete	
Berries	Leak	<i>Mucor</i> spp.	Zygomycete	
	Leather rot	<i>Phytophthora</i> spp.	Oomycete	
	Alternaria rot	<i>Alternaria</i> spp.	Hyphomycete	
Citrus	Anthracnose	<i>C. gloeosporioides</i>	Coelomycete	
	Bacterial canker	<i>Xanthomonas campestris</i>	Bacterium	
	Black pit	<i>Pseudomonas syringae</i>	Bacterium	
	Black spot	<i>Phyllosticta citricarpa</i>	Coelomycete	
	Blue mold	<i>Penicillium italicum</i>	Hyphomycete	
	Brown rot	<i>Phytophthora</i> spp.	Oomycete	
	Greasy spot	<i>Mycosphaerella citri</i>	Loculoascomycete	
	Green mold	<i>P. digitatum</i>	Hyphomycete	
	Scab	<i>Elsinoe fawcettii</i>	Loculoascomycete	
	Sour rot	<i>Geotrichum candidum</i>	Hyphomycete	
	Stem-end rots	<i>D. gregaria</i> , <i>Phomopsis citri</i> , <i>B. theobromae</i>	Coelomycetes	
	Kiwifruit	Gray mold	<i>B. cinerea</i>	Hyphomycete
	Grape	Aspergillus rot	<i>Aspergillus niger</i>	Hyphomycete
Blue mold		<i>Penicillium</i> spp.	Hyphomycete	
Gray mold		<i>B. cinerea</i>	Hyphomycete	
Rhizopus rot		<i>Rhizopus</i> spp.	Zygomycete	
Mango	Anthracnose	<i>C. gloeosporioides</i>	Coelomycete	
	Botryodiplodia rot	<i>B. theobromae</i>	Coelomycete	
	Stem-end rots	<i>B. theobromae</i> , <i>Phomopsis</i> spp.	Coelomycete	
Papaya	Anthracnose	<i>C. gloeosporioides</i>	Coelomycete	

	Black rot	<i>Phoma caricae-papayae</i>	Coelomycete
	Phytophthora rot	<i>P. palmivora</i>	Oomycete
	Rhizopus rot	<i>R. stolonifer</i>	Zygomycete
	Stem-end rot	<i>B. theobromae</i> , <i>Phomopsis</i> spp.	Coelomycetes
Pineapple	Black rot	<i>Thielaviopsis paradoxa</i>	Hyphomycete
	Fruitlet core rot	<i>Fusarium moniliforme</i> <i>P. funiculosum</i>	Hyphomycete
Pome fruit (apple, pear)	Bitter rot	<i>C. gloeosporioides</i>	Coelomycete
	Black rot	<i>Sphaeropsis malorum</i>	Coelomycete
	Blue mold	<i>Penicillium expansum</i> , <i>Penicillium</i> spp.	Hyphomycetes
	Brown rot	<i>Monilinia</i> spp.	Hyphomycete
	Bull's-eye rot	<i>Cryptosporiopsis curvispora</i>	Hyphomycete
	Gray mold	<i>B. cinerea</i>	Hyphomycete
	Moldy core	<i>Alternaria</i> spp., others	Hyphomycetes
	Mucor rot	<i>Mucor piriformis</i>	Zygomycete
	White rot	<i>D. gregaria</i>	Coelomycete
Stone fruit (cherry etc.)	Alternaria rot	<i>A. alternata</i>	Hyphomycete
	Blue mold	<i>P. expansum</i>	Hyphomycete
	Brown rot	<i>Monilinia</i> spp.	Hyphomycetes
	Rhizopus rot	<i>Rhizopus</i> spp.	Zygomycete

Table 2. Important postharvest diseases of vegetables

Vegetable	Disease	Causal Agent	Fungal Class/Type
Bulbs (Onion, garlic)	Bacterial soft rot	<i>Erwinia caratovora</i>	Bacterium
	Black rot	<i>Aspergillus niger</i>	Hyphomycete
	Blue mold rot	<i>Penicillium</i> spp.	Hyphomycete
	Fusarium basal rot	<i>Fusarium oxysporum</i>	Hyphomycete
	Neck rot	<i>Botrytis</i> spp.	Hyphomycete
	Purple blotch	<i>Alternaria porri</i>	Hyphomycete
	Sclerotium rot	<i>Sclerotium rolfsii</i>	Agonomycete
	Smudge	<i>Colletotrichum circinans</i>	Coelomycete
Crucifers (Cabbage, etc.)	Alternaria leaf spot	<i>Alternaria</i> spp.	Hyphomycete
	Bacterial soft rot	<i>E. caratovora</i>	Bacterium
	Black rot	<i>Xanthomonas campestris</i>	Bacterium
	Downy mildew	<i>Peronospora parasitica</i>	Oomycete
	Rhizoctonia rot	<i>Rhizoctonia solani</i>	Agonomycete
	Ring spot	<i>Mycosphaerella brassicicola</i>	Loculoascomycete
	Virus deases	Cauliflower mosaic virus Turnip mosaic virus	Virus
	Watery soft rot	<i>Sclerotinia</i> spp.	Discomycete
Cucurbits (Cucumber etc.)	White blister	<i>Albugo candida</i>	Oomycete
	Anthracnose	<i>Colletotrichum</i> spp.	Coelomycete
	Bacterial soft rot	<i>Erwinia</i> spp.	Bacterium
	Black rot	<i>Didymella bryoniae</i>	Loculoascomycete
	Botryodiplodia rot	<i>Botryodiplodia theobromae</i>	Coelomycete
Charcoal rot	<i>Macrophomina phaseolina</i>	Coelomycete	



	Phoma rot	<i>Phoma lycopersici</i>	Hyphomycete
	Phomopsis rot	<i>Phomopsis</i> spp.	Coelomycetes
	Phytophthora rot	<i>Phytophthora</i> spp.	Oomycete
	Pleospora rot	<i>Stemphylium herbarum</i>	Hyphomycete
	Rhizopus rot	<i>Rhizopus</i> spp.	Zygomycetes
	Sclerotium rot	<i>S. rolfsii</i>	Agonomycete
	Soil rot	<i>R. solani</i>	Agonomycete
	Sour rot	<i>Geotrichum candidum</i>	Hyphomycete
	Watery soft rot	<i>Sclerotinia</i> spp.	Discomycetes
Miscellaneous			
- Artichokes	Gray mold	<i>Botrytis cinerea</i>	Hyphomycete
	Watery soft rot	<i>Sclerotinia sclerotiorum</i>	Discomycete
- Asparagus	Bacterial soft rot	<i>Erwinia</i> or <i>Pseudomonas</i> spp.	Bacteria
	Fusarium rot	<i>Fusarium</i> spp.	Hyphomycete
	Phytophthora rot	<i>Phytophthora</i> spp.	Oomycete
	Purple spot	<i>Stemphylium</i> spp.	Hyphomycete
- Celery	Bacterial soft rot	<i>Erwinia</i> or <i>Pseudomonas</i> spp.	Bacteria
	Brown spot	<i>Cephalosporium apii</i>	Hyphomycete
	Cercospora spot	<i>Cercospora apii</i>	Hyphomycete
	Gray mold	<i>Botrytis cinerea</i>	Hyphomycete
	Licorice rot	<i>Mycocentrospora acerina</i>	Hyphomycete
	Phoma rot	<i>Phoma apiicola</i>	Coelomycete
	Pink rot	<i>Sclerotinia</i> spp.	Discomycete
	Septoria spot	<i>Septoria apiicola</i>	Coelomycete
- Lettuce	Bacterial rot	<i>Erwinia</i> , <i>Pseudomonas</i> , <i>Xanthomonas</i> spp.	Bacteria
	Gray mold rot	<i>B. cinerea</i>	Hyphomycete
	Rhizoctonia rot	<i>R. solani</i>	Agonomycete
	Ringspot	<i>Microdochium panattonianum</i>	Hyphomycete
	Septoria spot	<i>S. lactucae</i>	Coelomycete
	Stemphylium spot	<i>Stemphylium herbarum</i>	Hyphomycete
	Watery soft rot	<i>Sclerotinia</i> spp.	Discomycete