

References for Fission Isomers

62Po09 Spontaneous Fission with an Anomalously Short Period. I.

S. M. Polikanov, V. A. Druin, V. A. Karnaukhov, V. L. Mikheev, A. A. Pleve, N. K. Skobelev, G. M. Ter-Akopyan, V. A. Fomichev, Zhur. Eksppl. i Teoret. Fiz. 42, 1464 (1962); Soviet Phys. JETP 15, 1016 (1962).

Nuclear Structure: Fission ^{242}Am ; measured not abstracted; deduced nuclear properties.

62Pe26 Spontaneous Fission with an Anomalously Short Period. II

V. P. Perelygin, S. P. Almazova, B. A. Gvozdev, Y. T. Chuburkov, Zhur. Eksppl. i Teoret. Fiz. 42, 1472 (1962); Soviet Phys. JETP 15, 1022 (1962).

63FI08 Formation of a Spontaneously Fissioning Isomer in Reactions Involving α Particles and Deuterons

G. N. Flerov, S. M. Polikanov, K. A. Gavrilov, V. L. Mikheev, V. P. Perelygin, A. A. Pleve, Zh. Ekspерим. i Teor. Fiz. 45, 1396 (1963); Soviet Phys. JETP 18, 964 (1964).

63Pe27 Half-Life of a Spontaneously Fissioning Isomer

V. G. Perelygin, S. P. Tretyakova, Zh. Ekspерим. i Teor. Fiz. 45, 863 (1963); Soviet Phys. JETP 18, 592 (1964).

Nuclear Structure: Fission ^{238}U ; measured not abstracted; deduced nuclear properties.

65FI04 The Excitation Function and the Isomeric Yield Ratio for the 14 msec Fissioning Isomer from Deuteron Irradiation of Plutonium

G. N. Flerov, A. A. Pleve, S. M. Polikanov, E. Ivanov, N. Martalugu, D. Poenaru, N. Vilcov, Rev. Roumaine Phys. 10, 217 (1965).

Nuclear Structure: ^{242}Am ; measured not abstracted; deduced nuclear properties.

65Le22 Decay of the Am 242m 14-msec Isomer

R. B. Leachman, B. H. Erkkila, Bull. Am. Phys. Soc. 10, No. 9, 1204, P12 (1965)

Nuclear Structure: ^{242}Am ; measured not abstracted; deduced nuclear properties.

65Li05 The Formation of a Spontaneously Fissioning Isomer in the Capture of Neutrons by Am

A. F. Linev, B. N. Markov, A. A. Pleve, S. M. Polikanov, Nucl. Phys. 63, 173 (1965).

Radioactivity: ^{242}Am ; measured $T_{1/2}$, SF. $^{243}\text{Am}(n, 2n)$, $E=14$ MeV; measured σ .

66Br23 A Study of Nuclear Isomers Which Decay by Spontaneous Fission

D. S. Brenner, L. Westgaard, S. Bjornholm, Nucl. Phys. 89, 267 (1966).

Radioactivity: ^{242}Am isomer [from $^{242}\text{Pu}(d,2n)$]; measured $T_{1/2}$ (SF), $E(\text{fragment})$ -spectrum. Enriched target. $^{242}\text{Pu}(d,2n)$, (d,F), $E = 12$ MeV; measured $\sigma(F)(\text{delayed})/\sigma(F)(\text{prompt})$. Enriched target. ^{232}Th , ^{235}U , ^{239}Pu , ^{241}Am , $^{243}\text{Am}(d, xn)(d,F)$, $E = 12$ MeV; $^{243}\text{Am}(p,xn)(p,F)$, $E = 13$ MeV; measured upper limits $\sigma(F)(\text{delayed})/\sigma(F)(\text{prompt})$. Enriched targets.

66Ma48 Structure of Spontaneously Fissionable Isomers

L. A. Malov, S. M. Polikanov, V. G. Solovev, Yadern. Fiz. 4, 528 (1966); Soviet J. Nucl. Phys. 4, 376 (1967).

67Bj03 Excitation Energy of the Spontaneously Fissioning Isomeric State in ^{240}Am

S. Bjornholm, J. Borggreen, L. Westgaard, V. A. Karnaukhov, Nucl. Phys. A95, 513 (1967).

Nuclear Reactions: $^{240}\text{Pu}(d,2n)$, $E = 12.1$ MeV; $^{240}\text{Pu}(p,n)$, $E = 10$, 3-11.3 MeV; measured σ (delayed fission). $^{241}\text{Pu}(p,2n)$, $E = 9.6$ -13.6 MeV; measured σ (delayed fission); deduced threshold. Enriched target. ^{240m}Am measured $T_{1/2}$ for spontaneous fission.

67Bo23 A New Spontaneously Fissioning Isomer: ^{238}Am

J. Borggreen, Y. P. Gangsksy, G. Sletten, S. Bjornholm, Phys. Letters 25B, 402 (1967).

Nuclear Structure: ^{238}Am ; measured not abstracted; deduced nuclear properties.

67FI03 Excitation Energy of Spontaneously Fissioning Isomer $242m\text{-Am}$

G. N. Flerov, A. A. Pleve, S. M. Polikanov, S. P. Tretyakova, N. Martalugu, D. Poenaru, M. Sezon, I. Vilcov, N. Vilcov, Nucl. Phys. A97, 444 (1967).

Nuclear Reactions: $^{243}\text{Am}(n,2nF)$, $E = 8$ -14.4 MeV; measured $\sigma(E)$, n, F-delay. ^{242}Am deduced level, $T_{1/2}$. Enriched target.

67FI08 A Study of the Spontaneously-Fissioning Isomer of ^{242}Am Through the $^{241}\text{Am}(n,\gamma)$ Reaction

G. N. Flerov, A. A. Pleve, S. M. Polikanov, S. P. Tretyakova, I. Boca, M. Sezon, I. Vilcov, N. Vilcov, Nucl. Phys. A102, 443 (1967).

Nuclear Reactions: Fission $^{241}\text{Am}(n,\gamma)$, $E=0$ -6.5 MeV; measured $\sigma(E)$.

Radioactivity: Fission ^{244m}Am [from $^{243}\text{Am}(n,\gamma)$]; measured $T_{1/2}$ (SF).

67Ga04 Investigation of the Reaction $^{238}\text{U} + \text{B}^{11}$, Which Leads to the Spontaneously-Fissioning Isomer ^{242}Am

Y. P. Gangskii, B. N. Markov, S. M. Polikanov, G. Jungclaussen, Yadern. Fiz. 5, 22 (1967); Soviet J. Nucl. Phys. 5, 16 (1967).

Nuclear Structure: ^{242}Am ; measured not abstracted; deduced nuclear properties.

67Vi01 On the Spin Value of the 14-msec Spontaneously Fissioning Isomer of ^{242}Am

N. Vilcov, Rev. Roumaine Phys. 12, 487 (1967).

Nuclear Structure: ^{242}Am ; measured not abstracted; deduced nuclear properties.

68Bj04 Investigation of (d,p) and (d,t) Reactions Leading to Spontaneously Fissile Isomeric States

S. Bjornholm, I. Borggreen, Y. P. Gangskii, G. Sletten, Yadern. Fiz. 8, 459 (1968); Soviet J. Nucl. Phys. 8, 267 (1969).

Nuclear Reactions: $^{241}, ^{243}\text{Am}(d,p)$, (d,t), $E=9$ -13 MeV; measured $\sigma(E)$; deduced isomeric ratio.

68Ca23 Autocorrelation Effects in the Neutron Induced Fission Cross Section of ^{235}U

M. G. Cao, E. Migneco, J. P. Theobald, Phys. Lett. 27B, 409 (1968).

Nuclear Reactions: Fission $^{235}\text{U}(\text{n},\text{F})$, $E=0.006\text{-}3$ keV; measured $\sigma(E)$. ^{236}U deduced resonance, autocorrelation, intermediate state, shape isomer. Reanalysis of data.

68Er01 Energy of ^{242}Am and ^{242m}Am Fission Fragments

B. H. Erkkila, R. B. Leachman, Nucl. Phys. A108, 689 (1968).

Radioactivity: Fission $^{242m}\text{Am}(\text{SF})$ [from $^{242}\text{Pu}(\text{d},2\text{n})$]; measured $T_{1/2}$, $E(\text{fragment})$. ^{252}Cf measured $E(\text{fragment})$.

Nuclear Reactions: Fission $^{240}\text{Pu}(\text{d},\text{F})$, $E=7.6\text{-}14$ MeV; measured $\sigma(E)$; $E(\text{fragment})$. ^{230}Th , ^{233}U , $^{242}\text{Pu}(\text{d},\text{F})$, $E=14$ MeV; measured $\sigma(E(\text{fragment}))$.

68Mi14 Resonance Grouping Structure in Neutron Induced Subthreshold Fission of ^{240}Pu

E. Migneco, J. P. Theobald, Nucl. Phys. A112, 603 (1968).

Nuclear Reactions: $^{240}\text{Pu}(\text{n},\text{F})$, $E=0.2$ to 8 keV; measured $\sigma(\text{nf})(E)$. ^{241}Pu resonances deduced F-width.

68WoZZ Short-Lived Spontaneous Fission Isomers

K. L. Wolf, R. Vandenbosch, Bull. Am. Phys. Soc. 13, No. 11, 1407, CF4(1968)

Nuclear Reactions: $^{238}\text{U}(\alpha,2\text{n})$, $E=21\text{-}42$ MeV; measured isomer ratio, $\sigma(\alpha)$. ^{240}Pu deduced $T_{1/2}$, spontaneous fission.

69Bj02 Intermediate States in Fission

S. Bjornholm, V. M. Strutinsky, Nucl. Phys. A136, 1 (1969).

69Bo25 Population of the Spontaneously Fissioning Isomer $^{244}\text{mf-Am}$ Through the (n,γ) Reaction

I. Boca, N. Martalogu, M. Sezon, I. Vilcov, N. Vilcov, G. N. Flerov, A. A. Pleve, S. M. Polikanov, S. P. Tretyakova, Nucl. Phys. A134, 541 (1969).

Nuclear Reactions: $^{243}\text{Am}(\text{n},\gamma)$, (n,F) , $E = 0.3\text{-}4$ MeV; measured $\sigma(E)$. ^{244}Am deduced $T_{1/2}$, spontaneous fission. Enriched target.

69El06 Discussion on Papers SM 122/110 and SM 122/29

A. J. Elwyn, A. T. G. Ferguson, 2nd Symp. Phys. Chem. of Fission, Vienna, Intern. At. Energy Agency, Vienna, p. 457 (1969).

69Ja01 Fission Components in ^{242}Pu Resonances

G. D. James, Nucl. Phys. A123, 24 (1969).

Nuclear Reactions: $^{242}\text{Pu}(\text{n},\text{F})$, $E=16$ eV-35 keV; measured $\sigma(E)$. ^{243}Pu deduced resonances, resonance parameters. Enriched target.

69JoZU

A. B. Jorgensen, S. M. Polikanov, G. Sletten, Priv. Comm., quoted by 70PO01, unpublished (1969)

69Ka27 Photofission of Even-Even Nuclei and Structure of the Fission Barrier

S. P. Kapitza, N. S. Rabotnov, G. N. Smirenkin, A. S. Soldatov, L. N. Usachev, Y. M. Tsipenyuk, ZhETF Pisma v Redaktsiyu 9, 128 (1969); JETP Letters 9, 73 (1969).

Nuclear Reactions: Fission ^{232}Th , ^{238}U , 240 , 242 , $^{238}\text{Pu}(\gamma, \text{F})$, $E < 5\text{-}8$ MeV; measured $\sigma(E; E(\text{fragment}), \theta(\text{fragment}))$. ^{232}Th , ^{238}U , 238 , 240 , ^{242}Pu deduced fission barrier structure.

69Kr12 The Moment of Inertia of the Fission Isomer

J. Krumlinde, Phys. Letters 30B, 221 (1969).

Nuclear Structure: Fission ^{238}U , ^{242}Pu , ^{246}Cm , $^{250}\text{Cf}(\text{SF})$; calculated moments of inertia. Cranking model.

69La14 Spontaneously Fissioning Isomers in U, Np, Pu and Am Isotopes

N. L. Lark, G. Sletten, J. Pedersen, S. Bjornholm, Nucl. Phys. A139, 481 (1969).

Radioactivity: Fission ^{236m}U , ^{239m}Np , ^{236m}Pu , ^{237m}Pu , ^{240m}Pu , ^{241m}Pu , ^{242m}Pu , ^{243m}Pu , ^{239m}Am , $^{241m}\text{Am}(\text{SF})$; measured $T_{1/2}$.

Nuclear Reactions: ^{235}U , 239 , 241 , $^{242}\text{Pu}(\text{d},\text{p})$, $^{240}\text{Pu}(\text{d},\text{X})$, $E=11\text{-}13$ MeV; measured σ delayed fission. $^{237}\text{Np}(\text{p},2\text{n})$, $E=9\text{-}14$ MeV; $^{240}\text{Pu}(\text{p},2\text{n})$, $E=10\text{-}13$ MeV; $^{242}\text{Pu}(\text{p},2\text{n})$, $E=8.8\text{-}13$ MeV; measured σ delayed fission; deduced thresholds. ^{238}U , $^{237}\text{Np}(\text{d},\text{X})$, ^{239}Pu , $^{241}\text{Pu}(\text{d},2\text{n})$, $E=13$ MeV; measured σ delayed fission. $^{237}\text{Np}(\text{p},2\text{n})$, $E=13$ MeV; measured σ ground state. Enriched targets.

69Na20 On the Detection of Spontaneously Fissioning Isomer States

L. Nagy, T. Nagy, I. Vinnay, KFKI Kozlemen. 17, 165 (1969).

69Me11 Fission Isomerism Induced by Helium Ions

V. Metag, R. Repnow, P. Von Brentano, J. D. Fox, Z. Physik 226, 1 (1969).

Nuclear Reactions: 233 , 235 , 236 , ^{238}U , ^{237}Np , $^{239}\text{Pu}(\alpha,2\text{n})$, $E=26.1$ MeV; measured α . 235 , 237 , 238 , ^{240}Pu , ^{239}Am , ^{241}Cm deduced $T_{1/2}$ (SF-isomer). $^{239}\text{Pu}({}^3\text{He}, 2\text{np})$, $E=30$ MeV; measured σ . ^{239}Am deduced $T_{1/2}$ (SF-isomer). $^{236}\text{U}(\alpha,2\text{n})$, $E=26$ MeV; measured σ . ^{239}Pu deduced $T_{1/2}$ (SF-isomer). $^{237}\text{Np}({}^3\text{He}, {}^3\text{He}, \text{np})$, $({}^3\text{He}, 2\text{np})$, $E=26, 30$ MeV; measured σ . 237 , 238 , ^{239}Pu deduced $T_{1/2}$ (SF-isomer).

69MeZX Charged-Particle Studies of Isomeric Fission

V. Metag, R. Repnow, P. von Brentano, J. D. Fox, Proc. Symp. Phys. Chem. Fission, 2nd, Vienna, Intern. At. En. Agency, p. 449 (1969).

69Ni13 On the Nuclear Structure and Stability of Heavy and Superheavy Elements

S. G. Nilsson, C. F. Tsang, A. Sobczewski, Z. Szymanski, S. Wycech, C. Gustafson, I. -L. Lamm, P. Moller, B. Nilsson, Nucl. Phys. A131, 1 (1969).

69SIZZ Discussion on Papers SM-122/110 and SM-122/29

G. Sletten, S. M. Polikanov, Symp. Phys. Chem. Of Fission, 2nd, Vienna, Intern. At. Energy Agency, Vienna, p. 461 (1969).

Radioactivity: Fission ^{237m}Am , ^{239m}Am , ^{240m}Am , ^{241m}Cm , ^{243m}Cm , ^{244m}Am ; measured $T_{1/2}$.

69VaZX Spontaneous Fission Isomers with Very Short Half-Lives

R. Vandenbosch, K. L. Wolf, Proc. Symp. Phys. Chem. Fission, 2nd, Vienna, Intern. At. En. Agency, Vienna, p. 439 (1969).

Radioactivity: Fission 236 , 237 , 238 , 239 , 240 Pu(SF); measured $T_{1/2}$.

Nuclear Reactions: 236 , 238 U(α ,3n), 238 U(α ,2n), E=21-42 MeV; measured $\sigma(E)$; deduced isomer ratios.

69Vo18 Analysis of Neutron Fission of the Odd-Even Nuclei Pa^{231} , Np^{237} , and Am^{241}

P. E. Vorotnikov, Yadern. Fiz. 9, 538 (1969); Soviet J. Nucl. Phys. 9, 308 (1969).

Nuclear Reactions: 231 Pa, 237 Np, 241 Am(n,γ), (n, F), E=0-1 MeV; calculated $\sigma(E)$. 232 Pa, 238 Np, 242 Am calculated level-width, fission barrier penetrability.

70AlZT On Vibrational Type Resonances in Fission

J. Almberger, S. Jagare, Ann. Rept., Research Inst. Phys., Stockholm, p. 217 (1970).

Nuclear Structure: Fission 239 , 242 Pu, 242 Am; calculated fission branching ratios. Vibrational-type resonances.

70Be44 Search for a Long-Lived Spontaneous Fission Isomer of 241 Pu

C. E. Bemis, Jr., R. J. Silva, J. E. Bigelow, A. M. Friedman, Inorg. Nucl. Chem. Lett. 6, 747 (1970); ORNL-4581, p. 36 (1970).

Nuclear Reactions: 240 Pu(n,γ), E=thermal, > 1 MeV; 242 Pu($n, 2n$), E > 6.2 MeV; 238 U(α, n), E=40 MeV; measured σ . 241 Pu deduced no 0.3-yr SF-isomer.

70Bj02 Search for New Islands of Fission Isomerism

S. Bjornholm, J. Borggreen, E. K. Hyde, Nucl. Phys. A156, 561 (1970).

Nuclear Reactions: 197 Au(HI,X), E=5-10 MeV/nucleon for HI= 11 B, 12 C, 14 N, 16 O; measured $\sigma(E)$ for SF-isomers. 200 , 201 , 202 , 203 , 204 Po, 201 , 202 , 203 , 204 , 205 At, 202 , 203 , 204 , 205 , 206 , 207 Rn, 202 , 203 , 204 , 205 , 206 , 207 , 208 , 209 , 210 Fr deduced no SF-isomer ($\sigma < 0.1 \mu b$) with $2ns < T_{1/2} < 2000s$.

70Br32 Fission of Odd-A Uranium and Plutonium Isotopes Excited by (d,p), (t,d), and (t,p) Reactions

H. C. Britt, J. D. Cramer, Phys. Rev. C2, 1758 (1970).

Nuclear Reactions: 234 , 236 , 238 U, 242 Pu(d,pF), 233 , 235 U, 239 Pu(t,pF), 236 U, 240 , 242 Pu(t,dF), E=18 MeV; measured (p)(fragment)(θ), (d)(fragment)(θ). 235 , 237 , 239 U, 241 , 243 Pu deduced fission probabilities.

70Bu02 Systematics of Plutonium Fission Isomers

S. C. Burnett, H. C. Britt, B. H. Erkkila, W. E. Stein, Phys. Lett. 31B, 523 (1970).

Radioactivity: Fission 233m Pu, 237m Pu, 238m Pu, 239m Pu, 240m Pu(SF); measured $T_{1/2}$.

Nuclear Reactions: 233 , 235 , 236 , 238 U($\alpha, 2n$), E=20-28 MeV; 234 U(α, xn), 236 U(α, n), 238 U(α, n), ($\alpha, 3n$), E=20-29 MeV; measured isomeric σ ratios(E); deduced thresholds for SF-isomer production.

70Da05 Production of Spontaneously Fissioning Isomers 242 Am and 244 Am by Slow Neutron Capture

B. Dalhsuren, G. N. Flerov, Y. P. Gangrsky, Y. A. Lazarev, B. N. Markov, Nguyen Cong Khanh, Nucl. Phys. A148, 492 (1970).

Nuclear Reactions: 241 , 243 Am(n,γ), (n, F), E=0.2-20 eV; measured delayed, prompt fission σ ratios, (n)(fission fragment)-delay. 242 , 244 Am [SF-isomers] deduced $T_{1/2}$.

70El03 Short-Lived Fission Isomers from Neutron Studies

A. J. Elwyn, A. T. G. Ferguson, Nucl. Phys. A148, 337 (1970).

Nuclear Reactions: 233 , 234 , 235 , 238 U, 239 Pu(n, γ), E=0.55, 2.2 MeV; measured σ for SF-isomer production; deduced isomeric σ ratios. 234 , 235 , 236 , 239 U, 240 Pu deduced SF-isomers, $T_{1/2}$.

70Ga04 Study of (γ, n) Reactions Leading to Formation of Spontaneously Fissile Isomers of Am

Y. P. Gangrskii, B. N. Markov, Y. M. Tsipenyuk, Yad. Fiz. 11, 54 (1970); Sov. J. Nucl. Phys. 11, 30 (1970).

Nuclear Reactions: 241 , 243 Am(γ, n), E < 9.5-13.5 MeV; measured $\sigma(E)$ for producing SF-isomers. 240 , 242 Am deduced energy of SF isomeric state.

70Ga10 Investigation of the Properties of the Spontaneously Fissioning Isomer 241 Pu in the Reaction (γ, n)

Y. P. Gangrsky, B. N. Markov, Y. M. Tsipenyuk, Phys. Lett. 32B, 182 (1970).

Nuclear Reactions: 242 Pu(γ, nF), E < 8-13 MeV; measured $\sigma(E)$, (γ)(fragment)-delay. 241 Pu deduced SF-isomer $T_{1/2}$.

70Ga34 Production of Spontaneously Fissioning Isomers of Uranium, Plutonium, and Americium in the Neutron Reactions

Y. P. Gangrsky, T. Nagy, I. Vinnay, I. Kovacs, JINR-P3-5528 (1970).

Nuclear Reactions: 232 Th, 235 , 238 U, 239 Pu, 243 Am($n, 2n$), 238 U, 242 Pu, 243 Am(n, n'), E not given; measured SF-isomer production σ .

70Ja16 Excitation Energies of Fissioning Shape Isomers

S. Jagare, Phys. Lett. 32B, 571 (1970).

Nuclear Reactions: 239 , 240 , 241 , 242 Pu($p, 2n$), E=10.9-13.5 MeV; calculated σ for SF-isomer production. 238 , 239 , 240 , 241 , 242 Am calculated SF-isomer excitation energies.

70KrZT

Report: IN-1407 P151

Radioactivity: 241 Pu; measured activity; deduced no SF-isomer.

70Ot02 Fragment Angular Distributions from Neutron-Induced Fission of 242 Pu

K. Otozai, J. W. Meadows, A. N. Behkami, J. R. Huizenga, Nucl. Phys. A144, 502 (1970).

Nuclear Reactions: Fission 242 Pu(n, F), En=500, 620, 730, 990, 1230 keV; measured $\sigma(En, \theta)$ (fragment)). 243 Pu deduced information on transition states.

70PaZU

Report: CEA-N-1339, D Paya, 7/12/71

Nuclear Reactions: Fission $^{237}\text{Np}(\text{n},\text{F})$, E not given; measured(fragment)(fragment)-coin, (fragment)(fragment)-delay. ^{238}Np deduced no SF-isomer.

70Po01 Spontaneously Fissioning Isomers in U, Pu, Am and Cm Isotopes

S. M. Polikanov, G. Sletten, Nucl. Phys. A151, 656 (1970).

Nuclear Reactions: $^{233}\text{U}(\text{d,p})$, $^{238}\text{U}(\text{d,pn})$, $^{237}\text{Np}(\text{d,2n})$, 238 , $^{244}\text{Pu}(\text{p,2n})$, 238 , $^{240}\text{Pu}(\text{d, p})$, 239 , $^{241}\text{Pu}(\text{d,pn})$, 241 , $^{243}\text{Am}(\text{p, 2n})$, 241 , $^{243}\text{Am}(\text{d,2n})$, $^{243}\text{Am}(\text{d,pn})$; E=9-14.2 MeV; measured $\sigma(E)$ delayed fission. $^{239}\text{Pu}(\text{p, 2n})$, E=12.1-14.0 MeV; measured $\sigma(E)$; deduced threshold. Enriched targets.

Radioactivity: Fission 237m , 239m , 240m , 241m , $^{243m}\text{Pu}(\text{SF})$, 243m , $^{237m}\text{Am}(\text{SF})$, 240m , 241m , 242m , $^{243m}\text{Cm}(\text{SF})$, $^{234m}\text{U}(\text{SF})$; measured $T_{1/2}$. ^{239}Np deduced misassignment of (SF) isomer. $^{238m}\text{U}(\text{SF})$ deduced $T_{1/2}$.

70Re05 Evidence for a Direct Reaction Mechanism in the Production of Fission Isomers

R. Repnow, V. Metag, J. D. Fox, P. von Brentano, Nucl. Phys. A147, 183 (1970).

Nuclear Reactions: $^{235}\text{U}(\text{d,p})$, E=13-20 MeV; measured σ delayed fission. Enriched target. $^{236}\text{U}(\text{d,pn})$, E=11-20 MeV; measured σ delayed fission. Enriched target. $^{238}\text{U}(\text{d,pn})$, E=11-20 MeV; measured σ delayed fission. Natural target. $^{233}\text{U}(\text{d,X})$, $^{236}\text{U}(\text{p, X})$, E = 14, 20 MeV; E upper limits σ delayed fission. Enriched targets. $^{239}\text{U}(\text{p,X})$, E=14-20 MeV; measured upper limits σ delayed fission. Natural target.

Radioactivity: Fission 236 , ^{238}U deduced $T_{1/2}$ (SF-isomer). 234 , ^{237}U deduced no SF-isomer.

70So06 Intermediate Structure Effects in the Fission of Some Actinide Nuclei

D. K. Sood, N. Sarma, Nucl. Phys. A151, 532 (1970).

Nuclear Reactions: Fission 233 , ^{235}U , 239 , 240 , ^{241}Pu , $^{242}\text{Am}(\text{n,F})$, E < 1 MeV; measured nothing; analyzed $\sigma(E)$ data; deduced spacing of second minimum levels.

70Vi05 Izomeri Spontan Fisionabili Ai Nucleelor Transuraniene

N. Vilcov, Stud. Cercet. Fiz. 22, 795 (1970).

Radioactivity: Fission ^{236}U , ^{238}Np , 236 , 240 , 241 , 242 , ^{243}Pu , 239 , $^{241}\text{Am}(\text{SF})$; measured $T_{1/2}$.

70Wo06 Spontaneous Fission Isomerism in Uranium Isotopes

K. L. Wolf, R. Vandenbosch, P. A. Russo, M. K. Mehta, C. R. Rudy, Phys. Rev. C1, 2096 (1970).

Radioactivity: Fission ^{236m}U , $^{238m}\text{U}(\text{SF})$; measured $T_{1/2}$.

Nuclear Reactions: 236 , $^{238}\text{U}(\text{d,X})$, (d,pn) , E=13-22 MeV; measured $\sigma(E;\text{Ep})$. 236 , ^{238}U deduced isomer ratios.

71Au06 Neutron-Induced Fission Cross Sections of ^{242}Pu and ^{244}Pu

G. F. Auchampaugh, J. A. Farrell, D. W. Bergen, Nucl. Phys. A171, 31 (1971).

Nuclear Reactions: Fission 242 , $^{244}\text{Pu}(\text{n,F})$, E=20 eV-10 MeV; measured $\sigma(E)$. 243 , ^{244}Pu deduced level spacings, resonance parameters, second barrier widths.

71Ba30 Fission of U, Np, Pu and Am Isotopes Excited in the (d,p) Reaction

B. B. Back, J. P. Bondorf, G. A. Otroschenko, J. Pedersen, B. Rasmussen, Nucl. Phys. A165, 449 (1971).

Nuclear Reactions: Fission 233 , ^{235}U , ^{237}Np , 238 , 239 , ^{241}Pu , 241 , $^{243}\text{Am}(\text{d,pF})$, E=13.0 MeV; measured $\sigma(\text{Ep,E(fragment)})$. 234 , ^{236}U , ^{238}Np , 239 , 240 , ^{242}Pu , 242 , ^{244}Am deduced fission probability, fission barrier heights, transparencies.

71Be12 Neutron-Induced Fission Cross Section of ^{242}Pu

D. W. Bergen, R. R. Fullwood, Nucl. Phys. A163, 577 (1971).

Nuclear Reactions: Fission $^{242}\text{Pu}(\text{n,F})$, E=50 eV-5 keV, 0.1-3 MeV; measured $\sigma(E)$. ^{243}Pu deduced resonances, F-width, fission barrier.

71Be62 Production of the Spontaneously Fissioning U^{236} Isomer in Thermal Neutron Radiative Capture

A. G. Belov, Y. P. Gangskii, B. Dalkhsuren, A. M. Kucher, Yad. Fiz. 14, 685 (1971); Sov. J. Nucl. Phys. 14, 385 (1972).

Nuclear Reactions: Fission $^{235}\text{U}(\text{n,f})$, E=thermal; measured σ , (fragment)(ce)-delay. ^{236m}U deduced $T_{1/2}$.

71Bo61 Study of the ^{236}mf -U Isomeric Fission Through the $^{235}\text{U}(\text{n,f})$ Reaction in the Energy Range 0.25 - 4 MeV

I. Boca, M. Sezon, I. Vilcov, N. Vilcov, Rev. Roum. Phys. 16, 473 (1971).

Radioactivity: $^{236m}\text{U}(\text{SF})$; measured $T_{1/2}$.

Nuclear Reactions: Fission $^{235}\text{U}(\text{n,f})$, E=0.25-4 MeV; measured $\sigma(E)$ for $^{236m}\text{U}(\text{SF})$ production.

71Br38 Population of Fission Isomers in ^{236}U by the (d,p) Reaction

H. C. Britt, B. H. Erkkila, Phys. Rev. C4, 1441 (1971).

Nuclear Reactions: Fission $^{235}\text{U}(\text{d,pf})$, (d,p) , E=12 MeV; measured σ ratios, $\sigma(\text{Ep})$, $(\text{d})(\text{fragment})$ -delay. ^{236m}U deduced $T_{1/2}$.

Radioactivity: Fission ^{236m}U ; measured $T_{1/2}$.

71Br39 Systematics of Spontaneously Fissioning Isomers

H. C. Britt, S. C. Burnett, B. H. Erkkila, J. E. Lynn, W. E. Stein, Phys. Rev. C4, 1444 (1971).

Radioactivity: Fission 235m , 237m , 238m , 239m , 240m , $^{241m}\text{Pu}(\text{SF})$, 241m , 242m , 243m , 244m , $^{245m}\text{Cm}(\text{SF})$, $^{236m}\text{U}(\text{SF})$, 239m , 240m , 242m , 243m , $^{244m}\text{Am}(\text{SF})$; measured $T_{1/2}$, lower limits.

71Ga19 Excitation of the Spontaneously Fissioning Isomeric States of ^{239}Pu and ^{243}Am at Inelastic γ -Quantum Scattering

Y. P. Gangrsky, B. N. Markov, I. F. Kharisov, Y. M. Tsipenyuk, JINR-P15-5959 (1971).

Nuclear Reactions: ^{239}Pu , $^{243}\text{Am}(\gamma, \gamma')$, E=7-11 MeV; measured $\sigma(E; E\gamma)$, (γ) (fragment)-delay. ^{239}Pu , ^{243}Am deduced SF-isomer excitation. ^{239m}Pu deduced $T_{1/2}$.

71Ga35 Spontaneously Fissioning Isomers of Uranium, Plutonium, and Americium from Neutron Reactions

Y. P. Gangrskii, T. Nad, I. Vinnai, I. Kovach, At. Energ. 31, 156 (1971); Sov. At. Energy 31, 874 (1972).

Nuclear Reactions: ^{232}Th , 235 , ^{238}U , ^{239}Pu , $^{243}\text{Am}(n, 2n)$, E=14.7 MeV; ^{238}U , ^{242}Pu , $^{243}\text{Am}(n, n')$, E=2-7 MeV; measured σ (SF isomers). ^{231}Th , 234 , ^{237}U , ^{238}Pu deduced no SF-isomer yield. ^{238}U , ^{242}U , ^{243}Am deduced SF isomer yield.

71Ga39 Excitation of Spontaneously Fissioning Isomer States ^{239}Pu and ^{243}Am in Inelastic Scattering of γ Quanta

Y. P. Gangrskii, B. N. Markov, I. F. Kharisov, Y. M. Tsipenyuk, Pisma Zh. Eksp. Teor. Fiz. 14, 370 (1971); JETP Lett. (USSR) 14, 249 (1971).

Nuclear Reactions: Fission ^{239}Pu , $^{243}\text{Am}(\gamma, \gamma'F)$, E < 11 MeV; measured (γ) (fragment)-delay. ^{239m}Pu deduced $T_{1/2}$. ^{239}Pu , ^{243}Am deduced isomer yields.

71MaZE

Thesis: , Univ Kansas,D E Maharry,DABBB 32B 5981,5/5/72

Nuclear Reactions: $^{92}\text{Mo}(d,p\gamma)$, measured $\sigma(E_p, E\gamma)$. ^{93}Mo deduced levels.

Nuclear Structure: A=230-256; ^{236}U ; calculated fission barriers, shape isomer excitation energies, equilibrium deformations, total energy surfaces.

71Me03 Correlation between Fission Isomer Half-Lives and Liquid-Drop Model Parameters

V. Metag, R. Repnow, P. von Brentano, Nucl. Phys. A165, 289 (1971).

71Mo11 Analysis of the Fission and Capture Cross Sections of the Curium Isotopes

M. S. Moore, G. A. Keyworth, Phys. Rev. C3, 1656 (1971).

Nuclear Reactions: 244 , 245 , 246 , 247 , $^{248}\text{Cm}(n, F)$, 244 , $^{246}\text{Cm}(n, \gamma)$, E=20 eV-3 MeV; measured $\sigma(E)$. 245 , 246 , 247 , 248 , ^{249}Cm deduced resonances, level-width.

71Na26 Investigations of the Radiative Capture of Fast Neutrons Producing the Spontaneously Decaying Isomers ^{242}Am and ^{244}Am

T. Nagy, A. G. Belov, Y. P. Gangrsky, B. N. Markov, I. V. Sizov, I. F. Harisov, Acta Phys. 30, 293 (1971).

Nuclear Reactions: 241 , $^{243}\text{Am}(n, \gamma)$, E < 16 MeV; measured σ ratios for 242 , ^{244}Am SF-isomer production.

71Pa33 Fission Threshold Energies in the Actinide Region

H. C. Pauli, T. Ledergerber, Nucl. Phys. A175, 545 (1971).

Nuclear Structure: Fission 232 , ^{234}Th , 234 , 236 , 238 , ^{240}U , 236 , 238 , 240 , 242 , ^{244}Pu ; calculated liquid-drop barriers, first, second saddle point energies.

71Re11 Fission Isomers in Cm and Bk Isotopes

R. Repnow, V. Metag, P. von Brentano, Z. Phys. 243, 418 (1971).

Radioactivity: Fission ^{245m}Bk , ^{242m}Cm , ^{241m}Cm , ^{243m}Am , ^{243m}Cm , ^{237m}Pu ; measured $T_{1/2}$. $^{243}\text{Am}(\alpha, 2n\gamma)$, E=26 MeV; $^{243}\text{Am}(p, 2n\gamma)$, $(p, 3n\gamma)$, E=14, 20 MeV; $^{243}\text{Am}(d, pn)$, $(d, 2n\gamma)$, E=13-20 MeV; $^{237}\text{Np}(d, 2n\gamma)$, E=12-18 MeV; measured delays, $\sigma(E)$.

71Ru03 Spin Isomers of the Shape Isomer ^{237m}Pu

P. A. Russo, R. Vandebosch, M. Mehta, J. R. Tesmer, K. L. Wolf, Phys. Rev. C3, 1595 (1971).

Radioactivity: Fission ^{237m}Pu (SF); measured $T_{1/2}$; deduced shape isomerism.

71Ta17 Search for Bremsstrahlung-Induced Fission Isomers of ^{238}U and ^{239}Pu

B. Tamain, B. Pfeiffer, H. Wollnik, E. Konecny, Nucl. Phys. A173, 465 (1971).

Radioactivity: Fission ^{238}U , ^{239}Pu (SF); measured $T_{1/2}$.

Nuclear Reactions: Fission ^{238}U , $^{239}\text{Pu}(\gamma, F)$, E < 53 MeV; measured (γ) (fragment)-delay. ^{238}U , ^{239}Pu deduced fission isomers, $T_{1/2}$.

71Te07 Spontaneously Fissioning Isomers in ^{237}Pu

J. K. Temperley, J. A. Morrissey, S. L. Bacharach, Nucl. Phys. A175, 433 (1971).

Radioactivity: Fission ^{237m}Pu (SF) [from $^{237}\text{Np}(d, 2n)$]; measured $T_{1/2}$, E(fragment). $^{237}\text{Np}(d, 2n)$, E=8.5-14.5 MeV; measured delayed, prompt fission σ ratios, (d)(fission-fragment)-delay; E=13.0 MeV, measured E(fragment).

72Bo48 Search for Spontaneously Fissioning Isomers Produced with 600 MeV Protons

A. H. Boos, R. Brandt, D. Molzahn, D. M. Montgomery, J. Inorg. Nucl. Chem. 34, 3309 (1972).

Nuclear Reactions: U, Th, Bi, Pb(p,X), E=600 MeV; measured fission activities; deduced σ for SF-isomer production.

72Br04 Investigation of γ -Ray Emission Preceding Isomeric Fission of ^{236}U

J. C. Browne, C. D. Bowman, Phys. Rev. Lett. 28, 617 (1972).

Nuclear Reactions: Fission $^{235}\text{U}(n, \gamma F)$, E=1-100 eV; measured (γ) (fragment)-delay; deduced limit on pre-fission γ -emission. ^{236}U deduced relative double barrier penetrabilities.

72Br35 Excitation Functions for the Production of Fission Isomers in Various Am Isotopes

H. C. Britt, B. H. Erkkila, B. B. Back, Phys. Rev. C6, 1090 (1972).

Radioactivity: Fission ^{239m}Am , ^{245m}Am ; measured $T_{1/2}$.

Nuclear Reactions: $^{239}, ^{240}, ^{242}$, $^{244}\text{Pu}(p,2n)$, $(t,2n)$, $(t,3n)$, $E=10\text{-}16 \text{ MeV}$; measured σ for SF-isomer production.

72Ga04 Measurement of the Excitation Energy of the Spontaneously Fissioning Isomer $\text{Pu}^{239}(\gamma,n)$

Y. P. Gangrskii, V. N. Maykov, I. F. Kharisov, Y. M. Tsvipenyuk, Yad. Fiz. 16, 271 (1972); Sov. J. Nucl. Phys. 16, 151 (1973).

Nuclear Reactions: $^{240}\text{Pu}(\gamma,n)$, $E < 15 \text{ MeV}$; measured $\sigma(E)$ for ^{239m}Pu (SF) production. ^{239m}Pu (SF) deduced excitation energy.

72Ga42 Production of Spontaneously Fissioning Isomers with Nanosecond Lifetimes in α -Particle Reactions

Y. P. Gangrskii, Nguen Kong Khan, D. D. Pulatov, At. Energ. 33, 829 (1972); Sov. At. Energy 33, 948 (1973).

Nuclear Reactions: $^{233}, ^{235}, ^{238}\text{U}$, $^{239}, ^{242}\text{Pu}$, $^{241}, ^{243}\text{Am}(\alpha,xn)$, $E=20\text{-}36 \text{ MeV}$; measured $\sigma(E)$ for SF-isomers. $^{235}, ^{237}, ^{240}\text{Pu}$, $^{241}, ^{243}\text{Cm}$, $^{242}, ^{243}, ^{244}, ^{245}\text{Bk}$ deduced SF-isomers, $T_{1/2}$.

72Ho11 Total Spontaneous and Isomer Fission Half-Lives of ^{234}U , ^{236}U and ^{240}Pu

M. A. Hooshyar, F. B. Malik, Phys. Lett. 38B, 495 (1972).

Nuclear Structure: Fission $^{234}, ^{236}\text{U}$, ^{240}Pu (SF); calculated total $T_{1/2}$, $T_{1/2}$ (SF), average fragment kinetic energies. Coupled-channel decay theory.

72Ho48 A Coupled Channel Approach to the Isomer Fission State

M. A. Hooshyar, F. B. Malik, Helv. Phys. Acta 45, 567 (1972).

72HoXQ Suche nach γ -Übergangen im Spaltungs-Isomer ^{236}U

F. Horsch, E. Konecny, K. E. G. Lobner, H. J. Specht, Univ., Tech. Univ. Munchen, Jahresbericht 1972, p. 104 (1973).

Nuclear Reactions: $^{235}\text{U}(n,\gamma)$; measured E_γ , I_γ . ^{236}U deduced isomer.

72Ka59 Search for γ -Branch in the ^{236m}U Fission Isomer Decay

E. Kashy, J. Hattula, J. Borggreen, V. Maarbjerg, Comment. Phys.-Math. 42, 266 (1972).

Radioactivity: ^{236m}U ; measured upper limit for γ -ray decay.

72Ko10 Search for Conversion Electrons Populating the ^{236}U Fission Isomer

E. Konecny, H. J. Specht, J. Weber, H. Weigmann, R. L. Ferguson, P. Osterman, M. Waldschmidt, G. Siegert, Nucl. Phys. A187, 426 (1972).

Nuclear Reactions: Fission $^{235}\text{U}(n,\gamma)$, $E=\text{thermal}$; measured (fragment)(ce)-coin, -delay; deduced upper limit for isomeric/prompt fission ratio.

72Ku26 Search for Fission Isomers in the Radium Region

I. M. Kuks, V. I. Matvienko, Y. A. Nemilov, Y. A. Selitskii, V. B. Funstein, Yad. Fiz. 16, 438 (1972); Sov. J. Nucl. Phys. 16, 244 (1973).

Nuclear Reactions: $^{226}\text{Ra}(d,X)$, $E=6.6, 11.3 \text{ MeV}$; $^{226}\text{Ra}(n,X)$, $E=0.7\text{-}10, 14.5 \text{ MeV}$; measured $\sigma(F)$. $^{224}, ^{225}, ^{226}, ^{227}\text{Ra}$, $^{225}, ^{226}, ^{227}\text{Ac}$ deduced no SF-isomer.

72La05 Fission Barriers and the Inclusion of Axial Asymmetry

S. E. Larson, I. Ragnarsson, S. G. Nilsson, Phys. Lett. 38B, 269 (1972).

Nuclear Structure: Fission $^{186}, ^{188}, ^{190}, ^{192}\text{W}$, ^{196}Pt , ^{196}Hg , $^{196}, ^{204}\text{Pb}$, ^{242}Pu , ^{246}Cm , ^{252}Fm , $^{258}\text{104}$, superheavy; calculated potential energy surfaces vs deformation parameters, fission barriers. Modified oscillator model, axial symmetry.

72Ma11 A Single-Particle Model Calculation of Total Energy Surfaces in Heavy Nuclei

D. E. Maharry, J. P. Davidson, Nucl. Phys. A183, 371 (1972).

Nuclear Structure: Fission ^{236}U , $^{230}, ^{232}\text{Th}$, $^{234}, ^{236}, ^{238}\text{U}$, $^{246}, ^{248}, ^{250}, ^{252}\text{Cf}$, $^{238}, ^{240}, ^{242}, ^{244}\text{Am}$, $^{236}, ^{238}, ^{240}, ^{242}, ^{244}\text{Pu}$, $^{240}, ^{242}, ^{244}, ^{246}, ^{248}, ^{250}\text{Cm}$; calculated total energy surfaces, fission barriers. Single-particle model.

72Mo27 Odd-Multipole Shape Distortions and the Fission Barriers of Elements in the Region $84 < Z < 120$

P. Moller, Nucl. Phys. A192, 529 (1972).

Nuclear Structure: Fission $Z=84\text{-}120$; ^{210}Po , ^{236}U , ^{256}Fm , ^{252}Fm ; calculated potential energy surfaces, fission barriers.

72NaYU

Thesis: T Nagy, Dubna

Nuclear Reactions: $^{241}, ^{243}\text{Am}(n,\gamma)$, $E=0.8\text{-}16 \text{ MeV}$; ^{235}U , $^{239}\text{Pu}(n,\gamma)$, $E=\text{th}$; ^{238}U , $^{239}, ^{242}\text{Pu}$, $^{243}\text{Am}(n,n')$, $E=3\text{-}7 \text{ MeV}$, 14.7 MeV ; $^{240}, ^{242}\text{Pu}$, $^{243}\text{Am}(n,2n)$, $E=14.7 \text{ MeV}$; measured $\sigma(E)$ for SF isomers. ^{232}Th , $^{233}, ^{235}\text{U}$, ^{237}Np , $^{239}\text{Pu}(n,2n)$, $E=14.7 \text{ MeV}$; measured no SF isomer.

72Pe01 An Investigation of the Population of the Shape Isomer ^{236m}U Through the (d,p) Reaction

J. Pedersen, B. Rasmussen, Nucl. Phys. A178, 449 (1972).

Nuclear Reactions: $^{235}\text{U}(d,pF)$, $E=11 \text{ MeV}$; measured (p) (fragment)-delay. ^{236m}U deduced $T_{1/2}$, fission barrier parameters.

72PiZR Fission Isomer in Uranium-236

J. V. Pilcher, F. D. Brooks, W. R. McMurray, INDC(SEC)-28/L, p. 249 (1972).

Radioactivity: Fission ^{236m}U (SF); measured $T_{1/2}$.

72Sp06 Identification of a Rotational Band in the ^{240}Pu Fission Isomer

H. J. Specht, J. Weber, E. Konecny, D. Heunemann, Phys. Lett. 41B, 43 (1972).

Radioactivity: Fission ^{240}Pu (SF) [from $^{238}\text{U}(\alpha,2n\gamma)$; $E=25 \text{ MeV}$]; measured $I(\text{ce})$, (α) (fragment)-delay, $E(\text{ce})$. ^{240m}Pu deduced levels, rotational band structure.

72Va08 Spontaneous-Fission-Isomer Excitation Energies from Threshold Measurements

R. Vandenbosch, Phys. Rev. C5, 1428 (1972).

72Va44 Searches for the Spontaneously Fissioning Isomer Pu^{240m} in the Thermal-Neutron Capture Reaction

G. V. Valskii, O. M. Mrachkovskii, G. A. Petrov, Y. S. Pleva, Yad. Fiz. 16, 667 (1972); Sov. J. Nucl. Phys. 16, 374 (1973).

Nuclear Reactions: Fission $^{239}Pu(n,F)$; E=thermal; measured σ production for ^{240m}Pu .

72Vi10 $^{233m}Pu(f)$ Double Fission Isomer Study Through the $^{237}Np(d,2n)$ Reaction in the E = 9-12 MeV Energy Range

N. Vilcov, G. Griffith, I. Vilcov, R. B. Leachman, Rev. Roum. Phys. 17, 1031 (1972).

Nuclear Reactions: $^{237}Np(d,2n)$, E=9.1-12.1 MeV; measured $\sigma(E)$ ratio for two isomers. ^{237}Pu deduced levels, $T_{1/2}$.

72Vy07 Excitation Energies of the Spontaneously Fissile Isomers of Pu^{240} , Cm^{241} , and Bk^{243} in Reactions with α -Particles

I. Vylkov, N. Vylkov, Y. P. Gangrskii, M. Marinescu, A. A. Pleve, D. Poenaru, I. F. Kharisov, Yad. Fiz. 16, 454 (1972); Sov. J. Nucl. Phys. 16, 253 (1973).

Nuclear Reactions: ^{238}U , ^{239}Pu , $^{241}Am(\alpha,2n)$, E=20-26 MeV; measured σ for SF-isomer production. ^{240}Pu , ^{241}Cm , ^{243}Bk deduced SF isomer excitation energies.

72We09 Evaluation of Fission Barrier Parameters from Near-Barrier Fission and Isomeric Half-Life Data

H. Weigmann, J. P. Theobald, Nucl. Phys. A187, 305 (1972).

Nuclear Structure: Fission 234 , 235 , 236 , 237 , 238 , ^{239}U , 237 , 238 , 239 , ^{240}Np , 235 , 236 , 237 , 238 , 239 , ^{240}Am , 241 , 242 , 243 , 244 , ^{245}Pu , 237 , 238 , 239 , 240 , 241 , 242 , 243 , ^{244}Am , 241 , 242 , 243 , ^{244}Cm , 246 , ^{247}Cm , 244 , 245 , ^{246}Bk ; calculated fission barriers, $T_{1/2}$.

72Wo07 Fissioning Isomers of Americium, Curium and Berkelium Isotopes

K. L. Wolf, J. P. Unik, Phys. Lett. 38B, 405 (1972).

Radioactivity: Fission ^{240m}Am , ^{243m}Am , ^{245m}Am , ^{246m}Am , ^{244m}Am , ^{239m}Pu , ^{242m}Bk , ^{244m}Bk , ^{243m}Cm , ^{245m}Cm ; measured $T_{1/2}$.

Nuclear Reactions: 242 , ^{244}Pu , 241 , $^{243}Am(\alpha,xF)$, E=25-46 MeV; measured $\sigma(E)$ for SF-isomer production.

73Al08 A New Two-Center Shell Model for Nuclear Fission

K. Albrecht, Nucl. Phys. A207, 225 (1973).

Nuclear Structure: ^{226}Ra , ^{232}Th , 236 , ^{238}U , 240 , ^{242}Pu , ^{244}Cm , 248 , ^{252}Cf , ^{252}Fm ; calculated deformation energies, isomer energies.

73Ba19 Fission and Decay of Excited Nuclei

V. S. Barashenkov, A. S. Iljinov, V. D. Toneev, F. G. Gereghi, Nucl. Phys. A206, 131 (1973).

Nuclear Structure: ^{140}Eu , ^{157}Ho , ^{175}Ta , ^{186}Os , ^{187}Os , ^{188}Os , ^{185}Ir , ^{189}Ir , ^{191}Ir , ^{194}Hg , ^{198}Hg , ^{210}Po , ^{211}Po , ^{212}Po , ^{213}At , ^{227}Ra , ^{233}U , ^{234}U , ^{235}U , ^{236}U , ^{237}U , ^{238}U , ^{239}U , ^{237}Np , ^{238}Np , ^{241}Am , ^{242}Am , ^{244}Am , ^{240}Cm , ^{242}Cm , ^{246}Cm , ^{250}Cm , ^{246}Cf , ^{248}Cf , ^{250}Cf , ^{252}Cf , ^{251}No , ^{252}No , ^{253}No , ^{254}No , ^{255}No , ^{256}No , ^{257}No , ^{259}No ; calculated fission barrier, level-width(n)/level-width(F).

73Be04 Production of Spontaneously Fissioning Isomers in Th, U, Np, Pu and Am Isotopes in Reactions Induced by 14.7 MeV Neutrons

A. G. Belov, Y. P. Gangrsky, B. Dalkhsuren, A. M. Kucher, T. Nagy, D. M. Nadkarni, Indian J. Phys. 47, 232 (1973).

Nuclear Reactions: ^{232}Th , 235 , ^{238}U , ^{237}Np , 240 , ^{242}Pu , 241 , $^{243}Am(n,2n)$, ^{239}Pu , 241 , $^{243}Am(n,n')$, E=14.7 MeV; measured production σ for SF isomers, nF(t). 239m , ^{241m}Pu , 240m , 241m , 242m , ^{243m}Am deduced $T_{1/2}$.

73Be05 Search for α Emission in the Decay of Spontaneously Fissionable Isomers

A. G. Belov, Y. P. Gangrskii, B. Dalkhsuren, A. M. Kucher, Nguen Kong Khan, Yad. Fiz. 17, 942 (1973); Sov. J. Nucl. Phys. 17, 493 (1974).

Radioactivity: Fission 240m , ^{242m}Am , $^{241m}Pu(SF)$; measured E α , I α . Deduced no α -emission.

73Be10 Search for γ -Rays Emitted in the Formation of a Fission Isomer

D. Benson, Jr., C. M. Lederer, E. Cheifetz, Nucl. Phys. A201, 445 (1973).

Nuclear Reactions: $^{238}U(\alpha,\gamma F)$; measured $\alpha f(t)$, $\gamma f(t)$, E γ . $^{240m}Pu(SF)$; deduced limits on pre-fission γ -ray photons.

73Br04 Fission Barriers Deduced from the Analysis of Fission Isomer Results

H. C. Britt, M. Bolsterli, J. R. Nix, J. L. Norton, Phys. Rev. C7, 801 (1973).

73Br38 Properties of Fission Isomers

H. C. Britt, At. Data Nucl. Data Tables 12, 407 (1973).

73BrWU

Report: USNDc-7 P106

Nuclear Reactions: $^{243}Pu(n,F)$; measured $\sigma(E\gamma)$. ^{243}Pu deduced fission isomer.

73Fl03 Excitation Functions for Spallation Products and Fission Isomers in $^{237}Np(^4He,xn)^{241-x}Am$ Reactions

A. Fleury, F. H. Ruddy, M. N. Namboodiri, J. M. Alexander, Phys. Rev. C7, 1231 (1973).

Nuclear Reactions: $^{237}Np(\alpha,2n)$, $(\alpha,3n)$, $(\alpha,4n)$, E=19-45 MeV; measured $\sigma(E)$, σ , isomer σ ratio. ^{239m}Am deduced $T_{1/2}$.

- 73HeYN** Search for Conversion Electrons from the Decay of Excited States in the Secondary Minimum of ^{236}U
R. Heffner, J. Pedersen, P. A. Russo, H. Swanson, RLO-1388-221, p. 123 (1973).
Radioactivity: ^{236}U ; measured I(ce).
- 73Kh06** Angular Distribution of Fragments of Spontaneously Fissioning Isomers
Fam Zui Khien, Yad. Fiz. 17, 489 (1973); Sov. J. Nucl. Phys. 17, 251 (1974).
Nuclear Reactions: $^{235}\text{U}(\alpha, 3n)$; calculated ^{236}Pu fission isomer angular distribution.
- 73Li01** A Subnanosecond and a Nanosecond Fission Isomer in ^{238}Pu
P. Limkilde, G. Sletten, Nucl. Phys. A199, 504 (1973).
Radioactivity: Fission ^{238m}Pu , ^{240m}Pu ; measured $T_{1/2}$.
Nuclear Reactions: $^{236}\text{U}(\alpha, 2n)$, E=21.0-27.0 MeV; measured $\sigma(1)(E)$, $\sigma(2)(E)$ delayed fission; deduced thresholds; $^{236}\text{U}(\alpha, F)$, E=20.0-28.0 MeV; measured $\sigma(E)$ prompt fission; $^{238}\text{U}(\alpha, 2n)$, E approx 25 MeV; measured σ delayed fission.
- 73Me23** Neutron-Fission Competition Near Threshold; The Influence of Shells and Pairing on the Decay of the ^{241}Cm Compound Nucleus
V. Metag, S. M. Lee, E. Liukkonen, G. Sletten, S. Bjornholm, A. S. Jensen, Nucl. Phys. A213, 397 (1973).
Nuclear Reactions: $^{238}\text{Pu}(\alpha, n)$, E=19.9-23 MeV; $^{238}\text{Pu}(\alpha, 2n)$, E=19.9-27 MeV; $^{241}\text{Am}(p, 2n)$, E=8.2-16 MeV; measured $\sigma(^{241}\text{Cm})$, $\sigma(^{240}\text{Cm})$, $\sigma(\text{fission})$. $^{241}, ^{242}\text{Cm}$ deduced n-width, F-width.
- 73Na03** Excitation Functions for the Fission Isomers ^{240m}Pu and ^{239m}Pu from $^{238}\text{U}(^4\text{He}, xn)$ Reactions
M. N. Namboodiri, F. H. Ruddy, J. M. Alexander, Phys. Rev. C7, 1222 (1973).
Nuclear Reactions: $^{238}\text{U}(\alpha, 2n)$, $(\alpha, 3n)$, E < 28 MeV; measured $\sigma(E)$, $\sigma(^{240m}\text{Pu}$ deduced $T_{1/2}$.
- 73Na35** Neutronokkal Letrehozott, Izomer Allapotbol Spontan Hasado Magok Keletkezesere Vezeto Reakciok Vizsgalata
T. Nagy, Magy. Fiz. Foly. 21, 555 (1973).
Radioactivity: Fission ^{238}U , $^{239}, ^{241}, ^{242}\text{Pu}$, $^{242}, ^{243}, ^{244}\text{Am}$, $^{236}\text{Np(SF)}$; measured $T_{1/2}$, ^{238}Pu , $^{232}, ^{234}, ^{237}\text{U}$, ^{231}Th measured $T_{1/2}$ limits.
Nuclear Reactions: $^{241}, ^{243}\text{Am}(n, \gamma)$, E=0.8-16 mev; $^{233}, ^{235}, ^{238}\text{U}$, $^{239}\text{Pu}(n, \gamma)$, E=thermal; ^{238}U , $^{239}, ^{242}\text{Pu}$, $^{243}\text{Am}(n, n')$, E=3-7, 14.7 MeV; ^{237}Np , $^{233}, ^{235}, ^{238}\text{U}$, ^{232}Th , $^{239}, ^{240}, ^{242}\text{Pu}$, $^{243}\text{Am}(n, 2n)$, E=14.7 MeV; measured $\sigma(E)$ for production of SF isomers.
- 73OtZX**
Report: RCN-203 P169
Nuclear Reactions: $^{235}\text{U}(n, F)$, E=1 MeV; measured fission isomer yield.
- 73Po05** Spontaneously Fissioning Isomer 236m Excited by Capture of Thermal Neutrons
L. A. Popeko, G. A. Petrov, E. F. Kochubei, T. K. Zvezdkina, Yad. Fiz. 17, 234 (1973); Sov. J. Nucl. Phys. 17, 120 (1974).
Nuclear Reactions: $^{235}\text{U}(n, F)$, E=thermal; measured (fragment)(ce), γ, X -delay. ^{236m}U deduced yield.
- 73Po08** Neutron Resonance Parameters of ^{242}Pu
F. Poortmans, G. Rohr, J. P. Theobald, H. Weigmann, G. J. Vanpraet, Nucl. Phys. A207, 342 (1973).
Nuclear Reactions: $^{242}\text{Pu}(n, n)$, $^{242}\text{Pu}(n, \gamma)$, E=20-1300 eV; measured $\sigma(^{243}\text{Pu}$ resonances deduced resonance parameters n-width, γ -width. Enriched target.
- 73PoZA** Fission Isomers, Eleven Years of Experimental Work
D. N. Poenaru, IFA-CRD-54-1973 (1973).
Compilation: Fission $^{234m}, ^{235m}, ^{236m}, ^{238m}\text{U(SF)}$, $^{237m}\text{Np(SF)}$, $^{235m}, ^{236m}, ^{237m}, ^{238m}, ^{239m}$, $^{240m}, ^{241m}, ^{242m}, ^{243m}$, $^{244m}, ^{245m}\text{Am}$, $^{240m}, ^{241m}, ^{242m}, ^{243m}, ^{245m}\text{Cm}$, $^{242m}, ^{243m}, ^{244m}, ^{245m}\text{Bk}$; compiled experimental $T_{1/2}$.
- 73Sp04** Statistical Theory of Isomer Ratios for Shape (Fission) Isomers in (n, γ) Reactions
D. Sperber, Nuovo Cim. 13A, 373 (1973).
Nuclear Reactions: $^{233}, ^{235}\text{U}$, ^{239}Pu , $^{241}\text{Am}(n, \gamma)$; calculated isomer ratios.
- 73Va16** Relative Excitations of the ^{237}Pu Shape Isomers
R. Vandebosch, P. A. Russo, G. Sletten, M. Mehta, Phys. Rev. C8, 1080 (1973).
Radioactivity: Fission $^{237m}\text{Pu(SF)}$, ^{237}Pu ; measured delayed yields. ^{237}Pu deduced levels, J, π , $T_{1/2}$.
- 73Va30** Probability of Formation of Spontaneously Fissioning Isomer States Following Thermal Neutron Capture by 235 and Pu^{239}
G. V. Valskii, O. M. Mrachkovskii, G. A. Petrov, Y. S. Pleva, Yad. Fiz. 18, 492 (1973); Sov. J. Nucl. Phys. 18, 253 (1974).
Nuclear Reactions: ^{235}U , $^{239}\text{Pu}(n, \gamma)$; measured $\sigma(\text{isomer})$.
- 73Wo03** The Fissioning Isomer ^{237m}Np
K. L. Wolf, J. P. Unik, Phys. Lett. 43B, 25 (1973).
Radioactivity: Fission $^{237m}\text{Np(SF)}$; measured $T_{1/2}$, excitation energy.
- 73Ze05** Search for a Spontaneously Fissioning Isomer Nucleus U^{236m} in the Reaction $U^{235}(n, \gamma)$
Zen Chang Bom, A. Lajtai, A. A. Omelyanenko, T. T. Panteleev, S. M. Polikanov, Y. V. Ryabov, Tang San Khak, Yad. Fiz. 18, 34 (1973); Sov. J. Nucl. Phys. 18, 18 (1974).
Nuclear Reactions: $^{235}\text{U}(n, \gamma)$, E approx 60 keV; measured σ for SF isomer. ^{236}U deduced no SF isomer.

74Ba73 *Fission of Odd-A and Doubly Odd Actinide Nuclei Induced by Direct Reactions*

B. B. Back, H. C. Britt, O. Hansen, B. Leroux, J. D. Garrett, Phys. Rev. C10, 1948 (1974).

Nuclear Reactions: $^{230}, ^{232}\text{Th}(\text{He},\alpha\text{F})$, $^{230}, ^{232}\text{Th}$, $^{233}, ^{234}, ^{235}, ^{236}, ^{238}\text{U}$, $^{239}, ^{240}, ^{242}\text{Pu}$, $^{248}\text{Cm}(\text{He},\text{dF})$, E=24 MeV; ^{230}Th , ^{231}Pa , ^{237}Np , $^{248}\text{Cm}(\text{d,pF})$, E=15 MeV; ^{243}Am , $^{239}\text{Pu}(\text{t},\text{pF})$, E=15 MeV; $^{248}\text{Cm}(\text{t},\alpha\text{F})$, E=16 MeV; measured fission probabilities. $^{229}, ^{231}\text{Th}$, $^{231}, ^{232}, ^{233}\text{Pa}$, $^{234}, ^{235}, ^{236}, ^{237}, ^{238}, ^{239}\text{Np}$, ^{241}Pu , $^{240}, ^{241}, ^{243}, ^{245}, ^{247}\text{Am}$, ^{249}Cm , ^{249}Bk deduced barrier heights.

74Ba28 *Fission of Doubly Even Actinide Nuclei Induced by Direct Reactions*

B. B. Back, O. Hansen, H. C. Britt, J. D. Garrett, Phys. Rev. C9, 1924 (1974).

Nuclear Reactions: $^{230}, ^{232}\text{Th}$, $^{234}, ^{236}, ^{238}\text{U}$, $^{238}, ^{240}, ^{242}\text{Pu}$, $^{248}\text{Cm}(\text{t,pF})$, E=15 MeV; $^{231}\text{Pa}(\text{t},\alpha\text{F})$, E=16 MeV; ^{231}Pa , ^{237}Np , $^{243}\text{Am}(\text{He},\text{dF})$, E=24 MeV; $^{233}\text{U}(\text{d,pF})$, E=13 MeV; $^{248}\text{Cm}(\text{p,pF})$, E=22.5 MeV; measured E(fragment), I(fragment). $^{230}, ^{232}, ^{234}\text{Th}$, $^{234}, ^{236}, ^{238}, ^{240}\text{U}$, $^{238}, ^{240}, ^{242}, ^{244}\text{Pu}$, $^{244}, ^{248}, ^{250}\text{Cm}$ deduced fission probability.

74Ba82 *Comparison of Fragment Kinetic Energies from Two ^{237}Pu Fission Isomers*

S. L. Bacharach, P. S. Hoeper, J. A. Morrissey, J. K. Temperley, Phys. Rev. C10, 2636 (1974).

Radioactivity: Fission $^{237m}\text{Pu}(\text{SF})$; measured $T_{1/2}$.

74Be52 *Attempted Coulomb Excitation of the Spontaneous-Fission Isomeric State In ^{239}Pu*

C. E. Bemis, Jr., F. Plasil, R. L. Ferguson, E. E. Gross, A. Zucker, Phys. Rev. C10, 1590 (1974).

Nuclear Reactions: $^{239}\text{Pu}(^{20}\text{Ne},^{20}\text{Ne}')$, E=100, 117 MeV; measured fission fragments. ^{239m}Pu deduced upper limit on yield.

74BeYO

Report: ORNL-4937 P26

Nuclear Reactions: $^{239}\text{Pu}(^{20}\text{Ne},^{20}\text{Ne}')$, E=100, 117 MeV; measured $\sigma(\text{fragment mass}, \theta)$. ^{239}Pu deduced fission isomer.

74Bo02 *Search for a γ -Branch from Shape Isomers in ^{236}U and ^{238}Np*

J. Borggreen, J. Hattula, E. Kashy, V. Maarbjerg, Nucl. Phys. A218, 621 (1974).

Nuclear Reactions: $^{235}\text{U}(\text{d,p})$, E=11 MeV; $^{238}\text{U}(\text{p,n})$, E=8 MeV; measured $\sigma(\text{delayed } \gamma)$, $T_{1/2}=130$ ns, $2 \mu\text{s} < T_{1/2} < 20$ ms. ^{236m}U , ^{238m}Np deduced limits on σ for delayed γ from shape isomer.

74Br05 *Investigation of the γ Decay of Subthreshold-Fission Resonances of ^{242}Pu to a Fission Isomeric State*

J. C. Browne, C. D. Bowman, Phys. Rev. C9, 1177 (1974).

Nuclear Reactions: $^{242}\text{Pu}(\text{n,F}\gamma)$, E=400-3000 eV; measured $\sigma(E)$, $\gamma(t)$. ^{243}Pu resonance deduced γ -branching.

74BrYE

Conference proceedings: Rochester(Phys. Chem of Fission), Vol2 P493

Nuclear Reactions: $^{242}\text{Pu}(\text{n,F})$, E=subthreshold; measured $E\gamma$. ^{243}Pu deduced no fission isomer.

74Ga41 *Investigation of Photonuclear Reactions Leading to Spontaneously Fissioning Isomers*

Y. P. Gangsksy, B. N. Markov, Y. M. Tsypenyuk, Fortschr. Phys. 22, 199 (1974).

Nuclear Reactions: ^{239}Pu , $^{243}\text{Am}(\gamma,\gamma')$, $^{240}, ^{242}\text{Pu}$, $^{241}, ^{243}\text{Am}(\gamma,\text{n})$, E=7-16 MeV; measured $\sigma(E)$ for the production of spontaneously fissioning isomers; deduced barrier parameters.

74GaZD *Delayed Fission Fragment Angular Distributions in Some Alpha-Particle-Induced Reactions*

D. Galeriu, M. Marinescu, D. Poenaru, I. Vilcov, N. Vilcov, Y. P. Gangsksy, P. Z. Hien, N. C. Khan, Proc. Symp. Phys. Chem. Fission, 3rd, Rochester, N. Y. (1973), Int. At. En. Agency, Vienna, Vol. 1, p. 297 (1974).

Nuclear Reactions: $^{235}, ^{238}\text{U}$, ^{239}Pu , $^{241}\text{Am}(\alpha, 2\text{n})$, ^{235}U , $^{242}\text{Pu}(\alpha, 3\text{n})$, E=26-33 MeV; measured $\sigma(\text{fragment mass}, \theta)$, fragment(t). $^{236m}, ^{237m}, ^{240m}\text{Pu}$, $^{241m}, ^{243m}\text{Cm}$ deduced anisotropies.

74HeZE *Experimental Study of the Deformation of the Fission Isomer in ^{236}U*

R. H. Heffner, Thesis, Univ. Washington (1973); Diss. Abstr. Int. B35, 435 (1974).

Radioactivity: Fission $^{236m}\text{U}(\text{SF})$; measured $\gamma\text{ce}(t)$; deduced $T_{1/2}$, β .

74LoZN *Gamma-Ray Transitions Preceding Isomeric Fission in ^{236}U*

K. E. G. Lobner, D. Harrach, E. Konecny, N. Nenoff, H. J. Specht, J. Weber, Contrib. Int. Symp. Neutron Capture Gamma Ray Spectroscopy and Related Topics, 2nd, Petten, p. 409 (1974)

Nuclear Reactions: $^{235}\text{U}(\text{n,F})$; measured (fragment) $\gamma(t)$. ^{236}U deduced transitions.

74Me10 *Detection of Fission Isomers with Half-Lives in the Picosecond Range by the Recoil-Distance Technique*

V. Metag, E. Liukkonen, G. Sletten, O. Glomset, S. Bjornholm, Nucl. Instrum. Methods 114, 445 (1974).

Nuclear Reactions: $^{237}\text{Np}(\text{p,F})$, $^{242}\text{Pu}(\text{d,pnF})$; measured recoil distance. ^{242}Pu level deduced $T_{1/2}$.

74MeYP *Half-Life Systematics of Fission Isomers in Even-Even Pu Isotopes*

V. Metag, E. Liukkonen, O. Glomset, A. Bergman, Proc. Symp. Phys. Chem. Fission, 3rd, Rochester, N. Y. (1973), Int. At. En. Agency, Vienna, Vol. 1, p. 317 (1974).

Nuclear Reactions: $^{238}, ^{240}, ^{242}, ^{244}\text{Pu}(\text{d,pn})$, $^{237}\text{Np}(\text{p,2n})$, $^{234}\text{U}(\alpha, 2\text{n})$; measured delayed fission. $^{236}, ^{238}, ^{242}, ^{244}\text{Pu}$ deduced fission isomers, $T_{1/2}$.

74MoYC *Calculation of Fission Barriers*

P. Moller, J. R. Nix, Proc. Symp. Phys. Chem. Fission, 3rd, Rochester, N. Y. (1973), Int. At. En. Agency, Vienna, Vol. 1, p. 103 (1974).

Nuclear Structure: $^{244}, ^{248}, ^{252}, ^{256}, ^{260}\text{No}$, $^{240}, ^{244}, ^{248}, ^{252}, ^{256}\text{Cf}$, $^{236}, ^{240}, ^{244}, ^{248}, ^{252}\text{Pu}$, $^{232}, ^{236}, ^{240}, ^{244}, ^{248}\text{Th}$; calculated fission barriers. A=242; calculated single particle energies.

74SpZS Fragment Anisotropy in Isomeric Fission

H. J. Specht, E. Konecný, J. Weber, C. Kozuharov, Proc. Symp. Phys. and Chem. Fission, Rochester, N. Y., 3rd, (1973), IAEA, Vienna, Vol. I, p. 285 (1974).

Nuclear Reactions: $^{235}, ^{236}\text{U}$, $^{239}\text{Pu}(\alpha, 2n)$, $E=25$ MeV; measured σ (fragment mass, θ), fragment(t). $^{237m}, ^{238m}\text{Pu}$, ^{241m}Cm deduced anisotropies, J .

74WoZW Measurements on the Fissioning Isomer ^{238m}U with the (n,n') and (d,pn) Reactions

K. L. Wolf, J. W. Meadows, Bull. Am. Phys. Soc. 19, No. 4, 595, KH1 (1974)

Nuclear Reactions: Fission $^{238}\text{U}(n,n'F)$, (d,pnF); measured $\sigma(E;E(\text{fragment}), t)$. ^{238m}U deduced $T_{1/2}$.

75Ch09 Investigation of Delayed Fission in ^{236}U

J. Christiansen, G. Hempel, H. Ingwersen, W. Klinger, G. Schatz, W. Witthuhn, Nucl. Phys. A239, 253 (1975).

Nuclear Reactions: $^{235}\text{U}(d,pF)$, $E=11$ MeV; measured prompt, delayed fission. $^{236m}\text{U}(\text{SF})$ deduced $T_{1/2}$, isomeric to prompt fission ratio. $^{232}\text{Th}(d,F)$, $E=11$ MeV; measured prompt fission.

75Gr16 Feasibility of Experimental Verification of the Shape-Isomerism Hypothesis in Heavy Nuclei

D. P. Grechukhin, Yad. Fiz. 21, 956 (1975); Sov. J. Nucl. Phys. 21, 491 (1976).

Nuclear Structure: $^{242}, ^{242m}\text{Am}$; calculated isomeric shift.

75Ha09 An Investigation of the Properties of Single-Particle-States in the Second Minimum of ^{237}Pu

I. Hamamoto, W. Ogle, Nucl. Phys. A240, 54 (1975).

Nuclear Reactions: $^{235}\text{U}(\alpha, 2n)$, $E=22-25$ MeV; analyzed data. ^{237}Pu levels deduced g, J, π, K .

75Kh06 Determination of the Spins of Spontaneously-Fissioning Isomers

P. Z. Hien, Yad. Fiz. 22, 938 (1975); Sov. J. Nucl. Phys. 22, 489 (1976).

Radioactivity: Fission $^{241}\text{Cm}(\text{SF})$, $^{235}, ^{237}, ^{238}\text{Pu}$ (SF); calculated spins of SF isomers.

75LoZT Gamma-Ray Transitions Preceding Isomeric Fission in ^{236}U

K. E. G. Lobner, D. Harrach, E. Konecný, N. Nenoff, H. J. Specht, J. Weber, Proc. Int. Symp. Neutron Capture Gamma Ray Spectroscopy and Related Topics, 2nd, Petten, The Netherlands (1974), K. Abrahams, F. Stecher-Rasmussen, P. Van Assche, Eds., Reactor Centrum Nederland, p. 665 (1975).

Nuclear Reactions: $^{235}\text{U}(n,\gamma)$, $E=\text{thermal}$; measured fragment $\gamma(t)$. ^{236}U deduced levels.

75Me28 Systematics of Fission Isomer Halflives

V. Metag, Nukleonika 20, 789 (1975).

Nuclear Structure: $^{236}, ^{238}, ^{242}, ^{244}\text{Pu}$, $^{242}, ^{244}\text{Cm}$; analyzed, reviewed fission isomer $T_{1/2}$. Systematics.

75Ru03 Gamma Decay of the ^{238}U Shape Isomer

P. A. Russo, J. Pedersen, R. Vandenberg, Nucl. Phys. A240, 13 (1975).

Nuclear Reactions: $^{238}\text{U}(d,np\gamma)$, $E=13, 18$ MeV; $^{238}\text{U}(p,p'\gamma)$, $E=13$ MeV; measured $\sigma(E\gamma, t)$. ^{238}U deduced levels, $J, \pi, T_{1/2}$, barrier parameters.

75Va21 Formation of the Spontaneously Fissile Isomer ^{242m}Am in Thermal-Neutron Capture

G. V. Valsky, V. L. Varentsov, G. A. Petrov, Y. S. Pleva, B. M. Alekseyev, A. S. Krivokhatsky, Yad. Fiz. 22, 701 (1975); Sov. J. Nucl. Phys. 22, 363 (1976).

Nuclear Reactions: $^{241}\text{Am}(n,\gamma)$, $E=\text{thermal}$; measured σ for production of $^{242}\text{Am}(\text{SF})$ isomer. ^{242m}Am deduced $T_{1/2}$.

76An11 The Shape Isomer in ^{236}U Populated by Thermal Neutron Capture

V. Andersen, C. J. Christensen, J. Borggreen, Nucl. Phys. A269, 338 (1976).

Nuclear Reactions: $^{235}\text{U}(n,\gamma)$, $E=\text{th}$; measured ce X-coin, fragment delay; obtained isomeric/prompt fission ratio. ^{236m}U shape isomer deduced γ/F branching ratio.

76Be55 Search for Conversion Electrons Emitted during the Decay of Spontaneously Fissile Isomers

A. G. Belov, Y. P. Gangarskii, B. Dalkhsuren, M. B. Miller, Izv. Akad. Nauk SSSR, Ser. Fiz. 40, 1109 (1976); Bull. Acad. Sci. USSR, Phys. Ser. 40, No. 6, 10 (1976).

Nuclear Reactions: ^{238}U , $^{239}, ^{242}\text{Pu}$, $^{241}, ^{242}\text{Am}(n, X)$, $E=14.7$ MeV; ^{238}U , $^{239}, ^{242}\text{Pu}$, $^{241}, ^{243}\text{Am}(\gamma, X)$, $E=9, 15$ MeV; measured $E(\text{ce}), I(\text{ce})$. ^{238}U deduced γ -decay for SF isomer.

76BeZM Search for the Conversion Electrons Emitted in the Decay of Spontaneously Fissioning Isomers

A. G. Belov, Y. P. Gangarsky, B. Dalkhsuren, M. B. Miller, JINR-P6-9397 (1976).

Radioactivity: Fission ^{238}U , $^{239}, ^{241}\text{Pu}$, $^{240}, ^{241}, ^{242}, ^{243}\text{Am}(\text{SF})$; measured ce spectra.

76Br38 Search for Fissile Isomers in the ($n,2n$) Reaction

J. S. Browne, R. E. Houve, At. Energ. 40, 491 (1976); Sov. At. Energy 40, 587 (1976).

Nuclear Reactions: ^{238}U , $^{242}, ^{244}\text{Pu}(n,2n)$, $E=14$ MeV; measured σ for production of SF isomers. ^{237}U , $^{241}, ^{243}\text{Pu}$ deduced no SF isomers.

76Ga11 $\Gamma n/\Gamma f$ for Actinide Nuclei Using ($^3\text{He}, df$) and ($^3\text{He}, tf$) Reactions

A. Gavron, H. C. Britt, E. Konecný, J. Weber, J. B. Wilhelmy, Phys. Rev. C13, 2374 (1976).

Nuclear Structure: $^{230}, ^{231}, ^{232}, ^{233}\text{Pa}$, $^{231}, ^{232}\text{U}$, $^{233}, ^{234}, ^{235}, ^{236}, ^{237}, ^{238}, ^{239}\text{Np}$, $^{237}, ^{238}\text{Pu}$, $^{239}, ^{240}, ^{241}, ^{242}, ^{243}\text{Am}$, $^{241}, ^{242}, ^{243}, ^{244}\text{Cm}$; measured fission probability in ^3He induced reactions; deduced barrier heights, average $\Gamma n/\Gamma f$.

Nuclear Reactions: $^{230}, ^{232}\text{Th}$, ^{231}Pa , $^{234}, ^{236}, ^{238}\text{U}$, ^{237}Np , $^{239}, ^{241}\text{Pu}$, $^{243}\text{Am}(\text{He}, df)$, $(^3\text{He}, tf)$; ^{232}U , $^{242}\text{Pu}(\text{He}, df)$; $E=25$ MeV; measured fission spectra; deduced barrier heights, average neutron-, fission-widths. $^{230}, ^{231}, ^{232}, ^{233}\text{Pa}$, $^{231}, ^{232}\text{U}$, $^{233}, ^{234}, ^{235}, ^{236}, ^{237}, ^{238}, ^{239}\text{Np}$, $^{237}, ^{238}\text{Pu}$, $^{239}, ^{240}, ^{241}, ^{242}, ^{243}\text{Am}$, $^{241}, ^{242}, ^{243}, ^{244}\text{Cm}$ deduced fission probability.

76Ga29 Study of the γ -Ray Spectra Emitted in Formation of the Spontaneously Fissile Isomer ^{236}U in the (n,γ) Reaction

Y. P. Gangrskii, A. Lajtai, B. N. Markov, *Yad. Fiz.* 24, 880 (1976); Sov. J. Nucl. Phys. 24, 460 (1976).

Nuclear Reactions: $^{235}\text{U}(n,\gamma)$, E=th; measured γ -spectrum from $^{236m}\text{U}(\text{SF})$, fragment γ -coin.

76Si01 Picosecond Fission Isomers in Even-Even Cm Isotopes

G. Sletten, V. Metag, E. Liukkonen, *Phys. Lett.* 60B, 153 (1976).

Radioactivity: Fission 240 , $^{242}\text{Cm}(\text{SF})$; measured $T_{1/2}$. $^{244}\text{Cm}(\text{SF})$; measured $T_{1/2}$ upper limit.

76We03 Mass and Kinetic Energy Measurements of Fragments from the Isomeric and Excited State Fission of ^{242}Am

J. Weber, B. R. Erdal, A. Gavron, J. B. Wilhelmy, *Phys. Rev.* C13, 189 (1976).

Radioactivity: Fission $^{242m}\text{Am}(\text{SF})$; measured $T_{1/2}$, $\sigma(E(\text{fragment mass}))$.

Nuclear Reactions: $^{241}\text{Am}(d,pF)$, E=15 MeV; measured $\sigma(E(\text{fragment mass}))$.

77Ar22 Excitation and Spontaneous Fission of ^{238m}U Isomer by Neutrons with 14 MeV Energy

R. Arlt, G. Muziol, D. Hoffman, Proc. Conf. Neutron Physics, Kiev, Part 3, p. 247 (1977).

Nuclear Reactions: $^{238}\text{U}(n,n')$, E=14 MeV; measured isomer excitation, $\sigma(\text{ratio})$.

Radioactivity: Fission $^{238m}\text{U}(\text{SF})$; measured $\sigma(\text{fragment})$ vs t.

77Bo20 On the Spontaneous Fission of ^{238}U Isomer

A. P. Bordulya, S. N. Ezhov, Proc. Conf. Neutron Physics, Kiev, Part 3, p. 244 (1977).

Radioactivity: ^{238}Pa [from $^{238}\text{U}(n,p)$, E=14.7 MeV]; measured β -delayed γ -decay. ^{238}U deduced isomer fission probability.

77Bo09 The Rotational Band of the ^{236}U Shape Isomer

J. Borggreen, J. Pedersen, G. Sletten, R. Heffner, E. Swanson, *Nucl. Phys.* A279, 189 (1977).

Nuclear Reactions: $^{235}\text{U}(d,p)$, E=12 MeV; measured ce-delayed fission coin, pce-coin. ^{236m}U deduced rotational constant.

77Di09 Near Threshold Neutron-Fission Cross Section

M. Di Toro, G. Russo, *Nucl. Phys.* A284, 177 (1977).

Nuclear Structure: ^{235}U , ^{238}Np , ^{243}Pu ; calculated fission parameters. ^{238}Np ; calculated, predicted isomer.

77Ga09 $\Gamma n/\Gamma f$ in Heavy Actinides

A. Gavron, H. C. Britt, P. D. Goldstone, R. Schoenmackers, J. Weber, J. B. Wilhelmy, *Phys. Rev.* C15, 2238 (1977).

Nuclear Reactions: ^{244}Pu , 245 , 246 , ^{248}Cm , 249 , ^{250}Cf ($^3\text{He},d$), ($^3\text{He},t$), E=8, 11 MeV; measured fission probability of compound systems 244 , ^{245}Am , 245 , 246 , 247 , 248 , ^{249}Bk , 249 , 250 , ^{251}Es .

77Go03 Cross Section for Fission of ^{244}Pu by Fast Neutrons

B. M. Gokhberg, S. M. Dubrovina, V. A. Shigin, *Yad. Fiz.* 25, 21 (1977); Sov. J. Nucl. Phys. 25, 11 (1977).

Nuclear Reactions: $^{244}\text{Pu}(n,F)$, E=fast; measured $\sigma(E)$; deduced fission threshold. ^{245}Pu deduced fission barrier height.

77GoZH Transmissionresonanzen und Winkelverteilungen der prompten Spaltung in der $^{239}\text{Pu}(d,pf)$ Reaktion

U. Goerlach, D. Habs, M. Just, V. Metag, E. Mosler, B. Neumann, P. Paul, J. Schukraft, P. Singer, H. J. Specht, G. Ulfert, C. O. Wene, Max-Planck Institut für Keinphysik (Heidelberg), Jahresbericht 1976, p. 49 (1977).

Nuclear Reactions: $^{239}\text{Pu}(d,p)$, E=11 MeV; measured fission yields; deduced transmission resonance. $^{238}\text{U}(\alpha,3n)$; measured $\gamma(\theta,H,t)$. ^{239m}Pu deduced g.

77GoYZ Messung der Energie- und Massenverteilung bei der Spaltung des ^{239m}Pu mit Hilfe des Magnetischen Rückstossionenseparator

U. Goerlach, D. Habs, M. Just, V. Metag, E. Mosler, J. Pedersen, J. Schukraft, P. Singer, H. J. Specht, G. Ulfert, C. O. Wene, Max-Planck Institut für Keinphysik (Heidelberg), Jahresbericht 1977, p. 51 (1977).

Radioactivity: Fission $^{239}\text{Pu}(\text{SF})$ [from $^{238}\text{U}(\alpha,3n)$]; measured fragment mass, kinetic energy distribution. Compared with neutron induced fission.

77Ha01 Quadrupole Moment of the 8- μs Fission Isomer in ^{239}Pu

D. Habs, V. Metag, H. J. Specht, G. Ulfert, *Phys. Rev. Lett.* 38, 387 (1977).

Nuclear Reactions: $^{238}\text{U}(\alpha,3n)$, E=33 MeV; measured charge distribution, activity by charge-plunger technique. ^{239}Pu fission isomer deduced quadrupole moment.

77Ke21 Investigation of $(n,\gamma F)$ Reaction

J. Kecskemeti, Gy. Kluge, A. Lajtai, INDC(SEC)-61/LN, p. 44 (1977).

Nuclear Reactions: $^{235}\text{U}(n,F)$, E=th; measured $\gamma\gamma(t)$. $^{236m}\text{U}(\text{SF})$ deduced transitions.

77Me08 The Quadrupole Moment of the 40 ps Fission Isomer in ^{236}Pu

V. Metag, G. Sletten, *Nucl. Phys.* A282, 77 (1977).

Nuclear Reactions: $^{234}\text{U}(\alpha,2n)$, E=25 MeV; measured delayed fission fragment(θ). ^{236}Pu shape isomer deduced $T_{1/2}$, Q_0 .

77Mi09 Fission Isomer of ^{237m}Np

E. Migneco, G. Russo, R. De Leo, A. Pantaleo, *Phys. Rev.* C16, 1919 (1977).

Nuclear Reactions: $^{238}\text{U}(n,2n)$, E=9.75, 11.6, 12.5 MeV; measured delayed/prompt fission ratios. ^{237m}Np deduced partial $T_{1/2}$ for γ , fission, branching ratio.

77Ta05 ^{239}Pu Fission Isomer in the Reaction with 3-5 MeV Neutrons

E. Takekoshi, Y. Tsukihashi, *J. Phys. Soc. Jap.* 42, 1773 (1977).

Nuclear Reactions: $^{239}\text{Pu}(n,n')$, (n,F), E=3-5 MeV; measured σ for isomer production/ σ prompt fission; deduced σ for isomer production/ σ ground state.

77VaYN Spontaneously Fissioning Isomers

R. Vandenbosch, Ann. Rev. Nucl. Sci. 27, 1 (1977).

Nuclear Structure: $^{236}, ^{238}\text{U}$, ^{237}Np , $^{235}, ^{237}, ^{239}, ^{241}, ^{243}\text{Pu}$, $^{237}, ^{238}, ^{239}, ^{240}, ^{241}, ^{242}, ^{243}, ^{245}\text{Am}$, $^{240}, ^{241}, ^{242}, ^{243}, ^{245}\text{Cm}$, $^{252}, ^{254}, ^{255}, ^{256}\text{Bk}$; compiled, reviewed isomer SF-decay $T_{1/2}$ data.

77VoZU Production of Fission Isomers in the Reaction $^{238}\text{U}(n, n')$

P. E. Vorotnikov, V. A. Vukolov, E. A. Koltypin, Yu. D. Molchanov, G. A. Otrouschenko, Proc. Conf. Neutron Physics, Kiev, Part 3, p. 239 (1977).

Nuclear Reactions: $^{238}\text{U}(n, n')$, $E=2.5\text{-}4.7$ MeV; measured fission isomer yield, $T_{1/2}$, reaction threshold.

78Ba47 Search for a γ -Decay of the ^{236}U Shape Isomer

H. Bartsch, W. Gunther, K. Huber, U. Kneissl, H. Krieger, H. J. Maier, Nucl. Phys. A306, 29 (1978).

Radioactivity: ^{236}U shape isomer [from $^{238}\text{U}(\gamma, 2n)$, $E=45$ MeV bremsstrahlung]; measured $E\gamma$, $I\gamma$; deduced $\Gamma\gamma/\Gamma f$.

78De07 Fission-Evaporation Competition in Pu Isotopes of Mass 235-239

H. Delagrange, A. Fleury, J. M. Alexander, Phys. Rev. C17, 1706 (1978).

Nuclear Reactions: $^{233}, ^{234}, ^{235}\text{U}(\alpha, xn)$, $X=1\text{-}4$, $E \leq 46$ MeV; measured fusion $\sigma(E)$.

78FI05 Statistical-Model Analysis of Fission Isomer Production for $^{237}, ^{238}\text{Pu}$ And ^{239}Am

A. Fleury, H. Delagrange, J. M. Alexander, Phys. Rev. C17, 1721 (1978).

Nuclear Reactions: $^{235}, ^{233}\text{U}$, $^{237}\text{Np}(\alpha, 2n)$, $E=22\text{-}28$ MeV; calculated $\sigma(E)$, isomer production $\sigma(E)$. Statistical model analysis.

78Go10 Resonances in the Isomeric and Prompt Fission Probabilities of ^{240}Pu

U. Goerlach, D. Habs, M. Just, V. Metag, P. Paul, H. J. Specht, H. J. Maier, Z. Phys. A287, 171 (1978).

Nuclear Reactions: $^{239}\text{Pu}(\text{d}, \text{p})$, $E=11$ MeV; measured proton-fragment time distributions, prompt, delayed fission σ ; deduced fission probability.

78Gu02 Population of the ^{236}U Shape Isomer in a Photonuclear Reaction

W. Gunther, K. Huber, U. Kneissl, H. Krieger, Nucl. Phys. A297, 254 (1978).

Nuclear Reactions: $^{238}\text{U}(\gamma, 2n)$, $E=45$ MeV bremsstrahlung; measured isomer/prompt yields; deduced σ for isomer production. ^{236m}U shape isomer deduced $T_{1/2}$, $\Gamma\gamma/\Gamma f$. Natural target.

78Po01 Properties of Fission Isomers

K. Pomorski, A. Sobiczewski, Acta Phys. Pol. B9, 61 (1978).

Nuclear Structure: ^{226}Ra , $^{230}, ^{232}\text{Th}$, $^{234}, ^{236}, ^{238}\text{U}$, $^{236}, ^{238}, ^{240}, ^{242}, ^{244}\text{Pu}$, $^{240}, ^{242}, ^{244}, ^{246}, ^{248}, ^{250}\text{Cm}$; calculated fission isomer properties: moment of inertia, pairing energy gap, g. Nilsson potential.

78SoZP Production of ^{235m}U Fission Isomer and ^{234}Pu in the Reactions $\alpha + ^{233}\text{U}$ and $^3\text{He} + ^{234}\text{U}$

L. P. Somerville, M. J. Nurmia, A. Ghiorso, G. T. Seaborg, LBL-8151, p. 39 (1978).

Nuclear Reactions: $^{234}\text{U}(^3\text{He}, 2n)$, $E=21.5\text{-}31.4$ MeV; $^{238}\text{U}(\alpha, 2n)$, $E=36.1$ MeV; measured production $\sigma(E)$. ^{235m}Pu level deduced $T_{1/2}$. Mica spontaneous fission detector.

78UI01 Lifetime Measurements of Nuclear Levels with the Charge Plunger Technique

G. Ulfert, D. Habs, V. Metag, H. J. Specht, Nucl. Instrum. Methods 148, 369 (1978).

Nuclear Reactions: $^{239}\text{Pu}(\alpha, 3n)$, $E=27, 33$ MeV; measured recoil distance. ^{240}Cm levels deduced $T_{1/2}$, Q.

79Ba02 Spectroscopy in the Second Minimum of the Potential Energy Surface of ^{239}Pu

H. Backe, L. Richter, D. Habs, V. Metag, J. Pedersen, P. Singer, H. J. Specht, Phys. Rev. Lett. 42, 490 (1979).

Radioactivity: ^{239m}Pu [from $^{238}\text{U}(\alpha, 3n)$, $E=33$ MeV]; measured E(ce), I(ce). ^{239}Pu deduced levels in second minimum, J, π , δ , rotational parameters. Nilsson assignments.

79Be33 Deep Subthreshold Photofission Yields Analysis

G. Bellia, A. Del Zoppo, E. Migneco, R. C. Barna, D. De Pasquale, Phys. Rev. C20, 1059 (1979).

Nuclear Reactions: ^{232}Th , $^{235}, ^{236}, ^{238}\text{U}(\gamma, F)$, $E=3.6, 4.1, 4.6, 5.1$ MeV (bremsstrahlung); measured σ . ^{232}Th deduced three-humped fission barrier. ^{232}Th , $^{235}, ^{236}, ^{238}\text{U}$ deduced energies, fission branching ratios for shape isomers. Double-humped fission barrier model.

79Be46 Optical Isomer Shift for the Spontaneous-Fission Isomer $^{240}\text{Am-m}$

C. E. Bernis, Jr., J. R. Beene, J. P. Young, S. D. Kramer, Phys. Rev. Lett. 43, 1854 (1979); Erratum Phys. Rev. Lett. 44, 500 (1980).

Radioactivity: ^{240}Am ; measured $T_{1/2}$, optical isomer shift. ^{240m}Am , ^{240}Am deduced difference in rms radii.

Atomic Physics: $+240)\text{Am(SF)}$; measured optical isomer shift. $^{240m}, ^{240}\text{Am}$ deduced difference in rms radii.

79Gr04 Excitation of a Spontaneously Fissile Isomer in Positron Annihilation In the K Shell of an Atom

D. P. Grechukhin, A. A. Soldatov, Yad. Fiz. 29, 296 (1979); Sov. J. Nucl. Phys. 29, 146 (1979).

Radioactivity: Fission $^{236}, ^{238}\text{U}$; calculated $T_{1/2}$ (SF).

79Gu03 Photonuclear Yields of the ^{237}Pu Fission Isomers

W. Gunther, K. Huber, U. Kneissl, H. Krieger, H. J. Maier, Phys. Rev. C19, 433 (1979).

Nuclear Reactions: $^{239}\text{Pu}(\gamma, 2n)$, $E=45$ MeV bremsstrahlung; measured $T_{1/2}$ isomeric yield ratio. ^{237m}Pu levels deduced isomeric ratio, spin. Nilsson assignments.

- 79UI01** Quadrupole Moment of the 200-ns Fission Isomer in ^{237}U
G. Ulfert, V. Metag, D. Habs, H. J. Specht, Phys. Rev. Lett. 42, 1596 (1979).
- Nuclear Reactions:** $^{238}\text{U}(\text{d},\text{pn})$, E=20 MeV; measured yield of fission-isomeric recoil. ^{238m}U level deduced quadrupole moment.
- 79Va25** On Gamma-Rays in the Population of the Spontaneously Fissioning Isomer in the Reaction $^{241}\text{Am}(\text{n},\gamma)^{242m}\text{Am}$
G. V. Valskii, V. L. Varentsov, G. A. Petrov, Y. S. Pleva, Y. A. Otschik, Pisma Zh. Eksp. Teor. Fiz. 29, 92 (1979); JETP Lett. 29, 84 (1979).
- Nuclear Reactions:** $^{241}\text{Am}(\text{n},\gamma)$, E=thermal; measured $\gamma(t)$. ^{242}Am deduced transition, E(SF) isomer.
- 80Bj02** The Double-Humped Fission Barrier
S. Bjornholm, J. E. Lynn, Rev. Mod. Phys. 52, 725 (1980).
- Nuclear Structure:** A=231-245; analyzed resonance structure, fission data; deduced fission features. Double-humped fission barrier concept.
- 80Bu13** Experimental Upper Limit for a γ Branch from the ^{236}U Shape Isomer
P. A. Butler, R. Daniel, A. D. Irving, T. P. Morrison, P. J. Nolan, V. Metag, J. Phys. (London) G6, 1165 (1980).
- Nuclear Reactions:** $^{235}\text{U}(\text{d},\text{p})$, E=11 MeV; measured $\sigma(E\gamma)$, $\gamma p(t)$. ^{236}U level deduced limit on Γ/Γ' .
- 80Bu2L** Experimental Upper Limit for a γ -Branch from the ^{236}U Shape Isomer
P. A. Butler, R. Daniels, A. D. Irving, T. P. Morrison, P. J. Nolan, V. Metag, R. Wadsworth, Univ. Liverpool, 1979-1980 Ann. Rept., p. 52 (1980)
- Nuclear Reactions:** $^{235}\text{U}(\text{d},\text{p})$, E=11 MeV; measured $E\gamma$, $I\gamma$, $\gamma p(t)$. ^{236}U deduced shape isomer Γ/Γ' upper limit.
- 80Gu20** Systematics of Photonuclear Yields and Cross Sections for Plutonium and Uranium Fission Isomers
W. Gunther, K. Huber, U. Kneissl, H. Krieger, H. Ries, H. Stroher, W. Wilke, H. J. Maier, Nucl. Phys. A350, 1 (1980).
- Nuclear Reactions:** Fission ^{240}Pu , $^{235}\text{U}(\gamma,\text{xn})$, $^{239}\text{Pu}(\gamma, 2n)$, $^{242}\text{Pu}(\gamma,\text{xn})$, E=45 MeV bremsstrahlung; measured $T_{1/2}$, isomeric to prompt yield ratios. ^{236}U , 237 , 239 , ^{241}Pu levels deduced $\sigma(\text{fission})$. Natural, enriched targets.
- 80Ku14** A Simple Description of Dependence of Fission Barriers and of the Ratio $\Gamma(n)/\Gamma(f)$ on the Nucleonic Composition for Transuranium Nuclei
V. M. Kupriyanov, K. K. Istekov, B. I. Fursov, G. N. Smirenkin, Yad. Fiz. 32, 355 (1980); Sov. J. Nucl. Phys. 32, 184 (1980).
- Nuclear Structure:** 225 , 226 , 227 , ^{228}Ra , 226 , 227 , ^{228}Ac , 227 , 228 , 229 , 230 , 231 , 232 , ^{233}Th , 234 , 231 , 232 , ^{233}Pa , 231 , 232 , 233 , 234 , 235 , 236 , 237 , 238 , 239 , ^{240}U , 233 , 234 , 235 , 236 , 237 , 238 , 239 , 240 , 241 , 242 , 243 , 244 , ^{245}Pu , 239 , 240 , 241 , 242 , 243 , 244 , 245 , 246 , ^{247}Am , 241 , 242 , 243 , 244 , 245 , 246 , 247 , 248 , 249 , ^{250}Cm , 241 , 242 , 247 , 248 , 249 , ^{250}Bk , 250 , ^{253}Cf , 249 , 250 , ^{251}Es ; calculated $\langle \text{Tn}/\text{Tf} \rangle$, fission barrier height dependences on neutron number. Phenomenological model.
- 80Li15** Spectroscopic Properties of 237 , ^{239}Pu Fission Isomers from Self-Consistent Calculations
J. Libert, M. Meyer, P. Quentin, Phys. Lett. B95, 175 (1980).
- Nuclear Structure:** 237 , ^{239}Pu ; calculated levels, $B(\lambda)$, fission isomer spectroscopic properties. Rotor plus quasiparticle model, self-consistent single particle states.
- 80Me15** Spectroscopic Properties of Fission Isomers
V. Metag, D. Habs, H. J. Specht, Phys. Rep. 65, 1 (1980).
- Nuclear Structure:** A=230-250; compiled, reviewed fission isomers data; deduced superdeformed related features.
- 80Pa16** Superprolate shape of the Spontaneous-Fission Isomer ^{240}Am
L. Pauling, Phys. Rev. C22, 1585 (1980).
- Nuclear Structure:** ^{240}Am ; calculated deformation parameter. Cluster, polyspheron theory.
- 80Ti03** Isomeric-to-Prompt Fission Ratios for the Uranium Fission Isomers ^{236m}U and ^{238m}U
R. Tischler, A. Kleinrahm, R. Kroth, C. Gunther, Phys. Rev. C22, 324 (1980).
- Nuclear Reactions:** $^{235}\text{U}(\text{d},\text{pF})$, 236 , $^{238}\text{U}(\text{d},\text{npF})$, E=17-25 MeV; measured delayed E(fission fragment). 236 , ^{238}U deduced isomeric to prompt fission ratio.
- 81Be48** Study of the Fission Isomer $^{240m}\text{Am}(\text{SF})$ using Laser-Induced Nuclear Polarization
J. R. Beene, C. E. Bemis, Jr., J. P. Young, S. D. Kramer, Hyperfine Interactions 9, 143 (1981).
- Radioactivity:** Fission $^{240m}\text{Am}(\text{SF})$ [from $^{238}\text{U}(\text{Li}, 5n)$, E=49 MeV]; measured optical isomer shift; deduced quadrupole moment. Laser induced nuclear polarization, optical pumping.
- 81Ga25** A Rotating Wheel System for the Detection of Spontaneously Fissioning Nuclides from Heavy Ion Reactions
H. Gaggeler, W. Bruchle, J. V. Kratz, M. Schadel, K. Summerer, W. Weber, G. Wirth, G. Herrmann, Nucl. Instrum. Methods 188, 367 (1981).
- Radioactivity:** Fission $^{252}\text{Cf}(\text{SF})$, $^{244}\text{Fm}(\text{SF})$ [from $^{207}\text{Pb}(\text{Ar}, 3n)$, E=199 MeV]; $^{242m}\text{Am}(\text{SF})$ [from $^{238}\text{U}(\text{Li}, X)$, E=7.6 MeV/nucleon]; measured $T_{1/2}$. Rotating wheel technique, catcher foil, fission tract detectors.
- Nuclear Reactions:** $^{238}\text{U}(\text{Li}, X)$, E=7.6 MeV/nucleon; $^{207}\text{Pb}(\text{Ar}, 3n)$, E=199 MeV; measured production $\sigma(E)$ for ^{244}Fm , ^{242}Am . Rotating wheel technique, catcher foil, fission track detectors.
- 81Gu04** Yield Ratio for the Two ^{241}Pu Fission Isomers in the $^{242}\text{Pu}(\gamma,\text{n})$ Reaction
W. Gunther, K. Huber, U. Kneissl, H. Krieger, H. Ries, H. Stroher, W. Wilke, Nucl. Phys. A359, 397 (1981).
- Nuclear Reactions:** $^{242}\text{Pu}(\gamma,\text{n})$, E=40-48 MeV bremsstrahlung; measured $T_{1/2}$, isomeric to prompt yield ratio. ^{241}Pu levels deduced isomeric ratio, J. Enriched target.

81Me19 New Results on the Spectroscopy and Dynamics of Fission

V. Metag, Nucl. Phys. A354, 271c (1981).

Nuclear Structure: $^{236, 238}\text{U}$, $^{236, 239}\text{Pu}$, ^{240}Am ; compiled, reviewed fission isomer, ground state quadrupole moment, deformation data. Other nuclei included in review.

81Re06 Analysis of Fissionability Data at High Excitation Energies I. The Level Density Problem

W. Reisdorf, Z. Phys. A300, 227 (1981).

Nuclear Structure: ^{208}Pb , $^{216, 212}\text{Th}$, ^{212}Rn ; calculated level density constant; $^{230, 231, 232, 233}\text{Pa}$, $^{231, 232}\text{U}$, $^{233, 234, 235, 236, 237, 238, 239}\text{Np}$, $^{237, 238}\text{Pu}$, $^{239, 240, 241, 242, 243}\text{Am}$, $^{241, 242, 243}\text{Cm}$; analyzed fission probabilities; deduced barrier parameters, shell correction effects. Balian-Bloch single particle level density, shell, pairing effect Ansatz.

Nuclear Reactions: Fission ^{232}Th , ^{237}Np , ^{236}U , ^{240}Pu (^3He , dF), E=25 MeV; calculated fission probability vs excitation energy. Balian-Bloch single particle level density, shell, pairing effect Ansatz.

81VaZQ Experiments on the Transfermium Element Production in Nuclear Reactions Induced by Mg Ions

V. M. Vasko, G. G. Gulbekyan, S. P. Tretyakova, E. A. Cherepanov, JINR-P7-81-863 (1981).

Radioactivity: $^{242}\text{Am}(\text{SF})$ [from ^{243}Am ($^{26}\text{Mg}, \text{X}$), E=110-140 MeV]; measured $T_{1/2}$.

Nuclear Reactions: ^{232}Th , ^{238}U , ^{243}Am ($^{24}\text{Mg}, \text{F}$), E=130 MeV; ^{232}Th , ^{238}U , ^{243}Am ($^{26}\text{Mg}, \text{F}$), E=110-140 MeV; measured fission production σ for ^{252}No , ^{265}No , ^{260}No , $T_{1/2}$ (SF).

82Fo08 Parameters of Fission Barriers for Compound Nuclei ^{245}Cm , ^{247}Cm and ^{249}Cm

E. F. Fomushkin, G. F. Novoselov, Yu. I. Vinogradov, V. V. Gavrilov, Yad. Fiz. 36, 582 (1982).

Nuclear Reactions: $^{248}\text{Cm}(\text{n,F})$, E=0.3-5.5 MeV; measured fission $\sigma(E)$. $^{245, 247, 249}\text{Cm}$ deduced fission barrier parameters. Underground nuclear explosion impulse neutron source.

82Go02 Lowest β -Vibrational Phonon in the Second Minima of $^{236, 238}\text{U}$

U. Goerlach, D. Habs, V. Metag, B. Schwartz, H. J. Specht, H. Backe, Phys. Rev. Lett. 48, 1160 (1982).

Nuclear Reactions: Fission $^{236, 238}\text{U}(\text{d,np})$, E=20 MeV; measured I(ce), ce(fragment)(t). $^{236, 238}\text{U}$ deduced shape isomer decay characteristics, K/L ratio, transition multipolarity, vibrational band characteristics.

82Ma34 Symmetry Considerations on the Fission Isomer Spectra

G. Maino, A. Ventura, Lett. Nuovo Cim. 34, 533 (1982).

Nuclear Structure: $^{236, 238}\text{U}$; calculated levels, B(E2), band structure. Interacting boson model, SU_3 limit.

82Ra04 Measurement of the g Factor of the ^{237}Pu Short-Lived Fission Isomer

M. H. Rafailovich, E. Dafni, G. Schatz, S. Y. Zhu, K. Dybdal, S. Vajda, C. Alonso-Arias, G. D. Sprouse, Phys. Rev. Lett. 48, 982 (1982); Erratum Phys. Rev. Lett. 49, 244 (1982).

Nuclear Reactions: $^{235}\text{U}(\alpha, 2n)$, E=25.2 MeV; measured $\gamma(\theta, \text{H,T})$. ^{237}Pu deduced fission isomer g, Nilsson configuration.

83Dm04 Yield of Fissionable Isomers from Reactions $^{234}\text{U}(n, n')$, $^{236}\text{U}(n,n')$, and $^{238}\text{U}(n,n')$

S. V. Dmitriev, G. A. Otroshchenko, S. M. Solovyev, Yad. Fiz. 38, 1394 (1983).

Nuclear Reactions: $^{234, 236, 238}\text{U}(n,n')$, E=2.6-4.7 MeV; measured fission isomer production $\sigma(E)$.

83Dr14 The Decay of Uranium Shape Isomers Investigated by Photonuclear Reactions

J. Drexler, R. Heil, K. Huber, U. Kneissl, G. Mank, R. Ratzek, H. Ries, H. Stroher, T. Weber, W. Wilke, Nucl. Phys. A411, 17 (1983).

Nuclear Reactions: $^{238}\text{U}(\gamma, \gamma')$, E=12 MeV bremsstrahlung; measured isomer $T_{1/2}$, isomeric to prompt yield ratio; deduced isomeric fission cross section. ^{238}U deduced isomer decay branching ratio. Natural target.

83Ka11 Observation of an E0 Isomeric Transition from the ^{238}U Shape Isomer

J. Kantele, W. Stoffl, L. E. Ussery, D. J. Decman, E. A. Henry, R. W. Hoff, L. G. Mann, G. L. Struble, Phys. Rev. Lett. 51, 91 (1983).

Radioactivity: ^{238m}U [from $^{238}\text{U}(\text{d,pn})$, E=18 MeV]; measured I(ce); deduced shape isomer E0 transition, J, π , $T_{1/2}$ assignment consistency. Reevaluation of Iy data, superconducting, solenoid type electron spectrometer.

83Po14 Identification of ^{246}Pu , ^{247}Pu , ^{246m}Am , and ^{247}Am and Determination of Their Half-Lives

Yu. S. Popov, P. A. Privalova, G. A. Timofeev, V. B. Mishenev, A. V. Mamelin, B. I. Levakov, V. M. Prokopev, Radiokhimiya 25, 482 (1983); Sov. Radiochemistry 25, 458 (1983).

Radioactivity: $^{246, 247}\text{Pu}$, $^{246m, 247}\text{Am}(\beta^-)$ [from Pu neutron irradiation]; measured E γ , I γ , E(X-ray), I(X-ray); deduced $T_{1/2}$, ^{246}Pu burnout σ . Isotope identification by α -, β -, γ -spectroscopy techniques.

83Ra36 g-Factor Measurements of Fission Isomers

M. H. Rafailovich, E. Dafni, G. Schatz, S. Y. Zhu, K. Dybdal, S. Vajda, C. Alonso-Arias, S. Rolston, G. D. Sprouse, Hyperfine Interactions 15/16, 43 (1983).

Radioactivity: ^{239m}Am , $^{237}\text{Pu}(\text{SF})$ [from ^{235}U , $^{237}\text{Np}(\alpha, 2n)$, E=25 MeV]; measured fission fragment anisotropy, isomer $T_{1/2}$, g.

Nuclear Reactions: ^{235}U , $^{237}\text{Np}(\alpha, 2n)$, E=25 MeV; measured $\gamma(\theta, \text{H,T})$. ^{239}Am , ^{237}Pu deduced fission isomer g, $T_{1/2}$.

83WeZT Search for Alpha Particle Emission from the 14-ms ^{242m}Am Shape Isomer

J. Weber, H. C. Britt, C. Fontenla, M. M. Fowler, Z. Fraenkel, A. Gavron, K. Rudolph, J. Van der Plicht, J. B. Wilhelmy, LA-9797-PR, p. 151 (1983); Isotope and Nucl. Chem. Div. Ann. Rept., 1981-1982, H. A. Lindberg Ed., Los Alamos Nat. Lab., p. 151 (1983).

Radioactivity: $^{242m}\text{Am}(\alpha, \text{SF})$ [from $^{242}\text{Pu}(\text{t,3n})$, E=17 MeV]; measured E α , I α ; deduced deexcitation shape dependence, $T_{1/2}$, long range α -particle to SF branching ratio.

83WeZU Messungen zum α -Zerfall des Formisomers ^{242m}Am

J. Weber, K. Rudolph, C. Ley, K. E. G. Lobner, S. J. Skorka, J. B. Wilhelmy, H. C. Britt, A. Gavron, Z. Fraenkel, Univ., Tech. Univ. Munich, Jahrestschift 1982, p. 16 (1983).

Radioactivity: $^{242}\text{Am}(\alpha)$ [from $^{242}\text{Pu}(\text{t,3n}), (\text{d,2n})$]; measured E α , I α .

84Bo33 Alpha Decay of Fission Isomers

N. M. Borstnik, ATOMKI Kozlem. 26, 100 (1984).

Nuclear Structure: ^{242m}Am ; calculated α -decay characteristics. Vibrational degrees, α -particle motion dynamical coupling.

84Du03 Theoretical Analysis of the Single-Particle States in the Secondary Minima of Fissioning Nuclei

J. Dudek, W. Nazarewicz, A. Faessler, Nucl. Phys. A412, 61 (1984).

Nuclear Structure: $^{239m}, ^{237m}\text{Pu}$, ^{239m}Am , $^{231}, ^{233}\text{Th}$; calculated g, single particle resonances, deformations near fission second minima. Deformed Woods-Saxon potential.

84Ka10 Reinvestigation of the Gamma Branch from the ^{238}U Shape Isomer

J. Kantele, W. Stoffl, L. E. Ussery, D. J. Decman, E. A. Henry, R. J. Estep, R. W. Hoff, L. G. Mann, Phys. Rev. C29, 1693 (1984).

Nuclear Reactions: ICPND $^{238}\text{U}(\text{d,np})$, $E=18.1$ MeV; measured $E\gamma$, $I\gamma$; deduced (isomeric/ground state) σ . ^{238}U deduced shape isomer SF, conversion decay characteristics, levels.

Radioactivity: $^{238m}\text{U}(\text{SF})$, (IT) [from $^{238}\text{U}(\text{d,np})$, $E=18.1$ MeV]; measured $E\gamma$, $I\gamma$; deduced isomer decay process relative probabilities.

84Ku05 Systematics of Neutron Cross Sections and Other Characteristics of Fission Probabilities of Transuranium Nuclei

V. M. Kupriyanov, G. N. Smirenkin, B. I. Fursov, Yad. Fiz. 39, 281 (1984).

Nuclear Structure: $^{228}, ^{229}, ^{230}, ^{231}, ^{232}, ^{233}, ^{234}, ^{235}, ^{236}, ^{237}, ^{238}, ^{239}, ^{240}, ^{241}, ^{242}\text{U}$, $^{230}, ^{231}, ^{232}, ^{233}, ^{234}, ^{235}, ^{236}, ^{237}, ^{238}, ^{239}, ^{240}, ^{241}, ^{242}\text{Np}$, $^{234}, ^{235}, ^{236}, ^{237}, ^{238}, ^{239}, ^{240}, ^{241}, ^{242}, ^{243}, ^{244}, ^{245}, ^{246}\text{Pu}$, $^{236}, ^{237}, ^{238}, ^{239}, ^{240}, ^{241}, ^{242}, ^{243}, ^{244}, ^{245}, ^{246}, ^{247}\text{Am}$, $^{238}, ^{239}, ^{240}, ^{241}, ^{242}, ^{243}, ^{244}, ^{245}, ^{246}, ^{247}, ^{248}, ^{249}, ^{250}\text{Cm}$, $^{242}, ^{243}, ^{244}, ^{245}, ^{246}, ^{247}, ^{248}, ^{249}, ^{250}, ^{251}, ^{252}, ^{253}, ^{254}, ^{255}, ^{256}\text{Cf}$, $^{244}, ^{245}, ^{246}, ^{247}, ^{248}, ^{249}, ^{250}, ^{251}, ^{252}, ^{253}, ^{254}, ^{255}\text{Es}$, $^{250}, ^{251}, ^{252}, ^{253}, ^{254}, ^{255}, ^{256}\text{Fm}$; calculated fast neutron induced fission σ ; analyzed fission data systematics; deduced fission barrier heights, $(\Gamma(n)/\Gamma(F))$. Statistical approach, two-hump fission barrier model.

84Ma44 α Decay of Fission Isomers

N. Mankoc-Borstnik, J. Phys. (London) G10, 1371 (1984).

Radioactivity: $^{242}\text{Am}(\text{EC})$, (β^-), (α), $^{242m}\text{Am}(\text{SF})$, (α); calculated α -decay constant. First-order perturbation theory.

84Ni04 On Connection between α Decay and Ternary Fission of Heavy Nuclei

A. M. Nikitin, Yad. Fiz. 39, 380 (1984).

Nuclear Structure: ^{252}Cf , $^{233}, ^{234}, ^{235}, ^{236}\text{U}$, $^{243}, ^{242m}\text{Am}$; analyzed ternary fission light fragment emission, α -decay characteristics systematics; deduced initial nucleus quasistationary α -particle state role.

84Oh09 Systematic Analysis of Fission Cross Sections of Actinides by Means of Double-Humped Barrier Model

T. Ohsawa, Y. Shigemitsu, M. Ohta, K. Kudo, J. Nucl. Sci. Technol. (Tokyo) 21, 887 (1984).

Nuclear Reactions: $^{231}\text{Pa}, ^{232}, ^{234}, ^{235}, ^{236}, ^{238}\text{U}$, ^{237}Np , $^{238}, ^{239}, ^{240}, ^{241}, ^{242}\text{Pu}$, $^{241}, ^{242}, ^{243}\text{Am}$, $^{243}, ^{244}, ^{245}, ^{246}, ^{247}, ^{248}\text{Cm}$, ^{249}Bk , $^{252}\text{Cf}(n, F)$, $E=0.5-6$ MeV; calculated $\sigma(E)$; deduced optical model parameters. ^{232}Pa , $^{233}, ^{235}, ^{236}, ^{237}, ^{239}\text{U}$, ^{238}Np , $^{239}, ^{240}, ^{241}, ^{242}, ^{243}, ^{245}\text{Pu}$, $^{242}, ^{243}, ^{244}\text{Am}$, $^{244}, ^{245}, ^{246}, ^{247}, ^{248}, ^{249}\text{Cm}$, ^{250}Bk , ^{253}Cf deduced fission barriers. Double-humped barrier model.

84Vo18 Energy Dependence of Yield of Fission Isomers in the Reactions $^{241}\text{Am}(n,\gamma)$ and $^{243}\text{Am}(n,\gamma)$

P. E. Vorotnikov, G. A. Otroshchenko, Yad. Fiz. 40, 1135 (1984).

Nuclear Reactions: $^{241}, ^{243}\text{Am}(n,\gamma)$, $E=0.2-1.3$ MeV; measured fission isomer, prompt fission product yield ratios.

85Ba20 On Measurement of the Angular Momenta of Fission Isomer

A. L. Barabanov, D. P. Grechukhin, Yad. Fiz. 41, 582 (1985).

Nuclear Reactions: $^{238}\text{U}(^7\text{Li}, 5n)$, $E=46.1$ MeV; analyzed data. ^{240m}Am deduced fission isomer J estimate. Residual orientation in laser radiation field.

85Be58 Laser Optical Pumping in Nuclear Physics: Fission isomers, oriented targets, and hyperfine pumping in single-electron atoms

C. E. Bemis, Jr., Hyperfine Interactions 24, 139 (1985).

Radioactivity: $^{240m}\text{Am}(\text{SF})$; measured optical isomer shift. Oriented targets, anisotropic fission decay from resonant laser optical pumping.

85Dr01 The 'Isomeric Shelf' in the Deep Subbarrier Photofission of ^{238}U

J. Drexler, R. D. Heil, K. Huber, U. Kneissl, G. Mank, R. Ratzek, H. Ries, T. Weber, W. Wilke, B. Fischer, H. Hollick, Nucl. Phys. A437, 253 (1985).

Nuclear Reactions: $^{238}\text{U}(\gamma, F)$, $E=3.9-4.3$ MeV bremsstrahlung; measured $T_{1/2}$ isomeric to prompt yield ratio. Depleted targets.

85Ig01 Analysis of Cross Sections of U and Pu Isotope Fission Induced by Neutrons in the Range of the First 'Plateau'

A. V. Ignatyuk, A. B. Klepatsky, V. M. Maslov, E. Sh. Sukhovitsky, Yad. Fiz. 42, 569 (1985).

Nuclear Reactions: $^{239}, ^{240}, ^{241}, ^{242}, ^{244}, ^{245}\text{Pu}$, $^{234}, ^{235}, ^{236}, ^{237}, ^{238}, ^{239}, ^{240}\text{U}(n, F)$, $E=1-5.5$ MeV; analyzed fission $\sigma(E)$. $^{240}, ^{241}, ^{242}, ^{243}, ^{245}, ^{246}\text{Pu}$, $^{235}, ^{236}, ^{237}, ^{238}, ^{239}, ^{240}, ^{241}\text{U}$ deduced fission barriers, transitional states statistical characteristics.

85Jo04 ^{241}Am and ^{243}Am Charge Distributions from Muonic X-Ray Spectroscopy and the Quadrupole Moment of the ^{240}Am Fission Isomer

M. W. Johnson, E. B. Shera, M. V. Hoehn, R. A. Naumann, J. D. Zumbro, C. E. Bemis, Jr., Phys. Lett. 161B, 75 (1985).

Nuclear Reactions: $^{241}, ^{243}\text{Am}(\mu, X)$, E at rest; measured muonic E X-ray, I X-ray. $^{241}, ^{243}\text{Am}$ deduced intrinsic quadrupole moment, Barrett radii. ^{240}Am deduced fission isomer quadrupole moment. Optical isotope shift data input.

Atomic Physics: esic-Atoms $^{241}, ^{243}\text{Am}(\mu, X)$, E at rest; measured muonic X-rays.

85Ku18 Excitation of Fission Isomer ^{242m}Am , $^{242}\text{Am}(f)$ by Electrons in the Energy Region 17.5-78 MeV

V. L. Kuznetsov, L. E. Lazareva, V. G. Nedorezov, N. V. Nikitina, A. S. Sudov, Yad. Fiz. 42, 29 (1985).

Nuclear Reactions: $^{243}\text{Am}(e, n)$, (γ, n) , $E=17.5-78$ MeV; measured residual fission isomer production $\sigma(E)$; $^{243}\text{Am}(e, F)$, (γ, F) , $E=17.5-78$ MeV; measured fission $\sigma(E)$; deduced fission isomer production mechanism. Virtual photon theory.

- 85Ra28** A α -Factor Measurement of the ^{239}Am Fission Isomer
 M. H. Rafailovich, S. Vajda, E. Dafni, G. Schatz, S. Rolston, S. Y. Zhu, G. D. Sprouse, Phys. Lett. 163B, 327 (1985).
- Nuclear Reactions:** $^{237}\text{Np}(\alpha, 2n)$, E=tandem; measured $\gamma(\theta, H, t)$. ^{239}Am deduced fission isomer g.
- 85Vo17** Anisotropy of Fission of ^{242m}Am by Fast Neutrons
 P. E. Vorotnikov, B. M. Gokberg, V. A. Shigin, E. F. Fomushkin, G. F. Novoselov, Yad. Fiz. 42, 1038 (1985); Sov. J. Nucl. Phys. 42, 656 (1985).
- Radioactivity:** $^{242m}\text{Am}(\text{SF})$; measured fission fragment decay $\sigma(\theta n=0^\circ)/\sigma(\theta n=90^\circ)$, anisotropy.
- 86Bl10** Intermediate Structure in the Fission Cross Sections of the Even Curium Isotopes
 R. C. Block, D. R. Harris, H. T. Maguire, Jr., C. R. S. Stopa, R. E. Slovacek, J. W. T. Dabbs, R. J. Dougan, R. W. Hoff, R. W. Lougheed, Radiat. Eff. 92, 305 (1986).
- Nuclear Reactions:** $^{244}, ^{246}, ^{248}\text{Cm}(n, F)$, E \leq 100 keV; analyzed fission $\sigma(E)$; deduced structure. $^{245}, ^{247}, ^{249}\text{Cm}$ deduced barrier parameter differences.
- 86De04** Excitation Function and Half-Life for the Fission Isomer ^{240m}Pu from the $^{238}\text{U}(\alpha, 2n)^{240m}\text{Pu}$ Reaction
 S. de Barros, S. D. de Magalhaes, H. Wolf, J. Barreto, J. Eichler, N. Lisboa, I. O. de Souza, D. M. Vianna, Z. Phys. A323, 101 (1986).
- Radioactivity:** $^{240m}\text{Pu}(\text{SF})$ [from $^{238}\text{U}(\alpha, 2n)$, E=20.1-27.3 MeV]; measured $T_{1/2}$.
- Nuclear Reactions:** ICPND $^{238}\text{U}(\alpha, 2n)$, E=20.1-27.3 MeV; measured residual fission isomer production $\sigma(E)$. ^{240m}Pu deduced delayed fission σ , isomeric σ ratio.
- 87Ah07** Search for the Shape-Isomeric Gamma Decay in Muonic Uranium
 S. Ahmad, G. A. Beer, B. H. Olaniyi, A. Olin, S. N. Kaplan, A. Mireshghi, J. A. Macdonald, O. Hausser, Can. J. Phys. 65, 753 (1987).
- Nuclear Reactions:** $^{236}, ^{238}, ^{239}\text{U}(\mu^-, \gamma)$, E at rest; measured $E\gamma, I\gamma, E$ X-ray, I X-ray, $\gamma(t)$; deduced no shape isomer excitation evidence. $^{235}, ^{236}, ^{238}\text{U}$ deduced μ -capture $T_{1/2}$.
- Atomic Physics:** esic-Atoms $^{236}, ^{238}, ^{239}\text{U}(\mu^-, \gamma)$, E at rest; measured $E\gamma, I\gamma, E$ X-ray, I X-ray, $\gamma(t)$.
- 87Gu03** A New Macroscopic-Microscopic Description of the Double-Humped Fission Barriers
 S. K. Gupta, L. Satpathy, Z. Phys. A326, 221 (1987).
- Nuclear Structure:** $^{228}\text{Ra}, ^{228}\text{Ac}, ^{228}\text{Th}, ^{229}\text{Pa}, ^{234}\text{U}, ^{238}\text{Np}, ^{239}\text{Pu}, ^{241}\text{Am}, ^{243}\text{Cm}, ^{248}\text{Bk}, ^{250}\text{Cf}, ^{254}\text{Es}, ^{256}\text{Fm}, ^{256}\text{Md}, ^{257}\text{No}, ^{259}\text{Lr}, ^{261}\text{No}$; calculated binding energies; Z=90-98; calculated doubled-humped fission barriers, shell energies. New mass relation.
- 87ScZP** On the γ -Decay of the Shape Isomer in ^{236}U
 J. Schirmer, D. Habs, D. Schwalm, H. J. Maier, GSI-87-1, p. 32 (1987).
- Nuclear Reactions:** $^{235}\text{U}(\text{d,p})$, E=11 MeV; measured γ -spectra, $\gamma(t)$. ^{236}U deduced shape isomer decay characteristics.
- 88Ma43** α -Decay Probability of Spontaneously Fissioning Isomer and Deformation Hindrance Factor
 V. E. Makarenko, V. G. Nosov, Yad. Fiz. 48, 73 (1988).
- Radioactivity:** $^{238m}\text{U}(\alpha)$; calculated α -decay probability; deduced deformation hindrance factor.
- 88Ma52** Triple Fission of the Spontaneously Fissioning Isomer ^{238}U
 V. E. Makarenko, Yu. D. Molchanov, G. A. Otroshchenko, G. B. Yan'kov, Pisma Zh. Eksp. Teor. Fiz. 47, 489 (1988); JETP Lett. (USSR) 47, 573 (1988).
- Radioactivity:** $^{238m}\text{U}(\alpha)$, (SF) [from $^{238}\text{U}(n, n')$, E=4.5 MeV]; measured decay $T_{1/2}$, triple fission branching ratio.
- Nuclear Reactions:** $^{238}\text{U}(n, n')$, E=4.5 MeV; measured isomer production yield.
- 89Eg01** Actinide Nuclei Fission Cross-Section Irregularities
 S. A. Egorov, V. A. Rubchenya, S. V. Khlebnikov, Nucl. Phys. A494, 75 (1989).
- Nuclear Reactions:** $^{227}\text{Ac}(n, F)$, E 2-16 MeV; $^{226}\text{Ra}(n, F)$, E 3-16 MeV; $^{244}\text{Cm}(n, F)$, E 1-3 MeV; $^{242}, ^{246}, ^{248}\text{Cm}(n, F)$, E 0.5-5 MeV; calculated fission $\sigma(E)$. $^{243}, ^{245}, ^{247}, ^{249}\text{Cm}, ^{226}, ^{228}, ^{227}\text{Ra}, ^{228}, ^{227}\text{Ac}$ deduced fission barrier parameters.
- 89Ha40** Spectroscopy of the Second Minimum
 D. Habs, Nucl. Phys. A502, 105c (1989).
- Nuclear Structure:** A 150; analyzed high spin level systematics; deduced comparison with fission second minimum in actinide region.
- 89HoZP** Second Minimum Spectroscopy Using Heavy Ion Transfer Reactions
 T. H. Hoare, P. A. Butler, G. D. Jones, R. J. Poynter, C. A. White, Daresbury Lab., 1988-1989 Ann. Rept., Appendix, p. 92 (1989).
- Nuclear Reactions:** $^{238}\text{U}(^{58}\text{Ni}, ^{60}\text{Ni})$, E=325 MeV; measured $\gamma\gamma$ -coin, $\gamma\gamma(t)$; deduced residue fission isomer σ .
- 89Ma54** Ternary Fission of Neutron Induced Uranium Fissioning Isomers
 V. E. Makarenko, Yu. D. Molchanov, G. A. Otroshchenko, G. B. Yan'kov, Nucl. Phys. A502, 363c (1989).
- Radioactivity:** $^{236m}, ^{238m}\text{U}(\text{SF})$ [from $^{238}, ^{236}\text{U}(n, n')$, E=4.5 MeV]; measured $T_{1/2}$, fission fragment; deduced relative fission probabilities.
- 89Ma57** Ternary Fission of Uranium Fissioning Isomers Excited by Neutrons
 V. E. Makarenko, Yu. D. Molchanov, G. A. Otroshchenko, G. B. Yan'kov, Yad. Fiz. 50, 928 (1989).
- Radioactivity:** $^{236m}, ^{238m}\text{U}(\text{SF})$ [from $^{236}, ^{238}\text{U}(n, n')$, E=4.5 MeV]; measured fission fragment spectra; deduced $T_{1/2}$, decay probability, fission mechanism.
- 89Ma64** Spontaneous Fission Isomers α -Decay Hindrance Factor
 V. E. Makarenko, V. G. Nosov, Izv. Akad. Nauk SSSR, Ser. Fiz. 53, 933 (1989); Bull. Acad. Sci. USSR, Phys. Ser. 53, No. 5, 105 (1989).
- Radioactivity:** $^{238m}\text{U}(\text{SF})$; calculated α -decay hindrance factor.

89Sc30 γ Decay of the Superdeformed Shape Isomer in ^{236}U

J. Schirmer, J. Gerl, D. Habs, D. Schwalm, Phys. Rev. Lett. 63, 2196 (1989).

Nuclear Reactions: $^{235}\text{U}(\text{d},\text{p})$, E=11 MeV; measured γ time spectra, missing energy vs delayed sum energy. ^{236}U deduced isomer, decay, superdeformation features, γ -decay to fission branching ratio.

89So2Z Production of the Fission Isomer ^{235m}Pu and ^{234}Pu in the Reactions $\alpha + ^{233}\text{U}$ and $^3\text{He} + ^{234}\text{U}$

L. P. Somerville, M. J. Nurmia, A. Ghiors, J. M. Nitschke, G. T. Seaborg, Bull. Am. Phys. Soc. 34, No. 1, 69, EG7 (1989)

Nuclear Reactions: ICPND $^{234}\text{U}(^3\text{He},2n)$, ($^3\text{He},3n$), E not given; measured $\sigma(E)$. $^{233}\text{U}(\alpha,3n)$, E=36 MeV; measured E(α), I(α); deduced reaction σ , 235 , ^{234}Pu production.

90Bh02 Test of the Adequacy of Using Smoothly Joined Parabolic Segments to Parametrize the Multihumped Fission Barriers in Actinides

B. S. Bhandari, Phys. Rev. C42, 1443 (1990).

Nuclear Structure: 236 , ^{238}U , ^{237}Np , 235 , 237 , 238 , 239 , 240 , 241 , 242 , 243 , 244 , ^{245}Pu , 239 , 240 , 241 , 242 , 243 , 244 , ^{245}Am , 241 , 242 , 243 , 244 , ^{245}Cm ; calculated fission $T_{1/2}$; deduced fission barrier parametrization.

90HoZU Second Minimum Spectroscopy Using Heavy Ion Reactions

T. H. Hoare, P. A. Butler, N. Clarkson, G. D. Jones, C. A. White, R. J. Poyer, R. A. Cunningham, Daresbury Labs., 1989-1990 Ann. Rept., Appendix, p. 84 (1990).

Nuclear Reactions: ICPND $^{238}\text{U}(^{58}\text{Ni},^{60}\text{Ni})$, E=325 MeV; $^{238}\text{U}(^{62}\text{Ni},^{64}\text{Ni})$, E=332 MeV; measured fission isomer production σ upper limit.

90Ku17 Energy of Alpha Particles in Triple Fission of the Fissile Isomer Uranium-238

I. A. Kukushkin, V. E. Makarenko, Yu. D. Molchanov, G. A. Otroschenko, G. B. Yankov, Pisma Zh. Eksp. Teor. Fiz. 51, 611 (1990); JETP Lett. (USSR) 51, 693 (1990).

Radioactivity: ^{238m}U [from $^{238}\text{U}(\text{n},\text{n}')$, E=4.5 MeV]; measured fission fragment, α -spectra; deduced $T_{1/2}$, triple fission α -distribution features, branching ratio relative to SF-decay.

90Ma59 Method of Half-Life Determination

V. E. Makarenko, G. A. Otroschenko, Yad. Fiz. 51, 1201 (1990); Sov. J. Nucl. Phys. 51, 765 (1990).

Radioactivity: 236m , ^{238m}U ; calculated $T_{1/2}$. Time spectrum processing method proposed.

91Ku23 Energies of Long-Range Particles in Ternary Fission of the ^{238}U Spontaneously Fissioning Isomer

I. A. Kukushkin, V. E. Makarenko, Yu. D. Molchanov, G. A. Otroschenko, G. B. Yankov, Yad. Fiz. 54, 8 (1991); Sov. J. Nucl. Phys. 54, 4 (1991).

Nuclear Reactions: $^{238}\text{U}(\text{n},\text{n}')$, E=4.5 MeV; measured (fragment)(fragment)-coin following SF-decay, ternary fission. ^{238m}U deduced $T_{1/2}$, fission branching ratio.

92Ba67 First Observation of a Resonance Ionization Signal on ^{242m}Am Fission Isomers

H. Backe, Th. Blonigen, M. Dahlinger, U. Doppler, P. Graffe, D. Habs, M. Hies, Ch. Illgner, H. Kunz, W. Lauth, H. Schoppe, P. Schwamb, W. Theobald, R. Zahn, Hyperfine Interactions 74, 47 (1992).

Radioactivity: $^{242m}\text{Am}(\text{SF})$ [from $^{242}\text{Pu}(\text{d},2\text{n})$, E=12 MeV]; measured resonance ionization followed by isomer fission decay. Buffer gas cell, two-step resonance ionization, excimer dye laser combination.

92Bh03 Systematics of the Deduced Fission Barriers for the Doubly Even Transactinum Nuclei

B. S. Bhandari, Y. B. Bendardaf, Phys. Rev. C45, 2803 (1992).

Nuclear Structure: 236 , 238 , 240 , 242 , ^{244}Pu , 240 , 242 , 244 , 246 , 248 , ^{250}Cm ; calculated isomer energies, $T_{1/2}$, SF-decay $T_{1/2}$, outer barrier heights. 230 , ^{232}Th , 230 , 232 , 234 , ^{236}U , 246 , 248 , 250 , 252 , 254 , ^{256}Cf , 242 , 244 , 246 , 248 , 250 , 252 , 254 , ^{256}Fm , 250 , 252 , 254 , 256 , ^{258}No , $^{250}104$, $^{252}104$, $^{254}104$, $^{256}104$, $^{258}104$; calculated SF-decay $T_{1/2}$, outer barrier height. Double humped fission barrier model. Other nuclei, other aspects discussed.

92BIZZ Search for Low Spin Superdeformed States by Transfer Reaction

J. Blons, D. Goutte, A. Lepretre, R. Lucas, V. Meot, D. Paya, X. H. Phan, G. Barreau, T. Doan, G. Pedemey, Contrib. Int. Conf. Nuclear Structure at High Angular Momentum, Ottawa, p. 57 (1992); AECL-10613 (1992)

Nuclear Reactions: $^{236}\text{U}(^{18}\text{O},^{16}\text{O})$, E=9 MeV/nucleon; $^{192}\text{Pt}(^{16}\text{O},^{14}\text{C})$, E not given; measured γ sum spectra, γ (particle)-coin. ^{194}Hg deduced superdeformed band population.

92Ch08 Limits on the Lifetime of the Shape Isomer of ^{238}U

C. R. Chinn, J. -F. Berger, D. Gogny, M. S. Weiss, Phys. Rev. C45, 1700 (1992).

Radioactivity: ^{238}U ; calculated fission isomer partial $T_{1/2}$. Constrained Hartree-Fock-Bogoliubov.

92De2Z Population of the 0.5ns Fission Isomer and Excited States in ^{238}Pu by Heavy-Ion Induced 1n-Transfer

M. Devlin, D. Cline, K. G. Helmer, R. Ibbotson, C. Y. Wu, A. Cresswell, P. A. Butler, G. D. Jones, M. A. Stoyer, J. O. Rasmussen, Bull. Am. Phys. Soc. 37, No. 2, 870, A8 1 (1992)

Nuclear Reactions: $^{239}\text{Pu}(^{117}\text{Sn},^{118}\text{Sn})$, E=630 MeV; measured E_γ , I_γ , γ -multiplicity, particle spectra, (fragment)(fragment)-coin. ^{239}Pu deduced levels, J, π . ^{238}Pu deduced levels, J, π , fission isomer population.

92Er01 Quasi-Stationary State Population Probability of the Actinide Nuclei Second Well

D. O. Eremenko, S. Yu. Platonov, O. A. Yuminov, Bull. Rus. Acad. Sci. Phys. 56, 70 (1992).

Nuclear Structure: 239 , 238 , 236 , ^{235}Np , 240 , 238 , ^{237}Pu , 238 , ^{232}Pa ; calculated quasistationary states population probability under induced fission, second potential. Fluctuation dissipation dynamics.

92Ma34 α and γ Spectroscopy of Spontaneous-Fission Isomers

V. E. Makarenko, Yad. Fiz. 55, 1759 (1992); Sov. J. Nucl. Phys. 55, 973 (1992).

Nuclear Structure: ^{238}U , 239 , ^{241}Pu , 240 , 241 , 242 , ^{243}Am ; compiled, reviewed fission isomer decay by α -, γ -emission.

92So10 *Intrinsic Structures and Associated Rotational Bands in Deformed Even-Even Nuclei of the Actinide Region*

P. C. Sood, D. M. Headly, R. K. Sheline, At. Data Nucl. Data Tables 51, 273 (1992).

Nuclear Structure: $Z \geq 88$; $N \geq 134$; $^{230}, ^{232}, ^{234}, ^{236}, ^{238}\text{U}$, $^{220}, ^{222}, ^{224}, ^{226}, ^{228}, ^{230}, ^{232}, ^{234}\text{Th}$, $^{218}, ^{220}, ^{222}, ^{224}, ^{226}, ^{228}, ^{230}\text{Ra}$; analyzed levels; deduced band structure, fission isomers superdeformation, hyperdeformation evidence.

92St05 *Fission and Gamma-Ray Decay of the ^{238}U Shape Isomer*

M. Steinmayer, K. E. G. Lobner, L. Corradi, U. Lenz, U. Quade, P. R. Pascholati, K. Rudolph, W. Schomburg, Z. Phys. A341, 145 (1992).

Radioactivity: $^{238\text{m}}\text{U}$ [from $^{238}\text{U}(\text{d,np})$, $E=18$ MeV]; measured $\gamma(\text{ce})$ -coin; deduced delayed fission $T_{1/2}$. ^{238}U deduced transitions.

93Ar03 *Fission of Heavy Hypernuclei Formed in Antiproton Annihilation*

T. A. Armstrong, J. P. Bocquet, G. Ericsson, T. Johansson, T. Krogulski, R. A. Lewis, F. Malek, M. Maurel, E. Monnand, J. Mougey, H. Nifenecker, J. Passaneau, P. Perrin, S. M. Polikanov, M. Rey-Campagnolle, C. Ristori, G. A. Smith, G. Tibell, Phys. Rev. C47, 1957 (1993).

Nuclear Reactions: $^{238}\text{U}(\bar{p},X)$, E at 105 MeV/c; measured hypernuclei yield, fission (fragment)(fragment)-coin; deduced fission hypernuclei $T_{1/2}$.

93Ku16 *Yield of the Fissioning Isomer in the Reaction $^{241}\text{Am}(n, n')$*

I. A. Kukushkin, V. E. Makarenko, Yu. D. Molchanov, G. A. Otroshchenko, Yad. Fiz. 56, No 9, 13 (1993); Phys. Atomic Nuclei 56, 1157 (1993).

Nuclear Reactions: $^{241}\text{Am}(n,n')$, (n,γ) , $E=4.5$ MeV; measured fission isomer yields; deduced reaction dependence.

Radioactivity: $^{242\text{m}}, ^{241\text{m}}\text{Am}(\text{SF})$ [from $^{241}\text{Am}(n,n')$, (n,γ) , $E=4.5$ MeV]; measured fission fragment spectra. ^{241}Am deduced isomeric state fission probability, $T_{1/2}$.

93Ro07 *The Study of Prompt and Delayed Muon Induced Fission III. The Ratios of Prompt to Delayed Fission Yields*

Ch. Rosel, H. Hanscheid, J. Hartfiel, R. von Mutius, J. F. M. d'Achard van Enschut, P. David, H. Janszen, T. Johansson, J. Konijn, T. Krogulski, C. T. A. M. de Laat, H. Paganetti, C. Petitjean, S. M. Polikanov, H. W. Reist, F. Rissee, L. A. Schaller, L. Schellenberg, W. Schrieder, A. K. Sinha, A. Taal, J. P. Theobald, G. Tibell, N. Trautmann, Z. Phys. A345, 89 (1993).

Nuclear Reactions: $^{233}, ^{234}, ^{235}, ^{236}, ^{238}\text{U}$, ^{237}Np , $^{242}, ^{244}\text{Pu}(\mu,\text{F})$, E not given; measured prompt to delayed fission yields ratios, absolute probabilities; deduced fission probabilities per muon capture.

94Kr06 *GCM Calculation of the E2 Decay Lifetimes of Shape Isomers*

S. J. Krieger, P. Bonche, H. Flocard, P. H. Heenen, M. S. Weiss, Nucl. Phys. A572, 384 (1994).

Nuclear Structure: $^{230}, ^{232}\text{Th}$, ^{238}U ; calculated deformation energy vs mass quadrupole moment, first barrier, second minimum, absolute minimum quadrupole moment, charge quadrupole transition matrix element between superdeformed, ground bands, isomer E2 decay $T_{1/2}$. Hartree-Fock BCS calculations.

94Ob02 *Intermediate Structure and the Shape Isomer in ^{233}Th*

S. Oberstedt, J. P. Theobald, H. Weigmann, J. A. Wartena, C. Burholz, Nucl. Phys. A578, 31 (1994).

Nuclear Reactions: $^{232}\text{Th}(\text{n},\gamma)$, $E=0.05-4.2$ keV; measured $\gamma\gamma(t)$. ^{233}Th deduced shape isomer $T_{1/2}$ range, decay features, resonances admixture, coupling width.

94ReAA

Gamma Spectroscopy of Superdeformed ^{236}U . Reiter et al., Annual Report GSI 94-1 (March 1994), p 75.

94PaAA

Gamma Spectroscopy in the Second Minimum of ^{240}Pu . Pansegrouw et al., Annual Report GSI 94-1 (March 1994), p 76.