

**Table 17. Adult/adolescent AIDS cases by single and multiple exposure categories, reported through December 2001, United States**

Exposure category	AIDS cases	
	No.	(%)
<b>Single mode of exposure</b>		
Men who have sex with men	351,612	(44)
Injecting drug use	156,869	(19)
Hemophilia/coagulation disorder	4,308	(1)
Heterosexual contact	88,163	(11)
Receipt of transfusion <sup>1</sup>	8,958	(1)
Receipt of transplant of tissues, organs, or artificial insemination <sup>2</sup>	13	(0)
Other <sup>3</sup>	256	(0)
<b>Single mode of exposure subtotal</b>	<b>610,179</b>	<b>(76)</b>
<b>Multiple modes of exposure</b>		
Men who have sex with men; injecting drug use	43,096	(5)
Men who have sex with men; hemophilia/coagulation disorder	208	(0)
Men who have sex with men; heterosexual contact	13,128	(2)
Men who have sex with men; receipt of transfusion/transplant	3,613	(0)
Injecting drug use; hemophilia/coagulation disorder	219	(0)
Injecting drug use; heterosexual contact	41,114	(5)
Injecting drug use; receipt of transfusion/transplant	1,766	(0)
Hemophilia/coagulation disorder; heterosexual contact	136	(0)
Hemophilia/coagulation disorder; receipt of transfusion/transplant	808	(0)
Heterosexual contact; receipt of transfusion/transplant	1,968	(0)
Men who have sex with men; injecting drug use; hemophilia/coagulation disorder	55	(0)
Men who have sex with men; injecting drug use; heterosexual contact	7,245	(1)
Men who have sex with men; injecting drug use; receipt of transfusion/transplant	656	(0)
Men who have sex with men; hemophilia/coagulation disorder; heterosexual contact	24	(0)
Men who have sex with men; hemophilia/coagulation disorder; receipt of transfusion/transplant	46	(0)
Men who have sex with men; heterosexual contact; receipt of transfusion/transplant	334	(0)
Injecting drug use; hemophilia/coagulation disorder; heterosexual contact	100	(0)
Injecting drug use; hemophilia/coagulation disorder; receipt of transfusion/transplant	38	(0)
Injecting drug use; heterosexual contact; receipt of transfusion/transplant	1,193	(0)
Hemophilia/coagulation disorder; heterosexual contact; receipt of transfusion/transplant	40	(0)
Men who have sex with men; injecting drug use; hemophilia/coagulation disorder; heterosexual contact	20	(0)
Men who have sex with men; injecting drug use; hemophilia/coagulation disorder; receipt of transfusion/transplant	16	(0)
Men who have sex with men; injecting drug use; heterosexual contact; receipt of transfusion/transplant	200	(0)
Men who have sex with men; hemophilia/coagulation disorder; heterosexual contact; receipt of transfusion/transplant	6	(0)
Injecting drug use; hemophilia/coagulation disorder; heterosexual contact; receipt of transfusion/transplant	27	(0)
Men who have sex with men; injecting drug use; hemophilia/coagulation disorder; heterosexual contact; receipt of transfusion/transplant	5	(0)
<b>Multiple modes of exposure subtotal</b>	<b>116,061</b>	<b>(14)</b>
<b>Risk not reported or identified<sup>4</sup></b>	<b>80,835</b>	<b>(10)</b>
<b>Total</b>	<b>807,075</b>	<b>(100)</b>

<sup>1</sup>Includes 41 adult/adolescents who developed AIDS after receiving blood screened negative for HIV antibody.

<sup>2</sup>Thirteen adults developed AIDS after receiving tissue, organs, or artificial insemination from HIV-infected donors. Four of the 13 received tissue or organs from a donor who was negative for HIV antibody at the time of donation. See *N Engl J Med* 1992;326:726-32.

<sup>3</sup>"Other" includes 221 persons who acquired HIV infection perinatally, but had AIDS diagnosed after age 13. See Technical Notes.

<sup>4</sup>See Technical Notes.