
Saturation & BFKL dynamics in the HERA data at small x

Kazunori Itakura
(SPhT, CEA/Saclay)

E.Iancu, K.I. & S.Munier, hep-ph/0310338

Plan of the talk:

- Introduction
- Dipole picture in DIS at small x
- Previous attempts with saturation models
- Physics at $x < 10^{-2}$, $Q^2 \lesssim 50\text{GeV}^2$
- A CGC fit to the HERA F_2 data
- Summary


Introduction

Deep Inelastic Scattering (DIS) at small x is the best place to see **Color Glass Condensate**.


- can directly probe the hadron wavefunction
- increase of gluon density at small x is observed in F_2
- can enter CGC regime with increasing energy
- relation to RHIC

Saturation scale $Q_s^2(x, A) \propto A^{1/3}(1/x)^{0.3}$

DIS @ HERA $x \sim 10^{-4} \leftrightarrow$ AuAu @ RHIC $x \sim 10^{-2}$



Dipole picture in DIS at small x



F_2 structure function

$$F_2(x, Q^2) = \frac{Q^2}{4\pi^2\alpha_{\text{EM}}} \sum_{T,L} \int_0^1 dz \int d^2r_{\perp} |\Psi_{T,L}(z, r_{\perp}, Q^2)|^2 \sigma_{\text{dipole}}(x, r_{\perp})$$

where $\Psi_{T,L}$: LC wavefunc of γ^* (perturbatively calculable)

$\sigma_{\text{dipole}}(x, r_{\perp})$: Dipole-proton cross section

- $\propto r_{\perp}^2 x G(x, 1/r_{\perp}^2)$ at very short distance r_{\perp} ,
- should be unitarized at large r_{\perp} (up to log effects)

$$\sigma_{\text{dipole}}(x, r_{\perp}) = 2 \int d^2b \mathcal{N}(x, r_{\perp}, b_{\perp}) < \sigma_0$$

- Parametrizations of σ_{dipole} with saturation models work relatively well to describe the HERA data.

\Rightarrow Aim: Provide a simple parametrization rooted in QCD

Previous attempts with saturation models

Golec-Biernat–Wüsthoff model [PRD59(99)014017,60(99)114023]

$$\sigma_{\text{dipole}}(x, r_{\perp}) = \sigma_0 \left[1 - \exp \left\{ -\frac{1}{4} r_{\perp}^2 Q_s^2(x) \right\} \right]$$

$$Q_s^2(x) = 1 \text{ GeV}^2 \left(\frac{x_0}{x} \right)^{\lambda}$$

- **Saturation (unitarization):** $\sigma_{\text{dipole}} \rightarrow \sigma_0$ as $r_{\perp} \rightarrow \infty$
- **Geometric scaling:** $\sigma_{\text{dipole}}(x, r_{\perp}) = \sigma_{\text{dipole}}(r_{\perp}^2 Q_s^2(x))$

⇒ Good fit to old HERA data (F_2, F_2^{Diff}) for $x < 10^{-2}$

[H1: NPB470(96)3,497(97)3, ZEUS:Z.P.C72(96)399,EPJ.C7(99)609]

3 parameters: $\sigma_0 = 23\text{mb}$, $x_0 = 3 \times 10^{-4}$, $\lambda \simeq 0.3$

But, doesn't work well for new HERA data at high Q^2

[H1: EPJ.C21(01)33, ZEUS:PLB487(00)53,EPJ.C21(01)443]

Improvement of the GBW model

- by adding DGLAP evolution at high Q^2 (small r_{\perp})


Bartels, Golec-Biernat, Kowalski, PRD66(02)014001

- by further adding impact parameter (b) dependence

Kowalski& Teaney, PRD68(03)114005

Physics at $x < 0.01$, $Q^2 \lesssim 50\text{GeV}^2$

But, **BFKL**, rather than DGLAP, should be the right physics in the transition regime!
 We focus on the region with not too large Q^2 .



CGC (saturation) for $Q^2 < Q_s^2(x)$

BFKL dynamics in $Q_s^2 < Q^2 < Q_s^4/\Lambda^2 \sim 50\text{GeV}^2$

Physics at $x < 0.01$, $Q^2 \lesssim 50\text{GeV}^2$

Saturation for $Q^2 < Q_s^2(x)$ [Levin-Tuchin, Iancu-McLerran]

A solution to BK and JIMWLK equations:

Scattering amplitude \mathcal{N} : $\sigma_{\text{dipole}} = 2\pi R^2 \mathcal{N}(x, r_\perp)$

$$\mathcal{N}(x, r_\perp) \simeq 1 - \exp \left\{ -\frac{1}{2c} \ln^2 r_\perp^2 Q_s^2(x) \right\}$$

coherent multiple scatterings (\neq Glauber with indep. scat.)

BFKL dynamics in $Q_s^2 < Q^2 < Q_s^4/\Lambda^2 \sim 50\text{GeV}^2$

[Iancu-KI-McLerran, Mueller-Triantafyllopoulos]

saddle point approx. + saturation boundary

$$\mathcal{N}(x, r_\perp) \simeq \underbrace{(r_\perp^2 Q_s^2)^\gamma}_{\text{geometric scaling}} \underbrace{\exp \left\{ -\frac{1}{2\kappa\lambda Y} \left(\ln \frac{1}{r_\perp^2 Q_s^2} \right)^2 \right\}}_{\text{diffusion, scaling violation}}$$

- $\gamma \simeq 0.63 < 1$: anomalous dimension
- $\kappa = \chi''(\gamma)/\chi'(\gamma) \simeq 9.9$: diffusion coefficient
- λ : saturation exponent $Q_s^2(Y) \propto e^{\lambda Y}$ ($Y = \ln 1/x$)

Diffusion effectively increases the anomalous dimension.

Data very sensitive to the precise value of λ

A CGC fit to the HERA F_2 data

Iancu, KI, Munier, hep-ph/0310338

$$\sigma_{\text{dipole}}(x, \mathbf{r}) = 2\pi R^2 \mathcal{N}(rQ_s, Y), \quad Q_s^2(x) = (x_0/x)^\lambda \text{GeV}^2,$$

$$\mathcal{N}(rQ_s, Y) = \mathcal{N}_0 \left(\frac{r^2 Q_s^2}{4} \right)^{\gamma + \frac{\ln(2/rQ_s)}{\kappa \lambda Y}} \quad \text{for } rQ_s \leq 2$$

$$\mathcal{N}(rQ_s, Y) = 1 - e^{-a \ln^2(b r Q_s)} \quad \text{for } rQ_s > 2$$

$\gamma = 0.63$, $\kappa = 9.9$ (fixed by LO-BFKL \simeq resummed NLO)

a, b fixed by continuity at $rQ_s = 2$ where $\mathcal{N} = \mathcal{N}_0$.

3 parameters R, x_0 and λ (The same as in GBW. $\sigma_0 = 2\pi R^2$)

R, x_0 : non-perturbative.

λ : saturation exponent. Perturbatively computable,
but left free to account for theoretical uncertainty.


Resummed NLO: $\lambda \sim 0.3 \pm 15\%$ [Triantafyllopoulos]

Fit to the new data within

$$x \leq 10^{-2} \text{ and } 0.045 \leq Q^2 \leq 45 \text{GeV}^2$$

A CGC fit to the HERA F_2 data

Iancu, KI, Munier, hep-ph/0310338



F_2 as a function of x in bins of $Q^2 \leq 15 \text{ GeV}^2$.

red line: CGC with $N_0 = 0.7$ and $m_q = 140 \text{ MeV}$


blue line: BFKL without saturation

$R = 0.641 \text{ fm}$, $x_0 = 0.267 \times 10^{-4}$, $\lambda = 0.253$

with $\chi^2/\text{dof} = 0.81$ (156 data points)

A CGC fit to the HERA F_2 data

Iancu, KI, Munier, hep-ph/0310338



Larger Q^2 : (The fit is extrapolated in bins $Q^2 \geq 60\text{GeV}^2$.)

- $\lambda \sim 0.25$ is in agreement with the resummed NLO BFKL result
- Scaling violation is essential (pure scaling fit doesn't work well)
- BFKL w/o saturation cannot reproduce the low Q^2 data
- Good agreement even for $Q^2 \ll 1 \text{ GeV}^2$ (quark-hadron duality)
- Deviation at high Q^2 : no DGLAP, no valence quarks

Summary

- Presented a new **analytic** parametrization of the dipole cross section which is **rooted in QCD**.

BFKL dynamics near saturation

- **geometric scaling** as leading effect
- its **violation** by the diffusion term at high Q^2

CGC, saturation

- correct asymptotic behaviour
(from BK or JIMWLK eqs.)

- This works very well in describing **HERA $F_2(x, Q^2)$ data** within $x \leq 10^{-2}$ and $0.045 \leq Q^2 \leq 45 \text{GeV}^2$.
3 parameter fit (R , x_0 and λ) gives $\chi^2/dof \sim 0.8$.
- Our analysis suggests that the CGC may have been seen in the HERA DIS data at small x .

The CGC fit

$\mathcal{N}_0/\text{model}$	0.5	0.6	0.7	0.8	0.9	GBW
χ^2	146.43	129.88	123.63	125.61	133.73	243.87
$\chi^2/\text{d.o.f}$	0.96	0.85	0.81	0.82	0.87	1.59
$x_0 (\times 10^{-4})$	0.669	0.435	0.267	0.171	0.108	4.45
λ	0.252	0.254	0.253	0.252	0.250	0.286
R (fm)	0.692	0.660	0.641	0.627	0.618	0.585

Table 1: The CGC fits for different values of \mathcal{N}_0 and 3 quark flavors with mass $m_q = 140$ MeV. Also shown is the fit obtained by using the GBW model.

The CGC fit



$\mathcal{N}_0/\text{model}$	$m_q = 50 \text{ MeV}$			$m_q = 10 \text{ MeV}$		
	0.5	0.7	0.9	0.5	0.7	0.9
χ^2	148.02	108.52	108.76	149.27	107.64	106.49
$\chi^2/\text{d.o.f}$	0.97	0.71	0.71	0.98	0.70	0.70
$x_0 (\times 10^{-4})$	2.77	0.898	0.333	3.32	1.06	0.382
λ	0.290	0.281	0.274	0.295	0.285	0.276
$R \text{ (fm)}$	0.604	0.574	0.561	0.593	0.566	0.554

Table 2: The CGC fits for three values of \mathcal{N}_0 and quark masses $m_q = 50 \text{ MeV}$ (left) and $m_q = 10 \text{ MeV}$ (right).

\mathcal{N}_0	χ^2	$\chi^2/\text{d.o.f}$	$x_0 (\times 10^{-4})$	λ	$R \text{ (fm)}$	γ
0.7	215.70	1.42	3.79	0.313	0.572	0.845

Table 3: A 4 parameter fit for geometric scaling + saturation.

The CGC fit



The CGC fit

