Heavy Bosons Other Than Higgs Bosons, Searches for

We list here various limits on charged and neutral heavy vector bosons (other than W's and Z's), heavy scalar bosons (other than Higgs bosons), vector or scalar leptoquarks, and axigluons.

W_R (Right-Handed W Boson) MASS LIMITS

Assuming a light right-handed neutrino, except for BEALL 82, LANGACKER 89B, and COLANGELO 91. $g_R = g_L$ assumed. [Limits in the section MASS LIMITS for W' below are also valid for W_R if $m_{\nu_R} \ll m_{W_R}$.] Some limits assume manifest left-right symmetry, *i.e.*, the equality of left- and right Cabibbo-Kobayashi-Maskawa matrices. For a comprehensive review, see LANGACKER 89B. Limits on the W_L-W_R mixing angle ζ are found in the next section. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
> 549		[1] BARENBOIM	97	RVUE	μ decay
• • • We do not use the following data for averages, fits, limits, etc. • • •					
> 220	95	[2] STAHL	97	RVUE	τ decay
> 220	90	[3] ALLET	96	CNTR	β^+ decay
> 281	90	[4] KUZNETSOV	95	CNTR	Polarized neutron decay
> 282	90	[5] KUZNETSOV	94B	CNTR	Polarized neutron decay
> 439	90	[6] BHATTACH...	93	RVUE	Z-Z' mixing
> 250	90	[7] SEVERIJNS	93	CNTR	β^+ decay
		[8] IMAZATO	92	CNTR	K^+ decay
> 475	90	[9] POLAK	92B	RVUE	μ decay
> 240	90	[10] AQUINO	91	RVUE	Neutron decay
> 496	90	[10] AQUINO	91	RVUE	Neutron and muon decay
> 700		[11] COLANGELO	91	THEO	$m_{K^0_L} - m_{K^0_S}$
> 477	90	[12] POLAK	91	RVUE	μ decay
[none 540–23000]		[13] BARBIERI	89B	ASTR	SN 1987A; light ν_R
> 300	90	[14] LANGACKER	89B	RVUE	General
> 160	90	[15] BALKE	88	CNTR	$\mu \rightarrow e\nu\overline{\nu}$
> 406	90	[16] JODIDIO	86	ELEC	Any ζ
> 482	90	[16] JODIDIO	86	ELEC	$\zeta = 0$
> 800		MOHAPATRA	86	RVUE	$SU(2)_L \times SU(2)_R \times U(1)$
> 400	95	[17] STOKER	85	ELEC	Any ζ
> 475	95	[17] STOKER	85	ELEC	$\zeta < 0.041$
		[18] BERGSMA	83	CHRM	$\nu_\mu e \rightarrow \mu\nu_e$
> 380	90	[19] CARR	83	ELEC	μ^+ decay
>1600		[20] BEALL	82	THEO	$m_{K^0_L} - m_{K^0_S}$
[> 4000]		STEIGMAN	79	COSM	Nucleosynthesis; light ν_R

[1] The quoted limit is from μ decay parameters. BARENBOIM 97 also evaluate limit from K_L-K_S mass difference.

[2] STAHL 97 limit is from fit to τ-decay parameters.

[3] ALLET 96 measured polarization-asymmetry correlaton in $^{12}\mathrm{N}\,\beta^+$ decay. The listed limit assumes zero L-R mixing.

[4] KUZNETSOV 95 limit is from measurements of the asymmetry $\langle \vec{p}_\nu \cdot \sigma_n \rangle$ in the β decay of polarized neutrons. Zero mixing assumed. See also KUZNETSOV 94B.

[5] KUZNETSOV 94B limit is from measurements of the asymmetry $\langle \vec{p}_\nu \cdot \sigma_n \rangle$ in the β decay of polarized neutrons. Zero mixing assumed.

[6] BHATTACHARYYA 93 uses Z-Z' mixing limit from LEP '90 data, assuming a specific Higgs sector of $\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R \times \mathrm{U}(1)$ gauge model. The limit is for m_t=200 GeV and slightly improves for smaller m_t.

[7] SEVERIJNS 93 measured polarization-asymmetry correlation in $^{107}\mathrm{In}\,\beta^+$ decay. The listed limit assumes zero L-R mixing. Value quoted here is from SEVERIJNS 94 erratum.

[8] IMAZATO 92 measure positron asymmetry in $K^+ \rightarrow \mu^+ \nu_\mu$ decay and obtain $\xi P_\mu > 0.990$ (90%CL). If W_R couples to $u\overline{s}$ with full weak strength (V^R_{us}=1), the result corresponds to $m_{W_R} >653$ GeV. See their Fig. 4 for m_{W_R} limits for general $|V^R_{us}|^2 = 1 - |V^R_{ud}|^2$.

[9] POLAK 92B limit is from fit to muon decay parameters and is essentially determined by JODIDIO 86 data assuming ζ=0. Supersedes POLAK 91.

[10] AQUINO 91 limits obtained from neutron lifetime and asymmetries together with unitarity of the CKM matrix. Manifest left-right symmetry assumed. Stronger of the two limits also includes muon decay results.

[11] COLANGELO 91 limit uses hadronic matrix elements evaluated by QCD sum rule and is less restrictive than BEALL 82 limit which uses vacuum saturation approximation. Manifest left-right symmetry assumed.

[12] POLAK 91 limit is from fit to muon decay parameters and is essentially determined by JODIDIO 86 data assuming ζ=0. Superseded by POLAK 92B.

[13] BARBIERI 89B limit holds for $m_{\nu_R} \leq 10$ MeV.

[14] LANGACKER 89B limit is for any ν_R mass (either Dirac or Majorana) and for a general class of right-handed quark mixing matrices.

[15] BALKE 88 limit is for $m_{\nu_{eR}} = 0$ and $m_{\nu_{\mu R}} \leq 50$ MeV. Limits come from precise measurements of the muon decay asymmetry as a function of the positron energy.

[16] JODIDIO 86 is the same TRIUMF experiment as STOKER 85 (and CARR 83); however, it uses a different technique. The results given here are combined results of the two techniques. The technique here involves precise measurement of the end-point e^+ spectrum in the decay of the highly polarized μ^+.

[17] STOKER 85 is same TRIUMF experiment as CARR 83. Here they measure the decay e^+ spectrum asymmetry above 46 MeV/c using a muon-spin-rotation technique. Assumed a light right-handed neutrino. Quoted limits are from combining with CARR 83.

[18] BERGSMA 83 set limit $m_{W_2}/m_{W_1} >1.9$ at CL = 90%.

[19] CARR 83 is TRIUMF experiment with a highly polarized μ^+ beam. Looked for deviation from $V-A$ at the high momentum end of the decay e^+ energy spectrum. Limit from previous world-average muon polarization parameter is $m_{W_R} >240$ GeV. Assumes a light right-handed neutrino.

[20] BEALL 82 limit is obtained assuming that W_R contribution to K^0_L–K^0_S mass difference is smaller than the standard one, neglecting the top quark contributions. Manifest left-right symmetry assumed.

Limit on W_L-W_R Mixing Angle ζ

Lighter mass eigenstate $W_1 = W_L\cos\zeta - W_R\sin\zeta$. Light ν_R assumed unless noted. Values in brackets are from cosmological and astrophysical considerations.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<$ 0.0333		[21] BARENBOIM	97	RVUE	μ decay
$<$ 0.04	90	[22] MISHRA	92	CCFR	νN scattering
-0.0006 to 0.0028	90	[23] AQUINO	91	RVUE	
[none 0.00001–0.02]		[24] BARBIERI	89B	ASTR	SN 1987A
$<$ 0.040	90	[25] JODIDIO	86	ELEC	μ decay
-0.056 to 0.040	90	[25] JODIDIO	86	ELEC	μ decay

[21] The quoted limit is from μ decay parameters. BARENBOIM 97 also evaluate limit from K_L-K_S mass difference.

[22] MISHRA 92 limit is from the absence of extra large-x, large-y $\overline{\nu}_\mu N \rightarrow \overline{\nu}_\mu X$ events at Tevatron, assuming left-handed ν and right-handed $\overline{\nu}$ in the neutrino beam. The result gives $\zeta^2(1-2m^2_{W_1}/m^2_{W_2}) < 0.0015$. The limit is independent of ν_R mass.

[23] AQUINO 91 limits obtained from neutron lifetime and asymmetries together with unitarity of the CKM matrix. Manifest left-right asymmetry is assumed.

[24] BARBIERI 89B limit holds for $m_{\nu_R} \leq 10$ MeV.

[25] First JODIDIO 86 result assumes $m_{W_R}=\infty$, second is for unconstrained m_{W_R}.

A REVIEW GOES HERE – Check our WWW List of Reviews

MASS LIMITS for W' (A Heavy-Charged Vector Boson Other Than W) in Hadron Collider Experiments

Couplings of W' to quarks and leptons are taken to be identical with those of W. The following limits are obtained from $p\overline{p} \rightarrow W' X$ with W' decaying to the mode indicated in the comments. New decay channels (*e.g.*, $W' \rightarrow WZ$) are assumed to be suppressed. UA1 and UA2 experiments assume that the $t\overline{b}$ channel is not open.

VALUE (GeV)	CL%	DOCUMENT ID		TECN	COMMENT
>720	95	[26] ABACHI	96C	D0	$W' \rightarrow e\nu_e$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
none 300–420	95	[27] ABE	97G	CDF	$W' \rightarrow q\overline{q}$
>610	95	[28] ABACHI	95E	D0	$W' \rightarrow e\nu_e$ and $W' \rightarrow \tau\nu_\tau \rightarrow e\nu\nu\overline{\nu}$
>652	95	[29] ABE	95M	CDF	$W' \rightarrow e\nu_e$
>251	90	[30] ALITTI	93	UA2	$W' \rightarrow q\overline{q}$
none 260–600	95	[31] RIZZO	93	RVUE	$W' \rightarrow q\overline{q}$
>520	95	[32] ABE	91F	CDF	$W' \rightarrow e\nu$, $\mu\nu$
none 101–158	90	[33] ALITTI	91	UA2	$W' \rightarrow q\overline{q}$
>220	90	[34] ALBAJAR	89	UA1	$W' \rightarrow e\nu$
>209	90	[35] ANSARI	87D	UA2	$W' \rightarrow e\nu$
>210	90	[36] ARNISON	86B	UA1	$W' \rightarrow e\nu$
>170	90	[37] ARNISON	83D	UA1	$W' \rightarrow e\nu$

[26] For bounds on W_R with nonzero right-handed mass, see Fig. 5 from ABACHI 96C.

[27] ABE 97G search for new particle decaying to dijets.

[28] ABACHI 95E assume that the decay $W' \rightarrow WZ$ is suppressed and that the neutrino from W' decay is stable and has a mass significantly less $m_{W'}$.

[29] ABE 95M assume that the decay $W' \rightarrow WZ$ is suppressed and the (right-handed) neutrino is light, noninteracting, and stable. If m_ν=60 GeV, for example, the effect on the mass limit is neglibible.

[30] ALITTI 93 search for resonances in the two-jet invariant mass. The limit assumes $\Gamma(W')/m_{W'} = \Gamma(W)/m_W$ and $B(W' \rightarrow jj) = 2/3$. This corresponds to W_R with $m_{\nu_R} > m_{W_R}$ (no leptonic decay) and $W_R \rightarrow t\overline{b}$ allowed. See their Fig. 4 for limits in the $m_{W'}$–$B(q\overline{q})$ plane.

[31] RIZZO 93 analyses CDF limit on possible two-jet resonances. The limit is sensitive to the inclusion of the assumed K factor.

[32] ABE 91F assume leptonic branching ratio of 1/12 for each lepton flavor. The limit from the $e\nu$ ($\mu\nu$) mode alone is 490 (435) GeV. These limits apply to W_R if $m_{\nu_R} \lesssim 15$ GeV and ν_R does not decay in the detector. Cross section limit $\sigma \cdot B <$ (1–10) pb is given for $m_{W'}$ = 100–550 GeV; see Fig. 2.

[33] ALITTI 91 search is based on two-jet invariant mass spectrum, assuming $B(W' \rightarrow q\overline{q})$ = 67.6%. Limit on $\sigma \cdot B$ as a function of two-jet mass is given in Fig. 7.

[34] ALBAJAR 89 cross section limit at 630 GeV is $\sigma(W')$ $B(e\nu) <$ 4.1 pb (90% CL).

[35] See Fig. 5 of ANSARI 87D for the excluded region in the $m_{W'}$–$[(g_{W'q})^2$ $B(W' \rightarrow e\overline{\nu})]$ plane. Note that the quantity $(g_{W'q})^2$ $B(W' \rightarrow e\overline{\nu})$ is normalized to unity for the standard W couplings.

[36] ARNISON 86B find no excess at large p_T in 148 $W \rightarrow e\nu$ events. Set limit $\sigma \times B(e\nu)$ <10 pb at CL = 90% at E_{cm} = 546 and 630 GeV.

[37] ARNISON 83D find among 47 $W \rightarrow e\nu$ candidates no event with excess p_T. Also set $\sigma \times B(e\nu)$ <30 pb with CL = 90% at E_{cm} = 540 GeV.

MASS LIMITS for Z' (Heavy Neutral Vector Boson Other Than Z)

A REVIEW GOES HERE – Check our WWW List of Reviews

Limits for Z'_{SM}

Z'_{SM} is assumed to have couplings with quarks and leptons which are identical to those of Z.

VALUE (GeV)	CL%	DOCUMENT ID		TECN	COMMENT
>990	95	[38] CVETIC	98	RVUE	Electroweak
>690	95	[39] ABE	97S	CDF	$p\overline{p}$; $Z'_{SM} \rightarrow e^+e^-$, $\mu^+\mu^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>490	95	ABACHI	96D	D0	$p\overline{p}$; $Z'_{SM} \rightarrow e^+e^-$
>505	95	[40] ABE	95	CDF	$p\overline{p}$; $Z'_{SM} \rightarrow e^+e^-$
>398	95	[41] VILAIN	94B	CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\overline{\nu}_\mu e \rightarrow \overline{\nu}_\mu e$
>237	90	[42] ALITTI	93	UA2	$p\overline{p}$; $Z'_{SM} \rightarrow q\overline{q}$
>119	90	[43] ALLEN	93	CALO	$\nu e \rightarrow \nu e$
none 490–560	95	[44] RIZZO	93	RVUE	$p\overline{p}$; $Z'_{SM} \rightarrow q\overline{q}$
>412	95	ABE	92B	CDF	$p\overline{p}$; $Z'_{SM} \rightarrow e^+e^-$, $\mu^+\mu^-$
>779	95	[45,46] LANGACKER	92B	RVUE	Repl. by CVETIC 98

>387	95	[47] ABE	91D	CDF	$p\overline{p}$; $Z'_{SM} \rightarrow e^+e^-$
>307	90	[48] GEIREGAT	91	CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\overline{\nu}_\mu e \rightarrow \overline{\nu}_\mu e$
>426	90	[49] ABE	90F	VNS	e^+e^-
>208	90	[50] HAGIWARA	90	RVUE	e^+e^-
>173	90	[51] ALBAJAR	89	UA1	$p\overline{p}$; $Z'_{SM} \rightarrow e^+e^-$
>180	90	[52] ANSARI	87D	UA2	$p\overline{p}$; $Z'_{SM} \rightarrow e^+e^-$
>160	90	[53] ARNISON	86B	UA1	$p\overline{p}$; $Z'_{SM} \rightarrow e^+e^-$

[38] CVETIC 98 give 95%CL limit on the Z-Z' mixing $-0.0023 < \theta < 0.0004$.

[39] ABE 97S limit is obtained assuming that Z' decays to known fermions only.

[40] ABE 95 limit is obtained assuming that Z' decays to known fermions only.

[41] VILAIN 94B assume $m_t = 150$ GeV.

[42] ALITTI 93 search for resonances in the two-jet invariant mass. The limit assumes B($Z' \rightarrow q\overline{q}$)=0.7. See their Fig. 5 for limits in the $m_{Z'}$–B($q\overline{q}$) plane.

[43] ALLEN 93 limit is from total cross section for $\nu e \rightarrow \nu e$, where $\nu = \nu_e, \nu_\mu, \overline{\nu}_\mu$.

[44] RIZZO 93 analyses CDF limit on possible two-jet resonances. The limit is sensitive to the inclusion of the assumed K factor.

[45] LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. $m_t >$89 GeV used.

[46] LANGACKER 92B give 95%CL limits on the Z-Z' mixing $-0.0086 < \theta < 0.0005$.

[47] ABE 91D give $\sigma(Z')\cdot$B(e^+e^-) $<$ 1.31 pb (95%CL) for $m_{Z'} >$ 200 GeV at E_{cm} = 1.8 TeV. Limits ranging from 2 to 30 pb are given for $m_{Z'}$ = 100–200 GeV.

[48] GEIREGAT 91 limit is from comparison of g_V^e from $\nu_\mu e$ scattering with $\Gamma(Z \rightarrow ee)$ from LEP. Zero mixing assumed.

[49] ABE 90F use data for R, $R_{\ell\ell}$, and $A_{\ell\ell}$. They fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.

[50] HAGIWARA 90 perform a fit to e^+e^- data at PEP, PETRA, and TRISTAN including $\mu^+\mu^-$, $\tau^+\tau^-$, and hadron cross sections and asymmetries.

[51] ALBAJAR 89 cross section limit at 630 GeV is $\sigma(Z')$ B(ee) $<$ 4.2 pb (90% CL).

[52] See Fig. 5 of ANSARI 87D for the excluded region in the $m_{Z'}$–$[(g_{Z'q})^2$ B($Z' \rightarrow e^+e^-$)] plane. Note that the quantity $(g_{Z'q})^2$B($Z' \rightarrow e^+e^-$) is normalized to unity for the standard Z couplings.

[53] ARNISON 86B find no excess e^+e^- pairs among 13 pairs from Z. Set limit $\sigma\times$B(e^+e^-) <13 pb at CL = 90% at E_{cm} = 546 and 630 GeV.

Limits for Z_{LR}

Z_{LR} is the extra neutral boson in left-right symmetric models. $g_L = g_R$ is assumed unless noted. Values in parentheses assume stronger constraint on the Higgs sector, usually motivated by superstring models. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
>1700	68	[54] BARENBOIM	98	RVUE	Z parameters
> 390	95	[55] CVETIC	98	RVUE	Electroweak
> 630	95	[56] ABE	97S	CDF	$p\overline{p}$; $Z'_{LR} \rightarrow e^+e^-$, $\mu^+\mu^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>190	95	[57] BARATE	97B	ALEP	$e^+e^- \rightarrow \mu^+\mu^-$ and hadronic cross section
>445	95	[58] ABE	95	CDF	$p\overline{p}$; $Z_{LR} \rightarrow e^+e^-$
>253	95	[59] VILAIN	94B	CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\overline{\nu}_\mu e \rightarrow \overline{\nu}_\mu e$
>130	95	[60] ADRIANI	93D	L3	Z parameters
(>1500)	90	[61] ALTARELLI	93B	RVUE	Z parameters
none 490–560	95	[62] RIZZO	93	RVUE	$p\overline{p}$; $Z_{LR} \rightarrow q\overline{q}$
>310	95	[63] ABE	92B	CDF	$p\overline{p}$
>230	95	[64] ABE	92B	CDF	$p\overline{p}$
(>900)	90	[65] DELAGUILA	92	RVUE	
>389	95	[66,67] LANGACKER	92B	RVUE	Repl. by CVETIC 98
(>1400)		[68] LAYSSAC	92B	RVUE	Z parameters
(>564)	90	[69] POLAK	92	RVUE	μ decay
>474	90	[70] POLAK	92B	RVUE	Electroweak
(>1340)		[71] RENTON	92	RVUE	
(>800)	90	[72] ALTARELLI	91B	RVUE	Z parameters
(>795)	90	[73] DELAGUILA	91	RVUE	
>382	90	[74] POLAK	91	RVUE	Electroweak
$[>2000]$		WALKER	91	COSM	Nucleosynthesis; light ν_R
$[>500]$		[75] GRIFOLS	90	ASTR	SN 1987A; light ν_R
(>460)	90	[76] HE	90B	RVUE	
$[>$ 2400–6800$]$		[77] BARBIERI	89B	ASTR	SN 1987A; light ν_R
>189		[78] DELAGUILA	89	RVUE	$p\overline{p}$
$[>10000]$		RAFFELT	88	ASTR	SN 1987A; light ν_R
>325	90	[79] AMALDI	87	RVUE	
>278	90	[80] DURKIN	86	RVUE	
>150	95	[81] ADEVA	85B	MRKJ	$e^+e^- \rightarrow \mu^+\mu^-$

[54] BARENBOIM 98 also gives 68% CL limits on the Z-Z's mixing $-5.1\times10^{-4} < \theta < 3.3\times10^{-3}$.

[55] CVETIC 98 give 95%CL limit on the Z-Z' mixing $-0.0013 < \theta < 0.0021$.

[56] ABE 97S limit is obtained assuming that Z' decays to known fermions only.

[57] BARATE 97B gives 95% CL limits on Z-Z' mixing $-0.0017 < \theta < 0.0035$. The bounds are computed with $\alpha_s = 0.120 \pm 0.003$, $m_t = 175 \pm 6$ GeV, and $M_H = 150^{+120}_{-\ 90}$ GeV. See their Fig. 4 for the limit contour in the mass-mixing plane.

[58] ABE 95 limit is obtained assuming that Z' decays to known fermions only. See their Fig. 3 for the mass bound of Z' decaying to all allowed fermions and supersymmetric fermions.

[59] VILAIN 94B assume m_t = 150 GeV and θ=0. See Fig. 2 for limit contours in the mass-mixing plane.

[60] ADRIANI 93D give limits on the Z-Z' mixing $-0.002 < \theta < 0.015$ assuming the ABE 92B mass limit.

[61] ALTARELLI 93B limit is from LEP data available in summer '93 and is for m_t = 110 GeV. m_H = 100 GeV and α_s = 0.118 assumed. The limit improves for larger m_t (see their Fig. 5). The 90%CL limit on the Z-Z' mixing angle is in Table 4.

[62] RIZZO 93 analyses CDF limit on possible two-jet resonances. The limit is sensitive to the inclusion of the assumed K factor.

[63] These limits assume that Z' decays to known fermions only.

[64] These limits assume that Z' decays to all E_6 fermions and their superpartners.

[65] See Fig. 7b and 8 in DELAGUILA 92 for the allowed region in $m_{Z'}$-mixing plane and $m_{Z'} - m_t$ plane from electroweak fit including '90 LEP data.

[66] LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. m_t >89 GeV used.

[67] LANGACKER 92B give 95%CL limits on the Z-Z' mixing $-0.0025 < \theta < 0.0083$.

[68] LAYSSAC 92B limit is from LEP data available spring '92. Specific Higgs sector is assumed. See also LAYSSAC 92.

[69] POLAK 92 limit is from m_{W_R} >477 GeV, which is derived from muon decay parameters assuming light ν_R. Specific Higgs sector is assumed.

[70] POLAK 92B limit is from a simultaneous fit to charged and neutral sector in $SU(2)_L \times SU(2)_R \times U(1)$ model using Z parameters, m_W, and low-energy neutral current data as of 1991. Light ν_R assumed and m_t=m_H=100 GeV used. Supersedes POLAK 91.

[71] RENTON 92 limits use LEP data taken up to '90 as well as m_W, νN, and atomic parity violation data. Specific Higgs structure is assumed.

[72] ALTARELLI 91B is based on Z mass, widths, and A_{FB}. The limits are for superstring motivated models with extra assumption on the Higgs sector. $m_t > 90$ GeV and $m_{H^0} < 1$ TeV assumed. For large m_t, the bound improves drastically. Bounds for Z-Z' mixing angle and Z mass shift without this model assumption are also given in the paper.

[73] DELAGUILA 91 bounds have extra assumption of superstring motivated Higgs sector. From νN neutral current data with $m_Z = 91.10 \pm 0.04$ GeV, $m_t > 77$ GeV, $m_{H^0} < 1$ TeV assumed.

[74] POLAK 91 limit is from a simultaneous fit to charged and neutral sector in $SU(2)_L \times SU(2)_R \times U(1)$ model using m_W, m_Z, and low-energy neutral current data as of 1990. Light ν_R assumed and m_t=m_H=100 GeV used. Superseded by POLAK 92B.

[75] GRIFOLS 90 limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also GRIFOLS 90D, RIZZO 91.

[76] HE 90B model assumes a specific Higgs sector. Neutral current data of COSTA 88 as well as m_Z is used. g_R is left free in the fit.

[77] BARBIERI 89B limit holds for $m_{\nu_R} \leq 10$ MeV.

[78] DELAGUILA 89 limit is based on $\sigma(p\overline{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+ e^-) < 1.8$ pb at CERN $p\overline{p}$ collider.

[79] A wide range of neutral current data as of 1986 are used in the fit.

[80] A wide range of neutral current data as of 1985 are used in the fit.

[81] ADEVA 85B measure asymmetry of μ-pair production, following formalism of RIZZO 81.

Limits for Z_χ

Z_χ is the extra neutral boson in $SO(10) \rightarrow SU(5) \times U(1)_\chi$. $g_\chi = e/\cos\theta_W$ is assumed unless otherwise stated. We list limits with the assumption $\rho = 1$ but with no further constraints on the Higgs sector. Values in parentheses assume stronger constraint on the Higgs sector motivated by superstring models. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
>451	95	[82] CHO	98B	RVUE	Electroweak
>595	95	[83] ABE	97S	CDF	$p\overline{p}$; $Z'_\chi \rightarrow e^+ e^-, \mu^+ \mu^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>470	95	[84] CHO	98	RVUE	
>330	95	[85] CVETIC	98	RVUE	Electroweak
>190	95	[86] ARIMA	97	VNS	Bhabha scattering
>236	95	[87] BARATE	97B	ALEP	$e^+e^- \to \mu^+\mu^-$ and hadronic cross section
>196	95	[88] BUSKULIC	96N	ALEP	Hadronic cross section
>425	95	[89] ABE	95	CDF	$p\overline{p}$; $Z'_\chi \to e^+e^-$
>147	95	[90] ABREU	95M	DLPH	Z parameters and $e^+e^- \to \mu^+\mu^-(n\gamma)$
		[91] NARDI	95	RVUE	Z parameters
		[92] BUSKULIC	94	ALEP	Z parameters
>262	95	[93] VILAIN	94B	CHM2	$\nu_\mu e \to \nu_\mu e$ and $\overline{\nu}_\mu e \to \overline{\nu}_\mu e$
>117	95	[94] ADRIANI	93D	L3	Z parameters
(>900)	90	[95] ALTARELLI	93B	RVUE	Z parameters
>340	95	[96] ABE	92B	CDF	$p\overline{p}$
>280	95	[97] ABE	92B	CDF	$p\overline{p}$
(>650)	90	[98] DELAGUILA	92	RVUE	
>321	95	[99,100] LANGACKER	92B	RVUE	Repl. by CVETIC 98
(>760)		[101] LAYSSAC	92B	RVUE	Z parameters
>148	95	[102] LEIKE	92	RVUE	Z parameters
(>700)		[103] RENTON	92	RVUE	
(> 500)	90	[104] ALTARELLI	91B	RVUE	Z parameters
(> 570)		[105] BUCHMUEL...	91	RVUE	Z parameters
(> 555)	90	[106] DELAGUILA	91	RVUE	
[>1470]		[107] FARAGGI	91	COSM	Nucleosynthesis; light ν_R
>320	90	[108] GONZALEZ-G..	91	RVUE	
>221		[109] MAHANTHAP..	91	RVUE	Cs
>231	90	[110,111] ABE	90F	VNS	e^+e^-
>206	90	[111,112] ABE	90F	RVUE	e^+e^-, $\nu_\mu e$
>335		[113] BARGER	90B	RVUE	$p\overline{p}$
(> 650)	90	[114] GLASHOW	90	RVUE	
[> 1140]		[115] GONZALEZ-G..	90D	COSM	Nucleosynthesis; light ν_R
[> 2100]		[116] GRIFOLS	90	ASTR	SN 1987A; light ν_R
none <150 or > 363	90	[117] HAGIWARA	90	RVUE	e^+e^-
>177		[118] DELAGUILA	89	RVUE	$p\overline{p}$
>280	95	[119] DORENBOS...	89	CHRM	$g_\chi = g_Z$
>352	90	[120] COSTA	88	RVUE	
>170	90	[121] ELLIS	88	RVUE	$p\overline{p}$
>273	90	[120] AMALDI	87	RVUE	
>266	90	[122] MARCIANO	87	RVUE	
>283	90	[123] DURKIN	86	RVUE	

[82] CHO 98B use various electroweak data to constrain Z' models. See their Eq. (4.8) for their fit in mass-mixing plane.

[83] ABE 97S limit is obtained assuming that Z' decays to known fermions only.

[84] CHO 98 limit is from constraints on four-Fermi contact interactions obrained from low-energy electroweak experiments.

[85] CVETIC 98 give 95%CL limit on the Z-Z' mixing $-0.0029 < \theta < 0.0011$.

[86] Z-Z' mixing is assumed to be zero.

[87] BARATE 97B gives 95% CL limits on Z-Z' mixing $-0.0016 < \theta < 0.0036$. The bounds are computed with $\alpha_s = 0.120 \pm 0.003$, $m_t = 175 \pm 6$ GeV, and $M_H = 150^{+120}_{-\ 90}$ GeV. See their Fig. 4 for the limit contour in the mass-mixing plane.

[88] BUSKULIC 96N limit is from a combined fit to the hadronic cross sections measured at $\sqrt{s}$=130, 136 GeV (ALEPH) and $\sqrt{s}$=58 GeV (TOPAZ). Zero mixing is assumed.

[89] ABE 95 limit is obtained assuming that Z' decays to known fermions only. See their Fig. 3 for the mass bound of Z' decaying to all allowed fermions and supersymmetric fermions.

[90] ABREU 95M limit is for α_s=0.123, m_t=150 GeV, and m_H=300 GeV. For the limit contour in the mass-mixing plane, see their Fig. 13.

[91] NARDI 95 give 90%CL limits on Z-Z' mixing $-0.0032 < \theta < 0.0031$ for $M_{Z'}$ >500 GeV, m_t=170 GeV, m_H=250 GeV, α_s=0.12. The bound is relaxed under the simultaneous presence of the mixing of the known fermions with new heavy states, $-0.0032 < \theta <$ 0.0079.

[92] BUSKULIC 94 give 95%CL limits on the Z-Z' mixing $-0.0091 < \theta < 0.0023$.

[93] VILAIN 94B assume m_t = 150 GeV and θ=0. See Fig. 2 for limit contours in the mass-mixing plane.

[94] ADRIANI 93D give limits on the Z-Z' mixing $-0.004 < \theta < 0.015$ assuming the ABE 92B mass limit.

[95] ALTARELLI 93B limit is from LEP data available in summer '93 and is for m_t = 110 GeV. m_H = 100 GeV and α_s = 0.118 assumed. The limit improves for larger m_t (see their Fig. 5). The 90%CL limit on the Z-Z' mixing angle is in their Fig. 2.

[96] These limits assume that Z' decays to known fermions only.

[97] These limits assume that Z' decays to all E_6 fermions and their superpartners.

[98] See Fig. 7a and 8 in DELAGUILA 92 for the allowed region in $m_{Z'}$-mixing plane and $m_{Z'} - m_t$ plane from electroweak fit including '90 LEP data.

[99] LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. m_t >89 GeV used.

[100] LANGACKER 92B give 95%CL limits on the Z-Z' mixing $-0.0048 < \theta < 0.0097$.

[101] LAYSSAC 92B limit is from LEP data available spring '92. Specific Higgs sector is assumed. See also LAYSSAC 92.

[102] LEIKE 92 is based on '90 LEP data published in LEP 92.

[103] RENTON 92 limits use LEP data taken up to '90 as well as m_W, νN, and atomic parity violation data. Specific Higgs structure is assumed.

[104] ALTARELLI 91B is based on Z mass, widths, and A_{FB}. The limits are for superstring motivated models with extra assumption on the Higgs sector. m_t > 90 GeV and m_{H^0} < 1 TeV assumed. For large m_t, the bound improves drastically. Bounds for Z-Z' mixing angle and Z mass shift without this model assumption are also given in the paper.

[105] BUCHMUELLER 91 limit is from LEP data. Specific assumption is made for the Higgs sector.

[106] DELAGUILA 91 bounds have extra assumption of superstring motivated Higgs sector. From νN neutral current data with $m_Z = 91.10 \pm 0.04$ GeV, m_t > 77 GeV, m_{H^0} < 1 TeV assumed.

[107] FARAGGI 91 limit assumes the nucleosynthesis bound on the effective number of neutrinos ΔN_ν < 0.5 and is valid for m_{ν_R} < 1 MeV.

[108] GONZALEZ-GARCIA 91 limit is based on low-energy neutral current data, Z mass and widths, m_W from ABE 90G. 100 < m_t < 200 GeV, m_{H^0} = 100 GeV assumed. Dependence on m_t is shown in Fig. 7.

[109] MAHANTHAPPA 91 limit is from atomic parity violation in Cs with m_W, m_Z.

[110] ABE 90F use data for R, $R_{\ell\ell}$, and $A_{\ell\ell}$.

[111] ABE 90F fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.

[112] e^+e^- data for R, $R_{\ell\ell}$, $A_{\ell\ell}$, and $A_{c\overline{c}}$ below Z as well as $\nu_\mu e$ scattering data of GEIREGAT 89 is used in the fit.
[113] BARGER 90B limit is based on CDF limit $\sigma(p\overline{p} \rightarrow Z')\cdot B(Z' \rightarrow e^+e^-) < 1$ pb (Nodulman, EPS Conf. '89). Assumes no new threshold is open for Z' decay.
[114] GLASHOW 90 model assumes a specific Higgs sector. See GLASHOW 90B.
[115] These authors claim that the nucleosynthesis bound on the effective number of light neutrinos ($\delta N_\nu < 1$) constrains Z' masses if ν_R is light ($\lesssim$ 1 MeV).
[116] GRIFOLS 90 limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also GRIFOLS 90D, RIZZO 91.
[117] HAGIWARA 90 perform a fit to e^+e^- data at PEP, PETRA, and TRISTAN including $\mu^+\mu^-$, $\tau^+\tau^-$, and hadron cross sections and asymmetries. The upper mass limit disappears at 2.7 s.d.
[118] DELAGUILA 89 limit is based on $\sigma(p\overline{p} \rightarrow Z')\cdot B(Z' \rightarrow e^+e^-) < 1.8$ pb at CERN $p\overline{p}$ collider.
[119] DORENBOSCH 89 obtain the limit $(g_\chi/g_Z)^2 \cdot (m_Z/m_{Z_\chi})^2 < 0.11$ at 95% CL from the processes $\overline{\nu}_\mu e \rightarrow \overline{\nu}_\mu e$ and $\nu_\mu e \rightarrow \nu_\mu e$.
[120] A wide range of neutral current data as of 1986 are used in the fit.
[121] Z' mass limits from non-observation of an excess of $\ell^+\ell^-$ pairs at the CERN $p\overline{p}$ collider [based on ANSARI 87D and GEER Uppsala Conf. 87]. The limits apply when Z' decays only into light quarks and leptons.
[122] MARCIANO 87 limit from unitarity of Cabibbo-Kobayashi-Maskawa matrix.
[123] A wide range of neutral current data as of 1985 are used in the fit.

Limits for Z_ψ

Z_ψ is the extra neutral boson in $E_6 \rightarrow SO(10) \times U(1)_\psi$. $g_\psi = e/\cos\theta_W$ is assumed unless otherwise stated. We list limits with the assumption $\rho = 1$ but with no further constraints on the Higgs sector. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
>170	95	[124] CVETIC	98	RVUE	Electroweak
>590	95	[125] ABE	97S	CDF	$p\overline{p}$; $Z'_\psi \rightarrow e^+e^-$, $\mu^+\mu^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>140	95	[126] CHO	98	RVUE	
>136	95	[127] CHO	98B	RVUE	Electroweak
>160	95	[128] BARATE	97B	ALEP	$e^+e^- \rightarrow \mu^+\mu^-$ and hadronic cross section
>148	95	[129] BUSKULIC	96N	ALEP	Hadronic cross section
>415	95	[130] ABE	95	CDF	$p\overline{p}$; $Z'_\psi \rightarrow e^+e^-$
>105	95	[131] ABREU	95M	DLPH	Z parameters and $e^+e^- \rightarrow \mu^+\mu^-(n\gamma)$
		[132] NARDI	95	RVUE	Z parameters
>135	95	[133] VILAIN	94B	CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\overline{\nu}_\mu e \rightarrow \overline{\nu}_\mu e$
>118	95	[134] ADRIANI	93D	L3	Z parameters
>320	95	[135] ABE	92B	CDF	$p\overline{p}$
>180	95	[136] ABE	92B	CDF	$p\overline{p}$

>160	95	[137,138] LANGACKER	92B	RVUE	Repl. by CVETIC 98
>122	95	[139] LEIKE	92	RVUE	Z parameters
>105	90	[140,141] ABE	90F	VNS	e^+e^-
>146	90	[141,142] ABE	90F	RVUE	e^+e^-, $\nu_\mu e$
>320		[143] BARGER	90B	RVUE	$p\overline{p}$
[> 160]		[144] GONZALEZ-G..	90D	COSM	Nucleosynthesis; light ν_R
[> 2000]		[145] GRIFOLS	90D	ASTR	SN 1987A; light ν_R
>136	90	[146] HAGIWARA	90	RVUE	e^+e^-
>154	90	[147] AMALDI	87	RVUE	
>146	90	[148] DURKIN	86	RVUE	

[124] CVETIC 98 give 95%CL limit on the Z-Z' mixing $-0.0022 < \theta < 0.0026$.

[125] ABE 97S limit is obtained assuming that Z' decays to known fermions only.

[126] CHO 98 limit is from constraints on four-Fermi contact interactions obrained from low-energy electroweak experiments.

[127] CHO 98B use various electroweak data to constrain Z' models. See their Eq. (4.9) for their fit in mass-mixing plane.

[128] BARATE 97B gives 95% CL limits on Z-Z' mixing $-0.0020 < \theta < 0.0038$. The bounds are computed with $\alpha_s = 0.120 \pm 0.003$, $m_t = 175 \pm 6$ GeV, and $M_H = 150^{+120}_{-\ 90}$ GeV. See their Fig. 4 for the limit contour in the mass-mixing plane.

[129] BUSKULIC 96N limit is from a combined fit to the hadronic cross sections measured at $\sqrt{s}$=130, 136 GeV (ALEPH) and $\sqrt{s}$=58 GeV (TOPAZ). Zero mixing is assumed.

[130] ABE 95 limit is obtained assuming that Z' decays to known fermions only. See their Fig. 3 for the mass bound of Z' decaying to all allowed fermions and supersymmetric fermions.

[131] ABREU 95M limit is for α_s=0.123, m_t=150 GeV, and m_H=300 GeV. For the limit contour in the mass-mixing plane, see their Fig. 13.

[132] NARDI 95 give 90%CL limits on Z-Z' mixing $-0.0056 < \theta < 0.0055$ for $M_{Z'}$ >500 GeV, m_t=170 GeV, m_H=250 GeV, α_s=0.12. The bound is relaxed under the simultaneous presence of the mixing of the known fermions with new heavy states, $-0.0066 < \theta < 0.0071$.

[133] VILAIN 94B assume m_t = 150 GeV and θ=0. See Fig. 2 for limit contours in the mass-mixing plane.

[134] ADRIANI 93D give limits on the Z-Z' mixing $-0.003 < \theta < 0.020$ assuming the ABE 92B mass limit.

[135] These limits assume that Z' decays to known fermions only.

[136] These limits assume that Z' decays to all E_6 fermions and their superpartners.

[137] LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. m_t >89 GeV used.

[138] LANGACKER 92B give 95%CL limits on the Z-Z' mixing $-0.0025 < \theta < 0.013$.

[139] LEIKE 92 is based on '90 LEP data published in LEP 92.

[140] ABE 90F use data for R, $R_{\ell\ell}$, and $A_{\ell\ell}$.

[141] ABE 90F fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.

[142] e^+e^- data for R, $R_{\ell\ell}$, $A_{\ell\ell}$, and $A_{c\overline{c}}$ below Z as well as $\nu_\mu e$ scattering data of GEIREGAT 89 is used in the fit.

[143] BARGER 90B limit is based on CDF limit $\sigma(p\overline{p} \rightarrow Z')\cdot B(Z' \rightarrow e^+e^-) < 1$ pb (Nodulman, EPS Conf. '89). Assumes no new threshold is open for Z' decay.

[144] These authors claim that the nucleosynthesis bound on the effective number of light neutrinos ($\delta N_\nu < 1$) constrains Z' masses if ν_R is light ($\lesssim$ 1 MeV).

[145] GRIFOLS 90D limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also RIZZO 91.

[146] HAGIWARA 90 perform a fit to e^+e^- data at PEP, PETRA, and TRISTAN including $\mu^+\mu^-$, $\tau^+\tau^-$, and hadron cross sections and asymmetries.

[147] A wide range of neutral current data as of 1986 are used in the fit.

[148] A wide range of neutral current data as of 1985 are used in the fit.

Limits for Z_η

Z_η is the extra neutral boson in E_6 models, corresponding to $Q_\eta = \sqrt{3/8}\ Q_\chi - \sqrt{5/8}\ Q_\psi$. $g_\eta = e/\cos\theta_W$ is assumed unless otherwise stated. We list limits with the assumption $\rho = 1$ but with no further constraints on the Higgs sector. Values in parentheses assume stronger constraint on the Higgs sector motivated by superstring models. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
>317	95	[149] CHO	98B	RVUE	Electroweak
>620	95	[150] ABE	97S	CDF	$p\overline{p}$; $Z'_\eta \rightarrow e^+e^-, \mu^+\mu^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>340	95	[151] CHO	98	RVUE	
>220	95	[152] CVETIC	98	RVUE	Electroweak
>173	95	[153] BARATE	97B	ALEP	$e^+e^- \rightarrow \mu^+\mu^-$ and hadronic cross section
>167	95	[154] BUSKULIC	96N	ALEP	Hadronic cross section
>440	95	[155] ABE	95	CDF	$p\overline{p}$; $Z'_\eta \rightarrow e^+e^-$
>109	95	[156] ABREU	95M	DLPH	Z parameters and $e^+e^- \rightarrow \mu^+\mu^-(n\gamma)$
		[157] NARDI	95	RVUE	Z parameters
>100	95	[158] VILAIN	94B	CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\overline{\nu}_\mu e \rightarrow \overline{\nu}_\mu e$
>100	95	[159] ADRIANI	93D	L3	Z parameters
(>500)	90	[160] ALTARELLI	93B	RVUE	Z parameters
>340	95	[161] ABE	92B	CDF	$p\overline{p}$
>230	95	[162] ABE	92B	CDF	$p\overline{p}$
(>450)	90	[163] DELAGUILA	92	RVUE	
>182	95	[164,165] LANGACKER	92B	RVUE	Repl. by CVETIC 98
(>315)		[166] LAYSSAC	92B	RVUE	Z parameters
>118	95	[167] LEIKE	92	RVUE	Z parameters
(>470)		[168] RENTON	92	RVUE	
(> 300)	90	[169] ALTARELLI	91B	RVUE	Z parameters
>120	90	[170] GONZALEZ-G...	91	RVUE	
>125	90	[171,172] ABE	90F	VNS	e^+e^-
>115	90	[172,173] ABE	90F	RVUE	e^+e^-, $\nu_\mu e$
>340		[174] BARGER	90B	RVUE	$p\overline{p}$
[> 820]		[175] GONZALEZ-G...	90D	COSM	Nucleosynthesis; light ν_R
[> 3300]		[176] GRIFOLS	90	ASTR	SN 1987A; light ν_R
>100	90	[177] HAGIWARA	90	RVUE	e^+e^-
[> 1040]		[175] LOPEZ	90	COSM	Nucleosynthesis; light ν_R
>173		[178] DELAGUILA	89	RVUE	$p\overline{p}$
>129	90	[179] COSTA	88	RVUE	
>156	90	[180] ELLIS	88	RVUE	
>167	90	[181] ELLIS	88	RVUE	$p\overline{p}$
>111	90	[179] AMALDI	87	RVUE	
>143	90	[182] BARGER	86B	RVUE	$p\overline{p}$
>130	90	[183] DURKIN	86	RVUE	
[> 760]		[175] ELLIS	86	COSM	Nucleosynthesis; light ν_R
[> 500]		[175] STEIGMAN	86	COSM	Nucleosynthesis; light ν_R

[149] CHO 98B use various electroweak data to constrain Z' models. See their Eq. (4.10) for their fit in mass-mixing plane.

[150] ABE 97S limit is obtained assuming that Z' decays to known fermions only.

[151] CHO 98 limit is from constraints on four-Fermi contact interactions obrained from low-energy electroweak experiments.

[152] CVETIC 98 give 95%CL limit on the Z-Z' mixing $-0.0055 < \theta < 0.0021$.

[153] BARATE 97B gives 95% CL limits on Z-Z' mixing $-0.021 < \theta < 0.012$. The bounds are computed with $\alpha_s = 0.120 \pm 0.003$, $m_t = 175 \pm 6$ GeV, and $M_H = 150^{+120}_{-\ 90}$ GeV. See their Fig. 4 for the limit contour in the mass-mixing plane.

[154] BUSKULIC 96N limit is from a combined fit to the hadronic cross sections measured at $\sqrt{s}$=130, 136 GeV (ALEPH) and $\sqrt{s}$=58 GeV (TOPAZ). Zero mixing is assumed.

[155] ABE 95 limit is obtained assuming that Z' decays to known fermions only. See their Fig. 3 for the mass bound of Z' decaying to all allowed fermions and supersymmetric fermions.

[156] ABREU 95M limit is for α_s=0.123, m_t=150 GeV, and m_H=300 GeV. For the limit contour in the mass-mixing plane, see their Fig. 13.

[157] NARDI 95 give 90%CL limits on Z-Z' mixing $-0.0087 < \theta < 0.0075$ for $M_{Z'}$ >500 GeV, m_t=170 GeV, m_H=250 GeV, α_s=0.12. The bound is relaxed under the simultaneous presence of the mixing of the known fermions with new heavy states, $-0.0087 < \theta < 0.010$.

[158] VILAIN 94B assume $m_t = 150$ GeV and θ=0. See Fig. 2 for limit contours in the mass-mixing plane.

[159] ADRIANI 93D give limits on the Z-Z' mixing $-0.029 < \theta < 0.010$ assuming the ABE 92B mass limit.

[160] ALTARELLI 93B limit is from LEP data available in summer '93 and is for $m_t = 110$ GeV. $m_H = 100$ GeV and $\alpha_s = 0.118$ assumed. The 90%CL limit on the Z-Z' mixing angle is in Fig. 2.

[161] These limits assume that Z' decays to known fermions only.

[162] These limits assume that Z' decays to all E_6 fermions and their superpartners.

[163] See Fig. 7d in DELAGUILA 92 for the allowed region in $m_{Z'}$-mixing plane from electroweak fit including '90 LEP data.

[164] LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. m_t >89 GeV used.

[165] LANGACKER 92B give 95%CL limits on the Z-Z' mixing $-0.038 < \theta < 0.002$.

[166] LAYSSAC 92B limit is from LEP data available spring '92. Specific Higgs sector is assumed. See also LAYSSAC 92.

[167] LEIKE 92 is based on '90 LEP data published in LEP 92.

[168] RENTON 92 limits use LEP data taken up to '90 as well as m_W, νN, and atomic parity violation data. Specific Higgs structure is assumed.

[169] ALTARELLI 91B is based on Z mass, widths, and A_{FB}. The limits are for superstring motivated models with extra assumption on the Higgs sector. $m_t > 90$ GeV and $m_{H^0} < 1$ TeV assumed. For large m_t, the bound improves drastically. Bounds for Z-Z' mixing angle and Z mass shift without this model assumption are also given in the paper.

[170] GONZALEZ-GARCIA 91 limit is based on low-energy neutral current data, LEP Z mass and widths, m_W from ABE 90G. 100 $< m_t <$ 200 GeV, $m_{H^0} = 100$ GeV assumed. Dependence on m_t is shown in Fig. 8.

[171] ABE 90F use data for R, $R_{\ell\ell}$, and $A_{\ell\ell}$.

[172] ABE 90F fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.

[173] e^+e^- data for R, $R_{\ell\ell}$, $A_{\ell\ell}$, and $A_{c\overline{c}}$ below Z as well as $\nu_\mu e$ scattering data of GEIREGAT 89 is used in the fit.

[174] BARGER 90B limit is based on CDF limit $\sigma(p\overline{p} \rightarrow Z')\cdot B(Z' \rightarrow e^+e^-) < 1$ pb (Nodulman, EPS Conf. '89). Assumes no new threshold is open for Z' decay.

[175] These authors claim that the nucleosynthesis bound on the effective number of light neutrinos ($\delta N_\nu < 1$) constrains Z' masses if ν_R is light ($\lesssim$ 1 MeV).
[176] GRIFOLS 90 limit holds for $m_{\nu_R} \lesssim$ 1 MeV. See also GRIFOLS 90D, RIZZO 91.
[177] HAGIWARA 90 perform a fit to $e^+ e^-$ data at PEP, PETRA, and TRISTAN including $\mu^+ \mu^-$, $\tau^+ \tau^-$, and hadron cross sections and asymmetries.
[178] DELAGUILA 89 limit is based on $\sigma(p\overline{p} \rightarrow Z')\cdot B(Z' \rightarrow e^+ e^-) <$ 1.8 pb at CERN $p\overline{p}$ collider.
[179] A wide range of neutral current data as of 1986 are used in the fit.
[180] Z_η mass limits obtained by combining constraints from non-observation of an excess of $\ell^+ \ell^-$ pairs at the CERN $p\overline{p}$ collider and the global analysis of neutral current data by COSTA 88. Least favorable spectrum of three (E_6 27) generations of particles and their superpartners are assumed.
[181] Z' mass limits from non-observation of an excess of $\ell^+ \ell^-$ pairs at the CERN $p\overline{p}$ collider [based on ANSARI 87D and GEER Uppsala Conf. 87]. The limits apply when Z' decays only into light quarks and leptons.
[182] BARGER 86B limit is based on UA1/UA2 limit on $p\overline{p} \rightarrow Z'$, $Z' \rightarrow e^+ e^-$ (Lepton Photon Symp., Kyoto, '85). Extra decay channels for Z' are assumed not be open.
[183] A wide range of neutral current data as of 1985 are used in the fit.

Limits for other Z'

$Z_\beta = Z_\chi \cos\beta + Z_\psi \sin\beta$

VALUE (GeV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>290	95	[184] CHO	98	RVUE	Z_β with $\tan\beta = \sqrt{15}$
>284	95	[185] CHO	98B	RVUE	$\tan\beta = \sqrt{15}$
		[186] DELAGUILA	92	RVUE	
>360		[187] ALTARELLI	91	RVUE	Z_β with $\tan\beta = \sqrt{3/5}$; Cs
>190		[188] MAHANTHAP...	91	RVUE	Z_β with $\tan\beta = \sqrt{3/5}$; Cs
		[189] GRIFOLS	90C	RVUE	
		[190] DELAGUILA	89	RVUE	$p\overline{p}$
>180	90	[191,192] COSTA	88	RVUE	Z_β with $\tan\beta = \sqrt{15}$
>158	90	[193] ELLIS	88	RVUE	Z_β ($\tan\beta = \sqrt{15}$), $p\overline{p}$

[184] CHO 98 limit is from constraints on four-Fermi contact interactions obrained from low-energy electroweak experiments.
[185] CHO 98B use various electroweak data to constrain Z' models. See their Eq. (4.11) for their fit in mass-mixing plane.
[186] Fig. 7c and 7e in DELAGUILA 92 give limits for $\tan\beta = -1/\sqrt{15}$ and $\sqrt{15}$ from electroweak fit including '90 LEP data.
[187] ALTARELLI 91 limit is from atomic parity violation in Cs together with LEP, CDF data. Z-Z' mixing is assumed to be zero to set the limit.
[188] MAHANTHAPPA 91 limit is from atomic parity violation in Cs with m_W, m_Z. See Table III of MAHANTHAPPA 91 (corrected in erratum) for limits on various Z' models.
[189] GRIFOLS 90C obtains a limit for Z' mass as a function of mixing angle β (his $\theta = \beta - \pi/2$), which is derived from a LAMPF experiment on $\sigma(\nu_e e)$ (ALLEN 90). The result is shown in Fig. 1.
[190] See Table I of DELAGUILA 89 for limits on various Z' models.
[191] $g_\beta = e/\cos\theta_W$ and $\rho = 1$ assumed.
[192] A wide range of neutral current data as of 1986 are used in the fit.

[193] Z' mass limits from non-observation of an excess of $\ell^+ \ell^-$ pairs at the CERN $p\overline{p}$ collider [based on ANSARI 87D and GEER Uppsala Conf. 87]. The limits apply when Z' decays only into light quarks and leptons.

A REVIEW GOES HERE – Check our WWW List of Reviews

MASS LIMITS for Leptoquarks from Pair Production

These limits rely only on the color or electroweak charge of the leptoquark.

VALUE (GeV)	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
>225	95		[194] ABBOTT	98E	D0	First generation
>202	95		[195] ABE	98S	CDF	Second generation
> 99	95		[196] ABE	97F	CDF	Third generation
> 45.5	95		[197,198] ABREU	93J	DLPH	First + second generation
> 44.4	95		[199] ADRIANI	93M	L3	First generation
> 44.6	95		[200] ADRIANI	93M	L3	Third generation
> 44	95		[199] DECAMP	92	ALEP	First or second generation
> 45	95		[199] DECAMP	92	ALEP	Third generation
> 44.2	95		[199] ALEXANDER	91	OPAL	First or second generation
> 41.4	95		[199] ALEXANDER	91	OPAL	Third generation
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 94	95		[201] ABBOTT	98J	D0	Third generation
>225	95		[202] ABBOTT	97B	D0	Result included in ABBOTT 98E
>213	95		[202] ABE	97X	CDF	First generation
>119	95		[203] ABACHI	95G	D0	Second generation
>131	95		[204] ABE	95U	CDF	Second generation
>116	95		[205] ABACHI	94B	D0	First generation
> 80	95		[206] ABE	93I	CDF	First generation
> 44.5	95		[199] ADRIANI	93M	L3	Second generation
> 42.1	95		[207] ABREU	92F	DLPH	Second generation
> 74	95		[208] ALITTI	92E	UA2	First generation
> 43.2	95		[199] ADEVA	91B	L3	First generation
> 43.4	95		[199] ADEVA	91B	L3	Second generation
none 8.9–22.6	95		[209] KIM	90	AMY	First generation
none 10.2–23.2	95		[209] KIM	90	AMY	Second generation
none 5–20.8	95		[210] BARTEL	87B	JADE	
none 7–20.5	95	2	[211] BEHREND	86B	CELL	

[194] ABBOTT 98E search for scalar leptoquarks using $e\nu jj$, $eejj$, and $\nu\nu jj$ events in $p\overline{p}$ collisions at E_{cm}=1.8 TeV. The limit above assumes B(eq)=1. For B(eq)=0.5 and 0, the bound becomes 204 and 79 GeV, respectively.

[195] ABE 98S search for scalar leptoquarks using $\mu\mu jj$ events in $p\overline{p}$ collisions at E_{cm}= 1.8 TeV. The limit is for B(μq)= 1. For B(μq)=B(νq)=0.5, the limit is > 160 GeV.

[196] ABE 97F search for third generation scalar and vector leptoquarks in $p\overline{p}$ collisions at E_{cm} = 1.8 TeV. The quoted limit is for scalar leptoquark with B(τb) = 1.

[197] Limit is for charge $-1/3$ isospin-0 leptoquark with B(ℓq) = 2/3.

[198] First and second generation leptoquarks are assumed to be degenerate. The limit is slightly lower for each generation.

[199] Limits are for charge $-1/3$, isospin-0 scalar leptoquarks decaying to $\ell^- q$ or νq with any branching ratio. See paper for limits for other charge-isospin assignments of leptoquarks.

[200] ADRIANI 93M limit for charge $-1/3$, isospin-0 leptoquark decaying to τb.

[201] ABBOTT 98J search for charge $-1/3$ third generation scalar and vector leptoquarks in $p\overline{p}$ collisions at $E_{\rm cm}$= 1.8 TeV. The quoted limit is for scalar leptoquark with B(νb)=1.

[202] ABBOTT 97B, ABE 97X search for scalar leptoquarks using $eejj$ events in $p\overline{p}$ collisions at $E_{\rm cm}$=1.8 TeV. The limit is for B(eq)=1.

[203] ABACHI 95G search for scalar leptoquarks using $\mu\mu$+jets and $\mu\nu_\mu$+jets events in $p\overline{p}$ collisions at $E_{\rm cm}$ = 1.8 TeV. The limit is for B(μq) = 1. For B(μq) = B(νq) = 0.5, the limit is > 97 GeV.

[204] ABE 95U search for scalar leptoquarks of charge Q=2/3 and $-1/3$ using $\mu\mu jj$ events in $p\overline{p}$ collisions at $E_{\rm cm}$=1.8 TeV. The limit is for B(μq) = 1. For B(μq) = B(νq) = 0.5, the limit is > 96 GeV.

[205] ABACHI 94B search for $eejj$ and $e\nu jj$ events in $p\overline{p}$ collisions at $E_{\rm cm}$=1.8 TeV. ABACHI 94B obtain the limit >120 GeV for B(eq)=B(νq)=0.5 and >133 GeV for B(eq)=1. A change in the DØ luminosity monitor constant reduces the first bound to >116 GeV quoted above (see FERMILAB-TM-1911). This limit does not depend on the electroweak quantum numbers of the leptoquark.

[206] ABE 93I search for $\ell\ell jj$ events in $p\overline{p}$ collisions at $E_{\rm cm}$=1.8 TeV. The limit is for B(eq) = B(νq) = 0.5 and improves to >113 GeV for B(eq) = 1. This limit does not depend on electroweak quantum numbers of the leptoquark.

[207] ABREU 92F limit is for charge $-1/3$ isosin-0 leptoquark with B(μq)=2/3. If first and second generation leptoquarks are degenerate, the limit is 43.0 GeV, and for a charge 2/3 second generation leptoquark 43.4 GeV. Cross-section limit for pair production of states decaying to ℓq is given in the paper.

[208] ALITTI 92E search for $\ell\ell jj$ and $\ell\nu jj$ events in $p\overline{p}$ collisions at $E_{\rm cm}$=630 GeV. The limit is for B(eq) = 1 and is reduced to 67 GeV for B(eq) = B(νq) = 0.5. This limit does not depend on electroweak quantum numbers of the leptoquark.

[209] KIM 90 assume pair production of charge 2/3 scalar-leptoquark via photon exchange. The decay of the first (second) generation leptoquark is assumed to be any mixture of de^+ and $u\overline{\nu}$ ($s\mu^+$ and $c\overline{\nu}$). See paper for limits for specific branching ratios.

[210] BARTEL 87B limit is valid when a pair of charge 2/3 spinless leptoquarks X is produced with point coupling, and when they decay under the constraint B(X $\rightarrow$ $c\overline{\nu}_\mu$) + B(X $\rightarrow$ $s\mu^+$) = 1.

[211] BEHREND 86B assumed that a charge 2/3 spinless leptoquark, χ, decays either into $s\mu^+$ or $c\overline{\nu}$: B($\chi \rightarrow s\mu^+$) + B($\chi \rightarrow c\overline{\nu}$) = 1.

MASS LIMITS for Leptoquarks from Single Production

These limits depend on the q-ℓ-leptoquark coupling g_{LQ}. It is often assumed that $g^2_{LQ}/4\pi$=1/137. Limits shown are for a scalar, weak isoscalar, charge $-1/3$ leptoquark.

VALUE (GeV)	CL%	DOCUMENT ID		TECN	COMMENT
>237	95	[212] AID	96B	H1	First generation
> 73	95	[213] ABREU	93J	DLPH	Second generation
• • • We do not use the following data for averages, fits, limits, etc. • • •					
		[214] DERRICK	97	ZEUS	Lepton-flavor violation
>230	95	[215] AHMED	94B	H1	Sup. by AID 96B
> 65	95	[213] ABREU	93J	DLPH	First generation
>181	95	[216] ABT	93	H1	First generation
>168	95	[217] DERRICK	93	ZEUS	First generation

[212] The quoted limit is for a left-handed scalar leptoquark which solely couples to the first generation with electromagnetic strength. AID 96B also search for leptoquarks with lepton-flavor violating couplings. For limits on states with different quantum numbers and the limits in the coupling-mass plane, see their Fig. 2, Fig. 3, and Table 2. AID 96B supersedes AHMED 94B.

[213] Limit from single production in Z decay. The limit is for a leptoquark coupling of electromagnetic strength and assumes $B(\ell q) = 2/3$. The limit is 77 GeV if first and second leptoquarks are degenerate.

[214] DERRICK 97 search for various leptoquarks with lepton-flavor violating couplings. See their Figs. 5–8 and Table 1 for detailed limits.

[215] AHMED 94B limit is for the left-handed leptoquark decaying to eq and νq with $B(eq) = B(\nu q)=1/2$. Electromagnetic coupling strength is assumed for the scalar leptoquark interaction. For limits on states with different quantum numbers and the limits in the coupling-mass plane, see their Table 2 and Fig. 6.

[216] ABT 93 search for single leptoquark production in ep collisions with the decays eq and νq. The limit is for a leptoquark coupling of electromagnetic strength and assumes $B(eq) = B(\nu q) = 1/2$. The limit for $B(eq) = 1$ is 178 GeV. For limits on states with different quantum numbers, see their Fig. 2. ABT 93 superseded by AHMED 94B.

[217] DERRICK 93 search for single leptoquark production in ep collisions with the decay eq and νq. The limit is for leptoquark coupling of electromagnetic strength and assumes $B(eq) = B(\nu q) = 1/2$. The limit for $B(eq) = 1$ is 176 GeV. For limits on states with different quantum numbers, see their Table 3.

Indirect Limits for Leptoquarks

VALUE (TeV)	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
		[218] ABBIENDI	99	OPAL	
		[219] ACCIARRI	98J	L3	$e^+e^- \rightarrow q\overline{q}$
		[220] ACKERSTAFF	98V	OPAL	$e^+e^- \rightarrow q\overline{q}$, $e^+e^- \rightarrow b\overline{b}$
$>$ 0.76	95	[221] DEANDREA	97	RVUE	$\widetilde{R}_2$ leptoquark
		[222] DERRICK	97	ZEUS	Lepton-flavor violation
		[223] GROSSMAN	97	RVUE	$B \rightarrow \tau^+\tau^-$(X)
		[224] JADACH	97	RVUE	$e^+e^- \rightarrow q\overline{q}$
$>$ 0.31	95	[225] AID	95	H1	First generation
$>$1200		[226] KUZNETSOV	95B	RVUE	Pati-Salam type
		[227] MIZUKOSHI	95	RVUE	Third generation scalar leptoquark
$>$ 0.3	95	[228] BHATTACH...	94	RVUE	Spin-0 leptoquark coupled to $\overline{e}_R t_L$
		[229] DAVIDSON	94	RVUE	
$>$ 18		[230] KUZNETSOV	94	RVUE	Pati-Salam type
$>$ 0.43	95	[231] LEURER	94	RVUE	First generation spin-1 leptoquark
$>$ 0.44	95	[231] LEURER	94B	RVUE	First generation spin-0 leptoquark
		[232] MAHANTA	94	RVUE	P and T violation
$>$ 350		[233] DESHPANDE	83	RVUE	Sup. by KUZNETSOV 95B
$>$ 1		[234] SHANKER	82	RVUE	Nonchiral spin-0 leptoquark
$>$ 125		[234] SHANKER	82	RVUE	Nonchiral spin-1 leptoquark

[218] ABBIENDI 99 limits are from $e^+e^- \rightarrow q\overline{q}$ cross section at 130–136, 161–172, 183 GeV. See their Fig. 8 and Fig. 9 for limits in mass-coupling plane.

[219] ACCIARRI 98J limit is from $e^+e^- \rightarrow q\overline{q}$ cross section at $\sqrt{s}$= 130–172 GeV which can be affected by the t- and u-channel exchanges of leptoquarks. See their Fig. 4 and Fig. 5 for limits in the mass-coupling plane.

[220] ACKERSTAFF 98V limits are from $e^+ e^- \rightarrow q\overline{q}$ and $e^+ e^- \rightarrow b\overline{b}$ cross sections at $\sqrt{s}$ = 130–172 GeV, which can be affected by the t- and u-channel exchanges of leptoquarks. See their Fig. 21 and Fig. 22 for limits of leptoquarks in mass-coupling plane.

[221] DEANDREA 97 limit is for $\widetilde{R}_2$ leptoquark obtained from atomic parity violation (APV). The coupling of leptoquark is assumed to be electromagnetic strength. See Table 2 for limits of the four-fermion interactions induced by various scalar leptoquark exchange. DEANDREA 97 combines APV limit and limits from Tevatron and HERA. See Fig. 1–4 for combined limits of leptoquark in mass-coupling plane.

[222] DERRICK 97 search for lepton-flavor violation in ep collision. See their Tables 2–5 for limits on lepton-flavor violating four-fermion interactions induced by various leptoquarks.

[223] GROSSMAN 97 estimate the upper bounds on the branching fraction $B \rightarrow \tau^+ \tau^-$ (X) from the absence of the B decay with large missing energy. These bounds can be used to constrain leptoquark induced four-fermion interactions.

[224] JADACH 97 limit is from $e^+ e^- \rightarrow q\overline{q}$ cross section at $\sqrt{s}$=172.3 GeV which can be affected by the t- and u-channel exchanges of leptoquarks. See their Fig. 1 for limits on vector leptoquarks in mass-coupling plane.

[225] AID 95 limit is for the weak isotriplet spin-1 leptoquark with the electromagnetic coupling strength. For the limits of leptoquarks with different quantum number, see their Table 2. AID 95 limits are from the measurements of the Q^2 spectrum measurement of $ep \rightarrow eX$.

[226] KUZNETSOV 95B use π, K, B, τ decays and μe conversion and give a list of bounds on the leptoquark mass and the fermion mixing matrix in the Pati-Salam model. The quoted limit is from $K_L \rightarrow \mu e$ decay assuming zero mixing. See also KUZNETSOV 94, DESHPANDE 83, and DIMOPOULOS 81.

[227] MIZUKOSHI 95 calculate the one-loop radiative correction to the Z-physics parameters in various scalar leptoquark models. See their Fig. 4 for the exclusion plot of third generation leptoquark models in mass-coupling plane.

[228] BHATTACHARYYA 94 limit is from one-loop radiative correction to the leptonic decay width of the Z. m_H=250 GeV, $\alpha_s(m_Z)$=0.12, m_t=180 GeV, and the electroweak strength of leptoquark coupling are assumed. For leptoquark coupled to $\overline{e}_L\, t_R$, $\overline{\mu}\, t$, and $\overline{\tau}\, t$, see Fig. 2 in BHATTACHARYYA 94B erratum and Fig. 3.

[229] DAVIDSON 94 gives an extensive list of the bounds on leptoquark-induced four-fermion insteractions from π, K, D, B, μ, τ decays and meson mixings, *etc.* See Table 15 of DAVIDSON 94 for detail.

[230] KUZNETSOV 94 gives mixing independent bound of the Pati-Salam leptoquark from the cosmological limit on $\pi^0 \rightarrow \overline{\nu}\nu$.

[231] LEURER 94, LEURER 94B limits are obtained from atomic parity violation and apply to any chiral leptoquark which couples to the first generation with electromagnetic strength. For a nonchiral leptoquark, universality in $\pi_{\ell 2}$ decay provides a much more stringent bound. See also SHANKER 82.

[232] MAHANTA 94 gives bounds of P- and T-violating scalar-leptoquark couplings from atomic and molecular experiments.

[233] DESHPANDE 83 used upper limit on $K^0_L \rightarrow \mu e$ decay with renormalization-group equations to estimate coupling at the heavy boson mass. See also DIMOPOULOS 81.

[234] From $(\pi \rightarrow e\nu)/(\pi \rightarrow \mu\nu)$ ratio. SHANKER 82 assumes the leptoquark induced four-fermion coupling $4g^2/M^2$ $(\overline{\nu}_{eL}\; u_R)$ $(\overline{d}_L\, e_R)$with g=0.004 for spin-0 leptoquark and g^2/M^2 $(\overline{\nu}_{eL}\; \gamma_\mu u_L)$ $(\overline{d}_R\; \gamma^\mu\, e_R)$ with $g \simeq$ 0.6 for spin-1 leptoquark.

MASS LIMITS for Diquarks

VALUE (GeV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •					
none 290–420	95	[235] ABE	97G	CDF	E_6 diquark
none 15–31.7	95	[236] ABREU	94O	DLPH	SUSY E_6 diquark

[235] ABE 97G search for new particle decaying to dijets.

[236] ABREU 94O limit is from $e^+ e^- \rightarrow \overline{c}\overline{s}cs$. Range extends up to 43 GeV if diquarks are degenerate in mass.

MASS LIMITS for g_A (axigluon)

Axigluons are massive color-octet gauge bosons in chiral color models and have axial-vector coupling to quarks with the same coupling strength as gluons.

VALUE (GeV)	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>365	95	[237] DONCHESKI	98	RVUE	$\Gamma(Z \rightarrow$ hadron)
none 200–980	95	[238] ABE	97G	CDF	$p\overline{p} \rightarrow g_A X$, $X \rightarrow$ 2 jets
none 200–870	95	[239] ABE	95N	CDF	$p\overline{p} \rightarrow g_A X$, $g_A \rightarrow q\overline{q}$
none 240–640	95	[240] ABE	93G	CDF	$p\overline{p} \rightarrow g_A X$, $g_A \rightarrow$ 2jets
> 50	95	[241] CUYPERS	91	RVUE	$\sigma(e^+ e^- \rightarrow$ hadrons)
none 120–210	95	[242] ABE	90H	CDF	$p\overline{p} \rightarrow g_A X$, $g_A \rightarrow$ 2jets
> 29		[243] ROBINETT	89	THEO	Partial-wave unitarity
none 150–310	95	[244] ALBAJAR	88B	UA1	$p\overline{p} \rightarrow g_A X$, $g_A \rightarrow$ 2jets
> 20		BERGSTROM	88	RVUE	$p\overline{p} \rightarrow \Upsilon X$ via $g_A g$
> 9		[245] CUYPERS	88	RVUE	Υ decay
> 25		[246] DONCHESKI	88B	RVUE	Υ decay

[237] DONCHESKI 98 compare α_s derived from low-energy data and that from $\Gamma(Z \rightarrow$ hadrons$)/\Gamma(Z \rightarrow$ leptons).

[238] ABE 97G search for new particle decaying to dijets.

[239] ABE 95N assume axigluons decaying to quarks in the Standard Model only.

[240] ABE 93G assume $\Gamma(g_A) = N\alpha_s m_{g_A}/6$ with $N = 10$.

[241] CUYPERS 91 compare α_s measured in Υ decay and that from R at PEP/PETRA energies.

[242] ABE 90H assumes $\Gamma(g_A) = N\alpha_s m_{g_A}/6$ with $N = 5\,(\Gamma(g_A) = 0.09 m_{g_A})$. For $N = 10$, the excluded region is reduced to 120–150 GeV.

[243] ROBINETT 89 result demands partial-wave unitarity of $J = 0$ $t\overline{t} \rightarrow t\overline{t}$ scattering amplitude and derives a limit $m_{g_A} > 0.5\ m_t$. Assumes $m_t > 56$ GeV.

[244] ALBAJAR 88B result is from the nonobservation of a peak in two-jet invariant mass distribution. $\Gamma(g_A) < 0.4\ m_{g_A}$ assumed. See also BAGGER 88.

[245] CUYPERS 88 requires $\Gamma(\Upsilon \rightarrow g g_A) < \Gamma(\Upsilon \rightarrow g g g)$. A similar result is obtained by DONCHESKI 88.

[246] DONCHESKI 88B requires $\Gamma(\Upsilon \rightarrow g q\overline{q})/\Gamma(\Upsilon \rightarrow g g g) < 0.25$, where the former decay proceeds via axigluon exchange. A more conservative estimate of < 0.5 leads to $m_{g_A} > 21$ GeV.

X^0 (Heavy Boson) Searches in Z Decays

Searches for radiative transition of Z to a lighter spin-0 state X^0 decaying to hadrons, a lepton pair, a photon pair, or invisible particles as shown in the comments. The limits are for the product of branching ratios.

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
		[247] BARATE	98U	ALEP	$X^0 \to \ell\overline{\ell}, q\overline{q}, gg, \gamma\gamma, \nu\overline{\nu}$
		[248] ACCIARRI	97Q	L3	$X^0 \to$ invisible particle(s)
		[249] ACTON	93E	OPAL	$X^0 \to \gamma\gamma$
		[250] ABREU	92D	DLPH	$X^0 \to$ hadrons
		[251] ADRIANI	92F	L3	$X^0 \to$ hadrons
		[252] ACTON	91	OPAL	$X^0 \to$ anything
$<1.1 \times 10^{-4}$	95	[253] ACTON	91B	OPAL	$X^0 \to e^+e^-$
$<9 \times 10^{-5}$	95	[253] ACTON	91B	OPAL	$X^0 \to \mu^+\mu^-$
$<1.1 \times 10^{-4}$	95	[253] ACTON	91B	OPAL	$X^0 \to \tau^+\tau^-$
$<2.8 \times 10^{-4}$	95	[254] ADEVA	91D	L3	$X^0 \to e^+e^-$
$<2.3 \times 10^{-4}$	95	[254] ADEVA	91D	L3	$X^0 \to \mu^+\mu^-$
$<4.7 \times 10^{-4}$	95	[255] ADEVA	91D	L3	$X^0 \to$ hadrons
$<8 \times 10^{-4}$	95	[256] AKRAWY	90J	OPAL	$X^0 \to$ hadrons

[247] BARATE 98U obtain limits on B($Z \to \gamma X^0$)B($X^0 \to \ell\overline{\ell}$, $q\overline{q}$, gg, $\gamma\gamma$, $\nu\overline{\nu}$). See their Fig. 17.

[248] See Fig. 4 of ACCIARRI 97Q for the upper limit on B($Z \to \gamma X^0$; $E_\gamma > E_{\rm min}$) as a function of $E_{\rm min}$.

[249] ACTON 93E give $\sigma(e^+e^- \to X^0\gamma)\cdot$B($X^0 \to \gamma\gamma$)$<$ 0.4 pb (95%CL) for m_{X^0}=60 $\pm$ 2.5 GeV. If the process occurs via s-channel γ exchange, the limit translates to $\Gamma(X^0)\cdot$B$(X^0 \to \gamma\gamma)^2$ $<$20 MeV for m_{X^0} = 60 $\pm$ 1 GeV.

[250] ABREU 92D give σ_Z $\cdot$ B($Z \to \gamma X^0$) $\cdot$ B($X^0 \to$ hadrons) $<$(3–10) pb for m_{X^0} = 10–78 GeV. A very similar limit is obtained for spin-1 X^0.

[251] ADRIANI 92F search for isolated γ in hadronic Z decays. The limit σ_Z $\cdot$ B($Z \to \gamma X^0$) $\cdot$ B($X^0 \to$ hadrons) $<$(2–10) pb (95%CL) is given for m_{X^0} = 25–85 GeV.

[252] ACTON 91 searches for $Z \to Z^* X^0$, $Z^* \to e^+e^-$, $\mu^+\mu^-$, or $\nu\overline{\nu}$. Excludes any new scalar X^0 with m_{X^0} $<$ 9.5 GeV/c if it has the same coupling to ZZ^* as the MSM Higgs boson.

[253] ACTON 91B limits are for m_{X^0} = 60–85 GeV.

[254] ADEVA 91D limits are for m_{X^0} = 30–89 GeV.

[255] ADEVA 91D limits are for m_{X^0} = 30–86 GeV.

[256] AKRAWY 90J give $\Gamma(Z \to \gamma X^0)\cdot$B($X^0 \to$ hadrons) $<$ 1.9 MeV (95%CL) for m_{X^0} = 32–80 GeV. We divide by $\Gamma(Z)$ = 2.5 GeV to get product of branching ratios. For nonresonant transitions, the limit is B($Z \to \gamma q\overline{q}$) $<$ 8.2 MeV assuming three-body phase space distribution.

MASS LIMITS for a Heavy Neutral Boson Coupling to e^+e^-

VALUE (GeV)	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
none 55–61		[257] ODAKA	89	VNS	$\Gamma(X^0 \to e^+e^-) \cdot B(X^0 \to \text{hadrons}) \gtrsim 0.2$ MeV
>45	95	[258] DERRICK	86	HRS	$\Gamma(X^0 \to e^+e^-)$=6 MeV
>46.6	95	[259] ADEVA	85	MRKJ	$\Gamma(X^0 \to e^+e^-)$=10 keV
>48	95	[259] ADEVA	85	MRKJ	$\Gamma(X^0 \to e^+e^-)$=4 MeV
		[260] BERGER	85B	PLUT	
none 39.8–45.5		[261] ADEVA	84	MRKJ	$\Gamma(X^0 \to e^+e^-)$=10 keV
>47.8	95	[261] ADEVA	84	MRKJ	$\Gamma(X^0 \to e^+e^-)$=4 MeV
none 39.8–45.2		[261] BEHREND	84C	CELL	
>47	95	[261] BEHREND	84C	CELL	$\Gamma(X^0 \to e^+e^-)$=4 MeV

[257] ODAKA 89 looked for a narrow or wide scalar resonance in $e^+e^- \to$ hadrons at E_{cm} = 55.0–60.8 GeV.

[258] DERRICK 86 found no deviation from the Standard Model Bhabha scattering at E_{cm}= 29 GeV and set limits on the possible scalar boson e^+e^- coupling. See their figure 4 for excluded region in the $\Gamma(X^0 \to e^+e^-)$-m_{X^0} plane. Electronic chiral invariance requires a parity doublet of X^0, in which case the limit applies for $\Gamma(X^0 \to e^+e^-)$ = 3 MeV.

[259] ADEVA 85 first limit is from 2γ, $\mu^+\mu^-$, hadrons assuming X^0 is a scalar. Second limit is from e^+e^- channel. E_{cm} = 40–47 GeV. Supersedes ADEVA 84.

[260] BERGER 85B looked for effect of spin-0 boson exchange in $e^+e^- \to e^+e^-$ and $\mu^+\mu^-$ at E_{cm} = 34.7 GeV. See Fig. 5 for excluded region in the $m_{X^0} - \Gamma(X^0)$ plane.

[261] ADEVA 84 and BEHREND 84C have E_{cm} = 39.8–45.5 GeV. MARK-J searched X^0 in $e^+e^- \to$ hadrons, 2γ, $\mu^+\mu^-$, e^+e^- and CELLO in the same channels plus τ pair. No narrow or broad X^0 is found in the energy range. They also searched for the effect of X^0 with $m_X > E_{\text{cm}}$. The second limits are from Bhabha data and for spin-0 singlet. The same limits apply for $\Gamma(X^0 \to e^+e^-)$ = 2 MeV if X^0 is a spin-0 doublet. The second limit of BEHREND 84C was read off from their figure 2. The original papers also list limits in other channels.

Search for X^0 Resonance in e^+e^- Collisions

The limit is for $\Gamma(X^0 \to e^+e^-) \cdot B(X^0 \to f)$, where f is the specified final state. Spin 0 is assumed for X^0.

VALUE (keV)	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<10^3$	95	[262] ABE	93C	VNS	$\Gamma(ee)$
<(0.4–10)	95	[263] ABE	93C	VNS	$f = \gamma\gamma$
<(0.3–5)	95	[264,265] ABE	93D	TOPZ	$f = \gamma\gamma$
<(2–12)	95	[264,265] ABE	93D	TOPZ	f = hadrons
<(4–200)	95	[265,266] ABE	93D	TOPZ	$f = ee$
<(0.1–6)	95	[265,266] ABE	93D	TOPZ	$f = \mu\mu$
<(0.5–8)	90	[267] STERNER	93	AMY	$f = \gamma\gamma$

[262] Limit is for $\Gamma(X^0 \to e^+e^-)$ m_{X^0} = 56–63.5 GeV for $\Gamma(X^0)$ = 0.5 GeV.

[263] Limit is for m_{X^0} = 56–61.5 GeV and is valid for $\Gamma(X^0) \ll$ 100 MeV. See their Fig. 5 for limits for Γ = 1,2 GeV.

[264] Limit is for m_{X^0} = 57.2–60 GeV.

[265] Limit is valid for $\Gamma(X^0) \ll$ 100 MeV. See paper for limits for Γ = 1 GeV and those for $J = 2$ resonances.

[266] Limit is for m_{X^0} = 56.6–60 GeV.

[267] STERNER 93 limit is for m_{X^0} = 57–59.6 GeV and is valid for $\Gamma(X^0)$<100 MeV. See their Fig. 2 for limits for Γ = 1,3 GeV.

Search for X^0 Resonance in Two-Photon Process

The limit is for $\Gamma(X^0) \cdot B(X^0 \to \gamma\gamma)^2$. Spin 0 is assumed for X^0.

VALUE (MeV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.6	95	[268] ACTON	93E	OPAL	m_{X^0}=60 ± 1 GeV
<2.9	95	BUSKULIC	93F	ALEP	$m_{X^0} \sim$ 60 GeV

[268] ACTON 93E limit for a $J = 2$ resonance is 0.8 MeV.

Search for X^0 Resonance in $e^+ e^- \to X^0 \gamma$

VALUE (GeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •				
	[269] ADAM	96C	DLPH	X^0 decaying invisibly

[269] ADAM 96C is from the single photon production cross at $\sqrt{s}$=130, 136 GeV. The upper bound is less than 3 pb for X^0 masses between 60 and 130 GeV. See their Fig. 5 for the exact bound on the cross section $\sigma(e^+ e^- \to \gamma X^0)$.

Search for X^0 Resonance in $Z \to f \overline{f} X^0$

The limit is for $B(Z \to f \overline{f} X^0) \cdot B(X^0 \to F)$ where f is a fermion and F is the specified final state. Spin 0 is assumed for X^0.

VALUE (MeV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •					
		[270] ABREU	96T	DLPH	f=e,μ,τ; F=$\gamma\gamma$
$<3.7 \times 10^{-6}$	95	[271] ABREU	96T	DLPH	f=ν; F=$\gamma\gamma$
		[272] ABREU	96T	DLPH	f=q; F=$\gamma\gamma$
$<6.8 \times 10^{-6}$	95	[271] ACTON	93E	OPAL	f=e,μ,τ; F=$\gamma\gamma$
$<5.5 \times 10^{-6}$	95	[271] ACTON	93E	OPAL	f=q; F=$\gamma\gamma$
$<3.1 \times 10^{-6}$	95	[271] ACTON	93E	OPAL	f=ν; F=$\gamma\gamma$
$<6.5 \times 10^{-6}$	95	[271] ACTON	93E	OPAL	f=e,μ; F=$\ell\overline{\ell}$, $q\overline{q}$, $\nu\overline{\nu}$
$<7.1 \times 10^{-6}$	95	[271] BUSKULIC	93F	ALEP	f=e,μ; F=$\ell\overline{\ell}$, $q\overline{q}$, $\nu\overline{\nu}$
		[273] ADRIANI	92F	L3	f=q; F=$\gamma\gamma$

[270] ABREU 96T obtain limit as a function of m_{X^0}. See their Fig. 6.

[271] Limit is for m_{X^0} around 60 GeV.

[272] ABREU 96T obtain limit as a function of m_{X^0}. See their Fig. 15.

[273] ADRIANI 92F give $\sigma_Z \cdot$ B($Z \rightarrow q\overline{q}X^0$) $\cdot$ B($X^0 \rightarrow \gamma\gamma$)<(0.75–1.5) pb (95%CL) for m_{X^0} = 10–70 GeV. The limit is 1 pb at 60 GeV.

Search for X^0 Resonance in $p\overline{p} \rightarrow WX^0$

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •				
	[274] ABE	97W	CDF	$X^0 \rightarrow b\overline{b}$

[274] ABE 97W search for X^0 production associated with W in $p\overline{p}$ collisions at E_{cm}=1.8 TeV. The 95%CL upper limit on the production cross section times the branching ratio for $X^0 \rightarrow b\overline{b}$ ranges from 14 to 19 pb for X^0 mass between 70 and 120 GeV. See their Fig. 3 for upper limits of the production cross section as a function of m_{X^0}.

Search for Resonance X, Y in $e^+e^- \rightarrow XY$

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •				
	[275] ACKERSTAFF	98X	OPAL	$X \rightarrow$ 2jets, $Y \rightarrow$ 2jets
	[276] ACKERSTAFF	98Y	OPAL	$X \rightarrow \gamma\gamma$, $Y \rightarrow f\overline{f}$
	[277] ALEXANDER	97B	OPAL	$X \rightarrow$ 2 jets, $Y \rightarrow$ 2 jets
	[278] BUSKULIC,D	96	ALEP	$X \rightarrow$ 2 jets, $Y \rightarrow$ 2 jets

[275] ACKERSTAFF 98X search for $e^+e^- \rightarrow XY \rightarrow$ 4jets at $\sqrt{s}$= 130–184 GeV. The upper limits on $\sigma(e^+e^- \rightarrow XY)$, which are well below the excess reported by BUSKULIC,D 96, are shown in their Fig. 5.

[276] ACKERSTAFF 98Y search for $e^+e^- \rightarrow XY$, with $X \rightarrow \gamma\gamma$, $Y \rightarrow f\overline{f}$ where $f\overline{f}$ may be $q\overline{q}$, $\ell\overline{\ell}$, or $\nu\overline{\nu}$ at $\sqrt{s}$= 183 GeV. The upper limits on $\sigma(e^+e^- \rightarrow XY)\times$B($X \rightarrow \gamma\gamma$) are shown in their Fig. 4.

[277] ALEXANDER 97B search for the associated production of two massive particles decaying into quarks in e^+e^- collisions at $\sqrt{s}$=130–136 GeV. The 95%CL upper limits on $\sigma(e^+e^- \rightarrow XY)$ range from 2.7 to 4.5 pb for 95<m_X+m_Y < 120 GeV.

[278] BUSKULIC,D 96 observed an excess of four-jet production cross section in e^+e^- collisions at $\sqrt{s}$=130–136 GeV and find an enhancement in the sum of two dijet masses around 105 GeV.

Heavy Particle Production in Quarkonium Decays

Limits are for branching ratios to modes shown.

VALUE	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.5 \times 10^{-5}$	90	[279] BALEST	95	CLE2	$\Upsilon(1S) \rightarrow X^0\gamma$, $m_{X^0} < 5$ GeV
$< 3 \times 10^{-5}$–6×10^{-3}	90	[280] BALEST	95	CLE2	$\Upsilon(1S) \rightarrow X^0\overline{X}^0\gamma$, $m_{X^0} < 3.9$ GeV
$<5.6 \times 10^{-5}$	90	[281] ANTREASYAN	90C	CBAL	$\Upsilon(1S) \rightarrow X^0\gamma$, $m_{X^0} < 7.2$ GeV
		[282] ALBRECHT	89	ARG	

[279] BALEST 95 two-body limit is for pseudoscalar X^0. The limit becomes $< 10^{-4}$ for $m_{X^0} < 7.7$ GeV.

[280] BALEST 95 three-body limit is for phase-space photon energy distribution and angular distribution same as for $\Upsilon \rightarrow g g \gamma$.

[281] ANTREASYAN 90C assume that X^0 does not decay in the detector.

[282] ALBRECHT 89 give limits for B($\Upsilon(1S), \Upsilon(2S) \rightarrow X^0 \gamma$)·B($X^0 \rightarrow \pi^+ \pi^-$, $K^+ K^-$, $p\overline{p}$) for $m_{X^0} < 3.5$ GeV.

REFERENCES FOR Searches for Heavy Bosons Other Than Higgs Bosons

ABBIENDI	99	EPJ C6 1	G. Abbiendi+	(OPAL Collab.)
ABBOTT	98E	PRL 80 2051	B. Abbott+	(D0 Collab.)
ABBOTT	98J	PRL 81 38	B. Abbott+	(D0 Collab.)
ABE	98S	PRL 81 4806	F. Abe+	(CDF Collab.)
ACCIARRI	98J	PL B433 163	M. Acciarri+	(L3 Collab.)
ACKERSTAFF	98V	EPJ C2 441	K. Ackerstaff+	(OPAL Collab.)
ACKERSTAFF	98X	PL B429 399	K. Ackerstaff+	(OPAL Collab.)
ACKERSTAFF	98Y	PL B437 218	K. Ackerstaff+	(OPAL Collab.)
BARATE	98U	EPJ C4 571	R. Barate+	(ALEPH Collab.)
BARENBOIM	98	EPJ C1 369	G. Barenboim	
CHO	98	EPJ C5 155	G. Cho, K. Hagiwara, S. Matsumoto	
CHO	98B	NP B531 65	G. Cho, K. Hagiwara, Y. Umeda	
CVETIC	98	hep-ph/9707451	M. Cvetic, P. Langacker	
Perspectives in Supersymmetry				
DONCHESKI	98	PR D58 097702	M.A. Doncheski, R.W. Robinett	
ABBOTT	97B	PRL 79 4321	+Abolins, Acharya+	(D0 Collab.)
ABE	97F	PRL 78 2906	+Akimoto, Akopian, Albrow, Amendolia+	(CDF Collab.)
ABE	97G	PR D55 R5263	+Akimoto, Akopian, Albrow, Amendolia+	(CDF Collab.)
ABE	97S	PRL 79 2192	+Akimoto, Akopian, Albrow, Amendolia+	(CDF Collab.)
ABE	97W	PRL 79 3819	F. Abe+	(CDF Collab.)
ABE	97X	PRL 79 4327	+Akimoto, Akopian, Albrow, Amadon+	(CDF Collab.)
ACCIARRI	97Q	PL B412 201	M. Acciarri+	(L3 Collab.)
ALEXANDER	97B	ZPHY C73 201	G. Alexander+	(OPAL Collab.)
ARIMA	97	PR D55 19	+Odaka, Ogawa, Shirai, Tsuboyama+	(VENUS Collab.)
BARATE	97B	PL B399 329	+Buskulic, Decamp, Ghez, Goy, Lees+	(ALEPH Collab.)
BARENBOIM	97	PR D55 4213	+Bernabeu, Prades, Raidal	(VALE, IFIC)
DEANDREA	97	PL B409 277		(MARS)
DERRICK	97	ZPHY C73 613	M. Derrick+	(ZEUS Collab.)
GROSSMAN	97	PR D55 2768	+Ligeti, Nardi	(REHO, CIT)
JADACH	97	PL B408 281	+Ward, Was	(CERN, INPK, TENN, SLAC)
STAHL	97	ZPHY C74 73	A. Stahl, H. Voss	(BONN)
ABACHI	96C	PRL 76 3271	+Abbott, Abolins, Acharya, Adam+	(D0 Collab.)
ABACHI	96D	PL B385 471	+Abbott, Abolins, Acharya, Adam+	(D0 Collab.)
ABREU	96T	ZPHY C72 179	+Adam, Adye, Agasi, Ajinenko, Aleksan+	(DELPHI Collab.)
ADAM	96C	PL B380 471	+Adye, Agasi, Ajinenko, Aleksan+	(DELPHI Collab.)
AID	96B	PL B369 173	+Andreev, Andrieu, Appuhn, Arpagaus+	(H1 Collab.)
ALLET	96	PL B383 139	+Bodek, Camps, Deutsch+	(VILL, LEUV, LOUV, WISC)
BUSKULIC	96N	PL B378 373	+De Bonis, Decamp, Ghez, Goy, Lees+	(ALEPH Collab.)
BUSKULIC,D	96	ZPHY C71 179	D. Buskulic+	(ALEPH Collab.)
ABACHI	95E	PL B358 405	+Abbott, Abolins, Acharya, Adam, Adams+	(D0 Collab.)
ABACHI	95G	PRL 75 3618	+Abbott, Abolins, Acharya, Adam, Adams+	(D0 Collab.)
ABE	95	PR D51 R949	+Albrow, Amidei, Antos, Anway-Wiese+	(CDF Collab.)
ABE	95M	PRL 74 2900	+Albrow, Amidei, Antos, Anway-Wiese+	(CDF Collab.)
ABE	95N	PRL 74 3538	+Albrow, Amendolia, Amidei, Antos+	(CDF Collab.)
ABE	95U	PRL 75 1012	+Albrow, Amendolia, Amidei, Antos+	(CDF Collab.)
ABREU	95M	ZPHY C65 603	+Adam, Adye, Agasi, Ajinenko+	(DELPHI Collab.)
AID	95	PL B353 578	+Andreev, Andrieu, Appuhn, Arpagaus+	(H1 Collab.)
BALEST	95	PR D51 2053	+Cho, Ford, Johnson+	(CLEO Collab.)
KUZNETSOV	95	PRL 75 794	+Serebrov, Stepanenko+	(PNPI, KIAE, HARV, NIST)
KUZNETSOV	95B	PAN 58 2113	+Mikheev	(YARO)
		Translated from YAF 58 2228.		

MIZUKOSHI	95	NP B443 20	+Eboli, Gonzalez-Garcia	(SPAUL, CERN)
NARDI	95	PL B344 225	+Roulet, Tommasini	(MICH, CERN)
ABACHI	94B	PRL 72 965	+Abbott, Abolins, Acharya, Adam+	(D0 Collab.)
ABREU	94O	ZPHY C64 183	+Adam, Adye, Agasi, Ajinenko, Aleksan+	(DELPHI Collab.)
AHMED	94B	ZPHY C64 545	+Aid, Andreev, Andrieu, Appuhn, Arpagaus+	(H1 Collab.)
BHATTACH...	94	PL B336 100	Bhattacharyya, Ellis, Sridhar	(CERN)
Also	94B	PL B338 522 (erratum)	Bhattacharyya, Ellis, Sridhar	(CERN)
BHATTACH...	94B	PL B338 522 (erratum)	Bhattacharyya, Ellis, Sridhar	(CERN)
BUSKULIC	94	ZPHY C62 539	+Casper, De Bonis, Decamp, Ghez, Goy+	(ALEPH Collab.)
DAVIDSON	94	ZPHY C61 613	+Bailey, Campbell	(CFPA, TNTO, ALBE)
KUZNETSOV	94	PL B329 295	+Mikheev	(YARO)
KUZNETSOV	94B	JETPL 60 315	+Serebrov, Stepanenko+	(PNPI, KIAE, HARV, NIST)
		Translated from ZETFP 60 311.		
LEURER	94	PR D50 536		(REHO)
LEURER	94B	PR D49 333		(REHO)
Also	93	PRL 71 1324	Leurer	(REHO)
MAHANTA	94	PL B337 128		(MEHTA)
SEVERIJNS	94	PRL 73 611 (erratum)	+	(LOUV, WISC, LEUV, ETH, MASA)
VILAIN	94B	PL B332 465	+Wilquet, Beyer, Flegel, Grote+	(CHARM II Collab.)
ABE	93C	PL B302 119	+Amako, Arai, Arima, Asano, Chiba+	(VENUS Collab.)
ABE	93D	PL B304 373	+Adachi, Awa, Aoki, Belusevic, Emi+	(TOPAZ Collab.)
ABE	93G	PRL 71 2542	+Albrow, Akimoto, Amidei, Anway-Wiese+	(CDF Collab.)
ABE	93I	PR D48 R3939	+Albrow, Amidei, Anway-Wiese, Apollinari+	(CDF Collab.)
ABREU	93J	PL B316 620	+Adam, Adye, Agasi, Aleksan, Alekseev+	(DELPHI Collab.)
ABT	93	NP B396 3	+Andreev, Andrieu, Appuhn, Arpagaus+	(H1 Collab.)
ACTON	93E	PL B311 391	+Akers, Alexander+	(OPAL Collab.)
ADRIANI	93D	PL B306 187	+Aguilar-Benitez, Ahlen, Alcaraz, Aloisio+	(L3 Collab.)
ADRIANI	93M	PRPL 236 1	+Aguilar-Benitez, Ahlen, Alcaraz, Aloisio+	(L3 Collab.)
ALITTI	93	NP B400 3	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
ALLEN	93	PR D47 11	+Chen, Doe, Hausammann+	(UCI, LANL, ANL, UMD)
ALTARELLI	93B	PL B318 139	+Casalbuoni+	(CERN, FIRZ, GEVA, PADO)
BHATTACH...	93	PR D47 R3693	Bhattacharyya+	(CALC, JADA, ICTP, AHMED, BOSE)
BUSKULIC	93F	PL B308 425	+De Bonis, Decamp, Chez, Goy, Lees+	(ALEPH Collab.)
DERRICK	93	PL B306 173	+Krakauer, Magill, Musgrave, Repond+	(ZEUS Collab.)
RIZZO	93	PR D48 4470		(ANL)
SEVERIJNS	93	PRL 70 4047	+Gimeno-Nogues+	(LOUV, WISC, LEUV, ETH, MASA)
Also	94	PRL 73 611 (erratum)	Severijns+	(LOUV, WISC, LEUV, ETH, MASA)
STERNER	93	PL B303 385	+Abashian, Gotow, Haim, Mattson, Morgan+	(AMY Collab.)
ABE	92B	PRL 68 1463	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ABREU	92D	ZPHY C53 555	+Adam, Adami, Adye, Akesson, Alekseev+	(DELPHI Collab.)
ABREU	92F	PL B275 222	+Adam, Adami, Adye, Akesson, Alekseev+	(DELPHI Collab.)
ADRIANI	92F	PL B292 472	+Aguilar-Benitez, Ahlen, Akbari, Alcarez+	(L3 Collab.)
ALITTI	92E	PL B274 507	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
DECAMP	92	PRPL 216 253	+Deschizeaux, Goy, Lees, Minard+	(ALEPH Collab.)
DELAGUILA	92	NP B372 3	del Aguila+	(CERN, GRAN, MPIM, BRUXT, MADE)
Also	91C	NP B361 45	del Aguila, Moreno, Quiros	(BARC, MADE)
IMAZATO	92	PRL 69 877	+Kawashima, Tanaka+	(KEK, INUS, TOKY, TOKMS)
LANGACKER	92B	PR D45 278	+Luo	(PENN)
LAYSSAC	92	ZPHY C53 97	+Renard, Verzegnassi	(MONP, LAPP)
LAYSSAC	92B	PL B287 267	+Renard, Verzegnassi	(MONP, TRSTT)
LEIKE	92	PL B291 187	+Riemann, Riemann	(BERL, CERN)
LEP	92	PL B276 247	+ALEPH, DELPHI, L3, OPAL	(LEP Collabs.)
MISHRA	92	PRL 68 3499	+Leung, Arroyo+	(COLU, CHIC, FNAL, ROCH, WISC)
POLAK	92	PL B276 492	+Zralek	(SILES)
POLAK	92B	PR D46 3871	+Zralek	(SILES)
RENTON	92	ZPHY C56 355		(OXF)
ABE	91D	PRL 67 2418	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ABE	91F	PRL 67 2609	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ACTON	91	PL B268 122	+Alexander, Allison, Allport+	(OPAL Collab.)
ACTON	91B	PL B273 338	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ADEVA	91B	PL B261 169	+Adriani, Aguilar-Benitez, Akbari, Alcaraz+	(L3 Collab.)
ADEVA	91D	PL B262 155	+Adriani, Aguilar-Benitez, Akbari, Alcaraz+	(L3 Collab.)
ALEXANDER	91	PL B263 123	+Allison, Allport, Anderson, Arcelli+	(OPAL Collab.)
ALITTI	91	ZPHY C49 17	+Ansari, Ansorge, Autiero, Bareyre+	(UA2 Collab.)
ALTARELLI	91	PL B261 146	+Casalbuoni, De Curtis+	(CERN, FIRZ, GEVA)
ALTARELLI	91B	PL B263 459	+Casalbuoni, De Curtis+	(CERN, FIRZ, GEVA)
Also	90	PL B245 669	Altarelli, Casalbuoni, Feruglio, Gatto	(CERN, LECE, GEVA)
AQUINO	91	PL B261 280	+Fernandez, Garcia	(CINV, PUEB)
BUCHMUEL...	91	PL B267 395	Buchmueller, Greub, Minkowski	(DESY, BERN)

COLANGELO	91	PL B253 154	+Nardulli	(BARI)
CUYPERS	91	PL B259 173	+Falk, Frampton	(DURH, HARV, UNCCH)
DELAGUILA	91	PL B254 497	del Aguila, Moreno, Quiros	(BARC, MADE, CERN)
FARAGGI	91	MPL A6 61	+Nanopoulos	(TAMU)
GEIREGAT	91	PL B259 499	+Vilain, Wilquet, Binder, Burkard+	(CHARM II Collab.)
GONZALEZ-G...	91	PL B259 365	Gonzalez-Garcia, Valle	(VALE)
Also	90C	NP B345 312	Gonzalez-Garcia, Valle	(VALE)
MAHANTHAP...	91	PR D43 3093	Mahanthappa, Mohapatra	(COLO)
Also	91B	PR D44 1616 erratum	Mahanthappa, Mohapatra	(COLO)
POLAK	91	NP B363 385	+Zralek	(SILES)
RIZZO	91	PR D44 202		(WISC, ISU)
WALKER	91	APJ 376 51	+Steigman, Schramm, Olive+	(HSCA, OSU, CHIC, MINN)
ABE	90F	PL B246 297	+Amako, Arai, Asano, Chiba+	(VENUS Collab.)
ABE	90G	PRL 65 2243	+Amidei, Apollinari, Atac+	(CDF Collab.)
ABE	90H	PR D41 1722	+Amidei, Apollinari, Ascoli, Atac+	(CDF Collab.)
AKRAWY	90J	PL B246 285	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ALLEN	90	PRL 64 1330	+Chen, Doe+	(UCI, LASL, UMD)
ANTREASYAN	90C	PL B251 204	+Bartels, Besset, Bieler, Bienlein+	(Crystal Ball Collab.)
BARGER	90B	PR D42 152	+Hewett, Rizzo	(WISC, ISU)
GLASHOW	90	PR D42 3224	+Sarid	(HARV)
GLASHOW	90B	PRL 64 725	+Sarid	(HARV)
GONZALEZ-G...	90D	PL B240 163	Gonzalez-Garcia, Valle	(VALE)
GRIFOLS	90	NP B331 244	+Masso	(BARC)
GRIFOLS	90C	MPL A5 2657		(BARC)
GRIFOLS	90D	PR D42 3293	+Masso, Rizzo	(BARC, CERN, WISC, ISU)
HAGIWARA	90	PR D41 815	+Najima, Sakuda, Terunuma	(KEK, DURH, YCC, HIRO)
HE	90B	PL B240 441	+Joshi, Volkas	(MELB)
Also	90C	PL B244 580 erratum	He, Joshi, Volkas	(MELB)
KIM	90	PL B240 243	+Breedon, Ko, Lander, Maeshima, Malchow+	(AMY Collab.)
LOPEZ	90	PL B241 392	+Nanopoulos	(TAMU)
ALBAJAR	89	ZPHY C44 15	+Albrow, Allkofer, Arnison, Astbury+	(UA1 Collab.)
ALBRECHT	89	ZPHY C42 349	+Boeckmann, Glaeser, Harder+	(ARGUS Collab.)
BARBIERI	89B	PR D39 1229	+Mohapatra	(PISA, UMD)
DELAGUILA	89	PR D40 2481	del Aguila, Moreno, Quiros	(BARC, MADE)
Also	90B	PR D41 134	del Aguila, Moreno, Quiros	(BARC, MADE)
Also	90C	PR D42 262 erratum	del Aguila, Moreno, Quiros	(BARC, MADE)
DORENBOS...	89	ZPHY C41 567	Dorenbosch, Udo, Allaby, Amaldi+	(CHARM Collab.)
GEIREGAT	89	PL B232 539	+Vilain, Wilquet, Bergsma, Binder+	(CHARM II Collab.)
LANGACKER	89B	PR D40 1569	+Uma Sankar	(PENN)
ODAKA	89	JPSJ 58 3037	+Kondo, Abe, Amako+	(VENUS Collab.)
ROBINETT	89	PR D39 834		(PSU)
ALBAJAR	88B	PL B209 127	+Albrow, Allkofer, Astbury, Aubert+	(UA1 Collab.)
BAGGER	88	PR D37 1188	+Schmidt, King	(HARV, BOST)
BALKE	88	PR D37 587	+Gidal, Jodidio+	(LBL, UCB, COLO, NWES, TRIU)
BERGSTROM	88	PL B212 386		(STOH)
COSTA	88	NP B297 244	+Ellis, Fogli+	(PADO, CERN, BARI, WISC, LBL)
CUYPERS	88	PRL 60 1237	+Frampton	(UNCCH)
DONCHESKI	88	PL B206 137	+Grotch, Robinett	(PSU)
DONCHESKI	88B	PR D38 412	+Grotch, Robinett	(PSU)
ELLIS	88	PL B202 417	Ellis, Franzini, Zwirner	(CERN, UCB, LBL)
RAFFELT	88	PRL 60 1793	+Seckel	(UCB, LLL, UCSC)
AMALDI	87	PR D36 1385	+Bohm, Durkin, Langacker+	(CERN, AACH3, OSU+)
ANSARI	87D	PL B195 613	+Bagnaia, Banner+	(UA2 Collab.)
BARTEL	87B	ZPHY C36 15	+Becker, Felst+	(JADE Collab.)
MARCIANO	87	PR D35 1672	+Sirlin	(BNL, NYU)
ARNISON	86B	EPL 1 327	+Albrow, Allkofer+	(UA1 Collab.)
BARGER	86B	PRL 56 30	+Deshpande, Whisnant	(WISC, OREG, FSU)
BEHREND	86B	PL B178 452	+Buerger, Criegee, Fenner, Field+	(CELLO Collab.)
DERRICK	86	PL 166B 463	+Gan, Kooijman, Loos+	(HRS Collab.)
Also	86B	PR D34 3286	Derrick, Gan, Kooijman, Loos, Musgrave+	(HRS Collab.)
DURKIN	86	PL 166B 436	+Langacker	(PENN)
ELLIS	86	PL 167B 457	+Enqvist, Nanopoulos, Sarkar	(CERN, OXFTP)
JODIDIO	86	PR D34 1967	+Balke, Carr, Gidal, Shinsky+	(LBL, NWES, TRIU)
Also	88	PR D37 237 erratum	Jodidio, Balke, Carr+	(LBL, NWES, TRIU)
MOHAPATRA	86	PR D34 909		(UMD)
STEIGMAN	86	PL B176 33	+Olive, Schramm, Turner	(BART, MINN+)
ADEVA	85	PL 152B 439	+Becker, Becker-Szendy+	(Mark-J Collab.)
ADEVA	85B	PRL 55 665	+Becker, Becker-Szendy+	(Mark-J Collab.)
BERGER	85B	ZPHY C27 341	+Deuter, Genzel, Lackas, Pielorz+	(PLUTO Collab.)

STOKER	85	PRL 54 1887	+Balke, Carr, Gidal+	(LBL, NWES, TRIU)
ADEVA	84	PRL 53 134	+Barber, Becker, Berdugo+	(Mark-J Collab.)
BEHREND	84C	PL 140B 130	+Burger, Criegee, Fenner+	(CELLO Collab.)
ARNISON	83D	PL 129B 273	+Astbury, Aubert, Bacci+	(UA1 Collab.)
BERGSMA	83	PL 122B 465	+Dorenbosch, Jonker+	(CHARM Collab.)
CARR	83	PRL 51 627	+Gidal, Gobbi, Jodidio, Oram+	(LBL, NWES, TRIU)
DESHPANDE	83	PR D27 1193	+Johnson	(OREG)
BEALL	82	PRL 48 848	+Bander, Soni	(UCI, UCLA)
SHANKER	82	NP B204 375		(TRIU)
DIMOPOUL...	81	NP B182 77	Dimopoulos, Raby, Kane	(STAN, MICH)
RIZZO	81	PR D24 704	+Senjanovic	(BNL)
STEIGMAN	79	PRL 43 239	+Olive, Schramm	(BART, EFI)