

**12036**  
Olivine Basalt  
75 grams




Figure 1: Photo of 12036,0 showing coarse grain, vuggy nature. NASA #S69-63852.

### **Introduction**

Olivine basalt 12036 looks a lot like 12035 (figure 1), but has been studied by a different group of investigators. It has the same high modal olivine and pyroxene and the same chemical composition (within sampling error).

Busche et al. (1972) term this rock feldspathic peridotite with olivine poikilitically enclosed in pyroxene megacrysts up to 5 mm (figure 2). The megacrysts are incorporated into a second stage assemblage of olivine, plagioclase, pyroxene, spinel and accessory minerals.

### **Petrography**

Keil et al. (1971) describe 12036 as a coarse-grained cumulate containing abundant amounts of olivine, pigeonite, augite and chromite as cumulous phases.

Residual glass with high silica and high potassium is found interstitially and is associated with K-feldspar, fluroapatite, whitlockite and baddelyite.

---

#### **Mineralogical Mode for 12036**

	Neal et al. 1994	Dungan and Brown 1977	Busche et al. 1971
Olivine	24	24	24
Pyroxene	58	58	58
Plagioclase	12	12	12
Opaque			5
Ilmenite			
Chromite +Usp	5	5	
mesostasis			
melt inclusions			0.7




Figure 2: Photomicrograph of 12036,12 showing coarse mineral texture. Scale is 1.5 cm. NASA #S70-49400.

### **Mineralogy**

**Olivine:** Olivine in 12036 is  $Fo_{64} - Fo_{36}$ . (This is more iron rich than would be calculated for initial olivine to crystallize, figure 4).

**Pyroxene:** Busche et al. (1971) give the composition of pyroxene in 12036 (figure 4). Dungan and Brown (1977) compare the pyroxene in 12036 with 12005 (another apparent cumulate).

**Plagioclase:** Plagioclase in 12036 has more sodium ( $An_{85}$ ) and potassium.

**K-spar:** Keil et al. (1971) reported 3.7 % BaO in potassium feldspar in 12036.

**Spinel:** Busche et al. (1972) found two different trends in Cr-spinel in 12036. Jedwab (1971) studied the crystal growth of Ti-rich chromite growing in vugs.

**Whitlockite:** Keil et al. (1971) give detailed analysis of whitlockite in 12036.

**Baddeleyite:** Keil et al. (1971) reported the composition of four grains of baddeleyite.

**Metallic Iron:** The Ni content of iron grains in 12036 is high (up to 10%) and variable (Keil et al. 1971, figure 5).




Figure 3: Photomicrographs of thin section 12036,12 (plane-polarized; crossed-nicols). Scale is 2.2 mm. NASA #S70-49433-434.

### **Chemistry**

Rhodes et al. (1977) determined the major and minor element composition of 12036 (Figures 6 and 7). The sample has very high MgO content.

### **Radiogenic age dating**

12036 has not been dated.

### **Cosmogenic isotopes and exposure ages**

Burnett et al. (1975) determined an exposure age of  $165 \pm 15$  m.y. by  $^{81}Kr/^{83}Kr$ .

### **Processing**

12036 was broken, rather than sawed (figure 8).




Figure 4: Composition of olivine in 12036 (from Keil et al 1971).




Figure 6: Normalized rare-earth-element diagram for 12036 (data from Nyquist et al. 1977 and Rhodes et al. 1977).




Figure 5: Composition of iron grains in 12036 (Keil et al. 1971).




Figure 7: Composition of 12036 compared with that of other lunar basalts.

**List of Photo #s for 12036**

- S69-61586-61609 B&W
- S69-62318-62329 B&W
- S70-19148 B&W group
- S69-63847-63852 color
- S76-26866-26868 12036,1
- S70-49396-49420 thin section
- S70-49254-49257
- S70-49431-49436




Figure 8 : Processing diagram for 12036.

**Table 1. Chemical composition of 12036.**

<i>reference weight</i>	Rhodes77		Nyquist77	
SiO <sub>2</sub> %	43.11	(c)		
TiO <sub>2</sub>	3.2	(c)		
Al <sub>2</sub> O <sub>3</sub>	6.16	(c)		
FeO	21.82	(c)		
MnO	0.3	(c)		
MgO	16.71	(c)		
CaO	7.46	(c)		
Na <sub>2</sub> O	0.18	(a)		
K <sub>2</sub> O	0.06	(c)	0.061	(b)
P <sub>2</sub> O <sub>5</sub>	0.02	(c)		
S %	0.07	(c)		
<i>sum</i>				
Sc ppm	42.6	(a)		
V				
Cr	4880	(a)		
Co	63	(a)		
Ni	60	(a)		
Cu				
Zn				
Ga				
Ge ppb				
As				
Se				
Rb			1.08	(b)
Sr	91	(c)	94.5	(b)
Y	36	(c)		
Zr	97	(c)		
Nb	6.6	(c)		
Mo				
Ru				
Rh				
Pd ppb				
Ag ppb				
Cd ppb				
In ppb				
Sn ppb				
Sb ppb				
Te ppb				
Cs ppm				
Ba	56	(b)	56.3	(b)
La			5.03	(b)
Ce	14	(a)	14	(b)
Pr				
Nd			11.1	(b)
Sm	4.03	(a)	3.89	(b)
Eu	0.75	(a)	0.861	(b)
Gd			5.5	(b)
Tb	0.95	(a)		
Dy			6.36	(b)
Ho				
Er			3.76	(b)
Tm				
Yb	3.5	(a)	3.22	(b)
Lu	0.51	(a)	0.469	(b)
Hf	4.7	(a)		
Ta				
W ppb				
Re ppb				
Os ppb				
Ir ppb				
Pt ppb				
Au ppb				
Th ppm				
U ppm				
<i>technique</i>	<i>(a) INAA, (b) IDMS, (c) XRF</i>			