



Autism, asthma, inflammation, and the hygiene hypothesis

Kevin G. Becker *

Gene Expression and Genomics Unit, RRB, TRIAD Technology Center, National Institute on Aging, National Institutes of Health, Room 208, 333 Cassell Drive, Baltimore, MD 21224, United States

Received 30 January 2007; accepted 7 February 2007

Summary Inflammation and the genes, molecules, and biological pathways that lead to inflammatory processes influence many important and disparate biological processes and disease states that are quite often not generally considered classical inflammatory or autoimmune disorders. These include development, reproduction, aging, tumor development and tumor rejection, cardiovascular pathologies, metabolic disorders, as well as neurological and psychiatric disorders. This paper compares parallel aspects of autism and inflammatory disorders with an emphasis on asthma. These comparisons include epidemiological, morphometric, molecular, and genetic aspects of both disease types, contributing to a hypothesis of autism in the context of the immune based hygiene hypothesis. This hypothesis is meant to address the apparent rise in the prevalence of autism in the population.

© 2007 Elsevier Ltd. All rights reserved.

Introduction

Autism is an enigmatic childhood disorder of unknown origin. It is characterized by developmental, language, and social deficits, ranging in severity from patients with profound deficits to individuals that are high functioning. Although the underlying etiological basis of autism has eluded researchers, the genetic heritability of autism is quite strong [1]. Specifically what genes are involved and how they contribute to the disease phenotype is unclear.

Many theories regarding the biological basis of autism have been suggested, including neurodevel-

opmental, exposure to environmental toxins, particularly to mercury [2], and immune [3] hypotheses. More recently, theories of hyper-systemizing and assortative mating [4,5] and hyperdopamine [6] have been proposed. At this time there is little definitive evidence to support any single theory of the fundamental biological nature of autism.

Numerous reports have described imbalances in immune and inflammatory processes in autistic patients, including aberrations in antibody levels, cytokines, and cellular subsets [7–12]. Additionally, recent reports have described an increased frequency of HLA-A2 [13] and HLA-DR4 [14] antigens in autism. Interestingly, epidemiological studies have provided evidence for the association of asthma and allergies [15] or autoimmune disorders in families with autistic children [16–19]. The ex-

* Tel.: +1 410 558 8360; fax: +1 410 558 8281.
E-mail address: beckerk@grc.nia.nih.gov

act significance of immune abnormalities and the relationship of infections, immunizations, allergies, inflammation, or other aspects of immune response to disease etiology are unclear and controversial. Alterations of immune and inflammatory processes in autism have recently been reviewed [3,20–24].

One of the challenges in the early study of the molecular basis of classical autoimmune disorders was the attempt to establish the relevance of highly variable and fluctuating immune serum proteins and cell populations to disease etiology. That is, are fluctuations in any set of cytokines, immune mediators, or T cell populations, causative or are they epiphenomena due to peripheral effects of target tissue destruction, transient common infections, or more importantly, are they echoes of long ago infections. There is an ever present “which came first, the chicken or the egg” nature in the study of highly variable immune mediators. Are oligoclonal antibody bands found in the CSF of multiple sclerosis patients [25] related to the etiology of the disease or are they end stage phenomena? Do alterations in cytokines from a patient with systemic lupus erythematosus have a role in disease etiology or are they late stage responses to tissue destruction brought on by other mechanisms? Similarly, are immune aberrations in autism disease causing or are they epiphenomena?

Other comparisons of autism to asthma and autoimmune/inflammatory disorders

In addition to imbalances in immune molecular mediators, there are other seemingly unrelated parallels in the study of immune and inflammatory disorders as compared to autism that, when viewed collectively, may provide additional support for shared aspects of disease etiology between immune and inflammatory disorders and autism. These include; sex bias, birth order, age-of-onset, neonatal head circumference, increasing prevalence in the population, rural versus urban disease comparisons, and shared molecular and genetic markers.

Disease onset and sex bias

In asthma and in autism presentation is in early childhood. Both disease types have an age of onset in early childhood; 2–4 years for children with autistic disorder [26] and 3–6 for wheeze and asthma [27]. In addition, both autism and asthma display

a skewed sex bias toward boys. This bias is approximately 4:1 boys to girls in autism [1] and approximately 2:1 in asthma [28,29]. It is well known that in most adult autoimmune and inflammatory disorders, including asthma, there is a predominance of adult women with the diagnosis. However, less well known is that prior to puberty this skewing is toward boys [30]. This male bias prior to puberty may be true in other immune mediated disorders as well such as multiple sclerosis [31], Type 1 diabetes, and thyroiditis.

Birth order

Some studies have shown birth order to be relevant in atopic disorders as well as autism. In both cases, being first born may carry a greater risk for disease than later births. In a large study of 11,371 Italian young men those with no siblings had the highest level of serum IgE sensitization. An inverse association was observed between number of siblings at time of testing and prevalence of high atopy $P < 0.0001$ [32]. Similar findings have been shown in for atopic disease in Crete [33], asthma, eczema-urticaria and hay fever in Scotland [34], asthma with allergic rhinitis in Denmark [35] and asthma, allergy, and eczema in the Netherlands [36]. These observations are thought to be related to increased transmission of childhood infections due to a growing family size in the context of the hygiene hypothesis (see below). Similarly, the risk of autism has been shown in some cases to be related to birth order in the same direction as asthma and atopic disorders, with risk decreasing with a greater number of older siblings in the United States [37–39], Western Australia [40] and England [41].

Increased neonatal head circumference

Increased neonatal head circumference has been found in both autism and asthma. Increases in neonatal head circumference have been associated with asthma and atopy. In particular, head circumference has been associated with elevated serum IgE levels and hay fever disorders [42–45]. Increased neonatal head circumference or macrocephaly is a robust finding in autism with the largest effect between the ages of 2–5 [46–51]. This brain size difference is largely back to normal by adolescence. The biological basis for this increase is unknown although genetic, infectious, and inflammatory mechanisms have been proposed [50] (see PTEN below).

Increase in prevalence in the population: parallel “epidemics”

Both autism and asthma have had reports of apparent increases in the population over the last 30 years. Numerous studies show general increases in prevalence in both asthma [52,53] and autism [54–58], at roughly similar rates over the last 30 years. In both disease types this has been often referred to as an “epidemic” [54,59]. In both disease types this apparent increase is controversial. Changes in diagnostic classifications and access to health care resources have confounded the interpretation of prevalence estimates in the study of asthma and autism. Significant increases in disease prevalence over a short time in evolutionary terms suggest that purely genetic mechanisms may not be solely responsible [60]. Given the strong heritability of autism, changing environmental modifiers in the context of the background genetics of autism may be important over the past 30 years. There have been similar increases in the prevalence in classical autoimmune diseases over the same time span as well, including Type 1 diabetes [61,60,62].

Rural vs urban disease distribution

Both autism and asthma appear to show uneven geographical distributions in disease prevalence. Differential susceptibility or resistance to asthma and allergies is found in urban environments versus rural or farm environments [63–65]. Although the exact mechanistic basis of the difference is not known, this distribution pattern of disease is thought to have an inverse relationship to infection and is central to the hygiene hypothesis (see below).

The geographical distribution of autism is less clear although there is evidence that there may be an urban versus rural distribution. This has been found in epidemiological studies from multiple countries including Denmark [66], the United States [67], England [55,68] and Japan [69]. Interestingly, in studies of autism that analyzed numerous familial risk factors, a major risk factor for autism was increasing degree of urbanization [55,68]. In a study from the US, the urban versus rural distribution was attributable to mercury exposure in the environment, however this may reflect an industrialized versus rural pattern as well [67].

The Inuit of northern Canada may provide an interesting population case study. This isolated rural population exists in crowded living conditions, with high levels of mercury and other environmental toxins in the diet [70]. However, autism is

essentially non-existent in the Inuit. In a recent report Fombonne et al. state; No case of autism has ever been reported in an Inuit child in the past 15 years [71]. In parallel, asthma and atopic disorders are uncommon in Inuit children, even with very high rates of lower respiratory infections prior to age 2 and particularly high rates of childhood smoking (31.9%) [72].

Molecular and genetic markers shared with inflammatory/autoimmune diseases

Like many common human disorders, autism, asthma, and autoimmune disorders have been studied using genetic linkage and genetic association approaches. The chromosomal regions identified in linkage studies and the specific variants of genes identified in genetic association studies are quite often not unique to any one disorder. Many, if not most, genes in the human genome have broad based effects influencing different cells and tissues at different times of development under the influence of different environmental modifiers. In the context of common human disease, important regulatory genes may effect disease susceptibility differently when found in combination with different disease associated alleles [73].

ADRB2, beta(2)-adrenergic receptor

The gene for the beta(2)-adrenergic receptor encodes a member of the G protein-coupled receptor superfamily and is expressed on epithelial and endothelial cells of the lung, mast cells, as well as airway smooth muscle cells. ADRB2 activation is thought to work through increased intracellular cAMP levels [74]. Polymorphisms in ADRB2, including the Glu27 allele, have been studied in multiple disease states including hypertension [75], atopic dermatitis [76], Graves disease [77], rheumatoid arthritis [78,79], obesity [80] and in particular, asthma [81]. ADRB2 is of major interest in asthma as it may be involved in lung function as well as response to beta(2)-adrenergic agonists [82,83]. ADRB2 polymorphisms may not influence asthma incidence or prevalence but may influence persistence of asthmatic symptoms [84].

Importantly, the Glu27 allele of ADRB2 has recently been associated with autism in twins [85] as well as in the AGRE autism cohort [86].

PTEN-phosphatase and tensin homolog

PTEN, phosphatase and tensin homolog, is central to phosphoinositide metabolism as an important

regulatory checkpoint in the PI3K/ATK signaling pathway, effecting multiple downstream processes including immune function, cell growth, cell survival, and differentiation [87–89]. PTEN has been shown to play a role in lymphocyte proliferation, systemic autoimmunity, and autoimmune disease [87,88], as well as in benign tumors of the gastrointestinal tract. In relation to disease, PTEN has been implicated in bronchial asthma and allergic inflammation [90].

Interestingly, PTEN has been implicated in macrocephaly (OMIM # #153480) and Cowden disease (OMIM #158350). PTEN has been implicated in autism as well, in particular, within a subset of autistic individuals with macrocephaly [91,92]. A recent report described a patient with a PTEN mutation having autistic features, macrocephaly as well as nodular lymphoid hyperplasia of the small and large intestinal mucosa [93]. Moreover, a mouse model with specific deletions of PTEN in selected neuronal cell types resulted in macrocephaly, changes in social interactions, and increased responses to sensory stimuli, suggesting a model for autistic spectrum disorder [94].

MET – met proto-oncogene

The proto-oncogene MET, also known as hepatocyte growth factor receptor, encodes a tyrosine-kinase receptor which has been shown to have pleiotropic effects, in myocardial infarction, ischemia, angiogenesis, and importantly in cancer progression. Recently, polymorphic variants that result in reduced expression of MET has been genetically associated with autism [95]. MET also has been shown to effect the immune system [96], in particular it suppresses immune dendritic cell function [97]. In addition, c-MET and its ligand HGF have been shown to be involved in multiple neuronal processes including synaptic plasticity in the hippocampus [98], development of cortical pyramidal dendrites [99] and synaptic organization [100].

Genome wide scans

Genome wide scans are genetic linkage studies that use evenly spaced polymorphic markers that span the entire human genome in an attempt to link disease phenotypes to specific regions in the human genome. In a comparison of genome wide linkage studies between autoimmune and inflammatory disorders and similar studies in autism and Tourette syndrome, overlap of polymorphic markers were found to be statistically significant ($P = 0.01$) in chromosomal regions originally independently

identified in autism and Tourette's, or in autoimmune and inflammatory disorders. <http://www.grc.nia.nih.gov/branches/rrb/dna/pubs/cgoatad.pdf> This comparison was performed using the approach originally taken for autoimmune disorders [101]. Fig. 1 shows sixteen selected regions of the genome where this marker overlap occurs. A more comprehensive listing of marker overlap between autoimmune/inflammatory disorders and autism and Tourette syndrome can be found here: <http://www.grc.nia.nih.gov/branches/rrb/dna/atmap.htm> Moreover, a subset of these markers found to be statistically significant in both disease classes is not due to simple coincidental overlap of genetic regions, but includes 144 identical polymorphic markers originally found to be statistically significant in both autism and autoimmune or inflammatory disorders, including asthma. For example, in the chromosomal region 17q25.3, the polymorphic marker D17S784 has been independently linked to psoriasis [102], Crohn's disease [103], Tourette syndrome [104], and autism [105]. A listing of markers independently found in both disease classes can be found here: <https://www.quickbase.com/db/8qsiujvy>.

Summary of disease comparisons

The epidemiological, morphometric, molecular, and genetic comparisons between autism and inflammatory disorders stated above highlight multiple lines of evidence in addition to humoral and cellular immune abnormalities with the goal to strengthen an etiological relationship between autism and autoimmune and inflammatory disorders. It is not suggested that these comparisons support any direct link between these disorders. However, these shared observations between autism and inflammatory disorders are used in support of the development of a hypothesis for the apparent rise in the prevalence of autism using the framework of the immune hygiene hypotheses.

The hygiene hypothesis

The hygiene hypothesis is a widely held theory of the etiology of asthma and atopic disorders which builds on observations of rural versus urban distribution of disease. It suggests that cleaner environmental conditions in westernized countries, as compared to developing countries, play a role in the increase of the prevalence of these disorders in western countries [106]. Moreover, low levels




Figure 1 Selected clusters of linkage data from Autism, Tourette syndrome, and Autoimmune/Inflammatory disorders. All polymorphic markers come from independent genetic linkage whole genome scans. Each marker is positioned on a common reference map based on the LDB maps. Chromosome band and centimorgan position are shown above and below respectively for each chromosomal region. Polymorphic markers from autoimmune and inflammatory disorders are in black. Markers from autism are in orange. Markers from Tourette syndrome are in purple. All markers are arbitrarily assigned a 10-centimorgan interval at the peak marker reported. Line weight is proportional to LOD score of P -value. References and other information for all polymorphic markers can be found here: <https://www.quickbase.com/db/8jp3dz49?a=q&qid=1>.



Fig. 1 (continued)

of asthma and allergies are found with early exposure to cats [107,108], being raised in a farm environment [109] larger family size [110,35] day-care attendance [111] and birth order [32–36].

Risk for asthma and atopy may be due to a lack of early immune challenge of the post-natal immune system by microbial or parasitic infection possibly including environmental saprophytes and gut commensal organisms, relative to the developing innate immune system [112]. Alteration in the immune repertoire early in thymic development may lead to the establishment of immune hyper-

sensitivity ultimately leading to inflammatory pathology.

In certain ways, the hygiene hypothesis is counterintuitive, in that less clean polluted environments were once thought to cause asthma. Moreover, it is common practice in western society to “protect” children from bacteria and microorganisms through isolation indoors and through overuse of antibacterial soaps. This practice may be harmful in not allowing robust immune challenge in early neo-natal development.

The hygiene hypothesis is not without criticism. The changes in the prevalence of atopic disorders may have more complex etiologies with regard to overall microbial load or helminth infection in the general population [113–116] rather than with simple notions of personal or community hygiene practices.

Autism and the hygiene hypothesis

As compared above, similarities between autism, asthma, and inflammatory disorders raise the possibilities of shared mechanisms between these disease types. These include altered immune function in both types of disorders, a similar sex bias at diagnosis, similar birth order relationships, unexplained increased neonatal head circumference, a similar increase in prevalence rates during the last quarter century, a possible rural–urban distribution of the diagnosis with disease being more prevalent in urban environments, and shared molecular and genetic factors between autism and asthma. This adds multiple lines of evidence that mechanisms important in the etiology of immune and inflammatory processes may contribute to the etiology of autism.

It is proposed here that the hygiene hypothesis, a viable theory in the etiology of asthma, should be considered in the etiology of autism. Underlying factors important in the hygiene hypothesis, whether they are truly related to hygiene practices or to overall microbial or parasitic load, thought to be relevant to the increase in asthma and atopy, may contribute to the rise in the incidence of autism as well. Altered patterns of infant immune stimulation may hypersensitize the early immune system not toward allergic sensitivity and bronchial hypersensitivity but to inflammatory or cytokine responses affecting brain structure and function leading to autism. It is well documented that immune cytokines play an important role in normal brain development as well as pathological injury in early brain development [117,118]. It is hypothesized that immune pathways altered by hygiene practices in western society may effect brain structure or function contributing to the development of autism.

Acknowledgements

The author thank M. Pazin, A. Wurster, F. Indig, and B. Schauer for critical reading of the manuscript. This research was supported by the Intramu-

ral Research Program of the NIH, National Institute on Aging.

References

- [1] Folstein SE, Rosen-Sheidley B. Genetics of autism: complex aetiology for a heterogeneous disorder. *Nat Rev Genet* 2001;2(12):943–55.
- [2] Bernard S, Enayati A, Redwood L, Roger H, Binstock T. Autism: a novel form of mercury poisoning. *Med Hypotheses* 2001;56(4):462–71.
- [3] Korvatska E, Van de Water J, Anders TF, Gershwin ME. Genetic and immunologic considerations in autism. *Neurobiol Dis* 2002;9(2):107–25.
- [4] Baron-Cohen S. The hyper-systemizing, assortative mating theory of autism. *Prog Neuropsychopharmacol Biol Psychiatry* 2006.
- [5] Constantino JN, Lajonchere C, Lutz M, et al. Autistic social impairment in the siblings of children with pervasive developmental disorders. *Am J Psychiatry* 2006;163(2):294–6.
- [6] Previc FH. Prenatal influences on brain dopamine and their relevance to the rising incidence of autism. *Med Hypotheses* 2007;68(1):46–60.
- [7] Gupta S, Aggarwal S, Roshanravan B, Lee T. Th1- and Th2-like cytokines in CD4+ and CD8+ T cells in autism. *J Neuroimmunol* 1998;85(1):106–9.
- [8] Croonenberghs J, Wauters A, Devreese K, et al. Increased serum albumin, gamma globulin, immunoglobulin IgG, and IgG2 and IgG4 in autism. *Psychol Med* 2002;32(8):1457–63.
- [9] Ashwood P, Wakefield AJ. Immune activation of peripheral blood and mucosal CD3(+) lymphocyte cytokine profiles in children with autism and gastrointestinal symptoms. *J Neuroimmunol* 2006;173(1–2):126–34.
- [10] Zimmerman AW, Jyonouchi H, Comi AM, et al. Cerebrospinal fluid and serum markers of inflammation in autism. *Pediatr Neurol* 2005;33(3):195–201.
- [11] Molloy CA, Morrow AL, Meinzen-Derr J, et al. Elevated cytokine levels in children with autism spectrum disorder. *J Neuroimmunol* 2006;172(1–2):198–205.
- [12] Singer HS, Morris CM, Williams PN, Yoon DY, Hong JJ, Zimmerman AW. Antibrain antibodies in children with autism and their unaffected siblings. *J Neuroimmunol* 2006;178(1–2):149–55.
- [13] Torres AR, Sweeten TL, Cutler A, et al. The association and linkage of the HLA-A2 class I allele with autism. *Hum Immunol* 2006;67(4–5):346–51.
- [14] Lee LC, Zachary AA, Leffell MS, et al. HLA-DR4 in families with autism. *Pediatr Neurol* 2006;35(5):303–7.
- [15] Croen LA, Grether JK, Yoshida CK, Odouli R, Van de Water J. Maternal autoimmune diseases, asthma and allergies, and childhood autism spectrum disorders: a case-control study. *Arch Pediatr Adolesc Med* 2005;159(2):151–7.
- [16] Money J, Bobrow NA, Clarke FC. Autism and autoimmune disease: a family study. *J Autism Child Schizophr* 1971;1(2):146–60.
- [17] Sweeten TL, Bowyer SL, Posey DJ, Halberstadt GM, McDougle CJ. Increased prevalence of familial autoimmunity in probands with pervasive developmental disorders. *Pediatrics* 2003;112(5):e420.
- [18] Comi AM, Zimmerman AW, Frye VH, Law PA, Peeden JN. Familial clustering of autoimmune disorders and evaluation of medical risk factors in autism. *J Child Neurol* 1999;14(6):388–94.
- [19] Molloy CA, Morrow AL, Meinzen-Derr J, et al. Familial autoimmune thyroid disease as a risk factor for regression

- in children with autism spectrum disorder: a CPEA study. *J Autism Dev Disord* 2006.
- [20] Pardo CA, Vargas DL, Zimmerman AW. Immunity, neuroglia and neuroinflammation in autism. *Int Rev Psychiatry* 2005;17(6):485–95.
- [21] van Gent T, Heijnen CJ, Treffers PD. Autism and the immune system. *J Child Psychol Psychiatry* 1997;38(3):337–49.
- [22] Ashwood P, Van de Water J. Is autism an autoimmune disease? *Autoimmun Rev* 2004;3(7–8):557–62.
- [23] Ashwood P, Van de Water J. A review of autism and the immune response. *Clin Dev Immunol* 2004;11(2):165–74.
- [24] Cohly HH, Panja A. Immunological findings in autism. *Int Rev Neurobiol* 2005;71:317–41.
- [25] Link H, Huang YM. Oligoclonal bands in multiple sclerosis cerebrospinal fluid: an update on methodology and clinical usefulness. *J Neuroimmunol* 2006;180(1–2):17–28.
- [26] Mandell DS, Novak MM, Zubritsky CD. Factors associated with age of diagnosis among children with autism spectrum disorders. *Pediatrics* 2005;116(6):1480–6.
- [27] Morgan WJ, Stern DA, Sherrill DL, et al. Outcome of asthma and wheezing in the first 6 years of life: follow-up through adolescence. *Am J Respir Crit Care Med* 2005;172(10):1253–8.
- [28] Yunginger JW, Reed CE, O'Connell EJ, Melton 3rd LJ, O'Fallon WM, Silverstein MD. A community-based study of the epidemiology of asthma. Incidence rates, 1964–1983. *Am Rev Respir Dis* 1992;146(4):888–94.
- [29] Osman M, Tagiyeva N, Wassall HJ, et al. Changing trends in sex specific prevalence rates for childhood asthma, eczema, and hay fever. *Pediatr Pulmonol* 2007;42(1):60–5.
- [30] Osman M. Therapeutic implications of sex differences in asthma and atopy. *Arch Dis Child* 2003;88(7):587–590.
- [31] Haliloglu G, Anlar B, Aysun S, et al. Gender prevalence in childhood multiple sclerosis and myasthenia gravis. *J Child Neurol* 2002;17(5):390–2.
- [32] Matricardi PM, Franzinelli F, Franco A, et al. Sibship size, birth order, and atopy in 11,371 Italian young men. *J Allergy Clin Immunol* 1998;101(4 Pt 1):439–44.
- [33] Zekveld C, Bibakis I, Bibaki-Liakou V, et al. The effects of farming and birth order on asthma and allergies. *Eur Respir J* 2006;28(1):82–8.
- [34] Kinra S, Davey Smith G, Jeffreys M, Gunnell D, Galobardes B, McCarron P. Association between sibship size and allergic diseases in the Glasgow Alumni Study. *Thorax* 2006;61(1):48–53.
- [35] Westergaard T, Rostgaard K, Wohlfahrt J, Andersen PK, Aaby P, Melbye M. Sibship characteristics and risk of allergic rhinitis and asthma. *Am J Epidemiol* 2005;162(2):125–32.
- [36] Bernsen RM, de Jongste JC, van der Wouden JC. Birth order and sibship size as independent risk factors for asthma, allergy, and eczema. *Pediatr Allergy Immunol* 2003;14(6):464–9.
- [37] Deykin EY, MacMahon B. Pregnancy, delivery, and neonatal complications among autistic children. *Am J Dis Child* 1980;134(9):860–4.
- [38] Piven J, Simon J, Chase GA, et al. The etiology of autism: pre-, peri- and neonatal factors. *J Am Acad Child Adolesc Psychiatry* 1993;32(6):1256–63.
- [39] Croen LA, Grether JK, Selvin S. Descriptive epidemiology of autism in a California population: who is at risk? *J Autism Dev Disord* 2002;32(3):217–24.
- [40] Glasson EJ, Bower C, Petterson B, de Klerk N, Chaney G, Hallmayer JF. Perinatal factors and the development of autism: a population study. *Arch Gen Psychiatry* 2004;61(6):618–27.
- [41] Bolton PF, Murphy M, Macdonald H, Whitlock B, Pickles A, Rutter M. Obstetric complications in autism: consequences or causes of the condition? *J Am Acad Child Adolesc Psychiatry* 1997;36(2):272–81.
- [42] Katz KA, Pocock SJ, Strachan DP. Neonatal head circumference, neonatal weight, and risk of hayfever, asthma and eczema in a large cohort of adolescents from Sheffield, England. *Clin Exp Allergy* 2003;33(6):737–45.
- [43] Oryszczyn MP, Annesi-Maesano I, Campagna D, Sahuquillo J, Huel G, Kauffmann F. Head circumference at birth and maternal factors related to cord blood total IgE. *Clin Exp Allergy* 1999;29(3):334–41.
- [44] Gregory A, Doull I, Pearce N, et al. The relationship between anthropometric measurements at birth: asthma and atopy in childhood. *Clin Exp Allergy* 1999;29(3):330–3.
- [45] Leadbitter P, Pearce N, Cheng S, et al. Relationship between fetal growth and the development of asthma and atopy in childhood. *Thorax* 1999;54(10):905–10.
- [46] Piven J, Arndt S, Bailey J, Haverkamp S, Andreasen NC, Palmer P. An MRI study of brain size in autism. *Am J Psychiatry* 1995;152(8):1145–9.
- [47] Lainhart JE, Piven J, Wzorek M, et al. Macrocephaly in children and adults with autism. *J Am Acad Child Adolesc Psychiatry* 1997;36(2):282–90.
- [48] Redcay E, Courchesne E. When is the brain enlarged in autism? A meta-analysis of all brain size reports. *Biol Psychiatry* 2005;58(1):1–9.
- [49] Courchesne E, Pierce K. Brain overgrowth in autism during a critical time in development: implications for frontal pyramidal neuron and interneuron development and connectivity. *Int J Dev Neurosci* 2005;23(2–3):153–70.
- [50] Herbert MR. Large brains in autism: the challenge of pervasive abnormality. *Neuroscientist* 2005;11(5):417–40.
- [51] Lainhart JE, Bigler ED, Bocian M, et al. Head circumference and height in autism: a study by the collaborative program of excellence in autism. *Am J Med Genet A* 2006;140(21):2257–74.
- [52] Akinbami LJ, Schoendorf KC, Parker J. US childhood asthma prevalence estimates: the Impact of the 1997 National Health Interview Survey redesign. *Am J Epidemiol* 2003;158(2):99–104.
- [53] Akinbami LJ, Schoendorf KC. Trends in childhood asthma: prevalence, health care utilization, and mortality. *Pediatrics* 2002;110(2 Pt 1):315–22.
- [54] Gillberg C, Cederlund M, Lamberg K, Zeijlon L. Brief Report: The Autism Epidemic. The Registered Prevalence of Autism in a Swedish Urban Area. *J Autism Dev Disord* 2006.
- [55] Williams JG, Higgins JP, Brayne CE. Systematic review of prevalence studies of autism spectrum disorders. *Arch Dis Child* 2006;91(1):8–15.
- [56] Dales L, Hammer SJ, Smith NJ. Time trends in autism and in MMR immunization coverage in California. *JAMA* 2001;285(9):1183–5.
- [57] Gurney JG, Fritz MS, Ness KK, Sievers P, Newschaffer CJ, Shapiro EG. Analysis of prevalence trends of autism spectrum disorder in Minnesota. *Arch Pediatr Adolesc Med* 2003;157(7):622–7.
- [58] Newschaffer CJ, Falb MD, Gurney JG. National autism prevalence trends from United States special education data. *Pediatrics* 2005;115(3):e277–82.
- [59] Kippes C, Garrison CB. Are we in the midst of an autism epidemic? A review of prevalence data. *Mo Med* 2006;103(1):65–8.

- [60] Pitkaniemi J, Onkamo P, Tuomilehto J, Arjas E. Increasing incidence of Type 1 diabetes-role for genes? *BMC Genet* 2004;5:5.
- [61] Gillespie KM, Bain SC, Barnett AH, et al. The rising incidence of childhood type 1 diabetes and reduced contribution of high-risk HLA haplotypes. *Lancet* 2004;364(9446):1699–700.
- [62] Gardner SG, Bingley PJ, Sawtell PA, Weeks S, Gale EA. Rising incidence of insulin dependent diabetes in children aged under 5 years in the Oxford region: time trend analysis. The Bart's-Oxford Study Group. *BMJ* 1997;315(7110):713–7.
- [63] Adler A, Tager I, Quintero DR. Decreased prevalence of asthma among farm-reared children compared with those who are rural but not farm-reared. *J Allergy Clin Immunol* 2005;115(1):67–73.
- [64] Naleway AL. Asthma and atopy in rural children: is farming protective? *Clin Med Res* 2004;2(1):5–12.
- [65] Gold DR, Wright R. Population disparities in asthma. *Annu Rev Public Health* 2005;26:89–113.
- [66] Lauritsen MB, Pedersen CB, Mortensen PB. The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychol Med* 2004;34(7):1339–46.
- [67] Palmer RF, Blanchard S, Stein Z, Mandell D, Miller C. Environmental mercury release, special education rates, and autism disorder: an ecological study of Texas. *Health Place* 2006;12(2):203–9.
- [68] Lauritsen MB, Pedersen CB, Mortensen PB. Effects of familial risk factors and place of birth on the risk of autism: a nationwide register-based study. *J Child Psychol Psychiatry* 2005;46(9):963–71.
- [69] Hoshino Y, Kumashiro H, Yashima Y, Tachibana R, Watanabe M. The epidemiological study of autism in Fukushima-ken. *Folia Psychiatr Neurol Jpn* 1982;36(2):115–24.
- [70] Hermanson MH, Brozowski JR. History of Inuit community exposure to lead, cadmium, and mercury in sewage lake sediments. *Environ Health Perspect* 2005;113(10):1308–12.
- [71] Fombonne E, Morel J, Macarthur J. No Autism Amongst Inuits From Northern Quebec? Fifth International Meeting for Autism Research 2006 (Abstract PS6.4).
- [72] Hemmelgarn B, Ernst P. Airway function among Inuit primary school children in far northern Quebec. *Am J Respir Crit Care Med* 1997;156(6):1870–5.
- [73] Becker KG. The common variants/multiple disease hypothesis of common complex genetic disorders. *Med Hypotheses* 2004;62(2):309–17.
- [74] Johnson M. Molecular mechanisms of beta(2)-adrenergic receptor function, response, and regulation. *J Allergy Clin Immunol* 2006;117(1):18–24. quiz 25.
- [75] Ge D, Huang J, He J, et al. Beta(2)-adrenergic receptor gene variations associated with stage-2 hypertension in northern Han Chinese. *Ann Hum Genet* 2005;69(Pt 1):36–44.
- [76] Roguedas AM, Audrezet MP, Scotet V, Dupre-Goetghebeur D, Ferec C, Misery L. Intrinsic atopic dermatitis is associated with a beta(2)-adrenergic receptor polymorphism. *Acta Derm Venereol* 2006;86(5):447–8.
- [77] Jazdzewski K, Bednarczuk T, Stepnowska M, et al. Beta(2)-adrenergic receptor gene polymorphism confers susceptibility to Graves disease. *Int J Mol Med* 2007;19(1):181–6.
- [78] Xu B, Arlehag L, Rantapaa-Dahlquist SB, Lefvert AK. Beta(2)-adrenergic receptor gene single-nucleotide polymorphisms are associated with rheumatoid arthritis in northern Sweden. *Scand J Rheumatol* 2004;33(6):395–8.
- [79] Xu BY, Arlehag L, Rantapaa-Dahlquist SB, Lefvert AK. Beta(2)-adrenoceptor gene single nucleotide polymorphisms are associated with rheumatoid arthritis in northern Sweden. *Ann Rheum Dis* 2005;64(5):773–6.
- [80] Masuo K, Katsuya T, Kawaguchi H, et al. Beta(2)-adrenoceptor polymorphisms relate to obesity through blunted leptin-mediated sympathetic activation. *Am J Hypertens* 2006;19(10):1084–91.
- [81] Thakkestian A, McEvoy M, Minelli C, et al. Systematic review and meta-analysis of the association between beta(2)-adrenoceptor polymorphisms and asthma: a HuGE review. *Am J Epidemiol* 2005;162(3):201–11.
- [82] Taylor DR, Kennedy MA. Genetic variation of the beta(2)-adrenoceptor: its functional and clinical importance in bronchial asthma. *Am J Pharmacogenomics* 2001;1(3):165–74.
- [83] Hawkins GA, Tantisira K, Meyers DA, et al. Sequence, haplotype, and association analysis of ADRbeta(2) in a multiethnic asthma case-control study. *Am J Respir Crit Care Med* 2006;174(10):1101–9.
- [84] Hall IP, Blakey JD, Al Balushi KA, et al. Beta(2)-adrenoceptor polymorphisms and asthma from childhood to middle age in the British 1958 birth cohort: a genetic association study. *Lancet* 2006;368(9537):771–9.
- [85] Connors SL, Crowell DE, Eberhart CG, et al. Beta(2)-adrenergic receptor activation and genetic polymorphisms in autism: data from dizygotic twins. *J Child Neurol* 2005;20(11):876–84.
- [86] Cheslack-Postava K, Fallin MD, Avramopoulos D, et al. Beta(2)-adrenergic receptor gene variants and risk for autism in the AGRE cohort. *Mol Psychiatry* 2007.
- [87] Deleris P, Gayral S, Breton-Douillon M. Nuclear Ptdlns(3,4,5)P(3) signaling: An ongoing story. *J Cell Biochem* 2006;98(3):469–85.
- [88] Patel RK, Mohan C. PI3K/AKT signaling and systemic autoimmunity. *Immunol Res* 2005;31(1):47–55.
- [89] Gil A, Andres-Pons A, Pulido R. Nuclear PTEN: a tale of many tails. *Cell Death Differ* 2006.
- [90] Lee YC. The role of PTEN in allergic inflammation. *Arch Immunol Ther Exp (Warsz)* 2004;52(4):250–4.
- [91] Goffin A, Hoefsloot LH, Bosgoed E, Swillen A, Frys JP. PTEN mutation in a family with Cowden syndrome and autism. *Am J Med Genet* 2001;105(6):521–4.
- [92] Butler MG, Dasouki MJ, Zhou XP, et al. Subset of individuals with autism spectrum disorders and extreme macrocephaly associated with germline PTEN tumour suppressor gene mutations. *J Med Genet* 2005;42(4):318–21.
- [93] Boccone L, Dessi V, Zappu A, et al. Bannayan–Riley–Ruvalcaba syndrome with reactive nodular lymphoid hyperplasia and autism and a PTEN mutation. *Am J Med Genet A* 2006.
- [94] Kwon CH, Luikart BW, Powell CM, et al. Pten regulates neuronal arborization and social interaction in mice. *Neuron* 2006;50(3):377–88.
- [95] Campbell DB, Sutcliffe JS, Ebert PJ, et al. A genetic variant that disrupts MET transcription is associated with autism. *Proc Natl Acad Sci USA* 2006;103(45):16834–9.
- [96] Skibinski G. The role of hepatocyte growth factor/c-MET interactions in the immune system. *Arch Immunol Ther Exp (Warsz)* 2003;51(5):277–82.
- [97] Okunishi K, Dohi M, Nakagome K, et al. A novel role of hepatocyte growth factor as an immune regulator through suppressing dendritic cell function. *J Immunol* 2005;175(7):4745–53.
- [98] Akimoto M, Baba A, Ikeda-Matsuo Y, et al. Hepatocyte growth factor as an enhancer of NMDA currents and

- synaptic plasticity in the hippocampus. *Neuroscience* 2004;128(1):155–62.
- [99] Gutierrez H, Dolcet X, Tolcos M, Davies A. HGF regulates the development of cortical pyramidal dendrites. *Development* 2004;131(15):3717–26.
- [100] Tyndall SJ, Walikonis RS. The receptor tyrosine kinase MET and its ligand hepatocyte growth factor are clustered at excitatory synapses and can enhance clustering of synaptic proteins. *Cell Cycle* 2006;5(14):1560–8.
- [101] Becker KG, Simon RM, Bailey-Wilson JE, et al. Clustering of non-major histocompatibility complex susceptibility candidate loci in human autoimmune diseases. *Proc Natl Acad Sci USA* 1998;95(17):9979–84.
- [102] Tomfohrde J, Silverman A, Barnes R, et al. Gene for familial psoriasis susceptibility mapped to the distal end of human chromosome 17q. *Science* 1994;264(5162):1141–5.
- [103] Ma Y, Ohmen JD, Li Z, et al. A genome-wide search identifies potential new susceptibility loci for Crohn's disease. *Inflamm Bowel Dis* 1999;5(4):271–8.
- [104] Zhang H, Leckman JF, Pauls DL, Tsai CP, Kidd KK, Campos MR. Genomewide scan of hoarding in sib pairs in which both sibs have Gilles de la Tourette syndrome. *Am J Hum Genet* 2002;70(4):896–904.
- [105] Auranen M, Vanhala R, Varilo T, et al. A genomewide screen for autism-spectrum disorders: evidence for a major susceptibility locus on chromosome 3q25–27. *Am J Hum Genet* 2002;71(4):777–90.
- [106] Strachan DP. Hay fever, hygiene, and household size. *BMJ* 1989;299(6710):1259–60.
- [107] Karjalainen J, Virta M, Pessi T, Hulkkonen J, Nieminen MM, Hurme M. Childhood cat exposure-related tolerance is associated with IL1A and IL10 polymorphisms. *J Allergy Clin Immunol* 2005;116(1):223–5.
- [108] Fasce L, Tosca MA, Silvestri M, Olcese R, Pistorio A, Rossi GA. "Early" cat ownership and the risk of sensitization and allergic rhinitis in Ligurian children with respiratory symptoms. *Ann Allergy Asthma Immunol* 2005;94(5):561–5.
- [109] Kilpelainen M, Terho EO, Helenius H, Koskenvuo M. Farm environment in childhood prevents the development of allergies. *Clin Exp Allergy* 2000;30(2):201–8.
- [110] Karmaus W, Botezan C. Does a higher number of siblings protect against the development of allergy and asthma? A review. *J Epidemiol Community Health* 2002;56(3):209–17.
- [111] Celedon JC, Wright RJ, Litonjua AA, et al. Day care attendance in early life, maternal history of asthma, and asthma at the age of 6 years. *Am J Respir Crit Care Med* 2003;167(9):1239–43.
- [112] von Hertzen L, Haahtela T. Disconnection of man and the soil: reason for the asthma and atopy epidemic? *J Allergy Clin Immunol* 2006;117(2):334–44.
- [113] Yazdanbakhsh M, Kreamsner PG, van Ree R. Allergy, parasites, and the hygiene hypothesis. *Science* 2002;296(5567):490–4.
- [114] Ramsey CD, Celedon JC. The hygiene hypothesis and asthma. *Curr Opin Pulm Med* 2005;11(1):14–20.
- [115] Flohr C, Pascoe D, Williams HC. Atopic dermatitis and the 'hygiene hypothesis': too clean to be true? *Br J Dermatol* 2005;152(2):202–16.
- [116] Bloomfield SF, Stanwell-Smith R, Crevel RW, Pickup J. Too clean, or not too clean: the hygiene hypothesis and home hygiene. *Clin Exp Allergy* 2006;36(4):402–25.
- [117] Silverstein FS, Barks JD, Hagan P, Liu XH, Ivacko J, Szaflarski J. Cytokines and perinatal brain injury. *Neurochem Int* 1997;30(4–5):375–83.
- [118] Saliba E, Henrot A. Inflammatory mediators and neonatal brain damage. *Biol Neonate* 2001;79(3–4):224–7.

Available online at www.sciencedirect.com

