### 4. LAKE HURON

Lake Huron has only one U.S. Great Lakes AOC, the Saginaw River and Bay AOC.

## 4.1 SAGINAW RIVER AND BAY AOC, ARENAC, BAY, CLARE, GENESEE, GLADWIN, GRATIOT, HURON, IOSCO, ISABELLA, LAPEER, LIVINGSTON, MECOSTA, MIDLAND, MONTCALM, OGEMAW, OSCEOLA, ROSCOMMON, SAGINAW, SANILAC, SHIAWASSEE, AND TUSCOLA COUNTIES, MI

The Saginaw River and Bay AOC includes all of Saginaw Bay out to its interface with open Lake Huron at an imaginary line drawn between Au Sable Point and Point Aux Barques, as well as the entire 35 km length of the Saginaw River, which flows into Saginaw Bay (see AOC map in the appendix).

#### 4.1.1 Hazardous Waste Sites Relevant to the Saginaw River and Bay AOC

ATSDR has evaluated the data for hazardous waste sites in the 21 counties relevant to this AOC, and reached conclusions regarding the public health threat posed by these sites. These conclusions, along with information regarding the type and location of the site, and the date and type of assessment document, are summarized in Table 4-1, for sites that had public heath hazard categories of 1-3 at some point during their assessment history, and all NPL sites. Not all counties had waste sites in these categories.

For hazardous waste sites in the relevant Michigan counties that *at any time* had Public Health Hazard Categories of 1-3, the number of contaminant records in HazDat that exceeded health based-screening values was 1550, as shown in Table 4-2. Most of the records were for the soil media group; the water media group had the next highest number of records.

The IJC Great Lakes critical pollutants accounted for 197 (13%) of these records, with the majority for the soil media group. The specific IJC critical pollutants whose concentrations exceeded health-based screening values are: PCBs, TCDD, B(A)P, DDT and metabolites, aldrin, dieldrin, lead, mercury, and hexachlorobenzene. Details are provided in Table 4-2.

Further evaluations of the data for the sites with Public Health Hazard Categories of 1-3, as conducted by ATSDR in the Public Health Assessment and other health-related documents listed in the table, are discussed in the following subsections.

#### 4.1.1.1 Bay City Middlegrounds

The Bay City Middlegrounds site is an abandoned 40-acre landfill located on Middleground Island in the Saginaw River in southwestern Bay City, Bay County, MI. It operated as a landfill from 1956 to 1984. The landfill is partially capped, and has a leachate collection system. The cap was not fully sealed to the lower cap, and leachate has seeped out into ditches along the nearby roads. It was fenced on three sides, but not on the fourth, which borders the river, at the time that the 1996 health assessment was prepared by ATSDR. Information on this site is taken from that health assessment and from the 2003 EPA NPL fact sheet for this site.

	Public Health Hazard			
Site Name, County	Category	EPA NPL Status	Site ID	City
Bay City Middlegrounds, Bay	2 (1996 HA)	Proposed	MID981092935	Bay City
Keit Property, Bay	3 (1998 HC)	Non NPL	MISFN0507867	Bay City
Clare Water Supply, Clare	3 (1989 HA) 3 (n.d. SR)	Final	MID980002273	Clare
Berlin and Farrow, Genesee	3 (1985 HA) 2 (1992 HA)	Deleted Post SARA	MID000605717	Swartz Creek
Forest Waste Products, Genesee	3 (1988 HA) 3 (1994 HA)	Final	MID980410740	Otisville
Gratiot County Landfill, Gratiot	3 (1982 HA) 4 (n.d. SR)	Final	MID980506281	St. Louis
Velsicol Chemical Corp., Gratiot	3 (1988 HA) 3 (n.d. SR)	Final	MID000722439	St. Louis
Hedblum Industries, losco	3 (1989 HA)	Final	MID980794408	Oscoda
Wurtsmith Air Force Base, losco	4 (2001 HA)	Proposed		
Metamora Landfill, Lapeer	3 (1989 HA) 3 (1992 HA) 4 (n.d. SR)	Final	MID980506562	Metamora
Spiegelberg and Rasmussen Dump Sites, Livingston	3 (1989 HA) 2 (1992 HA)	Final	MID980794481, MID095402210	Brighton
Shiawassee River, Livingston	3 (1989 HA) 2 (n.d. SR)	Final	MID980794473	Howell
Dow Chemical Co., MI Div., Midland Loc, Midland	3 (2002 HC)	Non NPL	MID000724724	Midland
Tittabawassee River, Saginaw	3 (2002 HC)	Non NPL	MID980994354	Midland
Lufkin Rule, Saginaw	2 (1997 HC)	Non NPL	MID985584598	Saginaw
Laingsburg, Shiawassee	3 (2000 HC)	Non NPL	MISFN0507944	Laingsburg

Table 4-1. Hazardous Waste Sites in Counties Relevant to the Saginaw River and Bay AOC

2 = Public Health Hazard, 3 = Indeterminate Public Health Hazard, 4 = No Apparent Public Health Hazard HA = Public Health Assessment, HC = Health Consultation SR = Site Review and Update n.d. = no date provided

**Category of Public Health Hazard:** This site was categorized as a *Public Health Hazard* (category 2) because the surface soil contains organic chemicals and metals at concentrations potentially of human health concern, and trespassing occurs. In addition, the PCB-containing discharge from the landfill contributes to PCB contamination in the fish of the Saginaw River.

**Contaminants of Concern in Completed Exposure Pathways:** Many organic and inorganic chemicals, including the IJC critical pollutants PCBs, B(a)P, aldrin, dieldrin, hexachlorobenzene, lead, and mercury, are found in on-site soil, groundwater, and sediment at concentrations exceeding healthbased screening values. Actual exposure doses, however, are not likely to be of health concern for trespassers on the site, except upon frequent or prolonged exposure, which is not likely to occur. Soil and sediment concentrations of PAHs, including B(a)P, were typical of urban soils. Lead concentrations in soil were well above background, but lower than 400 ppm. Groundwater contaminated with PCBs discharges from the site into the Saginaw River, and PCBs have been found in the river water and sediment, at higher concentrations downstream of the site than upstream. Methylene chloride, detected in air at concentrations of human health concern including upwind of the site, may not be site-related.

Bioaccumulation of the PCBs through the food chain into fish that are ingested by humans is considered a pathway of great concern. Although this site is not the only source of PCBs discharged to the river, it contributes to the contamination, and levels of PCBs in fish are high enough to pose a risk of adverse health effects. This site was proposed for the NPL in 1995.

**Demographics:** Demographic profile, from the 2000 U.S. Census, for vulnerable populations living within 1 mile of this site:

Children 6 years and younger	793
Females aged 15-44	1,662
Adults 65 and older	1,000

**Public Health Outcome Data:** The Michigan Department of Community Health evaluated cancer incidence data for the area because of community health concerns. Statistics for 1990-1992 showed no statistically significant difference in cancer incidence or mortality between Bay County and Michigan as a whole. Cancer incidence for the zip code area including the site and Bay City west of the Saginaw River (48706) and for the zip code area including Bay City east of the Saginaw River (48708) for 1990 through 1993 indicate a slight, statistically significant elevation in incidence and rate for the entire period 1990-1993 (but not for any single year) in 48706, as compared with age- and sex-specific incidence rats for Michigan. None of the incidences or rates for 48708 were statistically significantly increased.

**Conclusions:** This site has contributed and continues to contribute to the environmental burden of the IJC critical pollutants PCBs, which are discharging from the landfill into the Saginaw River. PCBs are the major concern. PCB concentrations in fish are high enough to pose a health threat, and although this site is not the only contributor, concentrations in the river water and sediment are higher downstream than upstream of the site.

#### 4.1.1.2 Keit Property

The Keit property is approximately 18 acres of wetlands, grasslands, and woods in southwest Bay City, Bay Count, MI. It was used for agriculture since 1886. A large portion of the property has been filled in with material reportedly generated during a sewer project in the 1980s. ATSDR performed a health consultation on this site in 1998, as part of a Brownfields project; the information regarding this site is taken from that report, and from HazDat.

**Category of Public Health Hazard:** This site was categorized as an *Indeterminate Public Health Hazard* (category 3) because of the potential threat to human health from exposure to asbestos if the asbestos panels are not been removed from the property before it is used for a park.

**Contaminants of Concern in Completed Exposure Pathways:** None identified. Soil concentrations of the IJC critical pollutant B(a)P exceeded health based screening values in a few locations, but were considered typical for urban soils. Subsurface soil in one location contained PCBs above health-based screening values, but this was not a generalized finding, and surface soil concentrations of PCBs were not of concern. The primary hazard was a pile of Transite panels,

containing 40% chrysotile asbestos. If the panels are allowed to weather or are handled improperly, they could release asbestos fibers.

**Demographics:** Not reported.

Public Health Outcome Data: Not reported.

**Conclusions:** This site does not appear to have contributed significantly to the environmental burden of IJC critical pollutants or other chemicals, or to direct human exposure at levels that currently pose a health risk.

#### 4.1.1.3 Clare Water Supply

This site, a municipal water supply wellfield, is located in Clare, Clare County, MI. Information regarding this site is taken from the 1989 ATSDR preliminary public health assessment, HazDat, and the 2003 EPA NPL fact sheet for this site.

**Category of Public Health Hazard:** ATSDR categorized this site as an *Indeterminate Public Health Hazard* (category 3) in 1989 because of the potential threat to human health from exposure to municipal water containing VOCs, and the lack of up-to-date data to determine whether cleanup efforts have ameliorated the hazard. A subsequent ATSDR site review and update (not provided for inclusion in this document) also concluded that the site was a category 3.

**Contaminants of Concern in Completed Exposure Pathways:** No IJC critical pollutants were discussed. In 1985, VOCs, including trichloroethylene and other chlorinated compounds, and also benzene and xylenes were present in the groundwater used as the municipal water supply at concentrations that pose a public health concern Past completed exposure pathways included ingestion, inhalation, and dermal contact with the water. Contaminated soil from the suspected industrial sites northwest of the wellfield was then removed, but updated monitoring data were not available at the time of the 1989 health assessment. Air strippers were installed in 1991 and are removing over 90% of the volatile contaminants from the water supply. Additional remedial action, including groundwater collection and extraction initiated in 1996, soil vapor extraction initiated in 1999, and *in situ* ozonation of groundwater hot spots, is expected to continue for an indefinite number of years until cleanup goals are achieved.

**Demographics:** Demographic profile, from the 2000 U.S. Census, for vulnerable populations living within 1 mile of this site:

Children 6 years and younger	323
Females aged 15-44	718
Adults 65 and older	640

Public Health Outcome Data: Not reported.

**Conclusions:** This site has contributed to the environmental burden of VOCs and to human exposure. Remediation efforts, which are ongoing, are minimizing current and future impacts.

#### 4.1.1.4 Berlin and Farrow

This 40-acre site, located in Gaines Township near Swartz Creek, Genesee County, MI, was used as a licensed waste incinerator from 1971 to 1978. Violations included construction and operation of unlicensed waste lagoons and underground storage tanks, and burial of liquid wastes. Information regarding this site is taken from the 1992 interim public health assessment performed by ATSDR, HazDat, and the 2003 EPA NPL fact sheet. Cleanup activities prior to 1992 included removal of contaminated lagoon sludges and soils, removal of underground tanks and their contents, and removal of barrels. The chemicals in these materials included organochlorine intermediates (hexachlorobenzene, hexachlorcyclopentadienes, and octachlorocyclopentene) used in the production of certain pesticides, and also PCBs, benzene, and ethylbenzene.

**Category of Public Health Hazard:** In an early (1985) health assessment (not provided for inclusion in this document), ATSDR categorized this site as an *Indeterminate Public Health Hazard* (category 3). In the 1992 interim public health assessment, this site was categorized as a *Public Health Hazard* (category 2) because of the risk to human health from exposure to contaminants that may result in adverse health effects. Since that time, however, remediation of the site has been completed.

**Contaminants of Concern in Completed Exposure Pathways:** Not explicitly discussed. Onsite soil and sediment contained high levels of the IJC critical pollutant hexachlorobenzene; on-site surface water also was contaminated. Off-site sediment in the Slocum Drain, a stream draining the site, was contaminated with high levels of hexachlorobenzene. Off-site garden soil also contained hexachlorobenzene, but at much lower concentrations. Comparisons with health-based screening values were not presented for hexachlorobenzene. On-site soil and groundwater were contaminated with VOCs, including vinyl chloride and benzene at levels of concern for human health. PCBs, found in drums removed from the site, were not detected in sampling of site media as reported in the 1992 ATSDR health assessment. None of the off-site residential wells were contaminated with any of these compounds. From 1995 to 1996 final cleanup of the site was accomplished by excavation and removal of all remaining contaminated materials (soils, sediments, and aquifer materials), for disposal in a RCRA landfill. The site meets standards for unrestricted use and was deleted from the NPL in 1998.

**Demographics:** Demographic profile, from the 2000 U.S. Census, for vulnerable populations living within 1 mile of this site:

Children 6 years and younger	54
Females aged 15-44	148
Adults 65 and older	55

**Public Health Outcome Data:** Health surveys of households within the approximately 2 square miles surrounding the site were conducted in 1981. Surveys of a random sampling of the population, followed by a survey of all 122 households (418 people) found that respiratory symptoms were statistically significantly higher among persons who reported exposure to incinerator smoke as compared with those who reported no exposure to incinerator smoke. This information may suggest that the former industrial activity (incineration of hazardous wastes) at the site was potentially linked to health problems, but it does not provide insight into the potential health hazard from waste site-related contaminants.

Laboratory analyses of blood samples from 52 local residents revealed the presence of PCBs, DDT, and DDE at concentrations within the ranges generally found in Michigan residents, and thus, do not indicate a specific impact from the waste site.

**Conclusions:** In the past, this site contributed to the environmental burden of the IJC critical pollutant hexachlorobenzene, as well as other contaminants including VOCs. Although PCBs had been found in barrels at the site, they were not detected in environmental media. As reported in the EPA fact sheet, remediation of the site through removal and proper disposal of all contaminated materials and media is complete, and the site was deleted from the NPL list in 1998.

#### 4.1.1.5 Forest Waste Products

This 112-acre site is located 2 miles northwest of Otisville, Genesee County, MI. It includes an 11-acre landfill, which accepted general refuse and industrial and liquid waste from 1972 to 1978. Nine lagoons in another area of the site, covering a total of about 1 acre, also were used for disposal of industrial waste. Wastes included sludge and residues from a chemical warehouse fire, roofing material contaminated with PCBs, and cattle feed contaminated with PBBs. In 1978, the landfill was covered with soil. An estimated 3,000 waste drums may have been buried in the landfill. As of ATSDR's 1994 public health assessment, the site had been fenced, the waste material in the lagoons excavated and disposed off-site, and some of the drums removed and disposed off-site. The information on this waste site is taken from the 1994 ATSDR public health assessment, HazDat, and the 2003 EPA NPL fact sheet for this site.

**Category of Public Health Hazard:** This site was categorized as an *Indeterminate Public Health Hazard* (category 3) in a 1988 health assessment (not provided for inclusion in this document) as well as in the 1994 health assessment. The rationale in the 1994 assessment was that although human exposure did not appear to be occurring at levels of concern, uncertainty exists regarding the large number of drums reported to be buried in the landfill, which may release their contents into the environment.

**Contaminants of Concern in Completed Exposure Pathways:** The only chemical found at levels of concern for human health in a completed exposure pathway was arsenic, found in residential wells near the site, but the arsenic was thought to be of natural origin, rather than site-related. Although the IJC critical pollutant PCBs were disposed at the site, and formerly were found in subsurface soil, the PCBs had been removed during earlier remediation actions. Several contaminants were found at levels of concern in potential exposure pathways, including the IJC critical pollutant B(a)P, but the concentration of B(a)P in soil was not significantly above that commonly found in agricultural soil. VOCs have been found in groundwater, but not in residential wells. The IJC critical pollutants DDT and metabolites were found in game captured at the site boundary, but these contaminants were not site-related.

**Demographics:** Demographic profile, from the 2000 U.S. Census, for vulnerable populations living within 1 mile of this site:

Children 6 years and younger	34
Females aged 15-44	81
Adults 65 and older	48

**Public Health Outcome Data:** Health outcome data were not evaluated in conducting the 1994 public health assessment because there were no indications that humans had been significantly exposed to site-related contaminants, and no record of community reports of illnesses or health effects associated with the site.

**Conclusions:** Whether this site contributed to environmental contamination with IJC critical pollutant PCBs in the past is uncertain. No current exposure of humans to site-related contaminants at levels of concern is known to be occurring. Additional remediation activities, as described in the EPA NPL fact

sheet, included excavation and removal of buried drums and associated contaminated soil, and installation of a landfill cap. Monitoring of groundwater continues, particularly of a plume of VOCs that is migrating northward off the property.

#### 4.1.1.6 Gratiot County Landfill

This 40-acre landfill site is located southeast of St. Louis, in Gratiot County, MI. This landfill accepted general refuse, but was owned by a chemical corporation (Michigan Chemical Company), and disposed of chemicals wastes, including 269,000 pounds of PBBs, prior to 1977. The information regarding this site is taken from the 1982 ATSDR health assessment, HazDat, and the 2003 EPA NPL fact sheet for this site.

**Category of Public Health Hazard:** This site was categorized as an *Indeterminate Public Health Hazard* (category 3) in the 1982 health assessment, which focused on PBBs because of the potential threat to human health from exposure to contaminants and the poor quality of the support document that presented the monitoring data. A subsequent site review and update by ATSDR (not provided for inclusion in this report) ranked the site as *No Apparent Health Hazard* (category 4), possibly because remedial activities had mitigated the hazard. Remedial actions in 1984 included construction of a slurry wall and clay cap, and regrading of the landfill to minimize migration of contaminants from the landfill. In 1992, monitoring of the effectiveness of these remedies indicated that the slurry wall was ineffective in halting groundwater flow, and that VOCs (but not PBBs) were detected outside the slurry wall. A groundwater extraction system, constructed in 1998, appears to be effective in containing the plume. Further options are being evaluated by the Michigan Department of Environmental Quality, according to the EPA NPL fact sheet.

**Contaminants of Concern in Completed Exposure Pathways:** Not reported. The 1982 health assessment by ATSDR was primarily a review of a technical report regarding potential control strategies for the PBB contamination at the site. PBBs were detected in groundwater at concentrations above health-based screening values. No IJC critical pollutants were mentioned in the health assessment or the NPL fact sheet, but VOCs were apparently released from the site.

**Demographics:** Demographic profile, from the 2000 U.S. Census, for vulnerable populations living within 1 mile of this site:

Children 6 years and younger	170
Females aged 15-44	390
Adults 65 and older	252

Public Health Outcome Data: None reported.

**Conclusions:** This site may have contributed to the environmental burden of PBBs and VOCs in the past. The PBBs and VOCs have not been removed, but rather are contained by a slurry wall, cap, and groundwater extraction system.

#### 4.1.1.7 Velsicol Chemical

This 52-acre site is located in the City of St. Louis, Gratiot County, MI, and is surrounded on three sides by the Pine River, which drains into the Tittabawasse River, which joins the Saginaw River near the city of Saginaw. Velsicol Chemical Corporation, previously known as the Michigan Chemical Company, produced a variety of chemicals, including PBBs and DDT, at the Velsicol Chemical site plant from 1936 to 1978. Velsicol completed construction of a containment system at this site in 1985. This system consisted of a slurry wall around the entire site and a clay cap over the site. Information regarding this site is taken from the 1988 ATSDR preliminary health assessment, HazDat, and the 2003 EPA NPL fact sheet.

**Category of Public Health Hazard:** In 1988, ATSDR categorized this site as an *Indeterminate Public Health Hazard* (category 3) because exposure to PBBs through the food chain (fish and wildlife) has occurred and may possibly still be occurring, even though a fish consumption advisory was issued. A subsequent site review and update (not provided for inclusion in this document) also placed the site in this health hazard category.

**Contaminants of Concern in Completed Exposure Pathways:** None identified in the 1988 health assessment. The potential exposure of concern was to PBBs bioaccumlated in fish and wildlife. ATSDR noted that fish and river water and sediment concentrations of PBBs were declining. Subsequent developments included deterioration of the slurry wall in 1994, admitting water into the containment system; discovery of very high levels of DDT and metabolites in sediment of the Pine River/St. Louis impoundment; and the migration of dense non-aqueous phase liquids (DNAPL) from the containment area into the glacial till underlying the river sediments. The sediment and DNAPL are being removed and treated, according to the EPA NPL fact sheet.

**Demographics:** Demographic profile, from the 2000 U.S. Census, for vulnerable populations living within 1 mile of this site:

Children 6 years and younger	365
Females aged 15-44	821
Adults 65 and older	676

**Public Health Outcome Data:** In 1976, the Michigan Department of Public Health recruited many Velsicol workers for a PBB health study, which placed workers and their families in a registry to study the long-term effects of PBB exposure. The study, conducted in cooperation with the CDC, FDA, and EPA, was in operation at the time of the 1988 health assessment. No conclusions were reported.

**Conclusions:** This site has contributed to the environmental burden of the IJC critical pollutant DDT and metabolites, and also PBBs, with particular impacts on the Pine River/St. Louis impoundment sediments and fish. According to the EPA NPL fact sheet, although PBB concentrations are declining, DDT and metabolite concentrations in sediment are not. Remediation is underway. In addition, dense non-aqueous phase liquids have migrated from the site into the glacial till under the river sediments and are also being remediated.

#### 4.1.1.8 Hedblum Industries

The Hedblum Industries site is a 10-acre parcel located in Oscoda, Iosco County, MI, 1.2 miles west of Lake Huron. The site was leased to a series of industrial forms that manufactured parts for the automotive industry. Waste chemicals, including an estimated 4,000 gallons of spent trichloroethylene from a degreasing operation, were dumped in a pit near the main building. A pipe connecting an underground storage tank for trichloroethylene leaked. A number of residential wells in the area were found to be contaminated in 1973-1977. Most of the residents in the area of contamination were connected to municipal water in 1978, but a number were not. Trichloroethylene also was found in the

bayou into which groundwater from the site discharges; the bayou feeds the Au Sable River. The information regarding this site is taken from the 1989 ATSDR health assessment and the 2003 EPA NPL fact sheet for this site.

**Category of Public Health Hazard:** This site was categorized as an *Indeterminate Public Health Hazard* (category 3) because of the potential threat to human health from exposure to trichloroethylene and other VOCs.

**Contaminants of Concern in Completed Exposure Pathways:** None identified. No IJC critical pollutants are associated with this site. TCE has been identified in residential well water; eight households were estimated to have used contaminated well water at their household for an indeterminate time before they were switched to municipal water, but data were not adequate to measure the risks. Some residents still have not switched to municipal water, and others use well water for gardens and lawns. Exposure pathways include ingestion, dermal contact, and inhalation of trichloroethylene volatilized from the water. As of 1990, no VOC contaminants were detected in residential wells. The groundwater is being treated by a system constructed in 1992.

**Demographics:** Demographic profile, from the 2000 U.S. Census, for vulnerable populations living within 1 mile of this site:

Children 6 years and younger	135
Females aged 15-44	331
Adults 65 and older	349

Public Health Outcome Data: None reported.

**Conclusions:** This site has contributed to human exposure and to the environmental burden of trichloroethylene in the past through contamination of groundwater used for household water and discharge of contaminated groundwater into a bayou feeding the Au Sable River. The groundwater, however, has been under remediation since 1992.

#### 4.1.1.9 Metamora Landfill

This 160-acre site, located near the village of Metamora, Lapeer County, MI, contains a 25-acre landfill and 2 drum disposal areas, which may have contained many thousands of drums, believed to contain primarily paint and solvents. Testing of the drum wastes revealed that they contained VOCs, SVOCs, PAHs, and metals, at concentrations as high as 15%, and PCBs at as much as 1,200,000 ppb. As of 1990, excavation and off-site disposal of the drums and associated contaminated soil was underway. The information regarding this site was taken from the 1992 ATSDR public health assessment, HazDat, and the 2003 EPA NPL fact sheet for this site.

**Category of Public Health Hazard:** This site was categorized in the 1992 health assessment (and in an earlier health assessment, not provided for inclusion) as an *Indeterminate Public Health Hazard* (category 3) because although no current exposures at levels of concern had been documented, there was the potential for future exposure through groundwater use as household water. A subsequent ATSDR site review and update (not provided for inclusion in this document) concluded that the site poses *No Apparent Public Health Hazard* (category 4).

**Contaminants of Concern in Completed Exposure Pathways:** None. In 1992, ATSDR was concerned about the potential for exposure to VOCs and metals (particularly arsenic) at concentrations that may result in adverse health effects if, in the future, the shallow groundwater plume extends as far private wells that tap the shallow aquifer. Although the IJC critical pollutant PCBs was found at high concentrations in drums at the site, no further mention was made of this pollutant in the health assessment, implying that it had not contaminated the environment significantly. Site remediation has included the incineration of approximately 35,000 drums and 10,000 tons of soil offsite, and inclusion of minimally contaminated soil under a landfill cap. Groundwater studies in 1997, 1999, and 2000 indicate the VOC groundwater plume is stabilized. Therefore, monitored natural attenuation has been adopted as the remedy for groundwater.

**Demographics:** Demographic profile, from the 2000 U.S. Census, for vulnerable populations living within 1 mile of this site:

Children 6 years and younger	98
Females aged 15-44	205
Adults 65 and older	93

**Public Health Outcome Data:** Local health authorities have said that they have received no complaints of adverse health effects that can be plausibly associated with the site. Because ATSDR's analysis of the potential for human exposure to site-related chemicals did not indicate that adverse health effects were likely, no further investigation of health outcome data was performed.

**Conclusions:** This site may have contributed to the environmental burden of VOCs, but it has been remediated. As reported in the EPA fact sheet, extensive remediation of the site, including on-site incineration of wastes and disposal of the resulting ash in the landfill, capping, vegetating, installation of a runoff treatment system, and installation of a groundwater extraction system, has largely eliminated releases of contaminants from the site.

#### 4.1.1.10 Spiegelberg and Rasmussen Dump Sites

The 115-acre Spiegelberg Site and the 33-acre Rasmussen Dump are two separate sites in Livingston County, MI, that share a common property line. They are considered together in ATSDR health assessments. Both sites were used for the disposal of municipal and industrial wastes. Paint wastes were disposed on the Spiegelberg Site, and drummed industrial wastes were disposed on the Rasmussen site. Many of the drums were removed, along with contaminated soil, in 1984. A few residences are located on the sites. Information regarding these sites is taken from the 1989 public health assessment and the 1992 public health assessment addendum prepared by ATSDR, and from the 2003 EPA NPL fact sheets for the sites.

**Category of Public Health Hazard:** These sites were categorized as an *Indeterminate Public Health Hazard* (category 3) in 1989 because of the potential threat to human health from exposure to contaminants at levels that may result in adverse health effects and incomplete monitoring data. In the 1992 health assessment, the sites were categorized as *Public Health Hazards* (category 4) because of the threat of exposure to contaminated groundwater that was likely to occur unless the remedial actions indicated for this site were carried out.

**Contaminants of Concern in Completed Exposure Pathways:** None. In 1989, chemicals of concern in potential exposure pathways included the IJC critical pollutants PCBs (groundwater and soil)

and lead (groundwater and soil). Other contaminants of concern in potential exposure pathways were VOCs, including vinyl chloride and methylene chloride, in groundwater. In 1992, additional testing results were found to support the findings from the 1989 health assessment. Subsequent remedial actions included the removal of additional drums and contaminated soil, with disposal offsite, installation of a landfill cap, and installation of groundwater treatment. These actions have been found to be protective of public health and the environment.

**Demographics:** Demographic profile, from the 2000 U.S. Census, for vulnerable populations living within 1 mile of these sites:

	Spiegelberg	Rasmussen
Children 6 years and younger	119	59
Females aged 15-44	223	121
Adults 65 years and older	73	54

**Public Health Outcome Data:** Health outcome data were not evaluated because of a lack of community health concerns and of evidence that humans had not been significantly exposed to site-related contaminants.

**Conclusions:** This site may have contributed to the environmental burden of the IJC critical pollutant PCBs and lead, as well as other contaminants including VOCs, in the past. Remediation of these sites, including removal of much of the contamination, groundwater treatment, and ongoing monitoring make it unlikely that there will be further releases of contaminants or exposure of human populations.

#### 4.1.1.11 Shiawassee River

The Shiawassee River site, Livingston County, MI was contaminated by the Cast Forge Company, which discharged wastewater contaminated by hydraulic fluids containing PCBs into the South Branch of the Shiawassee River from 1969 to 1973. From 1973 to 1977, waster was discharged into a 400,000 gallon lagoon on-site. Discharges and overflows from this lagoon contaminated nearby wetlands and the Shiawassee River. Starting in 1982, the company removed the lagoon, cleaned up the PCB-contaminated soil and sediment from its property, and provided funds for restoration of the river. Dredging of the South Branch began in 1982, but only the first mile downstream from the plant was treated, removing approximately 2,600 pounds of PCBs. Both the company property and the river were still contaminated as of the ATSDR 1989 health assessment, from which information on this site is taken. Additional more recent information is taken from HazDat and the 2003 EPA NPL fact sheet for this site.

**Category of Public Health Hazard:** In the 1989 health assessment, this site was categorized as an *Indeterminate Public Health Hazard* (category 3) because of the risk to human health that could result from potential exposure to PCBs at levels that may result in adverse health effects. A subsequent ATSDR site review and update (not provided for inclusion in this document) concluded that the site is a *Public Health Hazard* (category 2).

**Contaminants of Concern in Completed Exposure Pathways:** Not identified. The concern was for potential exposure pathways including direct contact with PCB-contaminated river sediments or by eating PCB-contaminated fish or wildlife. PCB levels in fish tissue downstream from Cast Forge were very high; advisories against consumptions of fish from the contaminated zone were issued in 1979. The NPL fact sheet reports that remediation of the flood plan and contaminated areas near Cast Forge to

10 ppm PCBs, and of the river to 5 ppm PCBs for the first mile only, is occurring, and is expected to be completed in fall 2003.

**Demographics:** Demographic profile, from the 2000 U.S. Census, for vulnerable populations living within 1 mile of this site:

Children 6 years and younger	397
Females aged 15-44	885
Adults 65 and older	615

#### Public Health Outcome Data: None reported.

**Conclusions:** This Cast Forge Company's releases of PCBs to the Shiawassee River greatly contributed to the environmental burden of the IJC critical pollutant PCBs. Remediation of the company property, floodplain, and first mile of the river will mitigate, but not eliminate, the contamination.

#### 4.1.1.12 Dow Chemical Co., Michigan Division, Midland Location

The Dow Chemical Company plant in the city of Midland, Midland County, MI was the subject of an ATSDR health consultation that was triggered by community concerns regarding high levels of PCDDs in soil in Midland and in fish in the nearby Tittabawassee River downstream of Midland. The Dow plant encompasses approximately 1,900 acres on the southern perimeter of the city. The Tittabawassee River forms the southern boundary of the plant site and flows southeast to join with the Saginaw River in the vicinity of the city of Saginaw. A variety of chemicals have been produced at this Dow plant, including Agent Orange [which contains 2,4,5-trichlorophenoxyacetic acid (2,4,5-T)], and 2,4,5-trichlorophenol. PCDDs and PCDFs are known to be impurities in some chlorinated phenolic chemicals, such as 2,4,5-trichlorophenol and 2,4,5-T. Chlorophenol production started in 1915. Wastes generated from this process were initially transferred to 600 acres of on-site waste ponds. During high flow periods in the early 1900s, wastes from these ponds were intentionally released to the Tittabawassee River. Some site waste has been and is taken by truck from the Dow plant to local landfills. Since that time, Dow has operated its own wastewater treatment plant on-site, but a significant flood in 1986 overwhelmed the wastewater treatment plant and flooded areas of the plant where soils were contaminated with PCDDs. The runoff and untreated or partially treated chemical wastes entered the Tittabawassee River. Two incinerators are used for treatment of liquid and solid hazardous and non-hazardous wastes generated from manufacturing at the plant. Incineration of chlorine-containing wastes also produces PCDDs and PCDFs. Information regarding this site is taken from the 2002 health consultation prepared by ATSDR. This health consultation focused on contamination of Midland soil. A separate health consultation was prepared regarding contamination in the Tittabawassee River floodplain near the city of Saginaw, in Saginaw County (see section 4.1.1.13).

**Category of Public Health Hazard:** This site was categorized as an *Indeterminate Public Health Hazard* (category 3) because the data necessary to determine if dioxin contaminated soil in the Midland area poses a public health risk are not available.

**Contaminants of Concern in Completed Exposure Pathways:** Not identified. The IJC critical pollutants PCDDs and PCDFs were found at very high concentrations (expressed as total toxic equivalent (TEQ) concentrations in soil at the Dow plant. The residential areas to the northeast are expected to have the highest impact from historical incinerator emissions, but no data are available concerning dioxin concentrations in these areas of Midland. Most of the TEQ concentration data for the community fall

within the range (>50 but <1000 ppt TEQs) that triggers additional ATSDR evaluation, including consideration of background and bioavailability data in order to evaluate the incremental contribution of soil exposure; this information was not available.

**Demographics:** Residential neighborhoods are located in close proximity to the northeast perimeter of the Dow plant and within a quarter of a mile from a soil sampling site where total TEQs were above the ATSDR action level of 1,000 ppt.

#### **Public Health Outcome Data:**

- An analysis of cancer incidence data for zip codes 48640 (southwest area of Midland including the Dow plant site) and 48642 (area northeast of the Dow plant) as compared with Midland County, Bay County, and the state of Michigan showed no elevated incidences of specific cancer types in these two zip code areas. There was a higher-than-expected incidence of all cancers combined in 48640 (but not 48642) as compared with Midland County, Bay County, and the state of Michigan for individual years 1994 through 1998 and all years combined, but interpretation of this data is difficult.
- A Dow Cohort mortality study of workers in the Midland plant compared 2,187 male employees who worked at any time between 1940 and 1983 in areas of the plant where there was potential exposure to dioxin, with exposure classified on the basis of job history. Causes of death were compared to those of the U.S. population and an internal "unexposed" group of employees. Rates for all causes of death were lower in the exposed cohort than in the U.S. population, but were slightly higher for some cancers than in the unexposed employees. The relevance of this study to the non-Dow-employee residents of the community was considered questionable by ATSDR.
- An analysis of birth defects data for 1992 through 1996 from the Michigan Birth Defects Registry did not show any consistent pattern of excesses in any particular category or for birth defects overall for Midland County (about 1,000 births/year). No excess was seen for types of birth defects, such as anecephaly, spina bifida, and cleft lip, which had been reported as related to dioxin exposure.

**Conclusions:** This site has contributed to the environmental burden of the IJC critical pollutants PCDDs and PCDFs. Whether residents of the community near the plant experienced a level of exposure sufficient to be considered a public health risk could not be determined due to the lack of soil monitoring data in the critical areas and other data deficiencies.

#### 4.1.1.13 Tittabawassee River

The Dow Chemical Company plant in the city of Midland, Midland County, MI was the subject of an ATSDR health consultation that was triggered by community concerns regarding high levels of PCDDs in soil in the city of Midland and in fish in the nearby Tittabawassee River downstream of Midland. An additional concern arose when sampling of the Tittabawassee floodplain near the confluence of the Tittabawassee and Saginaw Rivers revealed high levels of dioxin contamination. The soil contamination issue was considered in the ATSDR health consultation on the Dow Chemical Co. site, presented in Section 4.1.1.12, which provides a description of the plant location and releases to the environment. The issue of contamination of the floodplain of the Tittabawassee River is considered in a separate 2002 ATSDR health consultation, summarized below. The Tittabawassee floodplain area that is potentially of concern extends from the City of Midland in Midland County to the City of Saginaw in Saginaw County. The sampling sites were within Saginaw County.

**Category of Public Health Hazard:** This site was categorized as an *Indeterminate Public Health Hazard* (category 3) because of the potential threat to human health from exposure to PCDDs and PCDFs and the lack of monitoring data for the residential area.

**Contaminants of Concern in Completed Exposure Pathways:** Elevated dioxin TEQs (as high as 7,261 ppt, includes PCDDs and PCDFs) were found in soil samples from a floodplain area near the confluence of the Tittabawassee and Saginaw Rivers in Saginaw County, analyzed as part of a wetland mitigation project, and in other floodplain areas (golf course, wildlife refuge) upstream from the mitigation site. These levels were considered to be high enough to pose an urgent public health hazard if people were routinely exposed to soil at these locations, but ATSDR concluded that the level of exposure on these properties is not known, and was concerned regarding the lack of sampling on nearby residential properties. The only known source of dioxin contamination was the Dow Chemical Company plant upstream at Midland. ATSDR concluded that the contamination likely resulted from deposition of contaminated river sediments in the Tittabawassee River floodplain. As discussed in Section 4.1.1.12, fish in the Tittabawassee River below the city of Midland have elevated levels of PCDDs and PCBs. Based on the floodplain soil data together with the fish data, ATSDR concluded that dioxin contamination may be widespread throughout the Tittabawassee River watershed below Midland.

**Demographics:** Twelve homes are located adjacent to the river less than half a mile upstream from the mitigation site where very high TEQs were detected. Numerous other residential properties are located within the floodplain upstream of the wetland mitigation site.

#### Public Health Outcome Data: None reported.

**Conclusions:** This site is contaminated with the IJC critical pollutants PCDDs and PCDFs, probably from releases from the Dow Chemical Company plant upstream at Midland, Midland County.

#### 4.1.1.14 Lufkin Rule

The 14-acre Lufkin Rule property is a large abandoned industrial property in a mostly residential area of Saginaw, Saginaw County MI. After being sold, the property was rented out to a large number of tenants. In 1994, a dry cleaning establishment on the property burned, and the remnants were later demolished. Since that time, the entire property has been vacant. Drums of dry-cleaning solvers, transformers, capacitors, and other electrical equipment containing PCBs were found on the property. Some of the equipment had been scavenged, and the PCB-containing oil spilled on the ground. The PCB-containing oil and soil, drummed solvents, and other waste materials were removed in 1995 for disposal at an approved facility. Information regarding this site is taken from the ATSDR 1997 health consultation.

**Category of Public Health Hazard:** This site was categorized as a *Public Health Hazard* (category 3) because of the physical hazards in the abandoned and decrepit buildings on the property, and contaminants in soil that would soil pose health hazards to anyone working on the property for long periods. The site is not secured from trespassers, and there is evidence of extensive trespassing.

**Contaminants of Concern in Completed Exposure Pathways:** None identified. There are hot spots of soil contamination with the IJC critical pollutant PCBs and also of bis(2-3ethylhexyl)phthalate that could pose health hazards through inadvertent ingestion to anyone working in those areas for long periods or visiting those areas daily over a long period of time, but this exposure scenario was considered unlikely. Levels of the IJC critical pollutants B(a)P and lead in soil and storm sewer sediment exceeded health based screening values, but were within ranges typically found in urban areas. Groundwater was contaminated with trichloroethylene, but is not used as a drinking water source. Levels of trichloroethylene and other VOCs in storm sewer water were above drinking water standards, and indicate release from the site through runoff.

**Demographics:** Not reported, but the site is located in a residential area.

Public Health Outcome Data: None reported.

**Conclusions:** This site may have contributed to the environmental burden of the IJC critical pollutant PCBs, and also of VOCs, but the extent of on-site contamination is limited.

#### 4.1.1.15 Laingsburg

The Laingsburg property is a former gasoline and automotive service station located in the city of Laingsburg, Shiawassee County, MI, which stopped operations in 1984, and since then, has been used for automotive repair and body shop work. In 2000, a health consultation was performed by ATSDR as part of a Brownfields project; that document is the source of information regarding this site. Records indicate that there may have been three underground fuel storage tanks on the property, and there is no clear indication that the tans were removed from the property.

**Category of Public Health Hazard:** This site was categorized as an *Indeterminate Public Health Hazard* (category 3) because of the potential threat to human health from exposure to contaminants and the lack of adequate monitoring data.

**Contaminants of Concern in Completed Exposure Pathways:** Not reported. Access to the site was denied, so no on-site monitoring data are available. Subsurface soil sampled around the perimeter of the site contained trimethylbenzene and xylenes above screening values for industrial or commercial use. Shallow groundwater at the site perimeter was similarly contaminated had a floating oily layer liquid (one monitoring well) containing trimethylbenzenes and other VOCs. Concentrations exceeded drinking water standards or screening levels. The contamination was consistent with gasoline leaking from the underground storage tanks.

**Demographics:** Not reported, but there are eight private wells within 0.2 miles of the site, and Laingsburg has no municipal water system; residents use individual private wells.

Public Health Outcome Data: None reported.

**Conclusions:** This site may be releasing gasoline from underground storage tanks, but access to the site was denied and the available monitoring data are inadequate to assess the potential threat to public health.

#### 4.1.2 TRI Data for the Saginaw River and Bay AOC

The TRI on-site chemical releases for the 21 counties (combined) that are relevant to this AOC are summarized in Table 4-3. Total on-site releases for the 21 counties in 2001 were 7,831,200 pounds, the majority of which were released to air, followed by releases to soil. Considerably less was released to surface water.

The IJC critical pollutants accounted for 92,142 pounds or 1.2% of the total on-site releases. The IJC critical pollutants released were PCDDs and PCDFs (primarily to land), lead and lead compounds (primarily to land); and mercury and mercury compounds (primarily to air and land). The facilities that

released these pollutants are listed in Table 4-4. PCDDs (and PCDFs) were the focus of ATSDR health consultations for soil contamination by the Dow Chemical Co. in the city of Midland, Midland County, MI (Section 4.1.1.10) and for contamination of the Tittabawassee River Flood Plain south of Midland (Section 4.1.1.11). The major TRI releases of these chemicals in the counties relevant to the Saginaw River and Bay AOC were in Midland County, by the Dow Chemical Company (1,618 pounds total onsite releases, primarily to land). Much smaller amounts were reported released by other facilities in Bay County and Saginaw County.

The major releases ( $\geq$ 500,000 pounds) of non-IJC chemicals were of hydrochloric acid aerosols to air and barium compounds (primarily to land). Other non-IJC chemicals released in substantial on-site quantities (300,000-499,999 pounds) were toluene (primarily to air); and barium compounds, manganese compounds, and zinc compounds (primarily to land); and ammonia (to air, water and land).

Looking at total on-site releases of all chemicals combined, the counties with the highest reported releases, 500,000-1,000,000 pounds, were Midland and Saginaw Counties. Counties with total on-site releases of 250,000-499,999 pounds were Bay, Genesee, and Huron counties. Counties in the range of 100,000-249,000 pounds total on-site releases were Montcalm, Osceola, and Sanilac. Counties in the range of 10,000-99,000 pounds total on-site releases were Gratiot, Isabella, Lapeer, Livingston, Mecosta, Ogemaw, Shiawassee, and Tuscola. Counties in the range of 0-9,999 pounds total on-site releases were Arenac, Clare, Gladwin, Iosco, and Roscommon.

#### 4.1.3 County Demographics and Health Status Data for the Saginaw River and Bay AOC

The demographic profiles, from the 2000 U.S. Census, for vulnerable populations living in the 21 counties of this AOC are shown in Table 4-5.

According to the 2000 HRSA community health status reports, health status indicators that compared unfavorably with those of the U.S. and also with the median of the peer counties for the 21 counties of this AOC were as shown in Table 4-5.

#### 4.1.4 Summary and Conclusions for the Saginaw River and Bay AOC

#### 4.1.4.1 Hazardous Waste Sites

Sixteen sites in the counties relevant to the Saginaw River and Bay AOC have been categorized by ATSDR in health hazard categories 1-3 at some time in their assessment history. Several of these sites have completed exposure pathways to the IJC critical pollutants PCDDs, PCDFs, PCBs, and/or DDT and metabolites and/or released these pollutants into the rivers that ultimately feed the Saginaw River. Sites that have not yet been completely remediated (as of this writing) and may be continuing to serve as a source of exposure are:

- Bay City Middlegrounds (Section 4.1.1.1)
- Velsicol Chemical Corp. (Section 4.1.1.7)
- Shiawassee River (Section 4.1.1.11)
- Dow Chemical Co., Midland Location (Section 4.1.1.12)
- Tittabawassee River (Section 4.1.1.13)

A pathway of major concern for these chemicals is bioaccumulation through the food chain into fish that are ingested by humans. Incidental ingestion, direct dermal contact, and inhalation of soil and dust from PCDD- and PCDF-contaminated soil also were of concern.

Public health outcome data, available for four of the sites, generally did not indicate unusual rates of health conditions, or consisted of occupational data, which were considered of questionable relevance to the general population. An analysis of cancer incidence data for the Dow Chemical Co. site found no elevated incidences of specific cancer types in the two zip code areas studied, as compared with county and state. A higher-than-expected incidence of all cancers combined was seen in the zip code area upwind and including the site, but not the zip code area downwind of the site, which was considered more highly contaminated with PCDDs and PCDFs from the Dow Chemical Company's on-site incineration of chemical wastes. ATSDR considered that interpretation of these data was problematic.

#### **Issues for Follow-Up**

The sites listed as still possibly contributing to environmental contamination and human exposure may need follow-up to determine whether the potential hazards have been mitigated. Additional monitoring data and other data also were needed to more fully assess the hazard.

#### 4.1.4.2 TRI Data

On-site TRI releases in the 21 counties (combined) of the Saginaw River and Bay AOC totaled 7,831,200 pounds, the majority of which were released to air, followed by releases to soil. Considerably less was released to surface water.

The highest release counties, Midland and Saginaw Counties, accounted for 10.5 and 12.3%, respectively, of the total on-site releases. The lowest release counties, Arenac, Gladwin, and Roscommon, had zero reported releases.

The IJC critical pollutants accounted for 92,142 pounds or 1.2% of the total on-site releases. The IJC critical pollutants released were PCDDs and PCDFs (primarily to land); lead and lead compounds (primarily to land); and mercury and mercury compounds (primarily to air and land).

The major releases ( $\geq$ 500,000 pounds) of non-IJC chemicals were of hydrochloric acid aerosols, xylenes, certain glycol ethers, n-butyl alcohol, and toluene (primarily to air); and nickel compounds, selenium, and arsenic compounds (primarily to land).

#### 4.1.4.3 County Demographics and Health Status Indicators

There is considerably variation across counties of the Saginaw River and Bay AOC in vulnerable population demographics and in the pattern of health indicators that compared unfavorably with U.S. indicators and with the peer counties' indicators. The two counties with the highest vulnerable population demographics, Genesee and Saginaw, also had the greatest number of unfavorable health status indicators.

			Number of Records						
		IJC							
		Tracking			Human		- ··		
CAS No.	Chemical Name	Number	Air	Biota	Material	Other Media	Soil	Water	Total
011097-69-1	AROCLOR 1254	1					4		4
011096-82-5	AROCLOR 1260	1					4		4
001336-36-3	POLYCHLORINATED	1		15		4	31	9	59
004740 04 0	BIPHENYLS	0				2	2		0
001746-01-6		2				3	3		6
000050-32-8		4					12	з	15
000072-54-8		5		1			6	3	10
000072-55-9	DDE PP'-	5		2			3	1	6
000050-29-3	DDT. P.P'-	5		6		1	7	2	16
000309-00-2	ALDRIN	6		•		•	4	4	8
000060-57-1	DIELDRIN	6		2			5	3	10
007439-92-1	LEAD	8		2			24	12	38
007439-97-6	MERCURY	9		2			10	3	15
000118-74-1	HEXACHLOROBENZENE	11		1			2	3	6
		Total IJC	0	31	0	8	115	43	197
000071-55-6	1,1,1-TRICHLOROETHANE		1			2	11	8	22
000079-34-5	1,1,2,2-TETRACHLOROETHANE						1	1	2
000079-00-5	1,1,2-TRICHLOROETHANE					1	4	7	12
000075-34-3	1,1-DICHLOROETHANE		1			1	6	17	25
000075-35-4	1,1-DICHLOROETHENE					1	1	7	9
000634-66-2	1,2,3,4-						2	2	4
	TETRACHLOROBENZENE								
000087-61-6							2		2
000120-82-1						4	5	4	9
000095-65-6			1			1	1	0	3 15
000107-00-2			I			I	4	9	15
000156-60-5						1	5	11	17
000100 00 0	TRANS-						0		
000540-59-0	1.2-DICHLOROETHYLENE						8	12	20
000078-87-5	1.2-DICHLOROPROPANE						•	1	1
000108-67-8	1,3,5-TRIMETHYLBENZENE		1			1	1	1	4
000106-99-0	1,3-BUTADIENE		1						1
000541-73-1	1,3-DICHLOROBENZENE		1			1	4		6
010061-02-6	1,3-DICHLOROPROPENE,		1			1	1	1	4
	TRANS-								
000106-46-7	1,4-DICHLOROBENZENE		1			1	7	8	17
000105-67-9	2,4-DIMETHYLPHENOL							2	2
000078-93-3							6	4	10
000591-78-6						4	10	1	10
000091-57-6						I	12	3 1	10
000100-02-7							1	1	1
000208-96-8							3		3
000067-64-1	ACETONE						3	2	5
007429-90-5	ALUMINUM						1	1	2
000120-12-7	ANTHRACENE						6		6
007440-36-0	ANTIMONY						9	2	11
007440-38-2	ARSENIC						14	12	26
001332-21-4	ASBESTOS					1			1
007440-39-3	BARIUM						13	13	26
000071-43-2	BENZENE		2			1	9	19	31
000092-87-5	BENZIDINE						2		2
000056-55-3	BENZO(A)ANTHRACENE						12	2	14
000203-33-8	BENZO(A)FLUORANTHENE						2	_	2
000205-99-2	BENZO(B)FLUORANTHENE						11	2	13
000191-24-2							12	1	13
000207-08-9							10	3	13
0-69-60000								∠	2

## Table 4-2. Waste Site Contaminants that Exceeded Health-Based Screening Values Saginaw River and Bay AOC

			Number of Records						
CAS No.	Chemical Name	IJC Tracking Number	Air	Biota	Human Material	Other Media	Soil	Water	Total
007440-41-7	BERYLLIUM						13	4	17
000075-27-4	BROMODICHLOROMETHANE					1	1	3	5
000074-83-9	BROMOMETHANE						1	1	2
000085-68-7	BUTYL BENZYL PHTHALATE						7	3	10
007440-43-9	CADMIUM			2			9	4	15
000086-74-8	CARBAZOLE						10	2	12
000075-15-0	CARBON DISULFIDE							1	1
000056-23-5	CARBON TETRACHLORIDE		2					4	6
000057-74-9	CHLORDANE			1				1	2
016887-00-6 HZ0400-01-T	CHLORIDE CHLORINATED DIOXINS,						6	1	1 6
000108-90-7			З			1	12	7	23
000124-48-1			0				12	1	1
000075-00-3	CHI OROFTHANE		1					7	8
000067-66-3	CHLOROFORM		2			1	5	7	15
000074-87-3	CHLOROMETHANE		1			1	1	2	5
007440-47-3	CHROMIUM			2			15	11	28
000218-01-9	CHRYSENE						12	4	16
005103-71-9	CIS-CHLORDANE						3		3
007440-48-4	COBALT						14	3	17
007440-50-8	COPPER			2			15	6	23
000095-48-7	CRESOL, ORTHO-						1	2	3
000106-44-5	CRESOL, PARA-						5	5	10
000098-82-8	CUMENE						1		1
000057-12-5	CYANIDE						5		5
000117-81-7	DI(2-ETHYLHEXYL)PHTHALATE						8	5	13
000053-70-3	DIBENZO(A,H)ANTHRACENE						5	1	6
025321-22-6	DICHLOROBENZENE						2		2
001300-21-6	DICHLOROETHANE							1	1
000084-66-2	DIETHYL PHTHALATE							1	1
000107-99-3	DIMETHYLAMINOETHYL							1	1
	CHLORIDE						_		_
000084-74-2	DI-N-BUTYL PHTHALATE						5	2	7
000117-84-0	DI-N-OCTYL PHTHALATE			•			5	2	1
HZ0400-05-1	DIOXINS N.O.S.			3			1		4
HZ0400-04-1	DIOXINS/FURANS,						2		2
001031-07-8	ENDOSULEAN SULEATE						2		2
000959-98-8	ENDOSULEAN ALPHA						1		1
033213-65-9	ENDOSULFAN, BETA						1		1
000072-20-8	ENDRIN						2		2
007421-93-4	ENDRIN ALDEHYDE						4		4
053494-70-5	ENDRIN KETONE						3		3
000100-41-4	ETHYLBENZENE		3				12	12	27
000206-44-0	FLUORANTHENE						7	1	8
000086-73-7	FLUORENE						4		4
HZ0500-03-T	FURANS, UNSPECIFIED			2			6		8
HZ0900-02-T	HEAVY METALS, UNSPECIFIED						6		6
000076-44-8	HEPTACHLOR						3	2	5
001024-57-3	HEPTACHLOR EPOXIDE			1			5	2	8
000087-82-1	HEXABROMOBENZENE					1	3		4
000087-68-3	HEXACHLOROBUTADIENE						2	2	4
000319-84-6	HEXACHLOROCYCLOHEXANE,						3	2	5
	ALPHA-								
000319-85-7	HEXACHLOROCYCLOHEXANE,						1		1
000040 00 0							~	~	0
000319-86-8	DELTA-						6	3	9
000058-89-9	HEXACHLOROCYCLOHEXANE, GAMMA-						6	2	8

## Table 4-2. Waste Site Contaminants that Exceeded Health-Based Screening Values Saginaw River and Bay AOC

## Table 4-2. Waste Site Contaminants that Exceeded Health-Based Screening Values Saginaw River and Bay AOC

			Number of Records						
		IJC Tracking			Human				
CAS No.	Chemical Name	Number	Air	Biota	Material	Other Media	Soil	Water	Total
HZ1000-01-T	HYDROCARBONS,					4	5	1	10
	UNSPECIFIED								
000193-39-5	INDENO(1,2,3-CD)PYRENE						12	1	13
HZ0900-18-T	INORGANICS, N.O.S.							1	1
007439-89-6	IRON						1	2	3
007439-95-4	MAGNESIUM						1		1
001309-42-8						1			1
001309-48-4						1	11	4.4	1
UU7439-96-5						2	14	1	25 5
000072-43-5	METHOXYCHLOR					5	3	I	3
000072 40 0							1	1	2
000075-09-2	METHYLENE CHLORIDE		4			1	10	12	27
000091-20-3	NAPHTHALENE		•			1	15	8	24
007440-02-0	NICKEL						11	8	19
000103-65-1	N-PROPYL BENZENE					1	1	1	3
029082-74-4	OCTACHLOROSTYRENE			1					1
HZ0600-01-T	OIL/GREASE, UNSPECIFIED						1		1
027858-07-7	PBB'S (OCTOBROMOMIXTURE)			4	1		8	7	20
000059-50-7	P-CHLORO-M-CRESOL						1	3	4
000087-86-5	PENTACHLOROPHENOL						_	3	3
HZ1200-01-T	PESTICIDES N.O.S.					1	2	1	4
000085-01-8	PHENANIHRENE						12	4	16
000108-95-2	PHENOL						2	4	6
067774 22 7				2		1	1	4	11
00///4-32-/				3		I	3	4	11
130498-29-2						1	1		2
100400 20 2	HYDROCARBONS						•		2
000129-00-0	PYRENE						7	1	8
007782-49-2	SELENIUM						3		3
HZ1900-02-T	SEMIVOLATILE ORGANIC					1	1	2	4
	COMPOUNDS N.O.S.								
007440-22-4	SILVER						3		3
000100-42-5	STYRENE						6	3	9
014808-79-8	SULFATE			_		_	_	1	1
HZ0400-03-T	TCDD EQUIVALENTS			2		5	7	4.0	14
000127-18-4	TETRACHLOROETHYLENE		2			2	14	12	30
000109-99-9							4	4	4
007440-26-0			2			1	4	11	о 20
000100-00-3			3			I	6	2	29
005103-74-2							0	2	2
000079-01-6	TRICHLOROFTHYLENE		5			3	18	29	55
000077-86-1	TRIS		Ũ			1	2	20	3
007440-62-2	VANADIUM						14	1	15
000075-01-4	VINYL CHLORIDE					1	4	18	23
HZ1900-01-T	VOLATILE ORGANIC						1	4	5
	COMPOUNDS N.O.S.								
001330-20-7	XYLENES, TOTAL		3			1	11	11	26
007440-66-6	ZINC			2			14	10	26
000132-64-9	DIBENZOFURAN		-				11	<i>c</i> -	11
MEDEXP-00-0			9	14		1	16	26	66
			1	6	<u> </u>	1	12	6	26
		I Otal	49	45	1	52	707	499	1353
		Total	40	76	4	60	000	540	1550
		iulai	49	10		00	022	542	1550

Table 4-3. TRI Releases (in pounds, 2001) for the Saginaw River and Bay AOC

Chemical	IJC Tracking Number	Total Air Emissions	Surface Water Discharges	Under- ground Injection	Releases to Land	Total On- site Releases	Total Off- site Releases	Total On- and Off-site Releases
DIOXIN AND DIOXIN-LIKE								
COMPOUNDS	2	0.013254255	0.005655825	0	1.6102674	1.62917748	0.05532345	1.68450093
(PCDDs and PCDFs)	3							
LEAD	8	243.18	15.2	0	0	258.38	20665.9	20924.28
LEAD COMPOUNDS	8	4521.6	1.1	0	86883.8	91406.5	6029.6	97436.1
MERCURY	9	10.2	0	0	44	54.2	0.8	55
MERCURY COMPOUNDS	9	270.6	1	0	150	421.6	3.7	425.3
	Total IJC	5045.593254	17.30565583	0	87079.4103	92142.30918	26700.05532	118842.3645
1,1,1,2-TETRACHLOROETHAN	١E	588	0	0	0	588	0	588
1,1-DICHLORO-1-FLUOROETH	HANE	101906	0	0	0	101906	250	102156
1,2,4-TRIMETHYLBENZENE		97955	0	0	0	97955	0	97955
1,2-DICHLOROETHANE		472	0	0	0	472	0	472
1,2-DICHLOROPROPANE		7	0	0	0	7	0	7
1,3-BUTADIENE		2751	0	0	0	2751	0	2751
2,4,6-TRICHLOROPHENOL		41	64	0	1	106	0	106
2,4-D		167	0	0	0	167	0	167
2,4-D BUTYL ESTER		1	1	0	0	2	0	2
2,4-DICHLOROPHENOL		181	0	0	1	182	0	182
2-METHOXYETHANOL		4524	0	0	0	4524	0	4524
2-PHENYLPHENOL		1	0	0	0	1	0	1
3-CHLORO-2-METHYL-1-PROI	PENE	68	0	0	0	68	0	68
4,4'-ISOPROPYLIDENE-DIPHE	NOL	754	0	0	0	754	4215	4969
ACETONITRILE		40902	3252	0	7	44161	0	44161
ACROLEIN		1700	0	0	0	1700	0	1700
ACRYLAMIDE		18	7	0	0	25	0	25
ACRYLIC ACID		12018	2	0	0	12020	0	12020
ACRYLONITRILE		7453	0	0	0	7453	0	7453
ALLYL ALCOHOL		630	0	0	0	630	0	630
ALLYL CHLORIDE		2128	0	0	0	2128	0	2128
ALLYLAMINE		3	0	0	0	3	0	3
ALUMINUM (FUME OR DUST)		1060	5	0	0	1065	11497	12562
AMMONIA		168025	116434	0	60000	344459	0	344459
ANTIMONY		250	0	0	0	250	750	1000
ANTIMONY COMPOUNDS		568	0	0	11011	11579	0	11579
BARIUM COMPOUNDS		2494	249	0	947010	949753	41012	990765
BENZENE		37544	1	0	101	37646	0	37646
BENZO(G.H.I)PERYLENE		0.3898	0	0	0	0.3898	0	0.3898
BIPHENYL		146	0	0	0	146	0	146
BROMINE		388	0	0	0	388	0	388
BUTYL ACRYLATE		1	0	0	0	1	0	1
CARBON DISULFIDE		515	0	0	2	517	0 0	517
CERTAIN GLYCOL ETHERS		265755	3100	0	4000	272855	11670	284525
CHLORINE		33689	0.00	0	0	33689	0	33689
		34	0	0	0	34	0	34
CHLOROBENZENE		56	0	0	ů 0	56	0	56
CHLORODIELLIORO-METHAN	F	75572	0	0	ů 0	75572	0	75572
CHLOROETHANE	-	36655	0	0	ů 0	36655	0	36655
CHLOROFORM		273	0	0	ů 0	273	0	273
		10073	0	ů 0	Ő	10073	0	10073
	HER	10070	0	0	0	1091	0	1091
		1001	27	0	1	32	0	32
		+ 2777		0	6205	8083	110366	110340
CHROMIUM COMPOUNDS (F)	CEPT	2111	1	0	0205	0000	110500	110040
CHROMITE ORE MINED IN TH	IE							
TRANSVAAL REGION)		4973	130	0	53452	58555	33085	91640
COBALT		750	5	0	0	755	11265	12020
COBALT COMPOUNDS		754	250	0	41700	42704	0	42704
COPPER		2249	270	0	5	2524	37338	39862
COPPER COMPOUNDS		2675	2400	0	68400	73475	351	73826
CUMENE		116	0	0	0	116	0	116
CUMENE HYDROPEROXIDE		800	0	0	0	800	560	1360
CYCLOHEXANE		2584	0	0	0	2584	0	2584
DECABROMODIPHENYL OXID	DE	3741	0	0	23600	27341	0	27341
DICHLOROMETHANE		10667	1	0	0	10668	0	10668

Chemical	IJC Tracking Number	Total Air Emissions	Surface Water Discharges	Under- ground Injection	Releases to Land	Total On- site Releases	Total Off- site Releases	Total On- and Off-site Releases
DIISOCYANATES		1529	0	0	0	1529	12351	13880
DIMETHYLAMINE		2185	131	0	0	2316	0	2316
DINITROBUTYL PHENOL		36	0	0	0	36	0	36
EPICHLOROHYDRIN		6	0	0	0	6	0	6
ETHYL ACRYLATE		45	0	0	0	45	0	45
ETHYLBENZENE		54591	1	0	5	54597	0	54597
ETHYLENE		40179	0	0	0	40179	0	40179
ETHYLENE GLYCOL		4455	283	0	0	4738	51500	56238
ETHYLENE OXIDE		904	0	0	0	904	0	904
FORMALDEHYDE		28051	21	0	7000	35072	242	35314
FORMIC ACID		252	118	0	0	370	0	370
HYDROCHLORIC ACID (1995 A	ND AFTER	_0_		Ũ	Ū.	0.0	Ŭ	0.0
'ACID AEROSOLS' ONLY)		2755268	0	0	0	2755268	0	2755268
HYDROGEN FLUORIDE		200090	0	0	0	200090	0	200090
MANGANESE		1795	1	0	0	1796	108939	110735
MANGANESE COMPOUNDS		9347	9190	0	341800	360337	36335	396672
METHACRYLONITRILE		945	0	0	0	945	0	945
METHANOL		100222	0	0	2	100224	0	100224
METHYL ACRYLATE		759	0	0	0	759	0	759
METHYL ETHYL KETONE		102164	0	0	0	102164	0	102164
		24110	0	0	Ő	24110	0	24110
		1018	0	õ	Ő	1018	õ	1018
METHYL TERT-BUTYL ETHER		854	0	Ő	ů 0	854	Ő	854
		16	0	0	0	16	0	16
		20055	0	0	101	20246	0	20246
		20033	1	0	191	86572	0	86572
		2027	1	0	0	200272	0	20272
N-HEXANE		2902	0	0	1005	2902	169144	172200
		11200	200	0	22000	42224	100144	60242
		11309	10000	0	32000	43321	10922	02243
		2000	10000	0	0	10500	21	10021
		0766	5	0	5	0766	0	0766
N-METHYL-2-PYRROLIDONE		9700	0	0	1002	9700	2025	9/00
PHENOL		105740	0	0	1903	107643	3935	111576
PHOSGENE		23	0	0	0	23	0	23
PICLORAM	_	1	257	0	0	258	0	258
POLYCHLORINATED ALKANES	5	0	0	0	0	0	63100	63100
POLYCYCLIC AROMATIC COM	IPOUNDS	2.81	0	0	53	55.81	0.8	56.61
PROPYLENE		240	0	0	0	240	0	240
PROPYLENE OXIDE		1234	0	0	0	1234	0	1234
PYRIDINE		201	0	0	0	201	0	201
QUINOLINE		3	0	0	0	3	0	3
SODIUM NITRITE		0	0	0	0	0	720	720
STYRENE		188607	23	0	38959	227589	498.2	228087.2
SULFURIC ACID (1994 AND AF	TER 'ACID	444005	0	~	0	444005	~	444005
AEROSOLS' ONLY)		144005	0	0	0	144005	0	144005
TETRACHLORO-ETHYLENE		22458	0	0	1	22459	0	22459
IOLUENE		373364	2	0	32	3/3398	0	373398
TOLUENE-2,4-DIISOCYANATE		578	0	0	0	578	0	578
TRICHLOROETHYLENE		11319	0	0	0	11319	0	11319
TRIETHYLAMINE		18928	0	0	4	18932	0	18932
VANADIUM COMPOUNDS		12011	0	0	106700	118711	0	118711
VINYL ACETATE		92	0	0	0	92	0	92
VINYL CHLORIDE		969	0	0	1	970	0	970
VINYLIDENE CHLORIDE		21474	3	0	0	21477	0	21477
XYLENE (MIXED ISOMERS)		228329	33	0	0	228362	0	228362
ZINC (FUME OR DUST)		5400	0	0	0	5400	2900	8300
ZINC COMPOUNDS		15569	170	0	288880	304619	111490	416109
	Total							
	Non-IJC	5557404.2	146716	0	2034937	7739057.2	841457	8580514.2
	Total	5562449.793	146733.3057	0	2122016.41	7831199.509	868157.0553	8699356.564

Table 4-3. TRI Releases (in pounds, 2001) for the Saginaw River and Bay AOC

IJC Critical Pollutant	Number of Facilities	Facility Name	TRIF ID	City	
Dioxin and dioxin-like compounds	4	•			
(PCDDs and PCDFs)	-				
Bay County, MI	1	DE KARN - JC WEADOCK GENERATING	48732DKRNJNWEAD	ESSEXVILLE	
Midland County, MI	1	DOW CHEMICAL CO. MIDLAND OPS.	48667THDWCMICHI	MIDI AND	
Saginaw County, MI	2	ALCHEM ALUMINUM INC.	486011 CHMI 2600N	SAGINAW	
	_	GMC SAGINAW METAL CASTING OPS.	48605SGNWG1629N	SAGINAW	
Lead and lead compounds	27				
Bay County, MI	1	DE KARN - JC WEADOCK GENERATING PLANT	48732DKRNJNWEAD	ESSEXVILLE	
Genesee County, MI	6	DELPHI ENERGY & CHASSIS SYS. FLINT WEST	48555CFLNT300NO	FLINT	
		GMC GRAND BLANC METAL FAB	48439CDLLC10800	GRAND BLANC	
		GMC MFD FLINT METAL CENTER	48553GMCTRG2238	FLINT	
		GMC POWERTRAIN FLINT ENGINE SOUTH	48552GMPWR2100B	FLINT	
		GMC POWERTRAIN FLINT NORTH	48550BCFLN902EH	FLINT	
		GMVM - FLINT ASSEMBLY PLANT	48551GMCTRG3100	FLINT	
Gratiot County, MI	1	CONTECH DIV. OF SPX CORP.	48801CNTCH205NG	ALMA	
Huron County, MI	4	DETROIT EDISON CO. HARBOR BEACH POWER PLANT	48441DTRTD755NH	HARBOR BEACH	
		MICHIGAN SUGAR CO. SEBEWAING PLANT	48759MCHGN763BE	SEBEWAING	
		TOWER AUTOMOTIVE TECH. PRODS.	48759TWRTM249NC	SEBEWAING	
		TOWER AUTOMOTIVE TOOL L.L.C.	48731TWRTM81DRE	ELKTON	
losco County. MI	2	ITT INDS. FHS	48750TTNDS4700N	OSCODA	
·····,		NEW NGC INC.	48748NTNLG2375N	NATIONAL CITY	
Isabella County, MI	1	EP HILLSDALE TOOL DIVISION-MT.	48858PHLLS1799G	MOUNT PLEASANT	
		PLEASANT			
Livingston County MI	1	PROGRESSIVE METAL FORMING INC.	48139PRGRS10850	HAMBURG	
Montcalm County, MI	1	FEDERAL MOGUL GREENVILLE	48838FDRLM510EG	GREENVILLE	
Saginaw County, MI	3	GMC SAGINAW METAL CASTING OPS.	48605SGNWG1629N	SAGINAW	
		GMPT SAGINAW MALLEABLE IRON	48605GMCSG77WCE	SAGINAW	
		MICHIGAN SUGAR CO.	48724MCHGN341SU	CARROLLTON	
Sanilac County, MI	2	MICHIGAN SUGAR CROSWELL FACTORY	48422MCHGN159SO	CROSWELL	
		TRELLEBORG YSH INC. SANDUSKY PLANT	48471YLRBB180ND	SANDUSKY	
Shiawassee County, MI	1	MOTOR PRODS OWOSSO CORP.	48867MTRPR201SD	OWOSSO	
Tuscola County, MI	4	GENERAL CABLE INDS.	48726GNRLC6285G	CASS CITY	
		GREDE FOUNDRIES INC. VASSAR FNDY.	48768GRDVS700EH	VASSAR	
		MICHIGAN SUGAR CO CAROFACTORY	48723MCHGN725AL	CARO	
		WALBRO ENGINE MANAGEMENT	48726WLBRN6242G	CASS CITY	
Mercury and mercury compounds	5				
Bay County, MI	1	DE KARN - JC WEADOCK GENERATING PLANT	48732DKRNJNWEAD	ESSEXVILLE	
Genesee County, MI	2	DELPHI ENERGY & CHASSIS SYS. FLINT EAST	48556CSPRK1300N	FLINT	
		GMC GRAND BLANC METAL FAB	48439CDLLC10800	GRAND BLANC	
Huron County, MI	1	DETROIT EDISON CO. HARBOR BEACH POWER PLANT	48441DTRTD755NH	HARBOR BEACH	
Saginaw County, MI	1	GMC SAGINAW METAL CASTING OPS.	48605SGNWG1629N	SAGINAW	

Table 4-4. TRI Facilities Releasing IJC Critical Pollutants On-site for the Saginaw River and Bay AOC

#### Arenac Bay Clare Genesee Gladwin Gratiot **Demographic Profile** Children 6 years and younger 1333 9543 2656 45136 2034 3532 Females aged 15-44 3159 22560 5707 96320 4522 8420 Adults 65 years and older 2860 16170 5398 50607 4768 5723 Infant Mortality (per 1,000 births) Infant mortality Х Х Х Х X X Х Х White infant mortality Х Х Black infant mortality Х Neonatal infant mortality Х Х Х Х Post-neonatal infant mortality Х **Birth Measures (%)** Low Birth Wt Х Х Very Low Birth Wt **Premature Births Teen Mothers** Older Mothers Х Х Х **Unmarried Mothers** No care in 1<sup>st</sup> trimester Х Death measures (per 100,000) Х Х Х Х Breast cancer (female) **Colon Cancer** Х Х Х Х Х Х Х Х Х Coronary heart disease Homicide Lung cancer Х Х Х Stroke х Huron Livingston losco Isabella Lapeer **Demographic Profile** Children 6 years and younger 2871 1893 4614 8606 16313 Females aged 15-44 6400 4537 18980 18585 33324 Adults 65 years and older 7006 5897 5722 8399 13037 Infant Mortality (per 1,000 births) Infant mortality White infant mortality Black infant mortality Neonatal infant mortality Post-neonatal infant mortality Х **Birth Measures (%)** Low Birth Wt Very Low Birth Wt **Premature Births Teen Mothers** Older Mothers Х **Unmarried Mothers** No care in 1<sup>st</sup> trimester Х Х Death measures (per 100,000) X X X X Breast cancer (female) Х Х Colon Cancer Х Х Х Х Х Coronary heart disease Homicide Lung cancer Х Х Х Х Х Х Stroke

# Table 4-5. County Health Status Indicators that Compared Unfavorably with U.S. Indicators and with the Median of the Peer Counties\* Saginaw River and Bay AOC

# Table 4-5. County Health Status Indicators that Compared Unfavorably with U.S. Indicators and with the Median of the Peer Counties\* Saginaw River and Bay AOC

\*Sources: 2000 U.S. Census; 2000 HRSA Community Health Status Indicators Reports

X = Indicators that compared unfavorably with both the U.S. and the median of the peer counties

 $\mathbf{X}$  = Indicators that also were above the upper end of the peer county range