11/4/03 DRAFT

Fire Regime Condition Class (FRCC) Interagency Handbook Reference Conditions

Modeler: Doug Havlina Date: 8/15/03 PNVG Code: SAGE1

Potential Natural Vegetation Group: Sagebrush-Other (e.g., Silver Sagebrush,

Wyoming Big Sagebrush) Without Trees.

Geographic Area: Great Plains.

Description: PNVG commonly occurs in scattered patches in the Great Plains and short grass prairie. This type is primarily composed of the three subspecies of silver sagebrush, with lesser amounts of fringed sagebrush, Wyoming big sagebrush, budsage, and birdfoot sagebrush. Occurrence is tied to topoedaphic features related to soil moisture and texture, and sites are transitional from moist to dryland habitats. PNVG generally abuts plains or prairie grasslands; common associates include buffalograss, wheatgrass, needlegrass, grama, and fescue species. Silver and fringed sagebrush are facultative sprouters.

.

Fire Regime Description: Fire Regime II, primarily short-interval (e.g., <30 yr) stand replacement fires.

Vegetation Type and Structure

Class	Percent of	Description
	Landscape	•
A: post	25	Plains/prairie grass and forb dominated
replacement		community, with sprouting sagebrush shrubs emerging
B: mid-	20	Dense sagebrush stand established from
development		sprouting and seedlings, with plains/prairie
closed		grasses in interspaces
C: mid- open	30	Landscape mosaic of young sagebrush plants intermixed with plains and prairie grasses
D: late- open	10	Open, late-development fire-maintained
		sagebrush community with plains/prairie
		grasses in understory
E: late- closed	15	Dense, late-development sagebrush stand with
		infrequent plains and prairie grasses
Total	100	

Fire Frequency and Severity

Fire Frequency-	Modeled	Pct, All	Description
Severity	Probability	Fires	
Replacement Fire	.03	75	Stand replacement fire in stages A, B, and E
Non-Replacement Fire	.01	25	Mosaic fire in stages C and D
All Fire Frequency*	.04	100	

^{*}Sum of replacement fire and non-replacement fire probabilities.

References

Anderson, Hal E. 1982. Aids to Determining Fuel Models For Estimating Fire Behavior. Gen. Tech. Rep. INT-122. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station. 22 p.

Brown, James K.; Smith, Jane Kapler, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.

Eyre, F. H., ed. 1980. Forest cover types of the United States and Canada. Washington, DC: Society of American Foresters. 148 p.

Hardy, Colin C., Kirsten M. Schmidt, James P. Menakis, R. Neil Samson. 2001. Spatial data for national fire planning and fuel management. Int. J. Wildland Fire. 10(3&4): 353-372.

Kilgore, B.M. 1981. Fire in ecosystem distribution and structure: western forests and scrublands. p. 58-89. In: H.A. Mooney et al. (Technical Coordinators). Proceedings: Conference on Fire Regimes and Ecosystem Properties, Honolulu, 1978. Gen. Tech. Rep. WO-GTR-26.

Kuchler, A.W. 1964. Potential Natural Vegetation of the Conterminous United States. American Geographic Society Special Publication No. 36. 116 p.

Schmidt, Kirsten M, Menakis, James P., Hardy, Colin C., Hann, Wendel J., Bunnell, David L. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 41 p. + CD.

U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (2002, December). Fire Effects Information

System, [Online]. Available: http://www.fs.fed.us/database/feis/ [Accessed: 2/2/03].

Wright, Henry A., Neuenschwander, Leon F., and Britton, Carlton M. 1979. The role and use of fire in Sagebrush-Grass and Pinyon-Juniper Plant Communities. Gen. Tech. Rep. INT-GTR-58. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station. 48 p.

VDDT Results







