HTL-Resummation of the Thermodynamic Potential

A. Peshier

BNL, Bldg. 510A, Upton, NY 11973, USA

Presented by: A. Peshier

Abstract

Due to their asymptotic nature, the accuracy of perturbative approximations cannot be improved, at large coupling, by higher-order terms. Therefore, nonperturbative approaches are required to describe, e. g., the thermodynamics of the quark-gluon plasma, as considered here, in the regimes of physical interest. Diagrams of all orders in the coupling, organized in thermodynamically selfconsistent sets, can be resummed in the so-called Φ-derivable approximation scheme for the thermodynamical potential and the propagators. In the leading-loop order and approximating the propagators by their hard-thermal-loop (HTL) contributions, at the expense of exact selfconsistency, an expression for the QCD thermodynamical potential is derived which is gauge-invariant, UV-finite and consistent with the perturbative result to order $\mathcal{O}(g^2)$. By comparison to the scalar theory, which in the leading-loop approximation can be resummed selfconsistently, it is argued that the thermodynamical potential expressed in terms of the propagators comprises information not contained in a corresponding expression for the entropy derived recently by Blaizot et al. This explains, in particular, why the HTL-resummed approximation cannot reproduce the QCD lattice data below twice the transition temperature T_c. On the other hand, the agreement with the data above $2T_c$ is a remarkable improvement of the conventional perturbative results which are not predictive even orders of magnitude above T_c. This fact encourages the application of the approach for questions not yet tractable by lattice calculations, as for the equation of state at nonzero chemical potential.

Preprint submitted to Elsevier Preprint *20 December 2000*