BACKGROUND DOCUMENT

REPORT ON REVISIONS TO 5TH EDITION AP-42 CHAPTER 15 - ORDNANCE DETONATION

EMISSION FACTORS DEVELOPED BASED ON PHASE III TESTING CONDUCTED AT DUGWAY PROVING GROUND, UTAH

Prepared for:

U.S. Army Environmental Command ATTN: IMAE-ATT Aberdeen Proving Ground, Maryland 21010-5401

June 2007



NOTICE

The information in this document has been funded by the U.S. Army Environmental Command (USAEC) and prepared by MACTEC Federal Programs, Inc., under Contract No. GS-10F-0157K, Order No. W911SR-05-F-0023. It has been reviewed by the U.S. Environmental Protection Agency (EPA) Office of Air Quality Planning and Standards (OAQPS) and has been approved for publication. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

DRAFT

TABLE OF CONTENTS

1.0	INT	INTRODUCTION1		
2.0	CON 2.1 2.2 2.3 2.4 2.5 2.6 2.7 2.8 2.9	Carbon Mo Total Susp Particulate Metals Hydrochlo Volatile On Semivolati Dioxin/Fur	MEASURED AND EMISSION MEASUREMENT METHODS 2 onoxide, Carbon Dioxide, Oxides of Nitrogen, and Sulfur Dioxide 2 ended Particulate 2 Size Distribution 2 ric Acid and Chlorine 2 rganic Compounds 2 ran Compounds 2 on compounds 2	4 4 5 5 5 6
3.0	TES	T DATA A	NALYSIS AND RATING	6
	3.1	EPA Guida	ance Regarding Test Data Quality Ratings	б
	3.2	Analysis of	f Test Data	7
	3.3	Test Data (Quality Ratings14	4
4.0	EMI	SSION FA	CTOR CALCULATIONS17	7
	4.1	Incorporati	on of Analytical Detection-Limits to the Test Data	8
	4.2	Determinat	tion of Background Concentration	8
	4.3	Determinat	tion of Background Concentration	9
	4.4	Determinat	tion of Background-Corrected Concentration	9
	4.5	Determinat	tion of Dilution-Corrected Concentration	9
	4.6		tion of Mass of Compound Released	
	4.7		tion of Emission Factors	
	4.8		tion of Average Emission Factors2	
	4.9		tion of PM-10 and PM-2.5 Emission Factors	
- 0				
5.0			CTOR RATINGS	
	5.1		Factor Ratings Assigned – Based on Ordnance-Specific Test Data	
	5.2	Emission F	Factor Ratings Assigned – Based on All Available Test Data	5
6.0	REF	ERENCES		2
APP	ENDI	X A:	COMPOUNDS ANALYZED AND EMISSION FACTORS DEVELOPED FOR ORDNANCE INCLUDED IN PHASE III TESTING AT DUGWAY PROVING GROUND, UTAH	
APPENDIX B:		X B:	NEW AP-42 SECTIONS FOR ORDNANCE INCLUDED IN PHASE III TESTING AT DUGWAY PROVING GROUND, UTAH	

LIST OF TABLES

<u>No.</u>	<u>Title</u> <u>I</u>	age
1	Ordnance for Which Emission Factors were Developed	1
2	Ordnance Net Explosive Weight	
3	Sampling and Analytical Methods Used	3
4	Selected Sampling or Analytical Method for Compounds Measured Using Two Sampling or Analytical Methods	
5	Compounds for Which Large Relative Percent Differences Were Noted Between Test Runs or Sample Trains.	10
6	Compounds Found in the Method Blank at Levels Between 20 Percent and 50 Percent of Both Test Values	13
7	Compounds Found in the Method Blank at Levels Greater Than 50 Percent of Both Test Values	14
8	Downgraded Data Quality Ratings	. 14
9	Ordnance Categorization for Emission Factor Correlation Assessment	

LIST OF FIGURES

<u>No.</u>	<u>Title</u>	<u>Page</u>
1	Calculation of background-corrected concentration (BCC)	
	DRAFI	

1.0 INTRODUCTION

Due to the lack of credible data concerning emissions from training ordnance when used in their tactical configurations, the U.S. Army Environmental Command (USAEC) established a program to quantify emissions from the detonation of ordnance. This document presents background information concerning the development of air emission factors for four ordnance types used during training exercises at U.S. Army installations. The air emission factors were developed from test data collected by USAEC. Ordnance for which emission factors have been developed and their corresponding AP-42 sections are identified in Table 1. To help readers easily find those emission factors of interest, the ordnance are organized according to their Department of Defense Identification Code (DODIC).

DODIC	Ordnance Description	AP-42 Section
G940	M18 Green Smoke Hand Grenade	15.5.6
G945	M18 Yellow Smoke Hand Grenade	15.5.7
G950	M18 Red Smoke Hand Grenade	15.5.8
G955	M18 Violet Smoke Hand Grenade	15.5.9

TABLE 1 ORDNANCE FOR WHICH EMISSION FACTORS WERE DEVELOPED

The emission factors described in this document are based on data obtained during testing conducted at Dugway Proving Ground, Utah, as presented in the final test report titled *Sampling Results for AEC Phase III Training Ordnance Emission Characterization.*¹ This document was supplemented by additional data provided by the testing contractor.² For each ordnance, one or two test runs were conducted. One item was detonated per run. Source test protocols were developed by USAEC before any testing was conducted and were reviewed by the U.S. Environmental Protection Agency's (EPA's) Emission Measurement Center. The tests were conducted between December 7 and 9, 1998.

The compounds that were measured included carbon monoxide (CO), carbon dioxide (CO₂), oxides of nitrogen (NO_x), sulfur dioxide (SO₂), total suspended particulate (TSP), metals, hydrogen chloride and chlorine (HCl/Cl₂), volatile organic compounds (VOC), semivolatile organic compounds (SVOC), and dioxins/furans (PCDD/PCDF). Within each of the AP-42 sections, only emission factors for criteria pollutants, carbon dioxide, hazardous air pollutants (as defined by 112(b)(1) of the *Clean Air Act* [CAA]), and toxic chemicals (as defined by 313 of the *Emergency Planning and Community Right-to-Know Act* [EPCRA]) are presented.

The emission factors were developed on a "per item" basis and on a "per net explosive weight (NEW)" basis. Users should choose the appropriate emission factor to estimate emissions based upon the data available; either factor is equally valid. The NEW of each ordnance tested is provided in the corresponding AP-42 section and in Table 2.

This document includes five sections in addition to this Introduction. Section 2 of this document identifies the compounds measured during the test program and describes the emission measurement methods used. Section 3 includes a discussion of the emission factor final test report and ratings for the test data contained therein. Section 4 describes the calculations and methodologies used to develop emission factors for each type of compound measured. Section 5 describes the methodology used to rate the emission factors and provides emission factor ratings for each type of compound measured. Section 6 includes a complete list of the references cited in this document.

DODIC	Ordnance Description	NEW (lb/item) ^a
G940	M18 Green Smoke Hand Grenade	7.33 E-01
G945	M18 Yellow Smoke Hand Grenade	7.33 E-01
G950	M18 Red Smoke Hand Grenade	7.33 E-01
G955	M18 Violet Smoke Hand Grenade	7.33 E-01

TABLE 2 ORDNANCE NET EXPLOSIVE WEIGHT

^aNEW values were obtained from References 1 and 3.

There are two appendices included with this document. Appendix A identifies, by ordnance type, all of the compounds for which analyses were performed and the emission factors that were developed. [Note: Compounds present in the method blank at greater than 50 percent of test levels are excluded from Appendix A as described in Section 3.2.4.] Appendix A also identifies the minimum detection levels associated with all compounds that were not detected. Emission factors and minimum detection levels presented in Appendix A were determined from the most accurate method if two sampling or analytical methods were used to measure one compound. Appendix B presents the new AP-42 sections for the four ordnance that were tested.

In addition to this document, there are electronic databases available on the web (<u>http://www.epa.gov/ttn/chief/ap42/index.html</u>) that contain the data used in the development of the emission factors. The general procedures that were followed to develop these emission factors can be found at the same web address under the title *Procedures for Preparing Emission Factor Documents*.⁴

2.0 COMPOUNDS MEASURED AND EMISSION MEASUREMENT METHODS

The USAEC Phase III series testing was conducted in the Smoke Characterization Test Chamber (Smoke Chamber) located at Dugway Proving Ground, Utah. The Smoke Chamber is an aluminum-lined chamber with an interior volume of approximately 820 cubic feet. The chamber is approximately 7 feet wide, 20 feet long, and 6 feet tall for 2/3 of its length and 5 feet tall for the remainder. During sampling, fans inside the chamber keep the gases mixed while stainless steel probes extract gas samples from the chamber through 12 sample ports. Gases extracted from the chamber are replaced by ambient air that enters the chamber through six ½-inch diameter vent lines that are distributed along one side of the chamber. An electrical firing circuit is used to remotely detonate the ordnance and release the tracer gas.

A number of different test methods were employed to collect and analyze the emission data that were used to develop emission factors for detonation of ordnance. Table 3 identifies each emission test method used; bracketed information identifies the purpose of using the method. The emissions data were collected using EPA test methods published in Title 40 of the Code of Federal Regulations, Part 50 (40 CFR 50); 40 CFR 60; and in *Compendium of Methods for the Determination of Toxic Organic Compounds in Ambient Air*.⁵ Some of the sample analytical procedures used were from EPA Office of Solid Waste (OSW) publication SW-846, *Test Methods for Evaluating Solid Waste, Physical/Chemical Methods*.⁶ Where necessary, the test methods were adapted to reflect application to the unique testing of ordnance detonation in the Smoke Chamber.

The following sections identify and briefly describe the test methods used to measure each compound or group of compounds. Additional information regarding the operation of the Smoke Chamber and the test methods used is presented in Reference 1. EPA-approved methods were used by the laboratories that provided sampling and analysis data.

Compound Test Method CO 40 CFR 60, Appendix A, EPA Method 10 - Determination of Carbon Monoxide *Emissions from Stationary Sources* [sampling and analysis] 40 CFR 60, Appendix A, EPA Method 3A - Determination of Oxygen and Carbon CO_2 Dioxide Concentrations in Emissions from Stationary Sources (Instrumental Analyzer *Procedure*) [sampling and analysis] 40 CFR 60, Appendix A, EPA Method 7E - Determination of Nitrogen Oxides NO_x Emissions from Stationary Sources (Instrumental Analyzer Procedure) [sampling and analysis] 40 CFR 60, Appendix A, EPA Method 6C - Determination of Sulfur Dioxide SO_2 Emissions from Stationary Sources (Instrumental Analyzer Procedure) [sampling and analysis] 40 CFR 60, Appendix A, EPA Method 17 - Determination of Particulate Emissions TSP *from Stationary Sources (in-Stack Filtration Method)* [sampling and analysis] Particulate Cascade impactor [sampling] Size 40 CFR 60, Appendix A, EPA Method 17 - Determination of Particulate Emissions Distribution *from Stationary Sources (in-Stack Filtration Method)* [analysis] Metal sample was obtained from TSP sample [sampling] Metals 40 CFR 60, Appendix A, EPA Method 29 - Determination of Metals Emissions from Stationary Sources [analysis] SW-846 Method 6010A - Inductively Coupled Plasma-Atomic Emission Spectrometry [analysis for all metals except mercury] SW-846 Method 7470 - Mercury in Liquid Waste (Manual Cold-Vapor Technique) [analysis for mercury] HCl/Cl₂ 40 CFR 60, Appendix A, EPA Method 26 - Determination of Hydrogen Halide and Halogen Emissions from Stationary Sources (Non-Isokinetic Method) [sampling] SW-846 Method 9057 - Determination of Chloride from HCl/Cl₂ Emission Sampling Train (Methods 0050 and 0051) By Anion Chromatography [analysis] EPA Compendium Method TO-12 - Method for the Determination of Non-Methane VOC Organic Compounds (NMOC) in Ambient Air Using Cryogenic Preconcentration and Direct Flame Ionization Detection (FID) [sampling and analysis] EPA Compendium Method TO-14 - Determination of Volatile Organic Compounds Speciated (VOCs) in Ambient Air Using SUMMA Passivated Canister Sampling and Gas VOC Chromatographic Analysis [sampling and analysis] **SVOC** SW-846 Method 0010 - *Modified Method 5 Sampling Train* [sampling] SW-846 Method 8270 - Semivolatile Organic Compounds by Gas Chromatography/ *Mass Spectrometry (GC/MS)* [analysis]

TABLE 3 SAMPLING AND ALYTICAL METHODS USED

TABLE 3 (cont.)

Compound	Test Method
Dioxins and Furans	40 CFR 60, Appendix A, EPA Method 23 - Determination of Polychlorinated Dibenzo-p-dioxins and Polychlorinated Dibenzofurans from Municipal Waste Combustors [sampling]
	SW-846 Method 8290 - Polychlorinated Dibenzodioxins (PCDDs) and Polychlorinated Dibenzofurans (PCDFs) by High-Resolution Gas Chromatography/High-Resolution Mass Spectrometry (HRGC/HRMS) [analysis]
Tracer Compound (SF ₆)	Grab sample [sampling] Gas Chromatograph/Electron Capture Detector [analysis]

2.1 Carbon Monoxide, Carbon Dioxide, Oxides of Nitrogen, and Sulfur Dioxide

Real-time concentrations of CO, CO₂, NO_x, and SO₂ that resulted from the use of ordnance in the Smoke Chamber were measured using a continuous emissions measurement system (CEMS). CO sampling and analysis was conducted in accordance with 40 CFR Part 60, Appendix A, Method 10 - *Determination of Carbon Monoxide Emissions from Stationary Sources*, with an API 300 nondispersive infrared analyzer. CO₂ sampling and analysis was conducted in accordance with 40 CFR Part 60, Appendix A, Method 3A - *Determination of Oxygen and Carbon Dioxide Concentrations in Emissions from Stationary Sources*, with a TECO Model 41C infrared analyzer. NO_x sampling and analysis was conducted in accordance with 40 CFR Part 60, Appendix A, Method 7E - *Determination of Nitrogen Oxides Emissions from Stationary Sources*, with an API 200A chemiluminescent NO-NO₂ gas analyzer. SO₂ sampling and analysis was conducted in accordance with 40 CFR Part 60, Appendix A, Method 6C-*Determination of Sulfur Dioxide Emissions from Stationary Sources*, with an API 100A fluorescent analyzer. For each run, the target minimum sampling time was 20 minutes.

2.2 Total Suspended Particulate

The TSP concentration that resulted from the use of ordnance in the Smoke Chamber was determined using a high-volume sampling and analysis procedure based on 40 CFR 60, Appendix A, Reference Method 17 - *Determination of Particulate Emissions from Stationary Sources (in-Stack Filtration Method)*. During each run, duplicate samples were obtained using two high-volume samplers operating simultaneously. For each run, the target minimum sampling time was 20 minutes. The sampling rate was recorded continuously using a data acquisition system (DAS). The TSP concentration was computed, in accordance with Method 17, by dividing the mass of TSP collected by the volume of air sampled, corrected to standard conditions.

2.3 Particulate Size Distribution

The particulate size distribution that resulted from the use of ordnance in the Smoke Chamber was determined using a cascade impactor system which consisted of a multistage, multijet impactor that aerodynamically separated the particles into eight size groups. A short probe was used to convey the gas stream to the impactor and each intermediate size group was collected on a preweighed filter. The particles sizes collected in each group varied from diameters greater than 15 microns to diameters of approximately 0.5 microns. Particles smaller than 0.5 microns were collected on a backup filter.

2.4 Metals

Metal concentrations that resulted from the use of ordnance in the Smoke Chamber were determined using particulate matter from the TSP samples collected as described in Section 2.2. After the TSP total weight gain was determined in the laboratory, a portion of the TSP filter was digested with concentrated hydrogen fluoride and nitric acid per 40 CFR 60, Appendix A, Method 29 - *Determination of Metals Emissions from Stationary Sources*. Alternatively, if insufficient TSP material was present, the entire filter was digested. The digestate was then analyzed for metals (except mercury) using inductively coupled argon plasma (ICAP) emission spectroscopy in accordance with SW-846 Method 6010A - *Inductively Coupled Plasma-Atomic Emission Spectrometry*. Mercury was determined by cold vapor atomic absorption spectroscopy (CVAAS) in accordance with SW-846 Method 7470 - *Mercury in Liquid Waste (Manual Cold-Vapor Technique)*. The concentration of each target metal was computed by dividing the mass of metal collected by the volume of air sampled, corrected to standard conditions.

2.5 Hydrochloric Acid and Chlorine

Hydrochloric acid (HCl) and chlorine (Cl₂) concentrations that resulted from the use of ordnance in the Smoke Chamber were measured in accordance with 40 CFR Part 60, Appendix A, Method 26 -*Determination of Hydrogen Chloride Emissions from Stationary Sources*. During each run, chamber gases were pulled through two sets of impingers in series containing dilute sulfuric acid and sodium hydroxide solutions. Collected samples were analyzed using SW-846 Method 9057 - *Determination of Chloride from HCl/Cl₂ Emission Sampling Train (Methods 0050 and 0051) by Anion Chromatography*. The concentrations of HCl and Cl₂ were computed by dividing the mass collected by the volume of air sampled, corrected to standard conditions. HCl was also measured by a continuous analyzer, but these results were not used for development of emission factors. For each run, the target minimum sampling time was 20 minutes.

2.6 Volatile Organic Compounds

VOC concentrations that resulted from the use of ordnance in the Smoke Chamber were determined using two methods from the Second Supplement to Compendium of Methods for the Determination of Toxic Organic Compounds in Ambient Air: (1) Method TO-12 - Method for the Determination of Non-methane Organic Compounds in Ambient Air using Cryogenic Preconcentration and Direct Flame Ionization Detection and (2) Method TO-14 - Determination of Volatile Organic Compounds in Ambient Air Sampling and Gas Chromatographic Analysis. For both procedures, air samples were collected in stainless steel SUMMA[®] canisters. Two or three identical canisters were used for each test run. The minimum sampling time for each VOC canister was 10 minutes.

2.7 Semivolatile Organic Compounds

SVOC concentrations that resulted from the use of ordnance in the Smoke Chamber were determined based on procedures found in SW-846 Method 0010 - *Modified Method 5 Sampling Train*. During each run, duplicate samples were collected using two PS-1 samplers that contained special sampling inlets (i.e., aluminum sampling modules) designed to hold 100-mm diameter quartz fiber filters to collect particulate matter, followed by XAD-2 adsorbent resin cartridges for collection of vapor phase SVOCs. A 20-minute sampling time was targeted. Following sampling, the filters and resin cartridges underwent solvent extraction and the mass of SVOC collected was quantitatively determined by GC/MS analysis following procedures in SW-846 Method 8270 - *Semivolatile Organic Compounds by Gas Chromatography/Mass Spectrometry (GC/MS)*. Unknown compounds, if any, were tentatively identified using computerized mass spectral matching techniques of the highest non-target "peaks."

2.8 Dioxin and Furan Compounds

Dioxin and furan compound concentrations that resulted from the use of ordnance in the Smoke Chamber were determined based on procedures found in 40 CFR 60, Appendix A, EPA Method 23 -*Determination of Polychlorinated Dibenzo-p-dioxins and Polychlorinated Dibenzofurans from Municipal Waste Combustors*. During each run, duplicate samples were obtained using two modified PS-1 samplers. The modified samplers used standard quartz filters, but the adsorbent cartridges contained XAD-2 resin sandwiched between polyurethane foam (PUF) plugs. A minimum sampling time of 20 minutes was targeted. After sampling, the filters and adsorbent cartridges underwent extraction with the appropriate solvent(s). The mass of dioxin and furan compounds collected was quantitatively determined following SW-846 Method 8290 - *Polychlorinated Dibenzodioxins (PCDDs) and Polychlorinated Dibenzofurans (PCDFs) by High-Resolution Gas Chromatography/High-Resolution Mass Spectrometry (HRGC/HRMS)*.

2.9 Tracer Compound

Sulfur hexafluoride (SF₆) was used as a tracer compound during each run to estimate the amount of sample dilution that occurred as a result of ambient air entering the test chamber during the run. Grab samples were collected five times during each run using evacuated canisters. The canisters were analyzed for the tracer compound using a GC with an electron capture detector.

3.0 TEST DATA ANALYSIS AND RATING

3.1 EPA Guidance Regarding Test Data Quality Ratings

Prior to inclusion of emission factors in AP-42, the reliability of the underlying emission test data must be appraised in accordance with the rating system specified in Reference 4. Under this rating system, test data are assigned a rating from A to D, where an "A" rating is assigned to the highest quality data. The criteria used to assign a specific data quality rating are summarized below.

- A Tests are performed by using an EPA reference test method, or when not applicable, a sound methodology. Tests are reported in enough detail for adequate validation and raw data are provided that can be used to duplicate the emission results presented in the report.
- **B** Tests are performed by a generally sound methodology, but lacking enough detail for adequate validation. Data are insufficient to completely duplicate the emission result presented in the report.
- **C** Tests are based on an unproven or new methodology, or are lacking a significant amount of background information.
- **D** Tests are based on a generally unacceptable method, but the method may provide an order-of-magnitude value for the source.

Four specific criteria are identified in Reference 4 for consideration to assist in the assignment of a test data quality rating. These four criteria are:

1. <u>Source operation</u>. If the manner in which the source was operated is well documented in the report and the source was operating within typical parameters during the test, an "A" rating should be assigned. If the report stated parameters that were typical, but lacked detailed information, a "B" rating should be assigned. If there is reason to believe the operation was not typical, a "C" or "D" rating should be assigned.

- 2. <u>Test methods and sampling procedures</u>. In developing the ratings, the estimated accuracy and precision of the test method as well as the adequacy of the documentation should be considered. In general, if a current EPA reference test method, appropriate for the source, was followed, the rating should be higher ("A" or "B"). If other methods were used, an assessment should be made of their validity. If it is judged that the method was likely to be inaccurate or biased, a lower rating ("C" or "D") should be given. A complete report should indicate whether any procedures deviated from standard methods and explain any deviations. If deviations were reported, an evaluation should be made of whether these were likely to influence the test results.
- 3. <u>Process information</u>. During testing, many variations in the process can occur without warning and sometimes without being noticed. Such variations can induce wide deviations in sampling results. If a large variation between test run results cannot be explained by information contained in the site final test report or from test reports of other sources, the data are suspect and should be given a lower rating or excluded. However, it should be recognized that a process may have highly variable emissions and a lower rating may not be appropriate solely on the basis of wide deviations in sampling results.
- 4. <u>Analysis and calculations</u>. Ideally, final test reports should contain original raw data sheets and other documentation such as gas parameters (dry cubic feet per minute, oxygen percentage), calculation sheets, or example calculations describing how the calculated emission results were obtained. If there are data sheets, the nomenclature and equations used should be compared to those specified by EPA to establish equivalency. The depth of review of the calculations should be dictated by the reviewers' confidence in the ability and conscientiousness of the tester, based on such factors as consistency of results and completeness of other areas of the final test report. Reports may indicate that raw data sheets were available, but were not included. If the final test report is of high quality based on the other criteria, the quality rating should not be lowered due to a lack of data sheets.

An overall test data quality rating should be assigned based upon the ratings assigned for each of the four criteria.

3.2 Analysis of Test Data

Data included in the final test report¹ were rated in accordance with the rating system described above. Results for each of the four criteria are presented in the following sections.

3.2.1 Source Operations

The manner by which the ordnance were deployed (i.e., used) is documented in the final test report. Each of the ordnance that was tested was deployed in a manner similar to that which would occur in the field. The tests appear to have replicated typical ordnance operating parameters; consequently, the test data should be assigned an "A" rating based on this criterion.

3.2.2 Test Methods and Sampling Procedures

The test methods and sampling procedures were evaluated as being appropriate and consistent with EPA test methods or sound methodology. Except as noted below, no problems of any significance were identified; consequently, the test data should be assigned an "A" rating based on this criterion.

3.2.2.1 CEMS-Measured Data

Although summaries of the CEMS data were provided for the tests,¹ raw CEMS data were not provided for the tests or for the pre- and post-test quality control (QC) activities. Whether CEMS QC measures specified under 40 CFR 60, Appendix A, EPA Methods 3A, 7E, and 6C were followed could not be verified. Furthermore, none of the calibration gas certifications were supplied. There was no evidence of bias in the data; however, based on the issues noted above, the test data for the CEMS-measured compounds (i.e., CO, CO₂, NO_x, and SO₂) should be assigned a "B" rating based on this criterion.

3.2.2.2 Compounds Sampled or Analyzed Using More than One Test Method or Analytical Method

Twenty-one compounds were either sampled or analyzed using two methods; these compounds are identified in Table 4. For each of these compounds, emission factors were calculated based upon the data measured using the more appropriate test or analytical method; data measured using the less appropriate method were ignored. The more appropriate method was identified by reviewing the methods and the target compound lists associated with each method. If a specific compound appeared on the target compound list for one method but not the other, the method targeting the compound was selected. If a specific compound appeared on the target compound lists for both methods, the method judged to provide the most accurate data was selected.

For compounds analyzed using both the TO-12 (VOC) and TO-14 (VOC) methods, the TO-14 method analysis was judged to be more accurate and was selected. For compounds analyzed using both the SW8270 (SVOC) and TO-14 (VOC) methods, the TO-14 method analysis was judged to be more accurate and was selected. For hydrochloric acid, which was analyzed using CEMS and SW9057, the SW9057 method analysis was judged to be more accurate and was selected.

Occasionally, the compound measurement from the less accurate method was chosen because the compound had poor precision between test runs for the sampling method that would have been more accurate under normal circumstances. These cases are noted in the footnotes to Table 4.

Compound	Selected Method	Other Method Employed
Acetophenone	TO-14 (VOC)	SW8270 (SVOC)
Benzene	TO-14 (VOC)	TO-12 (VOC)
1,3-Butadiene	TO-14 (VOC)	TO-12 (VOC)
1,2-Dichlorobenzene	TO-14 (VOC)	SW8270 (SVOC)
1,3-Dichlorobenzene	TO-14 (VOC)	SW8270 (SVOC)
1,4-Dichlorobenzene ^a	TO-14 (VOC)	SW8270 (SVOC)
Ethylbenzene	TO-14 (VOC)	TO-12 (VOC)
Hexachlorobutadiene ^b	TO-14 (VOC)	SW8270 (SVOC)
Hexachloroethane ^c	TO-14 (VOC)	TO-12 (VOC)
Hydrochloric acid ^d	SW9057 (HCl)	CEMS (HCl)
Methyl tert-butyl ether ^e	TO-14 (VOC)	TO-12 (VOC)

TABLE 4 SELECTED SAMPLING OR ANALYTICAL METHOD FOR COMPOUNDS MEASURED USING TWO SAMPLING OR ANALYTICAL METHODS

Compound	Selected Method	Other Method Employed
Nitrobenzene ^f	TO-14 (VOC)	SW8270 (SVOC)
Styrene	TO-14 (VOC)	TO-12 (VOC)
Toluene	TO-14 (VOC)	TO-12 (VOC)
1,2,4-Trichlorobenzene ^g	TO-14 (VOC)	SW8270 (SVOC)
1,2,4-Trimethylbenzene	TO-14 (VOC)	TO-12 (VOC)
m-Xylene, p-Xylene	TO-14 (VOC)	TO-12 (VOC)
o-Xylene	TO-14 (VOC)	TO-12 (VOC)
p-Ethyltoluene	TO-14 (VOC)	TO-12 (VOC)
d-Limonene ^h	TO-14 (VOC)	TO-12 (VOC)
1,3,5-Trimethylbenzene	TO-14 (VOC)	TO-12 (VOC)

TABLE 4 (cont.)

^a For DODIC G955, data collected from the SVOC sampling method were used to develop emission factors because this compound had a relative percent difference greater than 100 percent between VOC test samples.

- ^b For DODICs G940 and G955, data collected from the SVOC sampling method were used to develop emission factors because this compound had a relative percent difference greater than 100 percent between VOC test samples.
- ^c For DODICs G950 and G955, data analyzed using the TO-12 analytical method were used to develop emission factors because this compound had a relative percent difference greater than 100 percent between TO-14 analytical results.
- ^d For DODICs G940 and G950, data analyzed using the CEMS method were used to develop emission factors because this compound had a relative percent difference greater than 100 percent between the SW9057 analytical results.
- ^e For DODIC G940, data analyzed using the TO-12 analytical method were used to develop emission factors because this compound had a relative percent difference greater than 100 percent between TO-14 analytical results.
- ^f For DODIC G950, data collected from the SVOC sampling method were used to develop emission factors because this compound had a relative percent difference greater than 100 percent between VOC test samples.
- ^g For DODIC G940, data collected from the SVOC sampling method were used to develop emission factors because this compound had a relative percent difference greater than 100 percent between VOC test samples.
- ^h For DODIC G950, data analyzed using the TO-12 analytical method were used to develop emission factors because this compound had a relative percent difference greater than 100 percent between TO-14 analytical results.

3.2.2.3 Tentatively Identified Compounds

During the analysis of the SVOC data, the highest nontarget "peaks" were tentatively identified using computerized mass spectral matching techniques. Emission factors were developed for these tentatively identified compounds (TICs) if all of the following criteria were met.

1. The TIC corresponded to a unique compound (e.g., fluorene). Emission factors were not developed if the TIC corresponded to a class of compounds (e.g., unknown alcohol).

- 2. The TIC was not identified using another analysis method that provided higher confidence data. Emission factors were developed based upon the higher confidence analysis method if such data were available.
- 3. The TIC was not present in the method blank. Emission factors were not developed if the TIC was found in the corresponding method blank.

The number of SVOC that were tentatively identified as unique compounds, were not identified using a higher confidence method, and were not present in the method blank varied from zero to one compounds per ordnance. Emission factors were developed for all of these TICs, but because of the uncertainty in the true identity of the TICs, the test data were assigned a "C" rating.

3.2.2.4 Particulate Size Distribution Data

Particulate size distribution data were not available for the first test run for DODIC G940 or for the single test run conducted for DODIC G945. As a result, emission factors for small particles (e.g., PM-10 and PM-2.5) were not developed for these test runs.

3.2.3 Process Information

Ordnance are manufactured to tight tolerances and are expected to deploy in a very repeatable fashion. Consequently, the test data should be assigned an "A" rating based upon this criterion. However, large relative percent differences (i.e., greater than 100 percent) between test runs or sample trains were noted for several compounds. Specific instances in which these differences were noted are identified in Table 5. The equation below illustrates calculation of relative percent difference:

 $relative \ percent \ difference = \frac{test \ 1 \ concentration \ - \ test \ 2 \ concentration}{average \ of \ test \ 1 \ and \ test \ 2 \ concentrations} \ x \ 100\%$

Due to the large relative percent differences between test runs, the test data specifically identified in Table 5 were assigned a "C" rating. The remainder of the data should be assigned an "A" rating based on this criterion.

Compound	Applicable DODIC
Sulfur dioxide	G940
Acrolein	G950
Acrylonitrile	G940
Barium	G945
Beryllium	G940, G945, G950
Cadmium	G945
Carbon tetrachloride	G940, G955
Chlorine	G940
Chloroethane	G940
Chromium	G940

TABLE 5COMPOUNDS FOR WHICH LARGE RELATIVE PERCENT DIFFERENCESWERE NOTED BETWEEN TEST RUNS OR SAMPLE TRAINS

Compound	Applicable DODIC
Copper	G940
Dichlorodifluoromethane	G950
Ethylbenzene	G945
1,2,3,4,6,7,8-Heptachlorodibenzofuran	G940, G950
1,2,3,4,7,8,9-Heptachlorodibenzofuran	G940
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin	G940, G950
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin	G940, G950
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin	G940, G950
1,2,3,4,7,8-Hexachlorodibenzofuran	G940
1,2,3,6,7,8-Hexachlorodibenzofuran	G940
1,2,3,7,8,9-Hexachlorodibenzofuran	G940
2,3,4,6,7,8-Hexachlorodibenzofuran	G940, G950
Hexane	G940
Manganese	G950
Mercury	G950
Methylene chloride	G940
Nickel	G940
1,2,3,4,6,7,8,9-Octachlorodibenzofuran	G940
1,2,3,7,8-Pentachlorodibenzo-p-dioxin	G940
1,2,3,7,8-Pentachlorodibenzofuran	G940
2,3,4,7,8-Pentachlorodibenzofuran	G940
Propionaldehyde	G955
Selenium	G940
Silver	G940, G945, G950
Styrene	G945
Tetrachloroethylene	G940, G955
Trichloroethylene	G955
Trichlorofluoromethane	G940, G955
Vinyl chloride	G940
m-Xylene, p-Xylene	G945, G955
o-Xylene	G945
Zinc	G950
Acetic acid	G950

TABLE 5 (cont.)

Compound	Applicable DODIC
Acetoxyacetone	G940
Benzofuran	G945
i-Butane	G950
2,3-Butanedione	G940
trans-2-Butenal	All DODICs
i-Butene	G940
cis-2-Butene	G950
trans-2-Butene	G950
3-Buten-2-one	G950
1,2-Dichloro-3-methylbenzene	G955
1,1-Dichloro-2-propanone	G940
Dimethyldisulfide	G945
2,5-Dimethylfuran	G940, G955
Dimethyltrisulfide	G945, G955
2-Furaldehyde	G945
Furan	G940
Heptanal	G940
2-Heptanone	G940
1-Hexene	G940
Isoprene	G940
Methacrolein	G945
Methane, nitro-	G940, G945
4-Methylbenzonitrile	G940
2-Methyl-1-butene	G940, G950
3-Methyl-1-butene	G940, G950
Methyl ethyl ketone	G945
5-Methyl-2-furaldehyde	G940
2-Methylfuran	G950
3-Methylfuran	G940, G945
Methylnitrite	G955
Nonanal	G940
Octanal	G945, G955
2,3-Pentanedione	G940

TABLE 5 (cont.)

Compound	Applicable DODIC
3-Pentanone	G940
1-Pentene	G940
trans-2-Pentene	G950
3-Penten-2-one	G955
Phenylacetylene	G940
2-Thiophenecarboxaldehyde	G940, G955

TABLE 5 (cont.)

3.2.4 Analysis and Calculations

The test report¹ and supporting documentation² were reviewed to determine whether they contained all of the original raw data, other documentation, and example calculations. Although the test report did not contain raw CEMS data, the missing information was judged insufficient to result in a downgrade of the test data quality rating.

The raw data and sample calculations presented in the final test report were reviewed to determine if the emission factors presented in the report could be duplicated. Excel spreadsheets were created to assist with this review. Where differences were found between the emission factors calculated using the Excel spreadsheets and those presented in the test report, an examination was made to determine the reason for the differences.

Several minor errors were noted in the calculation of the emission factors within the test report, particularly with respect to the incorporation of analytical detection limits into the emission factors (see Sections 4.1 and 4.2) and the net explosive weight assumed for each ordnance. The emission factors presented in AP-42 are based upon the corrected spreadsheets. Based upon the raw data, other documentation, and the Excel spreadsheet calculations, the test data should be assigned an "A" rating.

Emission factors developed for compounds present in the method blank at levels of 20 percent to 50 percent of both test values were assumed to be biased high. Several compounds met this criterion and are identified in Table 6. For these compounds, the test data were assigned a "B" rating.

TABLE 6	COMPOUNDS FOUND IN THE METHOD BLANK AT LEVELS BETWEEN
	20 PERCENT AND 50 PERCENT OF BOTH TEST VALUES

Compound	Applicable DODIC	
Aluminum	G940, G945	
Barium	G945, G955	
Chromium	G950	
Mercury	G940, G945	
Nickel	G940	
Zinc	G940	
Magnesium	G945	

When compounds were found in the method blank at levels greater than 50 percent of both test values, the data were assumed to be suspect and no emission factors were developed. Several compounds met this criterion and are identified in Table 7.

TABLE 7 COMPOUNDS FOUND IN THE METHOD BLANK AT LEVELS GREATER THAN
50 PERCENT OF BOTH TEST VALUES

Compound	Applicable DODIC	
Aluminum	G950, G955	
Antimony	G940, G945, G950	
Arsenic	G940, G945, G950	
Barium	G950	
Cobalt	G940, G945, G950	
Nickel	G945, G950, G955	
Thallium	G940, G945, G950	
Magnesium	G950, G955	

3.3 Test Data Quality Ratings

Upon completing the analysis described in the preceding section of this document, the test data quality ratings assigned as a result of the four criteria were reviewed. This review led to a downgrading of some of the test data from an "A" rating to either a "B" rating or a "C" rating. Table 8 identifies the data quality ratings for all compounds that did not receive an "A" rating.

Compound	Data Quality Rating	Applicable DODIC
Carbon dioxide	В	All DODICs
Carbon monoxide	В	All DODICs
Oxides of nitrogen	В	All DODICs
Sulfur dioxide	В	G945, G950, G955
Sulfur dioxide	С	G940
Acrolein	С	G950
Acrylonitrile	С	G940
Aluminum	В	G940, G945
Barium	В	G955
Barium	С	G945
Beryllium	С	G940, G945, G950
Cadmium	С	G945
Carbon tetrachloride	С	G940, G955

TABLE 8 DOWNGRADED DATA QUALITY RATINGS

Compound	Data Quality Rating	Applicable DODIC
Chlorine	С	G940
Chloroethane	С	G940
Chromium	В	G950
Chromium	С	G940
Copper	С	G940
Dichlorodifluoromethane	С	G950
Ethylbenzene	С	G945
1,2,3,4,6,7,8-Heptachlorodibenzofuran	С	G940, G950
1,2,3,4,7,8,9-Heptachlorodibenzofuran	С	G940
1,2,3,4,7,8-Hexachlorodibenzo-p-dioxin	С	G940, G950
1,2,3,6,7,8-Hexachlorodibenzo-p-dioxin	С	G940, G950
1,2,3,7,8,9-Hexachlorodibenzo-p-dioxin	С	G940, G950
1,2,3,4,7,8-Hexachlorodibenzofuran	С	G940
1,2,3,6,7,8-Hexachlorodibenzofuran	С	G940
1,2,3,7,8,9-Hexachlorodibenzofuran	C	G940
2,3,4,6,7,8-Hexachlorodibenzofuran	С	G940, G950
Hexane	С	G940
Manganese	С	G950
Mercury	В	G940, G945
Mercury	С	G950
Methylene chloride	С	G940
Nickel	С	G940
1,2,3,4,6,7,8,9-Octachlorodibenzofuran	С	G940
1,2,3,7,8-Pentachlorodibenzo-p-dioxin	С	G940
1,2,3,7,8-Pentachlorodibenzofuran	С	G940
2,3,4,7,8-Pentachlorodibenzofuran	С	G940
Propionaldehyde	С	G955
Selenium	С	G940
Silver	С	G940, G945, G950
Styrene	С	G945
Tetrachloroethylene	С	G940, G955
Trichloroethylene	С	G955

TABLE 8 (cont.)

Compound	Data Quality Rating	Applicable DODIC
Trichlorofluoromethane	С	G940, G955
Vinyl chloride	С	G940
m-Xylene, p-Xylene	С	G945, G955
o-Xylene	С	G945
Zinc	В	G940
Zinc	С	G950
Acetic acid	С	G950
Acetoxyacetone	С	G940
7H-Benz[de]anthracen-7-one	С	G945
Benzofuran	С	G945
i-Butane	С	G950
2,3-Butanedione	С	G940
trans-2-Butenal	С	All DODICs
i-Butene	С	G940
cis-2-Butene	СС	G950
trans-2-Butene	С	G950
3-Buten-2-one	С	G950
1,2-Dichloro-3-methylbenzene	С	G955
1,1-Dichloro-2-propanone	С	G940
Dimethyldisulfide	С	G945
2,5-Dimethylfuran	С	G940, G955
Dimethyltrisulfide	С	G945, G955
2-Furaldehyde	С	G945
Furan	С	G940
Heptanal	С	G940
2-Heptanone	С	G940
1-Hexene	С	G940
Isoprene	С	G940
Magnesium	В	G945
Methacrolein	С	G945
Methane, nitro-	С	G940, G945
4-Methylbenzonitrile	С	G940

TABLE 8 (cont.)

Compound	Data Quality Rating	Applicable DODIC
2-Methyl-1-butene	С	G940, G950
3-Methyl-1-butene	С	G940, G950
Methyl ethyl ketone	С	G945
5-Methyl-2-furaldehyde	С	G940
2-Methylfuran	С	G950
3-Methylfuran	С	G940, G945
Methylnitrite	С	G955
Nonanal	С	G940
Octanal	С	G945, G955
2,3-Pentanedione	С	G940
3-Pentanone	С	G940
1-Pentene	С	G940
trans-2-Pentene	С	G950
3-Penten-2-one	С	G955
Phenylacetylene	C	- G 940
2-Thiophenecarboxaldehyde	С	G940, G955

TABLE 8 (cont.)

4.0 EMISSION FACTOR CALCULATIONS

The methodologies and procedures that were used to develop emission factors from the test data are described in this section. A similar approach was used to calculate emission factors for TSP, metals, SVOC, HCl, Cl₂, and dioxin/furan compounds. The calculation steps that were performed for each sampling train and each run are summarized below.

- 1. For compounds for which more than one test sample was obtained, analytical detection limits were incorporated into the test data.
- 2. The background compound concentration was calculated by dividing the mass of compound detected during the background run by the background run sample volume.
- 3. The test compound concentration was calculated by dividing the mass of compound detected during the test run by the test run sample volume.
- 4. A background-corrected concentration was calculated by subtracting the background concentration from the test concentration.
- 5. A dilution-corrected concentration was calculated by dividing the background-corrected concentration by the dilution correction factor.
- 6. The mass of compound released during the test run was calculated by multiplying the dilutioncorrected concentration by the volume of the Smoke Chamber.

- 7. Emission factors for each sample and sampling train or test run were calculated by dividing the mass of compound released by the number of ordnance detonated during the test run or by the NEW detonated during the test run, as appropriate.
- 8. Average emission factors were calculated for each compound.

Because concentration data (i.e., milligrams per cubic meter [mg/m³], parts per million by volume [ppmv], or parts per billion by volume [ppbv]) were recorded for VOC and CEMS-measured compounds, it was not necessary to calculate background and test concentrations as described in steps 2 and 3. Detection limits were applied directly to test compound concentrations of VOC and CEMS-measured compounds, as described in step 1. Where present, ppmv and ppbv values were converted to mg/m³. Emission factors for VOC and CEMS-measured compounds were then estimated in accordance with steps 4 through 8 described above.

The following sections describe each of the eight emission factor calculation steps listed above in more detail. In addition, Section 4.9 discusses the calculation of PM-10 and PM-2.5 based on the particle size distribution data.

4.1 Incorporation of Analytical Detection-Limits to the Test Data

In many cases, more than one test sample was obtained for a specific compound (i.e., more than one sample was obtained for a given test run or more than one test run was conducted). When multiple samples were obtained for the same compound, a comparison was made of all the sample data collected. Based upon the results of the comparison, the following adjustments were made to the test data:

- 1. If all of the samples indicated that a compound was "not detected," the sample data were not adjusted.
- 2. If all of the samples indicated that a compound was detected, the sample data were not adjusted.
- 3. If one or more of the samples indicated that a compound was detected and one or more of the samples indicated that a compound was not detected, the "not detected" values were replaced with a value equal to one half of the compound's analytical detection limit. The assumption inherent to this adjustment was that the measured presence of a compound in one or more samples was indicative of the compound's presence in all samples. The analytical detection limits for each sample were obtained from the test report.

4.2 Determination of Background Concentration

For TSP, metals, SVOC, HCl, Cl_2 , and dioxin/furan compounds, the background compound concentration (BC) was calculated by dividing the mass of compound detected during the background run (Bkgd mass) by the background run sample volume (Bkgd V). This calculation is illustrated by the following equation:

$$BC = \frac{Bkgd \ mass}{Bkgd \ V}$$

For VOC compounds, the background run data were used directly. Background data for CEMSmeasured compounds were recorded for each test run between the time the CEMS began sampling and the time of detonation. The background concentrations were assumed to equal representative values over the sampling period.

4.3 Determination of Test Compound Concentration

For TSP, metals, SVOC, HCl, Cl_2 , and dioxin/furan compounds, the test compound concentration (TC) was calculated by dividing the mass of compound measured during the test run (Test mass) by the test run sample volume (Test V). This calculation is illustrated by the following equation:

$$TC = \frac{Test \ mass}{Test \ V}$$

For VOC compounds, the test run data were used directly. For CEMS-measured compounds, the test compound concentration was determined as the arithmetic mean of the test data collected from the initial steady-state point until the end of the test.

4.4 Determination of Background-Corrected Concentration

For all compounds, the calculation of the background-corrected concentration (BCC) was dependent on whether the background (BC) and test (TC) concentrations were detected and whether they were less than, equal to, or greater than one another. The procedures used to calculate the background-corrected concentration for each sampling train and compound are described below and are displayed graphically in Figure 1.

- 1. If the test concentration was not detected (ND), the background-corrected concentration equaled ND.
- 2. If the test concentration was detected and the background concentration was not detected, the background-corrected concentration equaled the test concentration.
- 3. If the test and background concentrations were detected and the test concentration was less than or equal to the background concentration, the background-corrected concentration equaled 0.
- 4. If the test and background concentrations were detected and the test concentration was greater than the background concentration, the background concentration was subtracted from the test concentration. This calculation is illustrated by the following equation:

$$BCC = TC - BC$$

4.5 Determination of Dilution-Corrected Concentration

The dilution-corrected concentration (DCC) was calculated by dividing the background-corrected concentration by the applicable dilution correction factor (DCF). This calculation is illustrated by the following equation:

$$DCC = \frac{BCC}{DCF}$$

4.6 Determination of Mass of Compound Released

The mass of compound released was calculated by multiplying the dilution-corrected concentration by the volume of the Smoke Chamber. This calculation is illustrated by the following equation:

Mass compound released =
$$DCC x$$
 Smoke Chamber volume




Figure 1 Calculation of background-corrected concentration (BCC).

4.7 Determination of Emission Factors

Once the mass of compound released was calculated, two emission factors were developed for each sample and for each sampling train or test run: the mass of compound released per item (i.e., per single ordnance) and the mass of compound released per pound NEW. The NEW for all ordnance were determined from Reference 1 and verified using Reference 3.

4.8 Determination of Average Emission Factors

Steps 1 through 7, as described in Sections 4.1 through 4.7, are applicable to individual samples within individual sampling trains or test runs. The final step in the emission factor calculation process was to calculate average emission factors for each compound in terms of mass released per item and mass released per pound NEW. The average emission factors for each compound were calculated as the arithmetic mean of the individual samples associated with the compound. If all samples indicated that the compound was not detected (ND), then the average emission factor was assigned a value of ND. [Note: The minimum detection levels associated with the compounds that were not detected are presented in Appendix A.] Total dioxin/furan emission factors were calculated by summing the average emission factors for all dioxin/furan compounds.

4.9 Determination of PM-10 and PM-2.5 Emission Factors

PM-10 and PM-2.5 emission factors were calculated by multiplying the TSP emission factors, as calculated in accordance with Step 7 (above), by the weight percent of the TSP that was classified as PM-10 or PM-2.5, as determined from the particulate size distribution data. Average PM-10 and PM-2.5 emission factors were then calculated in accordance with Step 8.

5.0 EMISSION FACTOR RATINGS

The emission factors were appraised in accordance with the rating system specified in Reference 4. Under this rating system, emission factors are assigned a rating from A to E, where an "A" rating is assigned to the highest quality factors. The criteria used to assign a specific emission factor rating are summarized below.

- A Excellent. The emission factor was developed primarily from A- and B-rated source test data taken from many randomly chosen facilities in the industry population. The source category population was sufficiently specific to minimize variability.
- **B** Above average. The emission factor was developed primarily from A- or B-rated test data from a moderate number of facilities. Although no specific bias was evident, it was not clear if the facilities tested represented a random sample of the industry. As with the "A" rating, the source category population was sufficiently specific to minimize variability.
- **C** Average. The emission factor was developed primarily from A-, B- and/or C-rated test data from a reasonable number of facilities. Although no specific bias was evident, it was not clear if the facilities tested represented a random sample of the industry. As with the "A" rating, the source category population was sufficiently specific to minimize variability.
- **D** Below average. The emission factor was developed primarily from A-, B-, and C-rated test data from a small number of facilities, and there may have been reason to suspect that these facilities did not represent a random sample of the industry. There also may have been evidence of variability within the source category population.

E Poor. The emission factor was developed from C- and D-rated test data from a very limited number of facilities, and there may have been reason to suspect that the facilities tested did not represent a random sample of the industry. There also may have been evidence of variability within the source category population.

Two analyses were conducted to assign ratings to the ordnance emission factors. First, an analysis was conducted on an ordnance-specific basis. Second, an analysis was conducted using all available ordnance emission factor data. The second analysis was conducted to determine whether a sufficient correlation existed between emission factors for different but similar ordnance to allow the number of test data points to be increased to the point that higher emission factor ratings could be assigned than were possible when using the ordnance-specific approach. Both analyses are described below.

5.1 Emission Factor Ratings Assigned – Based on Ordnance-Specific Test Data

As previously described, emission factor ratings are dependent upon the test data quality, the number of test data points, the amount of variability present within a source category population, and the randomness of the source category sample. The following test data facts pertain to these rating criteria:

- 1. As described in Section 3 of this Background Document, the ordnance test data was primarily rated A or B. The test data for a few compounds was rated C.
- 2. Two tests were conducted or two sampling trains were used per ordnance.
- 3. Ordnance are manufactured to very tight tolerance levels so there is little variability within a specific type of ordnance.
- 4. There was no evidence that suggested the tested items within each type of ordnance were specially selected.

Emission factor ratings were assigned based upon these facts. The rationale used to accept or reject specific emission factor ratings follow.

- A: Rejected. The number of test data points was deemed to be insufficient to assign an A emission factor rating.
- B: Rejected. The number of test data points was deemed to be insufficient to assign a B emission factor rating.
- C: Accepted for most ordnance. The emission factors were developed using A- and B-rated test data, there is little variability among items, and there was no evidence that suggested the tested items were specially selected. Because of the limited number of data points, a C rating was deemed appropriate for this set of circumstances.
- D: Accepted for some ordnance. The emission factors were developed using C-rated test data, there is little variability among items, and there was no evidence that suggested the tested items were specially selected. Because of the limited number of data points, a D rating was deemed appropriate for this set of circumstances.
- E: Rejected. The ordnance described in this report were developed primarily using A- and B-rated test data rather than C- or D-rated data, there is little variability among items, and there was no evidence that suggested the tested items were specially selected. Therefore, an E emission factor rating was deemed inappropriate.

5.2 Emission Factor Ratings Assigned – Based on All Available Test Data

The proceeding sections of this Background Document concern the emission measurement methods, data analysis, and calculations used to develop emission factors for specific ordnance. However, USAEC's ordnance emission factor development program includes more than 200 ordnance that have been tested under more than 25 separate test series. Because many of these ordnance are similar in size and/or chemical composition, a statistical analysis was conducted to assess the similarity of the emission factors developed for similar ordnance. The results of this analysis were used to reevaluate the emission factor ratings assigned on an ordnance-specific basis.

USAEC characterized individual ordnance as falling into one of 17 separate categories, depending upon the size and/or chemical composition of the ordnance. The ordnance and their respective categories are identified in Table 9 along with a comment field describing the number of data points.

Category	DODIC	Ordnance Description	Test Series	Comment
CS	G963 ^a	M73A CS Riot Control Agent Hand Grenade	DPG VI	Data not yet
	K765 ^a	CS Riot Control Agent Capsule	DPG VI	available
	G900	TH3 AN-M14 Incendiary Grenade	EO5	
	G911	MK3A2 Offensive Hand Grenade	EO2	
	G911	MK3A2 Offensive Hand Grenade	EO6	
	K010	M4 Field Incendiary Burster	EO5	
	K145	M18A1 Antipersonnel Mine	EO2	
	M023	M112 Demolition Block Charge	EO1	
	M030	1/4-Pound Demolition Block Charge	EO1	
	M030	1/4-Pound Demolition Block Charge	EO3	
	M031	1/2-Pound Demolition Block Charge	DPG IVA	
Demolition	M032	1-Pound Demolition Block Charge	EO2	20+ data points
	M032	1-Pound Demolition Block Charge	EO3	201 data points
	M130 ^a	M6 Electric Blasting Cap	DPG VII	
	M130 ^a	M6 Electric Blasting Cap	EO7	
	M131 ^a	M7 Blasting Cap	EO7	
	M241	M10 High Explosive Universal Destructor	DPG IVB	
	M456	PETN Type 1 Detonating Cord	DPG IVA	
	M500 ^a	M21 REEF Line Cutter	FP10	
	M591	M1 Military Dynamite Demolition Block Charge	EO1	
	M913	M58A3 Linear Demolition Charge	EO3	

TABLE 9ORDNANCE CATEGORIZATION FOR EMISSION FACTOR
CORRELATION ASSESSMENT

Test Category DODIC Ordnance Description Comment Series ML05^a MK24 High Explosive Cutter EO11 **ML09** DPG IVA Linear Demolition Charge, Shaped 20 gr/ft Linear Demolition Charge, Shaped ML15 DPG IVA 225 gr/ft ML47^a M11 Blasting Cap EO7 MM50^a M221 Shaped Charge EO11 MN02^a M12 Blasting Cap EO7 Demolition, 20+ data points continued MN03^a EO7 M13 Blasting Cap MN06^a M14 Blasting Cap EO7 MN07^a M15 Delay Blasting Cap EO11 MN08^a M81 Igniter EO9 MN68^a M151 Booster Demolition Charge DPG VIII None PAX-11, Granular Powder Burn EO4 None PAX-11, Molded Pellet Detonation EO4 G878^a M228 Practice Hand Grenade Fuse DPG VI K051 M604 Anti-Tank Practice Mine Fuze EO6 N278^a -M564 MTSO Fuze EO11 N285^a M577 Fuze EO9 Fuze <10 data points N286^a M582 Fuze EO7 N335 M557 Point Detonating Fuze EO5 N340 M739A1 Point Detonating Fuze EO5 N464^a EO9 M732 Fuze G881 M67 Fragmentation Grenade EO1 M82 Smoke Simulant Screening Grenade G978 DPG V Launcher Grenade <10 data points G982 DPG V Terephthalic Acid Smoke Hand Grenade GG09^a M84 Non-Lethal Stun Hand Grenade EO12 M583A1 40-mm White Star Parachute DPG IVB B535 Cartridge B536 M585 40-mm White Star Cluster Cartridge DPG IVB 20+ data points Illumination M83A3, M83A2, & M83A1 60-mm B627 DPG V Illuminating Cartridge with Fuze M485A2 155-mm Illumination Round D505 DPG I (projectile)

TABLE 9 (cont.)

TABLE 9 (cont.)

Category	DODIC	Ordnance Description	Test Series	Comment
	L305	M195 Green Star Parachute Signal Flare	DPG I	
	L306	M158 Red Star Cluster Signal Illumination	DPG II	
	L307	M159 White Star Cluster Signal Illumination	DPG II	
Illumination,	L311	M126A1 Red Star Parachute Signal Flare	DPG II	
continued	L312	M127A1 White Star Parachute Signal Flare	DPG I	20+ data points
	L314	M125A1 Green Star Cluster Signal Flare	DPG I	
	L367 ^a	M22 Anti-Tank, Guided Missile, and Rocket Launching Simulator	DPG VI	
	L410 ^a	M206 Aircraft Countermeasure Flare	DPG VI	
Inert	HA11 ^a	Rocket, 2.75-inch Flechette with M255A1 Warhead	DPG VIII	Data not yet available
	C511	M490 105-mm Target Practice Tracer Cartridge (tracer)	EO6	
Large	C784 ^a	M831 120-mm Target Practice Tracer Cartridge	EO12	Only 1 data point yet available
	C785 ^a	M865 120-mm Target Practice Discarding Sabot Tracer Cartridge	EO12	
Medium	BA11 ^a	M1001 40-mm HVCC Cartridge	EO12	Data not yet
Medium	BA15 ^a	M769 60-mm FRP Cartridge	EO12	available
	A652	M220 20-mm TP-T Cartridge	FP9	
	A940	M910 25-mm Target Practice Discarding Sabot Tracer Cartridge	FP8	
	A976	M793 25-mm Target Practice Tracer Cartridge	FP8	
Medium-FP	B129 ^a	M789 30 mm CTG	EO9	10+ data points
	B519	M781 40-mm Practice Cartridge	FP2	
	B584	M918 40-mm Practice Cartridge	FP2	
	B505 ^a	M662 40-mm Red Star Parachute Cartridge	DPG VIII	
Mine	K042 ^a	M88 Volcano Practice Canister Mine	DPG VIII	Data not yet available
Mortar	CA03	XM929 120 mm Smoke Cartridge with M7334A1 Fuze	DPG V	Only 1 data point

TABLE 9	(cont.)
---------	---------

Category	DODIC	Ordnance Description	Test Series	Comment
	B542	M430 40-mm High Explosive Dual Purpose (HEDP) Cartridge (projectile)	EO3	
	B571	M383 40-mm High Explosive Cartridge (projectile)	EO3	
	B632	M49A4 60-mm High Explosive Cartridge (projectile)	EO3	
	B642	M720 60-mm High Explosive Cartridge (projectile)	EO6	
Projectile	BZ-13 ^a	M888 60-mm Cartridge with M935 PD Fuze	EO11	10+ data points
	C995	M136 AT4 Recoilless Rifle, 84-mm Cartridge (projectile)	EO3	
	H557	M72A3 66-mm High Explosive Antitank Rocket (warhead)	EO1	
	H708 ^a	M73 35-mm Subcaliber Practice Rocket	DPG VIII	
	None ^a	PAX-21, 60-mm Mortar	EO8	
	PJ02	FIM-92A Stinger-Basic Guided Missile (warhead)	EO6	
	B642	M720 60-mm High Explosive Cartridge (propelling charge)	FP4	
	B653 ^a	M766 60-mm Short Range Practice Mortar Cartridge	FP10	
	C226	M301A3 81-mm Illuminating Cartridge (propelling charge)	FP4	
	C379	M934 120-mm High Explosive Cartridge (Zone 1 - propelling charge)	FP8	
Propellant	C511	M490 105-mm Target Practice Tracer Cartridge (propelling charge)	FP5	20 - data points
Propellant	C784	M831 120-mm Target Practice Tracer Cartridge (propelling charge)	FP5	20+ data points
	C785	M865 120-mm Target Practice Discarding Sabot Tracer Cartridge (propelling charge)	FP5	
	C868	M821 81-mm High Explosive Cartridge (propelling charge)	FP4	
	C876	M880 81-mm Target Practice Short Range Cartridge (propelling charge)	FP4	
	CA09	M931 120-mm Full Range Practice Cartridge (Zone 1 - propelling charge)	FP8	

h	i		i	i
Category	DODIC	Ordnance Description	Test Series	Comment
Propellant, continued	CA09	M931 120-mm Full Range Practice Cartridge (Zone 4 - propelling charge)	FP8	20+ data points
	D533	M119A2 155-mm Propelling Charge (Zone 7)	FP5	
	D540	M3 155-mm Propelling Charge (Zone 3, M199 Cannon)	FP1	
	D540	M3 155-mm Propelling Charge (Zone 3, M199 Cannon)	FP5	
	D540	M3 155-mm Propelling Charge (Zone 3, M284 Cannon)	FP1	
	D540	M3 155-mm Propelling Charge (Zone 5, M199 Cannon)	FP1	
	D540	M3A1 155-mm Propelling Charge (Zone 3, M199 Cannon)	FP1	
	D540	M3A1 155-mm Propelling Charge (Zone 3, M284 Cannon)	FP1	
	D541	M4A2 155-mm Propelling Charge (Zone 7)	FP5	
	M174 ^a	MK209 Impulse Cartridge	FP10	
	M842 ^a	M1 Squib	EO7	
	M842 ^a	M79 Igniter	EO9	
	MD73 ^a	M796 Impulse Cartridge	FP10	
	PJ02	FIM-92A Stinger-Basic Guided Missile (flight motor)	FP7	
	PJ02	FIM-92A Stinger-Basic Guided Missile (launch motor)	EO5	
Pyrotechnic	H975 ^a	M274 2.75-inch Signature Smoke with H872 Warhead	DPG VIII	20+ data points
	L366	M74A1 Projectile Air Burst Simulator	DPG IVB	
	L495	M49A1 Surface Trip Flare	DPG II	
	L508 ^a	M72 Red Railroad Warning Fuse	DPG VI	
	L592	TOW Blast Simulator	DPG V	
	L594	M115A2 Ground Burst Simulator	DPG I	
	L595 ^a	M9 Liquid Projectile Air Burst Simulator	EO12	
	L596	M110 Flash Artillery Simulator	DPG I	
	L598	M117 Flash Booby Trap Simulator	DPG I	

TABLE 9 (cont.)

Test Category DODIC Ordnance Description Comment Series L599 M118 Illuminating Booby Trap Simulator DPG II L600 M119 Whistling Booby Trap Simulator DPG II L601 M116A1 Hand Grenade Simulator DPG I L602 M21 Artillery Flash Simulator DPG IVB L709 EO2 M25 Target Hit Simulator L709 XM25 Target Hit Simulator DPG V L720 EO6 M26 Target Kill Simulator $M327^{a}$ **FP10 Coupling Base Firing Device** Pyrotechnic, 20+ data points continued $M448^{a}$ EO11 M2 Percussion Detonator M626^a M1 Push Igniter EO9 M627^a M5 Pressure Release Igniter FP10 M630 M1 Pull Igniter DPG V $M630^{a}$ M1 Pull Igniter EO9 M670^a M700 Blasting Fuse EO11 EO9 M766^a M60 Igniter ML03^a M142 Firing Device EO11 Rocket, 2.75-inch Flechette, MK40 Mod 3 H459 FP7 Motor (propelling rocket) M72A3 66-mm High Explosive Antitank H557 FP4 Rocket (propelling rocket) M72A3 66-mm High Explosive Antitank H557 FP7 Rocket/Missile 10+ data points Rocket (propelling rocket) M73 35-mm Subcaliber Practice Rocket H708 FP9 Motor Rocket, 2.75-inch M267 Practice H974 Warhead, MK66 Mod 3 Motor (propelling FP7 rocket) M220 10 Gage Blank/Subcaliber Salute A010 FP9 Cartridge A011 12 Gage #00 Shot Cartridge FP9 A017^a 12 Gage #9 Shot Cartridge FP10 Small Arm-FP 20+ data points M855 5.56-mm Ball Cartridge (fired from A059 FP3 the M16A1 Rifle) M855 5.56-mm Ball Cartridge (fired from A059 FP3 the M16A2 Rifle)

TABLE 9 (cont.)

TABLE 9 (cont.)

Category	DODIC	Ordnance Description	Test Series	Comment
	A059	M855 5.56-mm Ball Cartridge (No-Lead)	FP4	
	A063	M856 5.56-mm Tracer Cartridge	FP3	
	A065	M862 5.56-mm Practice Ball Cartridge	FP3	
	A066	M193 5.56-mm Ball Cartridge	FP6	
	A068	M196 5.56-mm Tracer Cartridge	FP6	
	A080	M200 5.56-mm Blank Cartridge	FP3	
	A086	.22 Caliber Long Rifle Ball Cartridge	FP4	
	A106	.22 Caliber Standard Velocity Long Rifle Ball Cartridge	FP4	
	A111	M82 7.62-mm Blank Cartridge	FP3	
	A131	M62 7.62-mm Tracer Cartridge	FP6	
	A136	M118 7.62-mm Ball Match Cartridge	FP6	
	A143	M80 7.62-mm Ball Cartridge	FP3	20+ data points
	A171 -	M852 7.62-mm Ball Match Cartridge	FP6	
	A182	M1 .30 Caliber Ball Cartridge	FP6	
Small Arm-FP, continued	A212	M2 .30 Caliber Ball Cartridge	FP6	
continued	A218 -	M25 .30 Caliber Tracer Cartridge	FP9	
	A247	M72 .30 Caliber Ball Match Cartridge	FP6	
	A363	M882 9-mm Ball Cartridge	FP3	
	A365	M181 14.5-mm Trainer-Spotter Cartridge with 3-sec Delay (Artillery)	DPG V	
	A366 ^a	M182 14.5-mm Cartridge	EO11	
	A400	M41 .38 Caliber Special Ball Cartridge	FP9	
	A403	.38 Caliber Special Blank Cartridge	FP9	
	A475	M1911 .45 Caliber Ball Cartridge	FP3	
	A518	M903 .50 Caliber SLAP Cartridge	FP9	
	A518a	M962 .50 Caliber SLAP/T	FP10	
	A525	M2 .50 Caliber Armor Piercing Cartridge	FP8	
	A557	M17 .50 Caliber Tracer Cartridge	FP3	
	A557	M33 .50 Caliber Ball Cartridge	FP3	
	A598	M1A1 .50 Caliber Blank Cartridge	FP3	

TABLE 9 (cont.)

Category	DODIC	Ordnance Description	Test Series	Comment
Smoke	C870 ^a	M819 (IUK) 81-mm Red Phosphorus Smoke	DPG VII	
	G815 ^a	Red Phosphorus Smoke Screening Grenade Launcher (UK)	DPG VII	
	G930	Hexachloroethane Smoke Grenade	DPG V	
	G940	M18 Green Smoke Hand Grenade	DPG III	
	G945	M18 Yellow Smoke Hand Grenade	DPG III	
	G950	M18 Red Smoke Hand Grenade	DPG III	10+ data points
	G950	M18 Red Smoke Hand Grenade (new formulation)	DPG V	
	G955	M18 Violet Smoke Hand Grenade	DPG III	
	G955	M18 Violet Smoke Hand Grenade (new formulation)	DPG V	
	K866 ^a	ABC-M5 HC Ground Smoke Pot (MILES)	DPG VII	
	K867 ^a –	M4A2 Floating Smoke Pot	DPG VII	

^a Although testing may have been completed, emission factors for this ordnance have not yet been analyzed for inclusion in AP-42; therefore, these data were not included when the data correlation was assessed.

Within each of the 17 ordnance categories identified by USAEC, emission factors for each compound were compared. To allow the comparison of emission factors for ordnance with similar constituents but significant differences in net explosive weight, the comparison was made using the normalized emission factor units of mass of compound released per pound NEW. Based upon information provided by EPA,⁷ the following procedures were used to assess the data correlation:

- 1. The relative standard deviation, defined as the standard deviation divided by the mean, was calculated for each compound within each ordnance category.
- 2. If the relative standard deviation was less than 1.0, the evaluated emission factors were considered to demonstrate good correlation. As such, the rating for these emission factors <u>could</u> be elevated to a maximum of an A, depending on the number of data points within the evaluated ordnance category.
- 3. If the relative standard deviation was between 1.0 and 2.0, the evaluated emission factors were considered to demonstrate fair correlation. As such, the rating for these emission factors <u>could</u> be elevated to a maximum of a B, depending on the number of data points within the evaluated ordnance category.
- 4. If the relative standard deviation was greater than 2.0, the evaluated emission factors were considered to demonstrate poor correlation. As such, the emission factor rating could not be elevated, regardless of the amount of data available.

A poor correlation between emission factors was not necessarily construed as being indicative of poor test data. Rather, a poor correlation was more likely to indicate that the ordnance included in the category were not as similar in nature as anticipated by USAEC when the ordnance categories were defined.

In addition to assessing the data correlation, an assessment was made of the number of test data points available within each of the 17 ordnance categories. Because each ordnance test consisted of two test data points (i.e., two test runs per ordnance or two independent sampling trains were used during an ordnance test), the number of test data points available in each of the ordnance categories varied from 2 to 68. Based upon information provided by EPA,⁷ the following assumptions were used to assess whether sufficient category-specific test data points were available to justify elevating the emission factor ratings based on ordnance-specific data only:

- 1. If 20 or more data points were available, the emission factor rating could be elevated to a maximum of an A, provided that the data also demonstrated a good correlation.
- 2. If at least 10 but less than 20 data points were available, the emission factor rating could be elevated to a maximum of a B, provided that the data also demonstrated a good correlation.
- 3. If less than 10 data points were available, the emission factor rating could not be elevated, regardless of the data correlation.
- 4. If the data demonstrated a fair correlation and 20 or more data points were available, the emission factor rating could be elevated to a maximum of a B.
- 5. If the data demonstrated a fair correlation and at least 10 but less than 20 data points were available, the emission factor rating could be elevated to a maximum of a C.

Using the criteria specified above, the emission factor ratings assigned to ordnance in each of the 17 ordnance categories were reevaluated. This evaluation indicated that some of the emission factor ratings associated with ordnance included in nine categories could be elevated from a C or D rating to an A or B rating. These nine categories are:

- 1. Demolition
- 2. Illumination
- 3. Medium Firing Point
- 4. Projectiles
- 5. Propellants
- 6. Pyrotechnics
- 7. Rocket/Missile
- 8. Small Arms Firing Point
- 9. Smokes

A final assessment was made as to the emission factor rating assigned based on ordnance-specific test data only. If the original emission factor data rating assigned was a C, then the emission factor rating was elevated to an A or B, as appropriate, based upon the data for the whole ordnance category. If the original emission factor data rating assigned was a D, then the emission factor rating was elevated to a B or C, as appropriate, based upon the data for the whole ordnance category. The analysis is documented in an Excel spreadsheet that is located on the EPA website at: http://www.epa.gov/ttn/chief/ap42/index.html.

All four ordnance contained in the current test series were included in the Smoke category, which includes more than 10 test data points. As a result, some emission factor ratings associated with these ordnance were elevated. The emission factor ratings assigned are presented in Appendix A.

6.0 REFERENCES

- 1. Sampling Results for AEC Phase III Training Ordnance Emission Characterization, URS Corporation, Oak Ridge, TN, July 2001.
- Supporting information including Excel spreadsheets supplied upon request by the U.S. Army Dugway Proving Ground test support contractor, URS Group, Inc., Oak Ridge, TN, January 2005.
- Hazard Classification of United States Military Explosives and Munitions, Revision 11, U.S. Army Defense Ammunition Center, Logistics Review and Technical Assistance Office, McAlester, OK, February 2001.
- 4. *Procedures for Preparing Emission Factor Documents*, EPA-454/R-95-015, U.S. Environmental Protection Agency, Research Triangle Park, NC, November 1997.
- Compendium of Methods for the Determination of Toxic Organic Compounds in Ambient Air, Second Supplement, EPA/600/4-89/017, U.S. Environmental Protection Agency, Research Triangle Park, NC, June 1988.
- 6. *Test Methods for Evaluating Solid Waste, Physical/Chemical Methods (SW-846)*, U.S. Environmental Protection Agency, <u>http://www.epa.gov/epaoswer/hazwaste/test/sw846.htm</u>.
- 7. Information regarding the relationship between emission factor data correlation, the number of data points available, and the resulting emission factor rating assigned supplied upon request by Mr. Ron Myers, Measurement Policy Group, Office of Air Quality Planning and Standards, U.S. Environmental Protection Agency, Research Triangle Park, NC, June 2006.

APPENDIX A

COMPOUNDS ANALYZED AND EMISSION FACTORS DEVELOPED FOR ORDNANCE INCLUDED IN PHASE III TESTING AT DUGWAY PROVING GROUND, UTAH

This page left blank intentionally.

		Emission Factor ^{b,c}		Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
	Carbon Dioxide, Criteria Pollutants,		ane Hydrocarbor	18,
	and Total Suspen			
124-38-9	Carbon dioxide ^f	8.4 E-02	1.1 E-01	
630-08-0	Carbon monoxide ^f	1.2 E-02	1.6 E-02	
7439-92-1	Lead	3.4 E-05	4.6 E-05	
10102-44-0	Nitrogen dioxide	1.7 E-05	2.3 E-05	
10102-43-9	Nitric oxide ^f	6.4 E-05	8.8 E-05	
	Oxides of nitrogen ^f	1.2 E-04	1.6 E-04	
	PM-2.5	1.0 E-01	1.4 E-01	
	PM-10	1.3 E-01	1.7 E-01	
7446-09-5	Sulfur dioxide ^g	1.6 E-04	2.2 E-04	
	Total nonmethane hydrocarbons	2.1 E-03	2.9 E-03	
12789-66-1	Total suspended particulate ^f	1.3 E-01	1.7 E-01	
	Hazardous Air Pollutant	s and Toxic Che	micals	
83-32-9	Acenaphthene	ND	ND	3.1
208-96-8	Acenaphthylene	ND	ND	5.6
75-07-0	Acetaldehyde	3.4 E-04	4.6 E-04	
75-05-8	Acetonitrile	6.9 E-06	9.4 E-06	
98-86-2	Acetophenone	1.1 E-05	1.5 E-05	
53-96-3	2-Acetylaminofluorene	ND	ND	5.6
107-02-8	Acrolein	8.4 E-05	1.2 E-04	
107-13-1	Acrylonitrile ^g	5.1 E-07	6.9 E-07	
107-05-1	Allyl chloride	ND	ND	3.2 E-04
7429-90-5	Aluminum	9.0 E-05	1.2 E-04	
92-67-1	4-Aminobiphenyl	ND	ND	34.2
62-53-3	Aniline	ND	ND	12.8
120-12-7	Anthracene	ND	ND	4.0
7440-39-3	Barium	1.8 E-06	2.4 E-06	
71-43-2	Benzene	3.3 E-04	4.5 E-04	
92-87-5	Benzidine	ND	ND	108.6
56-55-3	Benzo[a]anthracene	ND	ND	3.6
205-99-2	Benzo[b]fluoranthene	ND	ND	4.8
207-08-9	Benzo[k]fluoranthene	ND	ND	2.4
191-24-2	Benzo[g,h,i]perylene	ND	ND	1.7

TABLE A1COMPOUNDS ANALYZED AND EMISSION FACTORS DEVELOPED FOR
DODIC G940, M18 GREEN SMOKE HAND GRENADE

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
50-32-8	Benzo[a]pyrene	ND	ND	5.3
100-44-7	Benzyl chloride	ND	ND	5.3
7440-41-7	Beryllium ^g	1.1 E-08	1.5 E-08	
74-83-9	Bromomethane	ND	ND	4.0 E-04
101-55-3	4-Bromophenylphenylether	ND	ND	2.6
106-99-0	1,3-Butadiene ^f	1.8 E-05	2.6 E-05	
71-36-3	n-Butyl alcohol	ND	ND	3.1 E-04
85-68-7	Butylbenzylphthalate	ND	ND	3.3
123-72-8	Butyraldehyde	5.0 E-06	6.9 E-06	
7440-43-9	Cadmium	ND	ND	8.4 E-04
86-74-8	Carbazole	ND	ND	3.9
75-15-0	Carbon disulfide ^f	4.1 E-05	5.7 E-05	
56-23-5	Carbon tetrachloride ^g	8.6 E-08	1.2 E-07	
463-58-1	Carbonyl sulfide	ND	ND	2.5 E-04
7782-50-5	Chlorine ^g	1.6 E-06	2.2 E-06	
106-47-8	p-Chloroaniline	ND	ND	6.6
108-90-7	Chlorobenzene ^f	6.7 E-06	9.2 E-06	
510-15-6	Chlorobenzilate	ND	ND	6.1
75-00-3	Chloroethane	1.9 E-07	2.7 E-07	
111-91-1	bis(2-Chloroethoxy)methane	ND	ND	4.3
111-44-4	bis(2-Chloroethyl)ether	ND	ND	3.2
67-66-3	Chloroform ^f	9.7 E-06	1.3 E-05	
108-60-1	bis(2-Chloroisopropyl)ether	ND	ND	4.1
74-87-3	Chloromethane	ND	ND	2.1 E-04
91-58-7	2-Chloronaphthalene	ND	ND	4.0
7005-72-3	4-Chlorophenylphenyl ether	ND	ND	2.7
7440-47-3	Chromium ^g	1.5 E-05	2.0 E-05	
218-01-9	Chrysene	ND	ND	3.6
7440-50-8	Copper ^g	4.2 E-07	5.7 E-07	
98-82-8	Cumene	ND	ND	3.3
110-82-7	Cyclohexane	ND	ND	1.0 E-04
2303-16-4	Diallate	ND	ND	2.9
53-70-3	Dibenz[a,h]anthracene	ND	ND	4.7
132-64-9	Dibenzofuran	ND	ND	3.8
106-93-4	1,2-Dibromoethane	ND	ND	7.8 E-04

TABLE A1 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
84-74-2	Dibutyl phthalate	ND	ND	3.0
95-50-1	1,2-Dichlorobenzene ^f	2.9 E-06	3.9 E-06	
541-73-1	1,3-Dichlorobenzene	2.0 E-07	2.7 E-07	
106-46-7	1,4-Dichlorobenzene	4.9 E-07	6.7 E-07	
91-94-1	3,3'-Dichlorobenzidine	ND	ND	6.6
75-71-8	Dichlorodifluoromethane	ND	ND	5.0 E-04
75-34-3	1,1-Dichloroethane	ND	ND	4.1 E-04
107-06-2	1,2-Dichloroethane	ND	ND	4.1 E-04
540-59-0	1,2-Dichloroethene	ND	ND	4.0 E-04
120-83-2	2,4-Dichlorophenol	ND	ND	5.2
78-87-5	1,2-Dichloropropane	ND	ND	4.7 E-04
10061-02-6	trans-1,3-Dichloro-1-propene	ND	ND	4.6 E-04
76-14-2	Dichlorotetrafluoroethane	ND	ND	7.1 E-04
60-11-7	p-Dimethylaminoazobenzene	ND	ND	6.9
57-97-6	7,12-Dimethylbenz[a]anthracene	ND	ND	8.8
119-93-7	3,3'-Dimethylbenzidine	ND	ND	60.2
105-67-9	2,4-Dimethylphenol	ND	ND	11.8
131-11-3	Dimethyl phthalate	ND	ND	4.0
99-65-0	1,3-Dinitrobenzene	ND	ND	24.5
534-52-1	4,6-Dinitro-o-cresol	ND	ND	20.5
51-28-5	2,4-Dinitrophenol	ND	ND	158.1
121-14-1	2,4-Dinitrotoluene	ND	ND	16.8
606-20-2	2,6-Dinitrotoluene	ND	ND	15.2
	Total dioxin/furan compounds	5.6 E-10	7.6 E-10	
100-41-4	Ethylbenzene	6.3 E-06	8.6 E-06	
74-85-1	Ethylene	7.3 E-05	1.0 E-04	
117-81-7	bis(2-Ethylhexyl)phthalate	ND	ND	4.4
206-44-0	Fluoranthene	ND	ND	1.8
86-73-7	Fluorene	ND	ND	3.3
76-13-1	Freon 113	ND	ND	7.8 E-04
35822-46-9	1,2,3,4,6,7,8-Heptachlorodibenzo-p- dioxin ^g	3.8 E-11	5.2 E-11	
67562-39-4	1,2,3,4,6,7,8- Heptachlorodibenzofuran ^g	4.4 E-11	6.0 E-11	
55673-89-7	1,2,3,4,7,8,9- Heptachlorodibenzofuran ^g	9.8 E-12	1.3 E-11	

TABLE A1 (cont.)

		Emission	a Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
118-74-1	Hexachlorobenzene	ND	ND	5.0
87-68-3	Hexachlorobutadiene	ND	ND	4.3
77-47-4	Hexachlorocyclopentadiene	ND	ND	69.9
39227-28-6	1,2,3,4,7,8-Hexachlorodibenzo-p- dioxin ^g	1.6 E-12	2.2 E-12	
57653-85-7	1,2,3,6,7,8-Hexachlorodibenzo-p- dioxin ^g	2.2 E-12	3.0 E-12	
19408-74-3	1,2,3,7,8,9-Hexachlorodibenzo-p- dioxin ^g	3.2 E-12	4.3 E-12	
70648-26-9	1,2,3,4,7,8- Hexachlorodibenzofuran ^g	1.5 E-11	2.1 E-11	
57117-44-9	1,2,3,6,7,8- Hexachlorodibenzofuran ^g	7.0 E-12	9.5 E-12	
72918-21-9	1,2,3,7,8,9- Hexachlorodibenzofuran ^g	9.5 E-12	1.3 E-11	
60851-34-5	2,3,4,6,7,8- Hexachlorodibenzofuran ^g	9.2 E-12	1.3 E-11	
67-72-1	Hexachloroethane	ND	ND	9.8 E-04
110-54-3	Hexane ^g	0	0	
7647-01-0	Hydrochloric acid	8.0 E-04	1.1 E-03	
193-39-5	Indeno[1,2,3-cd]pyrene	ND	ND	2.9
78-59-1	Isophorone	ND	ND	3.4
67-63-0	Isopropyl alcohol	1.3 E-05	1.7 E-05	
120-58-1	Isosafrole	ND	ND	19.9
556-61-6	Isothiocyanatomethane	ND	ND	3.0 E-04
7439-92-1	Lead	3.4 E-05	4.6 E-05	
7439-96-5	Manganese	5.3 E-06	7.2 E-06	
7439-97-6	Mercury	4.4 E-09	6.0 E-09	
56-49-5	3-Methylcholanthrene	ND	ND	3.9
75-09-2	Methylene chloride ^g	4.1 E-05	5.6 E-05	
108-10-1	Methyl isobutyl ketone	ND	ND	4.2 E-04
91-57-6	2-Methylnaphthalene	ND	ND	4.2
95-48-7	2-Methylphenol	ND	ND	4.9
109-06-8	2-Methylpyridine	ND	ND	12.7
1634-04-4	Methyl tert-butyl ether	ND	ND	1.0 E-04
91-20-3	Naphthalene	ND	ND	2.5
130-15-4	1,4-Naphthoquinone	ND	ND	12.7

TABLE A1 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
134-32-7	1-Naphthylamine	ND	ND	13.8
91-59-8	2-Naphthylamine	ND	ND	272.8
7440-02-0	Nickel ^g	7.3 E-06	1.0 E-05	
100-01-6	4-Nitroaniline	ND	ND	16.6
98-95-3	Nitrobenzene	ND	ND	5.1 E-04
88-75-5	2-Nitrophenol	ND	ND	15.9
100-02-7	4-Nitrophenol	ND	ND	147.9
56-57-5	4-Nitroquinoline-1-oxide	ND	ND	85.2
924-16-3	N-Nitrosodibutylamine	ND	ND	13.9
55-18-5	N-Nitrosodiethylamine	ND	ND	19.2
62-75-9	N-Nitrosodimethylamine	ND	ND	16.1
621-64-7	N-Nitrosodipropylamine	ND	ND	8.4
59-89-2	N-Nitrosomorpholine	ND	ND	5.2
100-75-4	N-Nitrosopiperidine	ND	ND	4.8
99-55-8	5-Nitro-o-toluidine	ND	ND	10.7
3268-87-9	1,2,3,4,6,7,8,9-Octachlorodibenzo- p-dioxin ^f	2.3 E-10	3.2 E-10	
39001-02-0	1,2,3,4,6,7,8,9- Octachlorodibenzofuran ^g	5.5 E-11	7.5 E-11	
608-93-5	Pentachlorobenzene	ND	ND	4.4
40321-76-4	1,2,3,7,8-Pentachlorodibenzo-p- dioxin ^g	2.7 E-12	3.7 E-12	
57117-41-6	1,2,3,7,8-Pentachlorodibenzofuran ^g	3.4 E-11	4.7 E-11	
57117-31-4	2,3,4,7,8-Pentachlorodibenzofuran ^g	6.1 E-12	8.3 E-12	
76-01-7	Pentachloroethane	ND	ND	9.8
82-68-8	Pentachloronitrobenzene	ND	ND	25.5
87-86-5	Pentachlorophenol	ND	ND	125.4
85-01-8	Phenanthrene	ND	ND	4.3
108-95-2	Phenol	ND	ND	4.5
7723-14-0	Phosphorus	ND	ND	1.6 E-02
23950-58-5	Pronamide	ND	ND	5.9
123-38-6	Propionaldehyde	4.6 E-05	6.2 E-05	
115-07-1	Propylene ^f	5.9 E-05	8.1 E-05	
129-00-0	Pyrene	ND	ND	4.3
110-86-1	Pyridine	ND	ND	11.8
94-59-7	Safrole	ND	ND	19.9

TABLE A1 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
7782-49-2	Selenium ^g	1.4 E-07	1.9 E-07	
7440-22-4	Silver ^g	1.0 E-07	1.4 E-07	
100-42-5	Styrene	ND	ND	4.3 E-04
1746-01-6	2,3,7,8-Tetrachlorodibenzo-p- dioxin ^g	3.2 E-12	4.3 E-12	
51207-31-9	2,3,7,8-Tetrachlorodibenzofuran ^g	8.5 E-11	1.2 E-10	
79-34-5	1,1,2,2-Tetrachloroethane	ND	ND	7.0 E-04
127-18-4	Tetrachloroethylene ^g	5.1 E-07	6.9 E-07	
108-88-3	Toluene	2.9 E-04	3.9 E-04	
95-53-4	o-Toluidine	ND	ND	9.1
120-82-1	1,2,4-Trichlorobenzene	ND	ND	3.5
71-55-6	1,1,1-Trichloroethane	ND	ND	5.5 E-04
79-00-5	1,1,2-Trichloroethane	ND	ND	5.5 E-04
79-01-6	Trichloroethylene	1.0 E-05	1.4 E-05	
75-69-4	Trichloromonofluoromethane ^g	1.5 E-08	2.0 E-08	
95-95-4	2,4,5-Trichlorophenol	ND	ND	13.5
88-06-2	2,4,6-Trichlorophenol	ND	ND	10.3
96-18-4	1,2,3-Trichloropropane	ND	ND	6.1 E-04
95-63-6	1,2,4-Trimethylbenzene	ND	ND	5.0 E-04
540-84-1	2,2,4-Trimethylpentane	ND	ND	1.0 E-04
75-01-4	Vinyl chloride	6.7 E-07	9.1 E-07	
75-35-4	Vinylidene chloride	3.4 E-07	4.6 E-07	
106-42-3, 108-38-3	m-Xylene, p-Xylene	3.5 E-05	4.7 E-05	
95-47-6	o-Xylene	3.8 E-06	5.2 E-06	
7440-66-6	Zinc	9.1 E-07	1.2 E-06	
	Other Pe	ollutants		
64-19-7	Acetic acid	1.8 E-05	2.5 E-05	
67-64-1	Acetone ^f	5.5 E-04	7.5 E-04	
592-20-1	Acetoxyacetone ^g	3.0 E-05	4.0 E-05	
74-86-2	Acetylene ^f	6.8 E-05	9.3 E-05	
100-52-7	Benzaldehyde ^f	2.3 E-05	3.2 E-05	
103-65-1	Benzene, propyl-	ND	ND	
271-89-6	Benzofuran	7.7 E-06	1.1 E-05	1.0 E-04
65-85-0	Benzoic acid	ND	ND	

TABLE A1 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
100-47-0	Benzonitrile ^f	3.5 E-05	4.8 E-05	82.1
100-51-6	Benzyl alcohol	ND	ND	
107-04-0	1-Bromo-2-chloroethane	ND	ND	4.2
3017-95-6	2-Bromo-1-chloropropane	ND	ND	5.9 E-04
75-28-5	i-Butane ^f	1.8 E-07	2.5 E-07	6.5 E-04
106-97-8	n-Butane ^f	2.1 E-06	2.8 E-06	
431-03-8	2,3-Butanedione ^g	1.1 E-04	1.5 E-04	
123-73-9	trans-2-Butenal ^g	2.8 E-05	3.8 E-05	
106-98-9	1-Butene ^f	9.0 E-06	1.2 E-05	
590-18-1	cis-2-Butene	4.5 E-06	6.1 E-06	
115-11-7	i-Butene ^g	1.0 E-05	1.4 E-05	
624-64-6	trans-2-Butene ^f	8.7 E-06	1.2 E-05	
78-94-4	3-Buten-2-one	3.6 E-05	4.9 E-05	
13466-78-9	delta-3-Carene	ND	ND	
109-69-3	1-Chlorobutane	ND	ND	1.0 E-04
622-98-0	1-Chloro-4-ethylbenzene	ND	ND	3.9 E-04
95-49-8	1-Chloro-2-methylbenzene	ND	ND	5.8 E-04
108-41-8	1-Chloro-3-methylbenzene	ND	ND	5.3 E-04
59-50-7	4-Chloro-3-methylphenol	ND	ND	5.3 E-04
95-57-8	2-Chlorophenol	ND	ND	5.3
287-92-3	Cyclopentane	ND	ND	1.0 E-04
96-37-7	Cyclopentane, methyl-	ND	ND	1.0 E-04
120-92-3	Cyclopentanone	7.8 E-06	1.1 E-05	1.0 E-04
142-29-0	Cyclopentene	ND	ND	
930-30-3	2-Cyclopenten-1-one	ND	ND	1.0 E-04
124-18-5	n-Decane	ND	ND	3.4 E-04
616-21-7	1,2-Dichlorobutane	ND	ND	1.0 E-04
32768-54-0	1,2-Dichloro-3-methylbenzene	ND	ND	4.6 E-04
594-37-6	1,2-Dichloro-2-methylpropane	ND	ND	5.3 E-04
87-65-0	2,6-Dichlorophenol	ND	ND	7.4
513-88-2	1,1-Dichloro-2-propanone ^g	4.5 E-06	6.1 E-06	
10061-01-5	cis 1,3-Dichloro-1-propene	ND	ND	4.6 E-04
84-66-2	Diethylphthalate	ND	ND	2.5
75-83-2	2,2-Dimethylbutane	1.4 E-06	1.9 E-06	
79-29-8	2,3-Dimethylbutane	ND	ND	1.0 E-04

TABLE A1 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
624-92-0	Dimethyldisulfide	ND	ND	3.9 E-04
625-86-5	2,5-Dimethylfuran ^g	7.7 E-06	1.1 E-05	
1071-26-7	2,2-Dimethylheptane	ND	ND	1.0 E-04
584-94-1	2,3-Dimethylhexane	ND	ND	1.0 E-04
589-43-5	2,4-Dimethylhexane	ND	ND	1.0 E-04
592-13-2	2,5-Dimethylhexane	ND	ND	1.0 E-04
565-59-3	2,3-Dimethylpentane	ND	ND	1.0 E-04
108-08-7	2,4-Dimethylpentane	ND	ND	1.0 E-04
	Dimethylphenethylamine	ND	ND	235.8
463-82-1	2,2-Dimethylpropane	ND	ND	1.0 E-04
3658-80-8	Dimethyltrisulfide	ND	ND	5.2 E-04
117-84-0	Di-n-octylphthlalate	ND	ND	3.8
74-84-0	Ethane	4.6 E-05	6.3 E-05	
64-17-5	Ethanol ^f	4.6 E-06	6.2 E-06	
1678-91-7	Ethylcyclohexane	ND	ND	1.0 E-04
619-99-8	3-Ethylhexane	ND	ND	1.0 E-04
62-50-0	Ethyl methanesulfonate	ND	ND	5.6
620-14-4	m-Ethyltoluene	ND	ND	1.0 E-04
611-14-3	o-Ethyltoluene	ND	ND	1.0 E-04
622-96-8	p-Ethyltoluene	ND	ND	5.0 E-04
98-01-1	2-Furaldehyde	1.2 E-04	1.6 E-04	
498-60-2	3-Furaldehyde	2.0 E-05	2.7 E-05	
110-00-9	Furan ^g	1.0 E-05	1.4 E-05	
109-99-9	Furan, tetrahydro-	ND	ND	3.0 E-04
111-71-7	Heptanal ^g	5.3 E-07	7.2 E-07	
142-82-5	Heptane	ND	ND	1.0 E-04
110-43-0	2-Heptanone ^g	1.5 E-07	2.1 E-07	
1888-71-7	Hexachloropropene	ND	ND	13.5
66-25-1	Hexanal	2.2 E-06	3.1 E-06	
591-78-6	2-Hexanone	1.4 E-06	2.0 E-06	
592-41-6	1-Hexene ^g	7.7 E-07	1.1 E-06	
7688-21-3	cis-2-Hexene	ND	ND	1.0 E-04
4050-45-7	trans-2-Hexene	ND	ND	1.0 E-04
78-79-5	Isoprene ^g	2.1 E-06	2.9 E-06	
143-50-0	Kepone	ND	ND	178.5

TABLE A1 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
5989-27-5	d-Limonene	ND	ND	5.7 E-04
7439-95-4	Magnesium	4.9 E-05	6.7 E-05	
78-85-3	Methacrolein	2.2 E-05	3.1 E-05	
75-52-5	Methane, nitro- ^g	2.2 E-07	2.9 E-07	
91-80-5	Methapyrilene	ND	ND	275.4
104-85-8	4-Methylbenzonitrile ^g	4.3 E-06	5.8 E-06	
563-80-4	3-Methyl-2-butanone	5.1 E-06	6.9 E-06	
563-46-2	2-Methyl-1-butene ^g	1.3 E-06	1.7 E-06	
513-35-9	2-Methyl-2-butene	1.8 E-06	2.5 E-06	
563-45-1	3-Methyl-1-butene	2.2 E-07	3.0 E-07	
108-87-2	Methylcyclohexane	ND	ND	1.0 E-04
78-93-3	Methyl ethyl ketone	1.5 E-04	2.0 E-04	
620-02-0	5-Methyl-2-furaldehyde ^g	3.9 E-05	5.3 E-05	
534-22-5	2-Methylfuran	1.7 E-05	2.3 E-05	
930-27-8	3-Methylfuran ^g	1.5 E-06	2.0 E-06	
592-27-8	2-Methylheptane	ND	ND	1.0 E-04
591-76-4	2-Methylhexane	ND	ND	1.0 E-04
589-34-4	3-Methylhexane	ND	ND	1.0 E-04
66-27-3	Methyl methanesulfonate	ND	ND	6.5
624-91-9	Methylnitrite	ND	ND	2.5 E-04
107-83-5	2-Methylpentane	ND	ND	1.0 E-04
96-14-0	3-Methylpentane	ND	ND	1.0 E-04
763-29-1	2-Methyl-1-pentene	ND	ND	1.0 E-04
625-27-4	2-Methyl-2-pentene	ND	ND	1.0 E-04
691-37-2	4-Methyl-1-pentene	ND	ND	1.0 E-04
691-38-3	cis-4-Methyl-2-pentene	ND	ND	1.0 E-04
554-14-3	2-Methylthiophene	ND	ND	4.1 E-04
616-44-4	3-Methylthiophene	ND	ND	4.1 E-04
88-74-4	2-Nitroaniline	ND	ND	23.6
99-09-2	3-Nitroaniline	ND	ND	8.5
10595-95-6	N-Nitrosomethylethylamine	ND	ND	15.7
930-55-2	N-Nitrosopyrrolidine	ND	ND	17.0
124-19-6	Nonanal ^g	1.4 E-06	2.0 E-06	
111-84-2	n-Nonane	ND	ND	1.0 E-04
821-55-6	2-Nonanone	ND	ND	5.9 E-04

TABLE A1 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
124-13-0	Octanal ^f	1.9 E-06	2.6 E-06	
111-65-9	Octane	ND	ND	1.0 E-04
1600-37-9	1,1,2,3,3-Pentachloro-1-propene	ND	ND	8.9 E-04
110-62-3	Pentanal	ND	ND	3.6 E-04
78-78-4	i-Pentane	ND	ND	1.0 E-04
109-66-0	n-Pentane	ND	ND	1.0 E-04
600-14-6	2,3-Pentanedione ^g	1.7 E-05	2.3 E-05	
107-87-9	2-Pentanone	1.1 E-05	1.5 E-05	
96-22-0	3-Pentanone ^g	2.9 E-06	4.0 E-06	
109-67-1	1-Pentene ^g	7.8 E-07	1.1 E-06	
627-20-3	cis-2-Pentene	6.2 E-07	8.5 E-07	
646-04-8	trans-2-Pentene	1.2 E-06	1.6 E-06	
107-39-1	2,4,4-Trimethyl-1-pentene	ND	ND	1.0 E-04
107-40-4	2,4,4-Trimethyl-2-pentene	ND	ND	1.0 E-04
1629-58-9	1-Penten-3-one	1.3 E-05	1.8 E-05	
3102-33-8	Pent-3-en-2-one	8.3 E-06	1.1 E-05	
62-44-2	Phenacetin	ND	ND	4.2
536-74-3	Phenylacetylene ^g	1.5 E-06	2.1 E-06	
80-56-8	alpha-Pinene	ND	ND	1.0 E-04
127-91-3	beta-Pinene	ND	ND	1.0 E-04
74-98-6	Propane	1.1 E-05	1.5 E-05	
95-94-3	1,2,4,5-Tetrachlorobenzene	ND	ND	3.7
58-90-2	2,3,4,6-Tetrachlorophenol	ND	ND	22.8
110-02-1	Thiophene ^f	6.1 E-06	8.3 E-06	
98-03-3	2-Thiophenecarboxaldehyde ^g	1.3 E-06	1.8 E-06	
545-06-2	Trichloroacetonitrile	ND	ND	6.0 E-04
108-67-8	1,3,5-Trimethylbenzene	ND	ND	5.0 E-04
16747-26-5	2,2,4-Trimethylhexane	ND	ND	1.0 E-04
565-75-3	2,3,4-Trimethylpentane	ND	ND	1.0 E-04
99-35-4	sym-Trinitrobenzene	ND	ND	17.3

TABLE A1 (cont.)

TABLE A1 (cont.)

- ^a CASRN = Chemical Abstracts Service Registry Number.
- ^b ND = nondetected.
- ^c Emission factors rated C unless otherwise noted.
- ^d NEW = net explosive weight. The NEW for this ordnance is 7.20 E-01 pounds per item.
- ^e Data provided for compounds that were not detected. ^f Emission factor rated B because of correlation with emission factors for similar ordnance and number of test data points
- ^g Emission factor rated D because the factor is based upon C-rated test data.

		Emission Factor ^{b,c}		Minimum		
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}		
	Carbon Dioxide, Criteria Pollutants, Total Nonmethane Hydrocarbons, and Total Suspended Particulate					
124-38-9	Carbon dioxide ^f	7.7 E-02	1.1 E-01			
630-08-0	Carbon monoxide ^f	4.0 E-03	5.4 E-03			
7439-92-1	Lead	1.5 E-05	2.0 E-05			
10102-44-0	Nitrogen dioxide	0	0			
10102-43-9	Nitric oxide ^f	8.1 E-05	1.1 E-04			
	Oxides of nitrogen ^f	7.8 E-05	1.1 E-04			
7446-09-5	Sulfur dioxide	9.9 E-04	1.3 E-03			
	Total nonmethane hydrocarbons	2.0 E-04	2.8 E-04			
12789-66-1	Total suspended particulate ^f	6.5 E-02	8.8 E-02			
Hazardous Air Pollutants and Toxic Chemicals						
83-32-9	Acenaphthene	ND	ND	6.1 E-01		
208-96-8	Acenaphthylene	ND	ND	1.1		
75-07-0	Acetaldehyde	9.2 E-06	1.3 E-05			
75-05-8	Acetonitrile	2.3 E-07	3.1 E-07			
98-86-2	Acetophenone	ND	ND	5.0 E-04		
53-96-3	2-Acetylaminofluorene	ND	ND	1.1		
107-02-8	Acrolein	2.6 E-06	3.5 E-06			
107-13-1	Acrylonitrile	1.1 E-07	1.5 E-07			
107-05-1	Allyl chloride	ND	ND	3.2 E-04		
7429-90-5	Aluminum	4.6 E-05	6.3 E-05			
92-67-1	4-Aminobiphenyl	ND	ND	6.8		
62-53-3	Aniline	ND	ND	2.5		
120-12-7	Anthracene	ND	ND	7.9 E-01		
7440-39-3	Barium ^g	6.0 E-07	8.2 E-07			
71-43-2	Benzene	7.1 E-06	9.7 E-06			
92-87-5	Benzidine	ND	ND	21.7		
56-55-3	Benzo[a]anthracene	ND	ND	7.2 E-01		
205-99-2	Benzo[b]fluoranthene	ND	ND	9.5 E-01		
207-08-9	Benzo[k]fluoranthene	ND	ND	4.9 E-01		
191-24-2	Benzo[g,h,i]perylene	ND	ND	3.4 E-01		
50-32-8	Benzo[a]pyrene	ND	ND	1.1		
100-44-7	Benzyl chloride	ND	ND	5.3 E-04		

TABLE A2COMPOUNDS ANALYZED AND EMISSION FACTORS DEVELOPED FOR
DODIC G945, M18 YELLOW SMOKE HAND GRENADE

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
7440-41-7	Beryllium ^g	1.7 E-08	2.4 E-08	
74-83-9	Bromomethane	ND	ND	4.0 E-04
101-55-3	4-Bromophenylphenylether	ND	ND	5.2 E-01
106-99-0	1,3-Butadiene ^f	1.0 E-06	1.4 E-06	
71-36-3	n-Butyl alcohol	ND	ND	3.1 E-04
85-68-7	Butylbenzylphthalate	ND	ND	6.6 E-01
123-72-8	Butyraldehyde	4.2 E-07	5.7 E-07	
7440-43-9	Cadmium ^g	7.3 E-09	9.9 E-09	
86-74-8	Carbazole	ND	ND	7.8 E-01
75-15-0	Carbon disulfide ^f	4.9 E-04	6.7 E-04	
56-23-5	Carbon tetrachloride	2.6 E-06	3.6 E-06	
463-58-1	Carbonyl sulfide	3.7 E-06	5.1 E-06	
7782-50-5	Chlorine	5.1 E-07	7.0 E-07	
106-47-8	p-Chloroaniline	ND	ND	1.3
108-90-7	Chlorobenzene ^f	4.0 E-06	5.5 E-06	
510-15-6	Chlorobenzilate	ND	ND	1.2
75-00-3	Chloroethane ^f	4.6 E-07	6.3 E-07	
111-91-1	bis(2-Chloroethoxy)methane	ND	ND	8.6 E-01
111-44-4	bis(2-Chloroethyl)ether	ND	ND	6.4 E-01
67-66-3	Chloroform ^f	2.7 E-05	3.7 E-05	
108-60-1	bis(2-Chloroisopropyl)ether	ND	ND	8.2 E-01
74-87-3	Chloromethane	ND	ND	2.1 E-04
91-58-7	2-Chloronaphthalene	ND	ND	8.0 E-01
7005-72-3	4-Chlorophenylphenyl ether	ND	ND	5.3 E-01
7440-47-3	Chromium	2.5 E-07	3.4 E-07	
218-01-9	Chrysene	ND	ND	7.2 E-01
7440-50-8	Copper	ND	ND	2.0 E-03
98-82-8	Cumene	ND	ND	1.0 E-04
110-82-7	Cyclohexane	ND	ND	1.0 E-04
2303-16-4	Diallate	ND	ND	5.8 E-01
53-70-3	Dibenz[a,h]anthracene	ND	ND	9.5 E-01
132-64-9	Dibenzofuran	ND	ND	7.5 E-01
106-93-4	1,2-Dibromoethane	ND	ND	7.8 E-04
84-74-2	Dibutyl phthalate	ND	ND	6.0 E-01
95-50-1	1,2-Dichlorobenzene ^f	5.1 E-06	7.0 E-06	

TABLE A2 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
541-73-1	1,3-Dichlorobenzene	3.0 E-06	4.1 E-06	
106-46-7	1,4-Dichlorobenzene	6.5 E-07	8.9 E-07	
91-94-1	3,3'-Dichlorobenzidine	ND	ND	1.3
75-71-8	Dichlorodifluoromethane	7.9 E-08	1.1 E-07	
75-34-3	1,1-Dichloroethane	2.3 E-07	3.1 E-07	
107-06-2	1,2-Dichloroethane	6.4 E-07	8.7 E-07	
540-59-0	1,2-Dichloroethene	1.1 E-05	1.5 E-05	96.0
120-83-2	2,4-Dichlorophenol	ND	ND	1.0
78-87-5	1,2-Dichloropropane	ND	ND	4.7 E-04
10061-02-6	trans-1,3-Dichloro-1-propene	ND	ND	4.6 E-04
76-14-2	Dichlorotetrafluoroethane	ND	ND	7.1 E-04
60-11-7	p-Dimethylaminoazobenzene	ND	ND	1.4
57-97-6	7,12-Dimethylbenz[a]anthracene	ND	ND	1.8
119-93-7	3,3'-Dimethylbenzidine	ND	ND	12.0
105-67-9	2,4-Dimethylphenol	ND	ND	2.3
131-11-3	Dimethyl phthalate	ND	ND	7.9 E-01
99-65-0	1,3-Dinitrobenzene	ND	ND	4.9
534-52-1	4,6-Dinitro-o-cresol	ND	ND	4.1
51-28-5	2,4-Dinitrophenol	ND	ND	31.5
121-14-1	2,4-Dinitrotoluene	ND	ND	3.4
606-20-2	2,6-Dinitrotoluene	ND	ND	3.0
	Total dioxin/furan compounds	4.6 E-09	6.2 E-09	
100-41-4	Ethylbenzene ^g	3.1 E-07	4.3 E-07	
74-85-1	Ethylene	2.2 E-05	3.0 E-05	
117-81-7	bis(2-Ethylhexyl)phthalate	ND	ND	8.8 E-01
206-44-0	Fluoranthene	ND	ND	3.6 E-01
86-73-7	Fluorene	ND	ND	6.6 E-01
76-13-1	Freon 113	8.1 E-09	1.1 E-08	
35822-46-9	1,2,3,4,6,7,8-Heptachlorodibenzo-p- dioxin	9.1 E-11	1.2 E-10	
67562-39-4	1,2,3,4,6,7,8- Heptachlorodibenzofuran	2.4 E-10	3.2 E-10	
55673-89-7	1,2,3,4,7,8,9- Heptachlorodibenzofuran	8.9 E-11	1.2 E-10	
118-74-1	Hexachlorobenzene	ND	ND	1.0
87-68-3	Hexachlorobutadiene	ND	ND	1.1 E-03

TABLE A2 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
77-47-4	Hexachlorocyclopentadiene	ND	ND	13.9
39227-28-6	1,2,3,4,7,8-Hexachlorodibenzo-p- dioxin	1.9 E-11	2.6 E-11	
57653-85-7	1,2,3,6,7,8-Hexachlorodibenzo-p- dioxin	5.0 E-11	6.9 E-11	
19408-74-3	1,2,3,7,8,9-Hexachlorodibenzo-p- dioxin	4.7 E-11	6.4 E-11	
70648-26-9	1,2,3,4,7,8-Hexachlorodibenzofuran	3.5 E-10	4.7 E-10	
57117-44-9	1,2,3,6,7,8-Hexachlorodibenzofuran	1.8 E-10	2.5 E-10	
72918-21-9	1,2,3,7,8,9-Hexachlorodibenzofuran	3.3 E-11	4.5 E-11	
60851-34-5	2,3,4,6,7,8-Hexachlorodibenzofuran	1.8 E-10	2.4 E-10	
67-72-1	Hexachloroethane	ND	ND	9.8 E-04
110-54-3	Hexane	7.9 E-08	1.1 E-07	
7647-01-0	Hydrochloric acid	2.8 E-06	3.8 E-06	
193-39-5	Indeno[1,2,3-cd]pyrene	ND	ND	5.8 E-01
78-59-1	Isophorone	ND	ND	6.8 E-01
67-63-0	Isopropyl alcohol	4.5 E-07	6.1 E-07	
120-58-1	Isosafrole	ND	ND	4.0
556-61-6	Isothiocyanatomethane	ND	ND	3.0 E-04
7439-92-1	Lead	1.5 E-05	2.0 E-05	
7439-96-5	Manganese	1.9 E-07	2.6 E-07	
7439-97-6	Mercury	1.3 E-09	1.8 E-09	
56-49-5	3-Methylcholanthrene	ND	ND	7.8 E-01
75-09-2	Methylene chloride	5.5 E-06	7.5 E-06	
108-10-1	Methyl isobutyl ketone	ND	ND	4.2 E-04
91-57-6	2-Methylnaphthalene	ND	ND	8.4 E-01
95-48-7	2-Methylphenol	ND	ND	9.7 E-01
109-06-8	2-Methylpyridine	ND	ND	2.4
1634-04-4	Methyl tert-butyl ether	ND	ND	3.7 E-04
91-20-3	Naphthalene	ND	ND	5.1 E-01
130-15-4	1,4-Naphthoquinone	ND	ND	2.5
134-32-7	1-Naphthylamine	ND	ND	2.7
91-59-8	2-Naphthylamine	ND	ND	54.4
100-01-6	4-Nitroaniline	ND	ND	3.3
98-95-3	Nitrobenzene	ND	ND	5.1 E-04
88-75-5	2-Nitrophenol	ND	ND	3.2

TABLE A2 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
100-02-7	4-Nitrophenol	ND	ND	29.5
56-57-5	4-Nitroquinoline-1-oxide	ND	ND	17.0
924-16-3	N-Nitrosodibutylamine	ND	ND	2.8
55-18-5	N-Nitrosodiethylamine	ND	ND	3.8
62-75-9	N-Nitrosodimethylamine	ND	ND	3.2
621-64-7	N-Nitrosodipropylamine	ND	ND	1.7
59-89-2	N-Nitrosomorpholine	ND	ND	1.0
100-75-4	N-Nitrosopiperidine	ND	ND	9.6 E-01
99-55-8	5-Nitro-o-toluidine	ND	ND	2.1
3268-87-9	1,2,3,4,6,7,8,9-Octachlorodibenzo- p-dioxin ^f	1.9 E-10	2.6 E-10	
39001-02-0	1,2,3,4,6,7,8,9- Octachlorodibenzofuran	7.0 E-10	9.6 E-10	
608-93-5	Pentachlorobenzene	ND	ND	8.8 E-01
40321-76-4	1,2,3,7,8-Pentachlorodibenzo-p- dioxin	6.5 E-11	8.9 E-11	
57117-41-6	1,2,3,7,8-Pentachlorodibenzofuran	5.3 E-10	7.3 E-10	
57117-31-4	2,3,4,7,8-Pentachlorodibenzofuran	3.5 E-10	4.8 E-10	
76-01-7	Pentachloroethane	ND	ND	2.0
82-68-8	Pentachloronitrobenzene	ND	ND	5.1
87-86-5	Pentachlorophenol	ND	ND	25.0
85-01-8	Phenanthrene	ND	ND	8.6 E-01
108-95-2	Phenol	ND	ND	9.0 E-01
7723-14-0	Phosphorus	ND	ND	8.8 E-03
23950-58-5	Pronamide	ND	ND	1.2
123-38-6	Propionaldehyde	ND	ND	2.4 E-04
115-07-1	Propylene ^f	5.9 E-06	8.1 E-06	
129-00-0	Pyrene	ND	ND	8.5 E-01
110-86-1	Pyridine	ND	ND	2.4
94-59-7	Safrole	ND	ND	4.0
7782-49-2	Selenium	ND	ND	3.0 E-03
7440-22-4	Silver ^g	8.8 E-09	1.2 E-08	
100-42-5	Styrene ^g	1.7 E-07	2.4 E-07	
1746-01-6	2,3,7,8-Tetrachlorodibenzo-p-dioxin	2.1 E-11	2.8 E-11	
51207-31-9	2,3,7,8-Tetrachlorodibenzofuran	1.4 E-09	2.0 E-09	
79-34-5	1,1,2,2-Tetrachloroethane	4.4 E-07	6.0 E-07	

TABLE A2 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
127-18-4	Tetrachloroethylene	5.5 E-06	7.5 E-06	
108-88-3	Toluene	2.1 E-06	2.9 E-06	
95-53-4	o-Toluidine	ND	ND	1.8
120-82-1	1,2,4-Trichlorobenzene	ND	ND	7.5 E-04
71-55-6	1,1,1-Trichloroethane	5.9 E-08	8.1 E-08	
79-00-5	1,1,2-Trichloroethane	1.2 E-06	1.7 E-06	
79-01-6	Trichloroethylene	8.6 E-06	1.2 E-05	
75-69-4	Trichloromonofluoromethane	7.3 E-08	1.0 E-07	
95-95-4	2,4,5-Trichlorophenol	ND	ND	2.7
88-06-2	2,4,6-Trichlorophenol	ND	ND	2.1
96-18-4	1,2,3-Trichloropropane	ND	ND	6.1 E-04
95-63-6	1,2,4-Trimethylbenzene	8.9 E-07	1.2 E-06	
540-84-1	2,2,4-Trimethylpentane	ND	ND	1.0 E-04
75-01-4	Vinyl chloride ^f	5.7 E-06	7.8 E-06	
75-35-4	Vinylidene chloride	4.2 E-06	5.7 E-06	
106-42-3, 108-38-3	m-Xylene, p-Xylene ^g	6.8 E-07	9.2 E-07	
95-47-6	o-Xylene ^g	2.2 E-07	3.0 E-07	
7440-66-6	Zinc	8.6 E-07	1.2 E-06	
	Other Po	llutants		
64-19-7	Acetic acid	ND	ND	2.5 E-04
67-64-1	Acetone ^f	1.0 E-04	1.4 E-04	
592-20-1	Acetoxyacetone	ND	ND	4.8 E-04
74-86-2	Acetylene ^f	2.6 E-05	3.5 E-05	
100-52-7	Benzaldehyde ^f	5.4 E-07	7.4 E-07	
82-05-3	7H-Benz[de]anthracen-7-one ^h	3.6 E-02	4.9 E-02	
103-65-1	Benzene, propyl-	ND	ND	1.0 E-04
271-89-6	Benzofuran ^g	9.3 E-08	1.3 E-07	
65-85-0	Benzoic acid	ND	ND	16.3
100-47-0	Benzonitrile ^f	2.5 E-07	3.4 E-07	
100-51-6	Benzyl alcohol	ND	ND	8.5 E-01
107-04-0	1-Bromo-2-chloroethane	ND	ND	5.9 E-04
3017-95-6	2-Bromo-1-chloropropane	ND	ND	6.5 E-04
75-28-5	i-Butane ^f	5.6 E-08	7.7 E-08	
106-97-8	n-Butane ^f	1.2 E-07	1.7 E-07	

TABLE A2 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
431-03-8	2,3-Butanedione	1.5 E-06	2.1 E-06	
123-73-9	trans-2-Butenal ^g	4.3 E-07	5.9 E-07	
106-98-9	1-Butene ^f	5.5 E-07	7.6 E-07	
590-18-1	cis-2-Butene	1.9 E-07	2.6 E-07	
115-11-7	i-Butene	1.9 E-06	2.5 E-06	
624-64-6	trans-2-Butene ^f	4.9 E-07	6.7 E-07	
78-94-4	3-Buten-2-one	4.1 E-07	5.6 E-07	
13466-78-9	delta-3-Carene	ND	ND	1.0 E-04
109-69-3	1-Chlorobutane	ND	ND	3.9 E-04
622-98-0	1-Chloro-4-ethylbenzene	ND	ND	5.8 E-04
95-49-8	1-Chloro-2-methylbenzene	ND	ND	5.3 E-04
108-41-8	1-Chloro-3-methylbenzene	ND	ND	5.3 E-04
59-50-7	4-Chloro-3-methylphenol	ND	ND	1.1
95-57-8	2-Chlorophenol	ND	ND	6.6 E-01
287-92-3	Cyclopentane	ND	ND	1.0 E-04
96-37-7	Cyclopentane, methyl-	2.2 E-07	2.9 E-07	
120-92-3	Cyclopentanone	1.1 E-07	1.5 E-07	
142-29-0	Cyclopentene	ND	ND	1.0 E-04
930-30-3	2-Cyclopenten-1-one	ND	ND	3.4 E-04
124-18-5	n-Decane	ND	ND	1.0 E-04
616-21-7	1,2-Dichlorobutane	ND	ND	5.3 E-04
32768-54-0	1,2-Dichloro-3-methylbenzene	ND	ND	6.7 E-04
594-37-6	1,2-Dichloro-2-methylpropane	ND	ND	5.3 E-04
3018-12-0	Dichloroacetonitrile	ND	ND	4.6 E-04
87-65-0	2,6-Dichlorophenol	ND	ND	1.5
10061-01-5	cis 1,3-Dichloro-1-propene	ND	ND	4.6 E-04
513-88-2	1,1-Dichloro-2-propanone	ND	ND	5.3 E-04
84-66-2	Diethylphthalate	ND	ND	5.0 E-01
75-83-2	2,2-Dimethylbutane	ND	ND	1.0 E-04
79-29-8	2,3-Dimethylbutane	ND	ND	1.0 E-04
624-92-0	Dimethyldisulfide ^g	3.4 E-07	4.6 E-07	
625-86-5	2,5-Dimethylfuran	ND	ND	4.0 E-04
1071-26-7	2,2-Dimethylheptane	ND	ND	1.0 E-04
584-94-1	2,3-Dimethylhexane	ND	ND	1.0 E-04
589-43-5	2,4-Dimethylhexane	ND	ND	1.0 E-04

TABLE A2 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
592-13-2	2,5-Dimethylhexane	ND	ND	1.0 E-04
565-59-3	2,3-Dimethylpentane	ND	ND	1.0 E-04
108-08-7	2,4-Dimethylpentane	ND	ND	1.0 E-04
	Dimethylphenethylamine	ND	ND	47.0
463-82-1	2,2-Dimethylpropane	ND	ND	1.0 E-04
3658-80-8	Dimethyltrisulfide ^g	2.6 E-07	3.6 E-07	
117-84-0	Di-n-octylphthalate	ND	ND	7.7 E-01
74-84-0	Ethane	1.8 E-07	2.5 E-07	
64-17-5	Ethanol ^f	3.5 E-07	4.8 E-07	
1678-91-7	Ethylcyclohexane	ND	ND	1.0 E-04
619-99-8	3-Ethylhexane	ND	ND	1.0 E-04
62-50-0	Ethyl methanesulfonate	ND	ND	1.1
620-14-4	m-Ethyltoluene	5.5 E-07	7.6 E-07	
611-14-3	o-Ethyltoluene	ND	ND	1.0 E-04
622-96-8	p-Ethyltoluene	4.0 E-07	5.4 E-07	
98-01-1	2-Furaldehyde ^g	1.1 E-07	1.4 E-07	
498-60-2	3-Furaldehyde	ND	ND	4.0 E-04
110-00-9	Furan	3.0 E-06	4.0 E-06	
109-99-9	Furan, tetrahydro-	ND	ND	3.0 E-04
111-71-7	Heptanal	3.7 E-07	5.0 E-07	
142-82-5	Heptane	2.0 E-07	2.7 E-07	
110-43-0	2-Heptanone	ND	ND	4.7 E-04
1888-71-7	Hexachloropropene	ND	ND	2.7
66-25-1	Hexanal	4.0 E-07	5.5 E-07	
591-78-6	2-Hexanone	ND	ND	4.2 E-04
592-41-6	1-Hexene	ND	ND	1.0 E-04
7688-21-3	cis-2-Hexene	ND	ND	1.0 E-04
4050-45-7	trans-2-Hexene	ND	ND	1.0 E-04
78-79-5	Isoprene	1.8 E-07	2.4 E-07	
143-50-0	Kepone	ND	ND	35.6
5989-27-5	d-Limonene	ND	ND	5.7 E-04
7439-95-4	Magnesium	7.6 E-06	1.0 E-05	
78-85-3	Methacrolein ^g	3.1 E-07	4.2 E-07	
75-52-5	Methane, nitro- ^g	2.5 E-07	3.4 E-07	
91-80-5	Methapyrilene	ND	ND	54.6

TABLE A2 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
104-85-8	4-Methylbenzonitrile	ND	ND	4.9 E-04
563-80-4	3-Methyl-2-butanone	3.9 E-07	5.4 E-07	
563-46-2	2-Methyl-1-butene	ND	ND	1.0 E-04
513-35-9	2-Methyl-2-butene	6.1 E-07	8.3 E-07	
563-45-1	3-Methyl-1-butene ^f	7.2 E-08	9.8 E-08	
108-87-2	Methylcyclohexane	4.0 E-07	5.5 E-07	
78-93-3	Methyl ethyl ketone ^g	2.0 E-06	2.7 E-06	
620-02-0	5-Methyl-2-furaldehyde	ND	ND	4.6 E-04
534-22-5	2-Methylfuran	6.5 E-07	8.8 E-07	
930-27-8	3-Methylfuran ^g	2.8 E-06	3.8 E-06	
592-27-8	2-Methylheptane	ND	ND	1.0 E-04
591-76-4	2-Methylhexane	ND	ND	1.0 E-04
589-34-4	3-Methylhexane	ND	ND	1.0 E-04
66-27-3	Methyl methanesulfonate	ND	ND	1.3
624-91-9	Methylnitrite	ND	ND	2.5 E-04
107-83-5	2-Methylpentane	ND	ND	1.0 E-04
96-14-0	3-Methylpentane	ND	ND	1.0 E-04
763-29-1	2-Methyl-1-pentene	ND	ND	1.0 E-04
625-27-4	2-Methyl-2-pentene	ND	ND	1.0 E-04
691-37-2	4-Methyl-1-pentene	ND	ND	1.0 E-04
691-38-3	cis-4-Methyl-2-pentene	ND	ND	1.0 E-04
554-14-3	2-Methylthiophene	2.1 E-06	2.8 E-06	
616-44-4	3-Methylthiophene	2.6 E-06	3.5 E-06	
88-74-4	2-Nitroaniline	ND	ND	4.7
99-09-2	3-Nitroaniline	ND	ND	1.7
10595-95-6	N-Nitrosomethylethylamine	ND	ND	3.1
930-55-2	N-Nitrosopyrrolidine	ND	ND	3.4
124-19-6	Nonanal	ND	ND	5.9 E-04
111-84-2	n-Nonane	2.8 E-07	3.8 E-07	
821-55-6	2-Nonanone	ND	ND	5.9 E-04
124-13-0	Octanal	2.8 E-07	3.9 E-07	
111-65-9	n-Octane	5.1 E-08	7.0 E-08	
1600-37-9	1,1,2,3,3-Pentachloro-1-propene	ND	ND	8.9 E-04
110-62-3	Pentanal	ND	ND	3.6 E-04
78-78-4	i-Pentane	1.4 E-07	1.9 E-07	

TABLE A2 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
109-66-0	n-Pentane	1.4 E-07	2.0 E-07	
600-14-6	2,3-Pentanedione	ND	ND	4.2 E-04
107-87-9	2-Pentanone	7.4 E-07	1.0 E-06	
96-22-0	3-Pentanone	ND	ND	3.6 E-04
109-67-1	1-Pentene	ND	ND	1.0 E-04
627-20-3	cis-2-Pentene	4.1 E-06	5.6 E-06	
646-04-8	trans-2-Pentene	ND	ND	1.0 E-04
107-39-1	2,4,4-Trimethyl-1-pentene	ND	ND	1.0 E-04
107-40-4	2,4,4-Trimethyl-2-pentene	ND	ND	1.0 E-04
1629-58-9	1-Penten-3-one	ND	ND	3.5 E-04
3102-33-8	Pent-3-en-2-one	ND	ND	3.5 E-04
62-44-2	Phenacetin	ND	ND	8.5 E-01
536-74-3	Phenylacetylene	ND	ND	4.2 E-04
80-56-8	alpha-Pinene	ND	ND	1.0 E-04
127-91-3	beta-Pinene	ND	ND	1.0 E-04
74-98-6	Propane	2.7 E-07	3.6 E-07	
95-94-3	1,2,4,5-Tetrachlorbenzene	ND	ND	7.3 E-01
58-90-2	2,3,4,6-Tetrachlorophenol	ND	ND	4.5
109-99-9	Furan, tetrahydro-	ND	ND	3.0 E-04
110-02-1	Thiophene ^f	1.4 E-05	1.9 E-05	
98-03-3	2-Thiophenecarboxaldehyde	ND	ND	4.7 E-04
545-06-2	Trichloroacetonitrile	ND	ND	6.0 E-04
108-67-8	1,3,5-Trimethylbenzene	6.6 E-07	9.0 E-07	
16747-26-5	2,2,4-Trimethylhexane	ND	ND	1.0 E-04
565-75-3	2,3,4-Trimethylpentane	ND	ND	1.0 E-04
99-35-4	sym-Trinitrobenzene	ND	ND	3.5

TABLE A2 (cont.)

^a CASRN = Chemical Abstracts Service Registry Number.

^b ND = nondetected.

^c Emission factors rated C unless otherwise noted. ^d NEW = net explosive weight. The NEW for this ordnance is 7.20 E-01 pounds per item. ^e Data provided for compounds that were not detected.

^f Emission factor rated B because of correlation with emission factors for similar ordnance and number of test data points

^g Emission factor rated D because the factor is based upon C-rated test data. ^h Emission factor rated D because the factor is for a tentatively identified compound.

		Emission Factor ^{b,c}	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
	Carbon Dioxide, Criteria Pollutants, and Total Susper		ane Hydrocarbor	18,
124-38-9	Carbon dioxide ^f	7.7 E-02	1.1 E-01	
630-08-0	Carbon monoxide ^f	5.8 E-03	7.9 E-03	
7439-92-1	Lead	1.9 E-05	2.6 E-05	
10102-44-0	Nitrogen dioxide	0	0	
10102-43-9	Nitric oxide ^f	3.0 E-04	4.1 E-04	
	Oxides of nitrogen ^f	4.2 E-04	5.8 E-04	
	PM-2.5	1.2 E-01	1.7 E-01	
	PM-10	1.4 E-01	1.9 E-01	
7446-09-5	Sulfur dioxide	4.3 E-04	5.9 E-04	
	Total nonmethane hydrocarbons	5.1 E-04	6.9 E-04	
12789-66-1	Total suspended particulate ^f	1.4 E-01	1.9 E-01	
	Hazardous Air Pollutant	s and Toxic Che	micals	
83-32-9	Acenaphthene	ND	ND	1.4
208-96-8	Acenaphthylene	ND	ND	2.7
75-07-0	Acetaldehyde	4.6 E-05	6.2 E-05	
75-05-8	Acetonitrile	2.9 E-06	4.0 E-06	
98-86-2	Acetophenone	ND	ND	5.0 E-04
53-96-3	2-Acetylaminofluorene	ND	ND	2.7
107-02-8	Acrolein ^g	2.6 E-05	3.6 E-05	
107-13-1	Acrylonitrile	9.7 E-07	1.3 E-06	
107-05-1	Allyl chloride	ND	ND	3.2 E-04
92-67-1	4-Aminobiphenyl	ND	ND	16
62-53-3	Aniline	ND	ND	6.0
120-12-7	Anthracene	ND	ND	1.9
71-43-2	Benzene	1.7 E-05	2.3 E-05	
92-87-5	Benzidine	ND	ND	51.4
56-55-3	Benzo[a]anthracene	ND	ND	1.7
205-99-2	Benzo[b]fluoranthene	ND	ND	2.3
207-08-9	Benzo[k]fluoranthene	ND	ND	1.2
191-24-2	Benzo[g,h,i]perylene	ND	ND	8.0 E-01
50-32-8	Benzo[a]pyrene	ND	ND	2.5
100-44-7	Benzyl chloride	ND	ND	5.3 E-04

TABLE A3COMPOUNDS ANALYZED AND EMISSION FACTORS DEVELOPED FOR
DODIC G950, M18 RED SMOKE HAND GRENADE

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
7440-41-7	Beryllium ^g	4.1 E-08	5.6 E-08	
74-83-9	Bromomethane	ND	ND	4.0 E-04
101-55-3	4-Bromophenylphenylether	ND	ND	1.2
106-99-0	1,3-Butadiene ^f	7.8 E-08	1.1 E-07	
123-72-8	Butanal	1.3 E-06	1.8 E-06	
71-36-3	n-Butyl alcohol	8.2 E-07	1.1 E-06	
85-68-7	Butylbenzylphthalate	ND	ND	1.6
7440-43-9	Cadmium	ND	ND	7.5 E-04
86-74-8	Carbazole	ND	ND	1.9
75-15-0	Carbon disulfide ^f	3.6 E-04	4.9 E-04	
56-23-5	Carbon tetrachloride	5.3 E-07	7.2 E-07	
463-58-1	Carbonyl sulfide	3.4 E-06	4.6 E-06	
7782-50-5	Chlorine	2.5 E-07	3.5 E-07	
106-47-8	p-Chloroaniline	ND	ND	3.1
108-90-7	Chlorobenzene ^f	1.3 E-06	1.7 E-06	
510-15-6	Chlorobenzilate	ND	ND	2.9
75-00-3	Chloroethane ^f	1.5 E-07	2.1 E-07	
111-91-1	bis(2-Chloroethoxy)methane	ND	ND	2.1
111-44-4	bis(2-Chloroethyl)ether	ND	ND	1.5
67-66-3	Chloroform ^f	1.3 E-05	1.8 E-05	
108-60-1	bis(2-Chloroisopropyl)ether	ND	ND	2.0
74-87-3	Chloromethane	ND	ND	2.1 E-04
91-58-7	2-Chloronaphthalene	ND	ND	1.9
7005-72-3	4-Chlorophenylphenyl ether	ND	ND	1.3
7440-47-3	Chromium	3.9 E-07	5.4 E-07	
218-01-9	Chrysene	ND	ND	1.7
7440-50-8	Copper	9.7 E-08	1.3 E-07	
98-82-8	Cumene	ND	ND	1.0E-04
110-82-7	Cyclohexane	ND	ND	1.0 E-04
2303-16-4	Diallate	ND	ND	1.4
53-70-3	Dibenz[a,h]anthracene	ND	ND	2.2
132-64-9	Dibenzofuran	ND	ND	1.8
106-93-4	1,2-Dibromoethane	ND	ND	7.8 E-04
84-74-2	Dibutyl phthalate	ND	ND	1.4
95-50-1	1,2-Dichlorobenzene ^f	1.2 E-06	1.6 E-06	

TABLE A3 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
541-73-1	1,3-Dichlorobenzene	1.1 E-07	1.5 E-07	
106-46-7	1,4-Dichlorobenzene	8.1 E-08	1.1 E-07	
91-94-1	3,3'-Dichlorobenzidine	ND	ND	3.1
75-71-8	Dichlorodifluoromethane ^g	1.6 E-07	2.2 E-07	
75-34-3	1,1-Dichloroethane	ND	ND	4.1 E-04
107-06-2	1,2-Dichloroethane	ND	ND	4.1 E-04
540-59-0	1,2-Dichloroethene	4.8 E-07	6.6 E-07	
120-83-2	2,4-Dichlorophenol	ND	ND	2.4
78-87-5	1,2-Dichloropropane	ND	ND	4.7 E-04
10061-02-6	trans-1,3-Dichloro-1-propene	ND	ND	4.6 E-04
76-14-2	Dichlorotetrafluoroethane	ND	ND	7.1 E-04
60-11-7	p-Dimethylaminoazobenzene	ND	ND	3.3
57-97-6	7,12-Dimethylbenz[a]anthracene	ND	ND	4.2
119-93-7	3,3'-Dimethylbenzidine	ND	ND	28.5
105-67-9	2,4-Dimethylphenol	ND	ND	5.6
131-11-3	Dimethyl phthalate	ND	ND	1.9
99-65-0	1,3-Dinitrobenzene	ND	ND	11.6
534-52-1	4,6-Dinitro-o-cresol	ND	ND	9.7
51-28-5	2,4-Dinitrophenol	ND	ND	74.8
121-14-1	2,4-Dinitrotoluene	ND	ND	8.0
606-20-2	2,6-Dinitrotoluene	ND	ND	7.2
	Total dioxin/furan compounds	6.6 E-10	8.9 E-10	
100-41-4	Ethylbenzene	9.6 E-07	1.3 E-06	
74-85-1	Ethylene	5.8 E-05	8.0 E-05	
117-81-7	bis(2-Ethylhexyl)phthalate	ND	ND	2.1
206-44-0	Fluoranthene	ND	ND	8.4 E-01
86-73-7	Fluorene	ND	ND	1.6
76-13-1	Freon 113	0	0	
35822-46-9	1,2,3,4,6,7,8-Heptachlorodibenzo-p- dioxin	3.4 E-11	4.6 E-11	
67562-39-4	1,2,3,4,6,7,8- Heptachlorodibenzofuran ^g	6.3 E-11	8.6 E-11	
55673-89-7	1,2,3,4,7,8,9- Heptachlorodibenzofuran	1.6 E-11	2.2 E-11	
118-74-1	Hexachlorobenzene	ND	ND	2.4
87-68-3	Hexachlorobutadiene	1.3 E-06	1.7 E-06	

TABLE A3 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
77-47-4	Hexachlorocyclopentadiene	ND	ND	33.1
39227-28-6	1,2,3,4,7,8-Hexachlorodibenzo-p- dioxin ^g	1.2 E-12	1.6 E-12	
57653-85-7	1,2,3,6,7,8-Hexachlorodibenzo-p- dioxin ^g	1.9 E-12	2.6 E-12	
19408-74-3	1,2,3,7,8,9-Hexachlorodibenzo-p- dioxin ^g	2.1 E-12	2.8 E-12	
70648-26-9	1,2,3,4,7,8-Hexachlorodibenzofuran	3.8 E-11	5.2 E-11	
57117-44-9	1,2,3,6,7,8-Hexachlorodibenzofuran	1.9 E-11	2.6 E-11	
72918-21-9	1,2,3,7,8,9-Hexachlorodibenzofuran	5.4 E-12	7.3 E-12	
60851-34-5	2,3,4,6,7,8- Hexachlorodibenzofuran ^g	2.5 E-11	3.4 E-11	
67-72-1	Hexachloroethane	ND	ND	1.8
110-54-3	Hexane	ND	ND	1.0 E-04
7647-01-0	Hydrochloric acid	6.8 E-05	9.3 E-05	
193-39-5	Indeno[1,2,3-cd]pyrene	ND	ND	1.4
78-59-1	Isophorone	ND	ND	1.6
67-63-0	Isopropyl alcohol	1.9 E-07	2.6 E-07	
120-58-1	Isosafrole	ND	ND	9.4
556-61-6	Isothiocyanatomethane	6.3 E-06	8.6 E-06	
7439-92-1	Lead	1.9 E-05	2.6 E-05	
7439-96-5	Manganese ^g	4.0 E-07	5.5 E-07	
7439-97-6	Mercury ^g	1.9 E-07	2.6 E-07	
56-49-5	3-Methylcholanthrene	ND	ND	1.9
75-09-2	Methylene chloride	2.4 E-06	3.3 E-06	
108-10-1	Methyl isobutyl ketone	9.8 E-07	1.3 E-06	
91-57-6	2-Methylnaphthalene	ND	ND	2.0
95-48-7	2-Methylphenol	ND	ND	2.3
109-06-8	2-Methylpyridine	ND	ND	5.8
1634-04-4	Methyl tert-butyl ether	ND	ND	3.7 E-04
91-20-3	Naphthalene	ND	ND	1.2
130-15-4	1,4-Naphthoquinone	ND	ND	6.0
134-32-7	1-Naphthylamine	ND	ND	6.5
91-59-8	2-Naphthylamine	ND	ND	129.1
100-01-6	4-Nitroaniline	ND	ND	7.8
98-95-3	Nitrobenzene	ND	ND	2.3

TABLE A3 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
88-75-5	2-Nitrophenol	ND	ND	7.5
100-02-7	4-Nitrophenol	ND	ND	70.0
56-57-5	4-Nitroquinoline-1-oxide	ND	ND	40.3
924-16-3	N-Nitrosodibutylamine	ND	ND	6.6
55-18-5	N-Nitrosodiethylamine	ND	ND	9.1
62-75-9	N-Nitrosodimethylamine	ND	ND	7.6
621-64-7	N-Nitrosodipropylamine	ND	ND	4.0
59-89-2	N-Nitrosomorpholine	ND	ND	2.4
100-75-4	N-Nitrosopiperidine	ND	ND	2.3
99-55-8	5-Nitro-o-toluidine	ND	ND	5.1
3268-87-9	1,2,3,4,6,7,8,9-Octachlorodibenzo- p-dioxin ^f	1.6 E-10	2.1 E-10	
39001-02-0	1,2,3,4,6,7,8,9- Octachlorodibenzofuran	2.0 E-10	2.8 E-10	
608-93-5	Pentachlorobenzene	ND	ND	2.1
40321-76-4	1,2,3,7,8-Pentachlorodibenzo-p- dioxin		ND	1.9 E-08
57117-41-6	1,2,3,7,8-Pentachlorodibenzofuran	3.2 E-11	4.3 E-11	
57117-31-4	2,3,4,7,8-Pentachlorodibenzofuran	1.7 E-11	2.3 E-11	
76-01-7	Pentachloroethane	ND	ND	4.6
82-68-8	Pentachloronitrobenzene	ND	ND	12.1
87-86-5	Pentachlorophenol	ND	ND	59.4
85-01-8	Phenanthrene	ND	ND	2.0
108-95-2	Phenol	ND	ND	2.1
7723-14-0	Phosphorus	4.8 E-07	6.5 E-07	
23950-58-5	Pronamide	ND	ND	2.8
123-38-6	Propionaldehyde	ND	ND	2.4 E-04
115-07-1	Propylene ^f	1.5 E-05	2.0 E-05	
129-00-0	Pyrene	ND	ND	2.0
110-86-1	Pyridine	ND	ND	5.6
94-59-7	Safrole	ND	ND	9.4
7782-49-2	Selenium	ND	ND	4.8 E-03
7440-22-4	Silver ^g	1.1 E-08	1.5 E-08	
100-42-5	Styrene	ND	ND	4.3 E-04
1746-01-6	2,3,7,8-Tetrachlorodibenzo-p-dioxin		ND	1.9 E-08
51207-31-9	2,3,7,8-Tetrachlorodibenzofuran	4.2 E-11	5.8 E-11	

TABLE A3 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
79-34-5	1,1,2,2-Tetrachloroethane	ND	ND	7.0 E-04
127-18-4	Tetrachloroethylene	6.4 E-07	8.7 E-07	
108-88-3	Toluene	1.7 E-06	2.4 E-06	
95-53-4	o-Toluidine	ND	ND	4.3
120-82-1	1,2,4-Trichlorobenzene	3.6 E-08	4.9 E-08	
71-55-6	1,1,1-Trichloroethane	ND	ND	5.5 E-04
79-00-5	1,1,2-Trichloroethane	ND	ND	5.5 E-04
79-01-6	Trichloroethylene	4.4 E-07	6.0 E-07	
75-69-4	Trichloromonofluoromethane	0	0	
95-95-4	2,4,5-Trichlorophenol	ND	ND	6.4
88-06-2	2,4,6-Trichlorophenol	ND	ND	4.9
96-18-4	1,2,3-Trichloropropane	ND	ND	6.1 E-04
95-63-6	1,2,4-Trimethylbenzene	ND	ND	5.0 E-04
540-84-1	2,2,4-Trimethylpentane	ND	ND	1.0 E-04
75-01-4	Vinyl chloride ^f	2.1 E-06	2.8 E-06	
75-35-4	Vinylidene chloride	4.0 E-07	5.5 E-07	
106-42-3, 108-38-3	m-Xylene, p-Xylene	4.0 E-06	5.4 E-06	
95-47-6	o-Xylene	3.7 E-07	5.0 E-07	
7440-66-6	Zinc ^g	9.0 E-06	1.2 E-05	
	Other Po	llutants		
64-19-7	Acetic acid ^g	7.8 E-07	1.1 E-06	
67-64-1	Acetone ^f	8.5 E-05	1.2 E-04	
592-20-1	Acetoxyacetone	ND	ND	4.8 E-04
74-86-2	Acetylene ^f	8.8 E-05	1.2 E-04	
100-52-7	Benzaldehyde ^f	5.8 E-06	7.9 E-06	
103-65-1	Benzene, propyl-	ND	ND	1.0 E-04
271-89-6	Benzofuran	1.7 E-06	2.3 E-06	
65-85-0	Benzoic acid	ND	ND	38.9
100-47-0	Benzonitrile ^f	3.9 E-06	5.3 E-06	
100-51-6	Benzyl alcohol	ND	ND	2.0
107-04-0	1-Bromo-2-chloroethane	ND	ND	5.9 E-04
3017-95-6	2-Bromo-1-chloropropane	ND	ND	6.5 E-04
75-28-5	i-Butane	1.8 E-08	2.5 E-08	
106-97-8	n-Butane ^f	1.4 E-07	1.9 E-07	

TABLE A3 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
431-03-8	2,3-Butanedione	7.9 E-06	1.1 E-05	
123-73-9	trans-2-Butenal ^g	7.0 E-06	9.6 E-06	
106-98-9	1-Butene ^f	3.1 E-06	4.2 E-06	
590-18-1	cis-2-Butene ^g	3.3 E-07	4.5 E-07	
115-11-7	i-Butene	2.0 E-06	2.7 E-06	
624-64-6	trans-2-Butene	1.4 E-06	1.9 E-06	
78-94-4	3-Buten-2-one ^g	4.3 E-06	5.9 E-06	
13466-78-9	delta-3-Carene	ND	ND	1.0 E-04
109-69-3	1-Chlorobutane	ND	ND	3.9 E-04
622-98-0	1-Chloro-4-ethylbenzene	ND	ND	5.8 E-04
95-49-8	1-Chloro-2-methylbenzene	9.7 E-05	1.3 E-04	
108-41-8	1-Chloro-3-methylbenzene	8.0 E-06	1.1 E-05	
59-50-7	4-Chloro-3-methylphenol	ND	ND	2.5
95-57-8	2-Chlorophenol	ND	ND	1.6
287-92-3	Cyclopentane	ND	ND	1.0 E-04
96-37-7	Cyclopentane, methyl-	ND	ND	1.0 E-04
120-92-3	Cyclopentanone	ND	ND	3.5 E-04
142-29-0	Cyclopentene	ND	ND	1.0 E-04
930-30-3	2-Cyclopenten-1-one	ND	ND	3.4 E-04
124-18-5	n-Decane	1.0 E-06	1.4 E-06	
3018-12-0	Dichloroacetonitrile	1.2 E-06	1.6 E-06	
616-21-7	1,2-Dichlorobutane	ND	ND	5.3 E-04
594-37-6	1,2-Dichloro-2-methylpropane	ND	ND	5.3 E-04
32768-54-0	1,2-Dichloro-3-methylbenzene	ND	ND	6.7 E-04
87-65-0	2,6-Dichlorophenol	ND	ND	3.5
513-88-2	1,1-Dichloro-2-propanone	ND	ND	5.3 E-04
10061-01-5	cis 1,3-Dichloro-1-propene	ND	ND	4.6 E-04
84-66-2	Diethylphthalate	ND	ND	1.2
75-83-2	2,2-Dimethylbutane	ND	ND	1.0 E-04
79-29-8	2,3-Dimethylbutane	ND	ND	1.0 E-04
624-92-0	Dimethyldisulfide	8.7 E-06	1.2 E-05	
625-86-5	2,5-Dimethylfuran	ND	ND	4.0 E-04
1071-26-7	2,2-Dimethylheptane	ND	ND	1.0 E-04
584-94-1	2,3-Dimethylhexane	ND	ND	1.0 E-04
589-43-5	2,4-Dimethylhexane	ND	ND	1.0 E-04

TABLE A3 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
592-13-2	2,5-Dimethylhexane	ND	ND	1.0 E-04
565-59-3	2,3-Dimethylpentane	ND	ND	1.0 E-04
108-08-7	2,4-Dimethylpentane	ND	ND	1.0 E-04
	Dimethylphenethylamine	ND	ND	111.5
463-82-1	2,2-Dimethylpropane	ND	ND	1.0 E-04
3658-80-8	Dimethyltrisulfide	1.3 E-05	1.8 E-05	
117-84-0	Di-n-octylphthalate	ND	ND	1.8
74-84-0	Ethane	1.4 E-06	2.0 E-06	
64-17-5	Ethanol ^f	9.1 E-07	1.2 E-06	
1678-91-7	Ethylcyclohexane	ND	ND	1.0 E-04
619-99-8	3-Ethylhexane	ND	ND	1.0 E-04
62-50-0	Ethyl methanesulfonate	ND	ND	2.7
620-14-4	m-Ethyltoluene	ND	ND	1.0 E-04
611-14-3	o-Ethyltoluene	ND	ND	1.0 E-04
622-96-8	p-Ethyltoluene	ND	ND	5.0 E-04
98-01-1	2-Furaldehyde	1.5 E-05	2.1 E-05	
498-60-2	3-Furaldehyde	5.8 E-06	8.0 E-06	
110-00-9	Furan	6.5 E-06	8.9 E-06	
109-99-9	Furan, tetrahydro-	ND	ND	3.0 E-04
111-71-7	Heptanal	3.8 E-07	5.2 E-07	
142-82-5	Heptane	ND	ND	1.0 E-04
110-43-0	2-Heptanone	1.9 E-07	2.6 E-07	
1888-71-7	Hexachloropropene	ND	ND	6.4
66-25-1	Hexanal	1.5 E-06	2.1 E-06	
591-78-6	2-Hexanone	ND	ND	4.2 E-04
592-41-6	1-Hexene	1.9 E-06	2.5 E-06	
4050-45-7	trans-2-Hexene	ND	ND	1.0 E-04
78-79-5	Isoprene	ND	ND	1.0 E-04
143-50-0	Kepone	ND	ND	84.5
5989-27-5	d-Limonene	ND	ND	1.0 E-04
78-85-3	Methacrolein	2.2 E-06	ND	
75-52-5	Methane, nitro-	8.8 E-07	2.9 E-06	
91-80-5	Methapyrilene	ND	1.2 E-06	129.8
104-85-8	4-Methylbenzonitrile	ND	ND	4.9 E-04
563-80-4	3-Methyl-2-butanone	ND	ND	3.6 E-04

TABLE A3 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
563-46-2	2-Methyl-1-butene ^g	4.3 E-07	5.9 E-07	
513-35-9	2-Methyl-2-butene	ND	ND	1.0 E-04
563-45-1	3-Methyl-1-butene	2.5 E-07	3.5 E-07	
108-87-2	Methylcyclohexane	ND	ND	1.0 E-04
78-93-3	Methyl ethyl ketone	3.5 E-06	4.8 E-06	
620-02-0	5-Methyl-2-furaldehyde	ND	ND	4.6 E-04
534-22-5	2-Methylfuran ^g	1.9 E-07	2.6 E-07	
930-27-8	3-Methylfuran	ND	ND	3.4 E-04
592-27-8	2-Methylheptane	ND	ND	6.5 E-04
591-76-4	2-Methylhexane	ND	ND	1.0 E-04
589-34-4	3-Methylhexane	ND	ND	1.0 E-04
66-27-3	Methyl methanesulfonate	ND	ND	3.1
624-91-9	Methylnitrite	1.5 E-06	2.0 E-06	
107-83-5	2-Methylpentane	ND	ND	1.0 E-04
96-14-0	3-Methylpentane	ND	ND	1.0 E-04
763-29-1	2-Methyl-1-pentene	ND	ND	1.0 E-04
625-27-4	2-Methyl-2-pentene	ND	ND	1.0 E-04
691-37-2	4-Methyl-1-pentene	ND	ND	1.0 E-04
691-38-3	cis-4-Methyl-2-pentene	ND	ND	1.0 E-04
554-14-3	2-Methylthiophene	1.2 E-06	1.6 E-06	
616-44-4	3-Methylthiophene	9.9 E-07	1.3 E-06	
88-74-4	2-Nitroaniline	ND	ND	11.1
99-09-2	3-Nitroaniline	ND	ND	4.0
10595-95-6	N-Nitrosomethylethylamine	ND	ND	7.4
930-55-2	N-Nitrosopyrrolidine	ND	ND	8.0
124-19-6	Nonanal	4.8 E-07	6.5 E-07	1.0 E-04
111-84-2	n-Nonane	ND	ND	1.0 E-04
821-55-6	2-Nonanone	ND	ND	5.9 E-04
124-13-0	Octanal ^f	5.3 E-07	7.3 E-07	
111-65-9	n-Octane	ND	ND	1.0 E-04
1600-37-9	1,1,2,3,3-Pentachloro-1-propene	ND	ND	8.9 E-04
110-62-3	Pentanal	ND	ND	3.6 E-04
78-78-4	i-Pentane	ND	ND	1.0 E-04
109-66-0	n-Pentane	ND	ND	1.0 E-04
600-14-6	2,3-Pentanedione	ND	ND	4.2 E-04

TABLE A3 (cont.)

Emission Factor ^{b,c} Minimum				
CASRN ^a	Commonwed	Emission		Minimum
CASINI	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
107-87-9	2-Pentanone	4.7 E-07	6.4 E-07	
96-22-0	3-Pentanone	ND	ND	3.6 E-04
109-67-1	1-Pentene	1.5 E-06	2.1 E-06	
627-20-3	cis-2-Pentene	ND	ND	1.0 E-04
646-04-8	trans-2-Pentene ^g	4.3 E-07	5.9 E-07	
107-39-1	1-Pentene, 2,4,4-trimethyl-	ND	ND	1.0 E-04
107-40-4	2-Pentene, 2,4,4-trimethyl-	ND	ND	1.0 E-04
1629-58-9	1-Penten-3-one	ND	ND	3.5 E-04
62-44-2	Phenacetin	ND	ND	2.0
536-74-3	Phenylacetylene	1.1 E-06	1.4 E-06	
80-56-8	alpha-Pinene	ND	ND	1.0 E-04
127-91-3	beta-Pinene	ND	ND	1.0 E-04
74-98-6	Propane	3.7 E-07	5.1 E-07	
95-94-3	1,2,4,5-Tetrachlorbenzene	ND	ND	1.7
58-90-2	2,3,4,6-Tetrachlorophenol	ND	ND	10.8
110-02-1	Thiophene ^f	2.4 E-05	3.2 E-05	
98-03-3	2-Thiophenecarboxaldehyde	1.1 E-06	1.5 E-06	
545-06-2	Trichloroacetonitrile	ND	ND	6.0 E-04
108-67-8	1,3,5-Trimethylbenzene	ND	ND	5.0 E-04
16747-26-5	2,2,4-Trimethylhexane	ND	ND	1.0 E-04
565-75-3	2,3,4-Trimethylpentane	ND	ND	1.0 E-04
99-35-4	sym-Trinitrobenzene	ND	ND	8.2

TABLE A3 (cont.)

^a CASRN = Chemical Abstracts Service Registry Number.

^b ND = nondetected.

^a ND = nondetected.
^c Emission factors rated C unless otherwise noted.
^d NEW = net explosive weight. The NEW for this ordnance is 7.20 E-01 pounds per item.
^e Data provided for compounds that were not detected.
^f Emission factor rated B because of correlation with emission factors for similar ordnance and number of test data points

^g Emission factor rated D because the factor is based upon C-rated test data.

		Emission Factor ^{b,c}		Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
	Carbon Dioxide, Criteria Pollutants, and Total Susper		ane Hydrocarbor	15,
124-38-9	Carbon dioxide ^f	4.3 E-02	5.9 E-02	
630-08-0	Carbon monoxide ^f	1.4 E-02	1.9 E-02	
7439-92-1	Lead	1.6 E-05	2.2 E-05	
10102-44-0	Nitrogen dioxide	0	0	
10102-43-9	Nitric oxide ^f	3.9 E-04	5.3 E-04	
	Oxides of nitrogen ^f	4.9 E-04	6.7 E-04	
	PM-2.5	1.0 E-01	1.4 E-01	
	PM-10	1.2 E-01	1.6 E-01	
7446-09-5	Sulfur dioxide	1.6 E-04	2.2 E-04	
	Total nonmethane hydrocarbons	1.1 E-03	1.5 E-03	
12789-66-1	Total suspended particulate ^f	1.2 E-01	1.6 E-01	
	Hazardous Air Pollutant	s and Toxic Che	micals	
83-32-9	Acenaphthene	ND	ND	1.5
208-96-8	Acenaphthylene	ND	ND	2.8
75-07-0	Acetaldehyde	8.0 E-05	1.1 E-04	
75-05-8	Acetonitrile	2.8 E-05	3.8 E-05	
98-86-2	Acetophenone	ND	ND	5.0 E-04
53-96-3	2-Acetylaminofluorene	ND	ND	2.8
107-02-8	Acrolein	3.1 E-06	4.2 E-06	
107-13-1	Acrylonitrile	3.6 E-05	4.9 E-05	
107-05-1	Allyl chloride	ND	ND	3.2 E-04
92-67-1	4-Aminobiphenyl	ND	ND	17.3
62-53-3	Aniline	ND	ND	6.4
120-12-7	Anthracene	ND	ND	2.0
7440-36-0	Antimony	ND	ND	7.9 E-03
7440-38-2	Arsenic	ND	ND	5.4 E-03
7440-39-3	Barium	7.2 E-07	9.9 E-07	
71-43-2	Benzene	8.1 E-05	1.1 E-04	
92-87-5	Benzidine	ND	ND	54.9
56-55-3	Benzo[a]anthracene	ND	ND	1.8
205-99-2	Benzo[b]fluoranthene	ND	ND	2.4
207-08-9	Benzo[k]fluoranthene	ND	ND	1.2

TABLE A4COMPOUNDS ANALYZED AND EMISSION FACTORS DEVELOPED FOR
DODIC G955, M18 VIOLET SMOKE HAND GRENADE

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
191-24-2	Benzo[g,h,i]perylene	ND	ND	8.6 E-01
50-32-8	Benzo[a]pyrene	ND	ND	2.7
100-44-7	Benzyl chloride	ND	ND	5.3 E-04
7440-41-7	Beryllium	9.8 E-10	1.3 E-09	
74-83-9	Bromomethane	ND	ND	4.0 E-04
101-55-3	4-Bromophenylphenylether	ND	ND	1.3
106-99-0	1,3-Butadiene ^f	1.2 E-05	1.7 E-05	
123-72-8	Butanal	1.0 E-06	1.4 E-06	
71-36-3	n-Butyl alcohol	ND	ND	3.1 E-04
85-68-7	Butylbenzylphthalate	ND	ND	1.7
7440-43-9	Cadmium	ND	ND	6.7 E-04
86-74-8	Carbazole	ND	ND	2.0
75-15-0	Carbon disulfide ^f	1.7 E-04	2.3 E-04	
56-23-5	Carbon tetrachloride ^g	1.1 E-06	1.5 E-06	
463-58-1	Carbonyl sulfide	1.5 E-05	2.0 E-05	
7782-50-5	Chlorine	1.6 E-06	2.2 E-06	
106-47-8	p-Chloroaniline	ND	ND	3.3
108-90-7	Chlorobenzene ^f	2.4 E-06	3.2 E-06	
510-15-6	Chlorobenzilate	ND	ND	3.1
75-00-3	Chloroethane ^f	2.0 E-07	2.8 E-07	
111-91-1	bis(2-Chloroethoxy)methane	ND	ND	2.2
111-44-4	bis(2-Chloroethyl)ether	ND	ND	1.6
67-66-3	Chloroform ^f	4.2 E-06	5.7 E-06	
108-60-1	bis(2-Chloroisopropyl)ether	ND	ND	2.1
74-87-3	Chloromethane	ND	ND	2.1 E-04
91-58-7	2-Chloronaphthalene	ND	ND	2.0
7005-72-3	4-Chlorophenylphenyl ether	ND	ND	1.4
7440-47-3	Chromium	3.0 E-07	4.2 E-07	
218-01-9	Chrysene	ND	ND	1.8
7440-48-4	Cobalt	4.0 E-08	5.5 E-08	
7440-50-8	Copper	ND	ND	2.9 E-03
98-82-8	Cumene	ND	ND	1.0 E-04
110-82-7	Cyclohexane	ND	ND	1.0 E-04
2303-16-4	Diallate	ND	ND	1.5
53-70-3	Dibenz[a,h]anthracene	ND	ND	2.4

TABLE A4 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
132-64-9	Dibenzofuran	ND	ND	1.9
106-93-4	1,2-Dibromoethane	ND	ND	7.8 E-04
84-74-2	Dibutyl phthalate	ND	ND	1.5
95-50-1	1,2-Dichlorobenzene ^f	8.3 E-07	1.1 E-06	
541-73-1	1,3-Dichlorobenzene	1.1 E-07	1.5 E-07	
106-46-7	1,4-Dichlorobenzene	ND	ND	2.5
91-94-1	3,3'-Dichlorobenzidine	ND	ND	3.3
75-71-8	Dichlorodifluoromethane	4.9 E-08	6.6 E-08	
75-34-3	1,1-Dichloroethane	ND	ND	4.1 E-04
107-06-2	1,2-Dichloroethane	ND	ND	4.1 E-04
540-59-0	1,2-Dichloroethene	ND	ND	4.0 E-04
120-83-2	2,4-Dichlorophenol	ND	ND	2.6
78-87-5	1,2-Dichloropropane	ND	ND	4.7 E-04
10061-02-6	trans-1,3-Dichloro-1-propene	ND	ND	4.6 E-04
76-14-2	Dichlorotetrafluoroethane	ND	ND	7.1 E-04
60-11-7	p-Dimethylaminoazobenzene	ND	ND	3.5
57-97-6	7,12-Dimethylbenz[a]anthracene	ND	ND	4.5
119-93-7	3,3'-Dimethylbenzidine	ND	ND	30.4
105-67-9	2,4-Dimethylphenol	ND	ND	6.0
131-11-3	Dimethyl phthalate	ND	ND	2.0
99-65-0	1,3-Dinitrobenzene	ND	ND	12.4
534-52-1	4,6-Dinitro-o-cresol	ND	ND	10.4
51-28-5	2,4-Dinitrophenol	ND	ND	80.0
121-14-1	2,4-Dinitrotoluene	ND	ND	8.5
606-20-2	2,6-Dinitrotoluene	ND	ND	7.7
	Total dioxin/furan compounds	4.6 E-10	6.3 E-10	
100-41-4	Ethylbenzene	1.0 E-06	1.4 E-06	
74-85-1	Ethylene	3.0 E-04	4.2 E-04	
117-81-7	bis(2-Ethylhexyl)phthalate	ND	ND	2.2
206-44-0	Fluoranthene	ND	ND	9.0 E-01
86-73-7	Fluorene	ND	ND	1.7
76-13-1	Freon 113	ND	ND	7.8 E-04
35822-46-9	1,2,3,4,6,7,8-Heptachlorodibenzo-p- dioxin	8.0 E-11	1.1 E-10	

TABLE A4 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
67562-39-4	1,2,3,4,6,7,8- Heptachlorodibenzofuran	1.8 E-11	2.5 E-11	
55673-89-7	1,2,3,4,7,8,9- Heptachlorodibenzofuran	ND	ND	5.2 E-08
118-74-1	Hexachlorobenzene	ND	ND	2.6
87-68-3	Hexachlorobutadiene	ND	ND	2.2
77-47-4	Hexachlorocyclopentadiene	ND	ND	35.3
39227-28-6	1,2,3,4,7,8-Hexachlorodibenzo-p- dioxin	1.7 E-12	2.3 E-12	
57653-85-7	1,2,3,6,7,8-Hexachlorodibenzo-p- dioxin	1.4 E-11	1.9 E-11	
19408-74-3	1,2,3,7,8,9-Hexachlorodibenzo-p- dioxin	6.3 E-12	8.5 E-12	
70648-26-9	1,2,3,4,7,8-Hexachlorodibenzofuran	8.5 E-12	1.2 E-11	
57117-44-9	1,2,3,6,7,8-Hexachlorodibenzofuran	6.3 E-12	8.5 E-12	
72918-21-9	1,2,3,7,8,9-Hexachlorodibenzofuran	ND	ND	2.4 E-08
60851-34-5	2,3,4,6,7,8-Hexachlorodibenzofuran	5.7 E-12	7.8 E-12	
67-72-1	Hexachloroethane	ND	ND	1.9
110-54-3	Hexane	ND	ND	1.0 E-04
7647-01-0	Hydrochloric acid	0	0	
193-39-5	Indeno[1,2,3-cd]pyrene	ND	ND	1.5
78-59-1	Isophorone	ND	ND	1.7
67-63-0	Isopropyl alcohol	1.6 E-07	2.2 E-07	
120-58-1	Isosafrole	ND	ND	10.1
556-61-6	Isothiocyanatomethane	ND	ND	3.0 E-04
7439-92-1	Lead	1.6 E-05	2.2 E-05	
7439-96-5	Manganese	1.1 E-06	1.4 E-06	
7439-97-6	Mercury	1.2 E-08	1.7 E-08	
56-49-5	3-Methylcholanthrene	ND	ND	2.0
75-09-2	Methylene chloride	2.0 E-06	2.7 E-06	
108-10-1	Methyl isobutyl ketone	ND	ND	4.2 E-04
91-57-6	2-Methylnaphthalene	ND	ND	2.1
95-48-7	2-Methylphenol	ND	ND	2.5
109-06-8	2-Methylpyridine	ND	ND	6.2
1634-04-4	Methyl tert-butyl ether	ND	ND	3.7 E-04
91-20-3	Naphthalene	ND	ND	1.3

TABLE A4 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
130-15-4	1,4-Naphthoquinone	ND	ND	6.4
134-32-7	1-Naphthylamine	ND	ND	7.0
91-59-8	2-Naphthylamine	ND	ND	138.0
100-01-6	4-Nitroaniline	ND	ND	8.4
98-95-3	Nitrobenzene	ND	ND	2.5
88-75-5	2-Nitrophenol	ND	ND	8.0
100-02-7	4-Nitrophenol	ND	ND	74.8
56-57-5	4-Nitroquinoline-1-oxide	ND	ND	43.1
924-16-3	N-Nitrosodibutylamine	ND	ND	7.0
55-18-5	N-Nitrosodiethylamine	ND	ND	9.7
62-75-9	N-Nitrosodimethylamine	ND	ND	8.1
621-64-7	N-Nitrosodipropylamine	ND	ND	4.2
59-89-2	N-Nitrosomorpholine	ND	ND	2.6
100-75-4	N-Nitrosopiperidine	ND	ND	2.4
99-55-8	5-Nitro-o-toluidine	ND	ND	5.4
3268-87-9	1,2,3,4,6,7,8,9-Octachlorodibenzo- p-dioxin ^f	2.2 E-10	3.0 E-10	
39001-02-0	1,2,3,4,6,7,8,9- Octachlorodibenzofuran	4.6 E-11	6.3 E-11	
608-93-5	Pentachlorobenzene	ND	ND	2.2
40321-76-4	1,2,3,7,8-Pentachlorodibenzo-p- dioxin	ND	ND	2.6 E-08
57117-41-6	1,2,3,7,8-Pentachlorodibenzofuran	6.8 E-12	9.3 E-12	
57117-31-4	2,3,4,7,8-Pentachlorodibenzofuran	6.3 E-12	8.5 E-12	
76-01-7	Pentachloroethane	ND	ND	5.0
82-68-8	Pentachloronitrobenzene	ND	ND	12.9
87-86-5	Pentachlorophenol	ND	ND	63.4
85-01-8	Phenanthrene	ND	ND	2.2
108-95-2	Phenol	ND	ND	2.3
7723-14-0	Phosphorus	ND	ND	1.3 E-02
23950-58-5	Pronamide	ND	ND	3.0
123-38-6	Propionaldehyde ^g	7.1 E-07	9.6 E-07	
115-07-1	Propylene ^f	4.3 E-05	5.9 E-05	
129-00-0	Pyrene	ND	ND	2.2
110-86-1	Pyridine	ND	ND	6.0
94-59-7	Safrole	ND	ND	10.1

TABLE A4 (cont.)

		Emission	Factor ^{b,c}	Minimum
CASRN ^a	Compound	lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
7782-49-2	Selenium	8.8 E-09	1.2 E-08	
7440-22-4	Silver	2.3 E-08	3.1 E-08	
100-42-5	Styrene	ND	ND	4.3 E-04
1746-01-6	2,3,7,8-Tetrachlorodibenzo-p-dioxin	ND	ND	2.4 E-08
51207-31-9	2,3,7,8-Tetrachlorodibenzofuran	4.3 E-11	5.8 E-11	
79-34-5	1,1,2,2-Tetrachloroethane	ND	ND	7.0 E-04
127-18-4	Tetrachloroethylene ^g	3.8 E-06	5.2 E-06	
7440-28-0	Thallium	ND	ND	1.0 E-02
108-88-3	Toluene	9.3 E-06	1.3 E-05	
95-53-4	o-Toluidine	ND	ND	4.6
120-82-1	1,2,4-Trichlorobenzene	ND	ND	7.5 E-04
71-55-6	1,1,1-Trichloroethane	ND	ND	5.5 E-04
79-00-5	1,1,2-Trichloroethane	ND	ND	5.5 E-04
79-01-6	Trichloroethylene ^g	6.0 E-08	8.1 E-08	
75-69-4	Trichloromonofluoromethane ^g	0	0	
95-95-4	2,4,5-Trichlorophenol	ND	ND	6.8
88-06-2	2,4,6-Trichlorophenol	ND	ND	5.2
96-18-4	1,2,3-Trichloropropane	ND	ND	6.1 E-04
95-63-6	1,2,4-Trimethylbenzene	ND	ND	5.0 E-04
540-84-1	2,2,4-Trimethylpentane	ND	ND	1.0 E-04
75-01-4	Vinyl chloride ^f	2.7 E-06	3.7 E-06	
75-35-4	Vinylidene chloride	2.4 E-07	3.3 E-07	
106-42-3, 108-38-3	m-Xylene, p-Xylene ^g	1.2 E-06	1.6 E-06	
95-47-6	o-Xylene	9.3 E-07	1.3 E-06	
7440-66-6	Zinc	2.2 E-06	2.9 E-06	
	Other Po	llutants		
64-19-7	Acetic acid	4.0 E-06	5.5 E-06	
67-64-1	Acetone ^f	1.9 E-04	2.6 E-04	
592-20-1	Acetoxyacetone	ND	ND	4.8 E-04
74-86-2	Acetylene ^f	1.6 E-04	2.2 E-04	
100-52-7	Benzaldehyde ^f	7.2 E-06	9.9 E-06	
103-65-1	Benzene, propyl-	ND	ND	1.0 E-04
271-89-6	Benzofuran	2.4 E-06	3.3 E-06	
65-85-0	Benzoic acid	ND	ND	41.5

TABLE A4 (cont.)

	Compound	Emission Factor ^{b,c}		Minimum
CASRN ^a		lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
100-47-0	Benzonitrile ^f	4.4 E-05	6.0 E-05	
100-51-6	Benzyl alcohol	ND	ND	2.1
107-04-0	1-Bromo-2-chloroethane	ND	ND	5.9 E-04
3017-95-6	2-Bromo-1-chloropropane	ND	ND	6.5 E-04
75-28-5	i-Butane ^f	2.1 E-07	2.9 E-07	
106-97-8	n-Butane ^f	1.1 E-06	1.4 E-06	
431-03-8	2,3-Butanedione	1.0 E-05	1.4 E-05	
123-73-9	trans-2-Butenal ^g	7.7 E-06	1.0 E-05	
106-98-9	1-Butene	1.0 E-05	1.4 E-05	
115-11-7	i-Butene	4.4 E-06	6.0 E-06	
590-18-1	cis-2-Butene	1.3 E-06	1.8 E-06	
624-64-6	trans-2-Butene ^f	7.2 E-06	9.9 E-06	
78-94-4	3-Buten-2-one	2.5 E-06	3.4 E-06	
13466-78-9	delta-3-Carene	ND	ND	1.0 E-04
109-69-3	1-Chlorobutane	ND	ND	3.9 E-04
622-98-0	1-Chloro-4-ethylbenzene	ND	ND	5.8 E-04
95-49-8	1-Chloro-2-methylbenzene	ND	ND	5.3 E-04
108-41-8	1-Chloro-3-methylbenzene	ND	ND	5.3 E-04
59-50-7	4-Chloro-3-methylphenol	ND	ND	2.7
95-57-8	2-Chlorophenol	ND	ND	1.7
287-92-3	Cyclopentane	ND	ND	1.0 E-04
96-37-7	Cyclopentane, methyl-	ND	ND	1.0 E-04
120-92-3	Cyclopentanone	ND	ND	3.5 E-04
142-29-0	Cyclopentene	ND	ND	1.0 E-04
930-30-3	2-Cyclopenten-1-one	ND	ND	3.4 E-04
124-18-5	n-Decane	ND	ND	1.0 E-04
3018-12-0	Dichloroacetonitrile	5.7 E-07	7.7 E-07	
616-21-7	1,2-Dichlorobutane	ND	ND	5.3 E-04
32768-54-0	1,2-Dichloro-3-methylbenzene ^g	2.7 E-07	3.7 E-07	
594-37-6	1,2-Dichloro-2-methylpropane	ND	ND	5.3 E-04
87-65-0	2,6-Dichlorophenol	ND	ND	3.8
513-88-2	1,1-Dichloro-2-propanone	ND	ND	5.3 E-04
10061-01-5	cis 1,3-Dichloro-1-propene	ND	ND	4.6 E-04
84-66-2	Diethylphthalate	ND	ND	1.3
75-83-2	2,2-Dimethylbutane	ND	ND	1.0 E-04

TABLE A4 (cont.)

	Compound	Emission Factor ^{b,c}		Minimum
CASRN ^a		lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
79-29-8	2,3-Dimethylbutane	ND	ND	1.0 E-04
624-92-0	Dimethyldisulfide	3.9 E-06	5.3 E-06	
625-86-5	2,5-Dimethylfuran ^g	2.5 E-07	3.5 E-07	
1071-26-7	2,2-Dimethylheptane	ND	ND	1.0 E-04
584-94-1	2,3-Dimethylhexane	ND	ND	1.0 E-04
589-43-5	2,4-Dimethylhexane	ND	ND	1.0 E-04
592-13-2	2,5-Dimethylhexane	ND	ND	1.0 E-04
565-59-3	2,3-Dimethylpentane	ND	ND	1.0 E-04
108-08-7	2,4-Dimethylpentane	ND	ND	1.0 E-04
	Dimethylphenethylamine	ND	ND	119.2
463-82-1	2,2-Dimethylpropane	ND	ND	1.0 E-04
3658-80-8	Dimethyltrisulfide ^g	1.0 E-05	1.4 E-05	
117-84-0	Di-n-octylphthalate	ND	ND	1.9
74-84-0	Ethane	1.1 E-05	1.6 E-05	
64-17-5	Ethanol ^f	9.7 E-07	1.3 E-06	
1678-91-7	Ethylcyclohexane	ND	ND	1.0 E-04
619-99-8	3-Ethylhexane	ND	ND	1.0 E-04
62-50-0	Ethyl methanesulfonate	ND	ND	2.8
620-14-4	m-Ethyltoluene	ND	ND	1.0 E-04
611-14-3	o-Ethyltoluene	ND	ND	1.0 E-04
622-96-8	p-Ethyltoluene	ND	ND	5.0 E-04
98-01-1	2-Furaldehyde	2.4 E-05	3.3 E-05	
498-60-2	3-Furaldehyde	9.8 E-06	1.3 E-05	
110-00-9	Furan	1.4 E-05	1.9 E-05	
109-99-9	Furan, tetrahydro-	ND	ND	3.0 E-04
111-71-7	Heptanal	ND	ND	4.7 E-04
142-82-5	Heptane	ND	ND	1.0 E-04
110-43-0	2-Heptanone	ND	ND	4.7 E-04
1888-71-7	Hexachloropropene	ND	ND	6.8
66-25-1	Hexanal	2.0 E-06	2.7 E-06	
591-78-6	2-Hexanone	ND	ND	4.2 E-04
592-41-6	1-Hexene	7.2 E-06	9.8 E-06	
7688-21-3	cis-2-Hexene	ND	ND	1.0 E-04
4050-45-7	trans-2-Hexene	ND	ND	1.0 E-04
78-79-5	Isoprene	ND	ND	1.0 E-04

TABLE A4 (cont.)

	Compound	Emission Factor ^{b,c}		Minimum
CASRN ^a		lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
143-50-0	Kepone	ND	ND	90.3
5989-27-5	d-Limonene	ND	ND	5.7 E-04
78-85-3	Methacrolein	3.1 E-06	4.3 E-06	
75-52-5	Methane, nitro-	ND	ND	2.5 E-04
91-80-5	Methapyrilene	ND	ND	138.8
104-85-8	4-Methylbenzonitrile	ND	ND	4.9 E-04
563-80-4	3-Methyl-2-butanone	ND	ND	3.6 E-04
563-46-2	2-Methyl-1-butene	1.2 E-06	1.7 E-06	
513-35-9	2-Methyl-2-butene	ND	ND	1.0 E-04
563-45-1	3-Methyl-1-butene ^f	8.6 E-07	1.2 E-06	
108-87-2	Methylcyclohexane	ND	ND	1.0 E-04
78-93-3	Methyl ethyl ketone	9.2 E-06	1.2 E-05	
620-02-0	5-Methyl-2-furaldehyde	ND	ND	4.6 E-04
534-22-5	2-Methylfuran	2.9 E-06	4.0 E-06	
930-27-8	3-Methylfuran	5.0 E-07	6.8 E-07	
592-27-8	2-Methylheptane	ND	ND	1.0 E-04
591-76-4	2-Methylhexane	ND	ND	1.0 E-04
589-34-4	3-Methylhexane	ND	ND	1.0 E-04
66-27-3	Methyl methanesulfonate	ND	ND	3.3
624-91-9	Methylnitrite ^g	3.3 E-07	4.5 E-07	
107-83-5	2-Methylpentane	ND	ND	1.0 E-04
96-14-0	3-Methylpentane	ND	ND	1.0 E-04
763-29-1	2-Methyl-1-pentene	ND	ND	1.0 E-04
625-27-4	2-Methyl-2-pentene	ND	ND	1.0 E-04
691-37-2	4-Methyl-1-pentene	ND	ND	1.0 E-04
691-38-3	cis-4-Methyl-2-pentene	ND	ND	1.0 E-04
554-14-3	2-Methylthiophene	2.8 E-06	3.8 E-06	
616-44-4	3-Methylthiophene	2.4 E-06	3.3 E-06	
88-74-4	2-Nitroaniline	ND	ND	11.9
99-09-2	3-Nitroaniline	ND	ND	4.3
10595-95-6	N-Nitrosomethylethylamine	ND	ND	7.9
930-55-2	N-Nitrosopyrrolidine	ND	ND	8.6
124-19-6	Nonanal	ND	ND	5.9 E-04
111-84-2	n-Nonane	ND	ND	1.0 E-04
821-55-6	2-Nonanone	ND	ND	5.9 E-04

TABLE A4 (cont.)

CASRN ^a	Compound	Emission Factor ^{b,c}		Minimum
		lb per item	lb per lb NEW ^d	Detection Level mg/m ^{3,e}
124-13-0	Octanal	5.8 E-07	8.0 E-07	
111-65-9	n-Octane	ND	ND	1.0 E-04
1600-37-9	1,1,2,3,3-Pentachloro-1-propene	ND	ND	8.9 E-04
110-62-3	Pentanal	ND	ND	3.6 E-04
78-78-4	i-Pentane	ND	ND	1.0 E-04
109-66-0	n-Pentane	ND	ND	1.0 E-04
600-14-6	2,3-Pentanedione	ND	ND	4.2 E-04
107-87-9	2-Pentanone	ND	ND	3.6 E-04
96-22-0	3-Pentanone	ND	ND	3.6 E-04
1629-58-9	1-Penten-3-one	ND	ND	3.5 E-04
109-67-1	1-Pentene	5.0 E-06	6.9 E-06	
627-20-3	cis-2-Pentene	ND	ND	1.0 E-04
646-04-8	trans-2-Pentene	ND	ND	1.0 E-04
107-39-1	1-Pentene, 2,4,4-trimethyl-	ND	ND	1.0 E-04
107-40-4	2-Pentene, 2,4,4-trimethyl-	ND	ND	1.0 E-04
3102-33-8	3-Penten-2-one ^g	4.8 E-07	6.5 E-07	
62-44-2	Phenacetin	ND	ND	2.1
536-74-3	Phenylacetylene	3.7 E-06	5.0 E-06	
80-56-8	alpha-Pinene	ND	ND	1.0 E-04
127-91-3	beta-Pinene	ND	ND	1.0 E-04
74-98-6	Propane	3.3 E-06	4.5 E-06	
95-94-3	1,2,4,5-Tetrachlorbenzene	ND	ND	1.9
58-90-2	2,3,4,6-Tetrachlorophenol	ND	ND	11.6
110-02-1	Thiophene ^f	4.9 E-05	6.7 E-05	
98-03-3	2-Thiophenecarboxaldehyde ^g	2.5 E-06	3.4 E-06	
545-06-2	Trichloroacetonitrile	ND	ND	6.0 E-04
108-67-8	1,3,5-Trimethylbenzene	ND	ND	5.0 E-04
16747-26-5	2,2,4-Trimethylhexane	ND	ND	1.0 E-04
565-75-3	2,3,4-Trimethylpentane	ND	ND	1.0 E-04
99-35-4	sym-Trinitrobenzene	ND	ND	8.8

TABLE A4 (cont.)

TABLE A4 (cont.)

- ^a CASRN = Chemical Abstracts Service Registry Number.
- ^b ND = nondetected.
- ^c Emission factors rated C unless otherwise noted.
- ^d NEW = net explosive weight. The NEW for this ordnance is 7.20 E-01 pounds per item.
- ^e Data provided for compounds that were not detected. ^f Emission factor rated B because of correlation with emission factors for similar ordnance and number of test data points
- ^g Emission factor rated D because the factor is based upon C-rated test data.

APPENDIX B

NEW AP-42 SECTIONS FOR ORDNANCE INCLUDED IN PHASE III TESTING AT DUGWAY PROVING GROUND, UTAH

Electronic versions of the new AP-42 sections for ordnance included in Phase III testing at Dugway Proving Ground, Utah, are located on the EPA website at: <u>http://www.epa.gov/ttn/chief/ap42/index.html</u>.

This page left blank intentionally.