




STEELHEAD FISH HATCHERY EVALUATIONS—IDAHO

Project Progress Report



T. Dean Rhine Senior Fisheries Research Biologist

> Randall S. Osborne Senior Fisheries Technician

Kristy A. Stevens Fisheries Biological-Aide

IDFG Report Number 99-18 April 1999

Steelhead Fish Hatchery Evaluations—Idaho

Period Covered: October 1, 1992 to September 30, 1993

Ву

T. Dean Rhine Randall S. Osborne Kristy A. Stevens

Idaho Department of Fish and Game 600 South Walnut Street P.O. Box 25 Boise, ID 83707

То

U.S. Fish and Wildlife Service Lower Snake River Compensation Plan Office 1387 S. Vinnell Way, Suite 343 Boise, ID 83709

Contract Number 14-48-0001-93500

IDFG Report Number 99-18 April 1999

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT	1
INTRODUCTION	2
OBJECTIVES	2
Objective 1	
Sub-objective 1.1	
Objective 2.	
Sub-objective 2.1	
Sub-objective 2.2	3
METHODS	3
IDFG LSRCP Program Success Documentation—Objective 1	3
Hatchery Operations Documentation—Sub-objective 1.1	3
Migration Conditions	
Migration Timing and Juvenile Survival	
Adult Returns	
Fisheries Contribution	
Hatchery Weirs	
Smolt-to-Adult Return Rate	
Experimentation—Objective 2	
Clearwater Fish Hatchery	
Cover Experiment	
Hagerman National Fish Hatchery	
Size-at-Release Experiment	
Acclimation Experiment	0
RESULTS	6
Haller Oracle Decreased from O. L. H. C. and A.	•
Hatchery Operations Documentation - Sub-objective 1.1	
Clearwater Fish HatcheryBrood Year 1992	6
Brood Year 1992 Brood Year 1993	•
Hagerman National Fish Hatchery	
Brood Year 1992	
Brood Year 1993	
Magic Valley Fish Hatchery	
Brood Year 1992	
Brood Year 1993	
Migration Conditions	
Migration Timing and Juvenile Survival	
Adult Returns	
Fisheries Contribution	
Hatchery Weirs	
Sawtooth Fish Hatchery Weir	
East Fork Salmon River Weir	

(Table of Contents, Continued.)

		<u>Page</u>
Clea Hage Magi Experir Clea Co Hage Siz	ro-Adult Return Rates rwater Fish Hatchery erman National Fish Hatchery c Valley Fish Hatchery mentation - Objective 2 rwater Fish Hatchery over Experiment erman National Fish Hatchery ze-at-Release Experiment climation Experiment	10 12 12 12 13
LITERAT	URE CITED	13
APPEND	ICES	22
	LIST OF TABLES	
	LIGITOT TABLES	
Γable 1.	Number of unique PIT tag interrogations of LSRCP steelhead smolts, by PIT tag file, at Lower Granite Dam (GRJ), Little Goose Dam (GOJ), Lower Monumental Dam (LMJ), and McNary Dam (MCJ) for the 1993 migration period. A total of 2,200 PIT-tagged steelhead were released from Hagerman National, Magic Valley, and Clearwater fish hatcheries between April 8 and April 21, 1993. Median travel time is to Lower Granite Dam.	14
Γable 2.	Snake River mean daily in-flow (thousand cubic feet per second) at Lower Granite Dam, Washington, from 1977-1993 during the Peak and Extended chinook salmon smolt migration periods as defined by Petrosky (1991)	16
Γable 3.	Estimated number of LSRCP steelhead that returned to Idaho in 1992-1993. The adult return in 1992-1993 included fish from three age classes. Steelhead were reared at Hagerman National and Magic Valley fish hatcheries. These estimates were prepared by the Idaho Department of Fish and Game's Harvest Monitoring Project and only include steelhead harvested in Idaho's sport fisheries, steelhead that returned to hatchery racks, and steelhead that escaped to spawn naturally. These are minimum estimates and do not include all tributary and mainstem strays or in-river pre-spawning mortalities.	16
Γable 4.	Steelhead smolts released from Magic Valley and Hagerman National fish hatcheries that contributed to the 1992-1993 Idaho steelhead return. The number of steelhead smolts released and the estimated number of adults that returned were compared to facility design production targets and projected adult return goals.	17

(List of Tables, Continued.)

		<u>Page</u>
Table 5.	Summary of the 1993 A-strain steelhead trout return to the Sawtooth Fish Hatchery weir. The fish return included fish of hatchery and natural origin. Hatchery aging criteria, based on length, were used to determine age ^a . ND indicates and data" (i.e., data were not available)	21
Table 6.	Summary of the 1993 B-strain steelhead trout return to the East Fork Salmon River weir. The fish return included fish of hatchery and natural origin. Hatchery aging criteria, based on length, were used to determine age ^a . ND indicates "no data" (i.e., data were not available).	22
Table 7.	Smolt-to-adult return rates of coded-wire tagged steelhead smolts released from Hagerman National and Magic Valley fish hatcheries. The number of adults was estimated by Ball (1996) and only include steelhead harvested in Idaho's sport fisheries, steelhead that returned to hatchery racks, and steelhead that escaped to spawn naturally. These are minimum estimates and do not include all tributary and mainstem strays or in-river pre-spawning mortalities.	23
Table 8.	Total number ^a of steelhead recovered with coded-wire tags designating them as either large size ^b or regular size ^c . Recovered fish were released as smolts at the Sawtooth Fish Hatchery weir in 1991 (brood year 1990). ND indicates "no data" (i.e., data were not available).	20
	LIST OF FIGURES	
Figure 1.	Number of unique PIT tag interrogations of LSRCP juvenile steelhead at Lower Granite Dam, Washington, plotted with the average daily in-flow of the Snake River at Lower Granite Dam in 1993. A total of 2,200 PIT-tagged steelhead smolts were released from Hagerman National, Magic Valley, and Clearwater fish hatcheries between April 8 and April 21, 1993. Fifty-three percent (1,160) of the PIT-tagged fish were interrogated at Lower Granite Dam. Flow data were not available for all dates. Data for one fish falls outside of this date range and is not shown.	24

LIST OF APPENDICES

			<u>Page</u>
Appendix A.	Table 1.	Release data and estimated adult returns for Clearwater Fish Hatchery summer steelhead trout, brood year 1992. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.	26
Appendix A.	Table 2.	Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1992. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.	28
Appendix A.	Table 3.	Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1992. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.	32

(List of Appendices, Continued.)

<u>Page</u>

Appendix B. Table 1.	Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1988. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.	36
Appendix B. Table 2.	Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1989. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.	40
Appendix B. Table 3.	Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1990. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.	43

(List of Appendices, Continued.)

<u>Page</u>

Appendix C. Table 1.	Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1988. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.	46
Appendix C. Table 2.	Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1989. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.	49
Appendix C. Table 3.	Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1990. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.	52

ABSTRACT

This annual report summarizes activities associated with Idaho Lower Snake River Compensation Plan (LSRCP) hatcheries' activities from October 1, 1992 to September 30, 1993. Included in this report are all fall 1992 and spring 1993 adult steelhead trout *Oncorhynchus mykiss* returns and all releases of juvenile steelhead trout made within the reporting period. Information presented in this report supersedes that included in previous reports.

An estimated 11,594 LSRCP steelhead returned to Idaho during this reporting period. Magic Valley Fish Hatchery produced an estimated 6,005 adult steelhead, and Hagerman National Fish Hatchery produced an estimated 5,589 adult steelhead. Steelhead returns to Idaho remain well below the LSRCP program mitigation goal of 39,260 fish.

Adult returns to Idaho LSRCP hatchery racks included 1,591 A-strain steelhead to Sawtooth Fish Hatchery and 176 B-strain steelhead to East Fork Salmon River satellite facility. Approximately 73% of the steelhead that returned to Sawtooth Fish Hatchery were males, as compared to 56.3% of the fish that returned to the East Fork Salmon River satellite.

Coded-wire tags were used to determine smolt-to-adult return rates. Dworshak B-stock steelhead, brood year 1988, reared at Hagerman National Fish Hatchery and released into the East Fork Salmon River had a smolt-to-adult return rate of 0.05%. Sawtooth A-stock steelhead, brood year 1989, reared at Hagerman National Fish Hatchery and released at Sawtooth Fish Hatchery and at the Shoup Bridge (located on the Salmon River) had smolt-to-adult return rates of 0.64% and 0.31%, respectively. East Fork B-stock steelhead, brood year 1988, reared at Magic Valley Fish Hatchery and released into the East Fork Salmon River had a smolt-to-adult return rate of 0.10%. Pahsimeroi A-stock steelhead, brood year 1989, reared at Magic Valley Fish Hatchery and released at the Sawtooth Fish Hatchery had a smolt-to-adult return rate of 0.31%.

In April 1993, Idaho LSRCP hatcheries released 3,739,842 brood year 1992 steelhead smolts. A total of 599,678 of these fish were tagged with coded-wire tags. In addition, 2,200 steelhead smolts were released with passive integrated transponder (PIT) tags.

In September 1993, Clearwater Fish Hatchery released 50,027 brood year 1993 steelhead fingerlings into the South Fork Clearwater River for Idaho's Supplementation Studies program. Five thousand of the fingerlings were tagged with PIT tags.

INTRODUCTION

The Water Resources Development Act of 1976 (90 Stat. 2917) authorized the Lower Snake River Compensation Plan (LSRCP) to mitigate for fish losses caused by the construction and operation of Ice Harbor, Lower Monumental, Little Goose, and Lower Granite dams on the lower Snake River. Mitigation for anadromous fishery losses included the construction and operation of fish hatchery facilities and smolt passage improvements at the lower Snake River dams. The United States Fish and Wildlife Service was authorized to administer the operation and maintenance for 12 hatcheries and 11 satellite facilities in Idaho, Oregon, and Washington.

In Idaho, the Idaho Department of Fish and Game (IDFG) operates Clearwater Fish Hatchery, McCall Fish Hatchery, Magic Valley Fish Hatchery, Sawtooth Fish Hatchery, South Fork Salmon River Trap, East Fork Salmon River Trap, and Red River, Crooked River, and Powell satellite facilities. The United States Fish and Wildlife Service operates Dworshak National Fish Hatchery and Hagerman National Fish Hatchery. Adult return goals for the entire LSRCP are 8,000 summer chinook salmon *Oncorhynchus tshawytscha*, 50,700 spring chinook salmon, 18,300 fall chinook salmon, and 55,100 steelhead trout *O. mykiss* to the Snake River basin. Adult return goals for the Idaho portion of the LSRCP program call for the return of 8,000 adult summer chinook salmon, 40,432 adult spring chinook salmon, and 39,260 adult steelhead. Adult return goals for Idaho LSRCP steelhead hatcheries are as follows: Clearwater Fish Hatchery—14,000, Hagerman National Fish Hatchery—13,600, and Magic Valley Fish Hatchery—11,660.

The LSRCP program includes a Hatchery Evaluation Study component to monitor and evaluate the mitigation hatchery program. The primary objective of the Hatchery Evaluation Study is to determine the best hatchery management practices for mitigation hatcheries to meet LSRCP and IDFG anadromous fisheries goals. Only if we understand the effects of hatchery operations on adult return characteristics (e.g., return rates, sex ratios, age structure) can we prescribe effective management actions. Tasks defined to satisfy the primary objective are divided into two categories: 1) documentation, and 2) investigation. We document hatchery practices for each brood year, or cohort, of fish and mitigation status in terms of annual adult returns. Our success at achieving LSRCP and IDFG goals can then be related to hatchery practices through the documentation tasks. Investigation tasks are manipulative experiments involving modified or alternative hatchery practices that show potential for increasing adult returns and achieving LSRCP and IDFG goals.

OBJECTIVES

This report summarizes steelhead Hatchery Evaluation Study activities carried out from October 1, 1992 through September 30, 1993. Juvenile steelhead released from Clearwater, Hagerman National, and Magic Valley fish hatcheries during this reporting period are documented, as well as adult steelhead that returned to Idaho during the fall of 1992 and the spring of 1993 (hereafter referred to as the 1992-1993 return). Specific objectives identified in Cooperative Work Agreement 14-48-0001-93500 are covered in this report and are as follows.

Objective 1. Document the success of the IDFG LSRCP program in meeting specific adult return goals.

- **Sub-objective 1.1** Document LSRCP fish rearing and release practices and adult returns in Idaho.
- **Objective 2.** Identify factors limiting hatchery success and recommend possible improvements based on existing knowledge and experimentation.
 - **Sub-objective 2.1** Continue ongoing documentation and monitoring to determine the relationships between adult returns and hatchery practices, characteristics of hatchery products, and juvenile survival.
 - **Sub-objective 2.2** Conduct controlled studies (short-term experiments) to determine the relationships between adult returns and hatchery practices, characteristics of hatchery products, and juvenile survival.

The results of experiments performed under Sub-objective 2.2 are printed separately from this report. Some results from those experiments, such as juvenile migration characteristics and adult return rates for experimental groups, are included in this report.

METHODS

IDFG LSRCP Program Success Documentation—Objective 1

To document the overall success of the program, we compared the estimated number of adult steelhead that returned above Lower Granite Dam between October 1, 1993 and September 30, 1994 to the Idaho LSRCP goal of 25,260 adult steelhead. The 39,260 fish goal was not used because 14,000 of these fish were charged to Clearwater Fish Hatchery, which only became operational in 1992. Adults from the first smolt release from Clearwater Fish Hatchery will return in 1995, 1996, and 1997. The Harvest Monitoring Project estimated the total number of returning adults and partitioned the total return between Hagerman National Fish Hatchery and Magic Valley Fish Hatchery based on coded-wire tag (CWT) data. Results for Objective 1 are reported under *Results, Adult Returns*.

Hatchery Operations Documentation—Sub-objective 1.1

Hatchery operations between October 1, 1992 and September 30, 1993 are documented in this report. Pertinent rearing information affecting brood years 1992 and 1993 are discussed. Additional information which occurred before this reporting period may be included for brood year 1992 steelhead for completeness. Information was collected from Hatchery Brood Year and Run reports and memoranda and from verbal communications with hatchery personnel. Fish marking and tagging information was provided by IDFG's Coded-Wire Tag Laboratory.

Migration Conditions

Snake River discharge during smolt migration is a major factor affecting survival of Idaho's anadromous fishes. Flow conditions at Lower Granite Dam, Washington, for the 1993 emigration period, the year that brood year 1992 steelhead emigrated, are reported. Adults that returned during this reporting period were from the 1988, 1989, or 1990 broods (depending on the stock and age-at-return). Flow conditions for 1989, 1990, and 1991 are reported since steelhead smolts are reared on a one-year program and released the following spring. Water flow data were obtained from Fish Passage Center reports and the United States Geological Survey Internet site.

Petrosky (1991) defined two time periods that accounted for most of the chinook emigration past Lower Granite Dam. The Extended period runs from April 20 to May 30 and includes the time when most of the wild and natural yearling chinook salmon emigrate, whereas the Peak period runs from April 15 to May 5 and encompasses the time when approximately 50% of the yearling chinook salmon emigrate past the dam. Hatchery steelhead smolts are generally released in April and emigrate during the same time period as chinook. Therefore, flows during the extended and peak time periods are reported.

Migration Timing and Juvenile Survival

Passive Integrated Transponder (PIT) tags were used to evaluate downstream juvenile emigration. The interrogation rate of PIT-tagged juvenile salmonids at Columbia and Snake river dams serves as a minimum survival index because: 1) an unknown (but we believe small) number of PIT-tagged fish that die in the hatchery may go undetected, although we scan the dead fish, 2) not all fish pass through detectors, 3) some PIT tags fail (approximately 2%, Russell Kiefer, IDFG, personal communication) or are lost between tagging and arrival at detection sites, 4) some fish arrive while detection gear is not being operated, and 5) mortality occurs between dams.

Steelhead juveniles were PIT-tagged by IDFG Fish Marking personnel. The PIT tag data were submitted to PTAGIS, a computerized PIT tag database operated by Pacific States Marine Fisheries Commission (Columbia River Basin PIT Tag Information System 1997). Interrogation rates and median travel times for specific PIT-tagged groups of steelhead were calculated after retrieving relevant interrogation data from PTAGIS. Interrogation rates were calculated for each PIT tag file (or files depending on the purpose of the tagging), by dividing the number of unique interrogations at Lower Granite, Little Goose, Lower Monumental, and McNary dams by the number of PIT-tagged fish released, multiplied by 100. Median travel times were calculated for each PIT tag file or files to Lower Granite Dam.

Adult Returns

Adult return goals for Clearwater Fish Hatchery, Hagerman National Fish Hatchery, and Magic Valley Fish Hatchery are 14,000, 13,600, 11,660 adult steelhead above Lower Granite Dam, respectively. The Harvest Monitoring Project (i.e., Ball 1996) estimated the total number of LSRCP steelhead that returned to Idaho in the 1992-1993 return. This estimate included LSRCP-reared fish that were harvested in Idaho's sport fishery and LSRCP-reared fish that escaped to spawn naturally or returned to hatchery racks. Ball's (1996) estimate should be considered a minimum estimate, because tributary and mainstem strays were not accounted

for, nor were in-river pre-spawning mortalities. The number of steelhead smolts released and the estimated number of adults that returned were compared to facility design production targets and projected adult return goals for each facility.

Fisheries Contribution

Idaho Department of Fish and Game Fish Marking personnel tagged juvenile steelhead with coded-wire tags according to marking/tagging plans developed by fishery managers and research biologists. Steelhead tagged with a CWT had the left pelvic fin excised to indicate the presence of a tag. The snouts from tagged adult steelhead harvested in Idaho's sport fishery were sent to the Coded-Wire Tag Laboratory and processed. The Harvest Monitoring Project used these data, along with data from a statewide telephone survey, to estimate the total number of steelhead harvested in Idaho. The numbers of steelhead harvested from specific release groups were estimated by expanding CWT recoveries for specific groups. See Ball (1996) for methods.

Hatchery Weirs

The numbers of steelhead that returned to the East Fork Salmon River and Sawtooth Fish Hatchery weirs were documented by hatchery personnel. The length, sex, and disposition of each fish were recorded. Fish length and strain (A or B) were used to determine age-at-return. Snouts from coded-wire-tagged steelhead were sent to the Coded-Wire Tag Laboratory and processed. The Harvest Monitoring Project used these data to estimate the total number of LSRCP-reared steelhead that returned to hatchery racks or escaped to spawn naturally.

Smolt-to-Adult Return Rate

The Harvest Monitoring Project estimated the total number of LSRCP-produced steelhead that returned to Idaho. For each CWT code, we summed the estimated number of steelhead that returned to Idaho in the 1990-1991 (Ball 1992), 1991-1992 (Ball 1994), and the 1992-1993 (Ball 1996) harvest seasons. For specific groups of fish (i.e., individual CWT codes by release site), we calculated a smolt-to-adult return rate by dividing the estimated number of adults that returned (provided by the Harvest Monitoring Project) by the number of smolts released (provided by the Coded-Wire Tag Laboratory), multiplied by 100.

Experimentation—Objective 2

Interim progress reports, printed independent of this report, document the current status of Hatchery Evaluation Study experiments. Results for some experiments, particularly adult return rates and migration rates for experimental groups, are included in this report.

Clearwater Fish Hatchery

<u>Cover Experiment</u>—The purpose of this experiment was to test the effects of shade covers on adult return and juvenile emigration rates. The experiment was conducted using Dworshak B-stock steelhead from the 1992 brood. Floating shade structures made from PVC

pipe were placed into raceways 9 and 10 (Cover Group). Raceway 11 was not equipped with shade structures (No Cover Group). Steelhead from each group were tagged with coded-wire tags and PIT tags (Appendix A. Table 1). Adults will return between 1995 and 1997. The total number of adults that return from each group will be tested using chi-square analysis (α = 0.05) (SYSTAT Inc. 1992). PIT tags will be used to determine juvenile interrogation rates at Snake and Columbia river dams and median travel to Lower Granite Dam for each group. Chi-square analysis (α = 0.05) will be used to test for PIT tag interrogation rate differences between groups. The Mann-Whitney test (α = 0.05) (SYSTAT Inc. 1992) will be used to test for travel time (to Lower Granite Dam) differences between groups.

Hagerman National Fish Hatchery

<u>Size-at-Release Experiment</u>—The purpose of this study was to determine the optimal size (length) to rear steelhead juveniles at Hagerman National Fish Hatchery. The experiment was conducted for two consecutive years using A-strain steelhead from brood years 1990 and 1991. For each year of the experiment, steelhead were programmed by feeding regime to reach two different sizes: large—2.5 to 3.0 fish per pound (fpp) (256-241 mm fork length) and regular—4.0 to 4.5 fpp (219-210 mm fork length). The CWT and PIT tag data will be used to test dependent variables associated with returning adults and migrating smolts, respectively. See Cannamela (1992) for complete details of the experimental design.

Acclimation Experiment—This study compared steelhead that were trucked from Hagerman National Fish Hatchery and acclimated at Sawtooth Fish Hatchery for two weeks (Acclimated Group) to steelhead that were trucked from Hagerman National Fish Hatchery two weeks later and released directly into the Salmon River (Non-acclimated Group). The two groups of fish were released at the same time and at the same location. The study was initiated in 1992 (1991 brood) and continued in 1993 (1992 brood). Pahsimeroi A-stock steelhead were used for both years of the study. Adults from brood year 1991 will return in 1994 and 1995; adult returns from the 1992 brood will be complete in 1996. Adult return data and juvenile emigration data will be analyzed by brood year. The total number of adults that return from each group will be tested using chi-square analysis ($\alpha = 0.05$). The PIT tag data will be used to determine unique interrogation rates at Snake and Columbia river dams and median travel to Lower Granite Dam for each group. Chi-square analysis ($\alpha = 0.05$) will be used to test interrogation rates between groups. Travel times for the two groups will be tested for differences using the Mann-Whitney test ($\alpha = 0.05$) (SYSTAT Inc. 1992).

RESULTS

Hatchery Operations Documentation - Sub-objective 1.1

Clearwater Fish Hatchery

<u>Brood Year 1992</u>—Clearwater Fish Hatchery received 397,000 eyed eggs, Dworshak B-stock, from Dworshak National Fish Hatchery in May, 1992 (George et al. 1994). Eggs were collected in April at Dworshak National Fish Hatchery during the Number 11, 12, and 13 egg-takes. Survival to the fry stage was 93.6% (371,688 fish) and survival to release was 82.2%

(326,300). Adipose fins were excised from all fish in September and October 1992. Fish were moved to the chinook raceways in October 1992 as a result of a water leak in the main pipeline.

In January 1993, 69,323 fish were tagged with coded-wire tags and marked by excising the left pelvic fin. Fish were tagged with coded-wire tags to estimate adult contribution to the fishery and to test the effects of shade structures placed in raceways 9 and 10. In addition, 300 steelhead were tagged with PIT tags in January 1993. Only steelhead without coded-wire tags were tagged with PIT tags.

Clearwater Fish Hatchery released 326,300 steelhead smolts into the South Fork Clearwater River from April 12-14, 1993 (Appendix A. Table 1). There were 69,016 fish released with coded-wire tags and 300 fish released with PIT tags. See Results, Experimentation - Objective 2, Clearwater Fish Hatchery, Cover Experiment for PIT tag interrogation information.

It was discovered that all fish were inadvertently exposed to nitrogen supersaturated water just before being released. The water supply line used to fill the transport trucks at Clearwater Fish Hatchery was not equipped with a de-gassing station. Therefore, all fish were exposed to supersaturated water when they were transported to the release site. This problem was not discovered until fish released from Clearwater Fish Hatchery were later captured downstream at the Clearwater Smolt Monitoring Trap (Lewiston, Idaho) by IDFG personnel who reported that large numbers of fish had cataracts in one or both eyes. Clearwater Fish Hatchery personnel estimated that 33%-75% of the fish trapped in the smolt trap showed symptoms of gas bubble disease.

<u>Brood Year 1993</u>—Idaho Department of Fish and Game Fishery Management personnel in the Clearwater Region initiated a new program at Clearwater Fish Hatchery in 1993 that utilized wild Selway B-stock steelhead as the brood source (George et al. 1995). The purpose of the program was to develop a separate hatchery brood stock that would return to Crooked River. It was believed that Selway B-stock steelhead would return at a higher rate than Dworshak B-stock steelhead and would provide more harvest opportunities for anglers. Contact Fishery Management personnel in the Clearwater Region for program details.

Forty-two wild adult Selway B-stock steelhead were captured at Selway Falls between March 11 and April 2, 1993. Fish captured included 14 males (four one-ocean and 10 two-ocean) and 28 females (five one-ocean and 23 two-ocean). Fish were transported to Kamiah ponds for holding and were spawned between April 8 and May 5. Eggs from each female were separated into two to four buckets, and each bucket of eggs was fertilized with the milt from a different male. Eggs were disinfected and incubated at Clearwater Fish Hatchery. Ovarian samples were collected from each female and sent to the Eagle Fish Health Laboratory for infectious hematopoietic necrosis (IHN) virus testing. Test results for all samples were negative.

A total of 136,476 green eggs, Selway B-stock, were collected during spawning operations at the Kamiah ponds. Of these, 93.2% (127,162) developed to the eyed stage. Survival to the fry stage was 86.8% (118,515). All Selway B-stock steelhead had the right pelvic fin excised in September 1993.

Clearwater Fish Hatchery received 869,900 eyed eggs, Dworshak B-stock, from Dworshak National Fish Hatchery in April and May of 1993 (George et al. 1995). Eggs were collected at Dworshak National Fish Hatchery in April during the Number 10 through 13 egg-takes. Survival to the fry stage was 88.5% (769,862 fish).

A total of 50,027 Dworshak B-stock steelhead were released into the South Fork Red River as fingerlings in September for supplementation. These fish were marked by excising the right pelvic fin (i.e., right ventral fin-clip), and 5,000 were PIT-tagged. Steelhead used for supplementation were not adipose fin-clipped. Contact IDFG's Idaho Supplementation Studies (Alan Byrne) for program details and PIT tag interrogation results.

All Dworshak B-stock steelhead, except those used for supplementation, were adipose fin-clipped in August 1993. Also in August, a total of 200,051 steelhead were tagged with coded-wire tags and 784 fish were tagged with PIT tags.

Hagerman National Fish Hatchery

Brood Year 1992—Hagerman National Fish Hatchery received a total of 2,332,710 eyed eggs comprised of two different stocks of steelhead: Sawtooth A-stock (1,256,701 eggs) and Pahsimeroi A-stock (1,076,009 eggs) (Hagerman National Fish Hatchery 1992). Survival rates from eyed eggs to the fry stage for the Sawtooth A-stock and the Pahsimeroi A-stock were 97% (1,219,000 fish) and 97.6% (1,050,574 fish), respectively. A total of 402,383 steelhead fingerlings were transferred to Lynn Babington Fish Hatchery in September and October 1992. These steelhead were being reared at Hagerman National Fish Hatchery to assist Niagara Springs Fish Hatchery. Excluding the fish transferred to Lynn Babington Fish Hatchery, survival rates from eyed eggs to smolt release for Sawtooth A-stock and Pahsimeroi A-stock were 78% (898,948 fish) and 76% (588,894 fish), respectively. No major health problems were reported.

Adipose fins were excised from all fish in September and October of 1992. In November 1992, 228,499 steelhead were tagged with coded-wire tags and marked by excising the left pelvic fin (Appendix A. Table 2). Fish were tagged with coded-wire tags to estimate adult contribution to the fishery and to test the effects of acclimating smolts at Sawtooth Fish Hatchery. In February 1993, 600 steelhead were tagged with PIT tags as part of the Sawtooth Acclimation Experiment to test emigration timing and interrogation rates for acclimated and non-acclimated steelhead.

Hagerman National Fish Hatchery released 1,487,842 steelhead smolts into the Salmon and Little Salmon rivers between April 8 and April 28, 1993 (Appendix A. Table 2). The total release included 898,948 Sawtooth A-stock and 588,894 Pahsimeroi A-stock steelhead. Fish were released at three locations: Sawtooth Fish Hatchery weir (729,520), Hammer Creek (211,006), and Hazard Creek (547,316). There were 228,499 fish released with coded-wire tags and 600 fish released with PIT tags. See Results, Experimentation - Objective 2, Hagerman National Fish Hatchery, Acclimation Experiment for PIT tag results.

<u>Brood Year 1993</u>—Hagerman National Fish Hatchery received a total of 2,019,973 eyed steelhead eggs in May and June 1993 comprised of two different stocks: Sawtooth A-stock (1,014,960 eggs) and Pahsimeroi A-stock (1,005,013 eggs) (Hagerman National Fish Hatchery 1993). Survival rates from the eyed egg stage to the fry stage for the Sawtooth A-stock and Pahsimeroi A-stock were 96.4 and 95.6%, respectively.

Magic Valley Fish Hatchery

<u>Brood Year 1992</u>—Magic Valley Fish Hatchery received a total of 2,487,548 eyed eggs comprised of three different stocks of steelhead: Dworshak B-stock (1,322,740 eggs), Pahsimeroi A-stock (1,031,274 eggs), and East Fork B-stock (133,826 eggs) (Ainsworth et al. 1993). Overall survival to the fry stage was 86.1% (2,140,861 fish) and survival to release was 77.4% (1,925,700 fish).

In November 1992, all steelhead were marked with adipose fin clips and 318,231 fish were tagged with coded-wire tags and marked by excising the left pelvic fin. Coded-wire tags were used to determine fishery contribution. In December 1992, flexibacteria was detected during a monthly health exam; however, no treatment was administered due to the low incidence of infection. In February 1993, 1,300 steelhead not tagged with coded-wire tags were tagged with PIT tags.

Magic Valley Fish Hatchery released a total of 1,925,700 brood year 1992 steelhead at seven different locations between April 7 and April 22, 1993 (Appendix A. Table 3). The total release included 903,400 Dworshak B-stock, 915,900 Pahsimeroi A-stock, and 106,400 East Fork B-stock steelhead. A total of 302,163 fish were released with coded-wire tags and marked by excising the left pelvic fin. In addition, 1,300 steelhead were released with PIT tags (Appendix A. Table 3). Brood year 1992 fish had no major disease outbreaks and hatchery personnel reported that the fish were in excellent condition.

Interrogation rates for groups of PIT-tagged fish ranged from 51.0 to 78.0% (Table 1). Median travel times to Lower Granite Dam for groups of PIT-tagged fish ranged from 15.5 to 38.0 days (Table 1).

Brood Year 1993—Magic Valley Fish Hatchery received a total of 2,767,613 eyed steelhead eggs comprised of three different stocks: Dworshak B-stock (1,507,033 eggs), Pahsimeroi A-stock (1,081,500 eggs), and East Fork B-stock (179,080 eggs) (Ainsworth et al. 1995). Survival rates from the eyed egg stage to the fry stage were as follows: Dworshak B-stock—97.1%, Pahsimeroi A-stock—99.3%, and East Fork B-stock—99.4%. In July 1993, 392,300 surplus fry, Dworshak B-stock, were shipped to Salmon Falls (227,600 fry) and Oakley (164,700 fry) reservoirs.

Migration Conditions

Snake River in-flows (mean in-flow) at Lower Granite Dam during the peak period in 1993 were the largest since 1989 (Table 2). Moreover, the flows during the extended period were the largest since 1984.

Comparing Snake River flow conditions for the three brood years of steelhead that returned to Idaho as adults during this reporting period (broods 1988, 1989, and 1990), the 1988 brood (which emigrated in 1989) had the largest river discharge during both the peak and extended migration periods (Table 2). The 1990 brood (which emigrated in 1991) had the lowest river discharge during the peak migration period.

Migration Timing and Juvenile Survival

A total of 2,200 steelhead smolts were released with PIT tags in 1993: Clearwater Fish Hatchery—300, Hagerman National Fish Hatchery—600, and Magic Valley Fish Hatchery—1,300 (Table 1). Overall, 65% (1,429) of the fish PIT-tagged were interrogated at downstream dams. Interrogation rates of PIT-tagged steelhead, by PIT tag file, ranged from 51.0 to 78.0% (Table 1). Median travel times to Lower Granite Dam for PIT-tagged steelhead, by PIT tag file, ranged from 15.5 to 38.0 days (Table 1). Most of the steelhead tagged with PIT tags were interrogated at Lower Granite Dam between April 20 and May 30 (Figure 1). Flow conditions for the Snake River at Lower Granite Dam during this time period ranged from approximately 55 to 180 thousand cubic feet per second.

Adult Returns

The Harvest Monitoring Project (Ball 1996) estimated that Magic Valley Fish Hatchery and Hagerman National Fish Hatchery returned 11,594 steelhead to Idaho in 1992-1993 (Table 3). Ball (1996) estimated that 8,747 steelhead were harvested in Idaho's sport fishery, and 2,847 steelhead returned to hatchery racks or escaped to spawn naturally. These estimates do not include tributary and mainstem strays or pre-spawning mortalities. The 1992-1993 steelhead return included 1-ocean, 2-ocean, and 3-ocean fish. The number of steelhead smolts released and the estimated number of adults that returned are compared to facility design production targets and projected adult return goals in Table 4. Hagerman National Fish Hatchery and Magic Valley Fish Hatchery achieved a minimum of 46% of their combined adult return goals (Table 4). Adult return estimates include only steelhead that returned to hatchery weirs, steelhead that were harvested in Idaho's sport fishery, and steelhead that escaped to spawn naturally.

Fisheries Contribution

Ball (1996) estimated that 8,747 LSRCP-reared hatchery steelhead were harvested during the 1992-1993 Idaho sport fishing season. See Ball (1996) for creel survey methods and results.

Hatchery Weirs

<u>Sawtooth Fish Hatchery Weir</u>—Hatchery steelhead returning to the Sawtooth Fish Hatchery in 1993 were A-strain fish released in 1990 and 1991 (brood years 1989 and 1990, respectively). Smolts were reared at Hagerman National and Magic Valley fish hatcheries prior to being trucked to Sawtooth Fish Hatchery and released.

A total of 1,591 adult steelhead (A-strain), comprised of 1,154 males (72.5%) and 437 females (27.5%), returned to the Sawtooth Fish Hatchery weir between March 18 and May 12, 1993 (Table 5) (Chapman 1993). The male component of the run was comprised of 1,151 hatchery origin fish and three natural origin (unmarked) fish; the female component was made up of 433 hatchery origin fish and four natural origin fish. All of the natural origin fish and 661 of the hatchery origin fish (487 males and 174 females) were released to spawn naturally. A total of 255 females and 530 males were spawned on 11 different dates to yield 1,131,877 green

eggs. A total of 1,031,635 eggs (91.1%) developed to the eyed stage. Disease samples collected during the spawning operations tested negative for IHN virus and bacteria.

<u>East Fork Salmon River Weir</u>—Hatchery steelhead returning to the East Fork Salmon River weir in 1993 were B-strain fish released in 1989, 1990, and 1991 (brood years 1988, 1989, and 1990, respectively). Smolts were reared at Hagerman National and Magic Valley fish hatcheries.

A total of 176 adult steelhead (B-strain), comprised of 99 males (56.3%) and 77 females (43.7%), returned to the East Fork Salmon River weir between March 30 and May 12, 1993 (Table 6) (Chapman 1993). The male component of the run was comprised of 91 hatchery origin fish (91.9%) and eight natural origin (unmarked) fish (8.1%), and the female component was made up of 68 hatchery origin fish (88.3%) and nine natural origin fish (11.7%). All of the natural origin fish and 56 of the hatchery origin fish (34 males and 22 females) were released to spawn naturally. Forty-six females and 57 males were spawned on nine different dates and produced 211,993 green eggs. A total of 178,925 eggs (84.4%) developed to the eyed stage.

Smolt-to-Adult Return Rates

Clearwater Fish Hatchery

Clearwater Fish Hatchery became operational in 1992. The first steelhead smolts were released in April 1993 (brood year 1992). Adults resulting from this release will return in 1995, 1996, and 1997.

Hagerman National Fish Hatchery

The 1992-1993 steelhead return included three age classes of fish which were released from Hagerman National Fish Hatchery in 1989 (brood year 1988), 1990 (brood year 1989), and 1991 (brood year 1990). Brood year 1988, 1989, and 1990 steelhead returned as 3-ocean, 2-ocean, and 1-ocean fish, respectively. The Harvest Monitoring Project estimated that 5,589 of the steelhead that returned to Idaho in 1992-1993 were reared at Hagerman National Fish Hatchery (Table 3). This equals 41.1% of the hatchery's adult return goal (13,600 fish). However, the number of smolts released in 1988, 1989, and 1990 was approximately 60% of the smolt production target for each year (Table 4).

A total of 1,472,623 steelhead were released in 1989 (brood year 1988). Adult return data were not available for the smolts released into Clear Creek and the South Fork Clearwater River. Excluding the Clear Creek and South Fork Clearwater River fish releases, an estimated 1,391 adults returned from 1,073,127 smolts released in 1989 to yield a SAR of 0.13% (Appendix B. Table 1). In 1990, 1,439,266 steelhead smolts (brood year 1989) were released (Appendix B. Table 2). The 2-ocean adult component for this brood returned during this reporting period. To date, an estimated 6,126 steelhead have returned to Idaho to yield a SAR of 0.43%. In 1991, Hagerman National Fish Hatchery released 1,436,910 brood year 1990 steelhead smolts (Appendix B. Table 3). Adult return data, specifically the 2- and 3-ocean components, for brood year 1990 are incomplete at this time.

The final adult steelhead from the 1988 (B-strain) and 1989 (A-strain) broods returned to Idaho during this reporting period. The smolt-to-adult return rates were calculated for coded-

wire-tagged smolts, by stock and release site (Table 7). Smolt-to-adult return rates for Sawtooth A-stock steelhead smolts, brood year 1989, released at the Sawtooth Fish Hatchery weir and at Shoup Bridge (Salmon River) were 0.64% and 0.31%, respectively (Table 7). The SAR for brood year 1988 Dworshak B-stock steelhead released into the East Fork Salmon River was 0.05%.

Magic Valley Fish Hatchery

The 1992-1993 steelhead return included three age classes of fish which were released from Magic Valley Fish Hatchery in 1989 (brood year 1988), 1990 (brood year 1989), and 1991 (brood year 1990). Brood year 1988, 1989, and 1990 steelhead returned as 3-ocean, 2-ocean, and 1-ocean fish, respectively. The Harvest Monitoring Project estimated that 6,005 of the adult steelhead that returned to Idaho in 1992-1993 were reared at Magic Valley Fish Hatchery (Table 3). This equals 51.5% of the hatchery's adult return goal (11,660 fish).

A total of 2,201,994 steelhead smolts were released in 1989 (brood year 1988) from Magic Valley Fish Hatchery. An estimated 3,830 of these smolts returned to Idaho as adults (Appendix C. Table 1). The SAR for brood year 1988 was 0.17%. A total of 2,122,900 steelhead smolts were released from Magic Valley Fish Hatchery in 1990 (brood year 1989). The 3-ocean adult component of brood year 1989 is not complete. To date, an estimated 6,881 steelhead have returned to Idaho to yield a SAR of 0.32% (Appendix C. Table 2). In 1991, Magic Valley Fish Hatchery released 2,062,000 steelhead smolts (brood year 1990). Adult return data, specifically the 2- and 3-ocean components, for brood year 1990 are incomplete at this time (Appendix C. Table 3).

The final adult steelhead from the 1988 (B-strain) and 1989 (A-strain) broods returned to Idaho during this reporting period. Smolt-to-adult return rates were calculated for coded-wire-tagged smolts, by stock and release site (Table 7). East Fork B-stock steelhead smolts released at the East Fork Salmon River had a SAR of 0.10% (Table 7). Sawtooth A-stock steelhead smolts released at the Sawtooth Fish Hatchery weir had a SAR of 0.31%.

Experimentation - Objective 2

Clearwater Fish Hatchery

<u>Cover Experiment</u>—The PIT tag interrogation rates to downstream dams were as follows: raceway 9 (Cover Group)—71%, raceway 10 (Cover Group)—70%, and raceway 11 (No Cover Group)—75%. The number of PIT-tagged steelhead interrogated at downstream dams was not significantly different ($\chi^2 = 0.0001$, P = 1.0) between the two raceways in the Cover Group (i.e., raceway 9 vs. raceway 10). Therefore, these data were pooled and tested against PIT tag data for raceway 11 (i.e., the No Cover Group). Chi-square analysis determined that interrogation rates were not significantly different ($\chi^2 = 0.47$, P = 0.495) between groups. Median travel time to Lower Granite Dam for steelhead with shade covers was approximately 17 days, whereas the median travel time for fish without shade covers was approximately 21 days (Table 1). However, there was no significant difference (P = 0.059) in travel times to Lower Granite Dam between groups.

As mentioned previously, all steelhead released from Clearwater Fish Hatchery in 1993 were inadvertently exposed to water supersaturated with nitrogen. Therefore, the integrity of this experiment may have been compromised. Adults will return in 1995, 1996, and 1997.

Hagerman National Fish Hatchery

<u>Size-at-Release Experiment</u>—The first adults from this experiment returned during this reporting period. A total of 179 adult steelhead returned to Idaho as 1-ocean fish. Of these, 102 were from the large size group and 77 were from the regular size group (Table 8). For adults that returned from the large size group, 35% (36) were females and 65% (66) were males. Adult returns from the regular size smolts were composed of 47% (36) females and 53% (41) males. Sixty-one percent of the adults that returned to Idaho were harvested in Idaho's sport fishery. Adult returns for brood years 1990 and 1991 will be complete in 1994 and 1995, respectively. Complete results of this study will be reported in a separate report.

Acclimation Experiment—For brood year 1992, a total of 188 (62.7%) of the PIT-tagged acclimated steelhead were interrogated at downstream dams as compared to 161 (53.7%) of the tagged non-acclimated fish (Table 1). Interrogation rates were not significantly different (P > 0.05) among replicate raceways for each group. Therefore, data from replicate raceways, within group, were pooled and an overall chi-square test was performed. There were significantly (χ^2 = 4.63, P = 0.031) more acclimated fish interrogated than non-acclimated fish. Among replicate raceways, median travel times to Lower Granite Dam ranged from 34.0 to 36.7 days for the acclimated group and from 31.3 to 36.3 days for the non-acclimated group (Table 1). Travel times to Lower Granite Dam were not significantly (P = 0.182) different between groups.

Adults from brood year 1992 will return in 1995 and 1996. Adult steelhead from the first year of the study, brood year 1991, will return in 1994 and 1995. Complete results of this study will be reported in a separate report.

LITERATURE CITED

- Ainsworth, B., M. A. Graham, and M. Baer. 1993. Magic Valley Hatchery 1992 brood year report. Idaho Department of Fish and Game. Boise, Idaho.
- Ainsworth, B., D. May, D. Aplanalp, and K. Hills. 1995. Magic Valley Hatchery 1993 brood year report. Idaho Department of Fish and Game. Boise, Idaho.
- Ball, K. 1992. Evaluation of the hatchery-wild composition of Idaho salmon and steelhead harvest. United States Fish and Wildlife Service Lower Snake River Fish and Wildlife Compensation Plan contract no. 14-16-0001-89501 (RWG) (October 1, 1989, to December 31, 1990). Idaho Department of Fish and Game. Boise, Idaho.
- Ball, K. 1994. Evaluation of the hatchery-wild composition of Idaho salmon and steelhead harvest. United States Fish and Wildlife Service Lower Snake River Fish and Wildlife Compensation Plan (October 1, 1991, to December 31, 1992). Idaho Department of Fish and Game. Boise, Idaho.
- Ball, K. 1996. Evaluation of the hatchery-wild composition of Idaho salmon and steelhead harvest. United States Fish and Wildlife Service-Lower Snake River Fish and Wildlife Compensation Plan (October 1, 1992, to December 31, 1993). Idaho Department of Fish and Game. Boise, Idaho.
- Cannamela, D. A. 1992. Fish hatchery evaluations—Idaho. United States Fish and Wildlife Service Lower Snake River Fish and Wildlife Compensation Plan contract no. 14-16-0001-90502 (October 1, 1989, to September 30, 1990). Idaho Department of Fish and Game. Boise, Idaho.
- Chapman, J. 1993. Sawtooth Fish Hatchery and East Fork Satellite 1993 steelhead run report. Idaho Department of Fish and Game. Boise, Idaho.
- Columbia River Basin Pit Tag Information System. 1997. Pacific States Marine Fisheries Commission. Gladstone, Oregon.
- George, B., D. Baker, J. McGehee, D. Burton, and J. Rankin. 1994. Clearwater Fish Hatchery 1991 chinook brood year and 1992 steelhead brood year. Idaho Department of Fish and Game. Boise, Idaho.
- George, B., J. McGehee, and D. Munson. 1995. Clearwater Fish Hatchery 1992 chinook brood year and 1993 steelhead brood year. Idaho Department of Fish and Game. Boise, Idaho.
- Hagerman National Fish Hatchery. 1992. Hagerman National Fish Hatchery annual report, fiscal year 1992. United States Fish and Wildlife Service. Hagerman, Idaho.
- Hagerman National Fish Hatchery. 1993. Hagerman National Fish Hatchery annual report, fiscal year 1993. United States Fish and Wildlife Service. Hagerman, Idaho.

Petrosky, C. E. 1991. Influence of smolt migration flows on recruitment and return rates of Idaho spring chinook. Idaho Department of Fish and Game. Boise, Idaho.

SYSTAT, Inc. 1992. SYSTAT for windows: statistics, version 5 edition. Evanston, Illinois.

Table 1. Number of unique PIT tag interrogations of LSRCP steelhead smolts, by PIT tag file, at Lower Granite Dam (GRJ), Little Goose Dam (GOJ), Lower Monumental Dam (LMJ), and McNary Dam (MCJ) for the 1993 migration period. A total of 2,200 PIT-tagged steelhead were released from Hagerman National, Magic Valley, and Clearwater fish hatcheries between April 8 and April 21, 1993. Median travel time is to Lower Granite Dam.

				Number / Percent Interrogated							Median			
				G	RJ	G	OJ	L	MJ	M	CJ	TO	TAL	Travel
File Name	Release Site	Rel. Date	No. Rel.	No.	%	No.	%	No.	%	No.	%	No.	%	Time (Days)
Magic Valley Hatch Contribution Study	<u>ery</u>													
Dworshak B-stock DAC93054.M01 DAC93054.M05	E.F. Salmon R. E.F. Salmon R.	4/8 4/8	100 100	33 42	33.0 42.0	9 6	9.0 6.0	7 4	7.0 4.0	2	2.0 1.0	51 53	51.0 53.0	38.0 37.3
East Fork B-stock DAC93054.M09	E.F. Salmon R.	4/9	100	34	34.0	11	11.0	3	3.0	3	3.0	51	51.0	36.9
Dworshak B-stock DAC93054.M04 DAC93054.M06 DAC93054.M08	Hazard Cr. Hazard Cr. Hazard Cr.	4/20 4/20 4/21	100 100 100	66 59 66	66.0 59.0 66.0	7 10 7	7.0 10.0 7.0	3 6 4	3.0 6.0 4.0	0 2 1	0.0 2.0 1.0	76 77 78	76.0 77.0 78.0	23.5 17.1 15.5
Contribution Total			600									386	64.3	
Pahsimeroi A-stock DAC93054.M10	N.F. Salmon R.	4/16	200	11 1	55.5	17	8.5	10	5.0	3	1.5	141	70.5	21.0
DAC93054.M11 DAC93054.M13 DAC93054.M15 DAC93054.M14 DAC93054.M16	Salmon R. Lemhi R. Lemhi R. Pahsimeroi weir Pahsimeroi weir	4/13 4/14 4/15 4/12 4/12	100 100 100 100 100	55 62 62 48 62	55.0 62.0 62.0 48.0 62.0	10 8 5 6 3	10.0 8.0 5.0 6.0 3.0	4 2 6 2 0	4.0 2.0 6.0 2.0 0.0	0 0 1 1 0	0.0 0.0 1.0 1.0 0.0	69 72 74 57 65	69.0 72.0 74.0 57.0 65.0	24.7 24.5 20.0 24.9 26.0
Total			700									478	68.3	

Magic V	alley Hatchery Grand	l Total
Table 1	Continued	

Clearwater Fish Hatchery Grand Total

1300

300

864 66.5

72.0

216

			Number / Percent Interrogated								Median			
				G	RJ	G	OJ	LI	MJ	M	CJ	TO	TAL	Travel
File Name	Release Site	Rel. Date	No. Rel.	No.	%	No.	%	No.	%	No.	%	No.	%	Time (Days)
Hagerman Nationa Acclimation Study, (
Pahsimeroi A-stock														
DAC93051.H69	Sawtooth weir	4/9	100	49	49.0	8	8.0	3	3.0	0	0.0	60	60.0	36.7
DAC93051.H70	Sawtooth weir	4/9	100	45	45.0	16	16.0	1	1.0	3	3.0	65	65.0	35.6
DAC93051.H71	Sawtooth weir	4/9	100	50	50.0	10	10.0	2	2.0	1	1.0	63	63.0	34.0
Acclimation Total			300									188	62.7	
Acclimation Study,	(Non-acclimated Gro	oup)												
Pahsimeroi A-stock	•													
DAC93051.H72	Sawtooth weir	4/9	100	37	37.0	10	10.0	2	2.0	2	2.0	51	51.0	32.9
DAC93051.H73	Sawtooth weir	4/9	100	43	43.0	4	4.0	4	4.0	1	1.0	52	52.0	36.3
DAC93051.H74	Sawtooth weir	4/9	100	45	45.0	7	7.0	4	4.0	2	2.0	58	58.0	31.3
Control Total			300									161	53.7	
Hagerman NFH Gr	and Total		600											
Clearwater Fish Ha	atchery													
Cover Study														
Dworshak B-stock														
DAC93021.C09	S.F. Clearwater R.	4/13	100	63	63.0	7	7.0	0	0.0	1	1.0	71	71.0	17.4
DAC93021.C10	S.F. Clearwater R.	4/13	100	60	60.0	7	7.0	3	3.0	0	0.0	70	70.0	17.0
DAC93021.C11	S.F. Clearwater R.	4/13	100	68	68.0	6	6.0	0	0.0	1	1.0	75	75.0	20.8

Table 2. Snake River mean daily in-flow (thousand cubic feet per second) at Lower Granite Dam, Washington, from 1977-1993 during the Peak and Extended chinook salmon smolt migration periods as defined by Petrosky (1991).

Year	Peak (04/15 - 05/05)	Extended (04/20—05/30)
1977	39.1	40.2
_		
1978	85.4	95.8
1979	64.8	89.9
1980	87.5	102.9
1981	76.2	86.7
1982	116.8	131.6
1983	85.6	111.3
1984	121.9	146.1
1985	86.9	87.2
1986	93.4	105.7
1987	59.0	62.4
1988	55.1	64.2
1989	93.6	87.2
1990	63.8	66.4
1991	44.0	70.5
1992	54.2	57.3
1993	69.8	114.0

Table 3. Estimated number of LSRCP steelhead that returned to Idaho in 1992-1993. The adult return in 1992-1993 included fish from three age classes. Steelhead were reared at Hagerman National and Magic Valley fish hatcheries. These estimates were prepared by the Idaho Department of Fish and Game's Harvest Monitoring Project and only include steelhead harvested in Idaho's sport fisheries, steelhead that returned to hatchery racks, and steelhead that escaped to spawn naturally. These are minimum estimates and do not include all tributary and mainstem strays or in-river pre-spawning mortalities.

HATCHERY	BROOD YEAR	3-OCEAN	2-OCEAN	1-OCEAN
Hagerman Hagerman Hagerman	1988 1989 1990	70 	 1,974 	 3,545
Estimated Return				5,589
Magic Valley Magic Valley Magic Valley	1988 1989 1990	0 	2,506 	 3,499
Estimated Return				6,005

Grand Total 11,594

Table 4. Steelhead smolts released from Magic Valley and Hagerman National fish hatcheries that contributed to the 1992-1993 Idaho steelhead return. The number of steelhead smolts released and the estimated number of adults that returned were compared to facility design production targets and projected adult return goals.

	Releases Contribut	ing to 1992-19	93 Adult Retur	ns	1992-93					
Brood Year	Fish Hatchery	Number Released	Design Target	Percent of Target	Adult Returns					
1988	Magic Valley	2,201,994	2,000,000	110%	0					
1988	Hagerman NFH	1,472,623	2,400,000	61%	70					
	Total	3,674,617	4,400,000	84%	70					
1989	Magic Valley	2,122,900	2,000,000	106%	2,506					
1989	Hagerman NFH	1,439,266	2,400,000	60%	1,974					
	Total	3,562,166	4,400,000	81%	4,480					
1990	Magic Valley	2,062,000	2,000,000	103%	3,499					
1990	Hagerman NFH	1,436,910	2,400,000	60%	3,545					
	Total	3,498,910	4,400,000	80%	7,044					
	Mean annual rele	ease as percer	nt of target:	82%						
		Total adult return ^a :								
		Adult return goal: Percent of goal achieved:								

^a Estimate includes only steelhead harvested in Idaho's sport fisheries, steelhead that returned to hatchery racks, and off-site escapement. These are minimum estimates and do not include all tributary and mainstem strays or in-river pre-spawning mortalities.

Summary of the 1993 A-strain steelhead trout return to the Sawtooth Fish Hatchery weir. The fish return included fish of hatchery and natural origin. Hatchery aging criteria, based on length, were used to determine age^a. ND indicates and Table 5. data (i.e., data were not available).

	HATCHERY ORIGIN n = 1584											
	Males n = 1151 Females n = 433											
Age ^b	Trapped	pped Released Spawned Morts Other Trapped Released Spawned Morts Other										
1-ocean	1010	ND	ND	0	ND	331	ND	ND	0	ND		
2-ocean	141	ND	ND	0	ND	102	ND	ND	0	ND		
Total	1151	487°	530	0	134 ^d	433	174°	255	0	4 ^d		

	NATURAL ORIGIN n = 7										
	Males n = 3 Females n = 4										
Age ^b	Trapped	apped Released Spawned Morts Other Trapped Released Spawned Morts Other									
1-ocean	0	0	0	0	0	1	1	0	0	0	
2-ocean	3	3	0	0	0	3	3	0	0	0	
Total	3	3 ^c	0	0	0	4	4 ^c	0	0	0	

Total number trapp	ed 1,591	Green egg number	1,131,877	
Trapping period	3/18 - 5/12/93	Eyed egg number	1,031,635	(91.1% eye-up)

^a Fish were aged using the following aging criteria:	RUN	SEX	<u>LENGTH</u>	AGE (years	in ocean)
	Α	·	male	≤ 68 cm	1-ocean
	Α		male	> 68 cm	2-ocean
	Α		female	≤ 65 cm	1-ocean
	Α		female	> 65 cm	2-ocean

b Hatchery fish classified as 1-ocean were released in 1991, brood year 1990. Hatchery fish classified as 2-ocean were released in 1990, brood year 1989.

Fish were released above the weir.

Fish were killed, but the eggs and milt were not used.

Summary of the 1993 B-strain steelhead trout return to the East Fork Salmon River weir. The fish return included fish of Table 6. hatchery and natural origin. Hatchery aging criteria, based on length, were used to determine agea. ND indicates "no data" (i.e., data were not available).

	HATCHERY ORIGIN n = 159											
	Males n = 91 Females n = 68											
Age ^b	Trapped	Released	Spawned	Morts	Trapped	Released	Spawned	Morts	Other			
1-ocean	57	ND	ND	0	0	2	ND	ND	0	0		
2, 3-ocean	34	ND	ND	0	0	66	ND	ND	0	0		
Total	91	34°	57	0	0	68	22 °	46	0	0		

			l	NATURA	L ORIGIN	n = 17					
	Males n = 8 Females n = 9										
Age ^b	Trapped	Released	Spawned	Morts	Trapped	Released	Spawned	Morts	Other		
1-ocean	5	5	0	0	0	1		0	0	0	
2, 3-ocean	3	3	0	0	0	8	8	0	0	0	
Total	8	8°	0	0	0	9	9°	0	0	0	

Total number trapped Trapping period	176 3/30 - 5/12/93		n egg number egg number	211,993 178,925	(84.4% eye-up)
^a Fish were aged using	the following aging criteria:		SEX LENGTH		ears in ocean)
		B B	male male	≤ 73 cm > 73 cm	1-ocean 2- or 3-ocean
		В	female	≤ 68 cm	1-ocean
		В	female	> 68 cm	2- or 3-ocean

b Hatchery fish classified as 1-ocean were released in 1991, brood year 1990. Hatchery fish classified as 2- or 3-ocean were released 1990 and 1989, respectively, (brood years 1989 and 1988, respectively).
c Fish were released above the weir.

Table 7. Smolt-to-adult return rates of coded-wire tagged steelhead smolts released from Hagerman National and Magic Valley fish hatcheries. The number of adults was estimated by Ball (1996) and only include steelhead harvested in Idaho's sport fisheries, steelhead that returned to hatchery racks, and steelhead that escaped to spawn naturally. These are minimum estimates and do not include all tributary and mainstem strays or in-river pre-spawning mortalities.

HATCHERY	STOCK	BROOD	RELEASE SITE	NUMBER CWT	NUMBER ADULTS	SAR (%)
Hagerman	DWOR B	1988	E.F. Salmon R.	43,569	23	0.05
Hagerman	SAW A	1989	Sawtooth FH	45,297	289	0.64
Hagerman	SAW A	1989	Salmon R.	45,828	144	0.31
Magic	E.F. B	1988	E.F. Salmon R.	44,064	42	0.10
Magic	PAH A	1989	Sawtooth FH.	39,620	123	0.31

Table 8. Total number^a of steelhead recovered with coded-wire tags designating them as either large size^b or regular size^c. Recovered fish were released as smolts at the Sawtooth Fish Hatchery weir in 1991 (brood year 1990). ND indicates "no data" (i.e., data were not available).

				Return Co	mpositio	n	
		Number	1-C	cean	2-0	Total	
Size Group	Brood Year	CWT	Male	Female	Male	Female	Return
Large	1990	53,245	66	36	ND	ND	102
Regular	1990	61,431	41	36	ND	ND	77
Large Regular	1991 1991	53,463 45,646	ND ND	ND ND	ND ND	ND ND	ND ND

^a Includes all Idaho fishery harvest returns, Idaho hatchery returns, and tributary stray recoveries

^b Large size steelhead averaged 241 mm (3.0 fish per pound) at time of release

^c Regular size steelhead averaged 221 mm (4.5 fish per pound) at time of release




Figure 1. Number of unique PIT tag interrogations of LSRCP juvenile steelhead at Lower Granite Dam, Washington, plotted with the average daily in-flow of the Snake River at Lower Granite Dam in 1993. A total of 2,200 PIT-tagged steelhead smolts were released from Hagerman National, Magic Valley, and Clearwater fish hatcheries between April 8 and April 21, 1993. Fifty-three percent (1,160) of the PIT-tagged fish were interrogated at Lower Granite Dam. Flow data were not available for all dates. Data for one fish falls outside of this date range and is not shown.

APPENDICES

Appendix A. Table 1. Release data and estimated adult returns for Clearwater Fish Hatchery summer steelhead trout, brood year 1992. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Clearwate	er FH	Brood Y	ear: 1992	lo	dentifying	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		eturn position	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
S.F. Clearwater R. Milept. 18 above Stites	9	43,050	DWOR B	CWT/LV/AD *LV/AD	102947	25,673 155	10.1	Shade/Cover Experiment	1 2	ND ND	ND ND	ND ND	ND ND
4/13/93				*AD CLIP *AD/PIT		17,122 100		Experimental Group	3	ND	ND	ND	ND
S.F. Clearwater R. Milept. 18 above Stites	11	45,180	DWOR B	CWT/LV/AD *LV/AD	104937	22,003 66	8.9	Shade/Cover Experiment	1 2	ND ND	ND ND	ND ND	ND ND
4/13-14/93				*AD *AD/PIT		23,011 100		Control Group	3	ND	ND	ND	ND
S.F. Clearwater R. Milept. 18 above Stites	10	38,350	DWOR B	CWT/LV/AD *LV/AD	104938	21,340 64	9.1	Shade/Cover Experiment	1 2	ND ND	ND ND	ND ND	ND ND
4/13/93				*AD *AD/PIT		16,846 100		Experimental Group	3	ND	ND	ND	ND
S.F. Clearwater R. Milept. 18 above Stites 4/12-13/93	ND	199,720	DWOR B	*AD		199,720	9.3	Production					
S.F. Clearwater R. Milept. 18 above Stites			DWOR B	NON-CWT (includes all *)		257,284	ND	Production (Includes all *)	1 2	ND ND	ND ND	ND ND	ND ND

Includes all release dates that have mark type indicated by *

ND ND

3

ND

ND

Appendix A. Table 1. Continued.

		RW Total	Stock ID	Identifying Marks									
Release Site/Date	RW NO.			Mark Type	CWT Code	Release Number	Size (FPP)	Marking Purpose	Age Ocean	Return Composition		Total	SAR
										Harvest	Hatchery	Returns	(%)
			TOTAL NO	/T RELEASE N-CWT RELE E RELEASE	ASE	69,016 257,284 326,300		TOTAL RETURN	l:	ND	ND	ND	ND
TOTAL CWT RELEASE FOR CLEARWATER FH TOTAL NON-CWT RELEASE FOR CLEARWATER FH TOTAL CLEARWATER FH RELEASE TOTAL PIT TAGS						69,016 257,284 326,300 300							

Appendix A. Table 2. Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1992. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Hagerman	n NFH	Brood '	Year: 1992										
				Identifying N	larks								
											eturn		
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		osition	Total	SAR
Site/Date	NO.	Total	<u>ID</u>	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Hammer Cr. Lower Salmon R. 4/28/93	95	67,581	SAW A	CWT/LV/AD *LV/AD *AD	104946	17,142 173 50,266	4.8	Production	1 2	ND ND	ND ND	ND ND	ND ND
Hammer Cr. Lower Salmon R. 4/28/93	96	77,389	SAW A	CWT/LV/AD *LV/AD *AD	104947	19,655 99 57,635	4.8	Production	1 2	ND ND	ND ND	ND ND	ND ND
Hammer Cr. Lower Salmon R. 4/28/93	97	66,036	SAW A	CWT/LV/AD *LV/AD *AD	104948	16,662 515 48,859	4.8	Production	1 2	ND ND	ND ND	ND ND	ND ND
Hammer Cr. Lower Salmon R.		157,547	SAW A	NON-CWT		157,547	ND	Production	1	ND	ND	ND	ND
Includes all release d		at have		(includes all *)				(Includes all *)	2	ND	ND	ND	ND
Mark Type indicated l	Dy		TOTAL N	WT RELEASE ON-CWT RELEAS ITE RELEASE	SE	53,459 157,547 211,006		TOTAL RETURN	:	ND	ND	ND	ND
Sawtooth weir 4/9/93	74	19,229	РАН А	CWT/LV/AD CWT/LV/AD/PIT *AD	104949 104949	19,196 (100) 33	4.9	Acclimation Control Group Trucked to SAWT on 4/9/93	1 2	ND ND	ND ND	ND ND	ND ND
Sawtooth weir 4/9/93	73 74	18,251	PAH A	CWT/LV/AD CWT/LV/AD/PIT	104950 104950	18,168 (100)	4.7	Acclimation Control Group	1 2	ND ND	ND ND	ND ND	ND ND

*LV/AD
*AD

51 32 Trucked to SAWT on 4/9/93

Appendix A.	Table 2	Continued

				Identifying N	larks								
Release Site/Date	RW NO.	RW Total	Stock ID	Mark Type	CWT Code	Release Number	Size (FPP)	Marking Purpose	Age Ocean	Comp	eturn position Hatchery	Total Returns	SAR (%)
Sawtooth weir 4/9/93	72 73	17,899	PAH A	CWT/LV/AD CWT/LV/AD/PIT *LV/AD *AD	104951 104951	17,818 (100) 50 31	4.9	Acclimation Control Group Trucked to SAWT on 4/9/93	1 2	ND ND	ND ND	ND ND	ND ND
Sawtooth weir 4/9/93	72	4,571	PAH A	CWT/LV/AD *AD	105034	4,563 8	4.9	Acclimation Control Group Trucked to SAWT on 4/9/93	1 2	ND ND	ND ND	ND ND	ND ND
Sawtooth weir 4/8/93	69	207,592	РАН А	CWT/LV/AD *LV/AD *AD *PIT	105020	20,262 246 187,084 (100)	5.1	Acclimation Experimental Group Trucked to SAWT on 3/18- 24/93	1 2	ND ND	ND ND	ND ND	ND ND
Sawtooth weir 4/8/93	69 70	191,791	РАН А	CWT/LV/AD *LV/AD *AD	105021	18,726 164 172,901	4.8	Acclimation Experimental Group Trucked to SAWT on 3/18- 24/93	1 2	ND ND	ND ND	ND ND	ND ND
Sawtooth weir 4/8/93	70 71	186,674	РАН А	CWT/LV/AD *LV/AD *AD *PIT	105022	18,235 71 168,368 (100)	5	Acclimation Experimental Group Trucked to SAWT on 3/18- 24/93	1 2	ND ND	ND ND	ND ND	ND ND
Sawtooth weir 4/8/93	71	83,513	PAH A	CWT/LV/AD *AD *PIT	105010	8,161 75,352 (100)	5.1	Acclimation Experimental Group Trucked to SAWT on 3/18-	1 2	ND ND	ND ND	ND ND	ND ND

24/93

Appendix A	Table 2	Continued
AUDEILUIX A	14018	COMMINICIA

				Identifying N	/larks								
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		eturn position	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Sawtooth weir 4/19/93&4/22/93	85- 91	140,626	SAW A	*AD		(140,626)	ND	Production					
Sawtooth Weir Includes all release of Mark Type indicated		at have	SAW A PAH A	NON-CWT (Includes all *)	140,626 463,765	604,391	ND	Production (Includes all *)	1 2	ND ND	ND ND	ND ND	ND ND
Mark Type muleated	IJ		TOTAL N	WT RELEASE ON-CWT RELEA ITE RELEASE	SE	125,129 604,391 729,520		TOTAL RETURN:	1				
Hazard Cr. L. Salmon R. 4/12/93	45	219,946	SAW A	CWT/LV/AD *LV/AD *AD	105017	20,045 340 199,561	5	Production	1 2	ND ND	ND ND	ND ND	ND ND
Hazard Cr. L. Salmon R. 4/12/93	45 46	112,344	SAW A	CWT/LV/AD *LV/AD *AD	105018	10,245 103 101,996	5	Production	1 2	ND ND	ND ND	ND ND	ND ND
Hazard Cr. L. Salmon R. 4/12/93	46	215,026	SAW A	CWT/LV/AD *LV/AD *AD	105019	19,621 65 195,340	4.8	Production	1 2	ND ND	ND ND	ND ND	ND ND
Hazard Cr. L.			SAW A	NON-CWT		497,405	ND	Production	1	ND	ND	ND	ND
Salmon R. Includes all release dates that have Mark Type				(Includes all *)				(Includes all *)	2	ND	ND	ND	ND
indicated by *			TOTAL N	WT RELEASE ON-CWT RELEA ITE RELEASE	SE	49,911 497,405 547,316		TOTAL RETURN:	:	ND	ND	ND	ND
TOTAL SAW A-STO TOTAL SAW A-STO TOTAL SAW A-STO	CK NC	N-CWT R				103,370 795,578 898,948							

Appendix A. Table 2. Continued.

				Identifying	Marks								
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		eturn position	Total	SAR
Site/Date	<u>NO.</u>	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
TOTAL PAH A-S	STOCK CW	T RELEA	SE			125,129							
TOTAL PAH A-S	STOCK NO	N-CWT R	ELEASE			463,765							
TOTAL PAH A-S	STOCK REL	EASE				588,894							
TOTAL CWT RE	LEASE FO	R HAGEI	RMAN HATO	HERY		228,499							
TOTAL NON-CV	VT RELEAS	E FOR H	IAGERMAN	HATCHERY		1,259,343							
TOTAL HAGERI	MAN HATC	HERY RE	ELEASE			1,487,842							
TOTAL PIT TAG	S					600							

Appendix A. Table 3. Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1992. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Magic V	alley FH	Brood	Year: 1992		ifying Mar	ks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Comp	eturn position	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
E.F. Salmon R. 4/7-8/93	1	212,881	DWOR B	CWT/LV/AD *LV/AD *LV/AD/PIT *AD	105007	18,924 792 100 193,065	6.6	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
E.F. Salmon R. 4/7-8/93	5	178,119	DWOR B	CWT/LV/AD *LV/AD *LV/AD/PIT *AD	105005	16,024 62 100 161,933	6.9	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
E.F. Salmon R. 4/7-9/93	9	106,400	E FK B	CWT/LV/AD *LV/AD *LV/AD/PIT *AD	105009	19,128 1,406 100 85,766	6.1	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND
E.F. Salmon R. 4/7-9/93		443,324	DWOR B E FK B	NON-CWT (Includes all *)	356,052 87,272	443,324	ND	Production (Includes all *)	1 2	ND ND	ND ND	ND ND	ND ND
			TOTAL NO	VT RELEASE ON-CWT RELE TE RELEASE	ASE	54,076 443,324 497,400		TOTAL RETUR	3 N:	ND ND	ND ND	ND ND	ND ND
Hazard Cr. L. Salmon R. 4/19-20/93	4	109,900	DWOR B	CWT/LV/AD *LV/AD *LV/AD/PIT *AD	105004	19,358 1,583 100 88,859	4.7	Contribution	1 2 3	ND ND ND	ND ND ND	ND ND ND	ND ND ND

Appendix A. Table 3. Continued.

				Ident	ifying Mar	ks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Comp	eturn position	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Hazard Cr.	6	126,500	DWOR B	CWT/LV/AD	105006	19,932	6.9	Contribution	1	ND	ND	ND	ND
L. Salmon R. 4/16/93				*LV/AD *LV/AD/PIT		1,016 100			2 3	ND ND	ND ND	ND ND	ND ND
Hazard Cr.	8	88,900	DWOR B	CWT/LV/AD	105008	19,909	6.6	Contribution	1	ND	ND	ND	ND
L. Salmon R.				*LV/AD		1,081			2	ND	ND	ND	ND
4/16-19/93				*LV/AD/PIT *AD		100 67,810			3	ND	ND	ND	ND
Hazard Cr.		266,101	DWOR B	NON-CWT		180,558	ND	Production	1	ND	ND	ND	ND
L. Salmon R.				(Includes all *)				(Includes all *)	2	ND	ND	ND	ND
4/16-20/93				,					3	ND	ND	ND	ND
			TOTAL NO	NT RELEASE ON-CWT RELE	ASE	59,199 266,101							
			TOTAL SI	TE RELEASE		325,300		TOTAL RETUR	N:	ND	ND	ND	ND
Lemhi R.	13	66,700	PAH A	CWT/LV/AD	105013	19,692	5.7	Contribution	1	ND	ND	ND	ND
4/14/93				*LV/AD *LV/AD/PIT *AD		1,046 100 45,862			2	ND	ND	ND	ND
Lemhi R.	15	131,800	PAH A	CWT/LV/AD	105015	21,390	5.9	Contribution	1	ND	ND	ND	ND
4/14-16/93				CWT/LV/AD *LV/AD *LV/AD/PIT *AD	105012	22,106 118 100 88,086			2	ND	ND	ND	ND
Lemhi R.		135,312	PAH A	NON-CWT		135,312	ND	Production	1	ND	ND	ND	ND
4/14-16/93				(Includes all *)		, -		(Includes all *)	2	ND	ND	ND	ND
				NT RELEASE ON-CWT RELE	ASE	63,188 135,312							
			TOTAL SI	TE RELEASE		198,500		TOTAL RETUR	N:	ND	ND	ND	ND

N.F. Salmon R. 4/16-22/93	10	190,500	PAH A	CWT/LV/AD *LV/AD *LV/AD/PIT *AD	104924	65,637 935 200 123,728	5.4	Contribution	1 2	ND ND	ND ND	ND ND	ND ND
Appendix A. Table 3	3. Cont	inued.											
				Ident	ifying Mar	KS	-			De	eturn		
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		osition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest		Returns	(%)
N.F. Salmon R. 4/16-22/93		124,863	PAH A	NON-CWT (Includes all *)		124,863	ND	Production (Includes all *)	1 2	ND ND	ND ND	ND ND	ND ND
			TOTAL N	WT RELEASE ON-CWT RELE ITE RELEASE	ASE	65,637 124,863 190,500		TOTAL RETUR	N:	ND	ND	ND	ND
	4.0	100.000	5411.4	014/7/11/45	105010			0 1 " "		ND		ND	ND
Upper Salmon R. Ellis Bridge 4/12-13/93	16	122,300	PAH A	CWT/LV/AD *LV/AD *LV/AD/PIT *AD	105016	20,361 210 100 101,629	5.5	Contribution	1 2	ND ND	ND ND	ND ND	ND ND
Upper Salmon R. Ellis Bridge 4/12/93	14	144,000	РАН А	CWT/LV/AD *LV/AD *LV/AD/PIT *AD	105014	19,778 1,051 100 123,071	5.5	Contribution	1 2	ND ND	ND ND	ND ND	ND ND
Upper Salmon R. Ellis Bridge		226,161	PAH A	NON-CWT (Includes all		226,161	ND	Production (Includes all *)	1 2	ND ND	ND ND	ND ND	ND ND
4/12/93				*)									
			TOTAL N	WT RELEASE ON-CWT RELE ITE RELEASE	ASE	40,139 226,161 266,300		TOTAL RETUR	N:	ND	ND	ND	ND
Salmon R. at Challis 4/13/93	11	260,600	PAH A	CWT/LV/AD *LV/AD *LV/AD/PIT *AD	105011	19,924 883 100 239,693	5.9	Contribution	1 2	ND ND	ND ND	ND ND	ND ND
Salmon R. at Challis		240,676	PAH A	NON-CWT includes all *		240,676	ND	Production (Includes all *)	1 2	ND ND	ND ND	ND ND	ND ND

TOTAL PIT TAGS

4/13/93													
11 10/00			TOTAL NO	T RELEASE N-CWT REL E RELEASE		19,924 240,676 260,600		TOTAL RETU	RN:	ND	ND	ND	ND
Slate Cr.	2	187,100	DWOR B	*AD		187,100	6.1	Production	1	ND	ND	ND	ND
Appendix A. Table 3	. Conti	nued.											
			_	Iden	tifying Ma	rks	-			De	eturn		
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		osition	Total	SAR
Site/Date	NO.	Total	<u>ID</u>	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Upper Salmon R. 4/15/93	3								2	ND ND	ND ND	ND ND	ND ND
			TOTAL NO	T RELEASE N-CWT REL E RELEASE	EASE	0 187,100 187,100		TOTAL RETU	RN:	ND	ND	ND	ND
TOTAL DWOR B-ST TOTAL DWOR B-ST TOTAL DWOR B-ST	OCK N	ION-CWT				94,147 809,253 903,400							
TOTAL PAH A-STOO TOTAL PAH A-STOO TOTAL PAH A-STOO	CK NO	N-CWT R				188,888 727,012 915,900							
TOTAL E FK B-STO TOTAL E FK B-STO TOTAL E FK B-STO	CK NO	N-CWT R				19,128 87,272 106,400							
TOTAL CWT RELEATOTAL NON-CWT R	ELEAS	SE FOR M	AGIC VALLE			302,163 1,623,537 1,925,700							

1,300

Appendix B. Table 1. Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1988. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Hagerman NFH		E	Brood Year:	1988	lder	ntifying Mar	·ks						
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		eturn position	Total	SAR
Site/Date	NO.	Tota I	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Sawtooth Weir 4/11-13/89	ND	ND	SAW A	CWT/LV/AD	104138	14,718	4.9	Contribution	1 2	4 6	0 2	12	0.08
Sawtooth Weir 4/11-13/89	ND	ND	SAW A	CWT/LV/AD	104139	14,585	4.9	Contribution	1 2	0 10	0 3	13	0.09
Sawtooth Weir 4/11-13/89	ND	ND	SAW A	CWT/LV/AD	104140	16,914	4.9	Contribution	1 2	5 0	1 0	6	0.04
Sawtooth Weir 4/11-13/89	ND	ND	SAW A	AD		590,334	4.9	Production	1 2	288 435	86 102	911	0.15
7777 10700			TOTAL NO	NT RELEASE ON-CWT RELE TE RELEASE	ASE	46,217 590,334 636,551		TOTAL RET		748	194	942	0.15
East Fork Salmon R. at	ND	ND	DWOR B	CWT/LV/AD	104132	14,939	5	Contribution	1	0	0	4	0.03
trap 4/17-19/89									2 3	4 0	0 0		
East Fork Salmon R. at	ND	ND	DWOR B	CWT/LV/AD	104133	14,911	5	Contribution	1	0	0	10	0.07
trap 4/17-19/89									2 3	2 7	1 0		
East Fork Salmon R. at	ND	ND	DWOR B	CWT/LV/AD	104134	13,719	5	Contribution	1	1	1	9	0.07

trap 4/17-19/89 East Fork Salmon R. at trap 4/17-19/89 Appendix B. Table 1. Cont	ND	ND	DWOR B	AD		393,007	5	Production	2 3 1 2 3	7 0 24 300 63	0 0 18 21 0	426	0.11
					ldeı	ntifying Mar	ks						
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		eturn position	Total	SAR
Site/Date	NO.	Tota	ID	Туре	Code	Number	(FPP)	Purpose	Ocean		Hatchery	Returns	(%)
	·		•				· <u></u>	-					
East Fork Salmon R. at trap 4/17-19/89													
				VT RELEASE ON-CWT RELE	ΔSF	43,569 393,007							
				TE RELEASE	AGE	436,576		TOTAL RET	JRN:	408	41	449	0.10
Clear Cr. at Kooskia NFH 5/8/89	ND	ND	DWOR B	CWT/LV/AD	104135	15,294	5	Contribution	1 2	ND ND	ND ND	ND	ND
3/0/09									3	ND	ND		
Clear Cr. at Kooskia NFH	ND	ND	DWOR B	CWT/LV/AD	104136	15,482	5	Contribution	1	ND	ND	ND	ND
5/8/89									2 3	ND ND	ND ND		
Clear Cr. at Kooskia NFH	ND	ND	DWOR B	CWT/LV/AD	104137	14,374	5	Contribution	1	ND	ND	ND	ND
5/8/89									2 3	ND ND	ND ND		
Clear Cr. at Kooskia NFH	ND	ND	DWOR B	AD		3,997	5	Production	1	ND	ND	ND	ND
5/8/89									2 3	ND ND	ND ND		
				VT RELEASE ON-CWT RELE	ASE	45,150 3,997							
				TE RELEASE		49,147		TOTAL RET	JRN:	ND	ND	ND	ND
S. Fork Clearwater R.	ND	ND	DWOR B	AD		83,431	5	Production	1	ND	ND	ND	ND
Crooked R. Bridge 4/24/89									2 3	ND ND	ND ND		
				VT RELEASE ON-CWT RELE	ASE	0 83,431							

			TOTAL SI	TE RELEASE		83,431		TOTAL RET	URN:	ND	ND	ND	ND
S. Fork Clearwater R., Mill Cr. Bridge	ND	ND	DWOR B	AD		60,372	5	Production	1	ND	ND	ND	ND
Mill Cr. Bridge 4/26/89									2	ND ND	ND ND		
Appendix B. Table 1. Cont	inued.												
					Ide	ntifying Ma	rks		_	D.	eturn		
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		position	Total	SAR
Site/Date	NO.	Tota	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest		Returns	(%)
			TOTAL C	NT RELEASE		0							
				ON-CWT RELE TE RELEASE	EASE	60,372 60,372		TOTAL RET	URN:	ND	ND	ND	ND
S. Fork Clearwater R.	ND	ND	DWOR B	AD		103,273	5	Production	1	ND	ND	ND	ND
Newsome Cr. Bridge, (towr))							2	ND	ND		
4/26,5/1/89			TOTAL CI	NT RELEASE		0			3	ND	ND		
				ON-CWT RELE	EASE	0 103,273							
				TE RELEASE		103,273		TOTAL RET	URN:	ND	ND	ND	ND
S. Fork Clearwater R.	ND	ND	DWOR B	AD		103,273	5	Production	1	ND	ND	ND	ND
Eldorado Cr. 2nd Bridge, 5	/1,3/89								2	ND	ND		
			TOTAL C	NT RELEASE		0			3	ND	ND		
				ON-CWT RELE	EASE	103,273							
			TOTAL SI	TE RELEASE		103,273		TOTAL RET	URN:	ND	ND	ND	ND
TOTAL SAW A STOCK OF	NT PEI	EASE				46 047							
TOTAL SAW A-STOCK CV						46,217 590,334		RETURN IS	COMPLE	TE			
TOTAL SAW A-STOCK RE						636,551							
TOTAL DWOR B-STOCK (`WT DI		: E			88,719		* RETURN GRAND TO	ΓΛΙ-	1.156	235	1,391	0.09
TOTAL DWOR B-STOCK O						747,353		* includes S.		,		1,391	0.09
TOTAL DWOR B-STOCK F						836,072							
TOTAL CWT RELEASE FO	OR HAG	SERMA	AN NFH			134,936		** RETURN GRAND TO	ΓΔΙ -	1,156	235	1,391	0.13
IOIAL OWI NELEASE FO	AL DAC		714 141 LI			104,500		SIVAIND IO	IAL.	1,150	200	1,391	0.13

Appendix B. Table 2. Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1989. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Hagern	nan NFI	H Broo	od Year: 19		dentifying	Marks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		eturn position	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Sawtooth Weir 4/5/90	70 71	99,878	SAW A	CWT/LV/AD *LV/AD *AD	104214	14,597 147 85,134	4.7	Contribution	1 2	76 12	19 6	113	0.77
Sawtooth Weir 4/5/90	70 71	100,772	SAW A	CWT/LV/AD *LV/AD *AD	104215	15,482 156 85,134	4.7	Contribution	1 2	48 33	12 5	98	0.63
Sawtooth Weir 4/5/90	70 71	100,506	SAW A	CWT/LV/AD *LV/AD *AD	104216	15,218 154 85,134	4.7	Contribution	1 2	60 3	15 0	78	0.51
Sawtooth Weir Includes all release Mark Type indicate		that have	SAW A	NON-CWT (Includes all *	·)	255,859	4.7	Production (Includes all	1 2	2,072 314	518 44	2,948	1.15
mant Type maioatt	ou by		TOTAL N	WT RELEASE ION-CWT RELI SITE RELEASE	EASE	45,297 255,859 301,156		TOTAL RETU	JRN:	2,618	619	3,237	1.07
Salmon R. at	92	67,001	SAW A	CWT/LV/AD	104227	15,528	5	Contribution	1	24	17	41	0.26
Shoup Bridge 4/12/90	93			*AD		51,473			2	0	0		
Salmon R. at	92	66,669	SAW A	CWT/LV/AD	104228	15,196	5	Contribution	1	30	23	61	0.40
Shoup Bridge 4/12/90	93			*AD		51,473			2	8	0		

Appendix B. Table 2. Continued.

Appendix B. Table 2	<u> 2. 001</u>	itiriaca.		le	dentifying	Marks				_			
Release Site/Date	RW NO.	RW Total	Stock ID	Mark Type	CWT Code	Release Number	Size (FPP)	Marking Purpose	Age Ocean	Com	eturn position Hatchery	Total Returns	SAR (%)
Salmon R. at	92	66,576	SAW A	CWT/LV/AD	104229	15,104	5	Contribution	1	19	14	42	0.28
Shoup Bridge 4/12/90	93			*AD		51,472			2	9	0		
Salmon R. at Shoup Bridge	92		SAW A	NON-CWT		154,418	5	Production	1	179	119	355	0.23
Includes all release dates that	93	h*		(Includes all *	")			(Includes all *)	2	57	0		
have Mark Type indi	icated	by	TOTAL N	WT RELEASE ON-CWT RELI ITE RELEASE	EASE	45,828 154,418 200,246		TOTAL RETU	IRN:	326	173	499	0.25
Little Salmon R. at Hazard Cr. 4/30/90	48 49	131,854	DWOR B	CWT/LV/AD *LV/AD *AD	104230	14,841 49 116,964	4.4	Contribution	1 2 3	0 258 ND	0 258 ND	516	3.48
Little Salmon R. at Hazard Cr. 4/30/90	48 49	131,075	DWOR B	CWT/LV/AD *LV/AD *AD	104231	14,065 47 116,963	4.4	Contribution	1 2 3	0 4 ND	0 4 ND	8	0.06
Little Salmon R. at Hazard Cr. 4/30/90	49	130,423	DWOR B	CWT/LV/AD *LV/AD *AD	104232	13,416 44 116,963	4.4	Contribution	1 2 3	0 9 ND	0 9 ND	18	0.13
Little Salmon R. at Hazard Cr.	ND	ND	SAW A	*AD		80,465	4.2	Production	1	33	33	126	0.16
4/23/90									2	30	30		
Little Salmon R. at Hazard Cr.	ND		DWOR B	NON-CWT		351,030	4.4	Production	1	0	0	332	0.09
Includes all release Mark Type indicated		that have		(Includes all *	")			(Includes all	2	166	166		
mant Type maleated	. Jy			WT RELEASE ON-CWT RELI	FASF	42,322 431,495		,	3	ND	ND		
			_	ITE RELEASE	LAJE	473,817		TOTAL RETU	IRN:	500	500	1,000	0.21

Appendix B. Table 2. Continued.

				lc	dentifying	g Marks				_			
Release Site/Date	RW NO.	RW Total	Stock ID	Mark Type	CWT Code	Release Number	Size (FPP)	Marking Purpose	Age Ocean	Com	eturn position Hatchery	Total Returns	SAR (%)
Site/Date	140.	Total		Туре	Code	- Hullibei	(177)	- ruipose	Ocean	i iai vest	Tiatchery	Neturns	(/0)
Salmon R. at Ellis Bridge	ND	ND	SAW A	*AD		200,295	4.4	Production	1	252	168	700	0.35
4/9-11/90				WT RELEASE		0			2	256	24		
			_	ION-CWT RELE SITE RELEASE	EASE	200,295 200,295		TOTAL RETU	JRN:	508	192	700	
Salmon R. at N.Fk Ramp	ND	ND	SAW A	*AD		199,602	4.3	Production	1	251	167	677	0.34
4/18-20/90			TOTAL C	WT RELEASE		0			2	250	9		
				ION-CWT RELE SITE RELEASE	EASE	199,602 199,602		TOTAL RETU	JRN:	501	176	677	0.34
East Fork Salmon R.	ND	ND	DWOR B	*AD		64,150	3.8	Production	1	3	0	13	
4/11/90			TOTAL C	WT RELEASE		0			2	9 ND	1 ND		
			TOTAL N	ION-CWT RELE SITE RELEASE	EASE	64,150		TOTAL RETU	JRN:	12	1	13	0.02
TOTAL SAW A-STO						91,125							
TOTAL SAW A-STO			RELEASE			890,639 981,764	_	RETURN IS I	NCOMPL	ETE			
TOTAL DWOR B-S						42,322		RETURN					
TOTAL DWOR B-S TOTAL DWOR B-S				E		415,180 457,502	_	GRAND TOT	AL:	4,465	1,661	6,126	0.43
TOTAL CWT RELE	ASE F	OR HAGE	ERMAN NF	н		133,447							
TOTAL NON-CWT						1,305,819							
TOTAL HAGERMA	N NFH	RELEAS	Ε			1,439,266							

Appendix B. Table 3. Release data and estimated adult returns for Hagerman National Fish Hatchery summer steelhead trout, brood year 1990. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Hage	rman	Brood Y	ear: 1990		I	dentifying I	Marks						
Release Site/Date	RW NO.	RW Total	Stock ID	Mark Type	CWT Code	Release Number	Size (FPP)	Marking Purpose	Age Ocean	Comp	eturn position Hatchery	Total Returns	SAR (%)
Sawtooth Weir 4/13/91	71- 74	22,652	SAW A	CWT/LV/AD *LV/AD *AD	104333	21,050 213 1,389	2.9	Size at release Large size group	1 2	45 ND	24 ND	69	0.33
Sawtooth Weir 4/13/91	71- 74	21,580	SAW A	CWT/LV/AD *LV/AD *AD	104334	20,129 122 1,329	2.9	Size at release Large size group	1 2	55 ND	27 ND	82	0.41
Sawtooth Weir 4/15/91	74- 75	12,864	SAW A	CWT/LV/AD *AD	104335	12,066 798	2.6	Size at release Large size group	1 2	32 ND	12 ND	44	0.36
Sawtooth Weir 4/13/91	76- 78	326,644	SAW A	CWT/LV/AD *LV/AD *AD	104336	21,775 131 304,738	4.5	Size at release Regular size group	1 2	55 ND	13 ND	68	0.31
Sawtooth Weir 4/13/91	76- 78	305,400	SAW A	CWT/LV/AD *LV/AD *AD	104337	20,318 143 284,939	4.5	Size at release Regular size group	1 2	49 ND	15 ND	64	0.31
Sawtooth Weir 4/16/91	78- 80	290,660	SAW A	CWT/LV/AD *LV/AD *AD	104338	19,338 156 271,166	4.4	Size at release Regular size group	1 2	52 ND	19 ND	71	0.37
Sawtooth Weir 4/13-16/91	71- 80	985	SAW A	*AD/PIT		985 (489) (496)	ND	Size at release (Lg) Size at release (Re	1 2 g)	2 ND	1 ND	3	0.30

Appendix B. Table 3. Continued.

Appendix B. Tab						Identifying N	/larks		_				
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		eturn position	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean		Hatchery	Returns	(%)
Sawtooth Weir Includes all releas have Mark Type i			SAW A	NON-CWT (Includes	all *)	865,124	2.9	Production (Includes all *)	1 2	2,173 ND	682 ND	2,855	0.33
Sawtooth Weir 10/5-17/90	ND	ND	SAW A	*AD VT RELEASE		304,907 114.676	39.6	Fall release	1 2	0 ND	0 ND	0	0.00
			TOTAL NO	ON-CWT RELE TE RELEASE	EASE	1,170,031 1,284,707		TOTAL RETURN:		2,463	793	3,256	0.25
Little Salmon R. at Hazard Cr. 4/22/91	19- 21	154,937	DWOR B	CWT/LV/AD *LV/AD *AD	104332	19,831 384 134,722	4.4	Contribution	1 2 3	6 ND ND	6 ND ND	12	0.06
Little Salmon R. at Hazard Cr. 4/19/91	51	154,379	DWOR B	CWT/LV/AD *LV/AD *AD	104339	19,813 241 134,325	4.4	Contribution	1 2 3	4 ND ND	4 ND ND	8	0.04
Little Salmon R. at Hazard Cr. 4/17/91	38	147,794	DWOR B	CWT/LV/AD *LV/AD *AD	104340	18,877 554 128,363	4.5	Contribution	1 2 3	3 ND ND	3 ND ND	6	0.03
Little Salmon R. at Hazard Cr. 4/24/91	44 45	424	DWOR B	AD/PIT		424	4.5	Contribution	1 2 3	0 ND ND	0 ND ND	0	ND
Little Salmon R. a Includes all releas have Mark Type i	se date	s that	DWOR B	NON-CWT (Includes	all *)	398,589	4-4.5	Production (Includes all *)	1 2	88 ND	88 ND	176	0.04
nate man Type I	. I a locato	~ <i>~</i> ,		NT RELEASE ON-CWT RELE	ASF	58,521 518,912			3	ND	ND		
			-	TE RELEASE	.AUL	577,433		TOTAL RETURN:		128	128	256	0.04
E. Fk Salmon R.	ND		UNK. B	AD		540,733	32.8	Excess Fingerling	1	29	4	33	0.01

 (above trap)
 Plant
 2
 ND
 ND

 9/5-7/90
 3
 ND
 ND

Appendix B. Table 3. Continued.

Appendix B. Tub						dentifying I	Marks						
Release	RW	RW	Stock	Mark	сwт	Release	Size	Marking	Age	Comp	eturn position	Total	SAR
Site/Date	NO.	Total	<u>ID</u> _	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Little Salmon R. at Hazard Cr.	ND		DWOR B	AD		120,323	22.5	Excess Fingerling	1	27	27	54	0.04
								Plant	2	ND ND	ND ND		
				T RELEASE		0							
				N-CWT RELE E RELEASE	ASE	540,733 540,733		TOTAL RETURN:		29	4	33	0.01
TOTAL SAW A-S	STOCK	CWT DE	LEASE			114.676							
TOTAL SAW A-S						1,170,031							
TOTAL SAW A-S	ТОСК	RELEAS	E			1,284,707	_						
TOTAL DWOD D	OTOO	/ OWT D	FLEAGE			50 504		RETURN IS INCOM	/IPLETE				
TOTAL DWOR B				F		58,521 518,912		RETURN					
TOTAL DWOR B				_		577,433	=	GRAND TOTAL:		2,620	925	3,545	0.15
						·		(includes fingerling		•		,	
TOTAL UNK B-S						0		and fry releases)					
TOTAL UNK B-S			_			540,733 540,733	_						
TOTAL ONK B-3	, i ock i	\LLLA5	_			J 4 0,733							
TOTAL CWT RE				- -		173,197							
TOTAL NON-CW			_	N NFH		2,229,676	_						
TOTAL HAGERN	IAN NF	H KELEA	ASE			2,402,873							
TOTAL PIT TAG	S					1,409							
TOTAL SMOLT	RELEAS	SE				1,436,910		RETURN (smolt releases only	y)	2,564	894	3,458	0.24

Appendix C. Table 1. Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1988. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Magic	Valley FH		Brood Ye	ar: 1988									
				Ident	ifying Mar	ks	=,						
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		turn position	Total	SAR
Site/Date	<u>NO.</u>	Total	<u>ID</u>	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Sawtooth Weir 4/10-19/89	ND	ND	PAH A	*AD		854,462	4.44	Production	1 2	421 627	126 147	1,321	0.15
Sawtooth Weir	ND	ND	PAH A	*AD/PIT		2,832	4.44	Water budget	1	1	0	6	0.21
4/10-19/89			TOTAL (WT RELEASE		0		migration	2	4	1		
			TOTAL N	ION-CWT REL	EASE	857,294							
			TOTAL S	SITE RELEASE		857,294		TOTAL RET	URN:	1,053	274	1,327	0.15
Hazard Cr. (L.	ND	ND	PAH A	*AD		401,052	4.23	Production	1	79	79	730	0.18
Salmon R.) 4/19-26/89									2	286	286		
Hazard Cr. (L.	ND	ND	PAH A	CWT/LV/AD	104141	15,209	4.23	Contribution	1	0	0	30	0.20
Salmon R.) 4/19-26/89									2	15	15		
Hazard Cr. (L.	ND	ND	PAH A	CWT/LV/AD	104142	15,154	4.23	Contribution	1	3	3	24	0.16
Salmon R.)						-, -			•	•			
4/19-26/89									2	9	9		
Hazard Cr. (L.	ND	ND	PAH A	CWT/LV/AD	104143	15,927	4.23	Contribution	1	0	0	18	0.11
Salmon R.) 4/19-26/89									2	9	9		
Hazard Cr. (L. Salmon R.)	ND	ND	PAH A	*AD/PIT		3,058	4.23	Contribution	1	ND	ND	0	0.00

4/19-26/89 2 ND ND

Appendix C. Table 1. Continued.

				ldent	ifying Mar	ks							
									_		eturn		
Release Site/Date	RW NO.	RW Total	Stock ID	Mark Type	CWT Code	Release Number	Size (FPP)	Marking Purpose	Age Ocean		osition Hatchery	Total Returns	SAR (%)
			TOTAL N	CWT RELEASE NON-CWT RELI SITE RELEASE	EASE	46,290 404,110 450,400	-	TOTAL RET	URN:	401	401	802	0.18
East Fork Salmon	ND	ND	E FK B	*AD		306,306	4.48	Production	1	19	14	656	0.21
R. 4/15-19/89									2	572 0	51 0		
East Fork Salmon	ND	ND	E FK B	CWT/LV/AD	104144	15,624	4.48	Contribution	1	3	2	27	0.17
R. 4/15-19/89									2	19 0	3 0		
East Fork Salmon R.	ND	ND	E FK B	CWT/LV/AD	104145	14,126	4.48	Contribution	1	1	1	5	0.04
4/15-19/89									2 3	3 0	0 0		
East Fork Salmon R.	ND	ND	E FK B	CWT/LV/AD	104146	14,314	4.48	Contribution	1	1	1	10	0.07
4/15-19/89									2	8	0		
									3	0	0		
East Fork Salmon R.	ND	ND	E FK B	*AD/PIT		2,930	4.48	Migration survival	1	0	0	7	0.24
4/15-19/89								and timing	2	6	1		
				OWT RELEASE	- 4 6 5	44,064			3	0	0		
				NON-CWT RELE SITE RELEASE	EASE	309,236 353,300		TOTAL RET	URN:	632	73	705	0.20
Slate Cr. Lower Sal. R.	ND	ND	PAH A	*AD		300,600	4.14	Production	1	60	60	549	0.18
4/24-27/89			TOTAL (OWT RELEASE		0			2	214	215		

				ON-CWT REL TE RELEASE		300,600 300,600		TOTAL RET	URN:	274	275	549	0.18
Yankee Fork Salmon R.	ND	ND	PAH A	*AD		104,400	4.33	Production	1	80	24	199	0.19
4/17-21/89									2	77	18		
Appendix C. Table	e 1. Conti	nued.											
			_	Ident	ifying Ma	rks				D,	eturn		
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		position	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean		Hatchery	Returns	(%)
			TOTAL NO	WT RELEASE ON-CWT REL TE RELEASE	EASE	0 104,400 104,400		TOTAL RET	URN:	157	42	199	0.19
Hammer Cr. 4/28-29/89	ND	ND	PAH A	*AD		136,000	4.04	Production	1 2	27 97	27 97	248	0.18
			TOTAL NO	WT RELEASE ON-CWT REL TE RELEASE	EASE	0 136,000 136,000		TOTAL RET	URN:	124	124	248	0.18
TOTAL PAH A-ST TOTAL PAH A-ST TOTAL PAH A-ST	OCK NO	N-CWT I				46,290 1,802,404 1,848,694		RETURN IS CO	MPLETE				
TOTAL E FK B-ST TOTAL E FK B-ST TOTAL E FK B-ST	госк по	N-CWT				44,064 309,236 353,300		RETURN GRAND TOTAL	:	2,641	1,189	3,830	0.17
TOTAL CWT REL TOTAL NON-CWT TOTAL MAGIC VA	RELEAS	SE FOR	MAGIC VAL			90,354 2,111,640 2,201,994							
TOTAL PIT TAGS						5,988							

Appendix C. Table 2. Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1989. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Magic	: Valley	FH Broo	d Year: 198		ying Mark	(S							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Comp	eturn position	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
Sawtooth Weir	1	1,159,080	PAH A	CWT/LV/AD/FB	104059	39,620	4.2	Contribution, timing	1	56	14	123	0.31
4/12-13/90				FB = LA-) (-1 *LV/AD *AD		2,804 1,156,276	4.2		2	53	0		
Sawtooth Weir 4/12-13/90			PAH A	NON-CWT (Includes all *)		1,159,080		Production (Includes all *)	1 2	3,312 1,424	828 198	5,762	0.50
Includes all releas								,					
Mark Type indica	ted by *	•	TOTAL N	WT RELEASE ON-CWT RELEAS ITE RELEASE	SE	39,620 1,159,080 1,198,700		TOTAL RE	TURN:	4,845	1,040	5,885	0.49
E. F. Salmon R.	10	152,110	E. FK. B	CWT/LV/AD/FB	104058	40,905	4.1	Contribution, timing	1	11	4	96	0.23
4/14/90				FB = RA-)	(-1			J	2	66	15		
				*LV/AD *AD		837 111,205			3	ND	ND		
E. F. Salmon R. 4/16/90	16	651,914	E. FK. B	CWT/LV/AD *LV/AD *AD	104236	15,474 366 42,068	4.2	Contribution	1 2 3	13 20 ND	0 4 ND	37	0.24
E. F. Salmon R. 4/16/90	11	115,013	E. FK. B	CWT/LV/AD *LV/AD *AD	104237	15,971 378 43,419	4.2	Contribution	1 2 3	5 21 ND	1 3 ND	30	0.19

Appendix C. Table 2. Continued.

				Identi	fying Mark	S							
Release Site/Date	RW NO.	RW Total	Stock ID	Mark Type	CWT Code	Release Number	Size (FPP)	Marking Purpose	Age Ocean	Comp	eturn position Hatchery	Total Returns	SAR (%)
E. F. Salmon R. 4/16/90	11	115,013	E. FK. B	CWT/LV/AD *LV/AD *AD	104238	14,958 354 40,665	4.2	Contribution	1 2 3	0 21 ND	3 3 ND	27	0.18
E. F. Salmon R. 4/16-18/90	16	651,914	DWOR B	CWT/LV/AD *LV/AD *AD	104233	14,964 1,079 183,731	5	Contribution	1 2 3	0 2 ND	0 0 ND	2	0.01
E. F. Salmon R. 4/16-18/90	16	651,914	DWOR B	CWT/LV/AD *LV/AD *AD	104234	15,157 1,093 186,101	5	Contribution	1 2 3	0 58 ND	0 0 ND	58	0.38
E. F. Salmon R. 4/16-18/90	16	651,914	DWOR B	CWT/LV/AD *LV/AD *AD	104235	14,642 1,056 179,777	5	Contribution	1 2 3	2 9 ND	0 1 ND	12	0.08
E. F. Salmon R. 4/14-18/90			DWOR B E. FK. B	NON-CWT (Includes all *)	552,837 239,292	792,129	4.6	Production (Includes all *)	1 2	112 535	14 73	734	0.09
Includes all releas Mark Type indicat		that have						,	3	ND	ND		
			TOTAL NO	NT RELEASE DN-CWT RELEAS TE RELEASE	SE	132,071 792,129 924,200		TOTAL RE	ΓURN:	875	121	996	0.11
Slate Cr.	ND	ND	DWOR B	*AD		162,700	60.8	Excess Fingerling	1	0	0	0	0
Upper Salmon R. 9/11/89				VT RELEASE		0		Plant	2	0 ND	0 ND		
				ON-CWT RELEAS TE RELEASE	SE	162,700 162,700		TOTAL RE	ΓURN:	0	0	0	0

Appendix C. Table 2. Continued.

лирения от п				lden	tifying Marl	KS							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		turn position	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
TOTAL PAH A	-STOCK C	WT RELE	ASE			39,620							
TOTAL PAH A	-STOCK N	ON-CWT F	RELEASE			1,159,080							
TOTAL PAH A	-STOCK R	ELEASE				1,198,700							
						, ,		RETURN IS IN	COMPLE	ETE			
TOTAL DWOR	B-STOCK	CWT REL	EASE			44,763							
TOTAL DWOR	B-STOCK	NON-CW	T RELEASE			715,537		RETURN					
TOTAL DWOR	B-STOCK	RELEASE	<u> </u>			760,300	•	GRAND TO	TAL:	5,720	1,161	6,881	0.30
								(includes fing	gerling				
TOTAL E FK B	S-STOCK C	WT RELE	ASE			87,308		and fry release	ases)				
TOTAL E FK B	S-STOCK N	ION-CWT	RELEASE			239,292	-						
TOTAL E FK B	S-STOCK F	RELEASE				326,600							
TOTAL CWT R	ELEASE F	OR MAGI	C VALLEY FH	1		171,691							
TOTAL NON-C	WT RELE	ASE FOR I	MAGIC VALL	EY FH		2,113,909							
TOTAL MAGIC	VALLEY	FH RELEA	SE			2,285,600	•						
TOTAL SMOL	T RELEAS	E				2,122,900		RETURN (smolt release	es only)	5,720	1,161	6,881	0.32

Appendix C. Table 3. Release data and estimated adult returns for Magic Valley Fish Hatchery summer steelhead trout, brood year 1990. An asterisk (*) indicates that the fish were counted in the NON-CWT category for the specified release site. A number enclosed in parentheses indicates that the fish were counted in an above category. Return data were tabulated (if available) for each CWT code and NON-CWT group. Release information is summarized by CWT/NON-CWT and stock at bottom of table. ND indicates data were not available. The harvest component of the return composition includes only fish that were harvested in Idaho's sport fishery. The hatchery component of the return composition includes fish that returned to hatchery weirs and off-site escapement.

Hatchery: Magic	valley	TH BIO	ou rear. I		fying Ma	rks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		turn osition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
E. F. Salmon R. 4/13/91	3	209,893	DWOR B	CWT/LV/AD *LV/AD *AD	104314	20,498 2,027 187,368	4.1	Contribution	1 2 3	200 ND ND	1 ND ND	201	0.98
E. F. Salmon R. 4/13/91	2	214,470	DWOR B	CWT/LV/AD *LV/AD *AD	104315	21,017 2,079 191,374	4.1	Contribution	1 2 3	5 ND ND	1 ND ND	6	0.03
E. F. Salmon R. 4/13/91	3	207,987	DWOR B	CWT/LV/AD *LV/AD *AD	104316	20,312 2,008 185,667	4.1	Contribution	1 2 3	0 ND ND	0 ND ND	0	0.00
E. F. Salmon R. 4/15-16/91	15	113,570	E. FK. B.	CWT/LV/AD *LV/AD *AD	104320	22,525 530 90,515	4.3	Contribution	1 2 3	4 ND ND	3 ND ND	7	0.03
E. F. Salmon R. 4/15-16/91	14	112,609	E. FK. B.	CWT/LV/AD *LV/AD *AD	104321	22,483 529 89,597	4.3	Contribution	1 2 3	0 ND ND	3 ND ND	3	0.01
E. F. Salmon R. 4/15-16/91	15	107,771	E. FK. B.	CWT/LV/AD *LV/AD *AD	104322	21,375 503 85,893	4.3	Contribution	1 2 3	0 ND ND	1 ND ND	1	0.00
E. F. Salmon R.	2	1,500	DWOR B	AD/PIT		1,500		Migration survival	1	0	0	0	0.00
4/13, 15-16/91	14		E. FK. B.					and timing	2	ND	ND		

3 ND ND

Appendix C. Table 3. Continued.

	<u> </u>	Continued.	•	Identi	fying Mai	rks							
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age	Comp	turn osition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean	Harvest	Hatchery	Returns	(%)
E. F. Salmon R. Includes all release	date	s that	DWOR B E FK B	NON-CWT (Includes all *)	570,523 267,567	838,090		Production (Includes all *)	1 2	337 ND	46 ND	383	0.05
have Mark Type inc	dicate	d by *		•				,	3	ND	ND		
			TOTAL N	WT RELEASE ON-CWT REL ITE RELEASE	.EASE	128,210 839,590 967,800		TOTAL RE	TURN:	546	55	601	0.06
	_											_	
Hazard Cr. L. Salmon R. 4/26/91	8	102,541	PAH A	CWT/LV/AD *LV/AD *AD	104317	21,809 744 79,988	3.6	Contribution	1 2	4 ND	4 ND	8	0.04
Hazard Cr.	7	105.146	PAH A	CWT/LV/AD	104318	22.704	3.6	Contribution	1	21	21	42	0.18
L. Salmon R. 4/26/91	8	,		*LV/AD *AD		774 81,668			2	ND	ND		
Hazard Cr.	8	101,013	PAH A	CWT/LV/AD	104319	21,484	3.6	Contribution	1	16	16	32	0.15
L. Salmon R. 4/26/91				*LV/AD *AD		733 78,796			2	ND	ND		
Hazard Cr.	7	1,600	PAH A	AD/PIT		1,600	3.6	Migration survival	1	1	1	2	0.13
L. Salmon R. 4/26/91	8							and timing	2	ND	ND		
Hazard Cr.			PAH A	NON-C	WT	242,703	3.6	Production	1	151	151	302	0.12
L. Salmon R.				(Includes	all *)			(Includes all	2	ND	ND		
Includes all release have Mark Type inc								*)					
7 ,		· · '		WT RELEASE		65,997							
			_	ON-CWT REL ITE RELEASE	_	244,303 310,300		TOTAL RE	TURN:	193	193	386	0.12
Sawtooth Weir 4/9-19/91	ND	ND	PAH A	*AD		364,700	3.9	Production	1 2	916 ND	288 ND	1204	0.33

TOTAL CWT RELEASE **TOTAL NON-CWT RELEASE TOTAL SITE RELEASE**

0 364,700 364,700

TOTAL RETURN:

988

288

1276 0.35

Appendix C. Tabl	le 3. C	ontinued	•	11. 4									
			_	Identi	irks	_			D.	turn			
Release	RW	RW	Stock	Mark	CWT	Release	Size	Marking	Age		osition	Total	SAR
Site/Date	NO.	Total	ID	Туре	Code	Number	(FPP)	Purpose	Ocean		Hatchery		(%)
Pahsimeroi R. at hatchery	ND	ND	PAH A	*AD		135,100	3.8	Production	1	445	411	856	0.63
4/18-19/91			-	T RELEASE N-CWT REL		0 135,100			2	ND	ND		
			-	E RELEASE	_	135,100		TOTAL RE	TURN:	445	411	856	0.63
Salmon R. at Shoup Br.	ND	ND	PAH A	*AD		97,800	3.9	Production	1	214	6	220	0.22
4/20-21/91				T RELEASE	_	0			2	ND	ND		
			-	N-CWT REL E RELEASE	_	97,800 97,800		TOTAL RE	TURN:	214	6	220	0.22
Hammer Cr. 4/22-25/91	ND	ND	PAH A	*AD		186,300	3.9	Production	1 2	116 ND	116 ND	232	0.12
			TOTAL NO	T RELEASE N-CWT REL	EASE	0 186,300 186.300		TOTAL RE	TIIRN:	116	116	232	0.12
TOTAL PAH A-STOCK CWT RELEASE TOTAL PAH A-STOCK NON-CWT RELEASE TOTAL PAH A-STOCK RELEASE					_	65,997 1,028,203 1,094,200		TOTALKL	· Orare	110	110	202	0.12
TOTAL DWOR BOTOTAL DWOTA DWOR BOTOTAL DWOTA	STOC	K NON-0	WT RELEAS	SE		61,827 571,273 633,100							
TOTAL E FK B-S TOTAL E FK B-S TOTAL E FK B-S	TOCK	NON-CV	VT RELEASE	i.		66,383 268,317 334,700		RETURN IS II			4.000	0.574	0.47
TOTAL CWT REL						194,207 1,867,793		GRAND TO	JIAL:	2,502	1,069	3,571	0.17

TOTAL MAGIC VALLEY FH RELEASE	2,062,000
TOTAL PIT TAGS	3,100

Prepared by:	Approved by:
	IDAHO DEPARTMENT OF FISH AND GAME
T. Dean Rhine	Virgil K. Moore, Chief
Senior Fisheries Research Biologist	Bureau of Fisheries
Randall S. Osborne	Steve Yundt
Senior Fisheries Technician	Fishery Research Manager

Kristy A. Stevens Fisheries Biological-Aide