

The Structure and Substructure of Cold Dark Matter Halos

Julio Navarro
University of Victoria

The Virgo Consortium

CDM halos: Main results

- CDM mass profiles are nearly **universal**
 - shape is independent of mass
- CDM density profiles are **cuspy**
 - no evidence for a constant-density central “core”
- CDM halos are **clumpy**
 - Abundant but non-dominant substructure
- CDM halos are **triaxial**
 - Preference for prolate configuration, asphericity increasing toward the center.

CDM halos: Outstanding issues

- **The Structure of the Central Cusp**
 - Power-law divergent slope ($\rho \propto r^{-1}$ or $\rho \propto r^{-1.2}$ or $\rho \propto r^{-1.5}$?)
 - Annihilation signal
 - Disk galaxy rotation curves (cusp vs core vs triaxiality)
- **The Structure of Substructure**
 - Mass profile and abundance of Local Group satellites
 - Annihilation signal from substructures and “boost factors”
 - Abundance, spatial distribution and kinematics
 - lensing flux ratio anomaly, satellite distribution + orbits
- **The Phase-Space Distribution of Dark Matter**
 - Implications for direct dark matter detection experiments
- **The Origin of a Universal Density Profile**
 - Theoretical interest
 - Important to understand baryon-induced transformations of dark halo structure

The Aquarius programme

6 different galaxy size halos simulated at varying resolution, allowing for a proper assessment of **numerical convergence** and **cosmic variance**

Numerical resolution	Particle number in halo (N_{50})	# of substructures	mass resolution
Aq-A-5	808,479	299	$3.14 \times 10^6 M_{\odot}$
Aq-A-4	6,424,399	1,960	$3.92 \times 10^5 M_{\odot}$
Aq-A-3	51,391,468	13,854	$4.91 \times 10^4 M_{\odot}$
Aq-A-2	184,243,536	45,024	$1.37 \times 10^4 M_{\odot}$
Aq-A-1	1,473,568,512	297,791	$1.71 \times 10^3 M_{\odot}$ (15 pc/h softening)

Springel et al '08

“Via Lactea I simulation”







84,700,000	~10,000	$2.18 \times 10^4 M_{\odot}$
------------	---------	------------------------------

“Via Lactea II simulation”

470,000,000	~100,000	$3.92 \times 10^3 M_{\odot}$
-------------	----------	------------------------------


Diemand et al '07, 08

Pictures of all Aquarius halos (level-2 resolution)



Aquarius: the Billennium simulation

500 kpc



The Aquarius
“Billennium”
halo simulation.
A dark matter
halo with 1
billion particles
within the virial
radius.

[Play Movie](#)

$z = 1.5$

400^3 run



$z = 1.5$

1200³ run



$z = 1.5$

2400³ run



$z = 0.1$

2400^3 run



$z = 0.1$

2400³ run




$z = 0.1$


2400³ run



The Density Profile: numerical convergence




The Mass Profile: numerical convergence




- Excellent numerical convergence down to radius where the collisional relaxation time approaches the age of the universe

Self-similarity in the mass profile?




- Slight but significant **deviations from similarity**.
- A “third parameter” is needed in order to describe accurately the mass profiles of CDM halos.

Velocity structure: convergence





- Excellent numerical convergence down to radius where the collisional relaxation time approaches the age of the universe

Velocity structure: self-similarity?





- Slight but significant **deviations from similarity**.
- Note that deviant systems in mass are also deviant in velocity
- Note similarity in shape between density and velocity dispersion

The Structure of the Cusp




- Logarithmic slope scales like a power-law of radius: the Sersic/Einasto profile
- Innermost profile shallower than r^{-1}

The Cusp: Maximum Asymptotic Slope




- Maximum asymptotic slope of the cusp: shallower than r^{-1}

The ‘Phase-Space Density’ Profile



- Remarkably, the ‘phase-space density’, ρ/σ^3 , scales like a power law of radius
- This is the same dependence as in Bertschinger’s secondary infall similarity solution

The ‘Phase-Space Density’ Profile



- All halos seem to share the same “phase-space density”, ρ / σ^3 , structure
- This seems to reflect a fundamental structural property of CDM halos

A blueprint for detecting halo the CDM annihilation signal in the Galactic halo

Springel et al, 2008 Nature (Nov 8 issue)

CDM particles may annihilate and lead to production of γ -rays which could be observable by GLAST/FERMI

Emission of annihilation radiation depends on:

$$\int \rho^2(\mathbf{x}) \langle \sigma v \rangle dV$$

halo density at \mathbf{x} cross-section

⇒ Theoretical expectation requires knowing $\rho(\mathbf{x})$

⇒ Need accurate high resolution N-body simulations of halo formation from CDM initial conditions



Myths about Cold Dark Matter halo substructure and annihilation signal

- Halo DM is mostly in small (e.g. Earth mass) clumps
- Small (Earth-mass) clumps should dominate DM annihilation signal observable from Earth
- Dwarf spheroidals/luminous satellites are the best targets for detecting DM annihilation signal
- Halo DM is in a self-similar (fractal) distribution of nested substructure halos (subhalos)
- Annihilation signal/detectability is significantly boosted by sub-substructure

The mass function of substructures

The subhalo mass function is **shallower** than M^2

- Most of the substructure mass is in the few most massive halos
 - The total mass in substructures (5-50% of the total) converges well even for moderate resolution




The substructure circular velocity function

CUMULATIVE NUMBER OF SUBSTRUCTURES AS A FUNCTION OF VMAX,

We find 3 times as many subhalos as Diemand et al find for Via Lactea I

- Cosmic variance? - **No**
- Substructure finding algorithm? - **No**
- Different cosmological parameters? - **unlikely**


Nevertheless, the difference in our conclusions stems from different assumptions about visibility of clumps




The number density profile of substructure halos

- The spatial distribution of subhalos (except for the few most massive ones) is independent of mass
- Most subhalos are at large radii -- subhalos are more effectively destroyed near the centre
- Most subhalos have completed only a few orbits; dynamical friction unimportant below a subhalo mass threshold
- Subhalos are far from the Sun

$n(r)/\langle n \rangle$




$df_n(<r)/d\log r$



How lumpy is the MW halo?

Mass fraction in subhalos as a function of the free-streaming cutoff mass in the CDM power spectrum




Substructure mass fraction within $R_{\text{sun}} < 0.1\%$

Annihilation radiation from the Milky Way halo and subhalos

- If small-scale clumping and angular variations in the background may be neglected, then for systems with similar density profiles:
 1. Luminosity $\propto V_{\max}^4/r_{\max}$
 2. Flux $\propto V_{\max}^4/(r_{\max}/d^2)$
 3. Signal-to-noise $\propto V_{\max}^4/(r_{\max}^2/d)$
- The known substructure with largest signal-to-noise is the LMC, but it is easy to show that
 - $(S/N)_{\text{MW}}/(S/N)_{\text{LMC}} \sim 134!$
- Substructures are easier to detect than the main halo **only** if the “boost factor” from small-scale clumping overwhelms this simple scaling.


A blueprint for detecting halo CDM annihilation radiation

To calculate L need contribution from 4 components:



1. Smooth emission from main halo (**MainSmooth**)
2. Smooth emission from resolved subhalos (**SubSmooth**)
3. Emission from unresolved subhalos in main halo (**MainUnres**)
4. Emission from substructure of subhalos (**SubSub**)

The radiation from the main halo (MainSmooth)



- Lack of steep central cusp means that the radiation from the smooth main halo component is well defined and constrained.

- Half of the total luminosity comes from within ~ 3 kpc, 95% from ~ 30 kpc

- $L \sim V_{\max}^4 / r_{\max}$

The radiation from substructures (SubSmooth)



- This depends on being able to estimate accurately V_{\max} and r_{\max} for subhalos

- Convergence in the size and maximum circular velocity for individual subhalos **cross-matched** between simulation pairs.


- Largest simulation gives convergent results for:

$$\begin{aligned} V_{\max} &> 1.5 \text{ km/s} \\ r_{\max} &> 165 \text{ pc} \end{aligned}$$

- Much smaller than the halos inferred for even the **faintest** dwarf galaxies




Enclosed mass and annihilation radiation profiles




$> 10^5 M_{\odot}$
 $> 10^6 M_{\odot}$
 $> 10^7 M_{\odot}$
 $> 10^8 M_{\odot}$

Projected annihilation radiation profile



- Extrapolating to $M_{\text{sub}} = 10^{-6} M_{\text{sun}}$ yields $L_{\text{SUBSMOOTH}} \sim 200 L_{\text{MAINSMOOTH}}$
- This is what would be seen by a distant observer
- The total flux from SUBSMOOTH and MAINSMOOTH are actually similar for an observer near the Sun.


Substructures within substructures



There are substructures embedded within other structures. We detect 4 generations of nested subhalos.


The hierarchy clearly is **NOT self-similar** and is heavily dependent on the degree of tidal stripping of the subhalo

A “fractal” distribution of nested substructures?




Tidal effects on sub-substructures

Tidal radius



Substructures within substructures


- Cumulative number of sub-subhalos within subhalos
- Substructure mass fraction in subhalos is **lower** than in the main halo



Substructures within substructures

- Sub-substructure abundance in subhalos is NOT, in general, a scaled-down version of that in the main halo

because:




- substructure abundance reduced by tidal truncation
- sub-subs continue to loose mass through tides
- sub-subs not replenished by infall of fresh halos

⇒ Distribution of sub-substructure is NOT self-similar


Emission from substructure within substructures (SUBSUB)

- Compute tidal radius at instantaneous position (conservative!)
- Assume all material beyond r_t is removed
- Scale from main halo (within scaled r_t)
- Correct for luminosity below (scaled) mass limit





Detectability of substructure

- $S/N = F / (\theta_h^2 + \theta_{psf}^2)^{1/2}$
- S/N for detecting subhalos in units of that for detecting the main halo.
- 30 highest S/N objects, assuming use of optimal filters





- Highest S/N subhalos have 1% of S/N of main halo
- Highest S/N subhalos have 5-10 times S/N of known satellites
- Substructure of subhalos has no influence on detectability

Mass and circular velocity of most detectable substructure



- Highest S/N subhalos have masses well below those inferred for known Milky Way satellites

Distance and angular scale of most detectable substructures




- Highest S/N subhalos have half-light radii below 10 arcmin and will not be resolved by FERMI/GLAST

The gamma-ray sky lit by annihilation radiation

smooth main halo emission (MainSm)


MAINSMOOTH



The gamma-ray sky lit by annihilation radiation

emission from resolved subhalos (SubSm+SubSub)

SUBSMOOTH
+ SUBSUB




-3.0  2.0 Log(Intensity)

The gamma-ray sky lit by annihilation radiation

unresolved subhalo emission (MainUn)

MAINUNRES




-0.50  2.0 $\text{Log}(\text{Intensity})$

The gamma-ray sky lit by annihilation radiation

TOTAL

total emission



Myths about Cold Dark Matter halo substructure and annihilation signal

- ~~• Halo DM is mostly in small (e.g. Earth mass) clumps~~
- ~~• Small (Earth-mass) clumps should dominate DM annihilation signal observable from Earth~~
- ~~• Dwarf spheroidals/luminous satellites are the best targets for detecting DM annihilation signal~~
- ~~• Halo DM is in a self-similar (fractal) distribution of nested substructure halos (subhalos)~~
- ~~• Annihilation signal/detectability is significantly boosted by sub-substructure~~

Λ CDM on small scales

- Predictions for galactic dark matter in Λ CDM well established
- N-body simulations of Λ CDM predict:
 - many small substructures, with convergent mass fraction
 - the distribution of DM is not fractal nor is it dominated by Earth-mass objects
- γ -ray annihilation may be detectable by FERMI which should:
 - First detect smooth halo (if background can be subtracted)
 - Then (perhaps) detect dark subhalos with no stars
 - Sub-substructure boost irrelevant for detection

The End

© Cartoonbank.com



"Great PowerPoint, Kevin, but the answer is no."