

# Evaluation of the 2004 Predications of the Run-size and Passage Distributions of Adult Chinook Salmon (*Oncorhynchus tshawytscha*) Returning to the Columbia and Snake Rivers

Technical Report 2002 - 2003

January 2006

OE/BP-00005396-1



This Document should be cited as follows:

*Beer, W., David Salinger, Susannah Iltis, James Anderson, "Evaluation of the 2004 Predications of the Run-size and Passage Distributions of Adult Chinook Salmon (Oncorhynchus tshawytscha) Returning to the Columbia and Snake Rivers", 2002-2003 Technical Report, Project No. 198910800, 22 electronic pages, (BPA Report DOE/BP-00005396-1)*

Bonneville Power Administration  
P.O. Box 3621  
Portland, OR 97208

This report was funded by the Bonneville Power Administration (BPA), U.S. Department of Energy, as part of BPA's program to protect, mitigate, and enhance fish and wildlife affected by the development and operation of hydroelectric facilities on the Columbia River and its tributaries. The views in this report are the author's and do not necessarily represent the views of BPA.

**Evaluation of the 2004 Predictions of Run-size and  
Passage Distributions of Adult  
Chinook Salmon [*Oncorhynchus tshawytscha*]  
Returning to the Columbia and Snake Rivers**

Annual Report

January 2004 – December 2004

Prepared by:

W. Nicholas Beer  
David Salinger  
Susannah Iltis  
James J. Anderson

Columbia Basin Research  
School of Aquatic and Fishery Sciences  
University of Washington  
Box 358218  
Seattle, WA 98195

Prepared for:

United States Department of Energy  
Bonneville Power Administratiotn  
Division of Fish and Wildlife  
P.O. Box 3621  
Portland, OR 97208

Project No. 1989-108-00  
Contract No. 00005396

January 2006

## Executive Summary

This report is a postseason analysis of the accuracy of the 2004 predictions from Escapement Forecaster / Adult Upstream Model. The effectiveness of these modeling efforts are compared to observations of passage and river conditions at the end of the season. A pattern matching routine forecasts total run-size and run timing (daily passage) by optimally correlating the shape of the current year's cumulative passage (to date) with truncations of historical cumulative passage data. At the end of the season, for each stock at each observation site, we compute the Mean Absolute Deviation (MAD) for the day ( $j$ ) on which the prediction was made. MAD for both spring and fall chinook was always less than 16.4% (the Maximum Absolute Daily Deviation) and generally much less. (Mean Absolute Deviation = 3.7% and 4.8% respectively for the season).

## Table of Contents

Executive Summary .....	i
Table of Contents .....	i
List of Figures .....	ii
List of Tables .....	ii
Introduction .....	1
Methods .....	1
Data .....	1
Escapement and travel time data .....	1
Flow and other system operations data .....	1
Temperature data .....	2
Archives of model predictions .....	2
Models .....	3
Escapement Forecaster .....	3
Adult Upstream Model .....	3
Schematic of data and modeling .....	5
Postseason Assessment of Predictions .....	5
Results .....	6
Run-size .....	6
Run Timing .....	8
Discussion .....	10
Management Implications .....	10
Future Work .....	12
References .....	13
Appendix .....	14

## List of Figures

Figure 1 Schematic of data, models and products. Brown is used for historical data, green is realtime up-to-date information, white boxes are modeling processes and the yellow frames are final products. ....	5
Figure 2 Time series of Spring chinook run-size predictions for 2004. These were updated daily. ....	7
Figure 3 Time series of Fall chinook run-size predictions for 2004. These were updated daily.....	8
Figure 4 Spring chinook passage prediction at Bonneville for 2004. Note: This is a comparison of the predicted percentage passed versus the actual percentage passed on each day of the passage season. ....	9
Figure 5 Fall chinook passage prediction at Bonneville for 2004. Note: This is a comparison of the predicted percentage passed versus the actual percentage passed on each day of the passage season. ....	9
Figure 6 Passage of chinook at Bonneville Dam (2004). ....	11
Figure 7 Passage of chinook at Ice Harbor Dam (2004). ....	11
Figure 8 Spring Chinook observations and predictions made during 2004.....	14
Figure 9 Spring Chinook observations and predictions made during 2004.....	14
Figure 10 Spring Chinook observations and predictions made during 2004.....	15
Figure 11 Spring Chinook observations and predictions made during 2004.....	15

## List of Tables

Table 1 U.S. Army Corps of Engineers fixed monitoring sites and USGS gaging stations used for temperature forecasts.....	2
Table 2 Earliest day, after which the final run-size was within the error specified (10% 20%, or 30%) .....	6
Table 3 Pacific Fishery Management Council 2004 predictions and results of ocean escapement. Numbers in thousands of fish.....	10

## **Introduction**

Visual counts of returning adult chinook have been made at Bonneville Dam each year since 1938. The detection of adult chinook at Bonneville and upstream dams provides a measure of the temporal distribution of the returning adult salmonid populations.

The adult upstream "RealTime" forecaster/passage model was developed to predict the current season's adult salmon run-size at Bonneville Dam and run timing from the Bonneville Dam Tailrace to the upstream dams on the Columbia and Snake Rivers. The forecaster consists of an Escapement Forecaster (EF) that predicts the arrival timing and run-size of adult salmon at Bonneville Dam and an Adult Upstream Model (AUM) that predicts the passage timing of the fish at dams above Bonneville Dam. Each day the predictions are updated on the web.

During the 2001 migration season, Columbia Basin Research launched a prototype run timing system, EF /AUM, to predict run timing with results updated on the World Wide Web. This project was launched in an effort to provide real-time in-season projections of adult salmon migration to managers of the Columbia-Snake River hydrosystem to inform decisions about mitigation efforts such as in-river harvest. The program EF uses an empirical pattern matching routine to predict the arrival distributions for adult chinook salmon stocks at the first detection point in the migratory route, Bonneville Dam. The AUM model takes the predictions from EF and uses hydrological, fish behavioral and dam geometry information to simulate the movement of the adult salmonid through Columbia and Snake River dams.

This report is a postseason analysis of the accuracy of the 2004 predictions from EF/AUM. The effectiveness of these modeling efforts are compared to observations of passage and river conditions at the end of the season. The analyses and graphic presentations herein demonstrate the accuracy of the model throughout the season.

## **Methods**

### ***Data***

#### ***Escapement and travel time data***

The fish analyzed in this report are adult spring and fall chinook salmon returning to spawn in tributaries (or hatcheries) of the Columbia and Snake Rivers above Bonneville Dam. For the escapement forecasts, the daily visual counts of returning adult chinook data come from Bonneville Dam. To assess our upstream run timing predictions, the daily visual counts come from additional detection sites at McNary and Lower Granite Dams. Data is from Columbia River DART database, provided as a courtesy by U.S. Army Corps of Engineers, NWD (<http://www.nwd.usace.army.mil>).

#### ***Flow and other system operations data***

Any forecast of fish movement relies critically on accurate forecasts of flow, and other key system operations. The U.S. Army Corps of Engineers generates operational forecasts at all projects on the Columbia and Snake Rivers where there is fish passage. Water supply forecasts are based on a number of factors: the National Weather Service's Northwest River Forecast Center predictions, flood control requirements from the Army Corps, electrical power demand forecasts, and other criteria. The substantial uncertainty associated with springtime conditions often results in frequent and marked changes in these forecasts during April and May. Moreover, attempts to reduce the biological impacts of dissolved gas generated from high spill levels also results in a shifting of spill between projects within as well as outside the basin. Although the forecasts covered as much as 90 days into the future,

it must be recognized that their intended use was in deciding operations for the next week. Forecast accuracy beyond even a few days was itself uncertain. Bonneville Power Administration processes the Army Corps forecasts and makes them available to CBR staff throughout the migration season.

Forecasts for flow, spill, and elevation were replaced with observations on a daily basis with a query to the Columbia River DART database (CBR 2005a), which downloads water quality data from the Army Corps for the majority of monitoring sites in the Columbia Basin. This method was begun in 2001 and was a significant improvement over the 2000 in-season forecasts that relied on the forecasts alone. Subsequent fish arrival predictions were therefore based on the forecasted values for flow and spill and the latest available observed data.

### *Temperature data*

The temperature time series is a combination of year-to-date temperature data and forecasted temperatures. The forecasts are based on observed year-to-date temperature and flow data, historical average temperature and flow profiles for 15 locations in the Snake and Columbia rivers, and the flow forecasts. Historic and observed year-to-date data was obtained from the Columbia River DART database. Temperature predictions are made by applying a three-day moving window to fit predicted temperature time series to historical average patterns of temperature change; this method is described in detail in Beer et al. (2003).

**Table 1 U.S. Army Corps of Engineers fixed monitoring sites and USGS gaging stations used for temperature forecasts.**

Monitoring Locations	AUM Model Input Locations
Chief Joseph Forebay	Columbia Headwater
Wells Forebay	Methow Headwater
Rock Island Forebay	Wenatchee Headwater
The Dalles Forebay	Deschutes Headwater
Anatone, WA USGS	Snake Headwater
Peck, ID USGS	Clearwater Headwater
Peck, ID USGS	North Fork Clearwater Headwater
Peck, ID USGS	Middle Fork Clearwater Headwater
Anatone, WA USGS	Salmon Headwater
Wells Forebay	Wells Pool
Rocky Reach Forebay	Rocky Reach Pool
Rock Island Forebay	Rock Island Pool
Wanapum Forebay	Wanapum Pool
Priest Rapids Forebay	Priest Rapids Pool
Lower Granite Forebay	Lower Granite Pool
Little Goose Forebay	Little Goose Pool
Lower Monumental Forebay	Lower Monumental Pool
Ice Harbor Forebay	Ice Harbor Pool
McNary Forebay	McNary Pool
John Day Forebay	John Day Pool
The Dalles Forebay	The Dalles Pool
Bonneville Forebay	Bonneville Pool

### *Archives of model predictions*

The results of EF/AUM runs are stored on the Columbia Basin Research web site (CBR 2005b). Graphs based on the results are available through web-based query tools at

[http://www.cbr.washington.edu/crisprt/index\\_adult.html](http://www.cbr.washington.edu/crisprt/index_adult.html). Runs are made daily and include daily passage distribution forecasts and run-size forecasts.

## *Models*

### *Escapement Forecaster*

The Escapement Forecaster predicts the arrival timing and run-size of adult salmon at Bonneville Dam. It consists of a Bayesian-type algorithm with lognormal predictive density (Hyun 2002) and a prior distribution based in the early season on the previous year's jack counts and in the later season on a pattern matching algorithm. There is also a blending routine to switch smoothly between the jack-based and pattern match methods.

The jack-based routine predicts run-size by using a linear regression of each year's adult return vs. the previous year's jack return for the years 1982-2002. The correlation is strong for spring chinook, but less for fall chinook ( $r^2 = 0.92$  and  $0.22$  respectively). The arrival distribution is taken as the historic daily mean scaled to produce the correct total run-size.

The pattern matching routine forecasts total run-size and run timing (daily passage) by optimally correlating the shape of the current year's cumulative passage (to date) with truncations of historical cumulative passage data. This returns the fraction of the run complete,  $f$ . Total run-size is then predicted by  $\tilde{r} = P_c / f$  where  $P_c$  is the total passage (current year) to date.

To compare the current year's passage to that of historic runs, the cumulative current passage data is partitioned into  $N$  time intervals. The pattern matching optimization is performed as least-squares minimization; comparing slopes  $S_i^c$  over each subinterval  $i$  of the current run with slopes  $S_i^h(f)$  of subintervals of each historic year run truncated after  $f$  fraction of the historic run has passed. The optimization to determine  $f$  is then performed as:

$$\text{minimize}_{f \in (0,1)} \sum_{h \in H} \sum_{i=1}^N (S_i^c - S_i^h(f))^2$$

where  $H$  is the set of historical data years being used.

After the pattern matching method determines the completed fraction  $f$  of the current run, the passage forecast for each remaining day of the season is produced by appending the historic daily mean passage for each day of the final  $1 - f$  fraction of the season – scaled to produce the correct total run-size. In this way, the forecast may be a forward or backward shift in time as compared to the historic average, thereby forecasting not just run-size, but also run timing.

### *Adult Upstream Model*

The Adult Upstream Model (AUM) describes in detail fish movement through reaches and dams and the effects of various river operations on their migration. For in-season forecasts we use the projected escapement at Bonneville as input to AUM and predict the arrival timing at the upstream dams. The model contains a temperature and flow based submodel for reservoir passage and submodels for dam passage, fallback and straying. In addition, it includes a bioenergetic model to predict fish migration energy consumption. River flow and temperature are modeled using portions of the CRiSP smolt passage model. Fish travel time has been calibrated using PIT-tag data of adult chinook detected at both Bonneville and Lower Granite dams. A paper describing the theory and calibration of the model is in preparation (Salinger and Anderson, b). The temperature and flows encountered by upstream migrating salmon are the main factors determining reach migration speed and a submodel controls




this process. The flow encountered should subtract directly from the swimming speed in order to compute net up-river velocity. Because oxygen metabolism of chinook is optimal at about 17°C, the sustainable swimming speed is also optimal at about 17°C. To represent this, we use a broken linear model for the net up-river velocity  $V_M$  in terms of temperature  $\theta$  and flow  $F$ :

$$V_M = \begin{cases} \beta_0 + \beta_1\theta + \beta_3F, & \text{where } \theta \leq \tilde{\theta} \\ \beta_0 + \beta_1\tilde{\theta} + \beta_2(\theta - \tilde{\theta}) + \beta_3F, & \text{where } \theta > \tilde{\theta} \end{cases}$$

where  $\beta_0$ ,  $\beta_1$ ,  $\beta_2$  and  $\beta_3$  are the coefficients and  $\tilde{\theta}$  is the break point (approximately 17°C). In each reach, the travel time distribution is determined by the migration velocity  $V_M$  and by the rate of spreading  $V_{VAR}$  (Zabel and Anderson, 1997). Salinger and Anderson (a) more fully develops the net up-river velocity submodel. The migration velocity parameters and the spread parameter ( $V_{VAR}$ ) are determined from historical data using an optimization routine that compares model predicted passage distributions to observed ones. The arrival distributions were constructed from PIT-tag data of fish detected at both Bonneville and Lower Granite Dams in 1998-2002. These are combined into weekly cohorts with known travel time median and standard deviation. The cohorts create a *release* distribution at Bonneville Dam, and the model results are compared to the observations using least-squares optimization to pick the best parameterization of the model. Fall-back and dam delay are components contributing to the distribution of travel times for the fish.

### Schematic of data and modeling

The relationship of the data and models is depicted in Figure 1.



**Figure 1 Schematic of data, models and products. Brown is used for historical data, green is realtime up-to-date information, white boxes are modeling processes and the yellow frames are final products.**

### Postseason Assessment of Predictions

To assess the performance of run-size predictions, we compute the first day when the run-size estimate was within 10, 20 and 30% of the true run-size, and we determine what percent of the run had been completed on that day. Run size predictions are important for catch allocations, and compliance with federal and state regulations on fishery management. There is no established standard by which these predictions are evaluated.

To assess the performance of passage timing predictions, we apply the same measure used to assess RealTime/CRiSP predictions (Beer et al. 2003). For each stock at each observation site, we compute the Mean Absolute Deviation (MAD) for the day ( $j$ ) on which the prediction was made. This measure is based on the average deviation between predicted and observed cumulative passage on prediction dates during the season. MAD is computed as:

$$MAD_j = \frac{1}{N} \sum_{t=1}^N \left| F_{Day_t} - \hat{F}_{Day_j} \right| \times 100$$

where:

$j$  = forecast day on which  $MAD_j$  is calculated;

$t$  = index of prediction day (from 1 to  $N$ );

$N$  = number of days on which a prediction and observation were made for the stock at the site during the season;

$Day$  = vector of length  $N$  which identifies the days of the year from first observation of the stock at the site until two weeks past last observation (this is fixed for each site and each stock);

$F$  = observed cumulative passage on  $Day_t$ ; and

$\hat{F}$  = predicted cumulative passage on  $Day_t$ .

The MAD summation is performed over each of the dates on which model predictions were implemented – approximately every day during the season. This provides a snapshot of how well the model performs as the season progresses based on the final, “true” data. Ideally, there would be general decrease in MAD as  $t$  goes from 1 to  $N$  because the true distribution of the run should be better known and the true state of the flow and spill profiles should be known.

A second measure for run timing is the Maximum Absolute Daily Deviation (MADD)

$$MADD = \max \left\{ \left| F_{Day_t} - \hat{F}_{Day_t} \right| \times 100 \right.$$

All estimates of the run passage percentage are as good or better than this estimate.

## Results


### Run-size

EF predicted within 20% of the final Spring Chinook run-size on day 123 (Figure 2). EF finally predicted within 20% of the final Fall Chinook run-size on day 255 (Figure 3) Daily predictions for 2004 can be seen at [http://www.cbr.washington.edu/crisprt/index\\_adult.html](http://www.cbr.washington.edu/crisprt/index_adult.html). Alternative targets (10%, 20% and 30%) were each evaluated and are summarized in Table 2.


**Table 2 Earliest day, after which the final run-size was within the error specified (10% 20%, or 30%) .**

Stock	Within 10%	Within 20%	Within 30%
Spring chinook	124	123	123
Fall chinook	260	255	251

Big changes in the run-size predictions in Figure 2 and Figure 3 result from delays in the arrivals at Bonneville, since the forecaster becomes more and more sensitive to the data as the season progresses. This year, the early season prediction of Spring Chinook was high, and mid-run fluctuations in the run made the predictions unstable. The Spring Chinook run prior was on the order of 50% greater than the resulting run. By contrast, the Fall run prediction was 50% too low.



**Figure 2 Time series of Spring chinook run-size predictions for 2004. These were updated daily.**




**Figure 3 Time series of Fall chinook run-size predictions for 2004. These were updated daily.**

### *Run Timing*


The EF/AUM model is run daily and upstream passage predictions are saved until they can be compared to observations of passage at the end of the migration season. See the figures in the Appendix (beginning on page 14) for passage predictions on several days and the end-of-year observations.

We track the success of these predictions by comparing the estimated percentage passed on each day with the observed passage percentage. For the Spring run at Bonneville, the Mean Absolute Deviation (MAD) was 3.7%. In 2003 it was 6.7%. The worst prediction, the maximum absolute daily deviation (MADD) was 16.4% on day 114 (In 2003 it was 18.4% on day 103). A comparison of the predicted Bonneville Dam passage percentage for each day and final observations is in Figure 4.

For the Fall run at Bonneville, MAD was 4.8% (3.9% in 2003). MADD was 14.2% on day 254. In 2003, MADD was 19.3% on day 253. A comparison of the predicted Bonneville Dam passage percentage for each day and final observations is in Figure 5.



**Figure 4 Spring chinook passage prediction at Bonneville for 2004. Note: This is a comparison of the predicted percentage passed versus the actual percentage passed on each day of the passage season.**



**Figure 5 Fall chinook passage prediction at Bonneville for 2004. Note: This is a comparison of the predicted percentage passed versus the actual percentage passed on each day of the passage season.**

## Discussion

### *Management Implications*

Predictions of the passage at Bonneville shaped the forecasts of passage at other dams so that all the predictions are sensitive to these important first observations. These errors end up effecting upstream passage predictions. Despite unmodeled delays in migration the MAD for both spring and fall chinook was always less than 16.4% (the Maximum Absolute Daily Deviation) and generally much less. (Mean Absolute Deviation = 3.7% and 4.8% respectively for the season).

In February of 2004, Pacific Fishery Management Council (PFMC 2004) issued a preseason forecast of ocean escapement. Their predictions are the only other forecasts that exist and are intended for fishery management purposes. Although their predictions are different from ours, the overall numbers are an index of the escapement that could be expected in the river. Of course, we expect Bonneville passage to be less than the ocean escapement due to turnoffs and harvest. The PFMC Fall Chinook are divided into 5 distinct stocks compared to our one stock. Three of them (MCB, URB and SCH) pass Bonneville Dam and the other two are Lower River stocks. Subsequent referrals to the PFMC predictions of fall chinook will ignore the lower river fish. Their predictions and postseason analysis (PFMC 2005) are in Table 3. Their estimate of run-size was 88% of the observed fall chinook passage. There are no other estimates of stock run-size for chinook entering the Columbia.

**Table 3 Pacific Fishery Management Council 2004 predictions and results of ocean escapement. Numbers in thousands of fish**

Stock	Pre-season	Post-season <sup>a</sup>	Bonneville Passage <sup>b</sup>	Pre/Postseason	Escapement targets
Spring chinook	360.7	NA	170	NA	> 115 over Bonneville
Summer chinook	102.8	NA	92		> 80 over Bonneville
Spring Creek Hatchery	138.0	175.3		0.77	
Upriver Brights (Columbia)	292.2	367.9		0.79	> 45 over McNary
Mid-Columbia Brights	90.4	117.6		0.77	
Fall chinook total	510.6	NA	583		

<sup>a</sup> The river escapement includes observations of fish passing dams and the estimated catch from all sources.

<sup>b</sup> Bonneville passage of a stock is determined by date alone. Springs: Mar 1 – May 31. Summers: June 1 – July 31. Falls: Aug. 1 – Nov. 15.

The predictions at all dams for all stocks vary slightly with the prediction day as we would expect; however, the observations of Spring Chinook at IHR and LWG and other Snake River dams (see Appendix) obviously show skewed arrivals with most fish arriving early. The Columbia River and Snake River fish are disproportionately distributed and enumerated. First, Snake River Spring Chinook are more numerous than their Fall counterparts. Compare passage at Bonneville (Figure 6) and Ice Harbor (Figure 7) to illustrate this. Second, they migrate earlier than their Columbia River Spring Run counterparts having completed 50% passage of IHR on Day 122 while the general spring run at MCN, which all Columbia and Snake River must pass, occurred on the same day.




Figure 6 Passage of chinook at Bonneville Dam (2004).




Figure 7 Passage of chinook at Ice Harbor Dam (2004).



Distinguishing the passage of individual stocks through the river system could help resolve these predictions more carefully. Endangered stocks can be protected from over harvest and unfavorable migration conditions only if they can be located in space and time.

### *Future Work*

A summer stock passes Bonneville between the spring and fall runs. In subsequent years, we hope to evaluate the efficacy of using the calendar-based method of determining run type. This analysis was not available this year. It has certainly been a convenience in the past, but several changes might warrant revisiting this enumeration method because:

- Modern counting and identification procedures (PIT tags) are available that identify individual adult fish and therefore their stock before they arrive at their spawning grounds.
- Possible preferred migration conditions which are correlated to stock type can be identified from river conditions databases such as DART and may be related to cueing and therefore provide a mechanism for delineating the differences between the stocks.
- Any and all methods for determining the shape of arrival distributions would allow returning stocks to be partitioned according to those rules as they pass Bonneville. At present, a fish passing Bonneville on July 31 is a summer chinook and one passing on August 1 is fall chinook, and therefore treated distinctly.

During both the Spring and Fall chinook runs, there were significant periods during the migration season when passage was reduced or accelerated. A more detailed understanding of the mechanisms that create this phenomenon will improve the forecasts since the pattern matching algorithm assumes that reductions in passage indicate an end to the run. It would be valuable to have mechanistic explanations for these anomalies in otherwise comparatively smooth return results and tailor the pattern matching algorithms to be less sensitive to short term noise in the return data while still being responsive to real-time daily conditions.

Identification of specific stocks via PIT-tag detections could provide higher resolution passage data during autumn 2004 and subsequent years. For example, it could help to distinguish the Snake and Columbia River stocks which are currently in-separable in the visual counts at lower dams. This will be very helpful in predicting the upstream passage movements since the destinations of individuals and therefore proportions of the daily observations will be known.

Daily run-size predictions also include confidence intervals; however, these bounds are not related to other predictions made with our system EF/AUM. Currently, the confidence bounds are based on the Bayesian posterior distribution.

## References

- Beer, W.N., S. Iltis, C. VanHolmes, and J.J. Anderson. 2003. Evaluation of the 2002 Predictions of the Run-Timing of Wild Migrant Yearling Chinook and Water Quality at Multiple Locations on the Snake and Columbia Rivers using CRiSP/RealTime. Columbia Basin Research, School of Aquatic and Fishery Sciences. University of Washington, Box 358218 Seattle, WA 98195.
- Columbia Basin Research (CBR). 2005a. Data Access in RealTime (DART) Available on-line 14 February 2005 at <http://www.cbr.washington.edu/dart/>.
- \_\_\_\_\_. 2005b. DART Adult Passage Composite Report. Available on-line 14 February 2005 at <http://www.cbr.washington.edu/dart/adult.html>.
- Hyun, S.Y. 2002. Inseason forecasts of sockeye salmon returning to the Bristol Bay districts of Alaska. Ph.D. dissertation. University of Washington, Seattle, WA. 167pp.
- Salinger, D.H. and J.J. Anderson. a. Under review. Effects of temperature and flow on adult salmon migration speed.
- Salinger, D.H. and J.J. Anderson. b. In Preparation. A mechanistic model for Columbia River adult salmon upriver migration.
- PFMC (Pacific Fishery Management Council) 2004. Preseason Report I. Stock Abundance Analysis for 2004 Ocean Salmon Fisheries. Available On-line 14 February 2005 at <http://www.pcouncil.org/salmon/salpreI04/salpre104.pdf>
- PFMC (Pacific Fishery Management Council) 2005. Preseason Report I. Stock Abundance Analysis for 2005 Ocean Salmon Fisheries: Available On-line 27 January 2006 at <http://www.pcouncil.org/salmon/salpreI05/salpre105.pdf>
- Zabel, R. and J.J. Anderson. 1997. A model of the travel time of migrating juvenile salmon, with an application to Snake River Sprint Chinook. N. Amer. J. Fish. Manag. 17:93-100.

## Appendix

The graphs that follow depict the observations of passage for a stock at the dam stipulated by the three letter abbreviation. Predictions made on various dates are drawn in different colors (not available in all distributions) and different line styles (not available in all distributions) and different line styles

### Spring chinook at BON




Figure 8 Spring Chinook observations and predictions made during 2004.

### Spring chinook at MCN




Figure 9 Spring Chinook observations and predictions made during 2004.

### Spring chinook at IHR




Figure 10 Spring Chinook observations and predictions made during 2004.

### Spring chinook at LWG






Figure 11 Spring Chinook observations and predictions made during 2004.


### Fall chinook at BON



### Fall chinook at MCN



### Fall chinook at IHR



### Fall chinook at LWG

