Fermi National Accelerator Laboratory

FERMILAB-Pub-90/161-E
[E-741/CDF]

91-1-31

A Measurement of the W-Boson Mass *

The CDF Collaboration
Fermi National Accelerator Laboratory
P.O. Box 500
Batavia, Illinois 60510

August 2, 1990

* Submitted to Phys. Rev. Lett.

Operated by Universities Research Association Inc. under contract with the United States Department of Energy

A Measurement of the W-Boson Mass

August 2, 1990

F. Abe,[8] D. Amidei,[4] G. Apollinari,[11] M. Atac,[4] P. Auchincloss,[14] A. R. Baden,[6] A. Bamberger,[4],[a] A. Barbaro-Galtieri,[9] V. E. Barnes,[12] F. Bedeschi,[11] S. Behrends,[2] S. Belforte,[11] G. Bellettini,[11] J. Bellinger,[18] J. Bensinger,[2] A. Beretvas,[4] J. P. Berge,[4] S. Bertolucci,[5] S. Bhadra,[7] M. Binkley,[4] R. Blair,[1] C. Blocker,[2] A. W. Booth,[4] G. Brandenburg,[6] D. Brown,[6] E. Buckley,[14] A. Byon,[12] K. L. Byrum,[18] C. Campagnari,[3] M. Campbell,[3] R. Carey,[6] W. Carithers,[9] D. Carlsmith,[18] J. T. Carroll,[4] R. Cashmore,[4],[a] F. Cervelli,[11] K. Chadwick,[4] G. Chiarelli,[5] W. Chinowsky,[9] S. Cihangir,[4] A. G. Clark,[4] D. Connor,[10] M. Contreras,[2] J. Cooper,[4] M. Cordelli,[5] D. Crane,[4] M. Curatolo,[5] C. Day,[4] S. Dell'Agnello,[11] M. Dell'Orso,[11] L. Demortier,[2] P. F. Derwent,[3] T. Devlin,[14] D. DiBitonto,[15] R. B. Drucker,[9] J. E. Elias,[4] R. Ely,[9] S. Errede,[7] B. Esposito,[5] B. Flaugher,[14] G. W. Foster,[4] M. Franklin,[6] J. Freeman,[4] H. Frisch,[3] Y. Fukui,[8] Y. Funayama,[16] A. F. Garfinkel,[12] A. Gauthier,[7] S. Geer,[6] P. Giannetti,[11] N. Giokaris,[13] P. Giromini,[5] L. Gladney,[10] M. Gold,[9] K. Goulianos,[13] H. Grassmann,[11] C. Grosso-Pilcher,[3] C. Haber,[9] S. R. Hahn,[4] R. Handler,[18] K. Hara,[16] R. M. Harris,[9] J. Hauser,[3] T. Hessing,[15] R. Hollebeek,[10] L. Holloway,[7] P. Hu,[14] B. Hubbard,[9] B. T. Huffman,[12] R. Hughes,[10] P. Hurst,[7] J. Huth,[4] M. Incagli,[11] T. Ino,[16] H. Iso,[16] H. Jensen,[4] C. P. Jessop,[6] R. P. Johnson,[4] U. Joshi,[4] R. W. Kadel,[4] T. Kamon,[15] S. Kanda,[16] D. A. Kardelis,[7] I. Karliner,[7] E. Kearns,[6] R. Kephart,[4] P. Kesten,[2] R. M. Keup,[7] H. Keutelian,[7] S. Kim,[16] L. Kirsch,[2] K. Kondo,[16] S. E. Kuhlmann,[1] E. Kuns,[14] A. T. Laasanen,[12] J. I. Lamoureux,[18] W. Li,[1] T. M. Liss,[7] N. Lockyer,[10] C. B. Luchini,[7] P. Maas,[4] M. Mangano[11] J. P. Marriner,[4] R. Markeloff,[18] L. A. Markosky,[18] R. Mattingly,[2] P. McIntyre,[15] A. Menzione,[11] T. Meyer,[15] S. Mikamo,[8] M. Miller,[3] T. Mimashi,[16] S. Miscetti,[5] M. Mishina,[8] S. Miyashita,[16] Y. Morita,[16] S. Moulding,[2] A. Mukherjee,[4] L. F. Nakae,[2] I. Nakano,[16] C. Nelson,[4] C. Newman-Holmes,[4] J. S. T. Ng,[6] M. Ninomiya,[16] L. Nodulman,[1] S. Ogawa,[16] R. Paoletti,[11] A. Para,[4] E. Pare,[6] J. Patrick,[4] T. J. Phillips,[6] R. Plunkett,[4] L. Pondrom,[18] J. Proudfoot,[1] G. Punzi,[11] D. Quarrie,[4] K. Ragan,[10] G. Redlinger,[3] J. Rhoades,[18] M. Roach,[17] F. Rimondi,[4],[a] L. Ristori,[11] T. Rohaly,[10] A. Roodman,[3] D. Saltzberg, [3] A. Sansoni,[5] R. D. Sard,[7] A. Savoy-Navarro,[4] V. Scarpine,[7] P. Schlabach,[7] E. E. Schmidt,[4] M. H. Schub,[12] R. Schwitters,[6] A. Scribano,[11] S. Segler,[4] Y. Seiya,[16] M. Sekiguchi,[16] P. Sestini,[11] M. Shapiro,[6] M. Sheaff,[18] M. Shochet,[3] J. Siegrist,[9] P. Sinervo,[10] J. Skarha,[18] K. Sliwa,[17] D. A. Smith,[11] F. D. Snider,[3] R. St. Denis,[6] A. Stefanini,[11] R. L. Swartz, Jr.,[7] M. Takano,[16] K. Takikawa,[16] S. Tarem,[2] D. Theriot,[4] M. Timko,[15] P. Tipton,[9] S. Tkaczyk,[4] A. Tollestrup,[4] G. Tonelli,[11] J. Tonnison,[12] W. Trischuk,[6] Y. Tsay,[3] F. Ukegawa,[16] D. Underwood,[1] R. Vidal,[4] R. G. Wagner,[1] R. L. Wagner,[4]

Submitted to Phys. Rev. Lett., August 10, 1990.

J. Walsh,[10] T. Watts,[14] R. Webb,[15] C. Wendt,[18] W. C. Wester, III,[9] T. Westhusing,[11] S. N. White,[13] A. B. Wicklund,[1] H. H. Williams,[10] B. L. Winer,[9] A. Yagil,[4] A. Yamashita,[16] K. Yasuoka,[16] G. P. Yeh,[4] J. Yoh,[4] M. Yokoyama,[16] J. C. Yun,[4] F. Zetti[11]

[1] *Argonne National Laboratory, Argonne, Illinois 60439*

[2] *Brandeis University, Waltham, Massachusetts 02254*

[3] *University of Chicago, Chicago, Illinois 60637*

[4] *Fermi National Accelerator Laboratory, Batavia, Illinois 60510*

[5] *Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, Frascati, Italy*

[6] *Harvard University, Cambridge, Massachusetts 02138*

[7] *University of Illinois, Urbana, Illinois 61801*

[8] *National Laboratory for High Energy Physics (KEK), Tsukuba, Ibaraki 305, Japan*

[9] *Lawrence Berkeley Laboratory, Berkeley, California 94720*

[10] *University of Pennsylvania, Philadelphia, Pennsylvania 19104*

[11] *Istituto Nazionale di Fisica Nucleare, University and Scuola Normale Superiore of Pisa, I-56100 Pisa, Italy*

[12] *Purdue University, West Lafayette, Indiana 47907*

[13] *Rockefeller University, New York, New York 10021*

[14] *Rutgers University, Piscataway, New Jersey 08854*

[15] *Texas A&M University, College Station, Texas 77843*

[16] *University of Tsukuba, Tsukuba, Ibaraki 305, Japan*

[17] *Tufts University, Medford, Massachusetts 02155*

[18] *University of Wisconsin, Madison, Wisconsin 53706*

Abstract

We have determined $m_W = 79.91 \pm 0.39$ GeV/c^2 from an analysis of W$\rightarrow e\nu$ and W$\rightarrow \mu\nu$ data from the CDF detector in $\bar{p}p$ collisions at $\sqrt{s} = 1.8$ TeV. This result, together with the world average Z mass, determines the weak mixing angle to be $\sin^2\theta_W = 0.232 \pm 0.008$. Bounds on the top quark mass are discussed.

PACS numbers: 13.85Qk, 14.80Er.

The masses of the W and Z vector bosons are fundamental parameters in the Standard Electroweak Model [1, 2, 3]. Together, they determine the weak mixing angle through its definition [4, 5], $\sin^2\theta_W \equiv 1 - m_W^2/m_Z^2$, and give an upper limit on the mass of the, as yet, unobserved top quark. The measured value of the W mass reported here is based on a sample of 1130 W$\rightarrow e\nu$ and 592 W$\rightarrow \mu\nu$ candidate events in the Collider Detector at Fermilab (CDF) from integrated luminosities of 4.4 pb^{-1} and 3.9 pb^{-1} respectively in

$\bar{p}p$ collisions at $\sqrt{s} = 1.8$ TeV. These data were taken during the 1988-1989 running period of the Tevatron Collider. Details of the W mass analysis, summarized here, may be found in Ref. [6].

The components of the CDF relevant for this analysis are described briefly here. A detailed description of the detector may be found in Ref. [7]. Charged tracks are measured with Vertex Time Projection Chambers (VTPC) and a Central Tracking Chamber (CTC) in a 1.4116 Tesla solenoidal magnetic field. Scintillator-based electromagnetic (EM) and hadronic (HAD) calorimeters in the central region, pseudorapidity $|\eta| < 1.1$, are arranged in a projective tower geometry with cell sizes of $\Delta\eta \times \Delta\phi = 0.1 \times 15°$. Muon drift chambers reside behind the calorimeters in the region $|\eta| < 0.63$. Gas-based calorimeters are used in the region $1.1 < |\eta| < 3.6$.

Charged particle momenta are determined in the CTC with an rms resolution of $\delta p_T/p_T = 0.0011\, p_T$ (p_T in GeV/c). An overall momentum scale uncertainty of 0.1% is determined from an analysis of muon pairs in J/ψ and Υ candidates. Electron transverse energies are measured with an accuracy of $\delta E_T/E_T = [(0.135/\sqrt{E_T})^2 + (0.020)^2]^{1/2}$ where E_T is in GeV. The cell-to-cell relative normalization of the EM calorimeters is obtained by analysing a large sample of inclusive electron events. The overall energy scale is normalized with an acuracy of 0.24% to the CTC momentum scale by fitting the energy to momentum ratio, E/p, of a sample of W electrons. [6]

The trigger for the electron sample required at least 12 GeV transverse electromagnetic energy in the central calorimeter, associated with a track in the CTC of transverse momentum $p_T > 6$ GeV/c. The muon trigger required a track stub in the muon drift chambers behind the central calorimeter modules, matched to a CTC track of $p_T > 9$ GeV/c. Electrons are restricted to the region $|\eta| < 1.0$ and muons to $|\eta| < 0.6$. The trigger is fully efficient in the kinematic range of interest.

The transverse momentum of the neutrino, p_T^ν, is inferred from the vector imbalance of the calorimeter E_T and charged lepton momentum. We do this by constructing the vector

$$\vec{u} = \sum_i E_i \sin\theta_i \, \hat{n}_i,$$

where E_i is the total (electromagnetic plus hadronic) energy in the i^{th} tower. The polar angle θ_i and the unit vector in the transverse plane $\hat{n}_i$ are calculated using the event vertex and the center of the tower. The cells containing the charged lepton energy are not included in the sum. The vector $\vec{u}$ and the charged lepton momentum, p_T^ℓ, determine the neutrino transverse momentum $\vec{p}_T^{\,\nu} = -k_u\,\vec{u} - \vec{p}_T^{\,\ell}$. The factor $k_u = 1.4$ multiplying $\vec{u}$ scales the calorimeter low energy response to that of the charged leptons [8]. The resolution

for each component of $\vec{u}$, determined from studies of minimum bias events, is $\sigma_u = 0.47\sqrt{\sum E_T}$ where $\sum E_T$ is the total, uncorrected, scalar E_T in the calorimeter, not including the charged leptons.

The event samples used to determine the W mass require $p_T^\ell > 25$ GeV/c and $p_T^\nu > 25$ GeV/c. Events were removed if any cluster [8] of raw calorimeter transverse energy greater than 5 GeV was within $\pm$ 30 degrees opposite in azimuth to the lepton. To minimize the impact of the resolution in the $\vec{u}$ measurement, we required no cluster anywhere in the calorimeter above 7 GeV transverse energy other than that containing the electron. To avoid mismeasured Z decays, events with any track above 15 GeV/c p_T other than the lepton track were eliminated from the samples. For the muon sample, the cosmic ray background was reduced by requiring no track with p_T above 10 GeV/c within $\pm$ 3 degrees opposite the muon and no other stub in the muon drift chambers consistent with a cosmic ray. A match consistent with multiple scattering was required between the central track and the muon stub. The electron was required to be within the calibrated fiducial region of the central EM calorimeter [9], to have E/p < 1.4 and to be inconsistent with a photon conversion. The final samples contain 1130 electron and 592 muon candidates.

The W mass is obtained from a maximum-likelihood fit of the transverse mass distributions to simulation predictions. The transverse mass is defined as $m_T = \sqrt{2p_T^\ell p_T^\nu\,(\,1-\cos\phi_{\ell\nu}\,)}$. The predictions are an interpolation of a grid in mass and width generated by Monte Carlo. There is no systematic offset attributable to the fitting procedure. When the width of the W is not constrained, there is a 20-40% correlation between the W width (or equivalently the detector resolution) and the W mass. We constrained the width to $\Gamma = 2.1$ GeV to remove some of this sensitivity.

The Monte Carlo model includes dynamics of W production and decay as well as detector response. The model assumes that the background underlying a W event is the sum of a cylindrically symmetric event, with a detector resolution determined with minimum-bias triggered events, plus energy flow balancing the transverse momentum of the W. The calorimeter response to the energy flow as well as its resolution is determined from a sample of Z events where the recoil energy flow is well determined from measurements of the two charged leptons. Using the resolution parameters, an input p_T^W distribution was chosen such that the observed p_T^W distribution was returned by the model, as shown in Fig. 1. Independent variation of each parameter indicated uncertainties in the W mass of 70 MeV/c^2 due to uncertainties in the electron energy resolution, and 80 MeV/c^2 from the uncertainties in the muon resolution. An additional uncertainty of 130 MeV/c^2 due to resolution modeling is common to both samples.

Uncertainty	Electrons	Muons	Common
STATISTICAL	350	530	
ENERGY SCALE	190	80	80
1. Tracking chamber	80	80	80
2. Calorimeter	175		
SYSTEMATICS	240	315	150
1. Proton structure	60	60	60
2. Resolution, W p_T	145	150	130
3. Parallel balance	170	240	
4. Background	50	110	
5. Fitting	50	50	50
OVERALL	465	620	

Table 1: Uncertainties, in units of MeV/c^2, in the W mass measurement. The uncertainties which are the same for both samples are listed as common.

A variety of parton distribution functions [10, 11, 12, 13] have been used to determine a possible bias in assumptions about the distribution in longitudinal momentum. The variations in the fitted W mass are of the order of 60 MeV/c^2. We use the MRSB set [13] as the nominal set and assign an associated systematic uncertainty on the mass of the W of 60 MeV/c^2.

As the Monte Carlo does not simulate all details of the component of $\vec{u}$ parallel to the charged lepton, $u_{\parallel}$, a constant offset, $u_{\parallel}^0$, is introduced to match the average value of the data. An accurate determination of $u_{\parallel}^0$ is important since its value enters directly into the calculation of $\not\!\!p_T$ and m_T. We use events with transverse mass above 50 GeV/c^2 to determine $u_{\parallel}^0$. The uncertainty in $u_{\parallel}^0$ leads to a systematic uncertainty in m_W of 170 and 240 MeV/c^2 for the electron and muon samples respectively.

Residual backgrounds in the electron sample are less than 1%. The rates from τ sequential decays are negligible. A 1% residual flat background due to cosmic rays is possible in the muon sample. There is a small ($< 4\%$) background in the muon sample due to Z's with a missing second track, but these events tend to have large rapidities and yield relatively soft leptons. We conclude 50 and 110 MeV/c^2 are the uncertainties in the W mass due to background in the electron and muon samples. Table 1 summarizes the uncertainties in our measurement.

The observed and fitted transverse mass distributions are shown in Fig. 2. The fitting range in m_T is 65-94 GeV/c^2. Corrected for wide angle radiative effects [14] (70 and 125 MeV/c^2 for the electron and muon samples), the results are $m_W^e = 79.91 \pm 0.35$ (stat.) ± 0.24 (sys.) ± 0.19 (scale) GeV/c^2

and $m_W^\mu = 79.90 \pm 0.53$ (stat.) ± 0.32 (sys.) ± 0.08 (scale) GeV/c^2. We have checked sensitivity to cutoffs, subdivided the samples, varied the selection, not constrained the width, and fit the transverse momenta of electrons, muons and neutrinos and find no evidence for additional systematic uncertainty. We have confirmed the statistical precision using multiple Monte Carlo samples of the size of the data. The combined result is $m_W = 79.91 \pm 0.39$ GeV/c^2, consistent with previous measurements [15, 16]. A division of the data into positively and negatively charged W's yields $m_W^+ - m_W^- = -0.19 \pm 0.58$ GeV/c^2, consistent with CPT invariance.

In order to determine the weak mixing angle we combine the W mass values from the electron and muon decays with the world average Z mass of 91.161 [17] to obtain $\sin^2\theta_W = 0.2317 \pm 0.0075$. Fig. 3 shows the relationship between the top quark mass and $\sin^2\theta_W$ with the Z mass constrained to the LEP value [18, 19, 20, 21]. For a Higgs mass lighter than 1000 GeV/c^2 the top quark mass is constrained, within the context of the minimal Standard Model, to be $m_{top} < 220$ GeV/c^2 (95% CL). Combining our value with that of UA2[16] yields $\sin^2\theta_W = 0.227 \pm 0.006$.

We thank the Fermilab Accelerator Division for their exceptional performance in the operation of the Tevatron and the Antiproton Source. This work was supported in part by the Department of Energy, the National Science Foundation, Istituto Nazionale di Fisica Nucleare, the Ministry of Science Culture, and Education of Japan, and the A. P. Sloan Foundation.

[a] Visitor.

[1] S.L.Glashow. *Nucl. Phys.*, 22:579, 1961.

[2] S. Weinberg. *Phys. Rev. Lett.*, 19(21):1264, 1967.

[3] A.Salam. page 367. Almqvist & Wiksell, Sweden, 1968.

[4] A. Sirlin. *Phys. Rev.*, D22(4):971, 1980.

[5] W. Marciano and A. Sirlin. *Phys. Rev.*, D22(11):2695, 1980.

[6] F. Abe *et al.* (CDF Collaboration). Submitted to *Phys. Rev.*

[7] F. Abe *et al.* (CDF Collaboration). *Nucl. Instrum. and Methods*, A(271):387, 1988.

[8] F. Abe *et al.* (CDF Collaboration). *Phys. Rev. Lett.*, 62(6):613, 1988. This reference provides a description of how we cluster energy into jet-like objects.

[9] F. Abe *et al.* (CDF Collaboration). *Phys. Rev. Lett.*, 63(7):720, 1989.

[10] E. Eichten, I. Hinchliffe, K. Lane, and C. Quigg. *Reviews of Modern Physics*, 56:579, 1984.

[11] D. Duke and J. F. Owens. *Phys. Rev.*, D(30):49, 1984.

[12] M.Diemos, F.Ferroni, E.Longo, and G.Martinelli. *Z. Phys.*, C(39):21, 1988.

[13] A.D. Martin, R.G. Robert, and W.J. Stirling. *Phys. Rev.*, D(37):1161, 1988.

[14] R. G. Wagner. Unpublished, based on calculations by F. Berends. *et al.*, Z. Phys. C (27):155, 1985; F. Behrends and R. Kleiss, Z. Phys. C (27):365, 1985.

[15] G. Arnison *et al.* (UA1 collaboration). *EuroPhys. Lett.*, 1(7):327, 1986. $Mw = 83.5^{+1.1}_{-1.0} \pm 2.7\,\mathrm{GeV/c^2}$.

[16] J. Alitti *et al.* (UA2 Collaboration). *Phys. Lett*, B(241):150, 1990. $Mw = 80.49 \pm 0.43 \pm 0.24\,\mathrm{GeV/c^2}$.

[17] The Particle Data Group. *Physics Letters*, B(239), 1990.

[18] B. Adeva *et. al. Phys. Lett.*, B(231):509, 1989.

[19] D. Decamp *et. al. Phys. Lett.*, B(231):519, 1989.

[20] M. Akrawy *et. al. Phys. Lett.*, B(231):530, 1989.

[21] P. Aarnio *et. al. Phys. Lett.*, B(231):539. 1989.

[22] We wish to thank Duncan Morris for the use of his computer program in generating figure 3. See also Hollik *et al* . Technical Report 88-188, DESY, 1988. (Unpublished).

[23] K. Sliwa. In *Proceedings of the XXVth Recontres de Moriond*, March 1990.



Figure 1: The observed p_T^W distribution. The curve is the prediction of the Monte Carlo simulation. We note that this is not the true p_T^W distribution because of data selection cuts and energy corrections.



Figure 2: a) The transverse mass distribution for W $\rightarrow$ e ν candidates. Overlaid is the best fit to the data. The range of transverse masses used in the fit is indicated with dashes . b) The transverse mass distribution for W $\rightarrow \mu\nu$ candidates.



Figure 3: The relationship between the top quark mass and $\sin^2\theta_W$ with the Z mass constrained to 91.16 GeV/c^2. The curves[22], from top to bottom, correspond to Higgs masses of 1000, 250 and 50 GeV/c^2. The horizontal dotted lines correspond to the 1 σ uncertainties. The vertical dotted line at 89 Gev/c^2 corresponds to the CDF top mass limit[23].