

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
All races 3/, both sexes										
2002.....	847.3	695.0	31.2	15.2	19.5	67.8	95.2	94.7	111.6	161.3
2001.....	848.5	683.4	33.3	15.3	19.2	66.9	95.0	96.2	113.5	165.9
2000.....	854.0	736.7	32.4	15.8	20.3	67.1	93.6	91.2	111.0	159.2
1999.....	857.0	736.0	34.2	16.9	20.4	68.6	90.8	90.4	113.4	160.8
All races 3/, male										
2002.....	846.6	761.5	35.2	16.7	23.1	94.0	140.8	134.4	149.4	206.6
2001.....	846.4	749.8	37.0	16.7	22.8	93.7	141.1	135.6	151.2	213.7
2000.....	853.0	806.5	35.9	17.6	24.2	93.3	138.0	128.8	147.9	205.4
1999.....	859.2	805.0	37.9	18.6	24.4	94.7	133.1	127.0	152.0	207.6
All races 3/, female										
2002.....	848.0	625.3	27.0	13.5	15.9	40.1	47.3	53.6	73.3	116.1
2001.....	850.4	613.9	29.5	13.9	15.3	38.5	46.8	55.8	75.2	118.2
2000.....	855.0	663.4	28.7	14.0	16.1	39.3	47.1	52.8	73.6	113.2
1999.....	854.9	663.6	30.3	15.0	16.2	41.0	46.6	53.2	74.4	114.4
White, both sexes										
2002.....	895.7	586.7	28.1	14.0	18.1	66.3	88.0	85.0	101.1	146.8
2001.....	895.1	570.9	30.7	14.4	17.9	64.2	87.1	87.0	102.7	151.2
2000.....	900.2	610.6	29.2	14.8	19.2	64.5	85.1	81.0	99.0	144.0
1999.....	901.4	606.1	30.8	15.3	19.2	65.3	81.6	80.8	101.4	144.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
White, male										
2002.....	884.0	650.9	31.5	15.4	21.2	90.5	128.9	119.7	136.0	188.7
2001.....	881.9	627.6	34.2	15.3	21.3	88.3	128.2	122.4	137.4	196.7
2000.....	887.8	667.6	32.6	16.5	23.1	88.1	124.6	113.9	133.6	188.2
1999.....	892.1	667.0	33.9	16.8	22.7	88.3	118.5	113.5	136.5	188.8
White, female										
2002.....	907.0	519.4	24.5	12.6	14.8	40.4	44.4	48.1	64.8	103.9
2001.....	907.9	511.5	27.1	13.3	14.4	38.5	43.2	49.6	66.6	104.9
2000.....	912.3	550.5	25.5	13.0	15.2	39.3	43.0	46.5	63.0	99.2
1999.....	910.4	542.0	27.5	13.8	15.4	40.8	42.3	46.7	65.0	99.5
Black, both sexes										
2002.....	768.4	1,263.6	47.1	21.3	27.5	82.0	146.8	172.1	200.7	284.0
2001.....	773.5	1,304.5	47.5	20.7	25.8	86.3	151.3	168.8	200.5	286.0
2000.....	781.1	1,426.1	49.9	21.8	26.6	87.4	155.0	168.4	206.9	283.1
1999.....	788.1	1,457.3	52.6	25.2	28.0	91.9	154.4	163.5	204.9	293.5
Black, male										
2002.....	816.7	1,351.5	54.4	24.1	33.4	121.7	227.6	255.7	272.9	366.5
2001.....	823.9	1,426.4	53.2	23.8	31.6	130.4	236.3	247.0	271.0	364.1
2000.....	834.1	1,567.6	54.5	24.5	32.1	130.1	240.6	246.8	275.0	361.6
1999.....	847.4	1,592.8	59.0	29.3	35.0	137.7	239.9	238.0	278.9	372.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
Black, female										
2002.....	724.4	1,172.0	39.5	18.5	21.3	41.0	68.0	95.7	135.7	210.6
2001.....	727.7	1,176.3	41.7	17.5	19.8	40.8	69.4	97.5	137.3	216.6
2000.....	733.0	1,279.8	45.3	19.0	21.0	43.7	73.9	97.3	145.6	213.4
1999.....	734.3	1,317.4	46.1	21.0	20.7	45.2	73.2	96.1	138.6	223.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
I. Certain infectious and parasitic diseases (A00-B99)										
All races 3/, both sexes										
2002.....	22.2	14.4	1.1	0.5	0.6	0.8	1.5	3.9	8.2	15.3
2001.....	21.2	14.0	1.5	0.5	0.5	1.0	1.7	4.1	9.0	15.5
2000.....	21.0	13.4	1.4	0.5	0.5	0.8	1.6	4.4	10.4	16.0
1999.....	21.5	14.8	1.5	0.7	0.6	0.7	1.8	5.1	11.6	16.5
All races 3/, male										
2002.....	24.2	14.8	1.2	0.5	0.6	0.7	1.6	4.4	10.5	21.6
2001.....	23.3	14.7	1.6	0.5	0.6	1.0	1.7	4.7	12.0	22.2
2000.....	23.0	14.1	1.5	0.5	0.5	0.9	1.8	4.9	13.8	23.1
1999.....	23.7	17.0	1.8	0.7	0.6	0.7	1.9	6.3	15.9	23.7
All races 3/, female										
2002.....	20.2	14.0	0.9	0.4	0.6	0.9	1.4	3.5	5.8	8.9
2001.....	19.2	13.2	1.3	0.5	0.5	0.9	1.7	3.5	6.0	8.9
2000.....	19.0	12.7	1.4	0.4	0.5	0.7	1.4	3.8	6.9	9.0
1999.....	19.3	12.5	1.2	0.7	0.5	0.7	1.8	3.9	7.2	9.3
White, both sexes										
2002.....	19.7	12.4	1.0	0.4	0.5	0.7	0.9	2.3	4.5	9.6
2001.....	18.6	10.6	1.3	0.5	0.4	0.7	1.0	2.1	5.1	9.8
2000.....	18.2	10.6	1.2	0.4	0.4	0.6	1.0	2.3	5.8	10.3
1999.....	18.6	11.9	1.3	0.6	0.4	0.6	1.1	2.9	7.1	10.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
I. Certain infectious and parasitic diseases (A00-B99)										
White, male										
2002.....	20.9	12.7	1.2	0.5	0.6	0.6	1.1	2.9	6.3	14.5
2001.....	19.8	11.3	1.5	0.5	0.5	0.7	1.1	2.7	7.3	15.0
2000.....	19.3	11.2	1.2	0.5	0.4	0.8	1.2	2.9	8.5	15.9
1999.....	19.8	14.3	1.6	0.6	0.5	0.6	1.3	3.8	10.6	16.7
White, female										
2002.....	18.6	12.1	0.8	0.4	0.5	0.8	0.8	1.6	2.6	4.6
2001.....	17.4	9.9	1.2	0.5	0.4	0.6	0.9	1.5	2.8	4.5
2000.....	17.2	10.0	1.2	0.4	0.5	0.5	0.7	1.8	2.9	4.6
1999.....	17.4	9.4	1.1	0.6	0.4	0.6	0.8	1.9	3.5	4.8
Black, both sexes										
2002.....	42.9	24.6	1.7	*	1.2	1.5	4.6	14.8	32.8	54.9
2001.....	42.3	31.2	2.2	0.6	1.1	2.4	5.9	16.6	35.7	55.4
2000.....	43.2	29.9	2.4	0.8	0.7	1.6	5.4	17.6	41.6	56.7
1999.....	44.5	31.1	2.5	1.3	1.1	1.3	6.0	19.6	42.4	57.5
Black, male										
2002.....	51.7	25.3	1.6	*	*	1.4	4.6	14.6	40.9	75.9
2001.....	51.2	33.2	2.6	*	*	2.4	5.6	18.6	46.4	76.4
2000.....	52.7	31.3	2.8	*	*	1.7	5.4	19.4	52.7	79.0
1999.....	54.6	33.2	3.2	1.3	1.4	1.3	5.1	23.8	56.2	78.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
I. Certain infectious and parasitic diseases (A00-B99)										
Black, female										
2002.....	35.0	23.9	1.7	*	1.4	1.7	4.6	14.9	25.5	36.4
2001.....	34.2	29.0	1.8	*	*	2.3	6.2	14.8	26.2	36.6
2000.....	34.5	28.4	2.1	*	*	1.4	5.4	16.1	31.7	36.9
1999.....	35.3	28.9	1.9	1.3	*	*	6.9	15.8	30.1	39.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Zoonotic and other bacterial diseases (A20-A49)										
All races 3/, both sexes										
2002.....	12.3	8.6	0.6	0.3	0.3	0.4	0.4	0.7	1.0	1.6
2001.....	11.9	8.9	0.9	0.2	0.3	0.5	0.5	0.6	1.0	1.5
2000.....	11.6	8.6	0.8	0.3	0.2	0.4	0.5	0.6	0.9	1.6
1999.....	11.7	9.1	0.8	0.3	0.2	0.3	0.6	0.7	1.0	1.6
All races 3/, male										
2002.....	11.1	8.6	0.7	0.3	0.4	0.4	0.5	0.8	1.0	1.6
2001.....	10.8	9.5	0.9	0.2	0.3	0.5	0.5	0.6	1.0	1.7
2000.....	10.4	8.9	0.9	0.3	*	0.5	0.5	0.7	1.0	1.6
1999.....	10.5	10.4	0.9	0.3	0.2	0.3	0.7	0.7	1.1	1.7
All races 3/, female										
2002.....	13.5	8.5	0.6	0.2	0.3	0.4	0.4	0.7	0.9	1.6
2001.....	12.9	8.3	0.8	*	0.3	0.5	0.4	0.6	1.0	1.4
2000.....	12.9	8.3	0.7	0.2	*	0.3	0.4	0.5	0.9	1.5
1999.....	12.9	7.7	0.7	0.3	0.3	0.3	0.5	0.6	0.9	1.5
White, both sexes										
2002.....	12.1	7.3	0.6	0.3	0.3	0.4	0.4	0.6	0.8	1.3
2001.....	11.7	6.8	0.8	0.2	0.3	0.5	0.3	0.5	0.7	1.2
2000.....	11.3	7.1	0.7	0.2	0.2	0.4	0.4	0.5	0.7	1.2
1999.....	11.3	7.2	0.8	0.3	0.2	0.3	0.5	0.5	0.8	1.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Zoonotic and other bacterial diseases (A20-A49)										
White, male										
2002.....	10.9	7.3	0.7	0.4	0.3	0.3	0.4	0.7	0.8	1.3
2001.....	10.5	7.5	0.9	*	0.3	0.5	0.4	0.5	0.7	1.4
2000.....	10.0	7.2	0.8	0.3	*	0.5	0.5	0.5	0.8	1.3
1999.....	10.0	8.5	0.8	0.3	*	0.3	0.6	0.6	0.9	1.4
White, female										
2002.....	13.4	7.4	0.5	*	*	0.4	0.4	0.5	0.8	1.3
2001.....	12.8	6.1	0.7	*	*	0.4	0.3	0.5	0.8	1.0
2000.....	12.6	7.0	0.7	*	*	0.3	0.3	0.5	0.6	1.1
1999.....	12.6	5.9	0.7	0.3	*	0.3	0.4	0.5	0.8	1.1
Black, both sexes										
2002.....	16.9	14.8	0.8	*	0.6	*	0.7	1.8	2.2	3.8
2001.....	16.5	20.3	1.1	*	*	0.9	1.0	1.2	2.8	3.9
2000.....	16.8	18.4	1.2	*	*	*	0.9	1.6	2.7	4.2
1999.....	17.0	19.0	0.9	*	*	*	1.2	1.7	2.2	4.2
Black, male										
2002.....	15.7	15.7	*	*	*	*	*	*	2.6	4.1
2001.....	15.6	20.7	*	*	*	*	1.4	*	2.8	4.0
2000.....	15.6	19.8	*	*	*	*	*	2.1	2.8	4.2
1999.....	16.3	20.5	*	*	*	*	*	*	2.8	4.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Zoonotic and other bacterial diseases (A20-A49)										
Black, female										
2002.....	18.0	13.9	*	*	*	*	*	2.2	2.0	3.5
2001.....	17.3	19.8	*	*	*	*	*	*	2.8	3.8
2000.....	18.0	16.9	*	*	*	*	*	*	2.6	4.2
1999.....	17.7	17.5	*	*	*	*	*	1.9	1.6	3.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Meningococcal infection (A39)										
All races 3/, both sexes										
2002.....	0.1	*	*	*	*	0.1	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	*	*	*	*
2000.....	0.1	0.6	*	*	*	0.2	0.1	*	*	*
1999.....	0.1	0.8	0.2	*	*	0.1	*	*	*	*
All races 3/, male										
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.2	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	*	*	*	*
1999.....	0.1	1.0	0.2	*	*	0.1	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Septicemia (A40-A41)										
All races 3/, both sexes										
2002.....	11.7	7.3	0.5	0.2	0.3	0.2	0.3	0.6	0.9	1.5
2001.....	11.3	7.7	0.7	0.1	0.2	0.3	0.4	0.5	0.9	1.4
2000.....	11.1	7.2	0.6	0.2	0.1	0.2	0.3	0.5	0.9	1.4
1999.....	11.0	7.5	0.6	0.2	0.1	0.2	0.5	0.5	0.8	1.4
All races 3/, male										
2002.....	10.6	7.7	0.5	0.3	0.2	0.2	0.4	0.7	1.0	1.5
2001.....	10.2	8.4	0.8	*	0.2	0.3	0.4	0.5	0.9	1.5
2000.....	9.8	7.1	0.7	0.2	*	0.2	0.4	0.6	0.9	1.4
1999.....	9.8	9.0	0.6	0.2	*	*	0.5	0.5	0.9	1.5
All races 3/, female										
2002.....	12.9	7.0	0.5	*	0.3	0.3	0.3	0.6	0.9	1.5
2001.....	12.4	7.0	0.6	*	*	0.3	0.3	0.5	0.9	1.2
2000.....	12.3	7.3	0.6	*	*	*	0.2	0.4	0.8	1.4
1999.....	12.2	5.8	0.5	0.2	*	*	0.4	0.5	0.7	1.3
White, both sexes										
2002.....	11.5	6.1	0.5	0.2	0.2	0.2	0.3	0.5	0.8	1.2
2001.....	11.1	5.4	0.7	0.2	0.2	0.2	0.3	0.4	0.7	1.1
2000.....	10.8	5.7	0.6	0.2	0.1	0.2	0.2	0.4	0.6	1.1
1999.....	10.6	5.5	0.5	0.2	0.1	0.2	0.4	0.4	0.7	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Septicemia (A40-A41)										
White, male										
2002.....	10.3	6.2	0.5	0.3	0.2	*	0.4	0.6	0.8	1.2
2001.....	9.9	6.2	0.8	*	*	0.3	0.3	0.4	0.6	1.3
2000.....	9.4	5.8	0.6	*	*	*	0.3	0.4	0.7	1.2
1999.....	9.3	6.9	0.5	*	*	*	0.5	0.5	0.8	1.2
White, female										
2002.....	12.8	6.0	0.4	*	*	0.3	*	0.4	0.8	1.2
2001.....	12.3	4.6	0.6	*	*	*	0.3	0.4	0.7	0.9
2000.....	12.1	5.6	0.5	*	*	*	*	0.4	0.5	1.0
1999.....	11.9	4.0	0.5	*	*	*	*	0.4	0.6	0.9
Black, both sexes										
2002.....	16.3	13.5	*	*	*	*	*	1.6	2.2	3.3
2001.....	15.8	19.3	0.9	*	*	*	0.7	1.1	2.5	3.7
2000.....	16.2	16.4	1.0	*	*	*	0.8	1.5	2.5	3.7
1999.....	16.1	17.7	*	*	*	*	1.0	1.4	1.7	3.7
Black, male										
2002.....	15.0	14.8	*	*	*	*	*	*	2.5	3.7
2001.....	14.9	20.1	*	*	*	*	*	*	2.5	3.7
2000.....	14.9	16.0	*	*	*	*	*	2.1	2.5	3.5
1999.....	15.3	19.8	*	*	*	*	*	*	2.1	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Septicemia (A40-A41)										
Black, female										
2002.....	17.4	12.1	*	*	*	*	*	1.9	2.0	3.1
2001.....	16.7	18.6	*	*	*	*	*	*	2.5	3.7
2000.....	17.4	16.9	*	*	*	*	*	*	2.5	3.9
1999.....	16.9	15.4	*	*	*	*	*	1.6	1.4	3.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral diseases (A80-B34)										
All races 3/, both sexes										
2002.....	7.4	4.1	0.3	0.1	0.2	0.3	0.9	2.9	6.8	13.1
2001.....	7.4	3.0	0.4	0.2	0.2	0.4	1.1	3.3	7.7	13.4
2000.....	7.5	2.9	0.4	0.2	0.2	0.3	1.0	3.6	9.1	13.9
1999.....	7.6	3.5	0.5	0.3	0.2	0.3	1.0	4.1	10.0	14.1
All races 3/, male										
2002.....	10.6	4.6	0.4	*	0.2	0.3	1.0	3.3	9.1	19.3
2001.....	10.5	3.1	0.5	0.2	*	0.4	1.0	3.8	10.7	19.7
2000.....	10.8	2.9	0.4	*	0.2	0.3	1.1	4.1	12.3	20.7
1999.....	10.9	4.0	0.6	0.3	0.3	0.3	1.0	5.3	14.2	21.0
All races 3/, female										
2002.....	4.4	3.6	*	*	0.2	0.4	0.8	2.6	4.5	6.9
2001.....	4.3	2.8	0.3	0.2	*	0.3	1.2	2.7	4.6	7.1
2000.....	4.4	2.9	0.4	*	0.3	0.3	0.9	3.1	5.8	7.1
1999.....	4.3	3.1	0.4	0.3	*	0.3	1.0	2.9	5.8	7.2
White, both sexes										
2002.....	5.1	3.7	0.3	*	0.2	0.2	0.4	1.5	3.4	7.9
2001.....	5.0	2.3	0.3	0.2	*	0.2	0.5	1.4	4.1	8.1
2000.....	5.1	2.1	0.3	0.2	0.2	0.2	0.4	1.7	4.8	8.7
1999.....	5.2	2.8	0.4	0.2	0.1	0.2	0.5	2.1	5.9	8.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral diseases (A80-B34)										
White, male										
2002.....	7.5	4.1	0.4	*	*	*	0.5	2.0	5.2	12.8
2001.....	7.4	2.4	0.4	*	*	*	0.6	1.9	6.3	13.0
2000.....	7.5	2.2	*	*	*	0.2	0.6	2.2	7.4	14.0
1999.....	7.6	3.4	0.6	*	*	*	0.6	2.9	9.2	14.5
White, female										
2002.....	2.8	3.3	*	*	*	0.3	0.3	0.9	1.6	2.9
2001.....	2.7	2.2	*	*	*	*	0.4	0.9	1.8	3.2
2000.....	2.8	2.1	*	*	*	*	*	1.2	2.2	3.3
1999.....	2.8	2.1	*	*	*	*	0.4	1.2	2.4	3.3
Black, both sexes										
2002.....	23.5	5.8	*	*	*	1.0	3.5	12.3	29.5	49.7
2001.....	23.7	6.6	0.8	*	0.6	1.3	4.7	15.0	31.8	50.1
2000.....	24.1	7.3	0.9	*	*	0.9	4.2	15.7	37.9	50.9
1999.....	24.3	7.3	1.1	0.9	0.6	0.8	4.3	17.0	38.3	51.0
Black, male										
2002.....	33.0	6.4	*	*	*	*	3.7	12.4	37.1	69.8
2001.....	33.1	7.5	*	*	*	1.4	3.9	17.1	42.7	70.6
2000.....	34.3	7.0	*	*	*	*	4.0	16.9	48.7	72.9
1999.....	35.0	7.5	*	*	*	*	3.8	21.5	51.2	71.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral diseases (A80-B34)										
Black, female										
2002.....	14.9	*	*	*	*	*	3.3	12.2	22.7	31.9
2001.....	15.0	*	*	*	*	*	5.4	13.0	22.1	31.9
2000.....	14.8	7.6	*	*	*	*	4.3	14.6	28.3	31.5
1999.....	14.6	7.0	*	*	*	*	4.9	13.0	26.7	33.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral hepatitis (B15-B19)										
All races 3/, both sexes										
2002.....	2.0	*	*	*	*	*	*	0.2	0.3	1.0
2001.....	2.0	*	*	*	*	*	*	*	0.3	0.9
2000.....	1.9	*	*	*	*	*	*	0.1	0.3	1.0
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9
All races 3/, male										
2002.....	2.7	*	*	*	*	*	*	0.2	0.4	1.4
2001.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.3
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.2
All races 3/, female										
2002.....	1.4	*	*	*	*	*	*	*	*	0.6
2001.....	1.4	*	*	*	*	*	*	*	0.2	0.6
2000.....	1.4	*	*	*	*	*	*	*	0.2	0.6
1999.....	1.3	*	*	*	*	*	*	*	0.2	0.6
White, both sexes										
2002.....	2.0	*	*	*	*	*	*	0.1	0.3	1.0
2001.....	1.9	*	*	*	*	*	*	*	0.2	0.9
2000.....	1.8	*	*	*	*	*	*	*	0.2	1.0
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease (B20-B24)										
All races 3/, both sexes										
2002.....	4.9	*	*	*	0.1	0.2	0.7	2.6	6.4	11.9
2001.....	5.0	*	*	*	0.1	0.2	0.9	3.1	7.3	12.3
2000.....	5.1	*	*	*	0.1	0.2	0.8	3.3	8.7	12.8
1999.....	5.3	*	0.2	0.2	0.1	0.2	0.9	3.9	9.5	13.0
All races 3/, male										
2002.....	7.4	*	*	*	*	*	0.7	3.0	8.5	17.7
2001.....	7.6	*	*	*	*	0.2	0.8	3.7	10.2	18.4
2000.....	7.9	*	*	*	*	*	0.8	3.8	11.9	19.2
1999.....	8.2	*	*	0.2	*	*	0.8	5.1	13.6	19.5
All races 3/, female										
2002.....	2.5	*	*	*	*	0.2	0.6	2.3	4.2	6.2
2001.....	2.5	*	*	*	*	0.2	1.0	2.5	4.4	6.3
2000.....	2.5	*	*	*	*	*	0.7	2.9	5.5	6.4
1999.....	2.5	*	*	*	*	*	1.0	2.7	5.4	6.6
White, both sexes										
2002.....	2.6	*	*	*	*	*	0.3	1.2	3.0	6.8
2001.....	2.7	*	*	*	*	*	0.3	1.3	3.8	7.1
2000.....	2.8	*	*	*	*	*	0.3	1.5	4.5	7.6
1999.....	3.0	*	*	*	*	*	0.4	2.0	5.4	7.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease (B20-B24)										
White, male										
2002.....	4.3	*	*	*	*	*	0.3	1.8	4.7	11.1
2001.....	4.5	*	*	*	*	*	0.4	1.8	5.8	11.6
2000.....	4.7	*	*	*	*	*	0.4	2.0	7.1	12.5
1999.....	5.0	*	*	*	*	*	0.4	2.8	8.6	13.0
White, female										
2002.....	0.9	*	*	*	*	*	*	0.7	1.3	2.3
2001.....	0.9	*	*	*	*	*	0.3	0.8	1.6	2.4
2000.....	1.0	*	*	*	*	*	*	1.0	1.9	2.6
1999.....	1.0	*	*	*	*	*	0.3	1.0	2.1	2.7
Black, both sexes										
2002.....	20.8	*	*	*	*	0.8	3.3	11.7	29.1	48.2
2001.....	21.1	*	*	*	*	1.1	4.4	14.5	31.5	48.8
2000.....	21.4	*	*	*	*	0.7	3.7	15.2	37.1	49.8
1999.....	21.8	*	*	0.7	*	0.7	4.1	16.6	37.8	49.9
Black, male										
2002.....	29.5	*	*	*	*	*	3.3	11.7	36.6	68.4
2001.....	30.1	*	*	*	*	*	3.5	16.8	42.3	69.2
2000.....	31.0	*	*	*	*	*	3.6	16.4	47.9	71.2
1999.....	31.9	*	*	*	*	*	3.5	21.1	50.6	69.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease (B20-B24)										
Black, female										
2002.....	12.8	*	*	*	*	*	3.3	11.7	22.2	30.3
2001.....	12.9	*	*	*	*	*	5.2	12.4	21.8	30.7
2000.....	12.8	*	*	*	*	*	3.8	14.1	27.4	30.8
1999.....	12.6	*	*	*	*	*	4.7	12.5	26.3	32.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)										
All races 3/, both sexes										
2002.....	1.9	*	*	*	*	*	0.3	1.1	2.5	4.9
2001.....	1.9	*	*	*	*	*	0.4	1.3	2.9	4.8
2000.....	2.0	*	*	*	*	*	0.4	1.4	3.4	5.2
1999.....	2.1	*	*	*	*	*	0.4	1.7	3.8	5.4
All races 3/, male										
2002.....	2.9	*	*	*	*	*	0.4	1.2	3.3	7.1
2001.....	2.9	*	*	*	*	*	0.4	1.6	3.9	7.1
2000.....	3.1	*	*	*	*	*	0.4	1.7	4.7	7.8
1999.....	3.3	*	*	*	*	*	0.4	2.2	5.3	8.0
All races 3/, female										
2002.....	1.0	*	*	*	*	*	0.2	1.0	1.8	2.7
2001.....	1.0	*	*	*	*	*	0.4	1.0	1.9	2.5
2000.....	1.0	*	*	*	*	*	0.4	1.1	2.1	2.6
1999.....	1.0	*	*	*	*	*	0.5	1.2	2.4	2.8
White, both sexes										
2002.....	1.1	*	*	*	*	*	0.1	0.5	1.2	2.9
2001.....	1.1	*	*	*	*	*	0.2	0.6	1.6	2.9
2000.....	1.1	*	*	*	*	*	0.2	0.7	1.8	3.1
1999.....	1.2	*	*	*	*	*	0.1	0.9	2.2	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)										
White, male										
2002.....	1.7	*	*	*	*	*	*	0.8	1.7	4.6
2001.....	1.8	*	*	*	*	*	*	0.9	2.4	4.6
2000.....	1.9	*	*	*	*	*	*	0.9	2.9	5.0
1999.....	2.0	*	*	*	*	*	*	1.3	3.3	5.3
White, female										
2002.....	0.4	*	*	*	*	*	*	0.3	0.7	1.2
2001.....	0.4	*	*	*	*	*	*	0.3	0.7	1.2
2000.....	0.4	*	*	*	*	*	*	0.5	0.8	1.2
1999.....	0.4	*	*	*	*	*	*	0.5	1.0	1.3
Black, both sexes										
2002.....	8.0	*	*	*	*	*	1.4	4.5	11.4	18.6
2001.....	8.0	*	*	*	*	*	1.8	5.6	12.0	18.1
2000.....	8.3	*	*	*	*	*	1.6	6.1	14.2	20.2
1999.....	8.6	*	*	*	*	*	2.1	6.9	15.1	20.1
Black, male										
2002.....	11.4	*	*	*	*	*	1.6	4.0	14.8	25.8
2001.....	11.3	*	*	*	*	*	1.9	6.1	15.2	26.2
2000.....	11.9	*	*	*	*	*	1.5	7.1	18.7	29.5
1999.....	12.6	*	*	*	*	*	1.7	8.3	19.7	28.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)										
Black, female										
2002.....	4.9	*	*	*	*	*	*	5.0	8.3	12.3
2001.....	4.9	*	*	*	*	*	1.7	5.2	9.1	11.0
2000.....	5.0	*	*	*	*	*	1.8	5.2	10.2	11.9
1999.....	5.0	*	*	*	*	*	2.5	5.6	11.0	12.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)										
All races 3/, both sexes										
2002.....	0.3	*	*	*	*	*	*	0.1	0.3	0.7
2001.....	0.3	*	*	*	*	*	*	0.2	0.5	0.7
2000.....	0.4	*	*	*	*	*	*	0.2	0.5	0.9
1999.....	0.4	*	*	*	*	*	*	0.2	0.8	1.0
All races 3/, male										
2002.....	0.5	*	*	*	*	*	*	*	0.5	1.2
2001.....	0.6	*	*	*	*	*	*	0.2	0.7	1.3
2000.....	0.6	*	*	*	*	*	*	0.2	0.8	1.5
1999.....	0.7	*	*	*	*	*	*	0.3	1.3	1.7
All races 3/, female										
2002.....	0.1	*	*	*	*	*	*	*	*	0.2
2001.....	0.1	*	*	*	*	*	*	*	0.2	*
2000.....	0.1	*	*	*	*	*	*	*	0.2	0.2
1999.....	0.1	*	*	*	*	*	*	*	0.2	0.3
White, both sexes										
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.5
2001.....	0.2	*	*	*	*	*	*	*	0.3	0.6
2000.....	0.3	*	*	*	*	*	*	*	0.4	0.6
1999.....	0.3	*	*	*	*	*	*	*	0.6	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)										
White, male										
2002.....	0.4	*	*	*	*	*	*	*	0.4	1.0
2001.....	0.4	*	*	*	*	*	*	*	0.5	1.1
2000.....	0.5	*	*	*	*	*	*	*	0.7	1.2
1999.....	0.5	*	*	*	*	*	*	*	1.1	1.5
White, female										
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002.....	1.1	*	*	*	*	*	*	*	1.0	2.1
2001.....	1.1	*	*	*	*	*	*	*	1.6	2.1
2000.....	1.2	*	*	*	*	*	*	0.7	1.5	2.6
1999.....	1.2	*	*	*	*	*	*	0.8	2.1	2.5
Black, male										
2002.....	1.6	*	*	*	*	*	*	*	*	3.4
2001.....	1.7	*	*	*	*	*	*	*	2.7	3.4
2000.....	1.9	*	*	*	*	*	*	*	2.2	4.3
1999.....	1.9	*	*	*	*	*	*	*	3.2	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)										
All races 3/, both sexes										
2002.....	0.5	*	*	*	*	*	*	0.3	0.7	1.0
2001.....	0.5	*	*	*	*	*	*	0.3	0.8	1.3
2000.....	0.6	*	*	*	*	*	*	0.3	1.1	1.5
1999.....	0.6	*	*	*	*	*	*	0.4	1.1	1.5
All races 3/, male										
2002.....	0.8	*	*	*	*	*	*	0.4	1.1	1.7
2001.....	0.8	*	*	*	*	*	*	0.4	1.2	2.2
2000.....	0.9	*	*	*	*	*	*	0.4	1.6	2.3
1999.....	0.9	*	*	*	*	*	*	0.6	1.6	2.5
All races 3/, female										
2002.....	0.2	*	*	*	*	*	*	0.3	0.4	0.4
2001.....	0.2	*	*	*	*	*	*	*	0.4	0.5
2000.....	0.2	*	*	*	*	*	*	0.2	0.6	0.7
1999.....	0.2	*	*	*	*	*	*	0.2	0.5	0.6
White, both sexes										
2002.....	0.3	*	*	*	*	*	*	0.1	0.4	0.6
2001.....	0.3	*	*	*	*	*	*	0.1	0.5	0.9
2000.....	0.4	*	*	*	*	*	*	0.2	0.7	1.0
1999.....	0.4	*	*	*	*	*	*	0.2	0.7	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)										
White, male										
2002.....	0.5	*	*	*	*	*	*	*	0.8	1.2
2001.....	0.5	*	*	*	*	*	*	*	0.8	1.5
2000.....	0.6	*	*	*	*	*	*	0.3	1.1	1.8
1999.....	0.6	*	*	*	*	*	*	0.3	1.1	1.9
White, female										
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	0.2
1999.....	0.1	*	*	*	*	*	*	*	*	0.3
Black, both sexes										
2002.....	1.7	*	*	*	*	*	*	1.4	2.6	3.6
2001.....	1.9	*	*	*	*	*	*	1.3	3.1	4.7
2000.....	2.1	*	*	*	*	*	*	1.6	4.0	4.7
1999.....	2.1	*	*	*	*	*	*	1.5	3.8	5.1
Black, male										
2002.....	2.5	*	*	*	*	*	*	*	3.5	5.5
2001.....	2.8	*	*	*	*	*	*	1.7	4.1	7.3
2000.....	3.2	*	*	*	*	*	*	1.8	5.4	6.2
1999.....	3.2	*	*	*	*	*	*	2.1	5.7	7.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)										
Black, female										
2002.....	1.0	*	*	*	*	*	*	1.4	1.9	1.8
2001.....	1.1	*	*	*	*	*	*	*	2.1	2.5
2000.....	1.2	*	*	*	*	*	*	*	2.8	3.5
1999.....	1.1	*	*	*	*	*	*	*	2.2	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)										
All races 3/, both sexes										
2002.....	1.2	*	*	*	*	*	0.1	0.6	1.4	2.6
2001.....	1.2	*	*	*	*	*	0.3	0.8	1.6	2.9
2000.....	1.2	*	*	*	*	*	0.1	0.7	1.9	2.9
1999.....	1.2	*	*	*	*	*	0.2	0.8	2.0	2.8
All races 3/, male										
2002.....	1.7	*	*	*	*	*	*	0.6	1.9	3.9
2001.....	1.8	*	*	*	*	*	*	0.8	2.2	4.2
2000.....	1.8	*	*	*	*	*	*	0.7	2.4	4.1
1999.....	1.8	*	*	*	*	*	*	1.1	2.7	3.9
All races 3/, female										
2002.....	0.7	*	*	*	*	*	*	0.5	0.9	1.4
2001.....	0.7	*	*	*	*	*	0.4	0.7	1.0	1.7
2000.....	0.7	*	*	*	*	*	*	0.7	1.4	1.6
1999.....	0.7	*	*	*	*	*	0.2	0.6	1.3	1.7
White, both sexes										
2002.....	0.5	*	*	*	*	*	*	0.2	0.5	1.3
2001.....	0.6	*	*	*	*	*	*	0.2	0.7	1.5
2000.....	0.6	*	*	*	*	*	*	0.3	0.8	1.5
1999.....	0.6	*	*	*	*	*	*	0.3	0.9	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)										
White, male										
2002.....	0.9	*	*	*	*	*	*	*	0.8	2.2
2001.....	0.9	*	*	*	*	*	*	*	1.1	2.4
2000.....	0.9	*	*	*	*	*	*	0.3	1.2	2.5
1999.....	0.9	*	*	*	*	*	*	0.5	1.4	2.2
White, female										
2002.....	0.2	*	*	*	*	*	*	*	*	0.5
2001.....	0.2	*	*	*	*	*	*	*	0.4	0.6
2000.....	0.2	*	*	*	*	*	*	*	0.5	0.6
1999.....	0.2	*	*	*	*	*	*	*	0.4	0.6
Black, both sexes										
2002.....	5.7	*	*	*	*	*	*	3.0	7.1	11.7
2001.....	5.7	*	*	*	*	*	1.2	4.1	7.6	12.8
2000.....	5.5	*	*	*	*	*	*	3.9	9.1	12.3
1999.....	5.8	*	*	*	*	*	0.9	4.1	9.2	12.7
Black, male										
2002.....	8.0	*	*	*	*	*	*	3.3	9.2	17.0
2001.....	8.0	*	*	*	*	*	*	4.7	10.4	17.9
2000.....	7.8	*	*	*	*	*	*	4.1	11.0	16.7
1999.....	8.3	*	*	*	*	*	*	5.4	11.9	17.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)										
Black, female										
2002.....	3.6	*	*	*	*	*	*	2.6	5.2	7.0
2001.....	3.6	*	*	*	*	*	1.9	3.6	5.1	8.2
2000.....	3.5	*	*	*	*	*	*	3.7	7.3	8.4
1999.....	3.6	*	*	*	*	*	*	2.9	6.7	8.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Unspecified human immunodeficiency virus (HIV) disease (B24)										
All races 3/, both sexes										
2002.....	1.0	*	*	*	*	*	0.2	0.6	1.4	2.7
2001.....	1.0	*	*	*	*	*	0.2	0.6	1.5	2.5
2000.....	1.0	*	*	*	*	*	0.1	0.7	1.8	2.4
1999.....	1.0	*	*	*	*	*	0.2	0.8	1.9	2.3
All races 3/, male										
2002.....	1.5	*	*	*	*	*	*	0.7	1.7	3.9
2001.....	1.5	*	*	*	*	*	*	0.7	2.2	3.6
2000.....	1.5	*	*	*	*	*	*	0.7	2.4	3.6
1999.....	1.5	*	*	*	*	*	*	0.9	2.7	3.5
All races 3/, female										
2002.....	0.5	*	*	*	*	*	*	0.5	1.1	1.5
2001.....	0.5	*	*	*	*	*	0.2	0.4	0.9	1.4
2000.....	0.5	*	*	*	*	*	*	0.6	1.2	1.2
1999.....	0.5	*	*	*	*	*	*	0.6	1.1	1.2
White, both sexes										
2002.....	0.5	*	*	*	*	*	*	0.3	0.6	1.3
2001.....	0.5	*	*	*	*	*	*	0.2	0.7	1.2
2000.....	0.5	*	*	*	*	*	*	0.3	0.8	1.3
1999.....	0.5	*	*	*	*	*	*	0.4	1.1	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Unspecified human immunodeficiency virus (HIV) disease (B24)										
White, male										
2002.....	0.8	*	*	*	*	*	*	0.4	0.9	2.2
2001.....	0.8	*	*	*	*	*	*	0.3	1.1	2.1
2000.....	0.8	*	*	*	*	*	*	0.5	1.3	2.1
1999.....	0.9	*	*	*	*	*	*	0.5	1.7	2.2
White, female										
2002.....	0.2	*	*	*	*	*	*	*	0.3	0.4
2001.....	0.1	*	*	*	*	*	*	*	0.2	0.3
2000.....	0.2	*	*	*	*	*	*	*	0.4	0.4
1999.....	0.2	*	*	*	*	*	*	*	0.4	0.4
Black, both sexes										
2002.....	4.3	*	*	*	*	*	0.8	2.4	6.9	12.2
2001.....	4.4	*	*	*	*	*	0.9	2.8	7.2	11.1
2000.....	4.3	*	*	*	*	*	0.8	2.9	8.4	10.1
1999.....	4.1	*	*	*	*	*	0.7	3.3	7.5	9.5
Black, male										
2002.....	6.0	*	*	*	*	*	*	2.8	7.8	16.7
2001.....	6.3	*	*	*	*	*	*	3.5	9.9	14.5
2000.....	6.3	*	*	*	*	*	*	2.5	10.6	14.6
1999.....	5.9	*	*	*	*	*	*	4.0	10.1	13.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Unspecified human immunodeficiency virus (HIV) disease (B24)										
Black, female										
2002.....	2.8	*	*	*	*	*	*	2.1	6.2	8.2
2001.....	2.7	*	*	*	*	*	*	2.2	4.8	8.1
2000.....	2.6	*	*	*	*	*	*	3.3	6.3	6.1
1999.....	2.4	*	*	*	*	*	*	2.7	5.2	6.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)										
All races 3/, both sexes										
2002.....	0.4	3.2	0.3	*	*	*	0.1	*	0.1	0.2
2001.....	0.3	2.1	0.3	0.1	*	0.1	*	*	*	0.1
2000.....	0.4	1.8	0.3	*	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.4	2.7	0.3	0.1	*	*	*	*	0.1	0.2
All races 3/, male										
2002.....	0.4	3.6	0.4	*	*	*	*	*	*	*
2001.....	0.3	2.1	0.4	*	*	*	*	*	*	*
2000.....	0.4	1.9	0.3	*	*	*	*	*	*	*
1999.....	0.4	2.8	0.4	*	*	*	*	*	*	0.2
All races 3/, female										
2002.....	0.4	2.8	*	*	*	*	*	*	*	*
2001.....	0.3	2.0	0.3	*	*	*	*	*	*	*
2000.....	0.4	1.7	*	*	*	*	*	*	*	*
1999.....	0.4	2.5	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.4	3.1	0.3	*	*	*	*	*	0.1	0.1
2001.....	0.3	1.7	0.3	*	*	*	*	*	*	*
2000.....	0.4	1.5	0.3	*	*	*	*	*	*	0.1
1999.....	0.4	2.3	0.3	*	*	*	*	*	*	0.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

All races 3/, both sexes

2002.....	1.3	1.3	*	*	*	*	0.1	0.2	0.3	0.4
2001.....	1.1	1.6	*	*	*	*	0.1	0.2	0.3	0.5
2000.....	1.0	1.3	*	*	*	*	*	0.1	0.3	0.4
1999.....	1.4	1.6	0.1	*	*	*	0.1	0.3	0.5	0.7

All races 3/, male

2002.....	1.5	1.5	*	*	*	*	*	0.2	0.4	0.6
2001.....	1.2	1.6	*	*	*	*	*	0.2	0.3	0.6
2000.....	1.1	1.5	*	*	*	*	*	*	0.3	0.6
1999.....	1.6	1.9	*	*	*	*	*	0.4	0.5	0.8

All races 3/, female

2002.....	1.0	1.2	*	*	*	*	*	*	0.3	0.3
2001.....	1.0	1.6	*	*	*	*	*	*	0.2	0.4
2000.....	0.9	1.1	*	*	*	*	*	*	0.2	0.3
1999.....	1.3	1.2	*	*	*	*	*	0.3	0.4	0.5

White, both sexes

2002.....	1.3	1.1	*	*	*	*	*	0.2	0.2	0.3
2001.....	1.1	1.2	*	*	*	*	*	0.2	0.2	0.4
2000.....	1.0	1.0	*	*	*	*	*	*	0.2	0.3
1999.....	1.3	1.2	*	*	*	*	*	0.2	0.3	0.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)										
White, male										
2002.....	1.5	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.2	*	*	*	*	*	*	*	*	0.5
2000.....	1.1	1.3	*	*	*	*	*	*	*	0.5
1999.....	1.5	1.6	*	*	*	*	*	0.3	0.3	0.7
White, female										
2002.....	1.0	*	*	*	*	*	*	*	*	0.2
2001.....	1.0	1.4	*	*	*	*	*	*	*	0.3
2000.....	0.9	*	*	*	*	*	*	*	*	*
1999.....	1.2	*	*	*	*	*	*	*	0.3	0.3
Black, both sexes										
2002.....	1.5	*	*	*	*	*	*	*	0.8	1.0
2001.....	1.3	3.7	*	*	*	*	*	*	0.8	1.0
2000.....	1.3	3.3	*	*	*	*	*	*	0.8	1.2
1999.....	2.2	3.8	*	*	*	*	*	0.9	1.7	1.9
Black, male										
2002.....	1.8	*	*	*	*	*	*	*	*	*
2001.....	1.5	*	*	*	*	*	*	*	*	*
2000.....	1.5	*	*	*	*	*	*	*	*	1.7
1999.....	2.3	*	*	*	*	*	*	*	1.9	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

Black, female

2002.....	1.1	*	*	*	*	*	*	*	*	*
2001.....	1.1	*	*	*	*	*	*	*	*	*
2000.....	1.1	*	*	*	*	*	*	*	*	*
1999.....	2.2	*	*	*	*	*	*	*	1.4	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Mycoses (B35-B49)										
All races 3/, both sexes										
2002.....	0.4	1.2	*	*	*	*	*	*	0.2	0.2
2001.....	0.4	1.4	*	*	*	*	*	*	*	0.1
2000.....	0.4	1.3	*	*	*	*	*	*	0.1	0.2
1999.....	0.7	1.4	*	*	*	*	*	0.2	0.2	0.3
All races 3/, male										
2002.....	0.5	1.2	*	*	*	*	*	*	0.2	0.2
2001.....	0.4	1.4	*	*	*	*	*	*	*	*
2000.....	0.5	1.5	*	*	*	*	*	*	*	0.3
1999.....	0.8	1.6	*	*	*	*	*	0.2	0.3	0.4
All races 3/, female										
2002.....	0.3	1.1	*	*	*	*	*	*	*	*
2001.....	0.4	1.5	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*
1999.....	0.6	1.1	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.4	0.9	*	*	*	*	*	*	*	0.1
2001.....	0.4	1.0	*	*	*	*	*	*	*	*
2000.....	0.4	1.0	*	*	*	*	*	*	*	0.2
1999.....	0.6	1.0	*	*	*	*	*	0.1	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
II. Neoplasms (C00-D48)										
All races 3/, both sexes										
2002.....	197.9	3.5	3.0	2.9	2.7	3.7	5.2	7.0	13.0	24.1
2001.....	199.2	2.9	3.1	2.7	2.7	3.8	5.2	7.7	13.0	25.3
2000.....	201.3	4.0	3.1	2.7	2.7	3.9	5.4	7.4	12.9	25.2
1999.....	201.8	3.3	3.1	2.8	2.7	4.0	5.5	7.6	13.1	25.8
All races 3/, male										
2002.....	208.5	3.5	3.1	3.1	3.0	4.2	6.0	7.4	11.7	20.5
2001.....	210.1	2.8	3.3	2.7	2.9	4.5	6.1	7.6	11.6	21.7
2000.....	212.0	4.2	3.4	2.9	3.0	4.6	6.3	7.4	11.8	22.0
1999.....	213.7	3.4	3.3	2.9	2.8	4.6	6.4	8.0	12.2	22.3
All races 3/, female										
2002.....	187.6	3.6	2.8	2.7	2.4	3.2	4.4	6.5	14.2	27.8
2001.....	188.7	3.0	2.9	2.7	2.6	3.2	4.3	7.7	14.5	28.8
2000.....	191.1	3.8	2.8	2.5	2.4	3.2	4.4	7.4	14.1	28.4
1999.....	190.3	3.2	2.9	2.6	2.5	3.3	4.6	7.2	13.9	29.3
White, both sexes										
2002.....	210.6	3.7	2.9	3.0	2.7	3.8	5.4	6.8	12.6	23.3
2001.....	211.7	2.9	3.4	2.7	2.7	3.9	5.2	7.5	12.6	24.4
2000.....	213.9	4.1	3.1	2.8	2.8	3.8	5.3	7.4	12.6	24.0
1999.....	213.9	3.2	3.3	2.8	2.6	3.8	5.4	7.6	12.6	24.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
II. Neoplasms (C00-D48)										
White, male										
2002.....	220.6	3.9	3.0	3.1	3.0	4.4	6.2	7.2	11.6	19.9
2001.....	221.7	2.7	3.5	2.7	2.6	4.5	6.1	7.5	11.6	21.2
2000.....	223.3	4.4	3.5	2.9	3.1	4.4	6.3	7.4	11.8	21.2
1999.....	224.8	3.4	3.5	3.0	2.8	4.3	6.2	7.9	11.9	21.7
White, female										
2002.....	200.9	3.5	2.7	2.9	2.5	3.2	4.6	6.4	13.6	26.9
2001.....	201.9	3.1	3.2	2.8	2.7	3.2	4.3	7.6	13.7	27.8
2000.....	204.8	3.9	2.7	2.6	2.5	3.1	4.4	7.3	13.4	26.9
1999.....	203.3	2.9	3.2	2.6	2.4	3.3	4.5	7.2	13.4	27.5
Black, both sexes										
2002.....	168.9	3.1	3.5	2.8	2.8	3.8	4.8	8.5	17.0	32.4
2001.....	170.2	*	2.1	2.8	3.2	4.0	5.4	9.4	16.8	33.6
2000.....	172.4	3.3	3.2	2.4	2.5	4.6	6.2	8.6	16.3	35.3
1999.....	174.5	4.3	2.5	2.8	3.2	4.9	7.0	8.9	16.6	36.9
Black, male										
2002.....	184.3	*	3.3	3.6	3.0	4.3	5.4	8.6	14.3	26.2
2001.....	187.5	*	2.6	2.9	3.8	4.7	6.3	8.7	13.0	27.8
2000.....	191.5	*	3.3	2.7	2.8	5.7	7.3	7.7	13.3	28.6
1999.....	194.1	*	2.8	2.6	3.5	6.1	7.8	9.4	14.9	28.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
II. Neoplasms (C00-D48)										
Black, female										
2002.....	155.0	*	3.7	1.9	2.5	3.3	4.2	8.5	19.3	37.9
2001.....	154.6	*	1.6	2.6	2.5	3.2	4.5	10.0	20.3	38.8
2000.....	155.0	*	3.0	2.2	2.3	3.6	5.2	9.5	19.0	41.2
1999.....	156.8	*	2.1	2.9	2.8	3.7	6.3	8.4	18.1	44.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
All races 3/, both sexes										
2002.....	193.2	1.8	2.6	2.7	2.5	3.5	5.0	6.6	12.5	23.5
2001.....	194.4	1.6	2.7	2.4	2.5	3.6	4.9	7.3	12.6	24.6
2000.....	196.5	2.4	2.7	2.4	2.6	3.7	5.1	7.0	12.5	24.5
1999.....	197.0	1.8	2.7	2.5	2.5	3.7	5.3	7.3	12.5	25.2
All races 3/, male										
2002.....	203.8	2.0	2.7	2.9	2.8	4.0	5.7	7.0	11.3	19.8
2001.....	205.3	1.5	2.9	2.4	2.6	4.3	5.7	7.2	11.2	21.0
2000.....	207.2	2.6	3.0	2.6	2.9	4.3	5.9	7.1	11.3	21.2
1999.....	208.9	1.9	2.9	2.6	2.7	4.3	6.1	7.6	11.6	21.6
All races 3/, female										
2002.....	183.0	1.6	2.4	2.5	2.2	3.0	4.2	6.2	13.7	27.2
2001.....	183.9	1.8	2.5	2.4	2.3	2.9	4.1	7.4	14.0	28.1
2000.....	186.2	2.3	2.5	2.2	2.2	3.0	4.2	7.0	13.7	27.8
1999.....	185.6	1.7	2.5	2.3	2.3	3.0	4.4	7.0	13.4	28.7
White, both sexes										
2002.....	205.5	1.8	2.5	2.8	2.5	3.6	5.2	6.5	12.1	22.7
2001.....	206.4	1.7	3.0	2.5	2.4	3.7	5.0	7.2	12.2	23.7
2000.....	208.6	2.5	2.8	2.5	2.6	3.6	5.1	7.0	12.1	23.3
1999.....	208.7	1.8	2.9	2.5	2.4	3.6	5.2	7.3	12.1	24.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
White, male										
2002.....	215.5	2.2	2.7	2.9	2.7	4.2	5.9	6.9	11.1	19.2
2001.....	216.4	1.5	3.1	2.4	2.4	4.4	5.8	7.2	11.2	20.5
2000.....	218.1	2.8	3.1	2.6	2.9	4.2	5.9	7.1	11.2	20.5
1999.....	219.5	2.0	3.2	2.7	2.6	4.1	5.9	7.6	11.3	21.1
White, female										
2002.....	195.8	1.4	2.3	2.6	2.4	3.0	4.4	6.1	13.2	26.3
2001.....	196.7	2.0	2.8	2.5	2.4	2.9	4.1	7.2	13.3	27.1
2000.....	199.4	2.3	2.4	2.3	2.3	3.0	4.2	7.0	13.0	26.3
1999.....	198.2	1.5	2.7	2.2	2.2	3.0	4.3	7.0	12.9	27.1
Black, both sexes										
2002.....	165.9	*	2.9	2.6	2.5	3.6	4.5	8.0	16.3	31.7
2001.....	167.1	*	1.9	2.4	2.8	3.6	5.1	9.1	16.4	32.7
2000.....	169.3	*	2.8	2.1	2.3	4.4	5.9	8.0	15.8	34.5
1999.....	171.3	*	2.0	2.4	3.0	4.5	6.6	8.6	15.8	35.8
Black, male										
2002.....	181.5	*	2.6	3.3	2.9	4.1	5.0	8.2	14.1	25.6
2001.....	184.5	*	2.4	2.5	3.4	4.3	5.9	8.5	12.4	26.8
2000.....	188.5	*	2.7	2.4	2.7	5.2	7.1	7.3	12.9	27.6
1999.....	191.0	*	2.1	2.3	3.3	5.6	7.3	8.8	14.0	27.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
Black, female										
2002.....	151.7	*	3.2	1.9	2.1	3.1	4.0	7.9	18.3	37.1
2001.....	151.3	*	*	2.2	2.2	2.9	4.3	9.7	20.0	37.9
2000.....	151.8	*	2.9	1.8	2.0	3.5	4.7	8.5	18.3	40.5
1999.....	153.3	*	1.9	2.6	2.7	3.4	5.9	8.3	17.5	43.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
All races 3/, both sexes										
2002.....	2.7	*	*	*	*	*	*	0.1	0.2	0.4
2001.....	2.7	*	*	*	*	*	*	*	0.2	0.4
2000.....	2.7	*	*	*	*	*	*	*	0.2	0.4
1999.....	2.7	*	*	*	*	*	*	*	0.1	0.4
All races 3/, male										
2002.....	3.7	*	*	*	*	*	*	*	0.2	0.4
2001.....	3.7	*	*	*	*	*	*	*	0.2	0.5
2000.....	3.6	*	*	*	*	*	*	*	*	0.5
1999.....	3.6	*	*	*	*	*	*	*	*	0.5
All races 3/, female										
2002.....	1.7	*	*	*	*	*	*	*	*	0.3
2001.....	1.8	*	*	*	*	*	*	*	*	0.3
2000.....	1.8	*	*	*	*	*	*	*	*	0.2
1999.....	1.8	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002.....	2.7	*	*	*	*	*	*	*	0.1	0.3
2001.....	2.7	*	*	*	*	*	*	*	0.1	0.3
2000.....	2.7	*	*	*	*	*	*	*	0.1	0.3
1999.....	2.7	*	*	*	*	*	*	*	0.1	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of digestive organs (C15-C26)										
All races 3/, both sexes										
2002.....	45.7	*	0.2	*	*	0.1	0.5	1.0	2.0	4.6
2001.....	46.0	*	*	*	*	0.2	0.4	0.9	2.1	5.0
2000.....	46.4	*	0.2	*	*	0.2	0.3	0.9	2.1	4.5
1999.....	46.4	*	0.2	*	*	0.2	0.4	1.0	2.0	4.6
All races 3/, male										
2002.....	50.7	*	0.3	*	*	*	0.5	1.2	2.3	5.3
2001.....	50.5	*	*	*	*	*	0.4	0.9	2.4	6.0
2000.....	50.8	*	*	*	*	0.2	0.4	1.1	2.5	5.5
1999.....	50.7	*	0.3	*	*	0.2	0.4	1.2	2.3	5.3
All races 3/, female										
2002.....	40.8	*	*	*	*	*	0.4	0.8	1.7	3.8
2001.....	41.7	*	*	*	*	*	0.3	0.8	1.8	4.0
2000.....	42.3	*	*	*	*	*	0.3	0.8	1.6	3.4
1999.....	42.3	*	*	*	*	*	0.4	0.8	1.7	3.8
White, both sexes										
2002.....	47.3	*	0.2	*	*	0.1	0.4	0.9	1.8	4.3
2001.....	47.6	*	*	*	*	0.1	0.3	0.8	1.9	4.5
2000.....	48.0	*	*	*	*	0.2	0.3	0.8	1.8	4.0
1999.....	48.0	*	0.2	*	*	0.1	0.4	0.8	1.7	4.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of digestive organs (C15-C26)										
White, male										
2002.....	52.3	*	*	*	*	*	0.4	1.0	2.0	4.8
2001.....	52.1	*	*	*	*	*	0.3	0.8	2.1	5.4
2000.....	52.3	*	*	*	*	*	0.3	0.9	2.2	5.0
1999.....	52.2	*	*	*	*	*	0.4	1.0	2.0	4.8
White, female										
2002.....	42.4	*	*	*	*	*	0.5	0.8	1.6	3.7
2001.....	43.2	*	*	*	*	*	0.4	0.8	1.6	3.5
2000.....	43.9	*	*	*	*	*	*	0.6	1.4	3.1
1999.....	43.9	*	*	*	*	*	0.4	0.6	1.5	3.4
Black, both sexes										
2002.....	43.6	*	*	*	*	*	*	1.4	2.7	5.9
2001.....	44.1	*	*	*	*	*	*	1.2	3.1	7.7
2000.....	44.5	*	*	*	*	*	*	1.3	3.3	6.9
1999.....	44.6	*	*	*	*	*	*	1.8	3.1	7.2
Black, male										
2002.....	48.5	*	*	*	*	*	*	1.9	3.5	7.6
2001.....	48.1	*	*	*	*	*	*	*	3.7	9.5
2000.....	49.2	*	*	*	*	*	*	*	4.1	8.5
1999.....	49.4	*	*	*	*	*	*	2.3	3.7	7.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of digestive organs (C15-C26)										
Black, female										
2002.....	39.2	*	*	*	*	*	*	*	2.0	4.4
2001.....	40.5	*	*	*	*	*	*	*	2.6	6.1
2000.....	40.3	*	*	*	*	*	*	1.4	2.5	5.5
1999.....	40.4	*	*	*	*	*	*	*	2.5	6.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of esophagus (C15)										
All races 3/, both sexes										
2002.....	4.4	*	*	*	*	*	*	*	*	0.4
2001.....	4.4	*	*	*	*	*	*	*	0.1	0.3
2000.....	4.3	*	*	*	*	*	*	*	0.1	0.3
1999.....	4.3	*	*	*	*	*	*	*	0.1	0.3
All races 3/, male										
2002.....	6.9	*	*	*	*	*	*	*	*	0.6
2001.....	6.9	*	*	*	*	*	*	*	*	0.5
2000.....	6.7	*	*	*	*	*	*	*	0.2	0.6
1999.....	6.6	*	*	*	*	*	*	*	0.2	0.6
All races 3/, female										
2002.....	2.0	*	*	*	*	*	*	*	*	*
2001.....	2.0	*	*	*	*	*	*	*	*	*
2000.....	2.1	*	*	*	*	*	*	*	*	*
1999.....	2.0	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	4.6	*	*	*	*	*	*	*	*	0.4
2001.....	4.6	*	*	*	*	*	*	*	*	0.3
2000.....	4.5	*	*	*	*	*	*	*	0.2	0.4
1999.....	4.4	*	*	*	*	*	*	*	0.1	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of stomach (C16)										
All races 3/, both sexes										
2002.....	4.2	*	*	*	*	*	0.1	0.3	0.4	0.7
2001.....	4.3	*	*	*	*	*	*	0.2	0.4	0.8
2000.....	4.5	*	*	*	*	*	*	0.2	0.4	0.7
1999.....	4.6	*	*	*	*	*	*	0.2	0.4	0.8
All races 3/, male										
2002.....	5.0	*	*	*	*	*	*	0.3	0.5	0.7
2001.....	5.1	*	*	*	*	*	*	0.2	0.5	0.9
2000.....	5.3	*	*	*	*	*	*	0.2	0.5	0.9
1999.....	5.5	*	*	*	*	*	*	0.2	0.5	0.9
All races 3/, female										
2002.....	3.5	*	*	*	*	*	*	*	0.4	0.7
2001.....	3.5	*	*	*	*	*	*	*	0.4	0.7
2000.....	3.7	*	*	*	*	*	*	*	0.4	0.6
1999.....	3.7	*	*	*	*	*	*	0.3	0.3	0.8
White, both sexes										
2002.....	4.0	*	*	*	*	*	*	0.2	0.4	0.7
2001.....	4.1	*	*	*	*	*	*	0.2	0.4	0.6
2000.....	4.2	*	*	*	*	*	*	0.1	0.3	0.6
1999.....	4.3	*	*	*	*	*	*	0.2	0.4	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of colon, rectum and anus (C18-C21)										
All races 3/, both sexes										
2002.....	19.7	*	*	*	*	*	0.2	0.5	1.0	2.0
2001.....	20.0	*	*	*	*	*	0.2	0.4	1.1	2.3
2000.....	20.4	*	*	*	*	*	0.1	0.4	0.9	2.1
1999.....	20.5	*	*	*	*	*	0.2	0.4	1.0	1.9
All races 3/, male										
2002.....	20.1	*	*	*	*	*	0.3	0.5	1.1	2.1
2001.....	20.2	*	*	*	*	*	*	0.4	1.1	2.6
2000.....	20.6	*	*	*	*	*	*	0.4	1.1	2.4
1999.....	20.6	*	*	*	*	*	0.2	0.6	1.0	2.0
All races 3/, female										
2002.....	19.2	*	*	*	*	*	*	0.4	0.9	1.9
2001.....	19.8	*	*	*	*	*	*	0.5	1.0	1.9
2000.....	20.2	*	*	*	*	*	*	0.4	0.8	1.8
1999.....	20.3	*	*	*	*	*	*	0.3	0.9	1.9
White, both sexes										
2002.....	20.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2001.....	21.0	*	*	*	*	*	0.2	0.4	1.0	2.1
2000.....	21.5	*	*	*	*	*	*	0.4	0.9	1.9
1999.....	21.6	*	*	*	*	*	0.2	0.4	0.9	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Colon (C18)										
All races 3/, both sexes										
2002.....	16.5	*	*	*	*	*	0.2	0.3	0.7	1.5
2001.....	16.8	*	*	*	*	*	0.1	0.3	0.8	1.7
2000.....	17.3	*	*	*	*	*	*	0.3	0.8	1.6
1999.....	17.4	*	*	*	*	*	0.2	0.3	0.7	1.5
All races 3/, male										
2002.....	16.6	*	*	*	*	*	0.2	0.3	0.8	1.5
2001.....	16.7	*	*	*	*	*	*	0.3	0.9	1.9
2000.....	17.1	*	*	*	*	*	*	0.3	0.9	1.7
1999.....	17.2	*	*	*	*	*	*	0.4	0.8	1.5
All races 3/, female										
2002.....	16.4	*	*	*	*	*	*	0.3	0.7	1.4
2001.....	16.9	*	*	*	*	*	*	0.3	0.8	1.5
2000.....	17.4	*	*	*	*	*	*	0.3	0.6	1.4
1999.....	17.5	*	*	*	*	*	*	0.3	0.7	1.5
White, both sexes										
2002.....	17.3	*	*	*	*	*	0.2	0.3	0.7	1.4
2001.....	17.6	*	*	*	*	*	0.1	0.3	0.8	1.6
2000.....	18.1	*	*	*	*	*	*	0.3	0.7	1.4
1999.....	18.2	*	*	*	*	*	*	0.3	0.7	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Rectosigmoid junction and rectum (C19-C20)										
All races 3/, both sexes										
2002.....	3.0	*	*	*	*	*	*	0.1	0.2	0.5
2001.....	3.0	*	*	*	*	*	*	0.1	0.2	0.5
2000.....	3.0	*	*	*	*	*	*	0.1	0.2	0.5
1999.....	3.0	*	*	*	*	*	*	0.1	0.2	0.3
All races 3/, male										
2002.....	3.4	*	*	*	*	*	*	*	0.3	0.5
2001.....	3.4	*	*	*	*	*	*	*	0.2	0.6
2000.....	3.4	*	*	*	*	*	*	*	*	0.6
1999.....	3.3	*	*	*	*	*	*	0.2	0.2	0.4
All races 3/, female										
2002.....	2.6	*	*	*	*	*	*	*	*	0.4
2001.....	2.6	*	*	*	*	*	*	*	*	0.3
2000.....	2.6	*	*	*	*	*	*	*	*	0.3
1999.....	2.7	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002.....	3.2	*	*	*	*	*	*	0.1	0.2	0.4
2001.....	3.2	*	*	*	*	*	*	0.1	0.2	0.4
2000.....	3.2	*	*	*	*	*	*	*	0.1	0.4
1999.....	3.2	*	*	*	*	*	*	0.1	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of liver and intrahepatic bile ducts (C22)										
All races 3/, both sexes										
2002.....	4.9	*	0.2	*	*	*	*	0.2	0.3	0.6
2001.....	4.7	*	*	*	*	*	*	0.2	0.2	0.6
2000.....	4.6	*	0.1	*	*	*	*	0.1	0.3	0.5
1999.....	4.4	*	0.2	*	*	*	0.1	0.2	0.2	0.6
All races 3/, male										
2002.....	6.4	*	*	*	*	*	*	0.2	0.4	0.8
2001.....	6.1	*	*	*	*	*	*	*	0.3	0.9
2000.....	5.9	*	*	*	*	*	*	*	0.4	0.7
1999.....	5.6	*	0.3	*	*	*	*	0.2	0.3	0.8
All races 3/, female										
2002.....	3.4	*	*	*	*	*	*	*	0.3	0.3
2001.....	3.4	*	*	*	*	*	*	*	*	0.3
2000.....	3.3	*	*	*	*	*	*	*	*	0.3
1999.....	3.3	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002.....	4.8	*	0.2	*	*	*	*	*	0.2	0.4
2001.....	4.6	*	*	*	*	*	*	*	0.2	0.5
2000.....	4.5	*	*	*	*	*	*	*	0.2	0.4
1999.....	4.4	*	0.2	*	*	*	*	*	*	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Liver (C22.0,C22.2-C22.9)										
All races 3/, both sexes										
2002.....	3.8	*	0.2	*	*	*	*	0.2	0.2	0.4
2001.....	3.7	*	*	*	*	*	*	0.1	0.2	0.5
2000.....	3.6	*	0.1	*	*	*	*	0.1	0.2	0.4
1999.....	3.5	*	0.2	*	*	*	0.1	0.1	0.2	0.4
All races 3/, male										
2002.....	5.3	*	*	*	*	*	*	0.2	0.3	0.6
2001.....	5.1	*	*	*	*	*	*	*	0.2	0.8
2000.....	5.0	*	*	*	*	*	*	*	0.3	0.6
1999.....	4.7	*	0.3	*	*	*	*	*	0.2	0.6
All races 3/, female										
2002.....	2.4	*	*	*	*	*	*	*	*	0.2
2001.....	2.4	*	*	*	*	*	*	*	*	0.2
2000.....	2.3	*	*	*	*	*	*	*	*	0.3
1999.....	2.4	*	*	*	*	*	*	*	*	0.2
White, both sexes										
2002.....	3.7	*	0.2	*	*	*	*	*	0.1	0.3
2001.....	3.5	*	*	*	*	*	*	*	0.2	0.3
2000.....	3.4	*	*	*	*	*	*	*	0.1	0.3
1999.....	3.4	*	0.2	*	*	*	*	*	*	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of pancreas (C25)										
All races 3/, both sexes										
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.7
2001.....	10.5	*	*	*	*	*	*	*	0.2	0.8
2000.....	10.4	*	*	*	*	*	*	*	0.2	0.6
1999.....	10.4	*	*	*	*	*	*	*	0.2	0.8
All races 3/, male										
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.8
2001.....	10.3	*	*	*	*	*	*	*	0.2	0.9
2000.....	10.3	*	*	*	*	*	*	*	0.2	0.8
1999.....	10.4	*	*	*	*	*	*	*	0.3	0.9
All races 3/, female										
2002.....	10.5	*	*	*	*	*	*	*	*	0.6
2001.....	10.6	*	*	*	*	*	*	*	*	0.6
2000.....	10.5	*	*	*	*	*	*	*	*	0.5
1999.....	10.5	*	*	*	*	*	*	*	0.2	0.6
White, both sexes										
2002.....	11.1	*	*	*	*	*	*	*	0.1	0.7
2001.....	11.1	*	*	*	*	*	*	*	0.2	0.7
2000.....	11.0	*	*	*	*	*	*	*	0.2	0.6
1999.....	11.0	*	*	*	*	*	*	*	0.2	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)										
All races 3/, both sexes										
2002.....	56.3	*	*	*	*	*	0.2	0.2	0.7	2.9
2001.....	56.5	*	*	*	*	*	0.1	0.2	0.7	3.1
2000.....	57.0	*	*	*	*	*	0.1	0.2	0.9	3.1
1999.....	56.2	*	*	*	*	*	0.1	0.3	0.8	3.1
All races 3/, male										
2002.....	66.1	*	*	*	*	*	0.3	0.3	0.8	3.0
2001.....	67.2	*	*	*	*	*	*	0.2	0.7	3.1
2000.....	68.1	*	*	*	*	*	*	0.4	0.8	3.5
1999.....	68.0	*	*	*	*	*	*	0.3	0.8	3.2
All races 3/, female										
2002.....	46.8	*	*	*	*	*	*	*	0.7	2.8
2001.....	46.1	*	*	*	*	*	*	0.3	0.7	3.1
2000.....	46.2	*	*	*	*	*	*	*	0.9	2.7
1999.....	45.0	*	*	*	*	*	*	0.2	0.7	3.1
White, both sexes										
2002.....	60.7	*	*	*	*	*	0.1	0.2	0.7	2.8
2001.....	60.8	*	*	*	*	*	0.1	0.2	0.7	3.1
2000.....	61.2	*	*	*	*	*	0.1	0.2	0.9	2.9
1999.....	60.2	*	*	*	*	*	*	0.3	0.8	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of trachea, bronchus and lung (C33-C34)										
All races 3/, both sexes										
2002.....	54.7	*	*	*	*	*	*	0.2	0.6	2.7
2001.....	54.8	*	*	*	*	*	*	0.2	0.6	2.9
2000.....	55.3	*	*	*	*	*	*	0.2	0.8	2.9
1999.....	54.5	*	*	*	*	*	*	0.2	0.7	2.9
All races 3/, male										
2002.....	63.7	*	*	*	*	*	*	*	0.7	2.6
2001.....	64.7	*	*	*	*	*	*	*	0.6	2.8
2000.....	65.5	*	*	*	*	*	*	0.3	0.7	3.3
1999.....	65.4	*	*	*	*	*	*	0.2	0.7	2.8
All races 3/, female										
2002.....	46.0	*	*	*	*	*	*	*	0.6	2.7
2001.....	45.3	*	*	*	*	*	*	*	0.6	2.9
2000.....	45.4	*	*	*	*	*	*	*	0.8	2.6
1999.....	44.1	*	*	*	*	*	*	*	0.6	3.0
White, both sexes										
2002.....	59.1	*	*	*	*	*	*	0.2	0.6	2.6
2001.....	59.1	*	*	*	*	*	*	0.2	0.6	2.9
2000.....	59.5	*	*	*	*	*	*	0.2	0.8	2.7
1999.....	58.5	*	*	*	*	*	*	0.2	0.7	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of bone and articular cartilage (C40-C41)										
All races 3/, both sexes										
2002.....	0.4	*	*	0.1	0.3	0.5	0.4	0.3	0.2	0.2
2001.....	0.5	*	*	*	0.3	0.5	0.5	0.3	0.2	0.2
2000.....	0.4	*	*	*	0.3	0.5	0.4	0.2	0.2	0.2
1999.....	0.4	*	*	0.1	0.3	0.6	0.4	0.2	0.2	0.2
All races 3/, male										
2002.....	0.5	*	*	*	0.3	0.5	0.5	0.4	0.2	0.2
2001.....	0.5	*	*	*	0.4	0.6	0.7	0.3	0.2	0.2
2000.....	0.5	*	*	*	0.4	0.6	0.5	0.3	0.3	0.2
1999.....	0.5	*	*	*	0.3	0.7	0.5	0.3	0.2	0.2
All races 3/, female										
2002.....	0.4	*	*	*	0.3	0.4	0.3	*	*	*
2001.....	0.4	*	*	*	0.3	0.4	0.3	0.2	0.2	0.2
2000.....	0.4	*	*	*	0.2	0.4	0.3	*	*	*
1999.....	0.4	*	*	*	0.3	0.5	0.3	*	0.2	*
White, both sexes										
2002.....	0.4	*	*	*	0.3	0.5	0.4	0.3	0.2	0.2
2001.....	0.5	*	*	*	0.4	0.6	0.5	0.3	0.2	0.2
2000.....	0.5	*	*	*	0.3	0.5	0.4	0.2	0.2	0.2
1999.....	0.5	*	*	*	0.3	0.7	0.4	0.2	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Melanoma and other malignant neoplasms of skin (C43-C44)										
All races 3/, both sexes										
2002.....	3.4	*	*	*	*	*	0.2	0.4	0.6	1.1
2001.....	3.5	*	*	*	*	*	0.2	0.4	0.6	1.0
2000.....	3.4	*	*	*	*	*	0.2	0.3	0.6	1.1
1999.....	3.4	*	*	*	*	*	0.2	0.3	0.7	1.2
All races 3/, male										
2002.....	4.5	*	*	*	*	*	*	0.4	0.8	1.2
2001.....	4.6	*	*	*	*	*	*	0.5	0.6	1.3
2000.....	4.5	*	*	*	*	*	*	0.4	0.7	1.1
1999.....	4.5	*	*	*	*	*	*	0.4	0.8	1.5
All races 3/, female										
2002.....	2.4	*	*	*	*	*	*	0.4	0.4	0.9
2001.....	2.5	*	*	*	*	*	*	0.3	0.5	0.8
2000.....	2.5	*	*	*	*	*	*	0.3	0.5	1.1
1999.....	2.4	*	*	*	*	*	*	0.2	0.6	0.8
White, both sexes										
2002.....	4.1	*	*	*	*	*	0.2	0.5	0.8	1.3
2001.....	4.2	*	*	*	*	*	0.2	0.5	0.7	1.2
2000.....	4.0	*	*	*	*	*	0.2	0.4	0.8	1.3
1999.....	4.0	*	*	*	*	*	0.2	0.4	0.8	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant melanoma of skin (C43)										
All races 3/, both sexes										
2002.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2001.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2000.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0
1999.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0
All races 3/, male										
2002.....	3.3	*	*	*	*	*	*	0.4	0.8	1.1
2001.....	3.4	*	*	*	*	*	*	0.4	0.6	1.2
2000.....	3.3	*	*	*	*	*	*	0.3	0.7	1.0
1999.....	3.3	*	*	*	*	*	*	0.4	0.7	1.3
All races 3/, female										
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.9
2001.....	1.9	*	*	*	*	*	*	0.3	0.5	0.7
2000.....	2.0	*	*	*	*	*	*	0.3	0.5	1.0
1999.....	1.9	*	*	*	*	*	*	0.2	0.6	0.8
White, both sexes										
2002.....	3.1	*	*	*	*	*	0.2	0.5	0.7	1.2
2001.....	3.2	*	*	*	*	*	0.2	0.5	0.7	1.2
2000.....	3.1	*	*	*	*	*	0.2	0.4	0.8	1.2
1999.....	3.1	*	*	*	*	*	0.2	0.4	0.8	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of mesothelial and soft tissue (C45-C49)										
All races 3/, both sexes										
2002.....	2.4	*	0.1	0.1	0.2	0.3	0.4	0.4	0.4	0.7
2001.....	2.4	*	*	0.1	0.2	0.3	0.3	0.4	0.6	0.6
2000.....	2.4	*	0.1	*	0.1	0.4	0.4	0.5	0.6	0.7
1999.....	2.4	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.7
All races 3/, male										
2002.....	2.8	*	*	*	0.2	0.3	0.4	0.5	0.5	0.8
2001.....	2.8	*	*	*	*	0.4	0.4	0.5	0.7	0.6
2000.....	2.9	*	*	*	*	0.4	0.5	0.5	0.6	0.8
1999.....	2.8	*	*	*	0.2	0.4	0.4	0.5	0.7	0.9
All races 3/, female										
2002.....	1.9	*	*	*	0.2	0.3	0.3	0.4	0.3	0.5
2001.....	2.0	*	*	*	*	0.3	0.2	0.4	0.4	0.6
2000.....	2.0	*	*	*	*	0.4	0.3	0.4	0.6	0.6
1999.....	1.9	*	*	*	*	*	0.4	0.4	0.4	0.6
White, both sexes										
2002.....	2.6	*	*	0.1	0.2	0.3	0.4	0.4	0.4	0.7
2001.....	2.6	*	*	*	0.2	0.3	0.3	0.4	0.5	0.6
2000.....	2.7	*	*	*	0.1	0.4	0.3	0.5	0.6	0.6
1999.....	2.6	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of other mesothelial and soft tissue (C47-C49)										
All races 3/, both sexes										
2002.....	1.5	*	0.1	0.1	0.2	0.3	0.4	0.4	0.4	0.6
2001.....	1.5	*	*	0.1	0.2	0.3	0.3	0.4	0.5	0.5
2000.....	1.6	*	0.1	*	0.1	0.4	0.4	0.4	0.5	0.6
1999.....	1.5	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.7
All races 3/, male										
2002.....	1.4	*	*	*	0.2	0.3	0.4	0.5	0.5	0.7
2001.....	1.4	*	*	*	*	0.4	0.4	0.4	0.6	0.5
2000.....	1.4	*	*	*	*	0.4	0.5	0.5	0.5	0.7
1999.....	1.5	*	*	*	0.2	0.4	0.4	0.5	0.6	0.7
All races 3/, female										
2002.....	1.6	*	*	*	0.2	0.3	0.3	0.4	0.3	0.5
2001.....	1.6	*	*	*	*	0.3	0.2	0.4	0.4	0.5
2000.....	1.7	*	*	*	*	0.4	0.3	0.4	0.5	0.5
1999.....	1.6	*	*	*	*	*	0.4	0.4	0.4	0.6
White, both sexes										
2002.....	1.6	*	*	0.1	0.2	0.3	0.4	0.4	0.4	0.6
2001.....	1.6	*	*	*	0.2	0.3	0.3	0.4	0.5	0.6
2000.....	1.7	*	*	*	0.1	0.3	0.3	0.5	0.6	0.6
1999.....	1.6	*	*	0.1	0.2	0.3	0.3	0.4	0.4	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of breast (C50)										
All races 3/, both sexes										
2002.....	14.5	*	*	*	*	*	*	0.3	1.7	4.1
2001.....	14.7	*	*	*	*	*	*	0.5	1.8	4.3
2000.....	15.0	*	*	*	*	*	*	0.4	1.9	4.3
1999.....	14.9	*	*	*	*	*	*	0.4	1.7	4.3
All races 3/, male										
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	28.3	*	*	*	*	*	*	0.7	3.5	8.2
2001.....	28.6	*	*	*	*	*	*	0.9	3.6	8.6
2000.....	29.2	*	*	*	*	*	*	0.8	3.8	8.5
1999.....	28.9	*	*	*	*	*	*	0.9	3.4	8.5
White, both sexes										
2002.....	15.1	*	*	*	*	*	*	0.3	1.5	3.6
2001.....	15.2	*	*	*	*	*	*	0.4	1.5	3.9
2000.....	15.7	*	*	*	*	*	*	0.4	1.6	3.9
1999.....	15.5	*	*	*	*	*	*	0.4	1.5	3.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of breast (C50)										
Black, female										
2002.....	28.2	*	*	*	*	*	*	1.6	6.9	15.1
2001.....	28.3	*	*	*	*	*	*	2.2	7.6	14.2
2000.....	27.9	*	*	*	*	*	*	*	7.1	14.3
1999.....	28.2	*	*	*	*	*	*	1.8	6.9	16.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of female genital organs (C51-C58)

All races 3/, both sexes

2002.....	9.4	*	*	*	*	*	0.2	0.4	1.2	2.2
2001.....	9.4	*	*	*	*	*	0.2	0.5	1.2	2.3
2000.....	9.4	*	*	*	*	*	0.2	0.6	1.2	2.2
1999.....	9.3	*	*	*	*	*	0.2	0.5	1.1	2.2

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	18.5	*	*	*	*	*	0.5	0.9	2.5	4.5
2001.....	18.5	*	*	*	*	*	0.3	1.1	2.5	4.5
2000.....	18.4	*	*	*	*	*	0.4	1.2	2.4	4.4
1999.....	18.2	*	*	*	*	*	0.4	1.1	2.2	4.4

White, both sexes

2002.....	9.9	*	*	*	*	*	0.2	0.4	1.2	2.2
2001.....	9.9	*	*	*	*	*	0.1	0.5	1.2	2.2
2000.....	9.9	*	*	*	*	*	0.2	0.7	1.1	2.1
1999.....	9.8	*	*	*	*	*	0.2	0.5	1.1	2.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of female genital organs (C51-C58)										
Black, female										
2002.....	16.4	*	*	*	*	*	*	*	3.3	6.3
2001.....	16.2	*	*	*	*	*	*	1.7	3.2	5.6
2000.....	16.4	*	*	*	*	*	*	*	3.7	5.8
1999.....	16.1	*	*	*	*	*	*	*	2.5	5.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of cervix uteri (C53)										
All races 3/, both sexes										
2002.....	1.4	*	*	*	*	*	*	0.2	0.8	1.3
2001.....	1.4	*	*	*	*	*	*	0.3	0.8	1.3
2000.....	1.5	*	*	*	*	*	*	0.4	0.7	1.3
1999.....	1.5	*	*	*	*	*	*	0.3	0.8	1.3
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	2.7	*	*	*	*	*	*	0.5	1.5	2.5
2001.....	2.8	*	*	*	*	*	*	0.5	1.5	2.6
2000.....	2.9	*	*	*	*	*	*	0.8	1.5	2.6
1999.....	3.0	*	*	*	*	*	*	0.6	1.5	2.5
White, both sexes										
2002.....	1.3	*	*	*	*	*	*	0.2	0.8	1.2
2001.....	1.3	*	*	*	*	*	*	0.3	0.7	1.3
2000.....	1.4	*	*	*	*	*	*	0.4	0.7	1.2
1999.....	1.4	*	*	*	*	*	*	0.3	0.7	1.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of cervix uteri (C53)										
Black, female										
2002.....	4.2	*	*	*	*	*	*	*	2.0	3.9
2001.....	4.0	*	*	*	*	*	*	*	2.1	3.5
2000.....	4.5	*	*	*	*	*	*	*	2.7	4.0
1999.....	4.4	*	*	*	*	*	*	*	1.9	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of ovary (C56)										
All races 3/, both sexes										
2002.....	5.1	*	*	*	*	*	0.1	0.2	0.3	0.7
2001.....	5.1	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	5.0	*	*	*	*	*	*	0.2	0.3	0.6
1999.....	4.9	*	*	*	*	*	*	0.1	0.2	0.7
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	10.0	*	*	*	*	*	0.3	0.4	0.6	1.4
2001.....	9.9	*	*	*	*	*	*	0.4	0.6	1.4
2000.....	9.8	*	*	*	*	*	*	0.4	0.6	1.3
1999.....	9.6	*	*	*	*	*	*	0.3	0.5	1.4
White, both sexes										
2002.....	5.6	*	*	*	*	*	0.1	0.2	0.3	0.7
2001.....	5.6	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	5.5	*	*	*	*	*	*	0.2	0.3	0.6
1999.....	5.4	*	*	*	*	*	*	0.1	0.2	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of male genital organs (C60-C63)										
All races 3/, both sexes										
2002.....	10.8	*	*	*	*	*	0.2	0.2	0.3	0.3
2001.....	11.0	*	*	*	*	*	0.1	0.2	0.2	0.2
2000.....	11.3	*	*	*	*	*	0.2	0.2	0.2	0.3
1999.....	11.6	*	*	*	*	*	0.2	0.3	0.3	0.3
All races 3/, male										
2002.....	21.9	*	*	*	*	*	0.3	0.4	0.5	0.5
2001.....	22.4	*	*	*	*	*	0.3	0.5	0.4	0.5
2000.....	22.9	*	*	*	*	*	0.4	0.4	0.4	0.6
1999.....	23.6	*	*	*	*	*	0.3	0.5	0.5	0.6
All races 3/, female										
2002.....
2001.....
2000.....
1999.....
White, both sexes										
2002.....	10.9	*	*	*	*	*	0.2	0.2	0.3	0.3
2001.....	11.0	*	*	*	*	*	0.2	0.3	0.2	0.3
2000.....	11.2	*	*	*	*	*	0.2	0.3	0.2	0.3
1999.....	11.6	*	*	*	*	*	0.2	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of testis (C62)										
All races 3/, both sexes										
2002.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
2001.....	0.1	*	*	*	*	*	0.1	0.2	0.2	0.2
2000.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
1999.....	0.1	*	*	*	*	*	0.1	0.2	0.3	0.2
All races 3/, male										
2002.....	0.3	*	*	*	*	*	0.3	0.4	0.5	0.4
2001.....	0.2	*	*	*	*	*	0.3	0.4	0.4	0.4
2000.....	0.2	*	*	*	*	*	0.3	0.4	0.4	0.5
1999.....	0.3	*	*	*	*	*	0.3	0.5	0.5	0.5
All races 3/, female										
2002.....
2001.....
2000.....
1999.....
White, both sexes										
2002.....	0.2	*	*	*	*	*	0.2	0.2	0.3	0.2
2001.....	0.1	*	*	*	*	*	0.2	0.3	0.2	0.2
2000.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.3
1999.....	0.2	*	*	*	*	*	0.2	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of urinary tract (C64-C68)										
All races 3/, both sexes										
2002.....	8.8	*	*	0.1	*	*	*	*	0.2	0.6
2001.....	8.7	*	0.1	0.2	*	*	*	0.1	0.2	0.5
2000.....	8.7	*	*	0.1	*	*	*	*	0.2	0.6
1999.....	8.5	*	*	0.1	*	*	*	0.2	0.2	0.6
All races 3/, male										
2002.....	11.6	*	*	*	*	*	*	*	0.3	0.8
2001.....	11.6	*	*	*	*	*	*	*	0.2	0.6
2000.....	11.4	*	*	*	*	*	*	*	0.2	0.7
1999.....	11.2	*	*	*	*	*	*	*	0.3	0.8
All races 3/, female										
2002.....	6.1	*	*	*	*	*	*	*	*	0.4
2001.....	6.0	*	*	0.2	*	*	*	*	*	0.3
2000.....	6.0	*	*	*	*	*	*	*	*	0.4
1999.....	5.9	*	*	*	*	*	*	*	*	0.4
White, both sexes										
2002.....	9.8	*	*	0.1	*	*	*	*	0.2	0.5
2001.....	9.7	*	*	0.1	*	*	*	*	0.2	0.4
2000.....	9.6	*	*	*	*	*	*	*	0.2	0.5
1999.....	9.3	*	*	*	*	*	*	*	0.2	0.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes

2002.....	4.4	*	0.7	1.0	0.6	0.5	0.5	0.7	1.2	1.7
2001.....	4.4	*	0.7	0.8	0.6	0.5	0.6	0.7	1.4	1.8
2000.....	4.5	0.5	0.7	0.8	0.7	0.5	0.6	0.8	1.2	2.0
1999.....	4.6	*	0.7	0.9	0.6	0.6	0.6	0.9	1.3	1.9

All races 3/, male

2002.....	5.0	*	0.8	1.1	0.7	0.6	0.7	0.7	1.5	2.0
2001.....	5.0	*	0.7	0.8	0.6	0.6	0.7	0.9	1.7	2.2
2000.....	5.1	*	0.7	0.8	0.8	0.6	0.7	1.0	1.6	2.6
1999.....	5.1	*	0.7	0.9	0.6	0.7	0.7	1.1	1.7	2.3

All races 3/, female

2002.....	3.9	*	0.7	0.9	0.5	0.4	0.3	0.7	1.0	1.4
2001.....	3.9	*	0.6	0.9	0.6	0.4	0.6	0.5	1.1	1.4
2000.....	3.9	*	0.6	0.8	0.6	0.4	0.5	0.7	0.8	1.4
1999.....	4.1	*	0.8	0.9	0.7	0.5	0.5	0.8	0.9	1.5

White, both sexes

2002.....	5.0	*	0.7	1.0	0.6	0.5	0.5	0.7	1.4	1.9
2001.....	5.0	*	0.7	0.9	0.6	0.6	0.6	0.8	1.5	2.1
2000.....	5.1	*	0.7	0.8	0.7	0.5	0.7	0.9	1.3	2.2
1999.....	5.2	*	0.8	0.9	0.6	0.5	0.6	1.1	1.4	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of brain (C71)										
All races 3/, both sexes										
2002.....	4.4	*	0.7	1.0	0.6	0.5	0.5	0.7	1.2	1.7
2001.....	4.3	*	0.6	0.8	0.6	0.5	0.6	0.7	1.3	1.8
2000.....	4.4	0.5	0.7	0.8	0.7	0.5	0.6	0.8	1.2	2.0
1999.....	4.5	*	0.7	0.9	0.6	0.5	0.5	0.9	1.3	1.9
All races 3/, male										
2002.....	4.9	*	0.8	1.1	0.7	0.6	0.7	0.7	1.4	2.0
2001.....	4.9	*	0.6	0.8	0.6	0.6	0.6	0.9	1.6	2.2
2000.....	5.0	*	0.7	0.8	0.8	0.6	0.7	0.9	1.6	2.6
1999.....	5.0	*	0.7	0.9	0.6	0.6	0.6	1.0	1.6	2.2
All races 3/, female										
2002.....	3.8	*	0.6	0.9	0.5	0.4	0.3	0.7	1.0	1.4
2001.....	3.8	*	0.6	0.9	0.6	0.4	0.6	0.5	1.0	1.4
2000.....	3.9	*	0.6	0.7	0.6	0.4	0.5	0.7	0.7	1.4
1999.....	4.0	*	0.7	0.9	0.6	0.4	0.5	0.8	0.9	1.5
White, both sexes										
2002.....	4.9	*	0.6	1.0	0.6	0.5	0.5	0.7	1.3	1.9
2001.....	4.9	*	0.7	0.9	0.6	0.5	0.6	0.8	1.5	2.1
2000.....	5.0	*	0.7	0.8	0.7	0.5	0.6	0.9	1.3	2.2
1999.....	5.1	*	0.8	0.8	0.6	0.5	0.6	1.0	1.4	2.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)										
All races 3/, both sexes										
2002.....	0.7	*	0.5	0.4	0.1	*	*	*	0.1	0.1
2001.....	0.7	*	0.5	0.3	0.1	*	*	0.1	*	0.2
2000.....	0.7	*	0.5	0.3	0.1	*	*	*	0.1	0.2
1999.....	0.7	*	0.5	0.4	0.1	*	*	*	0.1	0.2
All races 3/, male										
2002.....	0.6	*	0.4	0.4	*	*	*	*	*	*
2001.....	0.7	*	0.5	0.3	*	*	*	*	*	*
2000.....	0.7	*	0.5	0.3	*	*	*	*	*	*
1999.....	0.7	*	0.5	0.4	*	*	*	*	*	0.2
All races 3/, female										
2002.....	0.8	*	0.5	0.3	*	*	*	*	*	*
2001.....	0.8	*	0.4	0.2	*	*	*	*	*	*
2000.....	0.8	*	0.5	0.3	*	*	*	*	*	0.2
1999.....	0.8	*	0.4	0.4	*	*	*	*	*	0.2
White, both sexes										
2002.....	0.8	*	0.4	0.3	0.1	*	*	*	*	0.2
2001.....	0.8	*	0.5	0.3	*	*	*	*	*	0.1
2000.....	0.8	*	0.5	0.4	*	*	*	*	*	0.2
1999.....	0.8	*	0.5	0.4	*	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)										
All races 3/, both sexes										
2002.....	14.1	*	0.1	*	*	0.3	0.3	0.4	0.9	1.5
2001.....	14.1	*	*	*	0.1	0.2	0.3	0.5	0.8	1.6
2000.....	14.5	*	0.1	*	0.1	0.2	0.3	0.4	0.7	1.6
1999.....	15.7	*	*	*	0.1	0.2	0.4	0.5	0.9	1.7
All races 3/, male										
2002.....	14.7	*	*	*	*	0.3	0.4	0.4	0.9	1.6
2001.....	14.5	*	*	*	*	0.3	0.3	0.5	0.9	1.5
2000.....	14.8	*	*	*	*	0.2	0.3	0.5	0.7	1.5
1999.....	16.2	*	*	*	*	0.2	0.5	0.6	1.1	1.7
All races 3/, female										
2002.....	13.5	*	*	*	*	0.2	0.2	0.3	0.8	1.5
2001.....	13.6	*	*	*	*	*	0.2	0.5	0.6	1.6
2000.....	14.1	*	*	*	*	0.2	0.3	0.4	0.7	1.7
1999.....	15.3	*	*	*	*	*	0.3	0.4	0.8	1.7
White, both sexes										
2002.....	14.9	*	*	*	*	0.3	0.3	0.3	0.9	1.4
2001.....	14.8	*	*	*	*	0.2	0.3	0.4	0.7	1.5
2000.....	15.3	*	*	*	*	0.2	0.3	0.5	0.7	1.5
1999.....	16.6	*	*	*	*	0.2	0.4	0.5	0.9	1.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
All races 3/, both sexes										
2002.....	19.5	0.7	0.8	0.8	1.0	1.5	1.9	1.9	2.6	3.2
2001.....	19.8	0.7	1.0	0.8	1.0	1.5	2.0	2.3	2.7	3.5
2000.....	20.1	1.0	0.9	0.8	1.0	1.6	2.1	2.1	2.5	3.4
1999.....	20.2	0.8	0.9	0.7	1.0	1.5	2.2	2.2	2.6	3.7
All races 3/, male										
2002.....	21.4	*	0.8	0.9	1.2	1.8	2.2	2.3	3.0	3.7
2001.....	21.5	*	1.0	0.9	1.1	1.8	2.3	2.6	3.1	4.2
2000.....	21.6	1.1	1.0	0.9	1.2	1.9	2.6	2.3	3.0	3.9
1999.....	21.7	*	1.0	0.8	1.2	1.7	2.7	2.4	3.1	4.4
All races 3/, female										
2002.....	17.7	*	0.7	0.8	0.8	1.2	1.7	1.5	2.2	2.6
2001.....	18.1	*	1.0	0.7	0.8	1.2	1.7	2.0	2.2	2.7
2000.....	18.6	*	0.9	0.7	0.8	1.2	1.5	1.9	1.9	3.0
1999.....	18.7	*	0.8	0.6	0.9	1.3	1.8	1.9	2.2	3.0
White, both sexes										
2002.....	21.3	*	0.8	0.9	1.0	1.5	2.1	1.9	2.5	3.1
2001.....	21.5	0.7	1.1	0.8	1.0	1.6	2.0	2.1	2.6	3.4
2000.....	21.9	1.2	1.0	0.9	1.1	1.6	2.0	2.1	2.4	3.3
1999.....	21.9	0.8	1.0	0.7	1.0	1.5	2.3	2.1	2.5	3.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
White, male										
2002.....	23.4	*	0.9	0.9	1.2	1.9	2.4	2.3	2.9	3.5
2001.....	23.3	*	1.1	0.9	1.0	1.9	2.3	2.4	3.2	4.0
2000.....	23.6	*	1.0	0.9	1.2	1.9	2.5	2.3	2.9	3.8
1999.....	23.6	*	1.0	0.9	1.1	1.7	2.8	2.3	2.9	4.3
White, female										
2002.....	19.2	*	0.8	0.9	0.8	1.1	1.7	1.5	2.2	2.6
2001.....	19.7	*	1.1	0.7	0.9	1.2	1.6	1.9	2.0	2.8
2000.....	20.3	*	1.0	0.8	0.9	1.2	1.4	1.8	1.9	2.9
1999.....	20.4	*	0.9	0.6	0.9	1.3	1.7	1.9	2.1	2.9
Black, both sexes										
2002.....	13.6	*	*	*	1.0	1.4	1.5	2.3	3.4	3.8
2001.....	14.1	*	*	0.9	1.0	1.5	2.1	3.2	3.5	4.5
2000.....	13.9	*	0.8	0.6	0.7	1.9	2.6	2.8	3.1	4.4
1999.....	14.1	*	*	0.6	1.3	1.7	2.9	2.6	3.2	4.6
Black, male										
2002.....	14.6	*	*	*	1.1	1.6	1.3	2.8	4.4	5.1
2001.....	15.1	*	*	*	1.7	1.8	2.2	3.8	4.0	6.5
2000.....	14.7	*	*	*	*	2.3	3.2	2.6	4.4	4.8
1999.....	15.1	*	*	*	1.7	2.1	3.1	3.3	4.2	5.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
Black, female										
2002.....	12.8	*	*	*	*	1.3	1.6	2.0	2.5	2.7
2001.....	13.1	*	*	*	*	*	2.1	2.7	3.1	2.8
2000.....	13.1	*	*	*	*	1.4	2.1	3.0	1.9	4.1
1999.....	13.2	*	*	*	*	1.4	2.8	2.0	2.3	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hodgkin's disease (C81)										
All races 3/, both sexes										
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.3
1999.....	0.5	*	*	*	*	0.2	0.3	0.4	0.3	0.4
All races 3/, male										
2002.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.5
2000.....	0.5	*	*	*	*	*	0.3	0.3	0.5	0.4
1999.....	0.6	*	*	*	*	0.2	0.4	0.4	0.4	0.5
All races 3/, female										
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3
2001.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.3
2000.....	0.4	*	*	*	*	*	0.2	0.3	0.4	0.3
1999.....	0.4	*	*	*	*	*	0.3	0.4	0.3	0.3
White, both sexes										
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4
1999.....	0.6	*	*	*	*	0.2	0.3	0.4	0.4	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)										
All races 3/, both sexes										
2002.....	7.6	*	*	*	0.2	0.2	0.5	0.6	0.8	1.2
2001.....	7.9	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4
2000.....	8.1	*	*	0.1	0.1	0.3	0.6	0.6	0.8	1.3
1999.....	8.2	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4
All races 3/, male										
2002.....	8.2	*	*	*	0.3	0.3	0.6	0.7	1.0	1.5
2001.....	8.4	*	*	*	0.2	0.4	0.5	0.8	1.0	1.8
2000.....	8.6	*	*	*	*	0.4	0.8	0.7	1.0	1.7
1999.....	8.6	*	*	*	0.2	0.3	0.6	0.7	1.1	1.8
All races 3/, female										
2002.....	7.1	*	*	*	*	*	0.4	0.4	0.6	0.8
2001.....	7.3	*	*	*	*	*	0.4	0.6	0.6	1.0
2000.....	7.6	*	*	*	*	*	0.4	0.6	0.6	0.9
1999.....	7.8	*	*	*	*	0.2	0.4	0.4	0.7	1.1
White, both sexes										
2002.....	8.5	*	*	*	0.2	0.2	0.5	0.6	0.7	1.1
2001.....	8.8	*	*	*	0.1	0.3	0.4	0.6	0.8	1.3
2000.....	9.1	*	*	*	0.1	0.3	0.6	0.6	0.8	1.2
1999.....	9.2	*	*	*	*	0.2	0.5	0.5	0.8	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

White, male

2002.....	9.1	*	*	*	0.3	0.3	0.6	0.8	0.9	1.4
2001.....	9.3	*	*	*	*	0.4	0.5	0.7	1.0	1.6
2000.....	9.6	*	*	*	*	0.5	0.8	0.6	1.0	1.6
1999.....	9.6	*	*	*	*	0.3	0.6	0.6	1.0	1.7

White, female

2002.....	8.0	*	*	*	*	*	0.3	0.4	0.6	0.8
2001.....	8.3	*	*	*	*	*	0.4	0.5	0.5	1.0
2000.....	8.7	*	*	*	*	*	0.4	0.5	0.6	0.8
1999.....	8.8	*	*	*	*	*	0.4	0.4	0.6	1.0

Black, both sexes

2002.....	3.9	*	*	*	*	*	*	0.8	1.5	1.4
2001.....	4.1	*	*	*	*	*	*	1.1	1.3	2.0
2000.....	3.8	*	*	*	*	*	1.0	1.1	0.9	1.9
1999.....	4.0	*	*	*	*	*	*	0.9	1.2	2.1

Black, male

2002.....	4.3	*	*	*	*	*	*	*	2.2	2.0
2001.....	4.6	*	*	*	*	*	*	*	*	3.1
2000.....	4.2	*	*	*	*	*	*	*	*	2.7
1999.....	4.6	*	*	*	*	*	*	*	1.7	2.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Non-Hodgkin's lymphoma (C82-C85)										
All races 3/, both sexes										
2002.....	7.6	*	*	*	0.2	0.2	0.5	0.6	0.8	1.2
2001.....	7.8	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4
2000.....	8.1	*	*	0.1	0.1	0.3	0.6	0.6	0.8	1.3
1999.....	8.2	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4
All races 3/, male										
2002.....	8.1	*	*	*	0.3	0.3	0.6	0.7	1.0	1.5
2001.....	8.4	*	*	*	0.2	0.4	0.5	0.8	1.0	1.8
2000.....	8.6	*	*	*	*	0.4	0.8	0.7	1.0	1.7
1999.....	8.6	*	*	*	0.2	0.3	0.6	0.7	1.1	1.8
All races 3/, female										
2002.....	7.1	*	*	*	*	*	0.4	0.4	0.6	0.8
2001.....	7.3	*	*	*	*	*	0.4	0.6	0.6	1.0
2000.....	7.6	*	*	*	*	*	0.4	0.6	0.6	0.9
1999.....	7.7	*	*	*	*	0.2	0.4	0.4	0.7	1.0
White, both sexes										
2002.....	8.5	*	*	*	0.2	0.2	0.5	0.6	0.7	1.1
2001.....	8.8	*	*	*	0.1	0.3	0.4	0.6	0.8	1.3
2000.....	9.1	*	*	*	0.1	0.3	0.6	0.6	0.8	1.2
1999.....	9.2	*	*	*	*	0.2	0.5	0.5	0.8	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Non-Hodgkin's lymphoma (C82-C85)										
White, male										
2002.....	9.1	*	*	*	0.3	0.3	0.6	0.8	0.9	1.4
2001.....	9.3	*	*	*	*	0.4	0.5	0.7	1.0	1.6
2000.....	9.6	*	*	*	*	0.5	0.8	0.6	1.0	1.6
1999.....	9.6	*	*	*	*	0.3	0.6	0.6	1.0	1.7
White, female										
2002.....	8.0	*	*	*	*	*	0.3	0.4	0.6	0.8
2001.....	8.2	*	*	*	*	*	0.4	0.5	0.5	1.0
2000.....	8.6	*	*	*	*	*	0.4	0.5	0.6	0.8
1999.....	8.8	*	*	*	*	*	0.4	0.4	0.6	1.0
Black, both sexes										
2002.....	3.9	*	*	*	*	*	*	0.8	1.5	1.4
2001.....	4.0	*	*	*	*	*	*	1.1	1.2	2.0
2000.....	3.7	*	*	*	*	*	1.0	1.1	0.9	1.9
1999.....	4.0	*	*	*	*	*	*	0.9	1.2	2.1
Black, male										
2002.....	4.3	*	*	*	*	*	*	*	2.2	2.0
2001.....	4.5	*	*	*	*	*	*	*	*	3.0
2000.....	4.2	*	*	*	*	*	*	*	*	2.7
1999.....	4.6	*	*	*	*	*	*	*	1.7	2.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
All races 3/, both sexes										
2002.....	7.5	0.5	0.7	0.7	0.8	1.2	1.2	1.0	1.3	1.5
2001.....	7.5	0.6	1.0	0.8	0.8	1.1	1.3	1.2	1.4	1.6
2000.....	7.6	1.0	0.9	0.7	0.8	1.2	1.2	1.2	1.2	1.6
1999.....	7.5	0.7	0.8	0.6	0.9	1.1	1.4	1.2	1.3	1.6
All races 3/, male										
2002.....	8.5	*	0.7	0.7	0.9	1.4	1.4	1.2	1.5	1.6
2001.....	8.5	*	1.0	0.8	0.9	1.3	1.5	1.4	1.6	1.8
2000.....	8.5	1.1	0.9	0.8	1.0	1.4	1.5	1.3	1.5	1.6
1999.....	8.4	*	0.8	0.7	0.9	1.3	1.7	1.3	1.4	1.8
All races 3/, female										
2002.....	6.5	*	0.6	0.8	0.7	1.0	1.0	0.8	1.1	1.3
2001.....	6.6	*	1.0	0.7	0.7	0.9	1.0	1.0	1.2	1.3
2000.....	6.7	*	0.9	0.6	0.6	0.9	0.9	1.0	0.9	1.6
1999.....	6.7	*	0.7	0.5	0.8	0.9	1.1	1.1	1.2	1.5
White, both sexes										
2002.....	8.2	*	0.7	0.8	0.8	1.2	1.3	1.0	1.3	1.5
2001.....	8.3	0.6	1.1	0.7	0.8	1.2	1.3	1.1	1.4	1.5
2000.....	8.3	1.2	1.0	0.7	0.9	1.2	1.2	1.1	1.1	1.6
1999.....	8.3	0.8	0.9	0.7	0.8	1.1	1.4	1.2	1.3	1.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
White, male										
2002.....	9.3	*	0.8	0.7	0.9	1.4	1.5	1.2	1.5	1.6
2001.....	9.3	*	1.1	0.8	0.8	1.4	1.6	1.3	1.6	1.7
2000.....	9.4	*	0.9	0.8	1.1	1.4	1.5	1.3	1.4	1.7
1999.....	9.2	*	0.9	0.8	0.9	1.2	1.8	1.3	1.4	1.8
White, female										
2002.....	7.1	*	0.6	0.8	0.7	1.0	1.1	0.8	1.1	1.4
2001.....	7.3	*	1.1	0.7	0.8	0.9	1.0	0.9	1.2	1.4
2000.....	7.4	*	1.0	0.7	0.7	0.9	0.9	0.9	0.9	1.6
1999.....	7.3	*	0.8	0.6	0.8	1.0	1.0	1.1	1.2	1.5
Black, both sexes										
2002.....	4.7	*	*	*	0.7	0.9	0.8	1.1	1.4	1.5
2001.....	4.7	*	*	0.8	0.9	0.9	1.2	1.4	1.4	1.8
2000.....	4.8	*	*	*	0.6	1.5	1.3	1.4	1.8	1.7
1999.....	4.7	*	*	*	1.1	1.2	1.7	1.1	1.4	1.8
Black, male										
2002.....	5.1	*	*	*	*	*	*	*	1.6	1.9
2001.....	5.2	*	*	*	1.5	*	*	1.7	1.8	2.3
2000.....	5.3	*	*	*	*	2.0	1.8	*	2.5	1.4
1999.....	5.2	*	*	*	1.5	1.5	1.7	*	1.7	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
Black, female										
2002.....	4.3	*	*	*	*	*	*	*	*	*
2001.....	4.3	*	*	*	*	*	*	*	*	1.4
2000.....	4.2	*	*	*	*	*	*	1.5	*	1.9
1999.....	4.3	*	*	*	*	*	1.8	*	*	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Lymphoid leukemia (C91)										
All races 3/, both sexes										
2002.....	2.2	*	0.2	0.4	0.4	0.5	0.5	0.3	0.4	0.4
2001.....	2.2	*	0.3	0.4	0.4	0.5	0.5	0.4	0.4	0.4
2000.....	2.2	*	0.3	0.4	0.5	0.5	0.4	0.3	0.2	0.3
1999.....	2.2	*	0.3	0.3	0.4	0.5	0.5	0.4	0.3	0.3
All races 3/, male										
2002.....	2.6	*	*	0.4	0.5	0.7	0.6	0.4	0.4	0.5
2001.....	2.5	*	0.3	0.4	0.5	0.7	0.7	0.4	0.5	0.5
2000.....	2.6	*	0.3	0.4	0.6	0.7	0.7	0.4	0.3	0.4
1999.....	2.6	*	0.3	0.4	0.4	0.6	0.7	0.4	0.4	0.4
All races 3/, female										
2002.....	1.8	*	*	0.4	0.4	0.4	0.4	*	0.3	0.3
2001.....	1.8	*	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.3
2000.....	1.8	*	0.3	0.3	0.3	0.4	0.2	0.2	*	0.2
1999.....	1.9	*	0.3	0.2	0.3	0.3	0.4	0.3	0.2	0.3
White, both sexes										
2002.....	2.4	*	0.2	0.4	0.5	0.5	0.6	0.3	0.4	0.4
2001.....	2.4	*	0.3	0.4	0.4	0.6	0.6	0.4	0.4	0.3
2000.....	2.4	*	0.4	0.4	0.5	0.5	0.5	0.3	0.2	0.3
1999.....	2.5	*	0.4	0.4	0.4	0.5	0.6	0.4	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Myeloid leukemia (C92)										
All races 3/, both sexes										
2002.....	3.4	*	0.3	0.2	0.2	0.4	0.4	0.5	0.7	0.8
2001.....	3.5	*	0.5	0.2	0.2	0.4	0.5	0.6	0.8	1.0
2000.....	3.5	*	0.4	0.2	0.3	0.5	0.5	0.6	0.8	1.1
1999.....	3.3	*	0.4	0.2	0.3	0.4	0.6	0.6	0.7	1.0
All races 3/, male										
2002.....	3.9	*	0.4	*	0.2	0.4	0.4	0.6	0.8	0.9
2001.....	3.9	*	0.5	0.2	0.2	0.4	0.5	0.7	0.8	1.0
2000.....	3.9	*	0.4	*	0.3	0.5	0.6	0.6	0.9	0.9
1999.....	3.7	*	0.4	*	0.3	0.4	0.7	0.6	0.7	1.0
All races 3/, female										
2002.....	3.0	*	*	0.2	0.2	0.4	0.4	0.4	0.6	0.8
2001.....	3.1	*	0.5	*	0.3	0.4	0.5	0.5	0.7	0.9
2000.....	3.1	*	0.3	*	0.2	0.4	0.4	0.6	0.6	1.2
1999.....	3.0	*	0.3	*	0.3	0.4	0.5	0.6	0.6	0.9
White, both sexes										
2002.....	3.8	*	0.3	0.2	0.2	0.4	0.4	0.5	0.7	0.8
2001.....	3.8	*	0.5	0.2	0.2	0.4	0.5	0.6	0.8	0.9
2000.....	3.8	*	0.4	0.2	0.3	0.4	0.5	0.5	0.7	1.1
1999.....	3.6	*	0.4	0.1	0.3	0.3	0.6	0.6	0.7	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified leukemia (C94-C95)										
All races 3/, both sexes										
2002.....	1.8	*	*	0.2	0.1	0.2	0.2	0.2	0.2	0.2
2001.....	1.8	*	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2
2000.....	1.9	*	0.2	0.1	0.1	0.2	0.2	0.3	0.2	0.3
1999.....	1.9	*	*	0.2	0.2	0.2	0.3	0.2	0.3	0.3
All races 3/, male										
2002.....	2.0	*	*	*	*	0.3	0.3	0.2	0.3	0.2
2001.....	2.0	*	0.3	*	0.2	*	0.3	0.3	0.3	0.3
2000.....	2.0	*	*	*	*	0.2	0.2	0.4	0.2	0.3
1999.....	2.1	*	*	0.2	*	0.3	0.3	0.2	0.3	0.3
All races 3/, female										
2002.....	1.6	*	*	*	*	*	*	*	0.2	0.2
2001.....	1.6	*	*	*	*	0.2	*	*	*	0.2
2000.....	1.7	*	*	*	*	0.2	0.2	0.2	*	0.2
1999.....	1.8	*	*	*	*	0.2	*	*	0.3	0.3
White, both sexes										
2002.....	2.0	*	*	0.1	*	0.2	0.2	0.2	0.2	0.2
2001.....	2.0	*	0.3	0.1	0.2	0.2	0.2	0.2	0.2	0.3
2000.....	2.1	*	0.2	0.1	*	0.2	0.2	0.3	0.2	0.3
1999.....	2.1	*	*	0.2	0.1	0.3	0.3	0.2	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)										
All races 3/, both sexes										
2002.....	3.8	0.9	0.3	0.2	0.2	0.1	0.2	0.2	0.3	0.4
2001.....	4.0	0.5	0.3	0.2	0.2	0.1	0.2	0.2	0.3	0.5
2000.....	4.0	0.7	0.3	0.3	0.2	0.1	0.2	0.2	0.3	0.5
1999.....	4.0	0.8	0.4	0.3	0.2	0.2	0.2	0.2	0.4	0.4
All races 3/, male										
2002.....	4.0	*	0.3	*	0.2	*	0.2	0.2	0.3	0.4
2001.....	4.2	*	0.3	0.2	0.2	*	*	0.2	0.3	0.6
2000.....	4.2	*	0.4	0.3	*	*	0.3	0.3	0.3	0.6
1999.....	4.2	*	0.4	0.2	*	*	0.3	0.2	0.5	0.5
All races 3/, female										
2002.....	3.7	*	0.3	*	*	*	*	*	0.3	0.3
2001.....	3.9	*	0.3	0.2	0.2	*	*	*	0.3	0.5
2000.....	3.9	*	*	0.2	*	*	*	0.2	0.3	0.4
1999.....	3.8	*	0.3	0.3	*	0.2	*	*	0.3	0.4
White, both sexes										
2002.....	4.3	1.0	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.4
2001.....	4.5	*	0.3	0.2	0.2	0.1	0.1	0.2	0.3	0.5
2000.....	4.5	0.7	0.3	0.3	0.2	*	0.2	0.2	0.4	0.5
1999.....	4.4	0.7	0.3	0.3	0.2	0.2	0.2	0.2	0.4	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

All races 3/, both sexes

2002.....	3.4	2.0	0.5	0.2	0.2	0.4	0.5	0.7	0.9	1.2
2001.....	3.3	2.6	0.5	0.2	0.2	0.4	0.5	0.6	1.1	1.2
2000.....	3.3	1.8	0.5	0.2	0.2	0.3	0.5	0.6	0.9	1.3
1999.....	3.2	2.5	0.5	0.2	0.2	0.4	0.6	0.6	1.0	1.2

All races 3/, male

2002.....	2.9	2.5	0.5	*	0.2	0.4	0.6	0.9	1.0	1.2
2001.....	2.8	2.9	0.6	0.2	0.3	0.4	0.5	0.7	1.2	1.3
2000.....	2.8	2.1	0.5	0.3	0.2	0.3	0.5	0.6	0.9	1.2
1999.....	2.8	3.0	0.6	0.2	0.3	0.4	0.7	0.7	1.1	1.4

All races 3/, female

2002.....	3.8	1.5	0.5	0.2	*	0.3	0.5	0.6	0.8	1.2
2001.....	3.9	2.4	0.3	0.2	*	0.4	0.5	0.5	1.0	1.1
2000.....	3.8	1.6	0.6	*	0.2	0.3	0.5	0.6	1.0	1.3
1999.....	3.7	1.9	0.4	0.3	*	0.4	0.6	0.5	1.0	1.1

White, both sexes

2002.....	3.1	1.9	0.4	0.2	*	0.2	0.3	0.3	0.5	0.6
2001.....	3.1	2.3	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.5
2000.....	3.1	1.7	0.4	0.1	0.2	0.2	0.2	0.2	0.4	0.6
1999.....	3.0	1.8	0.4	0.2	0.2	0.3	0.2	0.2	0.4	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)										
White, male										
2002.....	2.6	2.5	0.5	*	*	0.3	0.3	0.4	0.6	0.7
2001.....	2.6	2.4	0.6	*	0.3	*	0.2	0.4	0.5	0.6
2000.....	2.5	1.7	0.4	*	*	*	*	*	0.4	0.6
1999.....	2.5	2.2	0.5	*	*	0.3	*	0.3	0.4	0.7
White, female										
2002.....	3.6	*	0.4	*	*	*	0.3	0.3	0.4	0.5
2001.....	3.6	2.1	*	*	*	*	*	*	0.5	0.4
2000.....	3.6	1.7	0.4	*	*	*	*	*	0.3	0.6
1999.....	3.4	1.4	*	*	*	0.3	*	*	0.4	0.4
Black, both sexes										
2002.....	6.0	*	0.8	*	*	0.9	2.1	3.2	3.8	5.5
2001.....	5.9	3.5	*	*	*	1.1	2.0	2.4	4.7	5.5
2000.....	5.8	*	1.2	0.6	0.6	0.7	2.1	3.0	4.7	5.7
1999.....	5.7	5.5	1.4	0.6	0.6	0.9	2.8	2.9	5.1	5.8
Black, male										
2002.....	5.5	*	*	*	*	*	2.7	4.1	4.3	5.2
2001.....	5.2	*	*	*	*	1.3	2.2	2.9	5.6	5.9
2000.....	5.1	*	*	*	*	*	2.1	3.0	4.4	5.8
1999.....	5.3	6.5	*	*	*	*	3.3	3.3	5.4	6.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

Black, female

2002.....	6.4	*	*	*	*	*	1.6	2.3	3.3	5.7
2001.....	6.4	*	*	*	*	*	1.8	1.9	3.8	5.0
2000.....	6.4	*	*	*	*	*	2.2	3.0	5.0	5.6
1999.....	6.2	*	*	*	*	*	2.3	2.5	4.7	5.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Anemias (D50-D64)										
White, male										
2002.....	1.1	*	*	*	*	*	*	*	*	*
2001.....	1.1	*	*	*	*	*	*	*	*	*
2000.....	1.1	*	*	*	*	*	*	*	*	*
1999.....	1.1	*	*	*	*	*	*	*	*	*
White, female										
2002.....	1.8	*	*	*	*	*	*	*	*	*
2001.....	1.9	*	*	*	*	*	*	*	*	*
2000.....	1.9	*	*	*	*	*	*	*	*	*
1999.....	1.9	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002.....	2.8	*	*	*	*	0.9	1.7	2.5	2.3	2.6
2001.....	2.7	*	*	*	*	0.9	1.5	1.9	2.5	2.1
2000.....	2.6	*	*	*	*	*	1.6	1.7	2.8	2.2
1999.....	2.8	*	0.8	*	*	0.7	2.2	2.1	3.0	2.2
Black, male										
2002.....	2.7	*	*	*	*	*	2.3	3.2	2.5	2.7
2001.....	2.5	*	*	*	*	*	1.6	2.5	3.3	2.2
2000.....	2.3	*	*	*	*	*	1.7	1.6	2.7	1.9
1999.....	2.7	*	*	*	*	*	2.8	2.3	3.3	2.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Anemias (D50-D64)										
Black, female										
2002.....	3.0	*	*	*	*	*	*	1.8	2.2	2.4
2001.....	2.8	*	*	*	*	*	1.5	*	1.8	2.1
2000.....	3.0	*	*	*	*	*	1.6	1.9	2.9	2.5
1999.....	3.0	*	*	*	*	*	1.7	1.8	2.8	2.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hemolytic anemias (D55-D59)										
White, male										
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
White, female										
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002.....	1.4	*	*	*	*	0.7	1.6	2.2	2.3	2.3
2001.....	1.4	*	*	*	*	0.7	1.4	1.6	2.3	1.8
2000.....	1.4	*	*	*	*	*	1.5	1.6	2.4	2.0
1999.....	1.5	*	0.8	*	*	0.6	1.9	1.9	2.8	2.1
Black, male										
2002.....	1.4	*	*	*	*	*	2.0	2.9	2.4	2.5
2001.....	1.4	*	*	*	*	*	1.5	2.0	3.1	1.9
2000.....	1.3	*	*	*	*	*	1.5	1.6	2.1	1.9
1999.....	1.7	*	*	*	*	*	2.6	2.2	3.1	2.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hemolytic anemias (D55-D59)										
Black, female										
2002.....	1.4	*	*	*	*	*	*	1.7	2.2	2.1
2001.....	1.3	*	*	*	*	*	*	*	1.5	1.8
2000.....	1.4	*	*	*	*	*	1.5	1.6	2.6	2.0
1999.....	1.3	*	*	*	*	*	*	1.8	2.5	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)										
All races 3/, both sexes										
2002.....	0.9	0.6	*	*	*	*	0.1	0.1	0.2	0.2
2001.....	0.9	0.8	*	*	*	*	*	0.1	0.2	0.3
2000.....	0.9	0.6	*	*	*	*	0.1	0.1	0.1	0.3
1999.....	0.9	1.1	*	*	*	*	0.1	0.1	0.2	0.3
All races 3/, male										
2002.....	0.8	*	*	*	*	*	*	*	0.3	0.2
2001.....	0.8	*	*	*	*	*	*	*	*	0.3
2000.....	0.8	*	*	*	*	*	*	*	*	0.2
1999.....	0.8	1.4	*	*	*	*	*	*	*	0.3
All races 3/, female										
2002.....	1.0	*	*	*	*	*	*	*	*	0.3
2001.....	1.0	*	*	*	*	*	*	*	0.4	0.2
2000.....	1.0	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.0	*	*	*	*	*	*	*	0.2	0.3
White, both sexes										
2002.....	1.0	*	*	*	*	*	*	0.1	0.2	0.2
2001.....	0.9	0.7	*	*	*	*	*	*	0.2	0.2
2000.....	0.9	*	*	*	*	*	*	*	*	0.3
1999.....	0.9	0.9	*	*	*	*	*	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Certain disorders involving the immune mechanism (D80-D89)

All races 3/, both sexes

2002.....	0.5	0.5	*	*	*	*	*	*	0.2	0.5
2001.....	0.5	0.8	*	*	*	*	*	*	0.3	0.5
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	0.5
1999.....	0.4	*	0.2	*	*	*	*	0.1	0.3	0.5

All races 3/, male

2002.....	0.5	*	*	*	*	*	*	*	0.3	0.5
2001.....	0.4	*	*	*	*	*	*	*	0.4	0.6
2000.....	0.5	*	*	*	*	*	*	*	0.3	0.6
1999.....	0.4	*	*	*	*	*	*	*	0.4	0.5

All races 3/, female

2002.....	0.6	*	*	*	*	*	*	*	*	0.5
2001.....	0.6	*	*	*	*	*	*	*	*	0.4
2000.....	0.5	*	*	*	*	*	*	*	0.2	0.5
1999.....	0.4	*	*	*	*	*	*	*	0.3	0.4

White, both sexes

2002.....	0.3	*	*	*	*	*	*	*	*	0.2
2001.....	0.3	0.8	*	*	*	*	*	*	0.2	0.2
2000.....	0.3	*	*	*	*	*	*	*	0.1	0.2
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Certain disorders involving the immune mechanism (D80-D89)										
White, male										
2002.....	0.3	*	*	*	*	*	*	*	*	0.3
2001.....	0.3	*	*	*	*	*	*	*	0.2	0.3
2000.....	0.3	*	*	*	*	*	*	*	*	0.2
1999.....	0.3	*	*	*	*	*	*	*	*	0.2
White, female										
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002.....	1.9	*	*	*	*	*	*	*	1.0	2.5
2001.....	1.8	*	*	*	*	*	*	*	1.4	2.6
2000.....	1.8	*	*	*	*	*	*	0.7	1.3	2.8
1999.....	1.5	*	*	*	*	*	*	*	1.5	2.5
Black, male										
2002.....	1.7	*	*	*	*	*	*	*	*	2.2
2001.....	1.6	*	*	*	*	*	*	*	1.9	3.0
2000.....	1.6	*	*	*	*	*	*	*	*	3.2
1999.....	1.3	*	*	*	*	*	*	*	1.7	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
IV. Endocrine, nutritional and metabolic diseases (E00-E88)										
All races 3/, both sexes										
2002.....	34.5	6.6	0.9	0.5	0.6	0.9	1.7	2.5	3.7	5.6
2001.....	34.0	7.2	1.2	0.5	0.6	1.0	1.5	2.3	3.2	5.3
2000.....	33.5	6.8	1.0	0.4	0.7	1.0	1.6	2.3	3.4	4.9
1999.....	33.3	6.9	1.0	0.5	0.6	1.1	1.6	2.2	3.1	4.9
All races 3/, male										
2002.....	31.9	7.4	0.8	0.6	0.7	0.9	1.9	2.7	4.4	6.3
2001.....	30.9	8.0	1.2	0.4	0.7	1.0	1.5	2.4	3.6	6.1
2000.....	30.2	7.4	1.1	0.4	0.7	1.0	1.7	2.6	4.1	5.6
1999.....	29.8	7.3	1.1	0.5	0.6	1.1	1.5	2.2	3.6	5.8
All races 3/, female										
2002.....	37.0	5.8	1.0	0.3	0.5	0.9	1.6	2.3	2.9	4.8
2001.....	37.0	6.4	1.2	0.5	0.5	1.0	1.6	2.2	2.8	4.5
2000.....	36.7	6.2	1.0	0.3	0.6	1.0	1.4	2.0	2.7	4.3
1999.....	36.6	6.5	0.9	0.5	0.6	1.1	1.7	2.1	2.7	4.1
White, both sexes										
2002.....	34.4	6.2	1.0	0.5	0.6	0.9	1.6	2.3	3.2	5.0
2001.....	34.0	6.8	1.2	0.5	0.6	0.9	1.4	2.0	2.8	4.7
2000.....	33.5	6.3	1.1	0.4	0.7	1.0	1.5	1.9	3.0	4.4
1999.....	33.1	6.0	1.1	0.4	0.6	1.1	1.6	2.0	2.7	4.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
IV. Endocrine, nutritional and metabolic diseases (E00-E88)										
White, male										
2002.....	32.2	7.6	0.8	0.7	0.7	0.9	1.6	2.5	3.9	5.7
2001.....	31.2	7.2	1.3	0.5	0.7	1.0	1.4	2.0	3.2	5.3
2000.....	30.6	6.8	1.2	0.4	0.8	1.0	1.7	2.3	3.7	5.0
1999.....	29.9	6.3	1.2	0.5	0.6	1.2	1.4	2.1	3.0	5.1
White, female										
2002.....	36.6	4.7	1.1	0.3	0.5	1.0	1.5	2.1	2.6	4.2
2001.....	36.8	6.3	1.2	0.5	0.5	0.9	1.5	2.0	2.5	4.0
2000.....	36.3	5.7	1.0	0.4	0.6	1.0	1.4	1.6	2.3	3.8
1999.....	36.2	5.7	0.9	0.4	0.6	1.1	1.7	2.0	2.4	3.4
Black, both sexes										
2002.....	42.6	8.3	0.8	*	0.7	1.0	2.9	4.9	7.5	10.7
2001.....	41.6	10.3	0.8	*	0.6	1.1	2.5	4.5	6.4	10.5
2000.....	41.4	8.9	0.8	*	*	1.1	2.5	5.3	6.8	9.6
1999.....	41.4	11.4	0.8	0.6	0.6	1.1	2.2	3.6	6.4	10.2
Black, male										
2002.....	36.6	7.0	*	*	*	1.2	3.5	5.5	9.3	12.0
2001.....	35.9	11.7	*	*	*	*	2.4	5.2	7.8	12.8
2000.....	34.6	9.9	*	*	*	*	2.7	5.5	8.0	10.7
1999.....	35.2	12.7	*	*	*	*	2.3	3.4	7.8	11.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
IV. Endocrine, nutritional and metabolic diseases (E00-E88)										
Black, female										
2002.....	48.0	9.7	*	*	*	*	2.4	4.3	5.8	9.6
2001.....	46.8	8.8	*	*	*	*	2.5	3.9	5.2	8.6
2000.....	47.6	7.9	*	*	*	*	2.2	5.1	5.7	8.6
1999.....	47.1	10.1	*	*	*	*	2.1	3.9	5.1	8.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Endocrine diseases (E00-E34)										
All races 3/, both sexes										
2002.....	26.4	1.0	*	*	0.1	0.3	0.7	1.3	2.1	3.7
2001.....	26.1	1.1	*	*	0.2	0.3	0.6	1.2	2.0	3.5
2000.....	25.6	1.2	0.1	*	0.1	0.3	0.6	1.2	2.2	3.2
1999.....	25.5	1.0	0.1	*	0.1	0.3	0.7	1.1	2.0	3.4
All races 3/, male										
2002.....	24.8	1.1	*	*	0.2	0.3	0.8	1.4	2.6	4.2
2001.....	24.1	1.3	*	*	0.2	0.3	0.7	1.2	2.3	4.1
2000.....	23.4	1.3	*	*	*	0.3	0.7	1.3	2.7	3.7
1999.....	23.3	*	*	*	*	0.3	0.7	1.1	2.3	4.0
All races 3/, female										
2002.....	28.0	*	*	*	*	0.3	0.6	1.2	1.7	3.1
2001.....	28.0	*	*	*	*	0.3	0.5	1.2	1.6	2.9
2000.....	27.7	1.1	*	*	*	0.2	0.6	1.1	1.7	2.8
1999.....	27.5	1.3	*	*	*	0.3	0.7	1.1	1.8	2.8
White, both sexes										
2002.....	26.0	1.1	*	*	*	0.3	0.6	1.1	1.9	3.2
2001.....	25.7	1.1	*	*	0.1	0.2	0.4	0.9	1.7	3.0
2000.....	25.1	1.0	*	*	*	0.3	0.5	0.9	1.9	2.8
1999.....	24.9	1.0	*	*	*	0.3	0.6	1.0	1.7	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Endocrine diseases (E00-E34)										
White, male										
2002.....	24.8	1.3	*	*	*	0.3	0.6	1.2	2.2	3.7
2001.....	24.1	*	*	*	*	0.3	0.5	0.9	1.9	3.5
2000.....	23.5	*	*	*	*	0.3	0.6	1.0	2.4	3.2
1999.....	23.2	*	*	*	*	0.3	0.6	1.0	1.9	3.5
White, female										
2002.....	27.1	*	*	*	*	0.3	0.5	1.0	1.5	2.7
2001.....	27.2	*	*	*	*	*	0.4	0.9	1.5	2.5
2000.....	26.8	*	*	*	*	*	0.4	0.8	1.3	2.5
1999.....	26.5	1.4	*	*	*	0.3	0.5	1.0	1.5	2.3
Black, both sexes										
2002.....	34.5	*	*	*	*	*	1.6	3.1	4.6	7.6
2001.....	34.0	*	*	*	*	0.6	1.4	3.4	4.4	7.3
2000.....	33.8	*	*	*	*	*	1.5	3.3	4.5	6.7
1999.....	34.0	*	*	*	*	*	1.2	2.1	4.4	7.1
Black, male										
2002.....	29.5	*	*	*	*	*	1.8	3.3	6.0	8.7
2001.....	29.2	*	*	*	*	*	1.5	4.0	5.8	9.3
2000.....	28.1	*	*	*	*	*	1.5	3.5	5.0	7.6
1999.....	28.5	*	*	*	*	*	*	1.8	5.2	8.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Endocrine diseases (E00-E34)										
Black, female										
2002.....	39.0	*	*	*	*	*	1.4	2.8	3.4	6.6
2001.....	38.3	*	*	*	*	*	1.3	2.8	3.1	5.5
2000.....	39.1	*	*	*	*	*	1.5	3.2	4.1	5.8
1999.....	39.0	*	*	*	*	*	*	2.5	3.8	6.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diabetes mellitus (E10-E14)										
All races 3/, both sexes										
2002.....	25.4	*	*	*	0.1	0.3	0.6	1.2	2.0	3.5
2001.....	25.1	*	*	*	0.2	0.2	0.5	1.1	1.8	3.3
2000.....	24.6	*	*	*	*	0.2	0.6	1.1	2.0	3.0
1999.....	24.5	*	*	*	0.1	0.3	0.6	1.0	1.9	3.2
All races 3/, male										
2002.....	24.2	*	*	*	*	0.3	0.7	1.3	2.4	4.1
2001.....	23.5	*	*	*	0.2	0.3	0.6	1.2	2.2	3.9
2000.....	22.9	*	*	*	*	0.3	0.7	1.2	2.5	3.5
1999.....	22.8	*	*	*	*	0.2	0.6	1.0	2.2	3.8
All races 3/, female										
2002.....	26.5	*	*	*	*	0.3	0.5	1.1	1.5	3.0
2001.....	26.6	*	*	*	*	0.2	0.5	1.1	1.5	2.7
2000.....	26.3	*	*	*	*	*	0.5	0.9	1.6	2.5
1999.....	26.2	*	*	*	*	0.3	0.5	1.0	1.6	2.6
White, both sexes										
2002.....	24.9	*	*	*	*	0.2	0.5	1.0	1.7	3.1
2001.....	24.6	*	*	*	*	0.2	0.4	0.8	1.6	2.8
2000.....	24.1	*	*	*	*	0.2	0.5	0.8	1.8	2.7
1999.....	23.9	*	*	*	*	0.2	0.5	0.9	1.6	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diabetes mellitus (E10-E14)										
White, male										
2002.....	24.2	*	*	*	*	*	0.5	1.1	2.1	3.7
2001.....	23.5	*	*	*	*	*	0.4	0.8	1.8	3.3
2000.....	22.9	*	*	*	*	0.3	0.5	1.0	2.3	3.1
1999.....	22.7	*	*	*	*	0.3	0.5	0.9	1.8	3.4
White, female										
2002.....	25.6	*	*	*	*	*	0.4	0.9	1.3	2.5
2001.....	25.7	*	*	*	*	*	0.3	0.8	1.4	2.3
2000.....	25.3	*	*	*	*	*	0.4	0.7	1.2	2.3
1999.....	25.0	*	*	*	*	*	0.4	0.9	1.4	2.1
Black, both sexes										
2002.....	33.6	*	*	*	*	*	1.4	2.8	4.3	7.3
2001.....	33.1	*	*	*	*	*	1.4	3.2	4.2	6.9
2000.....	32.8	*	*	*	*	*	1.5	2.8	4.1	6.0
1999.....	33.0	*	*	*	*	*	1.1	1.8	4.1	6.6
Black, male										
2002.....	29.0	*	*	*	*	*	1.7	3.0	5.7	8.5
2001.....	28.5	*	*	*	*	*	1.5	3.9	5.6	8.7
2000.....	27.4	*	*	*	*	*	1.5	3.1	4.5	7.1
1999.....	27.7	*	*	*	*	*	*	1.6	5.0	7.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diabetes mellitus (E10-E14)										
Black, female										
2002.....	37.8	*	*	*	*	*	*	2.6	3.1	6.3
2001.....	37.2	*	*	*	*	*	1.3	2.7	2.9	5.3
2000.....	37.8	*	*	*	*	*	1.5	2.5	3.8	5.0
1999.....	37.8	*	*	*	*	*	*	2.0	3.3	5.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nutritional diseases (E40-E68)										
All races 3/, both sexes										
2002.....	2.6	*	*	*	*	*	0.3	0.6	0.9	1.2
2001.....	2.4	*	*	*	*	*	0.3	0.4	0.7	1.2
2000.....	2.6	*	*	*	*	*	0.2	0.5	0.7	1.1
1999.....	2.5	*	*	*	*	0.1	0.2	0.4	0.6	0.9
All races 3/, male										
2002.....	2.2	*	*	*	*	*	0.4	0.6	1.1	1.4
2001.....	1.9	*	*	*	*	*	0.3	0.6	0.8	1.2
2000.....	2.0	*	*	*	*	*	0.2	0.6	0.9	1.2
1999.....	2.0	*	*	*	*	*	*	0.4	0.8	1.1
All races 3/, female										
2002.....	3.0	*	*	*	*	*	0.3	0.5	0.8	1.1
2001.....	2.9	*	*	*	*	*	0.3	0.3	0.7	1.2
2000.....	3.1	*	*	*	*	*	*	0.4	0.5	0.9
1999.....	2.9	*	*	*	*	*	0.3	0.4	0.4	0.8
White, both sexes										
2002.....	2.6	*	*	*	*	*	0.2	0.5	0.7	1.1
2001.....	2.4	*	*	*	*	*	0.2	0.4	0.6	1.0
2000.....	2.6	*	*	*	*	*	0.1	0.3	0.5	0.9
1999.....	2.5	*	*	*	*	*	0.2	0.3	0.5	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nutritional diseases (E40-E68)										
White, male										
2002.....	2.2	*	*	*	*	*	0.3	0.5	0.8	1.3
2001.....	1.9	*	*	*	*	*	*	0.5	0.7	1.0
2000.....	2.1	*	*	*	*	*	*	0.4	0.7	1.2
1999.....	2.0	*	*	*	*	*	*	0.3	0.7	0.9
White, female										
2002.....	3.0	*	*	*	*	*	*	0.4	0.6	0.9
2001.....	2.9	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	3.1	*	*	*	*	*	*	*	0.3	0.7
1999.....	3.0	*	*	*	*	*	*	0.3	0.3	0.6
Black, both sexes										
2002.....	3.3	*	*	*	*	*	0.7	1.4	2.4	2.3
2001.....	3.0	*	*	*	*	*	0.8	0.8	1.7	2.8
2000.....	3.1	*	*	*	*	*	*	1.3	1.6	2.1
1999.....	3.0	*	*	*	*	*	*	1.2	1.3	2.1
Black, male										
2002.....	2.6	*	*	*	*	*	*	1.6	2.9	2.3
2001.....	2.5	*	*	*	*	*	*	*	1.7	2.8
2000.....	2.5	*	*	*	*	*	*	*	2.1	2.2
1999.....	2.6	*	*	*	*	*	*	*	1.8	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Obesity and other hyperalimentation (E65-E68)										
All races 3/, both sexes										
2002.....	1.3	*	*	*	*	*	0.3	0.5	0.9	1.1
2001.....	1.1	*	*	*	*	*	0.3	0.4	0.7	1.1
2000.....	1.1	*	*	*	*	*	0.2	0.4	0.6	0.9
1999.....	0.9	*	*	*	*	*	0.2	0.4	0.6	0.8
All races 3/, male										
2002.....	1.2	*	*	*	*	*	0.3	0.5	1.1	1.3
2001.....	1.0	*	*	*	*	*	0.2	0.5	0.7	1.1
2000.....	1.0	*	*	*	*	*	0.2	0.5	0.8	1.1
1999.....	0.9	*	*	*	*	*	*	0.4	0.8	1.0
All races 3/, female										
2002.....	1.3	*	*	*	*	*	0.2	0.5	0.7	0.9
2001.....	1.2	*	*	*	*	*	0.3	0.3	0.6	1.1
2000.....	1.1	*	*	*	*	*	*	0.3	0.4	0.7
1999.....	1.0	*	*	*	*	*	0.2	0.4	0.4	0.7
White, both sexes										
2002.....	1.2	*	*	*	*	*	0.2	0.4	0.7	1.0
2001.....	1.0	*	*	*	*	*	0.2	0.4	0.6	0.9
2000.....	1.0	*	*	*	*	*	*	0.3	0.5	0.8
1999.....	0.9	*	*	*	*	*	0.1	0.3	0.5	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Obesity and other hyperalimentation (E65-E68)										
White, male										
2002.....	1.2	*	*	*	*	*	*	0.4	0.8	1.2
2001.....	1.0	*	*	*	*	*	*	0.5	0.7	1.0
2000.....	1.0	*	*	*	*	*	*	0.4	0.7	1.1
1999.....	0.9	*	*	*	*	*	*	0.3	0.6	0.8
White, female										
2002.....	1.2	*	*	*	*	*	*	0.4	0.5	0.8
2001.....	1.1	*	*	*	*	*	*	*	0.5	0.9
2000.....	1.0	*	*	*	*	*	*	*	0.3	0.6
1999.....	0.9	*	*	*	*	*	*	*	0.3	0.5
Black, both sexes										
2002.....	2.0	*	*	*	*	*	0.7	1.4	2.3	2.0
2001.....	1.8	*	*	*	*	*	0.7	0.8	1.6	2.6
2000.....	1.6	*	*	*	*	*	*	1.2	1.5	1.8
1999.....	1.4	*	*	*	*	*	*	1.2	1.3	1.9
Black, male										
2002.....	1.5	*	*	*	*	*	*	1.6	2.9	2.1
2001.....	1.3	*	*	*	*	*	*	*	1.6	2.6
2000.....	1.3	*	*	*	*	*	*	*	1.8	2.1
1999.....	1.1	*	*	*	*	*	*	*	1.8	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Metabolic disorders (E70-E88)										
All races 3/, both sexes										
2002.....	5.5	5.5	0.8	0.4	0.4	0.5	0.7	0.6	0.6	0.6
2001.....	5.5	6.0	1.0	0.4	0.4	0.6	0.7	0.6	0.5	0.7
2000.....	5.4	5.3	0.9	0.3	0.5	0.6	0.8	0.6	0.6	0.7
1999.....	5.3	5.6	0.8	0.4	0.5	0.7	0.7	0.6	0.5	0.6
All races 3/, male										
2002.....	4.9	6.1	0.8	0.5	0.5	0.5	0.7	0.7	0.7	0.7
2001.....	4.9	6.5	1.1	0.4	0.4	0.6	0.6	0.6	0.6	0.8
2000.....	4.8	5.5	0.9	0.3	0.6	0.6	0.8	0.8	0.6	0.7
1999.....	4.5	6.3	0.9	0.4	0.5	0.7	0.7	0.7	0.5	0.7
All races 3/, female										
2002.....	6.0	4.9	0.9	0.3	0.4	0.6	0.7	0.6	0.5	0.5
2001.....	6.1	5.5	1.0	0.4	0.4	0.6	0.7	0.6	0.5	0.5
2000.....	6.0	5.0	0.8	0.3	0.4	0.7	0.7	0.5	0.5	0.6
1999.....	6.1	4.9	0.7	0.4	0.4	0.7	0.8	0.6	0.5	0.5
White, both sexes										
2002.....	5.9	5.0	0.9	0.4	0.5	0.6	0.8	0.7	0.7	0.7
2001.....	5.9	5.6	1.1	0.4	0.4	0.7	0.8	0.7	0.6	0.7
2000.....	5.8	5.0	0.9	0.3	0.6	0.7	0.9	0.7	0.6	0.7
1999.....	5.7	4.9	0.9	0.4	0.5	0.8	0.9	0.8	0.5	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cystic fibrosis (E84)										
All races 3/, both sexes										
2002.....	0.2	*	*	*	0.1	0.3	0.5	0.4	0.3	0.2
2001.....	0.2	*	*	*	0.1	0.4	0.5	0.4	0.2	0.2
2000.....	0.2	*	*	*	0.2	0.4	0.5	0.4	0.3	0.3
1999.....	0.2	*	*	*	0.2	0.4	0.5	0.4	0.3	0.2
All races 3/, male										
2002.....	0.2	*	*	*	*	0.2	0.4	0.4	0.4	0.2
2001.....	0.2	*	*	*	*	0.3	0.4	0.4	0.3	0.2
2000.....	0.2	*	*	*	*	0.3	0.5	0.4	0.3	0.2
1999.....	0.2	*	*	*	*	0.3	0.5	0.4	0.3	*
All races 3/, female										
2002.....	0.2	*	*	*	*	0.4	0.5	0.3	0.2	0.2
2001.....	0.2	*	*	*	*	0.4	0.6	0.4	0.2	*
2000.....	0.2	*	*	*	0.2	0.4	0.5	0.3	0.4	0.3
1999.....	0.2	*	*	*	*	0.5	0.6	0.4	0.2	0.2
White, both sexes										
2002.....	0.2	*	*	*	0.2	0.4	0.5	0.5	0.4	0.2
2001.....	0.2	*	*	*	0.2	0.5	0.6	0.5	0.3	0.2
2000.....	0.2	*	*	*	0.2	0.4	0.6	0.4	0.4	0.3
1999.....	0.2	*	*	*	0.2	0.5	0.7	0.5	0.3	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other metabolic disorders (E70-E83,E85,E88)										
All races 3/, both sexes										
2002.....	2.8	3.4	0.7	0.3	0.3	0.2	0.2	0.2	0.2	0.3
2001.....	2.6	3.7	0.8	0.3	0.2	0.2	0.1	0.1	0.1	0.3
2000.....	2.4	3.2	0.8	0.2	0.3	0.2	0.2	0.2	0.2	0.2
1999.....	2.2	3.4	0.7	0.3	0.3	0.2	0.1	0.1	0.2	0.3
All races 3/, male										
2002.....	3.0	3.6	0.7	0.4	0.4	0.3	0.3	0.2	0.2	0.4
2001.....	2.9	4.1	0.9	0.3	0.3	0.3	*	*	*	0.4
2000.....	2.6	3.6	0.8	0.3	0.4	0.2	0.3	0.3	0.3	0.3
1999.....	2.4	3.7	0.8	0.3	0.3	0.3	*	*	*	0.3
All races 3/, female										
2002.....	2.5	3.2	0.7	*	0.2	*	*	*	*	*
2001.....	2.3	3.2	0.7	0.3	*	*	*	*	*	0.2
2000.....	2.3	2.8	0.7	0.2	*	*	*	*	*	0.2
1999.....	2.1	3.1	0.6	0.3	0.2	*	*	*	*	0.2
White, both sexes										
2002.....	3.0	3.3	0.7	0.3	0.3	0.2	0.2	0.2	0.2	0.3
2001.....	2.8	3.7	0.8	0.4	0.3	0.2	0.1	0.1	0.1	0.3
2000.....	2.7	3.4	0.8	0.3	0.3	0.2	0.2	0.2	0.1	0.3
1999.....	2.4	3.1	0.8	0.3	0.3	0.2	0.1	0.1	0.1	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
V. Mental and behavioral disorders (F01-F99)										
All races 3/, both sexes										
2002.....	19.3	*	*	*	0.1	0.3	0.8	1.3	2.0	4.0
2001.....	17.8	*	*	*	*	0.4	0.8	1.4	2.2	3.7
2000.....	16.4	*	*	*	*	0.4	0.9	1.3	2.3	3.9
1999.....	15.1	*	*	*	0.1	0.4	0.8	1.3	2.3	4.1
All races 3/, male										
2002.....	14.8	*	*	*	*	0.4	1.1	1.8	2.7	5.6
2001.....	13.8	*	*	*	*	0.5	1.2	2.1	3.2	5.2
2000.....	12.8	*	*	*	*	0.5	1.3	1.9	3.3	6.0
1999.....	12.3	*	*	*	*	0.5	1.1	2.0	3.3	6.0
All races 3/, female										
2002.....	23.7	*	*	*	*	0.2	0.5	0.7	1.3	2.4
2001.....	21.7	*	*	*	*	0.3	0.3	0.8	1.2	2.2
2000.....	19.8	*	*	*	*	0.3	0.5	0.7	1.3	1.8
1999.....	17.8	*	*	*	*	*	0.5	0.6	1.4	2.2
White, both sexes										
2002.....	21.3	*	*	*	*	0.4	0.9	1.3	2.1	3.9
2001.....	19.6	*	*	*	*	0.4	0.8	1.4	2.2	3.6
2000.....	17.9	*	*	*	*	0.4	0.9	1.4	2.2	3.7
1999.....	16.5	*	*	*	*	0.4	0.8	1.4	2.4	3.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
V. Mental and behavioral disorders (F01-F99)										
White, male										
2002.....	15.7	*	*	*	*	0.5	1.2	1.9	2.8	5.5
2001.....	14.5	*	*	*	*	0.5	1.2	2.0	3.2	5.1
2000.....	13.4	*	*	*	*	0.5	1.3	2.0	3.1	5.7
1999.....	12.8	*	*	*	*	0.6	1.1	2.1	3.4	5.6
White, female										
2002.....	26.7	*	*	*	*	0.3	0.5	0.7	1.3	2.3
2001.....	24.4	*	*	*	*	0.3	0.4	0.8	1.2	2.1
2000.....	22.2	*	*	*	*	0.3	0.6	0.8	1.2	1.7
1999.....	20.1	*	*	*	*	*	0.4	0.6	1.3	1.8
Black, both sexes										
2002.....	13.2	*	*	*	*	*	0.7	1.2	2.3	4.9
2001.....	12.6	*	*	*	*	*	0.8	1.4	2.3	4.9
2000.....	11.8	*	*	*	*	*	1.1	1.4	3.3	5.3
1999.....	10.7	*	*	*	*	*	0.9	1.3	2.4	6.8
Black, male										
2002.....	13.1	*	*	*	*	*	*	1.8	2.9	6.9
2001.....	13.0	*	*	*	*	*	1.4	2.5	3.4	7.1
2000.....	12.5	*	*	*	*	*	1.8	2.3	4.6	8.0
1999.....	11.9	*	*	*	*	*	*	2.0	3.1	9.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
V. Mental and behavioral disorders (F01-F99)										
Black, female										
2002.....	13.3	*	*	*	*	*	*	*	1.8	3.1
2001.....	12.1	*	*	*	*	*	*	*	1.4	3.0
2000.....	11.2	*	*	*	*	*	*	*	2.2	2.8
1999.....	9.6	*	*	*	*	*	*	*	1.8	4.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Mental and behavioral disorders due to psychoactive substance use (F10-F19)										
All races 3/, both sexes										
2002.....	3.3	*	*	*	*	0.2	0.7	1.1	1.8	3.7
2001.....	3.2	*	*	*	*	0.3	0.6	1.2	2.0	3.4
2000.....	3.1	*	*	*	*	0.2	0.7	1.1	2.1	3.6
1999.....	3.1	*	*	*	*	0.3	0.6	1.1	2.1	3.8
All races 3/, male										
2002.....	5.1	*	*	*	*	0.3	1.0	1.7	2.5	5.3
2001.....	4.9	*	*	*	*	0.4	1.1	1.8	2.9	4.9
2000.....	4.8	*	*	*	*	0.3	1.2	1.7	3.0	5.7
1999.....	4.8	*	*	*	*	0.4	1.0	1.7	3.0	5.6
All races 3/, female										
2002.....	1.6	*	*	*	*	*	0.3	0.5	1.0	2.2
2001.....	1.5	*	*	*	*	*	0.2	0.6	1.0	1.9
2000.....	1.5	*	*	*	*	*	0.3	0.6	1.1	1.6
1999.....	1.4	*	*	*	*	*	0.3	0.4	1.2	1.9
White, both sexes										
2002.....	3.2	*	*	*	*	0.3	0.7	1.1	1.8	3.6
2001.....	3.0	*	*	*	*	0.3	0.7	1.2	1.9	3.3
2000.....	3.0	*	*	*	*	0.3	0.8	1.2	2.0	3.5
1999.....	2.9	*	*	*	*	0.3	0.7	1.1	2.1	3.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Mental and behavioral disorders due to psychoactive substance use (F10-F19)										
White, male										
2002.....	4.9	*	*	*	*	0.4	1.1	1.7	2.7	5.2
2001.....	4.6	*	*	*	*	0.5	1.1	1.8	2.9	4.7
2000.....	4.6	*	*	*	*	0.4	1.2	1.8	2.9	5.5
1999.....	4.6	*	*	*	*	0.4	1.0	1.7	3.1	5.2
White, female										
2002.....	1.5	*	*	*	*	*	0.3	0.5	1.0	2.0
2001.....	1.5	*	*	*	*	*	0.3	0.6	1.0	1.8
2000.....	1.4	*	*	*	*	*	0.3	0.6	1.0	1.5
1999.....	1.3	*	*	*	*	*	0.3	0.4	1.1	1.6
Black, both sexes										
2002.....	4.4	*	*	*	*	*	*	1.1	2.0	4.7
2001.....	4.6	*	*	*	*	*	*	1.2	2.1	4.6
2000.....	4.6	*	*	*	*	*	0.9	1.1	3.1	4.8
1999.....	4.7	*	*	*	*	*	*	1.1	2.2	6.2
Black, male										
2002.....	7.0	*	*	*	*	*	*	1.6	2.7	6.6
2001.....	7.4	*	*	*	*	*	*	2.0	3.0	6.7
2000.....	7.3	*	*	*	*	*	1.5	2.0	4.3	7.2
1999.....	7.4	*	*	*	*	*	*	1.8	2.8	9.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Mental and behavioral disorders due to psychoactive substance use (F10-F19)										
Black, female										
2002.....	2.1	*	*	*	*	*	*	*	1.4	3.0
2001.....	2.1	*	*	*	*	*	*	*	*	2.7
2000.....	2.1	*	*	*	*	*	*	*	2.0	2.6
1999.....	2.3	*	*	*	*	*	*	*	1.7	3.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Mental and behavioral disorders due to use of alcohol (F10)										
All races 3/, both sexes										
2002.....	2.4	*	*	*	*	0.1	0.2	0.5	0.8	2.2
2001.....	2.3	*	*	*	*	0.1	0.2	0.5	1.0	2.2
2000.....	2.2	*	*	*	*	*	0.2	0.4	1.0	2.2
1999.....	2.2	*	*	*	*	*	0.2	0.4	1.0	2.3
All races 3/, male										
2002.....	3.8	*	*	*	*	*	0.3	0.8	1.3	3.3
2001.....	3.7	*	*	*	*	0.2	0.4	0.7	1.5	3.3
2000.....	3.6	*	*	*	*	*	0.3	0.7	1.5	3.6
1999.....	3.6	*	*	*	*	*	0.4	0.7	1.4	3.4
All races 3/, female										
2002.....	1.0	*	*	*	*	*	*	*	0.4	1.1
2001.....	1.0	*	*	*	*	*	*	0.2	0.5	1.0
2000.....	0.9	*	*	*	*	*	*	*	0.4	0.9
1999.....	0.9	*	*	*	*	*	*	*	0.6	1.1
White, both sexes										
2002.....	2.4	*	*	*	*	0.1	0.2	0.5	0.8	2.2
2001.....	2.3	*	*	*	*	0.1	0.2	0.5	1.0	2.2
2000.....	2.2	*	*	*	*	*	0.2	0.4	0.9	2.2
1999.....	2.1	*	*	*	*	*	0.3	0.4	1.0	2.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)										
All races 3/, both sexes										
2002.....	0.9	*	*	*	*	0.1	0.5	0.6	1.0	1.5
2001.....	0.9	*	*	*	*	0.1	0.4	0.7	0.9	1.3
2000.....	0.9	*	*	*	*	0.1	0.5	0.7	1.1	1.4
1999.....	0.9	*	*	*	*	0.2	0.4	0.6	1.1	1.5
All races 3/, male										
2002.....	1.3	*	*	*	*	*	0.7	0.9	1.3	1.9
2001.....	1.2	*	*	*	*	0.2	0.6	1.1	1.4	1.6
2000.....	1.2	*	*	*	*	*	0.8	1.0	1.5	2.1
1999.....	1.3	*	*	*	*	0.3	0.6	1.0	1.5	2.2
All races 3/, female										
2002.....	0.6	*	*	*	*	*	0.3	0.4	0.6	1.0
2001.....	0.5	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	0.6	*	*	*	*	*	0.2	0.4	0.7	0.8
1999.....	0.5	*	*	*	*	*	*	0.3	0.6	0.8
White, both sexes										
2002.....	0.9	*	*	*	*	0.1	0.5	0.7	1.0	1.4
2001.....	0.8	*	*	*	*	0.2	0.5	0.7	1.0	1.1
2000.....	0.8	*	*	*	*	0.2	0.6	0.7	1.0	1.3
1999.....	0.8	*	*	*	*	0.2	0.4	0.6	1.1	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)										
White, male										
2002.....	1.2	*	*	*	*	*	0.7	1.0	1.4	1.8
2001.....	1.1	*	*	*	*	0.3	0.7	1.1	1.4	1.5
2000.....	1.1	*	*	*	*	*	0.8	1.0	1.4	1.9
1999.....	1.1	*	*	*	*	0.3	0.6	1.0	1.6	2.0
White, female										
2002.....	0.6	*	*	*	*	*	*	0.3	0.6	1.0
2001.....	0.5	*	*	*	*	*	*	0.4	0.5	0.8
2000.....	0.5	*	*	*	*	*	*	0.5	0.6	0.7
1999.....	0.4	*	*	*	*	*	*	0.3	0.5	0.7
Black, both sexes										
2002.....	1.7	*	*	*	*	*	*	0.8	1.1	2.5
2001.....	1.6	*	*	*	*	*	*	0.9	1.2	2.5
2000.....	1.6	*	*	*	*	*	*	*	1.9	3.0
1999.....	1.7	*	*	*	*	*	*	0.8	1.4	3.0
Black, male										
2002.....	2.4	*	*	*	*	*	*	*	*	3.2
2001.....	2.3	*	*	*	*	*	*	*	1.6	3.0
2000.....	2.3	*	*	*	*	*	*	*	2.3	4.6
1999.....	2.5	*	*	*	*	*	*	*	1.6	4.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Mental retardation (F70-F79)										
All races 3/, both sexes										
2002.....	0.2	*	*	*	*	*	*	*	*	0.1
2001.....	0.1	*	*	*	*	*	*	*	0.1	0.1
2000.....	0.1	*	*	*	*	*	*	*	*	0.1
1999.....	0.2	*	*	*	*	*	*	0.1	0.1	0.2
All races 3/, male										
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	0.1	0.1
2000.....	0.2	*	*	*	*	*	*	*	*	0.1
1999.....	0.2	*	*	*	*	*	*	0.1	0.1	0.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
VI. Diseases of the nervous system (G00-G98)										
All races 3/, both sexes										
2002.....	35.7	9.5	2.1	1.1	1.3	1.8	2.2	2.1	2.3	3.3
2001.....	33.9	8.9	2.1	1.1	1.3	2.0	2.2	2.1	2.3	3.2
2000.....	32.3	10.2	2.1	1.1	1.4	1.8	2.0	1.9	2.3	3.2
1999.....	30.1	11.6	2.4	1.3	1.3	1.9	2.2	1.9	2.2	3.2
All races 3/, male										
2002.....	28.5	11.0	2.4	1.2	1.5	2.3	2.8	2.6	2.8	4.0
2001.....	27.4	10.0	2.2	1.1	1.4	2.7	2.9	2.5	2.3	3.8
2000.....	26.2	11.5	2.1	1.1	1.6	2.2	2.6	2.3	2.6	3.5
1999.....	25.0	12.2	2.6	1.4	1.4	2.5	3.0	2.3	2.5	3.7
All races 3/, female										
2002.....	42.8	7.8	1.7	1.0	1.1	1.2	1.6	1.7	1.8	2.5
2001.....	40.2	7.8	1.9	1.2	1.1	1.2	1.5	1.6	2.3	2.5
2000.....	38.2	8.8	2.0	1.1	1.2	1.3	1.3	1.5	2.1	2.8
1999.....	35.1	11.0	2.1	1.2	1.1	1.2	1.4	1.6	1.8	2.7
White, both sexes										
2002.....	40.2	9.3	1.8	1.1	1.2	1.7	2.0	2.0	2.2	3.1
2001.....	38.0	7.9	1.8	1.1	1.2	1.9	2.1	1.8	2.1	3.0
2000.....	36.3	9.4	1.8	1.1	1.3	1.7	2.0	1.8	2.2	3.0
1999.....	33.7	10.4	2.2	1.2	1.1	1.8	2.1	1.9	2.0	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
VI. Diseases of the nervous system (G00-G98)										
White, male										
2002.....	31.7	11.0	2.1	1.2	1.3	2.4	2.5	2.5	2.6	3.8
2001.....	30.5	8.7	2.1	1.1	1.4	2.6	2.8	2.2	2.1	3.7
2000.....	29.1	10.3	2.0	1.1	1.5	2.3	2.6	2.3	2.5	3.3
1999.....	27.7	11.8	2.5	1.2	1.2	2.5	2.8	2.3	2.3	3.6
White, female										
2002.....	48.6	7.5	1.5	0.9	1.0	1.0	1.5	1.5	1.7	2.3
2001.....	45.4	7.1	1.6	1.1	1.0	1.1	1.3	1.4	2.1	2.3
2000.....	43.2	8.4	1.6	1.0	1.1	1.1	1.3	1.3	1.9	2.6
1999.....	39.5	9.0	1.8	1.1	1.1	1.0	1.3	1.5	1.7	2.5
Black, both sexes										
2002.....	19.5	11.6	3.3	1.3	2.0	2.3	3.5	3.6	4.1	5.5
2001.....	19.0	14.4	3.1	1.5	1.8	3.0	3.0	3.9	4.2	5.4
2000.....	17.9	14.3	3.4	1.4	2.2	2.5	2.5	3.2	3.8	5.4
1999.....	16.7	17.8	3.3	1.9	1.8	2.5	3.5	2.8	3.5	5.3
Black, male										
2002.....	16.7	11.9	4.0	1.3	2.4	2.5	4.8	4.3	4.8	6.7
2001.....	16.1	17.1	2.9	1.2	1.7	4.0	3.5	5.0	4.5	6.5
2000.....	15.4	17.9	3.1	1.1	2.7	2.3	3.0	3.5	3.8	5.8
1999.....	14.6	15.9	3.2	2.3	2.4	2.9	5.0	3.3	3.9	5.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
VI. Diseases of the nervous system (G00-G98)										
Black, female										
2002.....	22.2	11.2	2.7	1.3	1.6	2.1	2.4	2.9	3.5	4.5
2001.....	21.5	11.7	3.4	1.8	1.8	1.9	2.5	2.9	4.0	4.4
2000.....	20.2	10.6	3.6	1.8	1.8	2.7	2.1	2.8	3.8	4.9
1999.....	18.5	19.8	3.4	1.5	1.2	2.0	2.1	2.3	3.1	5.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Meningitis (G00,G03)										
All races 3/, both sexes										
2002.....	0.2	1.8	*	*	0.1	*	*	*	0.1	0.1
2001.....	0.3	1.9	0.2	*	*	*	0.1	*	0.1	0.1
2000.....	0.3	2.4	0.2	*	*	*	*	*	0.1	0.2
1999.....	0.3	3.1	0.3	*	*	*	0.1	0.1	0.1	0.1
All races 3/, male										
2002.....	0.3	2.0	*	*	*	*	*	*	*	*
2001.....	0.3	1.9	*	*	*	*	*	*	*	0.2
2000.....	0.3	2.5	0.3	*	*	*	*	*	*	0.2
1999.....	0.3	3.1	0.3	*	*	*	*	*	*	*
All races 3/, female										
2002.....	0.2	1.7	*	*	*	*	*	*	*	*
2001.....	0.2	2.0	*	*	*	*	*	*	*	*
2000.....	0.2	2.3	*	*	*	*	*	*	*	*
1999.....	0.3	3.0	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.2	1.7	*	*	*	*	*	*	*	*
2001.....	0.2	1.5	*	*	*	*	*	*	0.1	*
2000.....	0.2	1.8	0.2	*	*	*	*	*	*	0.1
1999.....	0.3	2.3	0.3	*	*	*	*	*	*	0.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Multiple sclerosis (G35)										
All races 3/, both sexes										
2002.....	1.1	*	*	*	*	*	*	*	0.1	0.4
2001.....	1.1	*	*	*	*	*	*	*	0.2	0.4
2000.....	1.0	*	*	*	*	*	*	0.1	0.2	0.4
1999.....	1.0	*	*	*	*	*	*	*	0.2	0.4
All races 3/, male										
2002.....	0.8	*	*	*	*	*	*	*	*	0.4
2001.....	0.7	*	*	*	*	*	*	*	*	0.3
2000.....	0.7	*	*	*	*	*	*	*	*	0.3
1999.....	0.6	*	*	*	*	*	*	*	*	0.3
All races 3/, female										
2002.....	1.4	*	*	*	*	*	*	*	*	0.5
2001.....	1.4	*	*	*	*	*	*	*	0.2	0.5
2000.....	1.3	*	*	*	*	*	*	*	0.3	0.5
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.5
White, both sexes										
2002.....	1.2	*	*	*	*	*	*	*	*	0.4
2001.....	1.2	*	*	*	*	*	*	*	0.2	0.4
2000.....	1.1	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.0	*	*	*	*	*	*	*	0.2	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Epilepsy (G40-G41)										
All races 3/, both sexes										
2002.....	0.5	*	0.1	0.1	0.1	0.2	0.4	0.4	0.4	0.5
2001.....	0.5	*	*	0.1	*	0.2	0.3	0.4	0.4	0.5
2000.....	0.5	*	0.2	*	0.1	0.2	0.3	0.3	0.4	0.5
1999.....	0.5	*	0.2	0.1	0.1	0.2	0.3	0.4	0.4	0.5
All races 3/, male										
2002.....	0.5	*	*	*	*	*	0.4	0.5	0.5	0.6
2001.....	0.5	*	*	*	*	0.2	0.4	0.5	0.4	0.7
2000.....	0.5	*	*	*	*	*	0.4	0.4	0.5	0.5
1999.....	0.5	*	0.3	*	*	0.3	0.4	0.6	0.5	0.7
All races 3/, female										
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.4
2001.....	0.4	*	*	*	*	*	*	0.3	0.4	0.3
2000.....	0.4	*	*	*	*	0.2	*	0.2	0.4	0.4
1999.....	0.4	*	*	*	*	*	0.3	0.3	0.3	0.3
White, both sexes										
2002.....	0.4	*	*	*	*	0.2	0.3	0.4	0.4	0.5
2001.....	0.4	*	*	0.1	*	0.2	0.2	0.3	0.4	0.4
2000.....	0.4	*	0.2	*	*	0.1	0.3	0.3	0.4	0.4
1999.....	0.4	*	0.2	*	*	0.2	0.3	0.4	0.4	0.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Infantile cerebral palsy (G80)										
All races 3/, both sexes										
2002.....	0.4	*	0.6	0.5	0.5	0.6	0.7	0.5	0.2	0.2
2001.....	0.4	*	0.5	0.4	0.5	0.7	0.6	0.4	0.3	0.2
2000.....	0.4	*	0.4	0.5	0.6	0.6	0.6	0.3	0.3	0.2
1999.....	0.4	*	0.5	0.4	0.5	0.5	0.5	0.4	0.3	0.3
All races 3/, male										
2002.....	0.5	*	0.7	0.6	0.5	0.8	0.8	0.5	0.3	0.3
2001.....	0.5	*	0.5	0.4	0.6	0.8	0.7	0.4	0.4	0.2
2000.....	0.4	*	0.3	0.5	0.7	0.7	0.7	0.4	0.4	0.2
1999.....	0.4	*	0.6	0.5	0.5	0.6	0.6	0.4	0.3	0.2
All races 3/, female										
2002.....	0.4	*	0.5	0.5	0.5	0.5	0.5	0.4	0.2	0.2
2001.....	0.3	*	0.4	0.5	0.5	0.6	0.5	0.3	0.3	0.2
2000.....	0.3	*	0.4	0.6	0.4	0.5	0.5	0.3	0.3	0.2
1999.....	0.3	*	0.4	0.4	0.4	0.4	0.3	0.4	0.3	0.3
White, both sexes										
2002.....	0.4	*	0.5	0.5	0.4	0.6	0.5	0.4	0.2	0.2
2001.....	0.4	*	0.4	0.4	0.5	0.6	0.6	0.3	0.3	0.2
2000.....	0.4	*	0.3	0.5	0.5	0.6	0.5	0.3	0.3	0.2
1999.....	0.3	*	0.5	0.4	0.4	0.5	0.4	0.4	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)										
All races 3/, both sexes										
2002.....	6.9	6.9	1.1	0.3	0.6	0.8	1.0	1.0	1.3	1.8
2001.....	6.7	6.1	1.2	0.5	0.6	0.9	1.1	1.0	1.1	1.8
2000.....	6.7	7.1	1.2	0.4	0.5	0.9	1.0	1.0	1.2	1.8
1999.....	6.5	7.7	1.2	0.5	0.6	0.9	1.2	0.9	1.1	1.7
All races 3/, male										
2002.....	7.3	8.4	1.3	0.4	0.7	1.2	1.4	1.4	1.7	2.4
2001.....	7.1	7.1	1.2	0.5	0.7	1.5	1.6	1.4	1.1	2.3
2000.....	7.2	8.1	1.3	0.4	0.6	1.3	1.4	1.3	1.5	2.1
1999.....	7.0	8.6	1.2	0.6	0.6	1.5	1.7	1.1	1.3	2.2
All races 3/, female										
2002.....	6.5	5.4	0.9	0.3	0.4	0.4	0.6	0.7	0.9	1.3
2001.....	6.3	5.0	1.2	0.5	0.5	0.3	0.7	0.7	1.1	1.2
2000.....	6.3	6.0	1.1	0.4	0.5	0.5	0.6	0.6	1.0	1.4
1999.....	6.1	6.8	1.2	0.5	0.5	0.4	0.6	0.6	0.8	1.3
White, both sexes										
2002.....	7.3	6.9	1.0	0.3	0.6	0.8	1.0	1.0	1.3	1.8
2001.....	7.0	5.8	1.1	0.5	0.6	0.9	1.1	0.9	1.0	1.8
2000.....	7.1	6.9	1.1	0.4	0.5	0.9	1.0	0.9	1.2	1.7
1999.....	6.9	7.5	1.1	0.5	0.5	1.0	1.2	0.9	1.0	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)										
White, male										
2002.....	7.8	8.5	1.1	0.4	0.7	1.3	1.4	1.4	1.7	2.4
2001.....	7.5	6.6	1.1	0.5	0.7	1.5	1.6	1.2	1.0	2.4
2000.....	7.6	7.4	1.2	0.4	0.5	1.4	1.5	1.3	1.5	2.1
1999.....	7.4	8.4	1.1	0.5	0.5	1.6	1.8	1.2	1.2	2.1
White, female										
2002.....	6.9	5.2	0.8	0.3	0.4	0.3	0.5	0.6	0.8	1.3
2001.....	6.6	4.9	1.0	0.5	0.4	0.3	0.6	0.6	0.9	1.2
2000.....	6.6	6.3	1.0	0.5	0.4	0.4	0.6	0.5	0.9	1.3
1999.....	6.4	6.5	1.1	0.5	0.5	0.3	0.6	0.6	0.8	1.2
Black, both sexes										
2002.....	6.0	7.9	1.7	*	0.7	0.9	1.3	1.4	1.8	2.7
2001.....	6.1	8.0	2.0	*	0.7	1.1	1.4	1.7	2.2	2.4
2000.....	5.9	7.5	1.6	*	0.6	0.9	1.0	1.7	1.6	2.7
1999.....	5.7	9.1	1.4	0.8	0.8	0.8	1.4	0.9	1.7	2.6
Black, male										
2002.....	6.3	7.8	2.0	*	*	*	1.7	1.7	2.1	3.5
2001.....	6.3	10.2	1.9	*	*	1.6	1.5	2.5	2.2	2.9
2000.....	6.2	10.9	1.6	*	*	*	*	2.3	1.7	3.0
1999.....	5.7	10.4	*	*	1.2	*	2.3	*	2.1	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
IX. Diseases of the circulatory system (I00-I99)										
All races 3/, both sexes										
2002.....	320.2	16.5	1.5	0.6	1.1	2.5	4.0	7.3	13.1	25.9
2001.....	325.5	15.4	1.9	0.7	1.1	2.2	4.4	7.4	13.4	25.5
2000.....	334.6	17.4	1.5	0.7	1.1	2.5	4.4	7.0	12.5	24.7
1999.....	342.0	17.6	1.5	0.7	1.0	2.8	4.2	6.6	12.9	26.3
All races 3/, male										
2002.....	304.7	17.3	1.5	0.6	1.3	3.2	5.1	9.1	16.6	33.5
2001.....	307.6	15.8	2.0	0.6	1.1	2.7	5.4	8.9	16.8	33.5
2000.....	317.2	18.3	1.7	0.8	1.2	2.9	5.1	8.3	15.7	31.9
1999.....	324.2	18.5	1.6	0.8	1.1	3.4	4.9	7.9	16.4	34.6
All races 3/, female										
2002.....	335.2	15.7	1.4	0.7	0.9	1.7	2.9	5.4	9.5	18.2
2001.....	342.8	15.0	1.8	0.8	1.1	1.7	3.4	5.8	9.9	17.5
2000.....	351.3	16.5	1.4	0.6	1.0	2.1	3.7	5.6	9.4	17.6
1999.....	359.1	16.6	1.4	0.6	1.0	2.1	3.6	5.3	9.4	18.0
White, both sexes										
2002.....	341.0	14.1	1.3	0.6	0.9	2.2	3.4	5.9	10.6	22.1
2001.....	346.4	12.7	1.6	0.6	1.0	2.0	3.5	6.1	11.2	21.8
2000.....	356.4	15.3	1.4	0.6	0.9	2.1	3.6	5.7	10.4	21.1
1999.....	364.2	15.2	1.3	0.7	0.9	2.4	3.5	5.5	10.9	22.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
IX. Diseases of the circulatory system (I00-I99)										
White, male										
2002.....	322.2	15.2	1.5	0.5	1.1	2.8	4.4	7.6	13.7	29.0
2001.....	324.9	13.7	1.6	0.6	1.0	2.3	4.4	7.5	14.3	29.2
2000.....	335.6	16.1	1.6	0.7	1.1	2.5	4.3	7.0	13.5	28.1
1999.....	342.5	15.9	1.5	0.8	0.9	2.9	3.9	6.6	14.2	30.2
White, female										
2002.....	359.5	13.0	1.2	0.7	0.8	1.5	2.3	4.1	7.4	14.9
2001.....	367.4	11.7	1.5	0.7	0.9	1.6	2.6	4.6	7.9	14.2
2000.....	376.6	14.4	1.1	0.5	0.8	1.7	2.9	4.4	7.2	13.9
1999.....	385.2	14.4	1.2	0.6	0.9	1.9	3.0	4.3	7.4	14.4
Black, both sexes										
2002.....	278.2	28.6	2.0	0.8	2.0	4.1	8.1	17.2	31.0	54.4
2001.....	282.4	29.3	3.6	1.0	1.6	4.0	9.7	17.0	29.5	52.7
2000.....	287.0	28.6	2.6	1.0	1.9	4.8	9.4	15.9	28.7	52.4
1999.....	291.1	29.9	2.5	1.0	1.8	5.0	9.3	14.8	27.5	56.2
Black, male										
2002.....	270.5	28.8	2.0	*	2.3	5.3	9.8	21.1	38.5	68.3
2001.....	274.3	29.0	3.8	*	1.5	5.1	11.6	20.8	36.0	66.9
2000.....	277.6	29.7	2.2	*	1.8	5.3	10.5	18.7	33.7	63.2
1999.....	284.2	31.6	2.6	*	2.1	6.4	11.1	18.0	33.8	70.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
IX. Diseases of the circulatory system (I00-I99)										
Black, female										
2002.....	285.2	28.5	2.1	*	1.6	2.8	6.5	13.6	24.2	42.0
2001.....	289.9	29.6	3.4	1.3	1.6	2.9	8.0	13.4	23.5	40.1
2000.....	295.6	27.4	2.9	*	2.0	4.4	8.4	13.5	24.1	42.8
1999.....	297.4	28.2	2.4	*	1.5	3.5	7.6	11.8	21.9	43.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive diseases (I10-I13)										
All races 3/, both sexes										
2002.....	17.2	*	*	*	*	*	0.2	0.7	1.3	2.5
2001.....	16.4	*	*	*	*	*	0.2	0.6	1.3	2.2
2000.....	15.9	*	*	*	*	*	0.2	0.5	1.0	2.0
1999.....	15.4	*	*	*	*	*	0.2	0.4	1.1	2.2
All races 3/, male										
2002.....	14.5	*	*	*	*	*	0.3	1.0	1.8	3.5
2001.....	13.5	*	*	*	*	*	0.3	0.7	1.7	2.8
2000.....	13.0	*	*	*	*	*	0.3	0.6	1.3	2.5
1999.....	12.6	*	*	*	*	*	0.3	0.6	1.5	2.8
All races 3/, female										
2002.....	19.9	*	*	*	*	*	*	0.4	0.9	1.6
2001.....	19.2	*	*	*	*	*	*	0.5	0.9	1.6
2000.....	18.6	*	*	*	*	*	*	0.4	0.7	1.5
1999.....	18.1	*	*	*	*	*	*	0.2	0.7	1.5
White, both sexes										
2002.....	15.8	*	*	*	*	*	0.2	0.4	0.8	1.5
2001.....	15.0	*	*	*	*	*	*	0.3	0.8	1.4
2000.....	14.4	*	*	*	*	*	0.1	0.3	0.5	1.2
1999.....	13.9	*	*	*	*	*	*	0.2	0.7	1.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive diseases (I10-I13)										
White, male										
2002.....	12.7	*	*	*	*	*	0.2	0.6	1.0	2.2
2001.....	11.8	*	*	*	*	*	*	0.4	1.1	2.0
2000.....	11.3	*	*	*	*	*	*	0.4	0.8	1.7
1999.....	10.8	*	*	*	*	*	*	0.4	1.1	1.7
White, female										
2002.....	18.8	*	*	*	*	*	*	*	0.5	0.9
2001.....	18.2	*	*	*	*	*	*	*	0.4	0.8
2000.....	17.4	*	*	*	*	*	*	*	0.3	0.7
1999.....	16.9	*	*	*	*	*	*	*	0.3	0.6
Black, both sexes										
2002.....	30.7	*	*	*	*	*	0.7	2.4	5.3	9.3
2001.....	29.1	*	*	*	*	*	0.7	2.0	4.9	7.9
2000.....	28.9	*	*	*	*	*	0.8	1.8	4.4	7.5
1999.....	28.6	*	*	*	*	*	1.0	1.5	3.9	8.9
Black, male										
2002.....	29.3	*	*	*	*	*	*	3.6	7.0	13.1
2001.....	27.7	*	*	*	*	*	*	2.4	6.2	9.1
2000.....	26.8	*	*	*	*	*	*	2.0	5.2	8.5
1999.....	26.8	*	*	*	*	*	*	1.9	4.4	11.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive diseases (I10-I13)										
Black, female										
2002.....	31.9	*	*	*	*	*	*	*	3.8	6.0
2001.....	30.5	*	*	*	*	*	*	1.7	3.8	6.8
2000.....	30.8	*	*	*	*	*	*	1.6	3.6	6.5
1999.....	30.3	*	*	*	*	*	*	*	3.6	6.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
All races 3/, both sexes										
2002.....	9.2	*	*	*	*	*	0.1	0.4	1.0	1.8
2001.....	8.7	*	*	*	*	*	*	0.3	0.9	1.6
2000.....	8.4	*	*	*	*	*	0.1	0.4	0.7	1.4
1999.....	8.1	*	*	*	*	*	0.1	0.3	0.8	1.5
All races 3/, male										
2002.....	8.2	*	*	*	*	*	0.2	0.7	1.4	2.6
2001.....	7.5	*	*	*	*	*	*	0.4	1.2	2.1
2000.....	7.2	*	*	*	*	*	0.2	0.4	0.9	1.9
1999.....	6.8	*	*	*	*	*	*	0.4	1.0	2.0
All races 3/, female										
2002.....	10.2	*	*	*	*	*	*	*	0.6	1.0
2001.....	9.8	*	*	*	*	*	*	0.3	0.5	1.1
2000.....	9.7	*	*	*	*	*	*	0.3	0.5	1.0
1999.....	9.4	*	*	*	*	*	*	*	0.5	1.0
White, both sexes										
2002.....	8.3	*	*	*	*	*	*	0.3	0.6	1.3
2001.....	7.8	*	*	*	*	*	*	0.2	0.5	1.1
2000.....	7.6	*	*	*	*	*	*	0.2	0.4	0.9
1999.....	7.3	*	*	*	*	*	*	0.2	0.5	0.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
White, male										
2002.....	7.1	*	*	*	*	*	*	0.4	0.9	1.8
2001.....	6.4	*	*	*	*	*	*	0.3	0.9	1.6
2000.....	6.1	*	*	*	*	*	*	0.3	0.6	1.4
1999.....	5.8	*	*	*	*	*	*	0.3	0.8	1.3
White, female										
2002.....	9.5	*	*	*	*	*	*	*	0.3	0.7
2001.....	9.2	*	*	*	*	*	*	*	*	0.5
2000.....	9.0	*	*	*	*	*	*	*	*	0.5
1999.....	8.7	*	*	*	*	*	*	*	*	0.5
Black, both sexes										
2002.....	17.2	*	*	*	*	*	*	1.5	3.9	5.8
2001.....	16.0	*	*	*	*	*	*	1.1	3.1	5.1
2000.....	16.1	*	*	*	*	*	*	1.4	2.8	4.9
1999.....	15.5	*	*	*	*	*	*	1.0	2.6	5.7
Black, male										
2002.....	17.5	*	*	*	*	*	*	2.4	5.4	8.5
2001.....	15.9	*	*	*	*	*	*	*	4.0	5.9
2000.....	15.6	*	*	*	*	*	*	1.6	3.5	5.8
1999.....	15.2	*	*	*	*	*	*	*	2.8	7.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
Black, female										
2002.....	16.9	*	*	*	*	*	*	*	2.6	3.4
2001.....	16.2	*	*	*	*	*	*	*	2.3	4.4
2000.....	16.5	*	*	*	*	*	*	*	2.1	4.1
1999.....	15.9	*	*	*	*	*	*	*	2.5	4.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
All races 3/, both sexes										
2002.....	171.4	0.6	*	*	*	0.1	0.5	1.3	3.6	10.1
2001.....	176.3	0.6	*	*	*	0.2	0.7	1.5	4.0	9.9
2000.....	183.1	0.8	*	*	*	0.2	0.5	1.2	3.4	9.8
1999.....	189.8	0.7	*	*	*	0.3	0.5	1.3	3.7	10.4
All races 3/, male										
2002.....	178.4	*	*	*	*	*	0.7	2.0	5.2	15.2
2001.....	181.7	*	*	*	*	0.2	0.9	2.2	5.9	15.0
2000.....	188.7	*	*	*	*	0.2	0.6	1.8	5.1	14.7
1999.....	195.4	*	*	*	*	0.3	0.6	1.7	5.5	15.8
All races 3/, female										
2002.....	164.7	*	*	*	*	*	0.3	0.6	1.9	4.9
2001.....	171.2	*	*	*	*	*	0.4	0.7	2.1	4.9
2000.....	177.6	*	*	*	*	*	0.3	0.6	1.7	5.0
1999.....	184.5	*	*	*	*	*	0.3	0.8	1.9	5.1
White, both sexes										
2002.....	185.4	*	*	*	*	0.1	0.4	1.2	3.3	9.6
2001.....	190.5	*	*	*	*	0.1	0.5	1.3	3.8	9.5
2000.....	197.9	0.8	*	*	*	0.2	0.4	1.0	3.1	9.4
1999.....	205.1	*	*	*	*	0.2	0.3	1.2	3.5	9.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
White, male										
2002.....	192.5	*	*	*	*	*	0.6	1.9	5.0	14.6
2001.....	195.8	*	*	*	*	*	0.8	2.0	5.5	14.5
2000.....	203.6	*	*	*	*	*	0.6	1.6	4.8	14.3
1999.....	210.5	*	*	*	*	0.3	0.5	1.6	5.1	15.1
White, female										
2002.....	178.4	*	*	*	*	*	*	0.5	1.6	4.5
2001.....	185.3	*	*	*	*	*	0.3	0.6	1.9	4.5
2000.....	192.4	*	*	*	*	*	0.3	0.5	1.4	4.5
1999.....	200.0	*	*	*	*	*	*	0.7	1.7	4.6
Black, both sexes										
2002.....	132.9	*	*	*	*	*	0.9	2.1	5.6	15.1
2001.....	136.9	*	*	*	*	*	1.5	2.8	6.4	14.9
2000.....	140.1	*	*	*	*	*	0.7	2.4	6.0	14.6
1999.....	143.7	*	*	*	*	*	1.2	1.9	5.8	16.1
Black, male										
2002.....	135.3	*	*	*	*	*	*	2.7	7.5	22.0
2001.....	138.9	*	*	*	*	*	1.9	3.9	9.3	21.6
2000.....	141.5	*	*	*	*	*	*	3.7	8.0	20.3
1999.....	146.0	*	*	*	*	*	*	2.9	8.4	23.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
Black, female										
2002.....	130.8	*	*	*	*	*	*	1.5	4.0	9.0
2001.....	135.2	*	*	*	*	*	*	1.7	3.8	9.0
2000.....	138.8	*	*	*	*	*	*	*	4.3	9.7
1999.....	141.7	*	*	*	*	*	*	*	3.4	9.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
All races 3/, both sexes										
2002.....	62.3	*	*	*	*	*	0.2	0.5	1.5	4.0
2001.....	64.9	*	*	*	*	*	0.2	0.6	1.8	4.2
2000.....	68.5	0.5	*	*	*	*	0.2	0.6	1.3	4.0
1999.....	71.5	*	*	*	*	0.1	0.2	0.5	1.6	4.1
All races 3/, male										
2002.....	66.2	*	*	*	*	*	0.3	0.7	2.1	6.0
2001.....	68.6	*	*	*	*	*	0.3	0.7	2.5	6.2
2000.....	72.7	*	*	*	*	*	0.2	0.8	1.9	5.8
1999.....	76.1	*	*	*	*	*	0.3	0.7	2.3	6.2
All races 3/, female										
2002.....	58.4	*	*	*	*	*	*	0.3	0.8	2.1
2001.....	61.3	*	*	*	*	*	*	0.4	1.0	2.1
2000.....	64.6	*	*	*	*	*	*	0.3	0.7	2.2
1999.....	67.0	*	*	*	*	*	*	0.3	0.9	2.0
White, both sexes										
2002.....	67.3	*	*	*	*	*	0.2	0.4	1.4	4.0
2001.....	70.1	*	*	*	*	*	0.2	0.5	1.7	4.1
2000.....	74.1	*	*	*	*	*	0.2	0.5	1.2	4.0
1999.....	77.4	*	*	*	*	*	0.1	0.5	1.5	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
White, male										
2002.....	71.8	*	*	*	*	*	0.3	0.6	2.1	5.9
2001.....	74.4	*	*	*	*	*	0.3	0.6	2.5	6.1
2000.....	78.8	*	*	*	*	*	*	0.6	1.8	6.0
1999.....	82.7	*	*	*	*	*	*	0.6	2.1	6.2
White, female										
2002.....	62.8	*	*	*	*	*	*	*	0.7	1.9
2001.....	66.0	*	*	*	*	*	*	0.3	0.9	2.0
2000.....	69.6	*	*	*	*	*	*	0.3	0.6	2.0
1999.....	72.3	*	*	*	*	*	*	0.3	0.9	1.9
Black, both sexes										
2002.....	48.5	*	*	*	*	*	*	0.9	2.3	5.5
2001.....	50.0	*	*	*	*	*	*	1.1	2.5	5.8
2000.....	52.2	*	*	*	*	*	*	1.2	2.3	4.9
1999.....	53.3	*	*	*	*	*	*	0.8	2.4	5.5
Black, male										
2002.....	48.3	*	*	*	*	*	*	*	3.0	7.6
2001.....	49.2	*	*	*	*	*	*	*	3.4	8.0
2000.....	52.0	*	*	*	*	*	*	1.6	2.7	6.5
1999.....	53.4	*	*	*	*	*	*	*	3.4	7.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
Black, female										
2002.....	48.8	*	*	*	*	*	*	*	1.6	3.6
2001.....	50.6	*	*	*	*	*	*	*	1.7	3.8
2000.....	52.5	*	*	*	*	*	*	*	1.9	3.6
1999.....	53.2	*	*	*	*	*	*	*	1.4	3.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other forms of chronic ischemic heart disease (I20,I25)

All races 3/, both sexes

2002.....	108.0	*	*	*	*	*	0.3	0.8	2.0	6.0
2001.....	110.3	*	*	*	*	*	0.4	0.9	2.2	5.6
2000.....	113.3	*	*	*	*	*	0.3	0.6	2.0	5.7
1999.....	117.2	*	*	*	*	0.1	0.3	0.7	2.1	6.2

All races 3/, male

2002.....	110.9	*	*	*	*	*	0.4	1.3	3.0	9.2
2001.....	111.9	*	*	*	*	*	0.5	1.5	3.3	8.6
2000.....	114.8	*	*	*	*	*	0.4	1.1	3.1	8.7
1999.....	118.0	*	*	*	*	0.2	0.3	1.0	3.2	9.4

All races 3/, female

2002.....	105.2	*	*	*	*	*	*	0.4	1.0	2.8
2001.....	108.8	*	*	*	*	*	0.2	0.3	1.1	2.7
2000.....	111.9	*	*	*	*	*	*	0.2	1.0	2.8
1999.....	116.4	*	*	*	*	*	*	0.4	1.0	3.1

White, both sexes

2002.....	116.9	*	*	*	*	*	0.2	0.8	1.8	5.6
2001.....	119.1	*	*	*	*	*	0.3	0.8	2.0	5.3
2000.....	122.6	*	*	*	*	*	0.2	0.6	1.9	5.3
1999.....	126.5	*	*	*	*	*	0.2	0.7	1.9	5.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other forms of chronic ischemic heart disease (I20,I25)										
White, male										
2002.....	119.4	*	*	*	*	*	0.4	1.3	2.8	8.6
2001.....	120.1	*	*	*	*	*	0.4	1.4	3.0	8.2
2000.....	123.5	*	*	*	*	*	0.3	0.9	2.9	8.2
1999.....	126.5	*	*	*	*	*	*	0.9	3.0	8.8
White, female										
2002.....	114.4	*	*	*	*	*	*	0.3	0.8	2.5
2001.....	118.1	*	*	*	*	*	*	*	0.9	2.4
2000.....	121.6	*	*	*	*	*	*	*	0.8	2.4
1999.....	126.5	*	*	*	*	*	*	0.4	0.9	2.7
Black, both sexes										
2002.....	83.2	*	*	*	*	*	*	1.1	3.2	9.6
2001.....	85.8	*	*	*	*	*	0.8	1.7	3.8	8.9
2000.....	86.6	*	*	*	*	*	*	1.3	3.6	9.5
1999.....	89.3	*	*	*	*	*	0.7	1.1	3.4	10.1
Black, male										
2002.....	85.7	*	*	*	*	*	*	*	4.1	14.4
2001.....	88.2	*	*	*	*	*	*	2.4	5.9	13.3
2000.....	88.0	*	*	*	*	*	*	2.1	5.0	13.4
1999.....	91.1	*	*	*	*	*	*	1.7	5.0	14.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other forms of chronic ischemic heart disease (I20,I25)										
Black, female										
2002.....	81.0	*	*	*	*	*	*	*	2.3	5.3
2001.....	83.5	*	*	*	*	*	*	*	1.9	4.9
2000.....	85.3	*	*	*	*	*	*	*	2.4	5.9
1999.....	87.6	*	*	*	*	*	*	*	1.9	5.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Atherosclerotic cardiovascular disease, so described (I25.0)										
All races 3/, both sexes										
2002.....	23.6	*	*	*	*	*	*	0.2	0.7	1.9
2001.....	23.9	*	*	*	*	*	*	0.2	0.7	1.8
2000.....	24.7	*	*	*	*	*	*	0.2	0.5	1.7
1999.....	25.8	*	*	*	*	*	*	0.2	0.6	2.0
All races 3/, male										
2002.....	25.4	*	*	*	*	*	*	0.3	1.1	3.0
2001.....	25.3	*	*	*	*	*	*	0.3	1.0	2.8
2000.....	26.0	*	*	*	*	*	*	0.3	0.9	2.8
1999.....	26.8	*	*	*	*	*	*	0.2	0.9	3.1
All races 3/, female										
2002.....	21.9	*	*	*	*	*	*	*	0.3	0.8
2001.....	22.5	*	*	*	*	*	*	*	0.4	0.8
2000.....	23.5	*	*	*	*	*	*	*	0.2	0.7
1999.....	24.8	*	*	*	*	*	*	*	0.3	0.9
White, both sexes										
2002.....	24.1	*	*	*	*	*	*	0.2	0.6	1.7
2001.....	24.3	*	*	*	*	*	*	0.2	0.6	1.7
2000.....	25.1	*	*	*	*	*	*	*	0.5	1.6
1999.....	26.1	*	*	*	*	*	*	0.1	0.5	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Atherosclerotic cardiovascular disease, so described (I25.0)										
White, male										
2002.....	25.5	*	*	*	*	*	*	0.3	1.0	2.7
2001.....	25.3	*	*	*	*	*	*	0.3	0.9	2.7
2000.....	26.0	*	*	*	*	*	*	*	0.8	2.5
1999.....	26.6	*	*	*	*	*	*	*	0.8	2.9
White, female										
2002.....	22.7	*	*	*	*	*	*	*	*	0.7
2001.....	23.3	*	*	*	*	*	*	*	0.3	0.7
2000.....	24.3	*	*	*	*	*	*	*	*	0.6
1999.....	25.6	*	*	*	*	*	*	*	*	0.7
Black, both sexes										
2002.....	27.0	*	*	*	*	*	*	*	1.3	3.5
2001.....	27.7	*	*	*	*	*	*	*	1.0	3.0
2000.....	28.5	*	*	*	*	*	*	*	0.9	2.9
1999.....	29.9	*	*	*	*	*	*	*	1.2	3.7
Black, male										
2002.....	31.1	*	*	*	*	*	*	*	1.9	5.4
2001.....	31.5	*	*	*	*	*	*	*	*	4.6
2000.....	32.3	*	*	*	*	*	*	*	*	4.7
1999.....	33.8	*	*	*	*	*	*	*	1.7	5.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
All races 3/, both sexes										
2002.....	84.4	*	*	*	*	*	0.2	0.6	1.3	4.1
2001.....	86.4	*	*	*	*	*	0.3	0.7	1.5	3.8
2000.....	88.6	*	*	*	*	*	0.2	0.5	1.5	4.0
1999.....	91.4	*	*	*	*	0.1	0.2	0.6	1.5	4.3
All races 3/, male										
2002.....	85.5	*	*	*	*	*	0.3	0.9	1.9	6.2
2001.....	86.5	*	*	*	*	*	0.4	1.1	2.3	5.8
2000.....	88.8	*	*	*	*	*	0.2	0.8	2.2	5.9
1999.....	91.2	*	*	*	*	0.2	0.2	0.8	2.2	6.3
All races 3/, female										
2002.....	83.3	*	*	*	*	*	*	0.3	0.8	2.0
2001.....	86.2	*	*	*	*	*	*	0.2	0.7	1.9
2000.....	88.4	*	*	*	*	*	*	*	0.8	2.1
1999.....	91.6	*	*	*	*	*	*	0.3	0.7	2.2
White, both sexes										
2002.....	92.8	*	*	*	*	*	0.2	0.6	1.2	3.9
2001.....	94.8	*	*	*	*	*	0.2	0.6	1.4	3.6
2000.....	97.4	*	*	*	*	*	0.1	0.4	1.4	3.7
1999.....	100.4	*	*	*	*	*	0.1	0.5	1.4	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
White, male										
2002.....	93.9	*	*	*	*	*	0.3	0.9	1.8	5.9
2001.....	94.8	*	*	*	*	*	0.4	1.1	2.1	5.5
2000.....	97.5	*	*	*	*	*	*	0.7	2.1	5.6
1999.....	99.9	*	*	*	*	*	*	0.7	2.1	6.0
White, female										
2002.....	91.7	*	*	*	*	*	*	*	0.6	1.9
2001.....	94.9	*	*	*	*	*	*	*	0.6	1.7
2000.....	97.3	*	*	*	*	*	*	*	0.6	1.8
1999.....	100.8	*	*	*	*	*	*	0.3	0.6	2.0
Black, both sexes										
2002.....	56.2	*	*	*	*	*	*	0.9	1.9	6.1
2001.....	58.0	*	*	*	*	*	*	1.2	2.8	5.8
2000.....	58.1	*	*	*	*	*	*	0.8	2.8	6.5
1999.....	59.3	*	*	*	*	*	*	0.8	2.2	6.4
Black, male										
2002.....	54.5	*	*	*	*	*	*	*	2.3	9.0
2001.....	56.7	*	*	*	*	*	*	1.8	4.6	8.7
2000.....	55.8	*	*	*	*	*	*	*	3.8	8.7
1999.....	57.3	*	*	*	*	*	*	*	3.3	9.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
All races 3/, both sexes										
2002.....	58.8	11.7	1.0	0.4	0.7	1.8	2.3	3.8	5.2	8.1
2001.....	58.6	11.1	1.4	0.5	0.7	1.5	2.5	3.6	5.1	8.0
2000.....	58.8	12.0	1.1	0.5	0.7	1.8	2.6	3.5	5.2	7.5
1999.....	59.4	12.9	1.2	0.5	0.7	2.0	2.6	3.5	5.3	8.2
All races 3/, male										
2002.....	52.4	12.1	1.0	0.4	0.8	2.4	3.0	4.6	6.6	9.6
2001.....	51.8	10.8	1.5	0.4	0.8	1.9	3.1	4.3	6.0	9.9
2000.....	52.2	12.2	1.2	0.6	0.9	2.1	3.1	4.2	6.2	9.1
1999.....	53.1	13.1	1.3	0.5	0.8	2.4	3.0	4.0	6.4	10.3
All races 3/, female										
2002.....	65.0	11.4	1.0	0.5	0.5	1.1	1.7	3.0	3.8	6.6
2001.....	65.2	11.5	1.3	0.5	0.7	1.1	1.9	2.9	4.1	6.0
2000.....	65.1	11.8	0.9	0.4	0.6	1.4	2.1	2.9	4.1	5.9
1999.....	65.5	12.7	1.0	0.4	0.7	1.5	2.2	2.9	4.2	6.1
White, both sexes										
2002.....	62.6	10.0	0.9	0.4	0.6	1.5	1.9	3.0	4.1	6.6
2001.....	62.3	9.3	1.1	0.4	0.7	1.3	2.0	2.9	4.1	6.5
2000.....	62.5	10.6	1.0	0.4	0.6	1.4	2.1	2.8	4.3	6.1
1999.....	62.9	11.0	1.0	0.4	0.6	1.8	2.1	2.7	4.4	6.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
White, male										
2002.....	54.9	10.5	1.0	0.4	0.7	2.1	2.5	3.6	5.2	7.9
2001.....	54.1	9.6	1.2	0.4	0.7	1.5	2.4	3.5	4.8	8.1
2000.....	54.6	10.8	1.2	0.5	0.8	1.7	2.5	3.4	5.2	7.5
1999.....	55.1	11.4	1.1	0.5	0.7	2.1	2.4	3.3	5.4	8.5
White, female										
2002.....	70.2	9.6	0.8	0.4	0.5	1.0	1.3	2.3	2.9	5.3
2001.....	70.4	8.9	1.1	0.5	0.6	1.0	1.5	2.3	3.4	4.8
2000.....	70.2	10.4	0.7	0.3	0.4	1.2	1.6	2.2	3.3	4.6
1999.....	70.5	10.7	0.8	0.4	0.6	1.4	1.9	2.2	3.4	4.8
Black, both sexes										
2002.....	52.5	20.2	1.5	*	1.2	3.1	5.1	9.5	13.0	18.9
2001.....	52.8	20.1	2.9	0.8	1.1	2.9	5.5	8.7	12.3	18.4
2000.....	52.7	19.0	1.9	0.8	1.4	3.6	6.0	8.8	11.9	17.8
1999.....	54.4	22.3	2.0	0.7	1.2	3.5	5.6	8.6	12.4	19.5
Black, male										
2002.....	50.8	20.9	*	*	1.3	4.0	6.2	11.9	16.4	22.8
2001.....	51.7	17.7	3.2	*	1.2	3.8	7.0	10.7	14.9	23.1
2000.....	51.5	18.9	1.9	*	1.4	4.3	6.7	10.1	14.3	21.6
1999.....	54.3	21.8	2.1	*	1.5	4.8	6.8	10.2	15.2	24.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
Black, female										
2002.....	54.0	19.4	1.8	*	*	2.2	4.0	7.4	9.8	15.5
2001.....	53.8	22.7	2.5	*	*	2.0	4.1	6.9	10.0	14.1
2000.....	53.9	19.2	2.0	*	1.3	2.9	5.3	7.6	9.8	14.5
1999.....	54.5	22.8	1.8	*	*	2.2	4.5	7.1	9.9	15.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)										
All races 3/, both sexes										
2002.....	4.8	4.6	0.2	*	0.1	0.2	0.3	0.7	0.9	1.4
2001.....	4.7	4.2	0.1	*	*	0.2	0.4	0.8	1.0	1.4
2000.....	4.6	4.9	0.2	*	*	0.2	0.4	0.6	1.1	1.4
1999.....	4.7	5.7	*	0.1	*	0.3	0.5	0.7	1.0	1.5
All races 3/, male										
2002.....	3.8	4.7	*	*	*	*	0.2	0.7	0.7	1.3
2001.....	3.8	3.9	*	*	*	*	0.3	0.6	0.7	1.1
2000.....	3.8	5.1	*	*	*	*	0.3	0.5	0.9	1.2
1999.....	3.9	5.9	*	*	*	0.3	0.3	0.5	0.8	1.3
All races 3/, female										
2002.....	5.7	4.5	*	*	*	0.2	0.5	0.8	1.1	1.6
2001.....	5.5	4.6	*	*	*	*	0.6	0.9	1.3	1.7
2000.....	5.5	4.6	*	*	*	0.3	0.6	0.8	1.2	1.6
1999.....	5.5	5.6	*	*	*	0.3	0.7	0.9	1.2	1.7
White, both sexes										
2002.....	4.8	3.9	*	*	*	0.2	0.3	0.5	0.7	1.3
2001.....	4.7	3.6	*	*	*	0.1	0.4	0.6	0.8	1.1
2000.....	4.7	4.1	0.2	*	*	0.2	0.4	0.5	0.9	1.2
1999.....	4.7	5.2	*	*	*	0.3	0.4	0.5	0.8	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)										
White, male										
2002.....	3.9	4.1	*	*	*	*	*	0.5	0.6	1.1
2001.....	3.8	3.4	*	*	*	*	0.3	0.5	0.6	0.9
2000.....	3.8	4.3	*	*	*	*	0.3	0.5	0.7	1.1
1999.....	3.9	5.4	*	*	*	0.2	*	0.4	0.6	1.2
White, female										
2002.....	5.8	3.7	*	*	*	*	0.4	0.5	0.8	1.4
2001.....	5.6	3.9	*	*	*	*	0.5	0.7	1.1	1.4
2000.....	5.5	3.9	*	*	*	0.3	0.4	0.6	1.0	1.3
1999.....	5.5	4.9	*	*	*	0.4	0.6	0.6	1.0	1.3
Black, both sexes										
2002.....	5.9	8.2	*	*	*	*	0.8	2.2	2.5	3.1
2001.....	5.8	6.8	*	*	*	*	0.8	1.9	2.3	3.2
2000.....	5.7	8.0	*	*	*	*	1.0	1.6	2.5	3.1
1999.....	6.2	9.4	*	*	*	*	1.0	1.8	2.6	3.3
Black, male										
2002.....	4.5	9.0	*	*	*	*	*	2.2	1.8	2.9
2001.....	4.8	6.9	*	*	*	*	*	*	1.7	2.8
2000.....	4.6	7.7	*	*	*	*	*	*	2.5	2.5
1999.....	5.1	8.8	*	*	*	*	*	*	2.5	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)										
Black, female										
2002.....	7.2	7.3	*	*	*	*	*	2.2	3.1	3.3
2001.....	6.8	6.6	*	*	*	*	*	2.4	2.9	3.5
2000.....	6.8	8.3	*	*	*	*	1.5	2.2	2.6	3.6
1999.....	7.2	10.1	*	*	*	*	*	2.3	2.7	4.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pulmonary embolism (I26)										
All races 3/, both sexes										
2002.....	3.0	*	*	*	*	0.1	0.3	0.5	0.6	1.0
2001.....	3.0	*	*	*	*	*	0.3	0.5	0.7	0.9
2000.....	3.0	*	*	*	*	0.1	0.3	0.4	0.7	1.0
1999.....	3.2	*	*	*	*	0.2	0.3	0.4	0.7	0.9
All races 3/, male										
2002.....	2.5	*	*	*	*	*	*	0.5	0.5	0.9
2001.....	2.6	*	*	*	*	*	0.2	0.4	0.5	0.8
2000.....	2.6	*	*	*	*	*	*	0.3	0.6	0.8
1999.....	2.7	*	*	*	*	*	0.3	0.3	0.6	0.9
All races 3/, female										
2002.....	3.5	*	*	*	*	*	0.3	0.5	0.7	1.1
2001.....	3.5	*	*	*	*	*	0.4	0.6	0.9	1.1
2000.....	3.4	*	*	*	*	*	0.4	0.5	0.7	1.1
1999.....	3.7	*	*	*	*	*	0.5	0.6	0.8	1.0
White, both sexes										
2002.....	3.0	*	*	*	*	*	0.2	0.3	0.5	0.9
2001.....	3.1	*	*	*	*	*	0.3	0.4	0.6	0.7
2000.....	3.0	*	*	*	*	*	0.2	0.4	0.6	0.8
1999.....	3.2	*	*	*	*	0.1	0.3	0.3	0.5	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pulmonary embolism (I26)										
White, male										
2002.....	2.5	*	*	*	*	*	*	0.3	0.4	0.7
2001.....	2.6	*	*	*	*	*	*	0.3	0.4	0.6
2000.....	2.6	*	*	*	*	*	*	0.4	0.5	0.7
1999.....	2.7	*	*	*	*	*	*	*	0.5	0.8
White, female										
2002.....	3.5	*	*	*	*	*	0.3	0.4	0.6	1.0
2001.....	3.6	*	*	*	*	*	0.3	0.4	0.8	0.8
2000.....	3.4	*	*	*	*	*	0.3	0.3	0.6	0.8
1999.....	3.7	*	*	*	*	*	0.4	0.4	0.6	0.8
Black, both sexes										
2002.....	3.8	*	*	*	*	*	*	1.6	1.5	2.1
2001.....	3.8	*	*	*	*	*	*	1.4	1.7	2.4
2000.....	3.9	*	*	*	*	*	0.9	0.9	1.6	2.3
1999.....	4.3	*	*	*	*	*	0.8	1.3	2.0	2.0
Black, male										
2002.....	3.0	*	*	*	*	*	*	1.6	*	2.0
2001.....	3.3	*	*	*	*	*	*	*	*	2.1
2000.....	3.3	*	*	*	*	*	*	*	1.5	2.0
1999.....	3.8	*	*	*	*	*	*	*	2.0	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pulmonary embolism (I26)										
Black, female										
2002.....	4.4	*	*	*	*	*	*	1.5	1.6	2.2
2001.....	4.3	*	*	*	*	*	*	1.7	2.0	2.7
2000.....	4.4	*	*	*	*	*	*	1.4	1.6	2.6
1999.....	4.8	*	*	*	*	*	*	1.5	2.1	2.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

All races 3/, both sexes

2002.....	1.8	4.6	0.2	*	*	*	*	0.2	0.3	0.4
2001.....	1.6	4.2	0.1	*	*	*	0.2	0.3	0.3	0.5
2000.....	1.6	4.7	0.2	*	*	0.1	0.1	0.2	0.4	0.5
1999.....	1.5	5.7	*	0.1	*	0.1	0.2	0.2	0.3	0.5

All races 3/, male

2002.....	1.3	4.7	*	*	*	*	*	*	0.2	0.4
2001.....	1.2	3.9	*	*	*	*	*	*	0.2	0.3
2000.....	1.2	4.9	*	*	*	*	*	*	0.3	0.4
1999.....	1.2	5.9	*	*	*	*	*	*	*	0.4

All races 3/, female

2002.....	2.2	4.4	*	*	*	*	*	0.3	0.4	0.5
2001.....	2.0	4.5	*	*	*	*	0.2	0.4	0.4	0.6
2000.....	2.1	4.5	*	*	*	*	*	0.3	0.5	0.5
1999.....	1.8	5.5	*	*	*	*	0.3	0.3	0.5	0.7

White, both sexes

2002.....	1.8	3.8	*	*	*	*	*	0.1	0.2	0.4
2001.....	1.7	3.6	*	*	*	*	*	0.2	0.3	0.4
2000.....	1.7	4.0	0.2	*	*	*	*	0.2	0.3	0.4
1999.....	1.5	5.1	*	*	*	0.2	0.2	0.2	0.3	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other non-pulmonary forms of heart disease (I30-I51)										
All races 3/, both sexes										
2002.....	54.0	7.1	0.8	0.4	0.6	1.6	2.0	3.1	4.3	6.6
2001.....	53.9	6.9	1.3	0.4	0.6	1.3	2.0	2.9	4.1	6.6
2000.....	54.1	7.1	0.9	0.4	0.7	1.5	2.2	2.9	4.1	6.1
1999.....	54.7	7.2	1.0	0.4	0.7	1.7	2.1	2.8	4.3	6.7
All races 3/, male										
2002.....	48.6	7.3	0.9	0.3	0.7	2.2	2.7	4.0	5.9	8.3
2001.....	48.0	6.9	1.3	0.3	0.7	1.7	2.7	3.7	5.3	8.8
2000.....	48.5	7.1	1.0	0.4	0.8	2.0	2.8	3.6	5.3	7.9
1999.....	49.1	7.3	1.1	0.4	0.8	2.2	2.6	3.6	5.6	9.0
All races 3/, female										
2002.....	59.3	6.9	0.8	0.4	0.4	0.9	1.2	2.2	2.7	4.9
2001.....	59.7	7.0	1.2	0.5	0.6	1.0	1.3	2.0	2.8	4.4
2000.....	59.6	7.2	0.7	0.3	0.5	1.0	1.5	2.1	2.9	4.3
1999.....	60.0	7.1	0.9	0.4	0.6	1.2	1.5	2.0	3.0	4.5
White, both sexes										
2002.....	57.8	6.2	0.8	0.3	0.5	1.4	1.6	2.5	3.4	5.4
2001.....	57.6	5.7	1.0	0.4	0.6	1.1	1.6	2.3	3.3	5.3
2000.....	57.8	6.5	0.8	0.3	0.5	1.2	1.7	2.3	3.4	4.9
1999.....	58.1	5.9	0.9	0.3	0.6	1.5	1.7	2.2	3.6	5.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other non-pulmonary forms of heart disease (I30-I51)										
White, male										
2002.....	51.1	6.4	0.9	0.3	0.6	1.9	2.3	3.2	4.6	6.9
2001.....	50.3	6.3	1.0	0.3	0.6	1.4	2.1	3.0	4.3	7.2
2000.....	50.8	6.4	1.0	0.4	0.7	1.6	2.2	2.9	4.5	6.4
1999.....	51.1	6.0	1.0	0.4	0.6	1.8	2.2	2.9	4.8	7.4
White, female										
2002.....	64.4	5.9	0.7	0.4	0.4	0.8	0.9	1.8	2.1	3.9
2001.....	64.8	5.0	1.0	0.4	0.5	0.8	1.0	1.6	2.2	3.4
2000.....	64.6	6.6	0.6	0.3	0.4	0.9	1.1	1.6	2.3	3.3
1999.....	64.9	5.8	0.8	0.3	0.5	1.1	1.2	1.6	2.4	3.5
Black, both sexes										
2002.....	46.6	12.0	1.1	*	1.1	2.7	4.3	7.3	10.5	15.8
2001.....	47.0	13.4	2.7	0.7	1.0	2.5	4.7	6.8	10.0	15.2
2000.....	47.0	11.1	1.6	0.6	1.3	3.2	4.9	7.2	9.4	14.7
1999.....	48.2	12.9	1.8	0.7	1.2	3.2	4.6	6.8	9.8	16.2
Black, male										
2002.....	46.3	11.9	*	*	1.3	3.7	5.6	9.7	14.6	19.9
2001.....	46.9	10.8	3.0	*	*	3.3	6.4	9.3	13.2	20.4
2000.....	46.9	11.2	*	*	1.4	4.1	6.2	9.2	11.8	19.1
1999.....	49.2	13.0	2.0	*	1.4	4.5	5.9	9.0	12.7	21.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other non-pulmonary forms of heart disease (I30-I51)										
Black, female										
2002.....	46.8	12.1	*	*	*	1.7	3.0	5.1	6.8	12.2
2001.....	47.1	16.1	2.3	*	*	1.8	3.1	4.5	7.1	10.6
2000.....	47.1	10.9	1.6	*	1.2	2.1	3.8	5.4	7.2	10.8
1999.....	47.3	12.8	1.6	*	*	1.8	3.4	4.8	7.2	11.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races 3/, both sexes

2002.....	0.3	0.6	0.1	*	*	*	*	*	0.1	0.2
2001.....	0.3	0.8	0.2	*	*	*	0.1	0.1	0.1	0.2
2000.....	0.3	0.5	0.2	*	*	*	0.1	0.1	0.1	0.2
1999.....	0.3	0.7	0.2	*	*	*	0.1	*	0.2	0.2

All races 3/, male

2002.....	0.3	*	*	*	*	*	*	*	0.2	*
2001.....	0.3	*	*	*	*	*	*	*	*	0.2
2000.....	0.3	*	*	*	*	*	*	*	*	0.3
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.3

All races 3/, female

2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	0.2

White, both sexes

2002.....	0.3	*	*	*	*	*	*	*	*	0.1
2001.....	0.3	*	*	*	*	*	*	*	0.1	0.2
2000.....	0.3	*	*	*	*	*	*	0.1	*	0.1
1999.....	0.3	*	*	*	*	*	*	*	0.1	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of endocardium (I34-I38)										
All races 3/, both sexes										
2002.....	6.9	*	*	*	*	*	0.1	0.3	0.3	0.6
2001.....	6.9	*	*	*	*	*	0.1	0.3	0.3	0.6
2000.....	7.0	*	*	*	*	*	0.2	0.2	0.3	0.6
1999.....	7.0	*	*	*	*	0.1	0.2	0.2	0.5	0.6
All races 3/, male										
2002.....	5.4	*	*	*	*	*	*	0.2	0.4	0.7
2001.....	5.4	*	*	*	*	*	*	0.3	0.3	0.7
2000.....	5.5	*	*	*	*	*	*	0.3	0.3	0.7
1999.....	5.5	*	*	*	*	*	0.2	*	0.5	0.8
All races 3/, female										
2002.....	8.4	*	*	*	*	*	*	0.4	0.2	0.5
2001.....	8.2	*	*	*	*	*	*	0.2	0.3	0.5
2000.....	8.5	*	*	*	*	*	*	0.2	0.4	0.6
1999.....	8.5	*	*	*	*	*	*	0.3	0.5	0.5
White, both sexes										
2002.....	7.9	*	*	*	*	*	*	0.3	0.3	0.5
2001.....	7.8	*	*	*	*	*	*	0.2	0.3	0.5
2000.....	7.9	*	*	*	*	*	*	0.2	0.3	0.5
1999.....	7.9	*	*	*	*	0.1	0.2	0.2	0.5	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nonrheumatic mitral valve disorders (I34)										
All races 3/, both sexes										
2002.....	1.0	*	*	*	*	*	*	*	0.1	0.2
2001.....	1.0	*	*	*	*	*	*	0.1	0.1	0.2
2000.....	1.0	*	*	*	*	*	*	*	0.1	0.2
1999.....	1.0	*	*	*	*	*	*	*	0.2	0.2
All races 3/, male										
2002.....	0.7	*	*	*	*	*	*	*	*	*
2001.....	0.7	*	*	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*	*	*
1999.....	0.7	*	*	*	*	*	*	*	*	0.2
All races 3/, female										
2002.....	1.2	*	*	*	*	*	*	*	*	0.2
2001.....	1.3	*	*	*	*	*	*	*	*	0.2
2000.....	1.3	*	*	*	*	*	*	*	0.2	0.2
1999.....	1.3	*	*	*	*	*	*	*	0.2	*
White, both sexes										
2002.....	1.1	*	*	*	*	*	*	*	*	0.2
2001.....	1.1	*	*	*	*	*	*	0.1	0.2	0.2
2000.....	1.1	*	*	*	*	*	*	*	0.1	0.2
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases of endocardium (I36-I38)										
All races 3/, both sexes										
2002.....	1.7	*	*	*	*	*	*	0.1	0.1	0.2
2001.....	1.6	*	*	*	*	*	*	*	*	0.2
2000.....	1.6	*	*	*	*	*	*	*	0.1	0.3
1999.....	1.6	*	*	*	*	*	*	*	0.2	0.2
All races 3/, male										
2002.....	1.3	*	*	*	*	*	*	*	*	0.3
2001.....	1.3	*	*	*	*	*	*	*	*	0.2
2000.....	1.2	*	*	*	*	*	*	*	*	0.3
1999.....	1.3	*	*	*	*	*	*	*	0.2	0.2
All races 3/, female										
2002.....	2.0	*	*	*	*	*	*	*	*	0.2
2001.....	1.9	*	*	*	*	*	*	*	*	0.2
2000.....	1.9	*	*	*	*	*	*	*	*	0.2
1999.....	2.0	*	*	*	*	*	*	*	0.2	0.2
White, both sexes										
2002.....	1.8	*	*	*	*	*	*	*	0.1	0.2
2001.....	1.7	*	*	*	*	*	*	*	*	0.2
2000.....	1.7	*	*	*	*	*	*	*	*	0.2
1999.....	1.7	*	*	*	*	*	*	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cardiomyopathy (I42)										
All races 3/, both sexes										
2002.....	9.6	2.9	0.2	0.1	0.2	0.6	0.8	1.4	1.8	2.8
2001.....	9.4	3.4	0.4	0.1	0.2	0.6	0.8	1.1	1.7	2.6
2000.....	9.7	3.1	0.3	0.1	0.3	0.5	0.9	1.2	1.8	2.6
1999.....	9.8	3.5	0.2	0.1	0.2	0.6	0.7	1.2	1.8	3.0
All races 3/, male										
2002.....	11.3	2.7	*	*	0.3	0.8	1.1	2.0	2.6	3.7
2001.....	11.0	2.9	0.5	*	0.2	0.7	1.1	1.5	2.4	3.8
2000.....	11.4	2.8	0.3	*	0.4	0.7	1.2	1.6	2.6	3.6
1999.....	11.5	3.1	*	*	0.3	0.9	0.9	1.6	2.4	4.3
All races 3/, female										
2002.....	8.0	3.1	*	*	*	0.3	0.4	0.8	1.0	2.0
2001.....	8.0	3.9	0.3	*	*	0.4	0.4	0.7	0.9	1.5
2000.....	8.0	3.4	*	*	*	0.3	0.6	0.8	1.1	1.6
1999.....	8.1	3.8	*	*	*	0.3	0.5	0.8	1.2	1.7
White, both sexes										
2002.....	9.6	2.8	0.2	*	0.2	0.4	0.6	1.0	1.3	2.1
2001.....	9.3	2.8	0.3	*	0.2	0.4	0.6	0.8	1.3	2.0
2000.....	9.5	3.0	0.2	*	0.2	0.3	0.6	0.8	1.4	1.9
1999.....	9.6	3.1	0.2	*	0.2	0.4	0.5	0.9	1.4	2.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cardiomyopathy (I42)										
White, male										
2002.....	11.2	2.4	*	*	0.3	0.6	0.9	1.4	1.9	2.8
2001.....	10.8	2.7	0.4	*	*	0.5	0.8	1.2	1.8	2.9
2000.....	11.1	2.7	*	*	0.3	0.5	0.9	1.1	1.9	2.7
1999.....	11.1	2.6	*	*	*	0.6	0.7	1.2	2.0	3.2
White, female										
2002.....	8.0	3.2	*	*	*	0.3	*	0.5	0.7	1.4
2001.....	7.9	2.9	*	*	*	0.3	0.3	0.5	0.7	1.1
2000.....	8.0	3.4	*	*	*	*	0.3	0.5	0.8	1.1
1999.....	8.1	3.5	*	*	*	0.3	0.3	0.5	0.8	1.3
Black, both sexes										
2002.....	12.2	3.4	*	*	*	1.2	2.1	4.0	5.2	7.9
2001.....	12.4	6.8	0.9	*	*	1.3	2.1	3.2	4.6	7.1
2000.....	12.9	3.6	*	*	*	1.4	2.3	3.6	4.8	7.3
1999.....	13.2	5.1	*	*	0.6	1.7	1.9	3.2	4.4	8.0
Black, male										
2002.....	14.9	*	*	*	*	1.7	2.6	5.9	7.6	10.3
2001.....	14.9	*	*	*	*	1.7	2.9	4.1	6.7	9.9
2000.....	15.8	*	*	*	*	2.0	2.9	4.7	7.0	10.0
1999.....	16.4	*	*	*	*	2.7	2.5	4.5	5.8	11.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cardiomyopathy (I42)										
Black, female										
2002.....	9.8	*	*	*	*	*	1.6	2.2	3.1	5.8
2001.....	10.2	8.8	*	*	*	*	*	2.3	2.7	4.5
2000.....	10.3	*	*	*	*	*	1.7	2.7	2.9	4.9
1999.....	10.3	*	*	*	*	*	*	2.0	3.1	4.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Conduction disorders and cardiac dysrhythmias (I44-I49)										
All races 3/, both sexes										
2002.....	13.6	1.4	0.1	0.1	0.1	0.4	0.5	0.7	0.9	1.3
2001.....	13.5	0.9	0.2	*	0.2	0.3	0.5	0.6	0.8	1.3
2000.....	13.6	1.0	0.2	*	0.2	0.4	0.5	0.6	0.8	1.3
1999.....	14.3	1.0	0.3	*	0.2	0.4	0.5	0.7	0.9	1.3
All races 3/, male										
2002.....	12.2	1.6	*	*	*	0.6	0.7	0.8	1.1	1.6
2001.....	12.0	*	*	*	0.2	0.4	0.7	0.7	1.0	1.6
2000.....	12.1	1.1	*	*	0.2	0.5	0.6	0.8	0.9	1.6
1999.....	12.9	1.2	0.3	*	0.2	0.5	0.6	0.8	1.1	1.5
All races 3/, female										
2002.....	15.1	1.2	*	*	*	0.2	0.3	0.5	0.7	1.1
2001.....	15.1	*	*	*	0.2	*	0.3	0.4	0.7	1.0
2000.....	15.1	*	*	*	*	0.3	0.4	0.4	0.7	1.0
1999.....	15.7	*	*	*	*	0.4	0.4	0.5	0.7	1.0
White, both sexes										
2002.....	14.4	1.1	*	*	0.1	0.4	0.4	0.6	0.7	1.1
2001.....	14.3	0.9	0.2	*	0.2	0.3	0.4	0.5	0.7	1.0
2000.....	14.4	1.0	0.2	*	0.1	0.4	0.5	0.5	0.7	1.2
1999.....	15.2	0.8	0.3	*	0.2	0.4	0.4	0.6	0.8	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Conduction disorders and cardiac dysrhythmias (I44-I49)										
White, male										
2002.....	12.7	1.6	*	*	*	0.7	0.6	0.7	0.9	1.3
2001.....	12.4	*	*	*	*	0.4	0.6	0.6	0.9	1.3
2000.....	12.7	*	*	*	*	0.5	0.6	0.7	0.8	1.5
1999.....	13.5	*	0.3	*	*	0.5	0.5	0.7	1.0	1.3
White, female										
2002.....	16.1	*	*	*	*	*	*	0.4	0.6	0.9
2001.....	16.1	*	*	*	*	*	*	0.4	0.5	0.8
2000.....	16.1	*	*	*	*	*	0.4	0.3	0.5	0.9
1999.....	16.9	*	*	*	*	0.4	0.4	0.4	0.5	0.8
Black, both sexes										
2002.....	12.5	*	*	*	*	*	0.9	1.3	2.0	2.9
2001.....	12.6	*	*	*	*	*	1.0	1.0	1.9	3.2
2000.....	12.3	*	*	*	*	*	0.8	1.1	1.7	2.5
1999.....	13.0	*	*	*	*	*	1.1	1.4	2.0	3.1
Black, male										
2002.....	11.9	*	*	*	*	*	*	*	2.5	3.4
2001.....	12.2	*	*	*	*	*	1.5	*	2.1	3.8
2000.....	11.7	*	*	*	*	*	*	*	1.7	3.3
1999.....	12.9	*	*	*	*	*	*	1.6	2.2	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Conduction disorders and cardiac dysrhythmias (I44-I49)										
Black, female										
2002.....	13.1	*	*	*	*	*	*	*	1.5	2.4
2001.....	12.9	*	*	*	*	*	*	*	1.6	2.7
2000.....	12.8	*	*	*	*	*	*	*	1.7	1.8
1999.....	13.1	*	*	*	*	*	*	*	1.8	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Heart failure (I50)										
All races 3/, both sexes										
2002.....	19.6	0.6	*	*	*	*	*	0.2	0.3	0.5
2001.....	20.0	0.6	*	*	*	*	*	0.2	0.2	0.5
2000.....	19.8	0.8	*	*	*	0.1	*	0.1	0.2	0.3
1999.....	19.7	*	*	*	*	*	*	0.1	0.2	0.4
All races 3/, male										
2002.....	15.3	*	*	*	*	*	*	0.2	0.4	0.5
2001.....	15.5	*	*	*	*	*	*	0.3	0.3	0.7
2000.....	15.3	1.1	*	*	*	*	*	*	0.2	0.4
1999.....	15.2	*	*	*	*	*	*	*	*	0.5
All races 3/, female										
2002.....	23.7	*	*	*	*	*	*	*	0.3	0.4
2001.....	24.3	*	*	*	*	*	*	*	0.2	0.4
2000.....	24.1	*	*	*	*	*	*	*	*	0.3
1999.....	24.0	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002.....	21.6	*	*	*	*	*	*	*	0.2	0.3
2001.....	22.1	*	*	*	*	*	*	0.2	0.2	0.4
2000.....	21.8	*	*	*	*	*	*	*	0.1	0.2
1999.....	21.6	*	*	*	*	*	*	*	0.1	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)										
All races 3/, both sexes										
2002.....	0.3	0.8	0.1	*	*	0.2	0.1	0.1	0.2	0.3
2001.....	0.3	0.6	0.3	*	*	0.1	0.1	0.2	0.2	0.3
2000.....	0.2	0.6	0.1	*	*	0.1	0.1	0.2	0.2	0.2
1999.....	0.3	0.7	0.2	*	*	0.2	0.2	0.2	0.2	0.3
All races 3/, male										
2002.....	0.3	*	*	*	*	*	*	*	0.2	0.3
2001.....	0.3	*	0.3	*	*	*	*	*	*	0.4
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.3
1999.....	0.3	*	*	*	*	*	0.3	0.2	0.3	0.3
All races 3/, female										
2002.....	0.3	*	*	*	*	*	*	*	*	0.2
2001.....	0.2	*	*	*	*	*	*	0.2	0.2	0.2
2000.....	0.2	*	*	*	*	*	*	*	*	0.2
1999.....	0.2	*	*	*	*	*	*	*	*	0.2
White, both sexes										
2002.....	0.3	*	*	*	*	0.1	*	0.1	0.2	0.2
2001.....	0.2	*	0.2	*	*	*	*	0.1	0.1	0.3
2000.....	0.2	*	*	*	*	0.1	*	0.1	0.2	0.2
1999.....	0.2	*	*	*	*	0.2	0.2	0.1	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

All races 3/, both sexes

2002.....	3.3	0.6	*	*	*	0.2	0.3	0.3	0.5	0.8
2001.....	3.2	*	*	*	*	0.1	0.3	0.3	0.6	0.8
2000.....	3.1	0.8	*	*	*	0.2	0.3	0.3	0.5	0.6
1999.....	2.9	0.8	*	*	*	0.2	0.2	0.3	0.5	0.7

All races 3/, male

2002.....	3.3	*	*	*	*	0.3	0.4	0.4	0.7	1.2
2001.....	3.2	*	*	*	*	0.2	0.4	0.5	0.8	1.3
2000.....	3.1	*	*	*	*	0.3	0.5	0.5	0.8	0.9
1999.....	3.0	*	*	*	*	0.3	0.3	0.4	0.8	1.1

All races 3/, female

2002.....	3.2	*	*	*	*	*	*	*	*	0.5
2001.....	3.1	*	*	*	*	*	*	*	0.3	0.4
2000.....	3.1	*	*	*	*	*	*	*	*	0.4
1999.....	2.8	*	*	*	*	*	*	*	0.2	0.4

White, both sexes

2002.....	3.4	*	*	*	*	0.1	0.2	0.3	0.4	0.8
2001.....	3.2	*	*	*	*	0.1	0.2	0.3	0.5	0.7
2000.....	3.2	0.7	*	*	*	*	0.2	0.3	0.5	0.5
1999.....	3.0	0.7	*	*	*	0.2	0.2	0.2	0.4	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

White, male

2002.....	3.4	*	*	*	*	0.2	0.4	0.4	0.6	1.2
2001.....	3.2	*	*	*	*	*	0.3	0.5	0.7	1.1
2000.....	3.2	*	*	*	*	*	0.3	0.3	0.8	0.8
1999.....	3.0	*	*	*	*	*	0.3	0.4	0.7	1.0

White, female

2002.....	3.4	*	*	*	*	*	*	*	*	0.4
2001.....	3.3	*	*	*	*	*	*	*	0.3	0.3
2000.....	3.2	*	*	*	*	*	*	*	*	0.2
1999.....	2.9	*	*	*	*	*	*	*	*	0.3

Black, both sexes

2002.....	3.4	*	*	*	*	*	*	*	1.1	1.4
2001.....	3.4	*	*	*	*	*	*	*	1.2	1.5
2000.....	3.5	*	*	*	*	*	0.8	0.9	0.9	1.4
1999.....	3.3	*	*	*	*	*	*	*	1.2	1.5

Black, male

2002.....	3.8	*	*	*	*	*	*	*	1.5	1.8
2001.....	3.5	*	*	*	*	*	*	*	1.9	2.2
2000.....	3.9	*	*	*	*	*	*	*	*	1.7
1999.....	3.6	*	*	*	*	*	*	*	1.9	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cerebrovascular diseases (I60-I69)										
All races 3/, both sexes										
2002.....	56.4	2.9	0.3	0.2	0.3	0.3	0.6	1.0	1.8	3.6
2001.....	57.4	2.7	0.4	0.2	0.2	0.3	0.7	1.1	1.9	3.8
2000.....	59.6	3.3	0.3	0.1	0.2	0.3	0.7	1.1	1.9	3.9
1999.....	60.0	2.7	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.0
All races 3/, male										
2002.....	44.2	3.2	0.4	*	0.3	0.3	0.7	1.0	1.7	3.6
2001.....	45.2	3.1	0.3	0.2	0.2	0.3	0.7	1.1	2.0	3.9
2000.....	46.9	3.8	*	*	0.2	0.4	0.7	1.1	1.9	3.8
1999.....	47.1	3.3	0.3	*	*	0.3	0.6	1.0	1.9	3.9
All races 3/, female										
2002.....	68.2	2.5	0.3	*	0.3	*	0.5	0.9	1.9	3.7
2001.....	69.2	2.3	0.4	*	0.2	0.3	0.6	1.0	1.9	3.6
2000.....	71.8	2.7	0.4	*	0.3	0.3	0.7	1.0	1.9	4.0
1999.....	72.3	2.1	0.3	*	0.2	0.3	0.6	1.0	1.8	4.1
White, both sexes										
2002.....	59.5	2.3	0.3	0.2	0.2	0.2	0.6	0.8	1.5	2.9
2001.....	60.5	1.9	0.3	0.2	0.2	0.3	0.5	0.9	1.7	3.1
2000.....	62.8	2.5	0.3	0.1	0.2	0.3	0.6	1.0	1.6	3.0
1999.....	63.3	2.3	0.3	0.1	0.2	0.3	0.6	0.9	1.5	3.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cerebrovascular diseases (I60-I69)										
White, male										
2002.....	45.7	2.6	0.4	*	*	0.3	0.7	1.0	1.4	2.8
2001.....	46.6	2.2	*	*	*	0.3	0.7	0.9	1.8	3.1
2000.....	48.4	3.0	*	*	*	0.4	0.7	1.1	1.6	3.0
1999.....	48.7	2.8	*	*	*	0.3	0.6	0.9	1.7	3.2
White, female										
2002.....	73.0	2.0	*	*	0.3	*	0.5	0.7	1.6	2.9
2001.....	74.0	1.6	0.4	*	*	0.3	0.4	0.9	1.6	3.0
2000.....	76.9	2.0	*	*	0.3	*	0.6	0.9	1.5	3.0
1999.....	77.6	1.9	*	*	*	*	0.6	0.9	1.4	3.3
Black, both sexes										
2002.....	50.0	5.5	*	*	*	*	0.8	1.9	4.2	8.2
2001.....	51.1	6.6	*	*	*	*	1.3	2.2	3.6	8.1
2000.....	52.5	7.5	*	*	*	*	1.1	1.5	4.3	9.6
1999.....	52.2	4.6	*	*	*	*	0.8	1.8	3.6	9.1
Black, male										
2002.....	43.5	5.8	*	*	*	*	*	1.7	4.0	7.5
2001.....	44.6	8.1	*	*	*	*	*	2.8	3.4	8.8
2000.....	46.1	8.6	*	*	*	*	*	*	4.1	9.2
1999.....	45.9	*	*	*	*	*	*	1.9	3.7	8.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cerebrovascular diseases (I60-I69)										
Black, female										
2002.....	55.8	*	*	*	*	*	*	2.2	4.3	8.9
2001.....	56.9	*	*	*	*	*	1.7	1.7	3.8	7.4
2000.....	58.3	*	*	*	*	*	*	1.6	4.6	10.0
1999.....	57.9	*	*	*	*	*	*	1.8	3.6	9.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Subarachnoid hemorrhage (I60)										
All races 3/, both sexes										
2002.....	2.1	*	*	*	*	*	0.2	0.3	0.7	1.3
2001.....	2.2	*	*	*	*	0.1	0.2	0.4	0.7	1.3
2000.....	2.3	*	*	*	*	*	0.3	0.4	0.7	1.3
1999.....	2.3	*	*	*	*	0.1	0.2	0.4	0.7	1.6
All races 3/, male										
2002.....	1.5	*	*	*	*	*	0.3	0.3	0.6	1.1
2001.....	1.5	*	*	*	*	*	0.2	0.4	0.7	1.0
2000.....	1.6	*	*	*	*	*	0.3	0.4	0.7	1.0
1999.....	1.6	*	*	*	*	*	*	0.4	0.7	1.3
All races 3/, female										
2002.....	2.8	*	*	*	*	*	0.2	0.4	0.8	1.5
2001.....	2.8	*	*	*	*	*	0.2	0.3	0.7	1.5
2000.....	3.0	*	*	*	*	*	0.3	0.4	0.8	1.6
1999.....	3.0	*	*	*	*	*	0.2	0.4	0.8	1.8
White, both sexes										
2002.....	2.1	*	*	*	*	*	0.2	0.3	0.5	1.1
2001.....	2.1	*	*	*	*	*	0.2	0.3	0.6	1.1
2000.....	2.3	*	*	*	*	*	0.3	0.5	0.7	1.2
1999.....	2.3	*	*	*	*	*	0.2	0.4	0.7	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Subarachnoid hemorrhage (I60)										
Black, female										
2002.....	3.1	*	*	*	*	*	*	*	1.6	2.9
2001.....	3.2	*	*	*	*	*	*	*	*	3.0
2000.....	3.5	*	*	*	*	*	*	*	1.9	3.3
1999.....	3.3	*	*	*	*	*	*	*	1.4	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intracerebral and other intracranial hemorrhage (I61-I62)										
All races 3/, both sexes										
2002.....	9.4	1.0	*	*	*	0.1	0.2	0.3	0.6	1.3
2001.....	9.4	1.0	*	*	*	0.1	0.3	0.4	0.7	1.5
2000.....	9.7	1.1	*	*	*	0.1	0.3	0.3	0.7	1.5
1999.....	9.1	1.1	*	*	*	0.1	0.2	0.3	0.6	1.5
All races 3/, male										
2002.....	8.7	1.1	*	*	*	*	0.3	0.4	0.7	1.3
2001.....	8.8	1.2	*	*	*	0.2	0.3	0.5	0.8	1.8
2000.....	9.1	1.3	*	*	*	*	0.3	0.4	0.8	1.7
1999.....	8.5	1.3	*	*	*	*	0.2	0.3	0.7	1.7
All races 3/, female										
2002.....	10.0	1.0	*	*	*	*	*	0.3	0.6	1.3
2001.....	9.9	*	*	*	*	*	0.3	0.4	0.7	1.1
2000.....	10.4	*	*	*	*	*	0.3	0.2	0.6	1.3
1999.....	9.8	*	*	*	*	*	*	0.4	0.5	1.3
White, both sexes										
2002.....	9.5	0.9	*	*	*	*	0.2	0.3	0.5	1.0
2001.....	9.4	0.7	*	*	*	0.1	0.3	0.3	0.6	1.1
2000.....	9.8	0.7	*	*	*	*	0.3	0.3	0.5	1.0
1999.....	9.2	0.8	*	*	*	*	0.2	0.3	0.4	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intracerebral and other intracranial hemorrhage (I61-I62)										
White, male										
2002.....	8.7	*	*	*	*	*	0.3	0.4	0.5	1.0
2001.....	8.6	*	*	*	*	*	0.3	0.4	0.7	1.4
2000.....	8.9	*	*	*	*	*	0.3	0.4	0.6	1.2
1999.....	8.3	*	*	*	*	*	*	0.3	0.6	1.2
White, female										
2002.....	10.2	*	*	*	*	*	*	*	0.5	0.9
2001.....	10.1	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	10.7	*	*	*	*	*	*	*	0.4	0.9
1999.....	10.0	*	*	*	*	*	*	0.3	0.3	0.9
Black, both sexes										
2002.....	9.7	*	*	*	*	*	*	*	1.7	3.1
2001.....	10.0	*	*	*	*	*	*	1.1	1.6	3.3
2000.....	10.3	*	*	*	*	*	*	*	2.0	4.4
1999.....	9.5	*	*	*	*	*	*	*	1.4	4.4
Black, male										
2002.....	10.1	*	*	*	*	*	*	*	2.0	3.1
2001.....	10.3	*	*	*	*	*	*	*	1.5	4.2
2000.....	10.9	*	*	*	*	*	*	*	2.2	5.2
1999.....	9.8	*	*	*	*	*	*	*	1.5	5.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intracerebral and other intracranial hemorrhage (I61-I62)										
Black, female										
2002.....	9.4	*	*	*	*	*	*	*	1.4	3.2
2001.....	9.8	*	*	*	*	*	*	*	1.6	2.4
2000.....	9.8	*	*	*	*	*	*	*	1.7	3.8
1999.....	9.2	*	*	*	*	*	*	*	1.4	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Stroke, not specified as hemorrhage or infarction (I64)										
All races 3/, both sexes										
2002.....	30.1	*	*	*	*	*	*	0.1	0.3	0.6
2001.....	31.1	*	*	*	*	*	*	0.1	0.2	0.5
2000.....	32.1	*	*	*	*	*	*	0.2	0.2	0.5
1999.....	32.6	*	*	*	*	*	*	0.1	0.2	0.4
All races 3/, male										
2002.....	22.6	*	*	*	*	*	*	*	*	0.7
2001.....	23.5	*	*	*	*	*	*	*	*	0.5
2000.....	24.2	*	*	*	*	*	*	*	0.3	0.6
1999.....	24.7	*	*	*	*	*	*	*	0.2	0.4
All races 3/, female										
2002.....	37.4	*	*	*	*	*	*	*	0.4	0.5
2001.....	38.5	*	*	*	*	*	*	*	0.3	0.5
2000.....	39.7	*	*	*	*	*	*	*	*	0.5
1999.....	40.3	*	*	*	*	*	*	*	0.3	0.5
White, both sexes										
2002.....	32.1	*	*	*	*	*	*	*	0.2	0.4
2001.....	33.1	*	*	*	*	*	*	*	0.2	0.4
2000.....	34.2	*	*	*	*	*	*	*	0.2	0.4
1999.....	34.8	*	*	*	*	*	*	*	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other cerebrovascular diseases and their sequelae (I67,I69)										
All races 3/, both sexes										
2002.....	10.5	1.4	0.2	*	*	*	*	*	0.1	0.3
2001.....	10.3	1.2	0.2	*	*	*	*	*	0.2	0.3
2000.....	10.5	1.6	0.2	*	*	*	*	*	0.1	0.3
1999.....	10.6	1.1	0.1	*	*	*	*	*	0.1	0.2
All races 3/, male										
2002.....	7.9	1.7	0.3	*	*	*	*	*	*	0.3
2001.....	7.9	1.5	*	*	*	*	*	*	*	0.3
2000.....	8.1	1.8	*	*	*	*	*	*	*	0.2
1999.....	8.1	1.4	*	*	*	*	*	*	*	0.2
All races 3/, female										
2002.....	13.0	1.1	*	*	*	*	*	*	*	0.3
2001.....	12.7	*	*	*	*	*	*	*	*	0.3
2000.....	12.8	1.4	*	*	*	*	*	*	*	0.3
1999.....	12.9	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002.....	11.3	0.9	0.2	*	*	*	*	*	0.1	0.2
2001.....	11.1	0.9	0.2	*	*	*	*	*	0.2	0.3
2000.....	11.3	1.2	*	*	*	*	*	*	*	0.2
1999.....	11.4	0.9	*	*	*	*	*	*	0.1	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of arteries, arterioles and capillaries (I70-I78)										
All races 3/, both sexes										
2002.....	13.4	*	*	*	*	0.2	0.2	0.3	0.6	0.8
2001.....	13.8	*	*	*	*	0.1	0.2	0.4	0.5	0.9
2000.....	14.4	*	*	*	*	*	0.2	0.3	0.5	0.8
1999.....	14.6	*	*	*	*	*	0.2	0.3	0.5	0.8
All races 3/, male										
2002.....	13.0	*	*	*	*	*	0.3	0.3	0.7	0.8
2001.....	13.3	*	*	*	*	*	0.2	0.4	0.7	1.3
2000.....	14.1	*	*	*	*	*	0.3	0.4	0.7	0.9
1999.....	14.1	*	*	*	*	*	0.2	0.4	0.7	1.1
All races 3/, female										
2002.....	13.8	*	*	*	*	*	*	0.2	0.4	0.7
2001.....	14.4	*	*	*	*	*	*	0.3	0.3	0.6
2000.....	14.6	*	*	*	*	*	*	*	0.4	0.6
1999.....	15.2	*	*	*	*	*	*	*	0.3	0.6
White, both sexes										
2002.....	14.7	*	*	*	*	0.2	0.2	0.2	0.5	0.7
2001.....	15.1	*	*	*	*	*	0.2	0.3	0.5	0.8
2000.....	15.7	*	*	*	*	*	0.2	0.2	0.5	0.7
1999.....	16.0	*	*	*	*	*	0.2	0.3	0.4	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Aortic aneurysm and dissection (I71)										
All races 3/, both sexes										
2002.....	5.1	*	*	*	*	0.1	*	0.2	0.4	0.5
2001.....	5.3	*	*	*	*	*	0.1	0.2	0.3	0.6
2000.....	5.6	*	*	*	*	*	0.1	0.2	0.3	0.5
1999.....	5.7	*	*	*	*	*	0.1	0.2	0.3	0.6
All races 3/, male										
2002.....	6.4	*	*	*	*	*	*	0.3	0.5	0.6
2001.....	6.5	*	*	*	*	*	*	0.4	0.5	0.9
2000.....	7.0	*	*	*	*	*	*	0.3	0.5	0.6
1999.....	7.0	*	*	*	*	*	*	0.3	0.5	0.9
All races 3/, female										
2002.....	4.0	*	*	*	*	*	*	*	0.2	0.3
2001.....	4.2	*	*	*	*	*	*	*	*	0.3
2000.....	4.3	*	*	*	*	*	*	*	0.2	0.3
1999.....	4.4	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002.....	5.7	*	*	*	*	*	*	0.1	0.3	0.5
2001.....	5.9	*	*	*	*	*	0.1	0.2	0.3	0.5
2000.....	6.2	*	*	*	*	*	*	0.2	0.3	0.4
1999.....	6.3	*	*	*	*	*	*	0.2	0.3	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of arteries, arterioles and capillaries (I72-I78)										
All races 3/, both sexes										
2002.....	3.5	*	*	*	*	*	*	0.1	0.1	0.2
2001.....	3.5	*	*	*	*	*	*	0.1	0.2	0.3
2000.....	3.6	*	*	*	*	*	*	*	0.2	0.2
1999.....	3.6	*	*	*	*	*	*	*	0.1	0.2
All races 3/, male										
2002.....	3.0	*	*	*	*	*	*	*	*	*
2001.....	3.1	*	*	*	*	*	*	*	*	0.2
2000.....	3.2	*	*	*	*	*	*	*	*	0.2
1999.....	3.1	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	4.0	*	*	*	*	*	*	*	*	0.3
2001.....	4.0	*	*	*	*	*	*	*	*	0.3
2000.....	4.0	*	*	*	*	*	*	*	*	0.3
1999.....	4.1	*	*	*	*	*	*	*	*	0.2
White, both sexes										
2002.....	3.7	*	*	*	*	*	*	*	0.1	0.2
2001.....	3.7	*	*	*	*	*	*	*	0.1	0.2
2000.....	3.8	*	*	*	*	*	*	*	0.2	0.2
1999.....	3.8	*	*	*	*	*	*	*	0.1	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other disorders of circulatory system (I80-I99)										
All races 3/, both sexes										
2002.....	1.6	0.7	*	*	*	*	0.1	0.2	0.5	0.6
2001.....	1.6	0.5	*	*	*	*	0.1	0.2	0.4	0.4
2000.....	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.6
1999.....	1.4	0.6	*	*	*	*	*	0.2	0.3	0.4
All races 3/, male										
2002.....	1.4	*	*	*	*	*	*	*	0.5	0.6
2001.....	1.4	*	*	*	*	*	*	*	0.3	0.5
2000.....	1.4	*	*	*	*	*	*	*	0.4	0.6
1999.....	1.3	*	*	*	*	*	*	*	0.3	0.5
All races 3/, female										
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.6
2001.....	1.9	*	*	*	*	*	*	0.3	0.4	0.4
2000.....	1.8	*	*	*	*	*	*	0.4	0.4	0.5
1999.....	1.6	*	*	*	*	*	*	*	0.3	0.4
White, both sexes										
2002.....	1.6	0.6	*	*	*	*	*	0.2	0.4	0.5
2001.....	1.7	*	*	*	*	*	*	0.2	0.3	0.3
2000.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.5
1999.....	1.5	*	*	*	*	*	*	0.1	0.3	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)										
All races 3/, both sexes										
2002.....	1.2	*	*	*	*	*	0.1	0.2	0.4	0.5
2001.....	1.2	*	*	*	*	*	0.1	0.2	0.3	0.4
2000.....	1.2	*	*	*	*	*	0.1	0.3	0.3	0.5
1999.....	0.9	*	*	*	*	*	*	0.1	0.3	0.3
All races 3/, male										
2002.....	1.0	*	*	*	*	*	*	*	0.5	0.5
2001.....	1.0	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.0	*	*	*	*	*	*	*	0.3	0.5
1999.....	0.8	*	*	*	*	*	*	*	0.3	0.4
All races 3/, female										
2002.....	1.5	*	*	*	*	*	*	0.3	0.4	0.5
2001.....	1.5	*	*	*	*	*	*	0.3	0.4	0.4
2000.....	1.4	*	*	*	*	*	*	0.4	0.3	0.4
1999.....	1.1	*	*	*	*	*	*	*	0.3	0.3
White, both sexes										
2002.....	1.2	*	*	*	*	*	*	0.1	0.3	0.4
2001.....	1.2	*	*	*	*	*	*	0.2	0.3	0.3
2000.....	1.2	*	*	*	*	*	*	0.2	0.3	0.4
1999.....	0.9	*	*	*	*	*	*	*	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
X. Diseases of the respiratory system (J00-J98)										
All races 3/, both sexes										
2002.....	81.4	15.8	1.9	0.7	1.0	1.2	1.5	1.9	2.6	4.4
2001.....	80.8	17.3	1.8	0.6	0.8	1.0	1.6	1.9	2.8	4.7
2000.....	82.1	17.1	1.7	0.7	0.9	1.1	1.7	2.0	2.9	4.5
1999.....	82.4	18.1	2.1	0.7	0.9	1.3	1.6	2.1	2.9	4.6
All races 3/, male										
2002.....	78.9	18.8	2.2	0.7	0.9	1.4	1.6	1.9	2.8	4.6
2001.....	78.2	19.7	1.9	0.7	0.8	1.1	1.9	1.9	2.9	4.9
2000.....	80.2	18.9	1.7	0.8	0.9	1.3	1.9	2.2	3.1	4.7
1999.....	81.3	19.6	2.2	0.7	1.1	1.4	1.7	2.2	3.0	4.9
All races 3/, female										
2002.....	83.7	12.6	1.5	0.7	1.0	0.9	1.3	1.8	2.5	4.3
2001.....	83.2	14.7	1.8	0.6	0.7	0.9	1.4	1.9	2.6	4.4
2000.....	84.0	15.2	1.7	0.6	0.8	0.9	1.5	1.8	2.7	4.2
1999.....	83.4	16.6	1.9	0.6	0.8	1.1	1.4	2.1	2.8	4.4
White, both sexes										
2002.....	90.8	12.0	1.4	0.6	0.7	0.9	1.3	1.7	2.3	3.8
2001.....	89.7	12.3	1.5	0.5	0.7	0.8	1.3	1.5	2.2	3.8
2000.....	91.2	13.3	1.5	0.6	0.7	0.9	1.5	1.7	2.5	3.8
1999.....	91.2	13.3	1.6	0.6	0.8	1.1	1.2	1.6	2.3	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
X. Diseases of the respiratory system (J00-J98)										
White, male										
2002.....	86.7	14.6	1.7	0.6	0.6	1.1	1.4	1.6	2.5	3.7
2001.....	85.7	14.3	1.5	0.6	0.7	0.8	1.6	1.4	2.3	4.1
2000.....	87.7	15.1	1.3	0.7	0.7	1.0	1.7	1.7	2.5	3.9
1999.....	88.8	14.7	1.5	0.6	0.9	1.2	1.4	1.7	2.5	4.1
White, female										
2002.....	94.7	9.2	1.2	0.5	0.8	0.8	1.1	1.7	2.1	3.9
2001.....	93.7	10.2	1.6	0.4	0.6	0.8	1.0	1.6	2.2	3.5
2000.....	94.6	11.3	1.6	0.5	0.7	0.8	1.2	1.7	2.4	3.6
1999.....	93.6	11.8	1.7	0.6	0.6	1.0	1.1	1.5	2.2	3.7
Black, both sexes										
2002.....	46.7	34.1	4.1	1.2	2.3	2.5	2.9	3.5	5.6	9.4
2001.....	47.2	39.6	3.2	1.2	1.5	2.0	3.6	4.3	6.8	11.1
2000.....	48.0	37.7	3.1	1.2	1.9	2.5	3.4	4.4	6.1	9.6
1999.....	49.2	41.6	4.0	1.1	2.1	2.0	3.6	5.7	6.9	10.2
Black, male										
2002.....	50.2	39.2	4.7	*	2.4	3.4	3.2	4.0	6.3	10.7
2001.....	50.5	43.7	3.8	1.2	1.7	2.6	3.5	4.6	7.8	11.3
2000.....	52.4	39.3	3.9	1.4	2.2	3.1	3.7	5.9	7.7	10.7
1999.....	54.0	44.6	5.1	1.4	2.5	2.3	4.3	5.6	7.3	10.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
X. Diseases of the respiratory system (J00-J98)										
Black, female										
2002.....	43.5	28.8	3.5	1.4	2.2	1.5	2.7	3.1	4.9	8.3
2001.....	44.1	35.3	2.5	*	1.4	1.4	3.6	4.0	5.9	10.9
2000.....	44.1	36.0	2.2	*	1.5	1.9	3.2	3.0	4.6	8.6
1999.....	44.8	38.6	2.9	*	1.6	1.8	2.9	5.7	6.5	9.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of the respiratory system (J10-J98)										
All races 3/, both sexes										
2002.....	81.3	15.5	1.8	0.7	0.9	1.1	1.4	1.9	2.6	4.4
2001.....	80.7	17.0	1.7	0.6	0.8	1.0	1.6	1.9	2.8	4.7
2000.....	82.0	16.9	1.7	0.7	0.9	1.1	1.7	2.0	2.9	4.5
1999.....	82.3	17.8	2.0	0.7	0.9	1.2	1.6	2.1	2.9	4.6
All races 3/, male										
2002.....	78.9	18.6	2.1	0.7	0.9	1.4	1.6	1.9	2.8	4.5
2001.....	78.1	19.4	1.8	0.6	0.8	1.1	1.9	1.8	2.9	4.9
2000.....	80.1	18.7	1.7	0.8	0.9	1.3	1.9	2.2	3.1	4.7
1999.....	81.3	19.2	2.1	0.7	1.1	1.3	1.7	2.1	3.0	4.8
All races 3/, female										
2002.....	83.7	12.4	1.4	0.6	1.0	0.9	1.3	1.8	2.5	4.3
2001.....	83.2	14.6	1.7	0.6	0.7	0.9	1.4	1.9	2.6	4.4
2000.....	83.9	15.1	1.6	0.6	0.8	0.9	1.5	1.8	2.7	4.2
1999.....	83.3	16.4	1.9	0.6	0.7	1.1	1.4	2.0	2.8	4.4
White, both sexes										
2002.....	90.7	11.9	1.3	0.6	0.7	0.9	1.2	1.7	2.3	3.8
2001.....	89.7	12.1	1.4	0.5	0.7	0.8	1.3	1.5	2.2	3.8
2000.....	91.1	13.1	1.4	0.6	0.7	0.9	1.4	1.7	2.4	3.8
1999.....	91.2	13.1	1.5	0.6	0.7	1.1	1.2	1.6	2.3	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of the respiratory system (J10-J98)										
White, male										
2002.....	86.7	14.5	1.6	0.6	0.6	1.1	1.4	1.6	2.5	3.7
2001.....	85.6	14.1	1.4	0.5	0.7	0.8	1.6	1.4	2.3	4.1
2000.....	87.6	14.8	1.3	0.7	0.7	1.0	1.7	1.7	2.4	3.9
1999.....	88.7	14.5	1.4	0.6	0.9	1.2	1.4	1.7	2.5	4.1
White, female										
2002.....	94.7	9.0	1.1	0.5	0.8	0.8	1.1	1.7	2.1	3.9
2001.....	93.6	10.0	1.5	0.4	0.6	0.8	1.0	1.6	2.2	3.5
2000.....	94.5	11.3	1.6	0.5	0.7	0.7	1.2	1.7	2.4	3.6
1999.....	93.5	11.7	1.6	0.6	0.6	1.0	1.1	1.5	2.2	3.7
Black, both sexes										
2002.....	46.6	33.5	4.0	1.2	2.3	2.5	2.9	3.5	5.6	9.4
2001.....	47.1	39.1	3.0	1.2	1.5	1.9	3.6	4.3	6.8	11.1
2000.....	48.0	37.6	3.0	1.2	1.8	2.5	3.3	4.3	6.1	9.6
1999.....	49.1	41.0	3.9	1.0	2.1	2.0	3.6	5.7	6.7	10.1
Black, male										
2002.....	50.2	38.3	4.7	*	2.4	3.4	3.1	4.0	6.3	10.7
2001.....	50.4	42.8	3.6	1.2	1.7	2.5	3.5	4.6	7.8	11.2
2000.....	52.4	39.3	3.9	1.4	2.2	3.1	3.5	5.8	7.7	10.7
1999.....	53.9	43.9	4.8	1.3	2.5	2.2	4.3	5.6	7.0	10.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of the respiratory system (J10-J98)										
Black, female										
2002.....	43.4	28.5	3.4	1.4	2.2	1.5	2.6	3.1	4.9	8.3
2001.....	44.1	35.3	2.5	*	1.3	1.4	3.6	4.0	5.9	10.9
2000.....	44.0	35.7	2.2	*	1.5	1.9	3.1	3.0	4.6	8.6
1999.....	44.7	37.9	2.9	*	1.6	1.8	2.9	5.7	6.4	9.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
All races 3/, both sexes										
2002.....	22.5	6.3	0.7	0.2	0.2	0.3	0.4	0.7	1.0	1.7
2001.....	21.7	7.2	0.7	0.2	0.2	0.3	0.6	0.6	1.0	1.8
2000.....	22.6	7.4	0.6	0.2	0.2	0.3	0.6	0.7	1.1	1.7
1999.....	22.2	8.1	0.8	0.2	0.2	0.3	0.5	0.7	1.0	2.0
All races 3/, male										
2002.....	20.2	7.6	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8
2001.....	19.5	7.8	0.7	0.2	0.2	0.3	0.7	0.6	1.2	2.0
2000.....	20.3	8.1	0.6	0.2	*	0.3	0.7	0.8	1.2	1.8
1999.....	19.8	8.5	0.7	0.2	0.2	0.3	0.6	0.8	1.1	2.2
All races 3/, female										
2002.....	24.8	5.0	0.6	*	0.2	0.3	0.4	0.7	0.8	1.5
2001.....	23.8	6.7	0.7	*	0.2	0.3	0.5	0.6	0.8	1.6
2000.....	24.8	6.6	0.6	*	*	0.3	0.6	0.6	0.9	1.7
1999.....	24.6	7.7	0.8	*	*	0.3	0.5	0.6	0.8	1.8
White, both sexes										
2002.....	24.6	5.0	0.5	0.2	0.2	0.3	0.4	0.7	0.9	1.4
2001.....	23.5	5.1	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.5
2000.....	24.4	5.6	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.6
1999.....	24.1	5.9	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
White, male										
2002.....	21.7	6.4	0.5	*	*	0.4	0.5	0.6	1.0	1.5
2001.....	20.6	5.8	0.6	*	*	0.3	0.7	0.5	1.0	1.6
2000.....	21.5	6.1	0.4	0.3	*	0.3	0.6	0.6	1.0	1.7
1999.....	21.0	6.7	0.4	*	0.2	0.3	0.5	0.6	0.9	1.9
White, female										
2002.....	27.4	3.6	0.5	*	*	*	0.4	0.7	0.7	1.4
2001.....	26.2	4.4	0.6	*	*	0.3	0.3	0.6	0.7	1.4
2000.....	27.3	5.0	0.6	*	*	*	0.5	0.6	0.9	1.5
1999.....	27.1	5.2	0.7	*	*	0.3	0.4	0.4	0.6	1.6
Black, both sexes										
2002.....	15.5	12.7	1.5	*	*	*	0.7	1.0	2.0	3.5
2001.....	15.5	16.3	1.0	*	*	*	1.0	1.2	2.0	3.7
2000.....	16.1	16.4	1.2	*	*	*	1.0	1.4	2.0	2.9
1999.....	16.1	18.3	1.7	*	*	*	1.3	1.7	2.2	4.0
Black, male										
2002.....	15.3	13.7	1.6	*	*	*	*	*	2.4	4.0
2001.....	15.8	15.9	*	*	*	*	*	*	2.7	4.5
2000.....	16.5	17.9	1.7	*	*	*	*	2.0	2.5	3.0
1999.....	16.3	17.9	1.9	*	*	*	*	1.9	2.5	4.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
Black, female										
2002.....	15.6	11.8	*	*	*	*	*	*	1.6	3.1
2001.....	15.1	16.7	*	*	*	*	*	*	1.4	3.0
2000.....	15.8	14.9	*	*	*	*	*	*	1.6	2.7
1999.....	15.9	18.8	*	*	*	*	*	1.5	1.9	3.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia due to other or unspecified organisms (J16,J18)										
All races 3/, both sexes										
2002.....	20.8	3.9	0.5	0.1	0.2	0.3	0.4	0.6	0.8	1.4
2001.....	20.0	4.6	0.5	0.2	0.2	0.3	0.5	0.5	0.9	1.5
2000.....	20.8	5.1	0.4	0.2	0.1	0.2	0.5	0.6	0.9	1.4
1999.....	20.7	4.7	0.5	0.1	0.2	0.3	0.5	0.6	0.9	1.6
All races 3/, male										
2002.....	18.5	4.6	0.5	*	*	0.3	0.5	0.5	1.0	1.6
2001.....	17.8	4.9	0.5	*	0.2	0.3	0.6	0.5	1.0	1.7
2000.....	18.5	5.3	0.4	*	*	0.3	0.6	0.7	1.0	1.5
1999.....	18.3	5.0	0.5	*	*	0.3	0.5	0.6	1.0	1.8
All races 3/, female										
2002.....	23.0	3.1	0.4	*	*	0.3	0.3	0.6	0.7	1.3
2001.....	22.2	4.2	0.5	*	*	0.3	0.4	0.5	0.7	1.3
2000.....	23.1	4.9	0.4	*	*	0.2	0.5	0.4	0.8	1.4
1999.....	23.0	4.3	0.5	*	*	0.3	0.4	0.5	0.7	1.4
White, both sexes										
2002.....	22.7	2.9	0.4	0.1	0.2	0.3	0.4	0.5	0.7	1.2
2001.....	21.7	3.0	0.5	0.2	0.2	0.2	0.5	0.4	0.7	1.3
2000.....	22.6	3.9	0.3	0.2	0.1	0.2	0.5	0.5	0.8	1.3
1999.....	22.5	3.3	0.4	0.1	0.2	0.3	0.4	0.5	0.7	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia due to other or unspecified organisms (J16,J18)										
White, male										
2002.....	19.9	3.7	0.4	*	*	0.3	0.5	0.5	0.9	1.3
2001.....	18.9	3.5	0.5	*	*	*	0.6	0.4	0.8	1.4
2000.....	19.6	4.3	*	*	*	0.2	0.6	0.6	0.9	1.4
1999.....	19.5	3.8	*	*	*	0.3	0.4	0.5	0.8	1.6
White, female										
2002.....	25.5	2.0	0.3	*	*	*	0.3	0.5	0.6	1.2
2001.....	24.5	2.5	0.4	*	*	0.3	0.3	0.5	0.6	1.1
2000.....	25.5	3.5	0.4	*	*	*	0.4	0.4	0.7	1.2
1999.....	25.4	2.8	0.4	*	*	0.3	0.3	0.4	0.6	1.2
Black, both sexes										
2002.....	14.2	8.2	1.0	*	*	*	*	0.9	1.6	3.0
2001.....	14.1	11.8	0.8	*	*	*	1.0	1.1	1.7	3.3
2000.....	14.7	11.2	0.8	*	*	*	0.9	1.2	1.9	2.4
1999.....	14.7	11.4	1.1	*	*	*	1.1	1.4	1.9	3.4
Black, male										
2002.....	13.9	8.1	*	*	*	*	*	*	2.2	3.3
2001.....	14.3	11.1	*	*	*	*	*	*	2.4	3.8
2000.....	14.9	10.6	*	*	*	*	*	1.8	2.3	2.5
1999.....	14.8	12.0	*	*	*	*	*	1.6	2.4	3.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia due to other or unspecified organisms (J16,J18)										
Black, female										
2002.....	14.4	8.2	*	*	*	*	*	*	*	2.8
2001.....	14.0	12.6	*	*	*	*	*	*	*	2.8
2000.....	14.6	11.9	*	*	*	*	*	*	1.5	2.4
1999.....	14.6	10.7	*	*	*	*	*	*	1.5	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic lower respiratory diseases (J40-J47)										
All races 3/, both sexes										
2002.....	43.3	1.0	0.4	0.2	0.4	0.5	0.5	0.6	0.9	1.6
2001.....	43.2	1.0	0.3	0.2	0.3	0.4	0.5	0.5	0.9	1.6
2000.....	43.4	0.9	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
1999.....	44.5	0.9	0.4	0.2	0.4	0.5	0.5	0.7	0.9	1.3
All races 3/, male										
2002.....	42.9	1.1	0.6	0.2	0.5	0.6	0.5	0.7	0.9	1.5
2001.....	42.7	1.1	0.4	0.2	0.3	0.4	0.6	0.6	0.9	1.5
2000.....	43.5	1.2	0.4	0.2	0.5	0.6	0.6	0.7	0.9	1.3
1999.....	45.6	*	0.4	0.3	0.6	0.6	0.7	0.7	0.8	1.2
All races 3/, female										
2002.....	43.7	*	0.3	*	0.4	0.3	0.5	0.5	0.9	1.7
2001.....	43.7	*	*	*	0.2	0.3	0.4	0.5	0.9	1.7
2000.....	43.2	*	0.3	0.2	0.4	0.3	0.5	0.6	0.8	1.4
1999.....	43.4	*	0.3	0.2	0.3	0.5	0.4	0.8	1.0	1.4
White, both sexes										
2002.....	49.2	0.8	0.2	*	0.2	0.3	0.4	0.5	0.7	1.3
2001.....	49.0	0.7	0.2	*	0.2	0.3	0.3	0.4	0.6	1.2
2000.....	49.0	0.8	0.2	0.1	0.2	0.3	0.4	0.4	0.6	1.0
1999.....	50.2	*	0.3	0.2	0.3	0.4	0.4	0.5	0.6	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic lower respiratory diseases (J40-J47)										
White, male										
2002.....	47.8	*	0.3	*	*	0.4	0.4	0.5	0.8	1.1
2001.....	47.6	*	*	*	*	*	0.3	0.4	0.5	1.2
2000.....	48.3	*	*	*	0.3	0.3	0.5	0.5	0.6	0.9
1999.....	50.6	*	*	*	0.4	0.4	0.4	0.4	0.5	1.0
White, female										
2002.....	50.5	*	*	*	*	*	0.4	0.4	0.7	1.5
2001.....	50.3	*	*	*	*	0.3	*	0.3	0.7	1.2
2000.....	49.7	*	*	*	*	*	0.3	0.4	0.6	1.2
1999.....	49.8	*	*	*	*	0.4	0.4	0.5	0.7	1.2
Black, both sexes										
2002.....	20.7	*	1.2	0.6	1.6	1.4	1.2	1.6	2.2	3.8
2001.....	20.4	*	0.9	0.7	0.9	1.0	1.6	1.8	3.0	4.5
2000.....	20.8	*	0.8	0.7	1.4	1.4	1.3	1.9	2.4	3.5
1999.....	21.9	*	*	0.6	1.5	1.2	1.5	2.5	2.7	3.2
Black, male										
2002.....	24.1	*	1.6	*	1.7	2.2	1.5	2.2	2.3	4.5
2001.....	23.6	*	*	*	*	1.4	1.8	2.1	3.4	4.3
2000.....	24.3	*	*	*	1.7	2.0	1.5	2.4	3.1	3.7
1999.....	26.2	*	*	*	2.1	1.5	2.2	2.6	2.7	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic lower respiratory diseases (J40-J47)										
Black, female										
2002.....	17.7	*	*	*	1.4	*	*	*	2.1	3.2
2001.....	17.5	*	*	*	*	*	1.4	1.5	2.6	4.7
2000.....	17.6	*	*	*	*	*	*	1.4	1.7	3.3
1999.....	18.0	*	*	*	*	*	*	2.4	2.8	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
All races 3/, both sexes										
2002.....	42.9	1.0	0.4	0.2	0.4	0.5	0.5	0.6	0.9	1.5
2001.....	42.9	1.0	0.3	0.2	0.3	0.4	0.5	0.5	0.9	1.6
2000.....	43.0	0.9	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.3
1999.....	44.2	0.9	0.3	0.2	0.4	0.5	0.5	0.7	0.9	1.3
All races 3/, male										
2002.....	42.7	1.1	0.6	0.2	0.5	0.6	0.5	0.7	0.9	1.4
2001.....	42.5	1.1	0.4	0.2	0.3	0.4	0.5	0.6	0.9	1.5
2000.....	43.3	1.2	0.4	0.2	0.5	0.5	0.6	0.7	0.9	1.3
1999.....	45.4	*	0.4	0.3	0.6	0.6	0.6	0.7	0.8	1.2
All races 3/, female										
2002.....	43.2	*	0.3	*	0.4	0.3	0.5	0.5	0.8	1.6
2001.....	43.2	*	*	*	0.2	0.3	0.4	0.5	0.9	1.6
2000.....	42.8	*	0.3	0.2	0.4	0.3	0.4	0.5	0.7	1.4
1999.....	43.0	*	0.3	0.2	0.3	0.5	0.4	0.7	1.0	1.4
White, both sexes										
2002.....	48.8	0.8	0.2	*	0.2	0.3	0.4	0.4	0.7	1.3
2001.....	48.6	0.7	0.2	*	0.2	0.2	0.3	0.3	0.6	1.2
2000.....	48.7	0.8	0.2	0.1	0.2	0.2	0.4	0.4	0.6	1.0
1999.....	49.8	*	0.3	0.2	0.3	0.4	0.4	0.4	0.6	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
White, male										
2002.....	47.6	*	0.3	*	*	0.4	0.3	0.5	0.7	1.1
2001.....	47.4	*	*	*	*	*	0.3	0.4	0.5	1.2
2000.....	48.1	*	*	*	0.3	0.3	0.5	0.4	0.6	0.9
1999.....	50.3	*	*	*	0.4	0.4	0.4	0.4	0.5	0.9
White, female										
2002.....	49.9	*	*	*	*	*	0.4	0.4	0.7	1.5
2001.....	49.8	*	*	*	*	0.3	*	0.3	0.7	1.1
2000.....	49.2	*	*	*	*	*	0.3	0.4	0.6	1.2
1999.....	49.3	*	*	*	*	0.4	0.4	0.5	0.7	1.2
Black, both sexes										
2002.....	20.7	*	1.2	0.6	1.6	1.4	1.2	1.6	2.2	3.8
2001.....	20.3	*	0.9	0.7	0.9	1.0	1.6	1.7	2.9	4.5
2000.....	20.7	*	0.8	0.7	1.4	1.4	1.3	1.9	2.3	3.5
1999.....	21.8	*	*	0.6	1.5	1.2	1.5	2.4	2.7	3.2
Black, male										
2002.....	24.1	*	1.6	*	1.7	2.2	1.5	2.2	2.3	4.5
2001.....	23.6	*	*	*	*	1.4	1.8	2.1	3.4	4.3
2000.....	24.3	*	*	*	1.7	1.9	1.5	2.4	3.1	3.7
1999.....	26.1	*	*	*	2.1	1.5	2.2	2.6	2.7	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
Black, female										
2002.....	17.6	*	*	*	1.4	*	*	*	2.1	3.2
2001.....	17.4	*	*	*	*	*	1.4	1.4	2.5	4.6
2000.....	17.5	*	*	*	*	*	*	1.4	1.6	3.3
1999.....	17.9	*	*	*	*	*	*	2.2	2.8	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
All races 3/, both sexes										
2002.....	1.5	*	0.3	0.2	0.4	0.4	0.4	0.5	0.6	0.9
2001.....	1.5	*	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9
2000.....	1.6	*	0.2	0.2	0.4	0.4	0.5	0.5	0.7	0.8
1999.....	1.7	*	0.2	0.2	0.4	0.5	0.5	0.6	0.7	0.9
All races 3/, male										
2002.....	1.1	*	0.4	0.2	0.5	0.6	0.5	0.7	0.6	0.8
2001.....	1.1	*	0.3	0.2	0.3	0.3	0.4	0.5	0.6	0.8
2000.....	1.2	*	0.3	0.2	0.5	0.5	0.5	0.7	0.8	0.7
1999.....	1.2	*	*	0.2	0.6	0.5	0.6	0.6	0.6	0.8
All races 3/, female										
2002.....	1.8	*	*	*	0.4	0.2	0.4	0.4	0.6	1.0
2001.....	1.9	*	*	*	0.2	0.3	0.4	0.4	0.8	1.0
2000.....	2.0	*	*	0.2	0.3	0.3	0.4	0.4	0.6	0.9
1999.....	2.1	*	*	*	0.3	0.4	0.4	0.6	0.8	1.0
White, both sexes										
2002.....	1.3	*	*	*	0.2	0.3	0.3	0.4	0.5	0.7
2001.....	1.3	*	*	*	0.2	0.2	0.2	0.3	0.4	0.5
2000.....	1.4	*	*	*	0.2	0.2	0.4	0.3	0.5	0.5
1999.....	1.5	*	*	0.1	0.2	0.4	0.3	0.4	0.4	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
White, male										
2002.....	0.9	*	*	*	*	0.3	0.3	0.4	0.5	0.5
2001.....	0.8	*	*	*	*	*	0.2	0.3	0.3	0.5
2000.....	0.9	*	*	*	0.2	0.3	0.4	0.4	0.5	0.4
1999.....	0.9	*	*	*	0.3	0.4	0.4	0.4	0.4	0.6
White, female										
2002.....	1.7	*	*	*	*	*	0.3	0.4	0.6	0.9
2001.....	1.7	*	*	*	*	*	*	*	0.6	0.6
2000.....	1.8	*	*	*	*	*	0.3	0.3	0.5	0.6
1999.....	2.0	*	*	*	*	0.4	*	0.4	0.5	0.8
Black, both sexes										
2002.....	2.9	*	0.9	*	1.5	1.3	1.2	1.6	1.6	2.9
2001.....	3.0	*	0.8	0.7	0.8	0.9	1.4	1.7	2.6	3.4
2000.....	3.2	*	*	0.7	1.4	1.3	1.2	1.8	2.0	2.7
1999.....	3.2	*	*	0.6	1.4	1.1	1.4	2.3	2.4	2.6
Black, male										
2002.....	2.8	*	*	*	1.7	2.1	1.5	2.2	1.7	3.4
2001.....	2.6	*	*	*	*	*	1.5	1.9	3.0	3.0
2000.....	2.8	*	*	*	1.7	1.7	*	2.3	2.6	2.7
1999.....	2.8	*	*	*	2.0	1.3	2.0	2.6	2.6	2.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
Black, female										
2002.....	3.0	*	*	*	1.4	*	*	*	1.4	2.3
2001.....	3.3	*	*	*	*	*	*	1.4	2.3	3.7
2000.....	3.5	*	*	*	*	*	*	*	1.4	2.6
1999.....	3.5	*	*	*	*	*	*	2.0	2.3	2.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Lung diseases due to external agents (J60-J70)										
All races 3/, both sexes										
2002.....	6.5	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.5	*	*	*	*	*	0.1	0.2	0.2	0.3
2000.....	6.3	*	*	*	*	*	0.1	0.1	0.2	0.3
1999.....	5.9	*	*	*	*	*	0.1	0.2	0.2	0.3
All races 3/, male										
2002.....	7.0	*	*	*	*	*	*	*	0.2	0.4
2001.....	7.0	*	*	*	*	*	*	*	0.3	0.4
2000.....	6.9	*	*	*	*	*	*	*	0.3	0.4
1999.....	6.5	*	*	*	*	*	*	0.2	0.3	0.3
All races 3/, female										
2002.....	6.0	*	*	*	*	*	*	*	*	0.3
2001.....	6.1	*	*	*	*	*	*	*	0.2	0.2
2000.....	5.8	*	*	*	*	*	*	*	*	0.2
1999.....	5.3	*	*	*	*	*	*	*	0.2	0.2
White, both sexes										
2002.....	7.2	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	7.1	*	*	*	*	*	*	0.2	0.2	0.3
2000.....	7.0	*	*	*	*	*	*	*	0.2	0.2
1999.....	6.4	*	*	*	*	*	*	0.2	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonitis due to solids and liquids (J69)										
All races 3/, both sexes										
2002.....	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2000.....	5.9	*	*	*	*	*	0.1	0.1	0.2	0.3
1999.....	5.5	*	*	*	*	*	0.1	0.2	0.2	0.3
All races 3/, male										
2002.....	6.3	*	*	*	*	*	*	*	0.2	0.4
2001.....	6.1	*	*	*	*	*	*	*	0.2	0.4
2000.....	6.1	*	*	*	*	*	*	*	0.2	0.4
1999.....	5.7	*	*	*	*	*	*	0.2	0.3	0.3
All races 3/, female										
2002.....	5.9	*	*	*	*	*	*	*	*	0.3
2001.....	6.0	*	*	*	*	*	*	*	0.2	*
2000.....	5.8	*	*	*	*	*	*	*	*	0.2
1999.....	5.3	*	*	*	*	*	*	*	*	0.2
White, both sexes										
2002.....	6.7	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.6	*	*	*	*	*	*	0.2	0.2	0.2
2000.....	6.5	*	*	*	*	*	*	*	0.2	0.2
1999.....	5.9	*	*	*	*	*	*	0.1	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonitis due to food and vomit (J69.0)										
All races 3/, both sexes										
2002.....	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2000.....	5.9	*	*	*	*	*	0.1	0.1	0.2	0.3
1999.....	5.5	*	*	*	*	*	0.1	0.2	0.2	0.3
All races 3/, male										
2002.....	6.3	*	*	*	*	*	*	*	0.2	0.4
2001.....	6.1	*	*	*	*	*	*	*	0.2	0.4
2000.....	6.1	*	*	*	*	*	*	*	0.2	0.4
1999.....	5.7	*	*	*	*	*	*	0.2	0.3	0.3
All races 3/, female										
2002.....	5.9	*	*	*	*	*	*	*	*	0.3
2001.....	6.0	*	*	*	*	*	*	*	0.2	*
2000.....	5.8	*	*	*	*	*	*	*	*	0.2
1999.....	5.3	*	*	*	*	*	*	*	*	0.2
White, both sexes										
2002.....	6.7	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.6	*	*	*	*	*	*	0.2	0.2	0.2
2000.....	6.5	*	*	*	*	*	*	*	0.2	0.2
1999.....	5.9	*	*	*	*	*	*	0.1	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases of respiratory system (J80-J84,J93-J98)										
All races 3/, both sexes										
2002.....	7.9	6.1	0.4	0.1	0.1	0.2	0.3	0.3	0.5	0.7
2001.....	8.3	6.5	0.4	0.1	0.2	0.2	0.4	0.4	0.5	0.8
2000.....	8.3	6.4	0.4	0.1	0.2	0.3	0.3	0.4	0.6	0.9
1999.....	8.1	5.8	0.4	0.2	0.1	0.2	0.2	0.4	0.6	0.8
All races 3/, male										
2002.....	7.7	7.3	0.5	*	*	0.2	0.3	0.3	0.4	0.7
2001.....	8.1	8.1	0.4	*	0.2	0.2	0.4	0.3	0.4	0.8
2000.....	8.1	7.1	0.4	*	*	0.2	0.4	0.4	0.6	1.0
1999.....	7.9	6.6	0.4	*	*	0.3	0.3	0.3	0.6	0.8
All races 3/, female										
2002.....	8.0	4.9	0.3	*	*	0.2	0.3	0.4	0.5	0.8
2001.....	8.6	4.9	0.4	*	*	*	0.3	0.5	0.5	0.9
2000.....	8.4	5.7	0.3	*	*	0.3	0.2	0.4	0.6	0.8
1999.....	8.3	4.9	0.3	*	*	*	*	0.4	0.7	0.8
White, both sexes										
2002.....	8.6	4.2	0.3	*	*	0.2	0.2	0.3	0.4	0.6
2001.....	9.1	4.5	0.4	*	0.2	0.2	0.3	0.3	0.4	0.6
2000.....	9.0	4.9	0.4	0.1	0.2	0.2	0.3	0.4	0.5	0.7
1999.....	8.8	4.1	0.3	0.1	0.1	0.2	0.3	0.3	0.6	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases of respiratory system (J80-J84,J93-J98)										
Black, female										
2002.....	5.4	11.8	*	*	*	*	*	*	*	1.6
2001.....	6.2	11.7	*	*	*	*	*	*	*	2.6
2000.....	5.7	15.2	*	*	*	*	*	*	*	1.7
1999.....	5.9	12.1	*	*	*	*	*	*	1.5	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XI. Diseases of the digestive system (K00-K92)										
All races 3/, both sexes										
2002.....	30.1	12.6	0.8	0.2	0.3	0.4	0.9	1.4	3.2	7.3
2001.....	30.1	11.9	0.8	0.3	0.3	0.4	0.7	1.4	3.2	8.1
2000.....	29.9	13.1	0.7	0.3	0.3	0.4	1.0	1.4	3.3	8.0
1999.....	29.7	13.2	0.7	0.3	0.4	0.5	0.7	1.3	3.5	7.9
All races 3/, male										
2002.....	30.7	15.1	0.9	0.3	0.3	0.4	0.9	1.5	4.1	9.1
2001.....	30.6	13.4	0.9	0.3	0.4	0.4	0.8	1.6	4.0	10.3
2000.....	30.3	14.8	0.9	0.3	0.3	0.4	1.1	1.8	4.1	10.2
1999.....	30.4	16.5	0.7	0.2	0.4	0.5	0.7	1.5	4.3	9.9
All races 3/, female										
2002.....	29.5	10.0	0.7	*	0.4	0.4	0.8	1.3	2.3	5.6
2001.....	29.7	10.3	0.6	0.3	0.3	0.3	0.7	1.2	2.4	6.0
2000.....	29.4	11.4	0.5	0.3	0.4	0.3	0.8	1.0	2.5	5.8
1999.....	29.1	9.7	0.8	0.3	0.4	0.4	0.8	1.1	2.7	5.9
White, both sexes										
2002.....	32.2	10.1	0.7	0.2	0.3	0.4	0.8	1.2	3.0	7.2
2001.....	32.1	8.7	0.7	0.3	0.3	0.3	0.6	1.2	3.0	7.9
2000.....	31.7	10.0	0.5	0.2	0.3	0.3	0.9	1.2	3.1	7.6
1999.....	31.5	10.1	0.7	0.2	0.4	0.4	0.6	1.2	3.3	7.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XI. Diseases of the digestive system (K00-K92)										
White, male										
2002.....	32.4	12.1	0.7	0.3	0.3	0.4	0.9	1.2	4.0	9.0
2001.....	31.9	9.3	0.8	0.3	0.3	0.4	0.6	1.4	3.8	9.9
2000.....	31.7	11.2	0.6	0.3	0.3	0.4	1.0	1.7	3.9	9.8
1999.....	31.7	12.5	0.5	*	0.3	0.5	0.6	1.4	4.0	9.7
White, female										
2002.....	32.1	7.9	0.7	*	0.3	0.4	0.6	1.1	1.9	5.4
2001.....	32.2	8.1	0.5	0.3	0.3	0.3	0.5	1.1	2.3	5.9
2000.....	31.6	8.8	0.4	*	0.3	0.3	0.7	0.7	2.2	5.3
1999.....	31.2	7.5	0.8	0.3	0.4	0.4	0.6	1.0	2.5	5.5
Black, both sexes										
2002.....	23.7	25.9	1.4	*	*	0.6	1.5	2.5	4.3	8.0
2001.....	25.1	28.6	1.2	*	0.6	0.6	1.5	2.6	4.3	10.1
2000.....	25.2	29.6	1.5	*	*	*	1.6	2.9	4.7	10.8
1999.....	25.1	29.6	1.1	*	*	0.7	1.8	2.1	4.6	10.1
Black, male										
2002.....	26.0	31.4	1.7	*	*	*	*	2.9	4.9	9.1
2001.....	28.4	33.5	*	*	*	*	*	3.3	5.3	13.2
2000.....	27.7	33.9	2.0	*	*	*	2.1	3.0	5.1	13.6
1999.....	27.8	36.4	*	*	*	*	1.6	2.2	5.7	12.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XI. Diseases of the digestive system (K00-K92)										
Black, female										
2002.....	21.5	20.3	*	*	*	*	1.8	2.1	3.8	7.0
2001.....	22.1	23.3	*	*	*	*	1.7	2.0	3.4	7.3
2000.....	23.0	25.1	*	*	*	*	*	2.8	4.3	8.4
1999.....	22.7	22.5	*	*	*	*	1.9	2.0	3.6	8.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of other parts of digestive system (K20-K92)										
All races 3/, both sexes										
2002.....	30.0	12.5	0.8	0.2	0.3	0.4	0.9	1.4	3.2	7.3
2001.....	30.1	11.8	0.8	0.3	0.3	0.4	0.7	1.4	3.2	8.1
2000.....	29.8	13.0	0.7	0.3	0.3	0.4	1.0	1.4	3.3	8.0
1999.....	29.7	13.1	0.7	0.3	0.4	0.5	0.7	1.3	3.5	7.9
All races 3/, male										
2002.....	30.6	15.0	0.9	0.3	0.3	0.4	0.9	1.5	4.1	9.1
2001.....	30.5	13.3	0.9	0.3	0.4	0.4	0.8	1.6	4.0	10.3
2000.....	30.3	14.8	0.9	0.3	0.3	0.4	1.1	1.8	4.1	10.2
1999.....	30.3	16.4	0.7	0.2	0.4	0.5	0.7	1.5	4.3	9.9
All races 3/, female										
2002.....	29.4	9.9	0.7	*	0.4	0.4	0.8	1.3	2.3	5.6
2001.....	29.7	10.2	0.6	0.3	0.3	0.3	0.7	1.2	2.4	6.0
2000.....	29.4	11.2	0.5	0.3	0.4	0.3	0.8	1.0	2.5	5.8
1999.....	29.0	9.7	0.8	0.3	0.4	0.4	0.7	1.1	2.7	5.9
White, both sexes										
2002.....	32.2	10.0	0.7	0.2	0.3	0.4	0.8	1.2	3.0	7.2
2001.....	32.0	8.6	0.7	0.3	0.3	0.3	0.5	1.2	3.0	7.9
2000.....	31.6	9.9	0.5	0.2	0.3	0.3	0.9	1.2	3.0	7.6
1999.....	31.4	10.0	0.7	0.2	0.4	0.4	0.6	1.2	3.3	7.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of other parts of digestive system (K20-K92)										
White, male										
2002.....	32.4	12.0	0.7	0.3	0.3	0.4	0.9	1.2	4.0	9.0
2001.....	31.9	9.3	0.8	0.3	0.3	0.4	0.6	1.4	3.8	9.9
2000.....	31.7	11.2	0.6	0.3	0.3	0.4	1.0	1.6	3.9	9.8
1999.....	31.7	12.4	0.5	*	0.3	0.5	0.6	1.4	4.0	9.7
White, female										
2002.....	32.0	7.9	0.7	*	0.3	0.4	0.6	1.1	1.9	5.4
2001.....	32.1	8.0	0.5	0.3	0.3	0.3	0.5	1.0	2.3	5.9
2000.....	31.6	8.6	0.4	*	0.3	0.3	0.7	0.7	2.2	5.3
1999.....	31.2	7.5	0.8	0.3	0.4	0.4	0.6	1.0	2.5	5.5
Black, both sexes										
2002.....	23.6	25.9	1.4	*	*	0.6	1.4	2.5	4.2	7.9
2001.....	25.0	28.6	1.2	*	0.6	0.6	1.5	2.6	4.3	10.1
2000.....	25.2	29.4	1.5	*	*	*	1.6	2.9	4.6	10.8
1999.....	25.1	29.6	1.1	*	*	0.7	1.8	2.1	4.6	10.1
Black, male										
2002.....	25.9	31.4	1.7	*	*	*	*	2.9	4.8	9.0
2001.....	28.3	33.5	*	*	*	*	*	3.2	5.3	13.2
2000.....	27.7	33.9	2.0	*	*	*	2.1	3.0	5.1	13.6
1999.....	27.7	36.4	*	*	*	*	1.6	2.2	5.7	12.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of other parts of digestive system (K20-K92)										
Black, female										
2002.....	21.5	20.3	*	*	*	*	1.7	2.1	3.7	7.0
2001.....	22.0	23.3	*	*	*	*	1.7	2.0	3.4	7.3
2000.....	22.9	24.8	*	*	*	*	*	2.8	4.3	8.4
1999.....	22.6	22.5	*	*	*	*	1.9	2.0	3.6	8.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)										
All races 3/, both sexes										
2002.....	5.0	5.8	0.1	*	*	*	0.1	0.1	0.1	0.3
2001.....	5.0	5.8	0.1	*	*	*	*	0.1	0.1	0.2
2000.....	5.1	6.0	0.1	*	*	*	0.1	0.1	0.1	0.3
1999.....	5.0	6.1	0.1	*	*	*	*	0.1	0.2	0.2
All races 3/, male										
2002.....	3.6	6.5	*	*	*	*	*	*	0.2	0.3
2001.....	3.6	6.5	*	*	*	*	*	*	*	0.2
2000.....	3.6	6.3	*	*	*	*	*	*	*	0.3
1999.....	3.5	7.6	*	*	*	*	*	*	*	0.2
All races 3/, female										
2002.....	6.4	5.0	*	*	*	*	*	*	*	0.2
2001.....	6.4	5.1	*	*	*	*	*	*	*	0.2
2000.....	6.5	5.8	*	*	*	*	*	*	*	0.2
1999.....	6.4	4.4	*	*	*	*	*	*	0.2	0.3
White, both sexes										
2002.....	5.4	4.2	*	*	*	*	*	*	0.1	0.2
2001.....	5.4	4.1	*	*	*	*	*	*	*	0.2
2000.....	5.4	3.8	*	*	*	*	0.1	*	*	0.2
1999.....	5.4	4.2	*	*	*	*	*	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of intestines and peritoneum (K52,K58-K63)

All races 3/, both sexes

2002.....	1.7	2.0	0.2	*	*	*	*	*	*	0.2
2001.....	1.7	1.4	0.2	*	*	*	*	*	*	0.2
2000.....	1.6	1.7	*	*	*	*	*	*	0.1	0.1
1999.....	1.6	1.5	0.2	*	*	*	*	*	*	0.2

All races 3/, male

2002.....	1.3	2.4	*	*	*	*	*	*	*	*
2001.....	1.2	1.6	*	*	*	*	*	*	*	*
2000.....	1.2	1.8	*	*	*	*	*	*	*	*
1999.....	1.2	2.0	*	*	*	*	*	*	*	0.2

All races 3/, female

2002.....	2.0	1.5	*	*	*	*	*	*	*	0.2
2001.....	2.1	1.1	*	*	*	*	*	*	*	0.2
2000.....	2.0	1.6	*	*	*	*	*	*	*	*
1999.....	1.9	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	1.8	1.6	0.2	*	*	*	*	*	*	0.2
2001.....	1.8	0.8	*	*	*	*	*	*	*	0.1
2000.....	1.7	1.4	*	*	*	*	*	*	*	0.1
1999.....	1.7	1.2	0.2	*	*	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of liver (K70-K76)										
All races 3/, both sexes										
2002.....	12.4	2.2	0.2	*	*	0.1	0.3	0.7	1.9	5.3
2001.....	12.2	1.9	0.2	*	*	0.1	0.3	0.6	2.0	6.1
2000.....	12.0	2.0	0.2	*	*	*	0.4	0.6	2.1	6.0
1999.....	12.0	2.3	0.1	*	*	0.2	0.3	0.6	2.2	6.0
All races 3/, male										
2002.....	15.8	3.1	*	*	*	*	0.3	0.8	2.4	6.8
2001.....	15.7	2.3	*	*	*	*	0.3	0.7	2.5	7.8
2000.....	15.5	2.7	*	*	*	*	0.4	0.8	2.6	7.8
1999.....	15.6	2.9	*	*	*	*	*	0.6	2.8	7.6
All races 3/, female										
2002.....	9.0	1.3	*	*	*	*	0.3	0.6	1.4	3.9
2001.....	8.9	1.6	*	*	*	*	0.3	0.5	1.5	4.4
2000.....	8.7	1.3	*	*	*	*	0.3	0.4	1.6	4.2
1999.....	8.6	1.6	*	*	*	*	0.3	0.5	1.6	4.3
White, both sexes										
2002.....	13.1	1.9	*	*	*	*	0.2	0.6	1.9	5.4
2001.....	12.9	1.6	0.2	*	*	*	0.2	0.5	2.0	6.1
2000.....	12.6	1.6	*	*	*	*	0.3	0.5	2.0	5.8
1999.....	12.5	1.7	*	*	*	0.1	0.2	0.5	2.1	5.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of liver (K70-K76)										
White, male										
2002.....	16.8	2.7	*	*	*	*	0.3	0.7	2.4	6.9
2001.....	16.4	1.9	*	*	*	*	*	0.6	2.5	7.7
2000.....	16.2	2.0	*	*	*	*	0.3	0.7	2.6	7.7
1999.....	16.3	2.0	*	*	*	*	*	0.7	2.8	7.6
White, female										
2002.....	9.5	*	*	*	*	*	*	0.5	1.3	3.9
2001.....	9.4	1.3	*	*	*	*	*	0.4	1.5	4.5
2000.....	9.1	*	*	*	*	*	*	0.3	1.4	3.9
1999.....	8.9	1.4	*	*	*	*	*	0.4	1.5	4.1
Black, both sexes										
2002.....	9.9	3.7	*	*	*	*	*	1.0	1.9	4.5
2001.....	10.4	3.4	*	*	*	*	*	0.9	1.9	5.8
2000.....	10.4	4.2	*	*	*	*	0.7	1.2	2.3	6.9
1999.....	10.8	5.0	*	*	*	*	*	0.9	2.5	6.3
Black, male										
2002.....	12.6	*	*	*	*	*	*	*	2.2	4.8
2001.....	14.0	*	*	*	*	*	*	*	2.4	7.8
2000.....	13.5	6.4	*	*	*	*	*	*	2.4	8.9
1999.....	14.1	7.2	*	*	*	*	*	*	2.7	7.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of liver (K70-K76)										
Black, female										
2002.....	7.4	*	*	*	*	*	*	*	1.7	4.2
2001.....	7.2	*	*	*	*	*	*	*	1.5	4.0
2000.....	7.7	*	*	*	*	*	*	*	2.3	5.1
1999.....	7.9	*	*	*	*	*	*	*	2.3	5.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcoholic liver disease (K70)										
All races 3/, both sexes										
2002.....	4.2	*	*	*	*	*	*	0.2	0.9	2.8
2001.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3
2000.....	4.3	*	*	*	*	*	*	0.3	1.1	3.3
1999.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3
All races 3/, male										
2002.....	6.2	*	*	*	*	*	*	0.3	1.2	3.6
2001.....	6.4	*	*	*	*	*	*	0.3	1.4	4.4
2000.....	6.5	*	*	*	*	*	*	0.5	1.5	4.3
1999.....	6.5	*	*	*	*	*	*	0.2	1.4	4.3
All races 3/, female										
2002.....	2.3	*	*	*	*	*	*	*	0.7	2.0
2001.....	2.3	*	*	*	*	*	*	*	0.7	2.3
2000.....	2.2	*	*	*	*	*	*	*	0.6	2.2
1999.....	2.1	*	*	*	*	*	*	*	0.8	2.4
White, both sexes										
2002.....	4.4	*	*	*	*	*	*	0.2	0.9	2.8
2001.....	4.5	*	*	*	*	*	*	0.2	1.1	3.4
2000.....	4.5	*	*	*	*	*	*	0.3	1.1	3.2
1999.....	4.4	*	*	*	*	*	*	0.2	1.0	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Fibrosis and cirrhosis of liver (K74)										
All races 3/, both sexes										
2002.....	5.1	*	*	*	*	*	*	0.1	0.4	1.4
2001.....	5.0	*	*	*	*	*	*	0.1	0.4	1.6
2000.....	5.0	*	*	*	*	*	*	0.1	0.5	1.5
1999.....	5.0	*	*	*	*	*	*	0.1	0.5	1.4
All races 3/, male										
2002.....	6.0	*	*	*	*	*	*	*	0.5	1.9
2001.....	6.0	*	*	*	*	*	*	*	0.5	1.9
2000.....	5.9	*	*	*	*	*	*	*	0.6	2.0
1999.....	5.9	*	*	*	*	*	*	*	0.7	2.0
All races 3/, female										
2002.....	4.2	*	*	*	*	*	*	*	0.3	0.9
2001.....	4.2	*	*	*	*	*	*	*	0.3	1.2
2000.....	4.1	*	*	*	*	*	*	*	0.4	0.9
1999.....	4.1	*	*	*	*	*	*	*	0.4	0.9
White, both sexes										
2002.....	5.5	*	*	*	*	*	*	*	0.4	1.4
2001.....	5.4	*	*	*	*	*	*	*	0.4	1.6
2000.....	5.4	*	*	*	*	*	*	*	0.4	1.5
1999.....	5.3	*	*	*	*	*	*	*	0.6	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of liver (K71-K72,K75-K76)										
All races 3/, both sexes										
2002.....	2.9	2.1	0.2	*	*	*	0.2	0.3	0.5	1.1
2001.....	2.7	1.9	0.2	*	*	*	0.2	0.2	0.5	1.1
2000.....	2.6	1.9	0.2	*	*	*	0.2	0.2	0.5	1.2
1999.....	2.6	2.1	0.1	*	*	0.1	0.2	0.2	0.6	1.2
All races 3/, male										
2002.....	3.5	2.8	*	*	*	*	*	0.3	0.6	1.2
2001.....	3.2	2.2	*	*	*	*	*	*	0.5	1.4
2000.....	3.0	2.5	*	*	*	*	0.2	*	0.5	1.5
1999.....	3.1	2.7	*	*	*	*	*	0.2	0.7	1.4
All races 3/, female										
2002.....	2.3	1.3	*	*	*	*	*	0.3	0.4	1.0
2001.....	2.3	1.5	*	*	*	*	*	0.2	0.5	0.9
2000.....	2.2	1.3	*	*	*	*	*	*	0.5	0.9
1999.....	2.2	1.5	*	*	*	*	0.2	*	0.4	0.9
White, both sexes										
2002.....	3.0	1.8	*	*	*	*	*	0.3	0.5	1.1
2001.....	2.8	1.5	0.2	*	*	*	0.1	0.2	0.4	1.0
2000.....	2.6	1.6	*	*	*	*	0.2	0.1	0.5	1.1
1999.....	2.6	1.5	*	*	*	*	0.1	0.2	0.5	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Disorders of biliary tract and pancreas (K83-K86)										
All races 3/, both sexes										
2002.....	1.7	*	*	*	*	*	0.2	0.2	0.4	0.6
2001.....	1.7	*	*	*	*	*	0.1	0.2	0.5	0.7
2000.....	1.7	*	*	*	*	*	0.2	0.3	0.4	0.7
1999.....	1.6	*	*	*	*	*	0.2	0.3	0.4	0.6
All races 3/, male										
2002.....	1.9	*	*	*	*	*	0.2	0.3	0.6	0.8
2001.....	1.8	*	*	*	*	*	*	0.3	0.6	1.0
2000.....	1.8	*	*	*	*	*	0.2	0.4	0.6	0.9
1999.....	1.7	*	*	*	*	*	*	0.4	0.6	0.8
All races 3/, female										
2002.....	1.5	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.7	*	*	*	*	*	*	*	0.3	0.5
2000.....	1.6	*	*	*	*	*	*	*	0.2	0.4
1999.....	1.5	*	*	*	*	*	*	*	0.3	0.4
White, both sexes										
2002.....	1.8	*	*	*	*	*	0.1	0.2	0.4	0.6
2001.....	1.8	*	*	*	*	*	*	0.2	0.4	0.6
2000.....	1.7	*	*	*	*	*	0.2	0.2	0.4	0.6
1999.....	1.6	*	*	*	*	*	*	0.2	0.4	0.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pancreatitis (K85,K86.0-K86.1)										
All races 3/, both sexes										
2002.....	1.2	*	*	*	*	*	0.2	0.2	0.4	0.6
2001.....	1.3	*	*	*	*	*	0.1	0.2	0.4	0.6
2000.....	1.2	*	*	*	*	*	0.2	0.2	0.4	0.6
1999.....	1.2	*	*	*	*	*	0.1	0.2	0.4	0.5
All races 3/, male										
2002.....	1.4	*	*	*	*	*	0.2	0.3	0.6	0.8
2001.....	1.4	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	1.3	*	*	*	*	*	0.2	0.3	0.6	0.9
1999.....	1.3	*	*	*	*	*	*	0.3	0.6	0.7
All races 3/, female										
2002.....	1.1	*	*	*	*	*	*	*	*	0.4
2001.....	1.2	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.1	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.1	*	*	*	*	*	*	*	0.3	0.3
White, both sexes										
2002.....	1.3	*	*	*	*	*	0.1	0.2	0.3	0.5
2001.....	1.3	*	*	*	*	*	*	0.2	0.3	0.6
2000.....	1.2	*	*	*	*	*	0.2	0.2	0.3	0.6
1999.....	1.2	*	*	*	*	*	*	0.2	0.4	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)										
All races 3/, both sexes										
2002.....	5.0	*	*	*	0.1	0.2	0.5	0.7	0.7	1.2
2001.....	4.9	*	*	*	0.1	0.3	0.6	0.6	0.8	1.3
2000.....	4.9	*	*	*	0.1	0.2	0.6	0.7	0.8	1.1
1999.....	4.8	*	*	*	0.1	0.3	0.5	0.6	0.8	1.2
All races 3/, male										
2002.....	3.0	*	*	*	*	*	0.2	0.4	0.5	0.7
2001.....	2.9	*	*	*	*	*	0.2	0.3	0.5	1.0
2000.....	2.9	*	*	*	*	*	0.4	0.4	0.5	0.6
1999.....	2.8	*	*	*	*	*	0.2	0.3	0.5	0.8
All races 3/, female										
2002.....	7.0	*	*	*	*	0.4	0.8	1.0	1.0	1.6
2001.....	6.9	*	*	*	*	0.5	0.9	1.0	1.2	1.7
2000.....	6.8	*	*	*	*	0.3	0.8	1.1	1.2	1.6
1999.....	6.7	*	*	*	0.2	0.4	0.7	1.0	1.2	1.7
White, both sexes										
2002.....	5.2	*	*	*	*	0.2	0.3	0.4	0.4	0.8
2001.....	5.1	*	*	*	*	0.3	0.4	0.3	0.6	1.0
2000.....	5.1	*	*	*	*	*	0.3	0.5	0.6	0.7
1999.....	5.0	*	*	*	*	0.2	0.3	0.4	0.6	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

Black, female

2002.....	6.3	*	*	*	*	*	3.0	3.4	3.4	5.6
2001.....	6.3	*	*	*	*	*	2.7	3.6	3.6	4.8
2000.....	6.2	*	*	*	*	*	2.5	3.2	3.4	4.9
1999.....	6.2	*	*	*	*	1.4	2.4	3.1	3.4	5.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Systemic lupus erythematosus (M32)										
All races 3/, both sexes										
2002.....	0.4	*	*	*	*	0.1	0.3	0.3	0.3	0.5
2001.....	0.5	*	*	*	*	0.2	0.4	0.4	0.4	0.5
2000.....	0.5	*	*	*	*	0.1	0.3	0.4	0.3	0.5
1999.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.5
All races 3/, male										
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	0.8	*	*	*	*	0.2	0.5	0.5	0.5	0.9
2001.....	0.8	*	*	*	*	0.3	0.6	0.7	0.6	0.8
2000.....	0.8	*	*	*	*	*	0.5	0.6	0.6	0.9
1999.....	0.8	*	*	*	*	0.2	0.5	0.6	0.6	0.9
White, both sexes										
2002.....	0.3	*	*	*	*	*	*	*	0.1	0.2
2001.....	0.4	*	*	*	*	*	0.2	0.1	0.2	0.2
2000.....	0.4	*	*	*	*	*	0.2	0.2	0.2	0.3
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Systemic lupus erythematosus (M32)										
Black, female										
2002.....	2.1	*	*	*	*	*	2.1	2.3	2.0	3.8
2001.....	2.0	*	*	*	*	*	2.0	2.7	2.2	3.2
2000.....	2.0	*	*	*	*	*	1.8	2.5	2.0	2.8
1999.....	2.0	*	*	*	*	*	1.9	1.9	2.1	3.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)

All races 3/, both sexes

2002.....	1.7	*	*	*	*	*	0.1	0.2	0.3	0.5
2001.....	1.7	*	*	*	*	*	0.1	0.2	0.3	0.5
2000.....	1.7	*	*	*	*	*	0.2	0.2	0.4	0.4
1999.....	1.7	*	*	*	*	0.1	*	0.2	0.3	0.5

All races 3/, male

2002.....	1.0	*	*	*	*	*	*	*	0.2	0.4
2001.....	1.0	*	*	*	*	*	*	*	*	0.4
2000.....	1.1	*	*	*	*	*	*	0.2	0.3	0.2
1999.....	1.0	*	*	*	*	*	*	*	*	0.4

All races 3/, female

2002.....	2.3	*	*	*	*	*	0.2	0.3	0.4	0.5
2001.....	2.3	*	*	*	*	*	*	0.3	0.4	0.6
2000.....	2.3	*	*	*	*	*	*	0.3	0.5	0.6
1999.....	2.3	*	*	*	*	*	*	0.2	0.4	0.6

White, both sexes

2002.....	1.7	*	*	*	*	*	*	0.2	0.2	0.3
2001.....	1.8	*	*	*	*	*	*	0.1	0.2	0.4
2000.....	1.7	*	*	*	*	*	*	0.2	0.3	0.2
1999.....	1.7	*	*	*	*	*	*	*	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XIV. Diseases of the genitourinary system (N00-N98)										
All races 3/, both sexes										
2002.....	20.2	4.6	*	*	*	0.2	0.4	0.7	1.0	1.6
2001.....	20.0	3.9	0.2	*	*	0.1	0.4	0.6	0.9	1.7
2000.....	19.4	5.0	0.1	*	*	0.2	0.4	0.5	1.0	1.5
1999.....	19.0	5.1	0.2	0.1	0.1	0.2	0.3	0.7	0.9	1.5
All races 3/, male										
2002.....	18.1	5.6	*	*	*	*	0.4	0.6	1.0	1.6
2001.....	17.7	4.2	*	*	*	*	0.4	0.6	0.9	1.8
2000.....	17.2	6.1	*	*	*	0.2	0.3	0.5	1.0	1.5
1999.....	16.9	5.0	*	*	*	*	0.3	0.6	0.9	1.6
All races 3/, female										
2002.....	22.3	3.6	*	*	*	*	0.4	0.9	1.0	1.6
2001.....	22.2	3.6	*	*	*	*	0.5	0.6	1.0	1.6
2000.....	21.5	3.9	*	*	*	*	0.5	0.5	1.0	1.5
1999.....	21.1	5.1	*	*	*	0.2	0.4	0.8	0.9	1.4
White, both sexes										
2002.....	20.3	4.0	*	*	*	*	0.3	0.5	0.8	1.1
2001.....	20.0	3.0	*	*	*	*	0.4	0.4	0.7	1.2
2000.....	19.4	3.6	*	*	*	0.2	0.4	0.4	0.7	1.0
1999.....	18.9	4.0	*	*	*	0.1	0.2	0.5	0.7	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XIV. Diseases of the genitourinary system (N00-N98)										
White, male										
2002.....	18.1	4.7	*	*	*	*	0.3	0.4	0.8	1.1
2001.....	17.7	3.4	*	*	*	*	0.3	0.4	0.6	1.5
2000.....	17.2	4.6	*	*	*	*	0.3	0.4	0.7	1.0
1999.....	16.6	4.5	*	*	*	*	*	0.4	0.7	1.1
White, female										
2002.....	22.5	3.2	*	*	*	*	0.3	0.6	0.7	1.1
2001.....	22.3	2.6	*	*	*	*	0.5	0.5	0.8	1.0
2000.....	21.5	2.6	*	*	*	*	0.4	0.5	0.7	1.0
1999.....	21.2	3.4	*	*	*	*	0.3	0.6	0.6	1.0
Black, both sexes										
2002.....	25.2	8.3	*	*	*	*	1.1	2.3	2.6	4.8
2001.....	25.1	9.2	*	*	*	*	0.8	1.6	2.6	4.7
2000.....	24.4	12.7	*	*	*	*	*	1.2	2.9	4.9
1999.....	24.3	11.1	*	*	*	*	1.0	1.8	2.5	4.6
Black, male										
2002.....	23.2	10.5	*	*	*	*	*	1.9	2.4	4.6
2001.....	22.2	9.3	*	*	*	*	*	2.1	3.1	3.8
2000.....	21.9	14.1	*	*	*	*	*	*	3.1	5.6
1999.....	22.4	8.1	*	*	*	*	*	*	2.4	5.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XIV. Diseases of the genitourinary system (N00-N98)										
Black, female										
2002.....	27.0	6.1	*	*	*	*	1.3	2.7	2.8	5.0
2001.....	27.8	9.1	*	*	*	*	*	*	2.1	5.5
2000.....	26.6	11.2	*	*	*	*	*	*	2.8	4.3
1999.....	26.1	14.1	*	*	*	*	*	2.2	2.5	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of urinary system (N00-N39)										
All races 3/, both sexes										
2002.....	19.9	4.6	*	*	*	0.1	0.4	0.7	0.9	1.5
2001.....	19.7	3.7	0.2	*	*	0.1	0.4	0.6	0.9	1.7
2000.....	19.1	4.8	0.1	*	*	0.2	0.4	0.5	0.9	1.4
1999.....	18.7	4.7	0.1	0.1	0.1	0.2	0.3	0.7	0.9	1.4
All races 3/, male										
2002.....	17.7	5.6	*	*	*	*	0.4	0.6	1.0	1.5
2001.....	17.3	4.2	*	*	*	*	0.4	0.6	0.9	1.7
2000.....	16.8	6.0	*	*	*	0.2	0.3	0.5	1.0	1.5
1999.....	16.4	5.0	*	*	*	*	0.3	0.6	0.9	1.6
All races 3/, female										
2002.....	22.0	3.5	*	*	*	*	0.4	0.9	0.9	1.4
2001.....	22.0	3.2	*	*	*	*	0.5	0.6	1.0	1.6
2000.....	21.2	3.6	*	*	*	*	0.4	0.5	0.9	1.3
1999.....	20.9	4.3	*	*	*	*	0.3	0.7	0.8	1.3
White, both sexes										
2002.....	20.0	3.9	*	*	*	*	0.3	0.5	0.7	1.0
2001.....	19.7	2.8	*	*	*	*	0.4	0.4	0.7	1.2
2000.....	19.0	3.5	*	*	*	0.2	0.4	0.4	0.7	1.0
1999.....	18.6	3.7	*	*	*	0.1	0.2	0.5	0.6	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of urinary system (N00-N39)										
White, male										
2002.....	17.7	4.7	*	*	*	*	0.3	0.4	0.8	1.1
2001.....	17.3	3.3	*	*	*	*	0.3	0.4	0.6	1.5
2000.....	16.7	4.5	*	*	*	*	0.3	0.4	0.7	1.0
1999.....	16.2	4.5	*	*	*	*	*	0.4	0.7	1.1
White, female										
2002.....	22.2	3.0	*	*	*	*	*	0.6	0.6	1.0
2001.....	22.0	2.3	*	*	*	*	0.4	0.5	0.8	1.0
2000.....	21.3	2.4	*	*	*	*	0.4	0.4	0.7	0.9
1999.....	20.9	2.8	*	*	*	*	*	0.5	0.6	0.9
Black, both sexes										
2002.....	24.8	8.3	*	*	*	*	1.0	2.3	2.5	4.5
2001.....	24.8	8.9	*	*	*	*	0.8	1.6	2.6	4.4
2000.....	24.1	12.2	*	*	*	*	*	1.2	2.8	4.7
1999.....	23.9	9.9	*	*	*	*	0.9	1.7	2.4	4.5
Black, male										
2002.....	22.8	10.5	*	*	*	*	*	1.9	2.4	4.5
2001.....	21.8	9.3	*	*	*	*	*	2.1	3.1	3.6
2000.....	21.6	14.1	*	*	*	*	*	*	3.0	5.5
1999.....	22.0	8.1	*	*	*	*	*	*	2.4	5.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of urinary system (N00-N39)										
Black, female										
2002.....	26.6	6.1	*	*	*	*	*	2.6	2.6	4.4
2001.....	27.5	8.5	*	*	*	*	*	*	2.1	5.2
2000.....	26.3	10.3	*	*	*	*	*	*	2.6	4.0
1999.....	25.7	11.7	*	*	*	*	*	2.0	2.4	3.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
All races 3/, both sexes										
2002.....	13.9	4.1	*	*	*	*	0.3	0.5	0.7	1.2
2001.....	13.6	3.1	*	*	*	*	0.3	0.4	0.6	1.2
2000.....	13.0	4.0	*	*	*	0.1	0.3	0.4	0.7	1.2
1999.....	12.4	4.3	*	*	*	0.1	0.2	0.4	0.7	1.1
All races 3/, male										
2002.....	13.7	5.1	*	*	*	*	0.3	0.4	0.7	1.3
2001.....	13.2	3.5	*	*	*	*	0.3	0.4	0.6	1.3
2000.....	12.6	5.1	*	*	*	*	0.2	0.4	0.7	1.3
1999.....	12.1	4.5	*	*	*	*	*	0.3	0.7	1.3
All races 3/, female										
2002.....	14.2	3.0	*	*	*	*	0.3	0.6	0.7	1.1
2001.....	14.0	2.7	*	*	*	*	0.4	0.4	0.7	1.2
2000.....	13.3	2.8	*	*	*	*	0.3	0.3	0.7	1.0
1999.....	12.7	4.1	*	*	*	*	0.3	0.5	0.7	0.9
White, both sexes										
2002.....	13.6	3.4	*	*	*	*	0.2	0.3	0.5	0.8
2001.....	13.2	2.2	*	*	*	*	0.3	0.3	0.4	0.8
2000.....	12.6	2.7	*	*	*	*	0.2	0.3	0.5	0.7
1999.....	12.0	3.3	*	*	*	*	0.2	0.3	0.5	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
White, male										
2002.....	13.4	4.2	*	*	*	*	*	*	0.5	0.9
2001.....	13.0	2.6	*	*	*	*	*	0.3	0.4	1.0
2000.....	12.4	3.6	*	*	*	*	0.3	0.3	0.5	0.8
1999.....	11.8	3.9	*	*	*	*	*	*	0.5	0.8
White, female										
2002.....	13.8	2.6	*	*	*	*	*	0.4	0.5	0.7
2001.....	13.4	1.8	*	*	*	*	0.3	0.3	0.5	0.7
2000.....	12.8	1.7	*	*	*	*	*	*	0.5	0.7
1999.....	12.2	2.8	*	*	*	*	*	0.3	0.5	0.6
Black, both sexes										
2002.....	19.5	7.7	*	*	*	*	0.8	1.7	2.0	3.9
2001.....	19.3	8.1	*	*	*	*	0.7	1.2	2.1	3.9
2000.....	18.5	11.1	*	*	*	*	*	1.0	2.5	4.1
1999.....	18.2	9.6	*	*	*	*	*	1.4	2.2	3.8
Black, male										
2002.....	18.8	9.6	*	*	*	*	*	*	2.0	4.4
2001.....	17.7	8.7	*	*	*	*	*	1.6	2.6	3.3
2000.....	17.3	13.4	*	*	*	*	*	*	2.7	4.9
1999.....	17.1	8.1	*	*	*	*	*	*	2.2	4.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
Black, female										
2002.....	20.2	*	*	*	*	*	*	1.9	2.0	3.5
2001.....	20.7	7.6	*	*	*	*	*	*	1.7	4.4
2000.....	19.6	8.6	*	*	*	*	*	*	2.3	3.4
1999.....	19.1	11.1	*	*	*	*	*	1.6	2.2	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XV. Pregnancy, childbirth and the puerperium (000-099)										
All races 3/, both sexes										
2002.....	0.1	*	0.1	0.3	0.4	0.4	0.4
2001.....	0.1	*	0.2	0.4	0.5	0.5	0.3
2000.....	0.1	*	*	0.4	0.4	0.5	0.4
1999.....	0.1	*	0.2	0.3	0.5	0.4	0.4
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	0.3	*	0.3	0.6	0.9	0.9	0.7
2001.....	0.3	*	0.4	0.8	1.0	0.9	0.7
2000.....	0.3	*	*	0.8	0.9	0.9	0.8
1999.....	0.3	*	0.3	0.7	0.9	0.9	0.8
White, both sexes										
2002.....	0.1	*	*	0.2	0.3	0.3	0.2
2001.....	0.1	*	0.1	0.3	0.3	0.4	0.2
2000.....	0.1	*	*	0.3	0.3	0.3	0.3
1999.....	0.1	*	*	0.2	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XV. Pregnancy, childbirth and the puerperium (000-099)										
Black, female										
2002.....	0.8	*	*	2.0	2.8	1.9	2.3
2001.....	0.8	*	*	2.1	2.6	2.1	1.9
2000.....	0.7	*	*	2.2	2.2	2.3	1.7
1999.....	0.9	*	*	2.1	2.7	2.3	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

All races 3/, both sexes

2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

Black, female

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ectopic pregnancy (000)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
White, both sexes										
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ectopic pregnancy (000)										
Black, female										
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Spontaneous abortion (003)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Spontaneous abortion (003)

Black, female

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Medical abortion (004)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other abortion (005)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other abortion (005)										
Black, female										
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

Black, female

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other direct obstetric deaths (O10-O92)										
All races 3/, both sexes										
2002.....	0.1	*	0.1	0.2	0.3	0.3	0.3
2001.....	0.1	*	0.1	0.3	0.4	0.4	0.2
2000.....	0.1	*	*	0.3	0.3	0.4	0.3
1999.....	0.1	*	0.1	0.2	0.3	0.3	0.3
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	0.2	*	0.2	0.4	0.7	0.7	0.6
2001.....	0.2	*	0.3	0.5	0.8	0.7	0.5
2000.....	0.2	*	*	0.6	0.6	0.8	0.6
1999.....	0.2	*	0.3	0.5	0.7	0.7	0.6
White, both sexes										
2002.....	0.1	*	*	*	0.2	0.3	0.2
2001.....	0.1	*	*	0.2	0.3	0.3	0.1
2000.....	0.1	*	*	0.2	0.2	0.3	0.2
1999.....	0.1	*	*	0.2	0.2	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other direct obstetric deaths (O10-O92)										
Black, female										
2002.....	0.6	*	*	1.5	1.9	1.6	1.7
2001.....	0.6	*	*	1.5	2.0	1.6	*
2000.....	0.5	*	*	1.6	1.5	1.7	*
1999.....	0.6	*	*	*	2.0	1.7	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Eclampsia and pre-eclampsia (O11,O13-O16)										
All races 3/, both sexes										
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	0.1	*
1999.....	0.0	*	*	*	0.1	*	*
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	*
1999.....	0.1	*	*	*	0.2	*	*
White, both sexes										
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,072)

All races 3/, both sexes

2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Complications predominately related to the puerperium (085-092)

All races 3/, both sexes

2002.....	0.0	*	*	*	*	0.1	*
2001.....	0.0	*	*	*	0.1	0.1	*
2000.....	0.0	*	*	0.1	*	0.1	*
1999.....	0.0	*	*	*	*	0.1	*

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	0.1	*	*	*	*	0.2	*
2001.....	0.1	*	*	*	0.3	0.2	*
2000.....	0.1	*	*	0.2	*	0.3	*
1999.....	0.1	*	*	*	*	0.2	*

White, both sexes

2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Complications predominately related to the puerperium (085-092)

Black, female

2002.....	0.2	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Obstetric embolism (O88)										
All races 3/, both sexes										
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	0.1	*
1999.....	0.0	*	*	*	*	*	*
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	0.2	*
1999.....	0.0	*	*	*	*	*	*
White, both sexes										
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications predominately related to the puerperium (O85-O87,O89-O92)

All races 3/, both sexes

2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Black, female

2002.....	0.1	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)

All races 3/, both sexes

2002.....	0.0	*	*	0.1	0.1	0.1	0.1
2001.....	0.0	*	*	*	0.1	0.1	*
2000.....	0.0	*	*	0.1	*	0.1	0.1
1999.....	0.0	*	*	*	0.1	*	*

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	0.1	*	*	0.2	0.3	0.3	0.3
2001.....	0.1	*	*	*	0.3	0.2	*
2000.....	0.1	*	*	0.2	*	0.2	0.2
1999.....	0.1	*	*	*	0.2	*	*

White, both sexes

2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Obstetric death of unspecified cause (095)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Obstetric death of unspecified cause (095)

Black, female

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

All races 3/, both sexes

2002.....	0.0	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	0.0	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Black, female

2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Indirect obstetric deaths (098-099)										
All races 3/, both sexes										
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*
White, both sexes										
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Indirect obstetric deaths (098-099)

Black, female

2002.....	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVI. Certain conditions originating in the perinatal period (P00-P96)

All races 3/, both sexes

2002.....	4.9	349.7	0.4	*	*	*	*	*	*	*
2001.....	4.9	340.5	0.5	0.1	*	*	*	*	*	*
2000.....	5.0	365.7	0.5	0.1	*	*	*	*	*	*
1999.....	5.1	371.0	0.6	0.1	*	*	*	*	*	*

All races 3/, male

2002.....	5.7	385.7	0.5	*	*	*	*	*	*	*
2001.....	5.6	376.6	0.6	*	*	*	*	*	*	*
2000.....	5.8	403.9	0.7	*	*	*	*	*	*	*
1999.....	5.9	408.6	0.7	*	*	*	*	*	*	*

All races 3/, female

2002.....	4.2	312.0	0.3	*	*	*	*	*	*	*
2001.....	4.2	302.6	0.4	*	*	*	*	*	*	*
2000.....	4.3	325.6	0.4	*	*	*	*	*	*	*
1999.....	4.4	331.7	0.5	*	*	*	*	*	*	*

White, both sexes

2002.....	3.7	277.2	0.3	*	*	*	*	*	*	*
2001.....	3.7	268.5	0.4	*	*	*	*	*	*	*
2000.....	3.7	283.7	0.4	*	*	*	*	*	*	*
1999.....	3.8	285.9	0.5	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
All races 3/, both sexes										
2002.....	3.7	139.4	3.4	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001.....	3.7	136.7	3.6	0.9	0.9	1.3	1.3	1.2	1.1	1.2
2000.....	3.8	150.9	3.2	1.0	1.0	1.1	1.1	1.1	1.3	1.2
1999.....	3.7	144.2	3.6	1.0	1.1	1.1	1.1	1.1	1.2	1.2
All races 3/, male										
2002.....	3.9	142.6	3.5	1.0	1.1	1.4	1.3	1.4	1.4	1.2
2001.....	3.9	143.1	3.6	1.0	1.0	1.5	1.4	1.3	1.2	1.2
2000.....	4.0	155.4	3.3	1.0	1.0	1.3	1.4	1.2	1.5	1.4
1999.....	3.9	147.9	4.0	1.0	1.1	1.3	1.3	1.1	1.3	1.3
All races 3/, female										
2002.....	3.5	136.1	3.3	1.0	1.0	1.0	1.1	0.9	1.1	1.2
2001.....	3.4	129.9	3.7	0.8	0.8	1.0	1.1	1.1	1.0	1.1
2000.....	3.5	146.2	3.1	0.9	1.0	0.9	0.9	0.9	1.1	1.1
1999.....	3.5	140.3	3.2	1.0	1.1	0.9	1.0	1.1	1.2	1.1
White, both sexes										
2002.....	3.6	138.3	3.3	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001.....	3.6	135.0	3.2	0.9	0.9	1.2	1.2	1.2	1.2	1.1
2000.....	3.6	147.1	2.9	0.9	0.9	1.0	1.2	1.1	1.4	1.2
1999.....	3.6	141.3	3.2	0.9	1.0	1.1	1.1	1.1	1.3	1.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
White, male										
2002.....	3.8	143.3	3.4	0.9	1.0	1.3	1.3	1.4	1.5	1.2
2001.....	3.8	143.0	3.2	1.0	1.1	1.4	1.3	1.4	1.3	1.2
2000.....	3.9	151.9	2.9	0.9	0.9	1.1	1.4	1.2	1.6	1.4
1999.....	3.8	146.5	3.5	0.8	1.0	1.3	1.3	1.2	1.3	1.4
White, female										
2002.....	3.5	133.1	3.1	1.1	1.0	1.0	1.1	0.9	1.1	1.2
2001.....	3.4	126.7	3.3	0.8	0.7	0.9	1.1	1.0	1.1	1.0
2000.....	3.5	142.0	2.9	1.0	0.9	0.9	0.9	0.9	1.1	1.0
1999.....	3.5	135.7	2.9	1.0	1.0	0.8	0.9	1.1	1.2	1.1
Black, both sexes										
2002.....	4.7	155.2	4.5	1.2	1.2	1.7	1.5	1.3	1.4	1.4
2001.....	4.6	158.4	5.3	1.0	1.2	1.9	1.8	1.4	1.2	1.5
2000.....	4.9	179.2	5.0	1.3	1.4	1.8	1.3	1.4	1.4	1.6
1999.....	4.8	169.3	5.5	1.4	1.7	1.4	1.5	1.0	1.2	1.4
Black, male										
2002.....	5.1	151.7	4.6	1.4	1.4	2.1	1.7	1.7	1.6	*
2001.....	5.0	160.8	5.5	*	*	2.6	2.2	*	*	*
2000.....	5.4	181.4	5.5	1.6	1.4	2.5	*	1.9	1.7	1.8
1999.....	5.2	172.7	6.2	1.4	1.8	1.7	1.8	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Black, female

2002.....	4.3	158.9	4.4	*	*	1.3	1.3	*	*	1.4
2001.....	4.3	155.9	5.1	*	1.4	1.3	1.5	1.7	*	1.7
2000.....	4.4	176.9	4.6	*	1.5	*	*	*	*	1.5
1999.....	4.3	165.8	4.8	1.5	1.7	*	*	*	*	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)										
All races 3/, both sexes										
2002.....	0.3	9.3	0.8	0.3	0.2	0.2	0.1	*	*	0.1
2001.....	0.3	9.7	0.8	0.2	0.2	0.2	0.1	0.1	*	*
2000.....	0.3	10.7	0.6	0.3	0.2	0.1	0.1	*	*	*
1999.....	0.3	9.4	0.7	0.3	0.2	0.1	*	0.1	0.1	*
All races 3/, male										
2002.....	0.3	9.0	0.7	0.3	*	*	*	*	*	*
2001.....	0.3	9.4	0.8	0.2	0.2	*	*	*	*	*
2000.....	0.3	10.6	0.6	0.3	0.2	*	*	*	*	*
1999.....	0.3	8.6	0.7	0.3	0.2	*	*	*	*	*
All races 3/, female										
2002.....	0.3	9.6	0.9	0.3	0.3	*	*	*	*	*
2001.....	0.3	10.0	0.9	*	*	*	*	*	*	*
2000.....	0.3	10.9	0.7	0.3	0.2	*	*	*	*	*
1999.....	0.3	10.2	0.7	0.3	0.2	*	*	*	*	*
White, both sexes										
2002.....	0.3	9.4	0.7	0.3	0.2	0.2	0.2	*	*	*
2001.....	0.3	9.6	0.7	0.2	0.2	0.2	*	*	*	*
2000.....	0.3	10.5	0.6	0.3	0.2	*	*	*	*	*
1999.....	0.2	8.8	0.6	0.2	0.2	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Congenital malformations of heart (Q20-Q24)										
All races 3/, both sexes										
2002.....	1.1	37.4	1.2	0.3	0.4	0.6	0.5	0.5	0.6	0.5
2001.....	1.2	38.6	1.3	0.2	0.4	0.6	0.6	0.5	0.5	0.5
2000.....	1.2	42.9	1.2	0.3	0.4	0.6	0.5	0.5	0.7	0.5
1999.....	1.3	42.1	1.3	0.3	0.5	0.6	0.5	0.6	0.6	0.6
All races 3/, male										
2002.....	1.2	40.4	1.5	0.3	0.5	0.7	0.6	0.7	0.8	0.6
2001.....	1.3	41.8	1.4	0.3	0.4	0.9	0.7	0.6	0.7	0.6
2000.....	1.3	44.5	1.3	0.3	0.4	0.7	0.6	0.6	0.9	0.6
1999.....	1.4	45.7	1.5	0.3	0.6	0.8	0.7	0.6	0.7	0.7
All races 3/, female										
2002.....	1.0	34.4	1.0	0.3	0.3	0.4	0.3	0.4	0.5	0.5
2001.....	1.1	35.2	1.2	0.2	0.3	0.3	0.5	0.5	0.4	0.5
2000.....	1.1	41.3	1.2	0.3	0.3	0.4	0.4	0.4	0.5	0.4
1999.....	1.2	38.3	1.1	0.3	0.4	0.4	0.4	0.5	0.6	0.5
White, both sexes										
2002.....	1.1	36.4	1.3	0.3	0.4	0.6	0.4	0.5	0.7	0.6
2001.....	1.1	37.4	1.2	0.2	0.3	0.5	0.6	0.5	0.6	0.5
2000.....	1.2	41.5	1.1	0.3	0.3	0.5	0.5	0.5	0.7	0.5
1999.....	1.3	42.0	1.1	0.3	0.4	0.6	0.5	0.6	0.7	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Congenital malformations of heart (Q20-Q24)										
White, male										
2002.....	1.2	40.0	1.5	0.3	0.4	0.7	0.6	0.6	0.8	0.6
2001.....	1.2	41.3	1.3	0.3	0.4	0.7	0.6	0.6	0.7	0.6
2000.....	1.3	43.6	1.2	0.3	0.4	0.6	0.6	0.6	0.9	0.6
1999.....	1.3	46.2	1.3	0.3	0.5	0.8	0.6	0.6	0.7	0.7
White, female										
2002.....	1.0	32.7	1.0	0.3	0.3	0.4	0.3	0.4	0.6	0.5
2001.....	1.0	33.2	1.0	*	0.3	0.3	0.5	0.4	0.4	0.5
2000.....	1.1	39.3	1.1	0.3	0.3	0.4	0.4	0.4	0.6	0.4
1999.....	1.2	37.6	1.0	0.3	0.3	0.4	0.4	0.6	0.6	0.5
Black, both sexes										
2002.....	1.4	44.3	1.3	*	*	0.8	0.7	*	0.7	*
2001.....	1.6	47.7	1.9	*	*	1.2	1.0	*	*	*
2000.....	1.6	51.5	1.8	*	0.7	1.1	0.7	*	*	0.7
1999.....	1.6	47.9	2.2	*	0.8	0.7	0.9	*	*	0.8
Black, male										
2002.....	1.6	43.0	*	*	*	*	*	*	*	*
2001.....	1.7	48.8	1.7	*	*	1.9	*	*	*	*
2000.....	1.8	49.9	2.1	*	*	1.6	*	*	*	*
1999.....	1.8	50.1	2.5	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations of heart (Q20-Q24)

Black, female

2002.....	1.3	45.7	*	*	*	*	*	*	*	*
2001.....	1.4	46.6	2.0	*	*	*	*	*	*	*
2000.....	1.4	53.2	*	*	*	*	*	*	*	*
1999.....	1.4	45.6	2.0	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)										
All races 3/, both sexes										
2002.....	0.9	36.2	0.7	0.2	0.2	0.3	0.3	0.2	0.2	0.2
2001.....	0.9	36.3	0.8	0.2	0.2	0.2	0.2	0.2	0.2	0.3
2000.....	0.9	37.9	0.7	0.2	0.2	0.2	0.2	0.2	0.2	0.3
1999.....	0.9	38.3	0.9	0.2	0.1	0.1	0.2	0.2	0.2	0.2
All races 3/, male										
2002.....	1.0	39.2	0.6	0.2	0.2	0.3	0.3	0.2	*	0.2
2001.....	1.0	41.1	0.8	*	*	0.2	0.3	0.2	*	0.3
2000.....	1.0	42.1	0.7	0.2	*	0.2	0.2	0.2	0.3	0.3
1999.....	1.0	42.5	1.1	0.2	*	*	0.3	*	0.2	0.2
All races 3/, female										
2002.....	0.8	33.1	0.8	*	*	0.2	0.3	*	*	*
2001.....	0.8	31.3	0.8	*	*	*	*	0.2	*	0.3
2000.....	0.8	33.6	0.6	*	0.2	*	0.2	*	*	0.2
1999.....	0.8	33.8	0.8	*	*	*	*	*	*	0.2
White, both sexes										
2002.....	0.9	37.0	0.7	0.2	0.2	0.2	0.3	0.2	0.2	0.2
2001.....	0.9	36.0	0.7	0.2	0.1	0.2	0.2	0.2	0.2	0.3
2000.....	0.9	37.4	0.6	0.2	0.2	0.2	0.3	0.2	0.2	0.3
1999.....	0.9	36.8	0.9	0.2	0.1	0.2	0.2	0.2	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other congenital malformations and deformations (Q10-Q18,Q35-Q89)

White, male

2002.....	1.0	40.4	0.6	*	*	0.2	0.3	0.3	*	*
2001.....	1.0	40.6	0.6	*	*	0.3	0.3	*	*	0.3
2000.....	1.0	41.3	0.7	*	*	*	0.3	*	0.2	0.3
1999.....	1.0	42.1	1.0	*	*	*	0.3	*	*	0.2

White, female

2002.....	0.8	33.4	0.7	*	*	*	0.3	*	*	*
2001.....	0.8	31.1	0.7	*	*	*	*	*	*	0.3
2000.....	0.8	33.2	0.6	*	*	*	*	*	*	0.2
1999.....	0.8	31.2	0.7	*	*	*	*	*	*	0.2

Black, both sexes

2002.....	1.1	37.1	1.0	*	*	*	*	*	*	*
2001.....	1.2	41.9	0.9	*	*	*	*	*	*	*
2000.....	1.1	42.9	1.0	*	*	*	*	*	*	*
1999.....	1.2	46.3	1.4	*	*	*	*	*	*	*

Black, male

2002.....	1.3	40.1	*	*	*	*	*	*	*	*
2001.....	1.4	47.6	*	*	*	*	*	*	*	*
2000.....	1.4	48.0	*	*	*	*	*	*	*	*
1999.....	1.3	49.8	1.6	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)										
All races 3/, both sexes										
2002.....	0.4	24.3	0.2	*	*	*	*	*	*	*
2001.....	0.4	22.7	0.3	*	*	*	*	*	*	*
2000.....	0.4	24.2	0.3	*	*	*	*	*	*	*
1999.....	0.3	22.5	0.3	*	*	*	*	*	*	*
All races 3/, male										
2002.....	0.3	20.7	*	*	*	*	*	*	*	*
2001.....	0.3	20.1	*	*	*	*	*	*	*	*
2000.....	0.3	21.4	*	*	*	*	*	*	*	*
1999.....	0.3	18.7	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	0.4	28.0	*	*	*	*	*	*	*	*
2001.....	0.4	25.4	0.5	*	*	*	*	*	*	*
2000.....	0.4	27.2	0.3	*	*	*	*	*	*	*
1999.....	0.4	26.5	0.3	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.4	24.6	0.2	*	*	*	*	*	*	*
2001.....	0.3	22.9	0.3	*	*	*	*	*	*	*
2000.....	0.3	23.8	0.2	*	*	*	*	*	*	*
1999.....	0.3	22.7	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2002.....	10.4	85.7	1.6	0.3	0.4	1.3	2.4	3.0	3.6	4.6
2001.....	11.4	86.5	1.8	0.4	0.5	1.8	3.0	3.5	4.2	6.2
2000.....	11.3	95.6	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.7
1999.....	9.6	93.2	1.5	0.3	0.4	1.1	2.0	2.3	3.1	4.1

All races 3/, male

2002.....	10.1	98.5	1.8	0.4	0.4	1.6	3.2	4.1	4.9	6.1
2001.....	11.3	98.9	2.1	0.4	0.5	2.4	4.0	4.8	5.6	8.0
2000.....	11.5	110.6	1.7	0.5	0.5	1.9	4.0	4.7	5.9	7.6
1999.....	9.4	106.6	1.7	0.3	0.4	1.5	2.8	2.9	4.2	5.3

All races 3/, female

2002.....	10.7	72.3	1.4	0.2	0.3	0.8	1.6	1.8	2.3	3.1
2001.....	11.4	73.6	1.5	0.4	0.5	1.1	1.9	2.2	2.8	4.4
2000.....	11.2	79.9	1.5	0.4	0.4	1.2	1.8	2.2	2.8	3.7
1999.....	9.7	79.2	1.3	0.3	0.3	0.8	1.2	1.6	2.1	3.0

White, both sexes

2002.....	10.5	71.0	1.3	0.2	0.3	1.2	2.3	2.8	3.5	4.4
2001.....	11.6	70.9	1.5	0.3	0.4	1.8	3.0	3.5	4.1	6.0
2000.....	11.4	77.2	1.5	0.4	0.4	1.5	2.8	3.2	4.2	5.6
1999.....	9.4	74.7	1.3	0.3	0.3	1.0	1.8	2.0	2.9	3.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
White, male										
2002.....	9.8	82.5	1.4	0.3	0.4	1.6	3.2	3.9	4.7	5.8
2001.....	11.2	82.0	1.7	0.3	0.5	2.4	4.0	4.9	5.5	7.7
2000.....	11.3	89.4	1.7	0.4	0.4	1.9	3.9	4.4	5.7	7.7
1999.....	9.0	86.4	1.6	0.3	0.3	1.3	2.6	2.7	3.8	4.9
White, female										
2002.....	11.2	59.0	1.2	*	*	0.8	1.4	1.6	2.2	3.0
2001.....	11.9	59.3	1.4	0.4	0.4	1.2	1.9	1.9	2.7	4.3
2000.....	11.5	64.3	1.3	0.3	0.4	1.1	1.6	1.9	2.6	3.5
1999.....	9.9	62.4	1.1	0.3	*	0.7	1.0	1.3	2.1	2.5
Black, both sexes										
2002.....	12.3	162.3	2.6	0.7	0.7	1.8	3.2	4.6	5.6	7.5
2001.....	12.6	169.5	2.8	0.6	0.6	1.8	3.8	4.6	5.3	8.2
2000.....	13.7	197.1	2.3	0.6	0.8	2.2	4.3	5.3	6.3	7.3
1999.....	12.7	192.8	2.7	*	*	1.7	3.7	4.3	5.4	7.8
Black, male										
2002.....	14.1	183.0	3.5	*	*	2.4	3.6	5.8	7.8	10.3
2001.....	13.9	187.4	3.5	*	*	2.4	5.2	5.5	6.5	10.5
2000.....	15.5	227.1	2.3	*	*	2.0	5.6	7.1	8.2	8.3
1999.....	14.6	217.6	2.8	*	*	2.4	4.7	5.2	8.3	9.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, female

2002.....	10.6	140.8	1.8	*	*	*	2.7	3.5	3.7	5.1
2001.....	11.4	150.6	2.0	*	*	*	2.4	3.9	4.2	6.2
2000.....	12.0	166.0	2.3	*	*	2.3	3.0	3.6	4.6	6.3
1999.....	10.9	167.1	2.7	*	*	*	2.6	3.4	2.8	6.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ill-defined and unknown causes of mortality (R95-R99)										
All races 3/, both sexes										
2002.....	6.0	83.8	1.3	0.2	0.3	1.0	1.9	2.3	2.8	3.7
2001.....	7.1	84.8	1.5	0.3	0.3	1.5	2.5	3.0	3.6	5.4
2000.....	7.3	93.9	1.2	0.3	0.3	1.3	2.4	2.7	3.7	4.8
1999.....	5.9	91.3	1.3	0.2	0.3	0.9	1.7	1.7	2.6	3.3
All races 3/, male										
2002.....	6.8	96.5	1.5	0.3	0.3	1.3	2.6	3.2	3.8	5.1
2001.....	8.2	96.5	1.8	0.3	0.3	2.0	3.4	4.1	4.9	6.9
2000.....	8.5	108.9	1.4	0.3	0.4	1.6	3.3	3.7	5.1	6.5
1999.....	6.7	104.3	1.4	0.2	0.3	1.2	2.4	2.2	3.5	4.2
All races 3/, female										
2002.....	5.1	70.4	1.1	*	0.2	0.6	1.2	1.4	1.8	2.4
2001.....	6.1	72.4	1.2	0.2	0.4	1.0	1.5	1.7	2.3	3.8
2000.....	6.2	78.0	1.1	0.2	0.3	1.0	1.4	1.7	2.2	3.2
1999.....	5.2	77.7	1.1	0.3	0.2	0.6	0.9	1.2	1.6	2.4
White, both sexes										
2002.....	5.8	69.5	1.1	0.2	0.2	1.0	2.0	2.2	2.8	3.6
2001.....	7.1	69.3	1.3	0.2	0.3	1.5	2.5	3.0	3.6	5.4
2000.....	7.2	75.6	1.2	0.3	0.3	1.2	2.3	2.6	3.7	4.9
1999.....	5.6	73.1	1.1	0.2	0.2	0.8	1.6	1.6	2.4	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ill-defined and unknown causes of mortality (R95-R99)										
White, male										
2002.....	6.5	81.1	1.1	*	0.2	1.3	2.7	3.2	3.8	4.9
2001.....	8.1	79.9	1.5	*	0.3	2.0	3.5	4.3	4.8	6.9
2000.....	8.2	87.8	1.4	0.3	0.3	1.6	3.3	3.6	5.2	6.6
1999.....	6.2	84.5	1.3	*	0.2	1.1	2.3	2.1	3.3	4.0
White, female										
2002.....	5.1	57.4	1.0	*	*	0.6	1.2	1.2	1.8	2.4
2001.....	6.1	58.2	1.1	*	0.3	1.0	1.5	1.6	2.2	3.8
2000.....	6.2	62.7	0.9	*	0.3	0.9	1.3	1.6	2.1	3.1
1999.....	5.1	61.2	0.8	0.3	*	0.5	0.7	1.0	1.6	2.1
Black, both sexes										
2002.....	8.4	158.0	2.3	*	0.6	1.1	2.0	3.4	3.6	5.6
2001.....	8.8	167.3	2.3	*	*	1.2	2.5	3.4	4.4	6.3
2000.....	9.9	194.0	1.8	*	0.7	1.7	3.3	3.8	4.6	5.8
1999.....	9.1	189.1	2.3	*	*	1.3	2.7	2.8	4.0	5.8
Black, male										
2002.....	10.4	178.1	2.9	*	*	1.4	2.5	4.0	5.1	8.0
2001.....	10.4	184.7	2.9	*	*	1.7	3.5	4.2	5.3	7.9
2000.....	12.0	224.2	1.9	*	*	1.7	4.2	5.2	6.3	6.5
1999.....	11.2	212.7	2.3	*	*	1.7	3.6	3.7	6.2	7.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Ill-defined and unknown causes of mortality (R95-R99)

Black, female

2002.....	6.5	137.1	1.7	*	*	*	1.6	2.8	2.2	3.5
2001.....	7.4	149.0	1.6	*	*	*	1.5	2.6	3.5	4.9
2000.....	7.9	162.7	1.6	*	*	1.6	2.4	2.5	3.1	5.2
1999.....	7.2	164.8	2.4	*	*	*	1.9	2.0	2.1	4.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other ill-defined and unknown causes of mortality (R96-R99)

All races 3/, both sexes

2002.....	5.2	26.9	1.3	0.2	0.3	1.0	1.9	2.3	2.8	3.7
2001.....	6.3	29.4	1.5	0.3	0.3	1.5	2.5	3.0	3.6	5.4
2000.....	6.4	27.6	1.2	0.3	0.3	1.3	2.4	2.7	3.7	4.8
1999.....	5.0	21.6	1.3	0.2	0.3	0.9	1.7	1.7	2.6	3.3

All races 3/, male

2002.....	5.9	31.1	1.5	0.3	0.3	1.3	2.6	3.2	3.8	5.1
2001.....	7.3	32.7	1.8	0.3	0.3	2.0	3.4	4.1	4.9	6.9
2000.....	7.4	31.5	1.4	0.3	0.4	1.6	3.3	3.7	5.1	6.5
1999.....	5.5	24.4	1.4	0.2	0.3	1.2	2.4	2.2	3.5	4.2

All races 3/, female

2002.....	4.5	22.5	1.1	*	0.2	0.6	1.2	1.4	1.8	2.4
2001.....	5.5	25.8	1.2	0.2	0.4	1.0	1.5	1.7	2.3	3.8
2000.....	5.5	23.4	1.1	0.2	0.3	1.0	1.4	1.7	2.2	3.2
1999.....	4.5	18.6	1.1	0.3	0.2	0.6	0.9	1.2	1.6	2.4

White, both sexes

2002.....	5.2	21.8	1.1	0.2	0.2	1.0	2.0	2.2	2.8	3.6
2001.....	6.5	24.6	1.3	0.2	0.3	1.5	2.5	3.0	3.6	5.4
2000.....	6.5	21.5	1.2	0.3	0.3	1.2	2.3	2.6	3.7	4.9
1999.....	4.9	15.7	1.1	0.2	0.2	0.8	1.6	1.6	2.4	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other ill-defined and unknown causes of mortality (R96-R99)										
White, male										
2002.....	5.8	25.6	1.1	*	0.2	1.3	2.7	3.2	3.8	4.9
2001.....	7.3	27.6	1.5	*	0.3	2.0	3.5	4.3	4.8	6.9
2000.....	7.4	24.4	1.4	0.3	0.3	1.6	3.3	3.6	5.2	6.6
1999.....	5.3	18.2	1.3	*	0.2	1.1	2.3	2.1	3.3	4.0
White, female										
2002.....	4.6	17.9	1.0	*	*	0.6	1.2	1.2	1.8	2.4
2001.....	5.6	21.4	1.1	*	0.3	1.0	1.5	1.6	2.2	3.8
2000.....	5.7	18.4	0.9	*	0.3	0.9	1.3	1.6	2.1	3.1
1999.....	4.5	13.1	0.8	0.3	*	0.5	0.7	1.0	1.6	2.1
Black, both sexes										
2002.....	6.5	53.8	2.3	*	0.6	1.1	2.0	3.4	3.6	5.6
2001.....	6.8	53.0	2.3	*	*	1.2	2.5	3.4	4.4	6.3
2000.....	7.6	59.7	1.8	*	0.7	1.7	3.3	3.8	4.6	5.8
1999.....	6.8	50.5	2.3	*	*	1.3	2.7	2.8	4.0	5.8
Black, male										
2002.....	8.2	60.7	2.9	*	*	1.4	2.5	4.0	5.1	8.0
2001.....	8.0	58.4	2.9	*	*	1.7	3.5	4.2	5.3	7.9
2000.....	9.3	70.0	1.9	*	*	1.7	4.2	5.2	6.3	6.5
1999.....	8.5	57.6	2.3	*	*	1.7	3.6	3.7	6.2	7.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other ill-defined and unknown causes of mortality (R96-R99)										
Black, female										
2002.....	5.0	46.6	1.7	*	*	*	1.6	2.8	2.2	3.5
2001.....	5.7	47.3	1.6	*	*	*	1.5	2.6	3.5	4.9
2000.....	6.1	48.9	1.6	*	*	1.6	2.4	2.5	3.1	5.2
1999.....	5.3	43.3	2.4	*	*	*	1.9	2.0	2.1	4.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)										
All races 3/, both sexes										
2002.....	2.9	1.9	0.3	*	0.1	0.3	0.5	0.6	0.8	0.9
2001.....	2.7	1.8	0.3	0.1	0.1	0.3	0.5	0.5	0.6	0.8
2000.....	2.6	1.8	0.4	0.1	0.1	0.3	0.5	0.7	0.7	0.8
1999.....	2.5	1.9	0.3	*	*	0.3	0.4	0.6	0.6	0.8
All races 3/, male										
2002.....	2.4	1.9	0.4	*	*	0.4	0.5	0.8	1.0	1.1
2001.....	2.3	2.3	0.3	*	0.2	0.4	0.6	0.7	0.7	1.0
2000.....	2.3	1.7	0.3	*	*	0.3	0.7	0.9	0.7	1.1
1999.....	2.1	2.3	0.3	*	*	0.3	0.4	0.7	0.7	1.0
All races 3/, female										
2002.....	3.5	1.9	*	*	*	0.2	0.4	0.4	0.6	0.7
2001.....	3.2	1.2	0.3	*	*	*	0.4	0.4	0.5	0.6
2000.....	3.0	1.8	0.4	*	*	0.3	0.4	0.5	0.6	0.5
1999.....	2.9	1.5	*	*	*	*	0.3	0.5	0.5	0.6
White, both sexes										
2002.....	3.0	1.5	0.3	*	*	0.2	0.4	0.6	0.7	0.8
2001.....	2.8	1.7	0.2	*	0.1	0.3	0.4	0.5	0.6	0.7
2000.....	2.7	1.6	0.4	*	0.1	0.3	0.5	0.6	0.5	0.7
1999.....	2.5	1.6	0.3	*	*	0.2	0.3	0.5	0.5	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)										
White, male										
2002.....	2.4	1.4	*	*	*	0.2	0.4	0.8	0.8	0.9
2001.....	2.2	2.2	*	*	*	0.4	0.5	0.6	0.6	0.8
2000.....	2.2	1.6	0.3	*	*	0.3	0.6	0.8	0.6	1.0
1999.....	2.1	1.9	*	*	*	0.3	0.3	0.6	0.5	0.9
White, female										
2002.....	3.7	1.6	*	*	*	*	*	0.4	0.5	0.6
2001.....	3.4	*	*	*	*	*	0.4	0.3	0.5	0.5
2000.....	3.2	1.6	0.4	*	*	*	0.3	0.4	0.4	0.4
1999.....	3.0	*	*	*	*	*	*	0.3	0.5	0.4
Black, both sexes										
2002.....	3.2	4.3	*	*	*	0.7	1.1	1.2	2.1	1.9
2001.....	3.1	*	*	*	*	*	1.3	1.3	0.9	1.9
2000.....	3.0	*	*	*	*	*	1.0	1.5	1.7	1.4
1999.....	3.0	3.6	*	*	*	*	1.0	1.4	1.4	2.0
Black, male										
2002.....	3.3	*	*	*	*	*	*	1.7	2.8	2.3
2001.....	3.1	*	*	*	*	*	1.7	*	*	2.6
2000.....	3.0	*	*	*	*	*	*	2.0	1.9	1.8
1999.....	3.0	*	*	*	*	*	*	*	2.1	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

Black, female

2002.....	3.2	*	*	*	*	*	*	*	1.4	1.6
2001.....	3.0	*	*	*	*	*	*	*	*	1.3
2000.....	3.0	*	*	*	*	*	*	*	1.5	*
1999.....	3.0	*	*	*	*	*	*	1.4	*	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XX. External causes of mortality (*U01-*U03,V01-Y89)										
All races 3/, both sexes										
2002.....	56.9	33.8	13.7	6.8	9.9	52.6	71.9	60.6	55.5	61.1
2001 4/.....	56.2	35.1	14.3	7.2	9.9	51.0	70.6	60.9	55.6	62.4
2000.....	53.8	34.5	14.6	7.7	10.6	51.7	68.9	56.5	51.9	57.4
1999.....	54.2	33.9	15.3	8.1	10.7	52.7	67.2	56.4	54.0	57.6
All races 3/, male										
2002.....	79.8	37.6	16.3	7.9	12.7	76.5	114.0	95.6	84.8	90.3
2001 4/.....	79.3	38.9	16.8	8.5	12.9	75.1	113.0	96.1	85.3	92.4
2000.....	75.8	38.1	17.0	8.8	14.0	75.5	109.5	89.6	79.6	86.0
1999.....	76.2	38.3	17.3	9.5	14.1	76.3	106.5	88.9	82.6	86.0
All races 3/, female										
2002.....	34.8	29.8	11.1	5.7	6.9	27.5	27.6	24.4	25.7	31.8
2001 4/.....	34.0	31.2	11.8	5.8	6.7	25.4	26.2	24.7	25.4	32.5
2000.....	32.5	30.8	12.1	6.4	7.1	26.5	26.5	22.6	23.8	29.0
1999.....	33.0	29.2	13.3	6.6	7.2	27.8	26.2	23.3	25.2	29.4
White, both sexes										
2002.....	57.4	26.1	12.3	5.9	9.3	52.1	67.3	56.1	53.8	60.3
2001 4/.....	56.5	28.8	13.3	6.5	9.3	49.5	65.6	57.1	53.9	62.0
2000.....	53.6	27.6	13.1	7.0	10.3	50.4	63.2	51.7	49.6	56.6
1999.....	53.7	27.9	13.6	7.1	10.5	50.6	61.4	52.1	51.7	56.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XX. External causes of mortality (*U01-*U03,V01-Y89)										
White, male										
2002.....	79.0	30.7	14.5	6.8	11.8	74.1	104.5	86.0	80.8	87.9
2001 4/.....	78.3	31.8	15.8	7.4	12.1	71.1	102.8	88.2	81.6	91.2
2000.....	74.1	31.8	15.4	8.1	13.4	71.7	98.4	80.1	75.1	84.0
1999.....	74.2	32.6	15.4	8.2	13.6	71.1	95.2	80.8	77.8	83.3
White, female										
2002.....	36.4	21.3	10.1	5.0	6.6	28.9	27.7	24.3	25.6	32.2
2001 4/.....	35.2	25.7	10.5	5.6	6.5	26.5	26.0	24.3	25.2	32.2
2000.....	33.6	23.2	10.7	5.9	7.0	27.7	25.7	22.0	23.2	28.7
1999.....	33.9	23.1	11.7	5.9	7.1	28.9	25.4	22.2	24.6	29.0
Black, both sexes										
2002.....	63.7	72.8	21.0	11.2	12.9	59.9	106.7	100.3	78.6	78.5
2001 4/.....	64.2	68.9	21.2	10.6	12.6	62.5	107.7	94.8	76.8	76.9
2000.....	63.9	70.6	22.8	11.3	13.0	63.3	111.4	94.0	76.1	73.4
1999.....	65.2	67.1	24.8	13.3	13.2	68.4	108.3	91.2	76.9	75.1
Black, male										
2002.....	99.9	73.5	26.0	13.1	17.9	96.4	184.7	178.2	132.9	126.6
2001 4/.....	100.2	77.2	24.1	14.2	17.7	102.8	189.7	165.4	128.5	117.9
2000.....	100.2	72.3	25.5	13.3	18.2	103.1	193.6	165.9	126.7	117.9
1999.....	101.7	71.2	27.7	16.7	18.0	110.9	189.6	158.4	127.6	119.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XX. External causes of mortality (*U01-*U03,V01-Y89)										
Black, female										
2002.....	30.9	72.0	15.8	9.1	7.9	22.2	30.7	29.1	29.9	35.7
2001 4/.....	31.5	60.2	18.3	6.9	7.4	20.9	28.8	30.4	30.3	40.5
2000.....	30.9	68.8	19.9	9.2	7.6	22.6	33.5	29.0	30.5	34.0
1999.....	32.2	62.8	21.9	9.9	8.2	25.0	31.1	30.4	31.3	35.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
All races 3/, both sexes										
2002.....	37.0	23.5	10.5	5.9	7.3	35.0	40.9	32.3	30.8	34.9
2001.....	35.7	24.2	11.2	6.4	7.4	32.8	39.5	31.4	28.5	34.1
2000.....	34.8	23.1	11.9	6.8	7.7	33.4	38.8	30.4	28.6	32.9
1999.....	35.1	22.3	12.4	7.1	8.1	33.3	37.5	30.0	29.2	32.2
All races 3/, male										
2002.....	48.9	26.8	12.9	7.1	9.2	47.6	62.4	49.5	46.1	50.5
2001.....	47.2	26.2	13.3	7.6	9.5	45.3	60.7	48.0	43.1	49.5
2000.....	46.2	25.7	14.0	7.8	9.9	45.8	58.9	47.0	43.1	48.9
1999.....	46.4	25.4	14.4	8.5	10.4	44.8	57.1	46.1	43.9	47.4
All races 3/, female										
2002.....	25.6	20.0	8.1	4.7	5.3	21.8	18.4	14.5	15.2	19.2
2001.....	24.5	22.1	8.9	5.0	5.3	19.5	17.2	14.3	13.8	18.7
2000.....	23.8	20.5	9.7	5.7	5.5	20.3	17.8	13.5	13.9	17.0
1999.....	24.1	19.0	10.3	5.6	5.6	21.1	17.1	13.7	14.3	17.0
White, both sexes										
2002.....	38.7	18.6	9.9	5.2	7.0	37.9	43.8	33.5	32.2	35.5
2001.....	37.0	20.6	11.0	5.8	7.1	34.9	41.7	32.7	29.3	34.8
2000.....	35.9	19.1	11.1	6.2	7.6	36.0	40.7	30.9	29.1	33.4
1999.....	36.0	18.9	11.5	6.3	7.9	35.6	38.9	30.8	29.3	32.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
White, male										
2002.....	50.4	22.3	11.8	6.3	8.8	51.0	66.4	50.7	47.9	50.9
2001.....	48.4	22.1	13.3	6.7	9.0	47.8	64.2	49.6	44.0	50.3
2000.....	47.0	21.7	13.2	7.2	9.6	48.8	61.7	47.2	43.7	49.5
1999.....	46.9	21.8	13.2	7.4	10.1	47.2	58.9	47.1	43.8	47.5
White, female										
2002.....	27.3	14.7	7.8	4.2	5.2	23.9	19.6	15.1	15.9	19.7
2001.....	25.9	18.9	8.5	4.8	5.1	21.1	17.8	14.8	14.0	19.0
2000.....	25.1	16.4	9.0	5.3	5.5	22.3	18.3	13.8	13.8	17.0
1999.....	25.3	15.8	9.6	5.1	5.6	23.3	17.6	13.8	14.3	17.0
Black, both sexes										
2002.....	33.1	48.9	14.1	9.3	9.0	23.9	32.0	32.8	30.1	38.1
2001.....	33.5	43.1	13.6	9.4	9.4	24.7	33.0	30.5	30.0	35.8
2000.....	33.5	44.5	16.2	9.7	8.8	23.7	34.3	33.1	31.9	35.9
1999.....	35.2	41.6	17.7	11.3	9.6	25.0	34.6	31.5	32.9	36.6
Black, male										
2002.....	47.9	51.1	18.5	11.1	12.1	34.6	49.5	53.6	47.7	58.9
2001.....	48.2	48.5	15.4	12.6	12.9	36.1	51.0	48.3	46.6	53.3
2000.....	49.0	47.7	18.8	11.9	12.1	34.5	52.9	54.1	49.6	55.7
1999.....	51.0	45.9	21.0	14.5	13.0	36.6	54.7	51.0	51.7	55.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
Black, female										
2002.....	19.7	46.6	9.6	7.5	5.8	12.8	14.9	13.8	14.4	19.7
2001.....	20.1	37.5	11.9	6.2	5.8	12.9	15.6	14.2	15.1	20.2
2000.....	19.5	41.3	13.5	7.6	5.4	12.7	16.7	14.1	15.9	18.3
1999.....	20.8	37.3	14.3	8.0	6.1	13.1	15.6	13.8	16.0	19.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Transport accidents (V01-V99,Y85)										
All races 3/, both sexes										
2002.....	16.8	3.1	4.1	3.5	4.9	28.4	29.9	20.5	17.1	17.1
2001.....	16.6	3.7	4.4	3.8	5.0	26.6	29.2	20.9	16.9	17.4
2000.....	16.6	4.6	4.5	4.1	5.3	27.1	29.3	20.4	16.8	17.1
1999.....	16.6	5.0	4.6	4.4	5.3	27.1	28.2	20.0	17.0	16.9
All races 3/, male										
2002.....	23.7	3.5	4.8	4.0	5.9	37.0	44.4	31.0	25.4	24.7
2001.....	23.5	3.3	5.0	4.3	6.2	35.4	43.9	31.4	25.4	24.7
2000.....	23.3	4.8	5.0	4.4	6.4	35.5	43.3	30.8	24.7	24.8
1999.....	23.1	5.2	5.1	5.0	6.5	34.8	41.8	29.9	24.7	24.0
All races 3/, female										
2002.....	10.1	2.8	3.4	3.1	3.9	19.3	14.6	9.7	8.7	9.6
2001.....	10.0	4.2	3.8	3.3	3.8	17.3	13.8	10.1	8.3	10.2
2000.....	10.2	4.3	4.0	3.8	4.1	18.1	14.6	9.8	8.7	9.5
1999.....	10.4	4.8	4.2	3.6	4.1	19.0	14.0	10.0	9.3	9.8
White, both sexes										
2002.....	17.3	2.7	3.9	3.4	5.0	31.0	31.5	20.7	17.6	17.2
2001.....	17.0	3.7	4.3	3.6	5.0	28.6	30.4	21.6	17.1	17.6
2000.....	17.0	4.1	4.3	4.1	5.4	29.6	30.5	20.3	16.8	17.3
1999.....	17.0	4.6	4.1	4.1	5.5	29.4	29.2	20.4	17.2	16.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Transport accidents (V01-V99,Y85)										
White, male										
2002.....	24.2	3.1	4.5	3.8	6.0	40.2	46.7	31.0	25.9	24.5
2001.....	23.9	3.2	4.9	4.0	6.2	37.8	45.8	32.3	25.4	24.7
2000.....	23.6	4.5	4.7	4.4	6.5	38.4	45.0	30.3	24.6	24.8
1999.....	23.4	5.2	4.4	4.7	6.7	37.3	43.1	30.2	24.6	23.7
White, female										
2002.....	10.5	2.2	3.4	2.9	3.9	21.3	15.3	9.8	9.0	9.7
2001.....	10.3	4.1	3.6	3.1	3.8	18.9	14.0	10.4	8.4	10.4
2000.....	10.5	3.7	3.8	3.7	4.3	20.2	15.1	9.9	8.7	9.8
1999.....	10.7	3.9	3.8	3.5	4.3	21.0	14.4	10.1	9.6	9.8
Black, both sexes										
2002.....	15.4	5.3	5.0	4.6	4.6	17.9	24.6	23.0	18.0	19.2
2001.....	15.9	*	5.2	5.3	5.3	18.8	25.8	21.0	18.9	18.1
2000.....	16.0	6.5	5.6	4.5	4.8	17.3	26.6	23.5	19.4	17.9
1999.....	16.3	7.3	7.1	5.8	5.0	18.7	25.8	21.3	18.5	19.9
Black, male										
2002.....	23.1	*	6.4	5.0	5.6	24.5	37.1	37.5	28.9	29.8
2001.....	23.8	*	5.7	6.6	7.3	26.5	39.1	33.2	30.0	27.2
2000.....	24.1	6.7	6.2	4.9	6.5	24.2	40.3	38.2	30.7	28.6
1999.....	24.2	*	8.0	7.2	6.4	25.8	39.2	34.3	29.4	29.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Transport accidents (V01-V99,Y85)										
Black, female										
2002.....	8.3	*	3.4	4.2	3.5	11.2	12.5	9.8	8.2	9.8
2001.....	8.7	*	4.7	4.0	3.4	10.8	13.0	9.9	8.9	10.0
2000.....	8.5	*	5.0	4.1	3.0	10.3	13.7	10.2	9.2	8.4
1999.....	9.2	8.7	6.2	4.4	3.6	11.4	13.1	9.5	8.6	11.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Railway accidents (V05,V15,V80.6,V81.2-V81.9)										
All races 3/, both sexes										
2002.....	0.2	*	*	*	*	0.2	0.3	0.2	0.2	0.2
2001.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.3
2000.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.3
1999.....	0.2	*	*	*	*	0.2	0.3	0.2	0.2	0.3
All races 3/, male										
2002.....	0.3	*	*	*	*	0.3	0.5	0.3	0.4	0.4
2001.....	0.3	*	*	*	*	0.3	0.3	0.4	0.4	0.5
2000.....	0.3	*	*	*	*	0.4	0.4	0.4	0.3	0.5
1999.....	0.3	*	*	*	*	0.3	0.6	0.4	0.4	0.6
All races 3/, female										
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.2	*	*	*	*	0.2	0.3	0.2	0.2	0.2
2001.....	0.2	*	*	*	*	0.2	0.2	0.3	0.2	0.2
2000.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.3
1999.....	0.2	*	*	*	*	0.2	0.3	0.2	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
All races 3/, both sexes										
2002.....	15.7	3.0	3.9	3.3	4.5	27.6	28.8	19.7	16.0	15.8
2001.....	15.4	3.6	4.1	3.6	4.6	25.7	28.0	19.6	15.6	15.9
2000.....	15.4	4.4	4.2	3.8	4.8	26.0	28.0	19.2	15.5	15.6
1999.....	15.2	4.8	4.2	3.9	4.8	25.9	26.5	18.4	15.4	15.1
All races 3/, male										
2002.....	21.9	3.3	4.5	3.8	5.3	36.0	42.6	29.6	23.6	22.4
2001.....	21.5	3.2	4.5	4.0	5.7	34.0	42.1	29.5	23.1	22.3
2000.....	21.3	4.6	4.7	4.1	5.8	33.9	41.2	28.6	22.6	22.1
1999.....	20.9	5.0	4.7	4.4	5.7	33.1	39.1	27.2	22.2	21.2
All races 3/, female										
2002.....	9.8	2.8	3.3	2.9	3.6	18.9	14.3	9.4	8.3	9.2
2001.....	9.5	4.0	3.6	3.0	3.5	16.9	13.4	9.5	8.0	9.6
2000.....	9.7	4.2	3.8	3.5	3.8	17.5	14.2	9.5	8.2	9.1
1999.....	9.7	4.7	3.8	3.3	3.8	18.3	13.4	9.4	8.5	9.0
White, both sexes										
2002.....	16.2	2.6	3.7	3.2	4.6	30.2	30.4	19.8	16.4	15.8
2001.....	15.7	3.5	3.9	3.3	4.6	27.6	29.1	20.2	15.7	16.1
2000.....	15.7	3.9	4.0	3.7	5.0	28.4	29.1	19.0	15.4	15.8
1999.....	15.5	4.5	3.8	3.7	5.0	28.1	27.4	18.6	15.5	14.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
White, male										
2002.....	22.4	2.9	4.2	3.6	5.5	39.0	44.8	29.5	23.9	22.2
2001.....	21.9	3.1	4.4	3.7	5.6	36.2	43.8	30.1	23.1	22.3
2000.....	21.6	4.2	4.5	4.0	5.9	36.8	42.6	27.9	22.4	22.1
1999.....	21.1	5.0	4.1	4.2	5.9	35.5	40.4	27.4	22.1	20.8
White, female										
2002.....	10.1	2.2	3.2	2.7	3.7	20.9	15.0	9.5	8.6	9.3
2001.....	9.8	3.8	3.4	2.9	3.5	18.5	13.5	9.7	8.0	9.8
2000.....	10.0	3.5	3.6	3.4	4.0	19.5	14.6	9.6	8.2	9.3
1999.....	10.0	3.9	3.5	3.2	4.0	20.2	13.7	9.5	8.7	8.9
Black, both sexes										
2002.....	14.5	5.2	4.8	4.3	4.2	17.4	24.0	22.4	17.2	17.9
2001.....	14.8	*	5.0	4.9	4.8	18.1	25.2	20.1	17.7	16.7
2000.....	15.0	6.5	5.4	4.3	4.6	16.5	25.8	22.8	18.0	16.4
1999.....	15.1	7.1	6.4	5.3	4.6	17.8	24.8	20.1	17.0	18.2
Black, male										
2002.....	21.5	*	6.3	4.8	5.0	23.7	36.1	36.2	27.4	27.6
2001.....	22.0	*	5.3	6.2	6.5	25.5	38.0	31.9	28.1	24.9
2000.....	22.5	6.7	5.8	4.5	6.2	22.8	38.8	36.7	28.5	26.0
1999.....	22.2	*	7.0	6.3	5.7	24.5	37.7	32.4	27.0	27.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
Black, female										
2002.....	8.0	*	3.4	3.8	3.3	11.0	12.2	9.7	7.9	9.3
2001.....	8.2	*	4.6	3.6	3.1	10.5	12.9	9.5	8.4	9.5
2000.....	8.2	*	4.9	4.0	2.9	10.1	13.5	10.2	8.6	7.9
1999.....	8.7	8.7	5.8	4.2	3.5	11.0	12.6	9.0	8.1	10.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pedestrian involved in collision with motor vehicle (V02-V04)										
All races 3/, both sexes										
2002.....	1.3	*	1.0	0.6	0.5	1.0	1.1	1.0	1.0	1.3
2001.....	1.2	*	1.1	0.6	0.7	0.9	1.1	1.0	1.0	1.4
2000.....	1.2	*	1.2	0.7	0.6	0.8	1.0	1.0	1.0	1.2
1999.....	1.2	*	1.1	0.8	0.5	0.8	1.1	1.0	1.0	1.3
All races 3/, male										
2002.....	1.7	*	1.3	0.7	0.6	1.4	1.7	1.6	1.6	1.8
2001.....	1.8	*	1.4	0.8	1.0	1.2	1.8	1.6	1.5	1.9
2000.....	1.7	*	1.4	0.8	0.7	1.0	1.5	1.5	1.5	1.7
1999.....	1.7	*	1.2	0.9	0.7	1.2	1.6	1.5	1.5	1.9
All races 3/, female										
2002.....	0.8	*	0.6	0.5	0.4	0.6	0.5	0.5	0.5	0.8
2001.....	0.8	*	0.8	0.4	0.4	0.5	0.5	0.3	0.5	0.8
2000.....	0.8	*	0.9	0.6	0.4	0.6	0.4	0.4	0.5	0.7
1999.....	0.7	*	1.0	0.7	0.4	0.5	0.5	0.5	0.4	0.6
White, both sexes										
2002.....	1.2	*	0.9	0.5	0.5	0.9	1.1	1.0	1.0	1.1
2001.....	1.2	*	1.0	0.5	0.7	0.8	1.2	1.0	1.0	1.2
2000.....	1.1	*	1.0	0.6	0.6	0.8	1.0	0.9	0.9	1.1
1999.....	1.1	*	1.0	0.6	0.5	0.8	1.0	0.9	0.9	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pedestrian involved in collision with motor vehicle (V02-V04)										
White, male										
2002.....	1.6	*	1.2	0.7	0.6	1.3	1.8	1.5	1.5	1.5
2001.....	1.7	*	1.4	0.6	1.0	1.2	1.8	1.7	1.4	1.8
2000.....	1.6	*	1.2	0.7	0.7	1.0	1.6	1.3	1.4	1.5
1999.....	1.5	*	1.1	0.7	0.7	1.2	1.6	1.3	1.4	1.6
White, female										
2002.....	0.7	*	0.6	0.4	0.4	0.5	0.5	0.4	0.5	0.6
2001.....	0.7	*	0.7	0.3	0.4	0.5	0.5	0.3	0.5	0.7
2000.....	0.7	*	0.8	0.5	0.4	0.6	0.5	0.4	0.4	0.6
1999.....	0.7	*	1.0	0.5	0.4	0.5	0.5	0.4	0.4	0.5
Black, both sexes										
2002.....	1.6	*	1.3	1.0	0.6	1.3	1.0	1.4	1.4	2.4
2001.....	1.7	*	1.5	1.2	0.9	1.1	1.0	1.0	1.2	2.2
2000.....	1.7	*	1.9	1.2	0.7	0.8	0.8	1.3	1.1	2.1
1999.....	1.8	*	1.6	1.6	0.7	0.8	1.1	1.7	1.5	2.4
Black, male										
2002.....	2.4	*	1.7	*	*	1.7	1.5	2.1	2.2	3.6
2001.....	2.4	*	1.7	1.8	1.4	1.3	1.6	1.6	1.7	2.8
2000.....	2.5	*	2.3	1.5	*	*	*	2.1	1.6	3.4
1999.....	2.8	*	2.0	1.8	*	*	1.8	2.6	2.7	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)										
All races 3/, both sexes										
2002.....	0.2	*	*	0.2	0.3	0.2	0.1	0.1	0.2	0.2
2001.....	0.2	*	*	0.2	0.3	0.3	0.2	0.2	0.2	0.2
2000.....	0.2	*	*	0.2	0.5	0.2	0.1	0.1	0.2	0.2
1999.....	0.2	*	*	0.3	0.4	0.3	0.2	0.2	0.2	0.2
All races 3/, male										
2002.....	0.4	*	*	0.3	0.6	0.4	0.2	0.2	0.4	0.3
2001.....	0.4	*	*	0.3	0.5	0.5	0.3	0.4	0.3	0.4
2000.....	0.4	*	*	0.4	0.8	0.4	0.3	0.2	0.3	0.3
1999.....	0.4	*	*	0.5	0.6	0.6	0.4	0.3	0.4	0.4
All races 3/, female										
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	0.2	*	*	*	*	*
White, both sexes										
2002.....	0.2	*	*	0.1	0.3	0.2	0.2	0.2	0.2	0.2
2001.....	0.2	*	*	0.2	0.3	0.3	0.2	0.2	0.2	0.2
2000.....	0.2	*	*	0.2	0.4	0.3	0.2	*	0.2	0.2
1999.....	0.2	*	*	0.3	0.4	0.4	0.2	0.2	0.3	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)

All races 3/, both sexes

2002.....	1.1	*	*	*	0.1	0.8	2.3	2.0	1.8	1.7
2001.....	1.1	*	*	*	0.1	0.8	2.3	2.2	1.9	1.7
2000.....	1.0	*	*	*	0.1	0.8	2.1	2.1	1.6	1.5
1999.....	0.8	*	*	*	0.1	0.6	2.0	1.8	1.3	1.3

All races 3/, male

2002.....	2.1	*	*	*	*	1.5	4.1	3.7	3.3	3.0
2001.....	2.0	*	*	*	0.3	1.5	4.2	4.1	3.4	2.9
2000.....	1.8	*	*	*	0.2	1.5	3.8	3.9	2.9	2.7
1999.....	1.6	*	*	*	0.2	1.1	3.6	3.3	2.4	2.4

All races 3/, female

2002.....	0.2	*	*	*	*	*	0.3	0.3	0.3	0.5
2001.....	0.2	*	*	*	*	*	0.3	0.3	0.4	0.5
2000.....	0.2	*	*	*	*	*	*	0.2	0.2	0.4
1999.....	0.2	*	*	*	*	*	0.2	0.2	*	0.2

White, both sexes

2002.....	1.2	*	*	*	*	0.9	2.3	2.0	1.8	1.9
2001.....	1.1	*	*	*	0.2	1.0	2.4	2.2	1.8	1.8
2000.....	1.0	*	*	*	0.1	0.9	2.1	2.0	1.5	1.6
1999.....	0.9	*	*	*	0.2	0.7	2.0	1.6	1.3	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)										
White, male										
2002.....	2.2	*	*	*	*	1.7	4.2	3.6	3.1	3.1
2001.....	2.1	*	*	*	0.3	1.8	4.4	4.0	3.2	3.0
2000.....	1.9	*	*	*	*	1.6	3.9	3.6	2.6	2.8
1999.....	1.6	*	*	*	0.2	1.2	3.7	2.9	2.4	2.5
White, female										
2002.....	0.2	*	*	*	*	*	0.3	0.3	0.4	0.6
2001.....	0.2	*	*	*	*	*	0.3	0.3	0.4	0.6
2000.....	0.2	*	*	*	*	*	*	0.3	0.3	0.4
1999.....	0.2	*	*	*	*	*	*	0.3	*	0.2
Black, both sexes										
2002.....	0.9	*	*	*	*	*	2.2	2.8	2.4	1.5
2001.....	0.9	*	*	*	*	*	2.1	2.9	2.7	1.3
2000.....	0.9	*	*	*	*	*	2.2	3.2	2.9	1.5
1999.....	0.8	*	*	*	*	*	2.3	3.1	1.7	1.2
Black, male										
2002.....	1.8	*	*	*	*	*	4.3	5.5	5.1	3.2
2001.....	1.7	*	*	*	*	*	4.0	5.9	5.4	2.7
2000.....	1.9	*	*	*	*	*	4.2	6.6	5.8	3.1
1999.....	1.6	*	*	*	*	*	4.5	6.4	3.5	2.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accident involving collision with railway train (V25,V35,V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)

All races 3/, both sexes

2002.....	0.1	*	*	*	*	0.1	0.1	*	*	0.1
2001.....	0.1	*	*	*	*	*	0.2	0.1	*	0.1
2000.....	0.1	*	*	*	*	0.2	0.1	0.1	0.1	*
1999.....	0.1	*	*	*	*	0.2	0.2	*	*	*

All races 3/, male

2002.....	0.1	*	*	*	*	*	0.2	*	*	*
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	*
2000.....	0.1	*	*	*	*	0.2	0.2	*	0.2	*
1999.....	0.1	*	*	*	*	0.2	0.2	*	*	*

All races 3/, female

2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.1	*	*	*	*	0.1	0.2	*	*	*
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	0.1
2000.....	0.1	*	*	*	*	0.2	0.1	0.1	0.2	*
1999.....	0.1	*	*	*	*	0.2	0.2	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)

All races 3/, both sexes

2002.....	0.1	*	*	*	*	0.1	0.1	*	*	*
2001.....	0.1	*	*	*	*	*	0.2	0.1	*	0.1
2000.....	0.1	*	*	*	*	0.2	0.1	0.1	0.1	*
1999.....	0.1	*	*	*	*	0.2	0.2	*	*	*

All races 3/, male

2002.....	0.1	*	*	*	*	*	0.2	*	*	*
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	*
2000.....	0.1	*	*	*	*	0.2	*	*	0.2	*
1999.....	0.1	*	*	*	*	0.2	*	*	*	*

All races 3/, female

2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.1	*	*	*	*	0.1	0.2	*	*	*
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	0.1
2000.....	0.1	*	*	*	*	0.2	0.1	0.1	0.2	*
1999.....	0.1	*	*	*	*	0.2	0.2	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57,V60-V61,V66-V67,V70-V71,V76-V77)

All races 3/, both sexes

2002.....	1.8	*	0.2	0.2	0.4	4.5	4.6	2.8	1.9	1.8
2001.....	1.5	*	0.1	0.2	0.2	3.7	4.0	2.2	1.8	1.5
2000.....	1.4	*	0.2	0.2	0.3	3.4	3.5	2.0	1.5	1.6
1999.....	1.4	*	0.1	0.1	0.3	3.3	3.3	2.0	1.6	1.4

All races 3/, male

2002.....	2.6	*	*	0.2	0.4	6.3	7.0	4.4	2.9	2.7
2001.....	2.2	*	*	*	0.3	5.2	6.1	3.4	2.7	2.1
2000.....	2.0	*	*	*	0.3	4.8	5.7	3.2	2.2	2.3
1999.....	2.0	*	*	*	0.4	4.4	5.3	3.2	2.5	2.1

All races 3/, female

2002.....	0.9	*	*	*	0.3	2.6	1.9	1.1	0.8	0.9
2001.....	0.8	*	*	*	0.2	2.1	1.7	1.0	0.8	0.8
2000.....	0.8	*	*	*	0.3	1.9	1.3	0.8	0.8	1.0
1999.....	0.7	*	*	*	0.2	2.2	1.3	0.8	0.7	0.7

White, both sexes

2002.....	1.8	*	0.2	0.2	0.3	4.9	4.8	2.8	1.9	1.8
2001.....	1.5	*	*	0.1	0.3	3.8	3.9	2.2	1.7	1.5
2000.....	1.4	*	*	0.2	0.3	3.6	3.5	1.9	1.5	1.7
1999.....	1.3	*	*	0.1	0.3	3.6	3.3	2.0	1.6	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57,V60-V61,V66-V67,V70-V71,V76-V77)

White, male

2002.....	2.6	*	*	*	0.3	6.7	7.3	4.4	3.0	2.5
2001.....	2.2	*	*	*	0.3	5.3	6.1	3.4	2.6	2.1
2000.....	2.0	*	*	*	0.3	5.0	5.6	3.0	2.2	2.3
1999.....	2.0	*	*	*	0.4	4.7	5.3	3.1	2.4	2.0

White, female

2002.....	1.0	*	*	*	0.3	2.9	2.0	1.2	0.8	1.0
2001.....	0.8	*	*	*	*	2.3	1.5	1.0	0.8	0.8
2000.....	0.8	*	*	*	0.3	2.1	1.3	0.8	0.9	1.0
1999.....	0.7	*	*	*	*	2.4	1.2	0.8	0.7	0.6

Black, both sexes

2002.....	1.9	*	*	*	*	3.3	4.3	3.1	2.1	2.4
2001.....	1.8	*	*	*	*	3.6	4.7	2.6	2.3	1.7
2000.....	1.6	*	*	*	*	2.6	3.7	2.7	1.4	1.7
1999.....	1.6	*	*	*	*	2.8	4.1	2.5	2.1	1.8

Black, male

2002.....	3.0	*	*	*	*	5.1	6.7	5.8	3.3	4.0
2001.....	2.8	*	*	*	*	5.3	7.4	4.2	3.7	2.6
2000.....	2.5	*	*	*	*	3.9	5.9	4.8	2.6	2.7
1999.....	2.6	*	*	*	*	4.0	6.4	4.3	3.6	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51, V56-V57,V60-V61,V66-V67,V70-V71,V76-V77)

Black, female

2002.....	0.9	*	*	*	*	1.5	1.9	*	*	*
2001.....	0.8	*	*	*	*	1.7	2.1	*	*	*
2000.....	0.7	*	*	*	*	*	1.7	*	*	*
1999.....	0.8	*	*	*	*	1.6	1.9	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)										
All races 3/, both sexes										
2002.....	3.9	1.1	1.0	0.8	0.9	6.9	6.8	4.2	3.4	3.4
2001.....	3.7	1.6	1.0	0.8	1.0	6.3	5.9	4.2	3.1	3.3
2000.....	3.7	1.5	0.9	0.9	0.9	6.2	6.0	4.1	3.2	3.3
1999.....	3.7	1.6	0.9	0.9	1.0	6.0	5.7	4.0	3.5	3.0
All races 3/, male										
2002.....	4.8	1.5	1.0	0.9	0.9	8.3	9.1	6.0	4.3	4.2
2001.....	4.5	1.6	0.9	0.8	1.0	7.6	8.1	5.7	4.3	4.2
2000.....	4.5	1.5	0.9	0.9	0.9	7.1	8.0	5.3	4.3	4.1
1999.....	4.4	1.7	1.0	0.9	1.0	7.1	7.5	5.2	4.6	3.5
All races 3/, female										
2002.....	3.0	*	0.9	0.8	0.9	5.3	4.4	2.4	2.4	2.5
2001.....	2.9	1.6	1.1	0.9	1.0	4.9	3.6	2.7	2.0	2.4
2000.....	2.9	1.5	0.9	0.9	0.8	5.2	3.8	3.0	2.2	2.4
1999.....	2.9	1.4	0.7	0.9	1.1	4.8	3.8	2.7	2.4	2.4
White, both sexes										
2002.....	4.1	1.0	0.9	0.8	1.0	7.6	7.2	4.2	3.4	3.4
2001.....	3.8	1.5	1.0	0.8	1.1	6.8	6.3	4.4	3.2	3.3
2000.....	3.8	1.5	0.9	0.9	0.9	6.8	6.2	4.2	3.2	3.3
1999.....	3.8	1.5	0.8	0.9	1.1	6.6	5.9	4.1	3.5	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)										
White, male										
2002.....	5.0	1.4	1.0	0.9	1.0	9.0	9.7	5.9	4.3	4.2
2001.....	4.7	1.6	0.9	0.7	1.1	8.1	8.5	5.9	4.3	4.1
2000.....	4.6	1.5	0.9	1.0	0.9	7.8	8.1	5.3	4.2	4.1
1999.....	4.5	1.8	0.9	0.9	1.0	7.6	7.7	5.4	4.6	3.2
White, female										
2002.....	3.2	*	0.9	0.7	1.0	6.1	4.6	2.3	2.4	2.6
2001.....	3.0	1.5	1.1	0.9	1.0	5.3	3.9	2.9	2.1	2.4
2000.....	3.1	1.5	0.9	0.9	0.9	5.7	4.1	2.9	2.3	2.4
1999.....	3.1	*	0.7	0.9	1.2	5.4	3.9	2.7	2.4	2.5
Black, both sexes										
2002.....	3.3	*	1.2	0.8	0.7	4.1	5.5	5.5	4.1	3.4
2001.....	3.3	*	1.1	0.9	0.8	4.4	4.9	4.3	3.5	3.7
2000.....	3.3	*	1.0	0.7	0.8	3.7	5.5	4.7	3.7	3.3
1999.....	3.4	*	1.1	0.9	1.0	3.6	5.1	4.2	4.0	4.2
Black, male										
2002.....	4.4	*	1.6	*	*	5.7	7.0	7.8	5.7	4.5
2001.....	4.4	*	*	*	*	5.6	6.9	5.9	5.2	4.8
2000.....	4.4	*	*	*	*	4.5	8.0	6.0	5.7	4.0
1999.....	4.5	*	*	*	*	5.0	6.9	5.7	5.5	5.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

Black, female

2002.....	2.4	*	*	*	*	2.4	4.1	3.4	2.6	2.5
2001.....	2.3	*	*	*	*	3.2	3.1	2.8	2.1	2.8
2000.....	2.2	*	*	*	*	2.9	3.2	3.5	1.9	2.6
1999.....	2.3	*	*	*	*	2.2	3.4	2.9	2.7	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)										
All races 3/, both sexes										
2002.....	1.5	*	0.3	0.3	0.5	3.7	3.4	2.3	1.8	1.7
2001.....	1.4	*	0.2	0.2	0.5	2.8	3.3	1.9	1.7	1.6
2000.....	1.3	0.6	0.2	0.2	0.5	2.9	3.2	2.2	1.5	1.5
1999.....	1.3	*	0.3	0.3	0.3	3.0	3.1	2.0	1.5	1.5
All races 3/, male										
2002.....	2.2	*	0.4	0.3	0.5	4.6	5.0	3.4	2.6	2.4
2001.....	2.0	*	*	0.3	0.5	3.4	5.0	2.8	2.6	2.3
2000.....	1.9	*	*	0.2	0.6	3.9	4.7	3.4	2.2	2.1
1999.....	1.9	*	0.3	0.3	0.3	3.9	4.6	3.0	2.1	2.1
All races 3/, female										
2002.....	0.9	*	*	0.3	0.4	2.7	1.7	1.1	1.0	1.1
2001.....	0.8	*	*	0.2	0.5	2.1	1.5	1.0	0.8	0.9
2000.....	0.7	*	*	0.3	0.4	1.9	1.6	0.9	0.8	0.9
1999.....	0.8	*	0.3	0.3	0.3	2.2	1.5	0.9	0.8	0.9
White, both sexes										
2002.....	1.6	*	0.3	0.3	0.5	4.0	3.6	2.5	1.9	1.8
2001.....	1.4	*	0.2	0.2	0.5	3.1	3.5	1.9	1.7	1.7
2000.....	1.4	*	0.2	0.2	0.4	3.2	3.4	2.3	1.6	1.5
1999.....	1.3	*	0.3	0.3	0.3	3.3	3.3	2.0	1.5	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)										
White, male										
2002.....	2.2	*	0.3	0.3	0.6	5.1	5.3	3.5	2.7	2.4
2001.....	2.0	*	*	*	0.5	3.7	5.5	2.8	2.6	2.4
2000.....	2.0	*	*	*	0.5	4.3	5.0	3.5	2.4	2.2
1999.....	1.9	*	*	0.3	0.3	4.2	4.9	3.1	2.2	2.1
White, female										
2002.....	0.9	*	*	0.3	0.4	2.8	1.7	1.3	1.1	1.1
2001.....	0.8	*	*	*	0.5	2.4	1.5	0.9	0.8	1.0
2000.....	0.7	*	*	0.3	0.3	2.1	1.6	1.0	0.7	0.9
1999.....	0.8	*	*	0.3	0.3	2.4	1.5	1.0	0.8	0.9
Black, both sexes										
2002.....	1.2	*	*	*	*	1.8	2.2	1.7	1.2	1.4
2001.....	1.1	*	*	*	*	1.3	2.0	1.8	1.7	1.0
2000.....	1.1	*	*	*	0.6	1.4	2.5	1.6	1.1	1.3
1999.....	1.2	*	*	*	*	1.7	2.2	1.6	1.4	1.6
Black, male										
2002.....	1.8	*	*	*	*	2.2	3.2	3.0	2.2	1.8
2001.....	1.6	*	*	*	*	1.8	2.4	2.5	3.2	1.5
2000.....	1.6	*	*	*	*	1.8	3.5	3.0	1.7	2.3
1999.....	1.7	*	*	*	*	2.2	2.9	2.8	2.0	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

Black, female

2002.....	0.6	*	*	*	*	1.4	*	*	*	*
2001.....	0.6	*	*	*	*	*	1.6	*	*	*
2000.....	0.6	*	*	*	*	*	1.5	*	*	*
1999.....	0.7	*	*	*	*	*	1.5	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of special-use motor vehicle involved in any accident (V83-V86)

All races 3/, both sexes

2002.....	0.3	*	*	0.2	0.5	0.6	0.5	0.4	0.3	0.4
2001.....	0.4	*	*	0.2	0.4	0.6	0.5	0.4	0.4	0.3
2000.....	0.4	*	*	0.2	0.4	0.7	0.5	0.4	0.3	0.3
1999.....	0.4	*	*	0.2	0.4	0.4	0.5	0.4	0.4	0.3

All races 3/, male

2002.....	0.6	*	*	0.3	0.7	0.9	0.8	0.7	0.5	0.7
2001.....	0.7	*	*	0.3	0.7	1.1	0.8	0.7	0.7	0.6
2000.....	0.7	*	*	0.3	0.6	1.1	0.9	0.7	0.6	0.6
1999.....	0.7	*	*	0.3	0.6	0.7	0.8	0.6	0.7	0.6

All races 3/, female

2002.....	0.1	*	*	*	0.2	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	0.2	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.4	*	*	0.2	0.6	0.7	0.6	0.5	0.4	0.5
2001.....	0.4	*	*	0.3	0.5	0.8	0.5	0.5	0.5	0.4
2000.....	0.4	*	*	0.3	0.5	0.7	0.6	0.4	0.4	0.4
1999.....	0.4	*	*	0.2	0.4	0.5	0.6	0.4	0.5	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)										
All races 3/, both sexes										
2002.....	5.5	1.1	1.3	1.0	1.3	9.9	9.9	6.6	5.5	5.1
2001.....	5.8	1.3	1.5	1.2	1.3	10.2	10.6	7.2	5.4	5.8
2000.....	6.0	1.8	1.6	1.3	1.6	10.7	11.4	7.2	5.9	5.8
1999.....	6.0	2.3	1.6	1.3	1.7	11.1	10.5	7.0	5.7	6.0
All races 3/, male										
2002.....	7.3	1.1	1.4	1.1	1.4	12.4	14.3	9.3	7.7	7.1
2001.....	7.7	1.1	1.6	1.3	1.4	13.3	15.5	10.4	7.5	7.7
2000.....	8.0	1.9	1.8	1.3	1.7	13.7	16.0	10.2	8.2	8.0
1999.....	7.9	2.1	1.8	1.4	1.8	13.8	15.0	9.9	7.6	8.0
All races 3/, female										
2002.....	3.7	1.1	1.3	0.9	1.1	7.3	5.4	3.8	3.1	3.2
2001.....	3.9	1.5	1.3	1.1	1.1	6.9	5.5	4.0	3.4	4.0
2000.....	4.1	1.7	1.4	1.3	1.4	7.5	6.6	4.0	3.6	3.6
1999.....	4.2	2.5	1.5	1.2	1.5	8.2	5.8	4.1	3.8	4.0
White, both sexes										
2002.....	5.6	0.8	1.3	0.9	1.3	10.9	10.3	6.6	5.6	5.1
2001.....	5.9	1.1	1.4	1.1	1.2	11.0	10.9	7.4	5.5	5.8
2000.....	6.2	1.5	1.6	1.3	1.6	11.8	11.9	7.1	5.8	5.9
1999.....	6.2	2.2	1.4	1.2	1.7	12.0	11.0	7.3	5.7	6.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)										
White, male										
2002.....	7.4	*	1.3	0.9	1.4	13.5	14.7	9.2	7.8	7.0
2001.....	7.8	*	1.5	1.2	1.2	14.2	15.9	10.6	7.6	7.4
2000.....	8.2	1.6	1.8	1.2	1.8	15.0	16.8	10.0	8.2	8.1
1999.....	8.0	2.2	1.4	1.3	1.8	14.8	15.6	10.3	7.5	8.1
White, female										
2002.....	3.9	*	1.3	0.9	1.1	8.0	5.6	3.8	3.2	3.1
2001.....	4.0	1.3	1.3	1.0	1.1	7.6	5.5	4.1	3.3	4.2
2000.....	4.2	1.4	1.4	1.3	1.5	8.4	6.6	4.0	3.4	3.7
1999.....	4.3	2.3	1.4	1.2	1.5	9.0	6.0	4.1	3.9	4.1
Black, both sexes										
2002.....	5.1	*	1.5	1.4	1.2	6.3	8.7	7.7	5.6	5.9
2001.....	5.6	*	1.7	1.8	1.7	6.7	10.0	7.4	5.9	6.4
2000.....	5.9	3.6	1.9	1.5	1.3	7.0	10.8	8.7	7.4	6.0
1999.....	5.8	*	2.6	1.7	1.6	7.9	9.4	6.9	6.0	6.4
Black, male										
2002.....	7.3	*	1.8	1.9	1.4	7.9	13.1	11.7	8.3	8.8
2001.....	8.0	*	1.8	2.2	2.2	9.3	14.9	11.3	8.2	9.7
2000.....	8.5	*	2.0	1.4	1.5	9.4	15.3	13.0	10.2	9.2
1999.....	8.2	*	3.1	2.0	1.9	10.5	14.1	10.2	9.0	8.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)										
Black, female										
2002.....	3.1	*	*	*	*	4.7	4.4	4.0	3.3	3.4
2001.....	3.4	*	*	1.3	1.2	3.9	5.4	3.8	3.8	3.5
2000.....	3.6	*	1.8	1.6	*	4.5	6.6	4.8	5.0	3.2
1999.....	3.7	*	2.1	1.4	1.4	5.4	4.9	3.9	3.2	4.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2002.....	0.2	*	*	*	0.2	0.1	0.2	0.1	0.2	0.2
2001.....	0.3	*	0.2	0.1	0.2	0.3	0.3	0.2	0.2	0.4
2000.....	0.4	*	0.2	0.2	0.2	0.4	0.4	0.3	0.4	0.3
1999.....	0.6	*	0.3	0.3	0.2	0.7	0.7	0.5	0.6	0.6

All races 3/, male

2002.....	0.3	*	*	*	0.2	0.2	0.4	*	*	0.4
2001.....	0.5	*	*	*	0.2	0.4	0.5	0.3	0.4	0.6
2000.....	0.5	*	*	*	0.3	0.5	0.6	0.5	0.5	0.5
1999.....	0.8	*	0.3	0.3	0.3	0.9	1.1	0.9	0.7	0.8

All races 3/, female

2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	0.2	*	*	0.2	*
1999.....	0.3	*	0.3	*	*	0.5	0.3	*	0.4	0.3

White, both sexes

2002.....	0.2	*	*	*	0.2	0.1	0.2	*	0.2	0.2
2001.....	0.3	*	0.2	*	0.2	0.3	0.3	0.2	0.2	0.3
2000.....	0.4	*	*	0.2	0.2	0.4	0.4	0.3	0.4	0.3
1999.....	0.6	*	0.3	0.2	0.3	0.7	0.8	0.6	0.6	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Water transport accidents (V90-V94)										
All races 3/, both sexes										
2002.....	0.2	*	*	*	*	0.2	0.3	0.2	0.3	0.3
2001.....	0.2	*	*	*	*	0.3	0.3	0.3	0.2	0.2
2000.....	0.2	*	*	*	*	0.3	0.3	0.3	0.3	0.3
1999.....	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.3
All races 3/, male										
2002.....	0.4	*	*	*	*	0.3	0.4	0.4	0.6	0.6
2001.....	0.4	*	*	*	*	0.5	0.5	0.5	0.4	0.4
2000.....	0.4	*	*	*	*	0.5	0.5	0.6	0.4	0.6
1999.....	0.4	*	*	*	0.2	0.4	0.6	0.5	0.5	0.5
All races 3/, female										
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.2	*	*	*	*	0.2	0.3	0.2	0.4	0.4
2001.....	0.2	*	*	*	*	0.3	0.3	0.3	0.2	0.2
2000.....	0.2	*	*	*	*	0.3	0.3	0.3	0.3	0.3
1999.....	0.2	*	*	*	0.1	0.3	0.4	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Air and space transport accidents (V95-V97)										
All races 3/, both sexes										
2002.....	0.2	*	*	*	*	*	0.2	0.1	0.2	0.3
2001.....	0.3	*	*	*	*	*	0.2	0.4	0.4	0.4
2000.....	0.3	*	*	*	*	0.1	0.3	0.3	0.3	0.3
1999.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.3
All races 3/, male										
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.3	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.5	0.7	0.7
2000.....	0.5	*	*	*	*	*	0.5	0.5	0.6	0.6
1999.....	0.4	*	*	*	*	*	0.3	0.5	0.5	0.5
All races 3/, female										
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	0.2	*	0.2
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.3	*	*	*	*	*	0.2	0.2	0.2	0.3
2001.....	0.4	*	*	*	*	*	0.3	0.4	0.5	0.5
2000.....	0.3	*	*	*	*	*	0.3	0.4	0.4	0.4
1999.....	0.3	*	*	*	*	*	0.2	0.4	0.4	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified transport accidents and their sequelae (V98-V99,Y85)										
All races 3/, both sexes										
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	0.2	*	*	*	*	0.1	0.1	0.2	0.2	0.2
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.3
1999.....	0.2	*	*	*	*	0.1	0.1	0.2	0.2	0.3
All races 3/, male										
2002.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.4
2001.....	0.3	*	*	*	*	*	0.2	0.2	0.4	0.3
2000.....	0.3	*	*	*	*	*	*	0.3	0.3	0.4
1999.....	0.3	*	*	*	*	*	*	0.3	0.3	0.4
All races 3/, female										
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	0.2	*	*	*	*	*	*	0.2	0.2	0.2
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.3
1999.....	0.2	*	*	*	*	*	*	0.2	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Falls (W00-W19)										
All races 3/, both sexes										
2002.....	5.6	*	0.2	*	0.1	0.4	0.8	0.7	0.8	1.2
2001.....	5.3	0.6	0.2	0.2	0.2	0.4	0.9	0.8	0.9	1.2
2000.....	4.7	*	0.2	*	0.1	0.5	0.7	0.7	0.8	1.0
1999.....	4.7	*	0.4	0.1	0.1	0.6	0.7	0.8	0.9	1.2
All races 3/, male										
2002.....	6.0	*	0.4	*	*	0.6	1.4	1.2	1.4	1.7
2001.....	5.8	*	0.3	0.2	0.2	0.8	1.5	1.5	1.5	2.1
2000.....	5.2	*	0.3	*	*	0.8	1.2	1.2	1.5	1.8
1999.....	5.2	*	0.5	*	0.2	0.9	1.2	1.4	1.5	2.0
All races 3/, female										
2002.....	5.3	*	*	*	*	*	*	*	0.3	0.6
2001.....	4.8	*	*	*	*	*	*	*	0.3	0.4
2000.....	4.3	*	*	*	*	*	0.2	*	*	0.3
1999.....	4.3	*	*	*	*	*	*	*	0.3	0.4
White, both sexes										
2002.....	6.4	*	0.2	*	*	0.4	0.9	0.8	0.9	1.2
2001.....	6.0	*	0.2	0.2	0.2	0.5	1.0	0.9	0.9	1.3
2000.....	5.3	*	0.2	*	*	0.6	0.8	0.8	0.9	1.0
1999.....	5.2	*	0.4	*	0.2	0.6	0.8	0.9	0.9	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Falls (W00-W19)										
White, male										
2002.....	6.6	*	*	*	*	0.7	1.6	1.4	1.5	1.8
2001.....	6.4	*	0.3	0.2	0.2	0.9	1.8	1.6	1.6	2.1
2000.....	5.7	*	*	*	*	1.0	1.4	1.4	1.5	1.7
1999.....	5.6	*	0.5	*	*	1.1	1.3	1.5	1.5	2.0
White, female										
2002.....	6.2	*	*	*	*	*	*	*	0.3	0.7
2001.....	5.5	*	*	*	*	*	*	*	0.3	0.4
2000.....	5.0	*	*	*	*	*	0.3	*	*	0.3
1999.....	4.9	*	*	*	*	*	*	*	0.3	0.5
Black, both sexes										
2002.....	2.2	*	*	*	*	*	*	*	*	1.1
2001.....	2.3	*	*	*	*	*	*	*	*	1.1
2000.....	2.1	*	*	*	*	*	*	*	*	1.0
1999.....	2.4	*	*	*	*	*	*	*	1.1	0.9
Black, male										
2002.....	3.0	*	*	*	*	*	*	*	*	1.8
2001.....	3.1	*	*	*	*	*	*	*	*	2.0
2000.....	2.8	*	*	*	*	*	*	*	*	1.7
1999.....	3.3	*	*	*	*	*	*	*	1.8	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Fall from one level to another (W10-W17)										
All races 3/, both sexes										
2002.....	1.2	*	0.1	*	*	0.3	0.6	0.5	0.5	0.7
2001.....	1.2	*	0.1	*	*	0.3	0.6	0.6	0.5	0.7
2000.....	1.0	*	0.2	*	*	0.4	0.5	0.5	0.6	0.6
1999.....	1.1	*	0.2	*	*	0.4	0.4	0.6	0.6	0.6
All races 3/, male										
2002.....	1.7	*	*	*	*	0.4	1.0	0.9	0.9	1.1
2001.....	1.7	*	*	*	*	0.5	1.1	1.1	0.9	1.2
2000.....	1.6	*	*	*	*	0.6	0.9	0.9	1.1	1.0
1999.....	1.7	*	0.3	*	*	0.6	0.8	1.0	1.0	1.1
All races 3/, female										
2002.....	0.6	*	*	*	*	*	*	*	*	0.2
2001.....	0.6	*	*	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	1.3	*	*	*	*	0.3	0.7	0.6	0.6	0.7
2001.....	1.3	*	*	*	*	0.4	0.7	0.7	0.6	0.7
2000.....	1.1	*	*	*	*	0.4	0.6	0.6	0.7	0.6
1999.....	1.2	*	0.2	*	*	0.4	0.5	0.6	0.6	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Unspecified fall (W19)										
All races 3/, both sexes										
2002.....	2.7	*	*	*	*	*	*	*	0.1	0.3
2001.....	2.7	*	*	*	*	*	0.1	0.1	0.2	0.3
2000.....	2.5	*	*	*	*	*	*	0.2	0.1	0.3
1999.....	2.8	*	*	*	*	0.1	0.2	0.1	0.3	0.5
All races 3/, male										
2002.....	2.5	*	*	*	*	*	*	*	0.3	0.4
2001.....	2.6	*	*	*	*	*	*	*	0.3	0.5
2000.....	2.5	*	*	*	*	*	*	0.3	0.2	0.5
1999.....	2.8	*	*	*	*	*	0.3	0.2	0.4	0.7
All races 3/, female										
2002.....	2.8	*	*	*	*	*	*	*	*	0.2
2001.....	2.7	*	*	*	*	*	*	*	*	*
2000.....	2.6	*	*	*	*	*	*	*	*	*
1999.....	2.8	*	*	*	*	*	*	*	*	0.2
White, both sexes										
2002.....	3.0	*	*	*	*	*	*	*	*	0.3
2001.....	3.1	*	*	*	*	*	*	*	0.2	0.3
2000.....	2.9	*	*	*	*	*	*	0.2	0.1	0.3
1999.....	3.1	*	*	*	*	*	0.2	0.2	0.3	0.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental discharge of firearms (W32-W34)										
All races 3/, both sexes										
2002.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.2
2001.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.2	0.3
2000.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.3
1999.....	0.3	*	*	*	0.3	0.6	0.7	0.4	0.3	0.3
All races 3/, male										
2002.....	0.5	*	*	*	0.3	1.0	0.9	0.7	0.6	0.4
2001.....	0.5	*	*	*	0.3	1.0	0.9	0.7	0.4	0.5
2000.....	0.5	*	*	*	0.4	0.9	0.9	0.7	0.4	0.5
1999.....	0.5	*	*	*	0.5	1.1	1.2	0.6	0.6	0.5
All races 3/, female										
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.3	*	*	*	0.2	0.4	0.5	0.3	0.3	0.2
2001.....	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.4
2000.....	0.3	*	*	*	0.3	0.5	0.4	0.3	0.3	0.3
1999.....	0.3	*	*	*	0.3	0.6	0.6	0.4	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental discharge of firearms (W32-W34)

White, male

2002.....	0.4	*	*	*	0.3	0.8	0.8	0.6	0.5	0.4
2001.....	0.5	*	*	*	0.4	0.9	0.9	0.6	0.5	0.6
2000.....	0.5	*	*	*	0.5	0.9	0.8	0.6	0.4	0.5
1999.....	0.5	*	*	*	0.5	1.0	1.0	0.6	0.6	0.5

White, female

2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

Black, both sexes

2002.....	0.4	*	*	*	*	1.1	0.8	0.9	*	*
2001.....	0.3	*	*	*	*	0.8	0.8	*	*	*
2000.....	0.4	*	*	*	*	0.9	0.9	*	*	*
1999.....	0.4	*	*	*	*	1.2	1.1	*	*	*

Black, male

2002.....	0.8	*	*	*	*	2.2	1.5	1.8	*	*
2001.....	0.6	*	*	*	*	1.6	1.5	*	*	*
2000.....	0.7	*	*	*	*	1.6	1.5	*	*	*
1999.....	0.8	*	*	*	*	2.1	2.1	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
All races 3/, both sexes										
2002.....	1.2	1.6	2.9	0.8	0.8	1.6	1.5	1.2	1.0	1.2
2001.....	1.2	1.7	3.0	0.8	0.8	1.6	1.4	1.1	0.8	1.0
2000.....	1.2	2.0	3.2	1.0	0.8	1.8	1.5	1.1	1.0	1.1
1999.....	1.3	1.8	3.2	0.9	0.9	1.8	1.5	1.1	1.1	1.1
All races 3/, male										
2002.....	1.9	1.9	3.9	1.2	1.1	2.8	2.8	2.1	1.7	2.0
2001.....	1.8	1.5	3.7	1.2	1.1	2.8	2.5	1.9	1.4	1.6
2000.....	2.0	2.2	4.0	1.3	1.2	3.3	2.5	1.9	1.7	1.9
1999.....	2.0	2.0	4.0	1.3	1.3	3.2	2.8	2.0	2.0	1.9
All races 3/, female										
2002.....	0.5	1.2	1.9	0.3	0.4	0.3	*	0.3	0.3	0.4
2001.....	0.5	1.9	2.2	0.5	0.5	0.3	0.3	0.3	0.2	0.4
2000.....	0.5	1.7	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3
1999.....	0.5	1.6	2.4	0.5	0.4	0.3	0.3	*	0.3	0.3
White, both sexes										
2002.....	1.1	1.6	3.0	0.6	0.5	1.4	1.5	1.1	1.0	1.1
2001.....	1.1	1.7	3.3	0.6	0.5	1.3	1.4	1.0	0.7	1.0
2000.....	1.2	1.6	3.5	0.8	0.6	1.6	1.4	1.1	1.0	1.1
1999.....	1.2	1.5	3.5	0.7	0.5	1.6	1.5	1.0	1.1	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
White, male										
2002.....	1.8	1.9	4.0	0.9	0.6	2.4	2.7	1.9	1.7	1.9
2001.....	1.7	1.6	4.2	0.8	0.6	2.3	2.4	1.7	1.2	1.5
2000.....	1.9	1.8	4.4	1.0	0.8	2.8	2.4	1.8	1.7	1.8
1999.....	1.9	1.7	4.3	0.9	0.7	2.8	2.5	1.7	1.9	1.8
White, female										
2002.....	0.4	*	2.0	0.3	0.3	0.3	*	0.3	*	0.4
2001.....	0.5	1.8	2.4	0.5	0.3	*	0.3	0.3	*	0.4
2000.....	0.5	*	2.5	0.5	0.3	0.3	0.3	0.3	*	0.4
1999.....	0.5	1.4	2.6	0.4	*	0.3	0.3	*	0.3	0.4
Black, both sexes										
2002.....	1.6	*	2.5	1.8	2.2	2.3	1.9	1.6	1.2	1.5
2001.....	1.6	*	1.9	1.7	2.3	2.9	1.5	1.2	1.1	1.2
2000.....	1.6	4.2	2.2	1.9	2.1	2.8	1.7	1.3	0.9	1.2
1999.....	1.8	3.5	2.4	2.2	2.6	2.7	1.9	1.8	1.4	1.1
Black, male										
2002.....	2.8	*	3.4	2.8	3.5	4.3	3.7	3.1	2.0	2.8
2001.....	2.7	*	2.2	2.7	3.2	5.2	3.1	2.6	2.2	2.2
2000.....	2.7	*	2.8	2.7	3.2	5.2	3.3	2.3	1.8	2.3
1999.....	3.1	*	3.2	3.4	3.9	5.0	3.8	3.5	2.7	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)										
All races 3/, both sexes										
2002.....	1.4	1.5	0.5	*	*	*	0.2	0.2	0.2	0.3
2001.....	1.5	1.5	0.5	*	*	*	0.2	0.1	0.2	0.3
2000.....	1.5	2.0	0.6	*	*	0.1	0.1	0.2	0.3	0.3
1999.....	1.4	1.7	0.6	0.1	0.1	*	0.1	0.1	0.3	0.3
All races 3/, male										
2002.....	1.5	2.0	0.6	*	*	*	0.2	0.3	0.3	0.4
2001.....	1.5	2.0	0.6	*	*	*	0.2	*	0.2	0.4
2000.....	1.6	2.3	0.7	*	*	*	*	0.2	0.4	0.5
1999.....	1.4	1.9	0.8	*	*	*	*	*	0.3	0.3
All races 3/, female										
2002.....	1.4	1.0	0.3	*	*	*	*	*	*	0.3
2001.....	1.4	1.0	0.4	*	*	*	*	*	*	0.3
2000.....	1.5	1.8	0.5	*	*	*	*	*	0.2	0.2
1999.....	1.4	1.4	0.4	*	*	*	*	*	*	0.2
White, both sexes										
2002.....	1.5	1.4	0.4	*	*	*	0.1	0.2	0.2	0.3
2001.....	1.5	1.3	0.5	*	*	*	0.2	*	0.2	0.3
2000.....	1.6	1.7	0.5	*	*	*	*	0.2	0.3	0.4
1999.....	1.4	1.4	0.5	*	*	*	0.1	*	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other accidental and unspecified threats to breathing (W75-W77,W81-W84)										
All races 3/, both sexes										
2002.....	0.5	14.3	0.4	0.1	0.2	0.2	0.3	0.2	0.3	0.3
2001.....	0.5	13.7	0.4	0.1	0.2	0.2	0.2	0.2	0.3	0.4
2000.....	0.5	11.8	0.4	0.2	0.3	0.2	0.3	0.2	0.3	0.4
1999.....	0.6	10.8	0.5	0.1	0.3	0.2	0.3	0.3	0.2	0.3
All races 3/, male										
2002.....	0.7	15.7	0.5	*	0.4	0.4	0.5	0.4	0.5	0.5
2001.....	0.7	15.3	0.5	0.2	0.4	0.4	0.4	0.4	0.5	0.6
2000.....	0.7	13.2	0.5	*	0.5	0.4	0.5	0.4	0.5	0.7
1999.....	0.8	12.6	0.5	0.2	0.5	0.4	0.5	0.5	0.4	0.5
All races 3/, female										
2002.....	0.3	12.8	0.4	*	*	*	*	*	*	*
2001.....	0.3	12.0	0.3	*	*	*	*	*	*	*
2000.....	0.3	10.3	0.3	*	*	*	*	*	*	*
1999.....	0.4	8.9	0.5	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.4	10.7	0.4	*	0.3	0.3	0.3	0.3	0.3	0.4
2001.....	0.5	10.8	0.4	0.2	0.3	0.2	0.3	0.3	0.3	0.4
2000.....	0.5	9.4	0.4	0.2	0.3	0.3	0.3	0.3	0.3	0.4
1999.....	0.6	8.7	0.5	0.1	0.3	0.3	0.3	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to electric current (W85-W87)										
All races 3/, both sexes										
2002.....	0.1	*	*	*	*	0.2	0.2	0.3	0.2	0.2
2001.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.3
2000.....	0.1	*	*	*	*	0.1	0.2	0.2	0.2	0.2
1999.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.3
All races 3/, male										
2002.....	0.3	*	*	*	*	0.3	0.4	0.5	0.4	0.5
2001.....	0.3	*	*	*	*	*	0.4	0.4	0.5	0.5
2000.....	0.3	*	*	*	*	0.2	0.5	0.4	0.4	0.5
1999.....	0.3	*	*	*	*	0.3	0.5	0.4	0.4	0.5
All races 3/, female										
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.2	*	*	*	*	0.2	0.3	0.3	0.3	0.3
2001.....	0.2	*	*	*	*	*	0.2	0.2	0.3	0.3
2000.....	0.2	*	*	*	*	0.2	0.3	0.3	0.3	0.3
1999.....	0.2	*	*	*	*	0.2	0.3	0.3	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
All races 3/, both sexes										
2002.....	1.1	0.9	1.4	0.8	0.5	0.4	0.5	0.6	0.7	0.7
2001.....	1.2	1.2	1.5	0.8	0.4	0.4	0.7	0.6	0.6	0.9
2000.....	1.2	1.0	1.9	0.9	0.4	0.4	0.6	0.6	0.6	0.8
1999.....	1.2	1.1	2.0	0.8	0.5	0.4	0.6	0.7	0.7	0.9
All races 3/, male										
2002.....	1.4	1.0	1.6	0.9	0.6	0.6	0.7	0.8	0.9	0.9
2001.....	1.4	1.5	1.8	1.0	0.4	0.4	0.9	0.8	0.8	1.2
2000.....	1.5	1.0	2.2	1.0	0.3	0.4	0.8	0.8	0.8	1.2
1999.....	1.5	1.1	2.2	0.9	0.5	0.6	0.8	0.9	1.0	1.2
All races 3/, female										
2002.....	0.8	*	1.2	0.6	0.4	0.3	0.4	0.4	0.6	0.4
2001.....	0.9	1.0	1.1	0.6	0.4	0.3	0.4	0.4	0.4	0.5
2000.....	0.9	*	1.5	0.7	0.5	0.4	0.4	0.4	0.4	0.5
1999.....	0.9	1.1	1.7	0.7	0.4	0.3	0.4	0.5	0.4	0.5
White, both sexes										
2002.....	1.0	*	1.0	0.5	0.4	0.4	0.5	0.5	0.7	0.6
2001.....	1.0	1.1	1.2	0.7	0.4	0.3	0.7	0.5	0.6	0.8
2000.....	1.0	0.8	1.3	0.7	0.3	0.4	0.5	0.5	0.5	0.7
1999.....	1.0	0.9	1.5	0.6	0.4	0.4	0.6	0.7	0.7	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
White, male										
2002.....	1.2	*	1.1	0.6	0.5	0.5	0.6	0.7	0.9	0.8
2001.....	1.3	*	1.6	0.8	0.4	0.4	0.9	0.7	0.8	1.0
2000.....	1.3	*	1.6	0.7	0.3	0.4	0.8	0.7	0.7	1.0
1999.....	1.3	*	1.7	0.6	0.4	0.6	0.7	0.9	1.0	1.0
White, female										
2002.....	0.7	*	1.0	0.4	*	*	0.3	0.3	0.5	0.4
2001.....	0.8	*	0.8	0.5	0.3	0.3	0.4	*	0.4	0.5
2000.....	0.8	*	1.1	0.6	0.4	0.4	0.3	0.4	0.3	0.4
1999.....	0.8	*	1.3	0.6	0.3	0.3	0.4	0.4	0.3	0.5
Black, both sexes										
2002.....	2.1	*	3.5	1.9	0.9	0.7	0.9	1.2	1.1	1.0
2001.....	2.3	*	3.3	1.8	0.8	*	1.0	1.4	1.0	1.5
2000.....	2.5	*	4.8	2.1	0.8	*	1.0	1.1	1.2	1.5
1999.....	2.5	*	4.5	2.1	0.9	*	0.9	1.1	0.9	1.9
Black, male										
2002.....	2.6	*	4.1	2.2	*	*	*	*	*	1.5
2001.....	2.9	*	3.6	2.3	*	*	*	*	*	2.5
2000.....	3.2	*	5.6	2.6	*	*	*	*	1.5	2.2
1999.....	3.0	*	5.0	2.4	*	*	*	*	*	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental exposure to smoke, fire and flames (X00-X09)

Black, female

2002.....	1.6	*	2.8	1.7	*	*	*	*	*	*
2001.....	1.7	*	2.9	1.2	*	*	*	1.5	*	*
2000.....	2.0	*	4.1	1.6	*	*	*	*	*	*
1999.....	2.0	*	3.9	1.8	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
All races 3/, both sexes										
2002.....	6.1	0.6	0.2	*	0.1	2.4	5.9	6.9	8.6	11.9
2001.....	4.9	*	0.2	*	0.2	2.0	4.9	5.7	6.9	10.4
2000.....	4.5	*	0.2	*	0.1	1.7	4.3	5.1	6.8	9.5
1999.....	4.4	*	0.2	*	0.1	1.3	3.8	4.9	6.8	9.1
All races 3/, male										
2002.....	8.5	1.0	*	*	*	3.5	9.1	10.4	12.5	16.5
2001.....	7.1	*	0.3	*	*	3.0	7.6	8.7	9.9	14.6
2000.....	6.6	*	0.3	*	*	2.6	6.7	7.9	10.1	13.7
1999.....	6.5	*	*	*	*	1.9	5.9	7.7	10.3	13.3
All races 3/, female										
2002.....	3.7	*	*	*	*	1.2	2.6	3.3	4.7	7.2
2001.....	2.9	*	*	*	*	1.0	1.9	2.6	3.9	6.1
2000.....	2.5	*	*	*	*	0.8	1.7	2.1	3.5	5.3
1999.....	2.3	*	*	*	*	0.7	1.6	2.0	3.2	4.9
White, both sexes										
2002.....	6.3	*	*	*	0.2	2.8	6.9	7.8	9.5	12.5
2001.....	5.0	*	0.2	*	0.2	2.3	5.6	6.4	7.4	10.8
2000.....	4.6	*	*	*	0.1	2.0	4.9	5.5	7.2	9.8
1999.....	4.3	*	0.2	*	0.1	1.5	4.2	5.2	6.9	9.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
White, male										
2002.....	8.8	*	*	*	*	4.1	10.6	11.6	13.6	17.3
2001.....	7.2	*	*	*	*	3.5	8.7	9.5	10.7	15.2
2000.....	6.7	*	*	*	*	3.1	7.8	8.6	10.7	14.3
1999.....	6.4	*	*	*	*	2.1	6.5	8.3	10.6	13.7
White, female										
2002.....	3.9	*	*	*	*	1.4	3.0	3.8	5.1	7.5
2001.....	2.9	*	*	*	*	1.0	2.3	3.0	3.9	6.2
2000.....	2.5	*	*	*	*	0.9	1.8	2.3	3.6	5.2
1999.....	2.3	*	*	*	*	0.8	1.7	2.1	3.1	4.8
Black, both sexes										
2002.....	6.3	*	*	*	*	*	2.3	4.4	6.8	11.9
2001.....	5.8	*	*	*	*	0.9	2.3	4.0	6.3	11.1
2000.....	5.5	*	*	*	*	0.7	2.1	4.3	6.8	10.7
1999.....	5.7	*	*	*	*	*	2.8	4.2	8.0	10.3
Black, male										
2002.....	9.1	*	*	*	*	*	3.5	7.0	9.9	17.0
2001.....	8.4	*	*	*	*	*	3.8	6.7	8.2	15.4
2000.....	8.2	*	*	*	*	*	2.7	7.1	9.6	14.9
1999.....	8.7	*	*	*	*	*	4.7	6.7	11.5	14.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
Black, female										
2002.....	3.7	*	*	*	*	*	*	2.0	4.0	7.4
2001.....	3.5	*	*	*	*	*	*	1.6	4.6	7.3
2000.....	3.1	*	*	*	*	*	1.5	1.7	4.3	7.1
1999.....	3.0	*	*	*	*	*	*	2.0	4.9	6.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

All races 3/, both sexes

2002.....	5.7	*	*	*	*	2.2	5.6	6.6	8.2	11.4
2001.....	4.6	*	*	*	*	1.7	4.5	5.4	6.6	9.8
2000.....	4.2	*	*	*	*	1.4	4.0	4.7	6.5	9.1
1999.....	4.0	*	*	*	*	1.0	3.4	4.6	6.4	8.6

All races 3/, male

2002.....	7.9	*	*	*	*	3.2	8.6	9.9	11.8	15.7
2001.....	6.5	*	*	*	*	2.6	7.1	8.2	9.5	13.8
2000.....	6.1	*	*	*	*	2.1	6.2	7.5	9.4	13.0
1999.....	5.9	*	*	*	*	1.5	5.4	7.2	9.8	12.5

All races 3/, female

2002.....	3.5	*	*	*	*	1.1	2.5	3.2	4.6	7.0
2001.....	2.7	*	*	*	*	0.8	1.8	2.5	3.7	5.8
2000.....	2.3	*	*	*	*	0.7	1.6	2.0	3.4	5.2
1999.....	2.1	*	*	*	*	0.5	1.5	2.0	3.0	4.7

White, both sexes

2002.....	5.9	*	*	*	*	2.6	6.6	7.4	9.1	12.0
2001.....	4.7	*	*	*	*	2.0	5.2	6.0	7.1	10.2
2000.....	4.2	*	*	*	*	1.7	4.6	5.2	6.8	9.3
1999.....	4.0	*	*	*	*	1.2	3.8	5.0	6.6	8.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)										
White, male										
2002.....	8.2	*	*	*	*	3.7	10.0	11.0	13.0	16.5
2001.....	6.6	*	*	*	*	3.1	8.1	9.1	10.3	14.4
2000.....	6.1	*	*	*	*	2.5	7.3	8.1	10.0	13.5
1999.....	5.9	*	*	*	*	1.7	5.9	7.8	10.1	12.9
White, female										
2002.....	3.7	*	*	*	*	1.3	2.9	3.7	5.0	7.3
2001.....	2.7	*	*	*	*	0.9	2.2	2.8	3.8	6.0
2000.....	2.3	*	*	*	*	0.8	1.7	2.2	3.5	5.1
1999.....	2.1	*	*	*	*	0.7	1.6	2.1	3.0	4.7
Black, both sexes										
2002.....	5.8	*	*	*	*	*	2.2	4.2	6.4	11.2
2001.....	5.4	*	*	*	*	*	2.1	3.8	6.1	10.5
2000.....	5.2	*	*	*	*	*	2.0	3.9	6.6	10.3
1999.....	5.3	*	*	*	*	*	2.5	4.2	7.6	9.8
Black, male										
2002.....	8.4	*	*	*	*	*	3.4	6.7	9.3	15.8
2001.....	7.8	*	*	*	*	*	3.5	6.3	8.0	14.5
2000.....	7.7	*	*	*	*	*	2.6	6.6	9.3	14.0
1999.....	8.1	*	*	*	*	*	4.2	6.7	10.9	13.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/ year 2/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Black, female

2002.....	3.5	*	*	*	*	*	*	2.0	3.9	7.1
2001.....	3.2	*	*	*	*	*	*	1.4	4.4	6.9
2000.....	2.9	*	*	*	*	*	1.4	1.5	4.2	7.1
1999.....	2.8	*	*	*	*	*	*	2.0	4.6	6.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)

All races 3/, both sexes

2002.....	0.2	*	*	*	*	0.1	*	0.2	0.2	0.2
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.1	0.3
2000.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.2
1999.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.3

All races 3/, male

2002.....	0.3	*	*	*	*	*	*	0.3	0.4	0.4
2001.....	0.2	*	*	*	*	*	*	*	0.2	0.5
2000.....	0.2	*	*	*	*	*	*	*	0.3	0.4
1999.....	0.3	*	*	*	*	*	0.2	0.2	0.2	0.4

All races 3/, female

2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.2	*	*	*	*	0.1	*	0.2	0.2	0.2
2001.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3
2000.....	0.2	*	*	*	*	*	*	*	0.1	0.2
1999.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning by and exposure to other gases and vapors (X47)										
All races 3/, both sexes										
2002.....	0.2	*	*	*	*	0.1	0.2	0.2	0.2	0.3
2001.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.2
2000.....	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.2
1999.....	0.2	*	*	*	*	0.1	0.2	*	0.2	0.2
All races 3/, male										
2002.....	0.3	*	*	*	*	*	0.3	0.3	0.3	0.4
2001.....	0.3	*	*	*	*	0.2	0.4	0.3	0.2	0.4
2000.....	0.3	*	*	*	*	0.3	0.3	0.3	0.4	0.4
1999.....	0.3	*	*	*	*	0.2	0.3	*	0.3	0.4
All races 3/, female										
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.2	*	*	*	*	0.1	0.2	0.2	0.2	0.2
2001.....	0.2	*	*	*	*	0.2	0.3	0.2	0.2	0.2
2000.....	0.2	*	*	*	*	0.2	0.2	0.2	0.3	0.2
1999.....	0.2	*	*	*	*	0.2	0.2	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)										
All races 3/, both sexes										
2002.....	3.8	0.9	0.6	0.3	0.4	0.8	1.0	1.1	1.3	1.6
2001.....	4.0	1.3	0.8	0.3	0.3	0.8	1.2	1.1	1.3	1.8
2000.....	3.9	1.1	0.7	0.3	0.3	0.8	1.2	1.2	1.4	1.9
1999.....	4.3	1.2	0.8	0.4	0.4	0.9	1.2	1.3	1.4	1.7
All races 3/, male										
2002.....	4.3	1.2	0.7	0.4	0.4	1.2	1.8	1.8	2.3	2.6
2001.....	4.5	1.6	0.9	0.4	0.4	1.2	2.1	2.0	2.3	3.0
2000.....	4.5	1.2	0.8	0.4	0.4	1.3	2.2	2.2	2.4	3.3
1999.....	4.8	1.4	1.0	0.5	0.5	1.4	2.0	2.3	2.3	2.9
All races 3/, female										
2002.....	3.2	*	0.5	*	0.3	0.3	0.2	0.4	0.3	0.6
2001.....	3.5	1.1	0.7	0.2	*	0.4	0.3	0.3	0.4	0.6
2000.....	3.3	*	0.5	0.2	*	0.3	0.2	0.2	0.4	0.5
1999.....	3.7	*	0.5	0.3	0.2	0.4	0.3	0.4	0.4	0.6
White, both sexes										
2002.....	4.1	0.8	0.5	0.3	0.3	0.8	1.1	1.2	1.3	1.5
2001.....	4.3	1.3	0.8	0.3	0.3	0.8	1.3	1.2	1.3	1.8
2000.....	4.2	1.0	0.6	0.3	0.3	0.8	1.3	1.3	1.4	1.9
1999.....	4.5	1.0	0.8	0.4	0.4	0.9	1.2	1.3	1.4	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)										
White, male										
2002.....	4.5	*	0.6	0.5	0.3	1.2	2.0	2.0	2.3	2.5
2001.....	4.8	1.5	0.9	0.3	0.4	1.3	2.2	2.0	2.3	3.1
2000.....	4.6	*	0.7	0.4	0.4	1.3	2.2	2.3	2.5	3.2
1999.....	5.0	*	1.0	0.5	0.5	1.4	2.1	2.3	2.3	2.9
White, female										
2002.....	3.6	*	0.4	*	0.3	0.3	*	0.4	0.3	0.5
2001.....	3.9	*	0.6	*	*	0.4	0.3	*	0.4	0.5
2000.....	3.7	*	0.5	*	*	0.3	*	0.3	0.4	0.5
1999.....	4.1	*	0.5	0.3	0.3	0.4	0.3	0.3	0.4	0.6
Black, both sexes										
2002.....	2.8	*	1.0	*	*	0.8	*	0.9	1.4	2.3
2001.....	2.9	*	0.9	*	*	*	0.8	1.3	1.3	1.8
2000.....	3.0	*	0.9	*	*	0.7	1.2	1.1	1.5	2.4
1999.....	3.6	*	0.8	*	*	0.8	1.1	1.5	1.6	1.7
Black, male										
2002.....	3.7	*	*	*	*	*	*	*	2.6	4.1
2001.....	3.7	*	*	*	*	*	*	2.0	2.4	2.7
2000.....	4.4	*	*	*	*	*	2.2	2.2	2.5	4.4
1999.....	4.9	*	*	*	*	1.4	2.1	2.4	2.8	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
All races 3/, both sexes										
2002.....	11.0	*	1.2	7.4	12.4	12.8	12.5	14.3
2001 4/.....	10.8	*	1.3	7.9	12.0	12.6	13.0	14.3
2000.....	10.4	*	1.5	8.0	12.5	12.1	12.0	13.7
1999.....	10.5	*	1.2	8.0	12.3	12.3	13.1	14.4
All races 3/, male										
2002.....	17.9	*	1.8	12.2	20.8	20.8	20.2	22.7
2001 4/.....	17.6	*	1.9	12.9	20.5	21.0	21.0	22.7
2000.....	17.1	*	2.3	13.0	21.4	20.0	19.2	21.7
1999.....	17.1	*	1.9	13.0	20.8	20.3	21.1	22.7
All races 3/, female										
2002.....	4.3	*	0.6	2.4	3.5	4.5	4.7	6.0
2001 4/.....	4.1	*	0.6	2.7	3.1	4.0	4.8	5.9
2000.....	4.0	*	0.6	2.7	3.2	4.0	4.6	5.8
1999.....	4.0	*	0.5	2.7	3.4	4.1	5.1	6.1
White, both sexes										
2002.....	12.2	*	1.3	8.2	13.2	13.5	13.7	15.9
2001 4/.....	11.9	*	1.3	8.6	12.6	13.5	14.2	15.8
2000.....	11.5	*	1.5	8.5	13.1	12.7	13.0	15.1
1999.....	11.5	*	1.3	8.5	12.7	12.8	14.2	15.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
White, male										
2002.....	19.9	*	1.9	13.4	22.0	21.4	21.7	24.7
2001 4/.....	19.5	*	2.0	14.0	21.3	22.2	22.6	24.7
2000.....	18.8	*	2.4	13.7	22.3	20.8	20.7	23.6
1999.....	18.7	*	2.0	13.7	21.5	21.0	22.5	24.3
White, female										
2002.....	4.8	*	0.7	2.6	3.7	5.1	5.3	6.8
2001 4/.....	4.6	*	0.6	2.9	3.3	4.3	5.4	6.7
2000.....	4.4	*	0.7	2.9	3.3	4.1	5.1	6.5
1999.....	4.5	*	0.5	2.9	3.5	4.3	5.5	6.9
Black, both sexes										
2002.....	5.1	*	1.0	4.0	9.1	10.3	8.2	8.1
2001 4/.....	5.3	*	1.1	4.3	10.1	9.6	8.6	7.8
2000.....	5.4	*	1.2	5.5	11.0	9.9	7.6	7.3
1999.....	5.4	*	0.9	5.8	10.7	9.6	8.7	8.4
Black, male										
2002.....	9.1	*	1.6	6.9	16.1	19.5	15.2	14.5
2001 4/.....	9.2	*	1.7	7.3	19.4	17.7	15.4	13.8
2000.....	9.4	*	2.1	9.5	19.5	17.5	13.6	13.0
1999.....	9.6	*	1.3	9.9	19.5	17.9	15.7	15.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
Black, female										
2002.....	1.5	*	*	*	2.3	2.0	2.0	2.5
2001 4/.....	1.7	*	*	1.3	*	2.2	2.5	2.5
2000.....	1.7	*	*	1.4	3.0	3.0	2.2	2.2
1999.....	1.6	*	*	1.6	2.3	2.1	2.5	2.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

All races 3/, both sexes

2002.....	1.3	*	*	0.4	0.7	1.1	1.6	2.2
2001.....	1.2	*	*	0.4	0.7	1.0	1.5	2.1
2000.....	1.2	*	*	0.4	0.7	1.0	1.3	1.8
1999.....	1.1	*	*	0.3	0.7	1.0	1.3	1.9

All races 3/, male

2002.....	1.3	*	*	0.4	0.7	1.3	1.7	2.2
2001.....	1.2	*	*	0.4	0.7	1.1	1.6	2.4
2000.....	1.1	*	*	0.4	0.7	0.9	1.5	1.9
1999.....	1.1	*	*	0.3	0.7	1.0	1.2	2.0

All races 3/, female

2002.....	1.4	*	*	0.4	0.8	1.0	1.4	2.1
2001.....	1.3	*	*	0.5	0.8	0.9	1.4	1.8
2000.....	1.2	*	*	0.4	0.6	1.1	1.2	1.8
1999.....	1.2	*	*	0.4	0.6	1.1	1.4	1.8

White, both sexes

2002.....	1.5	*	*	0.4	0.9	1.3	1.7	2.5
2001.....	1.4	*	*	0.5	0.8	1.1	1.7	2.4
2000.....	1.3	*	*	0.4	0.7	1.1	1.5	2.1
1999.....	1.3	*	*	0.4	0.7	1.1	1.4	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

White, male

2002.....	1.5	*	*	0.5	0.8	1.4	1.9	2.5
2001.....	1.4	*	*	0.5	0.8	1.2	1.8	2.7
2000.....	1.2	*	*	0.4	0.7	1.0	1.7	2.1
1999.....	1.3	*	*	0.3	0.8	1.1	1.3	2.2

White, female

2002.....	1.6	*	*	0.4	0.9	1.2	1.5	2.4
2001.....	1.5	*	*	0.5	0.9	1.0	1.6	2.1
2000.....	1.4	*	*	0.4	0.7	1.2	1.4	2.0
1999.....	1.3	*	*	0.5	0.7	1.2	1.5	2.1

Black, both sexes

2002.....	0.4	*	*	*	*	*	1.0	0.8
2001.....	0.4	*	*	*	*	*	*	1.0
2000.....	0.4	*	*	*	*	*	*	0.8
1999.....	0.4	*	*	*	*	*	0.8	0.7

Black, male

2002.....	0.4	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

Black, female

2002.....	0.4	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

All races 3/, both sexes

2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.1	*	0.1
2000.....	0.1	*	*	*	*	*	*	0.1
1999.....	0.1	*	*	*	*	*	*	0.1

All races 3/, male

2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	0.2
1999.....	0.1	*	*	*	*	*	*	*

All races 3/, female

2002.....	0.0	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*

White, both sexes

2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	0.1
1999.....	0.1	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

White, male

2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	0.2
1999.....	0.1	*	*	*	*	*	*	*

White, female

2002.....	0.0	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*

Black, both sexes

2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Black, male

2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

Black, female

2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

All races 3/, both sexes

2002.....	0.5	*	*	0.2	0.2	0.4	0.7	0.8
2001.....	0.5	*	*	0.1	0.3	0.4	0.7	0.8
2000.....	0.5	*	*	0.2	0.3	0.6	0.7	0.9
1999.....	0.6	*	*	0.2	0.5	0.5	0.9	0.9

All races 3/, male

2002.....	0.8	*	*	0.3	0.4	0.7	1.0	1.2
2001.....	0.8	*	*	0.2	0.6	0.7	1.1	1.3
2000.....	0.8	*	*	0.3	0.5	0.9	1.1	1.4
1999.....	0.9	*	*	0.3	0.9	0.9	1.3	1.4

All races 3/, female

2002.....	0.2	*	*	*	*	*	0.3	0.3
2001.....	0.2	*	*	*	*	*	0.2	0.3
2000.....	0.2	*	*	*	*	0.2	0.3	0.4
1999.....	0.2	*	*	*	*	*	0.4	0.3

White, both sexes

2002.....	0.6	*	*	0.2	0.3	0.5	0.8	0.9
2001.....	0.6	*	*	0.2	0.4	0.5	0.8	0.9
2000.....	0.6	*	*	0.2	0.4	0.6	0.8	1.0
1999.....	0.6	*	*	0.2	0.6	0.6	1.0	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

White, male

2002.....	0.9	*	*	0.3	0.4	0.8	1.2	1.4
2001.....	0.9	*	*	0.2	0.7	0.8	1.3	1.5
2000.....	1.0	*	*	0.3	0.7	1.0	1.3	1.6
1999.....	1.0	*	*	0.3	1.1	0.9	1.6	1.6

White, female

2002.....	0.3	*	*	*	*	*	0.4	0.4
2001.....	0.3	*	*	*	*	*	*	0.4
2000.....	0.3	*	*	*	*	*	0.3	0.4
1999.....	0.3	*	*	*	*	*	0.5	0.4

Black, both sexes

2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*

Black, male

2002.....	0.2	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

Black, female

2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)										
All races 3/, both sexes										
2002.....	2.2	*	0.7	2.7	3.6	3.6	3.5	3.4
2001.....	2.2	*	0.8	2.7	3.5	3.5	3.5	3.6
2000.....	2.0	*	0.8	2.5	3.3	3.3	3.0	3.0
1999.....	1.9	*	0.6	2.2	2.9	3.1	3.4	3.2
All races 3/, male										
2002.....	3.8	*	1.1	4.3	6.2	6.0	6.0	5.9
2001.....	3.7	*	1.2	4.3	6.0	6.0	5.9	6.2
2000.....	3.4	*	1.3	4.0	5.7	5.6	5.2	5.2
1999.....	3.3	*	1.0	3.4	4.8	5.3	5.9	5.6
All races 3/, female										
2002.....	0.8	*	0.3	1.0	1.0	1.2	0.8	1.0
2001.....	0.7	*	0.3	1.0	0.9	0.9	1.0	1.1
2000.....	0.7	*	0.3	1.0	0.8	0.9	0.8	0.8
1999.....	0.7	*	0.2	0.9	0.8	0.9	0.9	0.8
White, both sexes										
2002.....	2.4	*	0.7	3.0	4.0	3.8	3.7	3.7
2001.....	2.3	*	0.8	2.9	3.8	3.6	3.6	4.0
2000.....	2.1	*	0.8	2.7	3.4	3.4	3.1	3.2
1999.....	2.0	*	0.6	2.2	3.0	3.3	3.6	3.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)										
White, male										
2002.....	4.0	*	1.1	4.9	6.7	6.3	6.3	6.2
2001.....	4.0	*	1.2	4.8	6.6	6.2	6.1	6.6
2000.....	3.6	*	1.2	4.3	6.0	6.0	5.4	5.5
1999.....	3.4	*	1.0	3.5	5.1	5.6	6.2	5.9
White, female										
2002.....	0.8	*	0.4	1.1	1.0	1.2	0.9	1.1
2001.....	0.7	*	0.3	1.0	0.9	0.9	1.0	1.2
2000.....	0.7	*	0.4	1.1	0.7	0.8	0.8	0.9
1999.....	0.7	*	*	0.9	0.8	0.9	0.9	0.9
Black, both sexes										
2002.....	1.1	*	0.7	1.0	2.0	2.3	2.3	1.9
2001.....	1.1	*	0.6	1.1	1.8	2.5	2.6	1.8
2000.....	1.2	*	1.0	1.3	2.3	2.3	1.9	1.7
1999.....	1.0	*	0.7	1.3	1.8	2.0	2.2	1.9
Black, male										
2002.....	2.0	*	1.1	1.6	3.1	4.0	4.3	3.6
2001.....	2.0	*	*	1.9	3.1	4.7	4.8	3.3
2000.....	2.1	*	1.8	2.1	4.1	3.9	3.7	3.0
1999.....	1.9	*	*	1.9	3.3	3.8	4.2	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
All races 3/, both sexes										
2002.....	5.9	*	0.4	3.6	6.7	6.4	5.6	6.6
2001.....	5.9	*	0.4	4.1	6.6	6.7	6.3	6.4
2000.....	5.9	*	0.5	4.4	7.2	6.3	6.0	6.7
1999.....	5.9	*	0.5	4.9	7.2	6.6	6.5	7.1
All races 3/, male										
2002.....	10.6	*	0.6	6.4	11.9	11.1	9.6	11.3
2001.....	10.6	*	0.6	7.1	11.8	11.7	10.8	10.8
2000.....	10.5	*	0.9	7.6	12.9	11.1	10.1	11.1
1999.....	10.6	*	0.8	8.4	12.8	11.5	10.9	11.8
All races 3/, female										
2002.....	1.4	*	*	0.7	1.2	1.6	1.7	2.0
2001.....	1.5	*	0.2	1.0	1.0	1.5	1.7	2.1
2000.....	1.5	*	0.2	1.1	1.3	1.5	1.9	2.3
1999.....	1.5	*	0.2	1.1	1.4	1.6	1.9	2.3
White, both sexes										
2002.....	6.7	*	0.4	3.9	7.0	6.8	6.2	7.4
2001.....	6.7	*	0.4	4.5	6.7	7.2	7.0	7.2
2000.....	6.6	*	0.6	4.6	7.6	6.6	6.6	7.5
1999.....	6.6	*	0.6	5.2	7.4	7.0	7.1	7.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
White, male										
2002.....	11.9	*	0.7	6.9	12.3	11.3	10.4	12.4
2001.....	11.8	*	0.7	7.6	11.9	12.3	11.7	11.9
2000.....	11.6	*	1.0	7.9	13.5	11.4	11.0	12.2
1999.....	11.7	*	0.9	8.9	13.0	11.9	11.8	12.7
White, female										
2002.....	1.6	*	*	0.8	1.3	2.0	1.9	2.3
2001.....	1.7	*	*	1.1	1.1	1.7	2.0	2.4
2000.....	1.7	*	*	1.1	1.3	1.6	2.0	2.7
1999.....	1.7	*	0.3	1.2	1.4	1.8	2.2	2.7
Black, both sexes										
2002.....	2.8	*	*	2.5	5.8	6.1	4.0	4.0
2001.....	2.9	*	*	2.8	7.1	5.8	4.1	3.7
2000.....	3.0	*	*	3.8	6.8	6.0	4.0	3.4
1999.....	3.1	*	*	3.9	7.6	5.7	4.2	4.2
Black, male										
2002.....	5.4	*	*	4.3	11.0	12.4	7.7	7.4
2001.....	5.5	*	*	5.0	13.9	10.9	8.0	7.1
2000.....	5.5	*	*	6.9	12.6	11.3	7.3	6.4
1999.....	5.9	*	*	7.0	14.2	11.2	8.1	8.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Black, female

2002.....	0.5	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by jumping from a high place (X80)										
All races 3/, both sexes										
2002.....	0.3	*	*	0.2	0.4	0.4	0.4	0.3
2001.....	0.2	*	*	0.2	0.2	0.3	0.3	0.3
2000.....	0.2	*	*	0.2	0.4	0.3	0.3	0.3
1999.....	0.2	*	*	0.2	0.4	0.3	0.3	0.4
All races 3/, male										
2002.....	0.4	*	*	0.2	0.6	0.6	0.6	0.5
2001.....	0.3	*	*	0.3	0.3	0.4	0.4	0.4
2000.....	0.3	*	*	0.3	0.5	0.4	0.3	0.5
1999.....	0.4	*	*	0.3	0.6	0.5	0.4	0.6
All races 3/, female										
2002.....	0.2	*	*	*	*	*	0.3	0.2
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	0.2	*	*	*
1999.....	0.1	*	*	*	*	*	*	0.2
White, both sexes										
2002.....	0.3	*	*	0.1	0.4	0.4	0.4	0.4
2001.....	0.2	*	*	0.2	0.3	0.2	0.3	0.3
2000.....	0.2	*	*	0.2	0.4	0.2	0.2	0.3
1999.....	0.2	*	*	0.2	0.4	0.3	0.3	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by jumping from a high place (X80)

White, male

2002.....	0.4	*	*	*	0.6	0.5	0.5	0.5
2001.....	0.3	*	*	0.3	0.4	0.3	0.4	0.4
2000.....	0.3	*	*	0.3	0.5	0.4	0.3	0.5
1999.....	0.4	*	*	0.2	0.6	0.4	0.4	0.5

White, female

2002.....	0.2	*	*	*	*	*	0.3	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	0.2

Black, both sexes

2002.....	0.2	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*

Black, male

2002.....	0.3	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by jumping from a high place (X80)

Black, female

2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)										
All races 3/, both sexes										
2002.....	0.6	*	*	0.3	0.6	0.7	0.7	0.9
2001 4/.....	0.6	*	*	0.3	0.6	0.7	0.7	0.9
2000.....	0.6	*	*	0.3	0.6	0.6	0.6	0.9
1999.....	0.6	*	*	0.3	0.6	0.6	0.7	0.8
All races 3/, male										
2002.....	1.0	*	*	0.6	1.0	1.0	1.2	1.5
2001 4/.....	0.9	*	*	0.5	1.0	1.1	1.0	1.4
2000.....	0.9	*	*	0.4	1.0	0.9	0.9	1.4
1999.....	0.8	*	*	0.5	1.0	0.9	1.2	1.1
All races 3/, female										
2002.....	0.3	*	*	*	0.3	0.4	0.3	0.4
2001 4/.....	0.3	*	*	*	0.2	0.2	0.3	0.4
2000.....	0.3	*	*	*	0.2	*	0.3	0.3
1999.....	0.3	*	*	*	0.2	*	0.3	0.5
White, both sexes										
2002.....	0.7	*	*	0.4	0.6	0.7	0.8	1.0
2001 4/.....	0.6	*	*	0.4	0.6	0.7	0.7	0.9
2000.....	0.6	*	*	0.3	0.6	0.6	0.6	0.9
1999.....	0.6	*	*	0.3	0.6	0.5	0.7	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)

White, male

2002.....	1.0	*	*	0.6	1.0	1.0	1.2	1.5
2001 4/.....	1.0	*	*	0.6	0.9	1.1	1.1	1.4
2000.....	0.9	*	*	0.4	0.9	1.0	0.9	1.4
1999.....	0.9	*	*	0.5	1.0	0.9	1.1	1.1

White, female

2002.....	0.3	*	*	*	0.3	0.4	0.3	0.4
2001 4/.....	0.3	*	*	*	*	*	0.4	0.4
2000.....	0.3	*	*	*	*	*	0.3	0.3
1999.....	0.3	*	*	*	*	*	0.3	0.5

Black, both sexes

2002.....	0.5	*	*	*	0.7	0.8	*	1.1
2001 4/.....	0.5	*	*	*	*	*	*	0.9
2000.....	0.5	*	*	*	1.0	*	*	0.9
1999.....	0.5	*	*	*	*	*	0.9	0.8

Black, male

2002.....	0.8	*	*	*	*	*	*	1.8
2001 4/.....	0.7	*	*	*	*	*	*	1.5
2000.....	0.8	*	*	*	*	*	*	1.9
1999.....	0.7	*	*	*	*	*	1.5	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)

Black, female

2002.....	0.2	*	*	*	*	*	*	*
2001 4/.....	0.2	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)										
All races 3/, both sexes										
2002.....	6.1	7.5	2.7	0.7	1.0	9.3	16.5	13.3	9.4	7.9
2001 4/.....	7.1	8.2	2.7	0.7	0.9	9.4	17.3	14.9	11.5	10.4
2000.....	6.0	9.2	2.3	0.7	1.1	9.5	16.0	11.9	9.0	7.6
1999.....	6.1	8.7	2.5	0.9	1.2	10.4	15.6	12.1	9.1	7.7
All races 3/, male										
2002.....	9.6	7.9	2.9	0.6	1.2	15.3	27.7	22.0	14.7	11.8
2001 4/.....	11.1	9.5	3.0	0.7	1.1	15.7	28.9	24.1	17.7	15.3
2000.....	9.3	10.4	2.5	0.7	1.4	15.5	26.7	19.6	14.0	10.9
1999.....	9.3	9.6	2.5	0.9	1.5	16.9	25.7	19.5	13.8	11.0
All races 3/, female										
2002.....	2.7	7.1	2.5	0.8	0.8	2.9	4.6	4.2	4.1	4.1
2001 4/.....	3.3	6.9	2.4	0.6	0.7	2.7	5.1	5.5	5.2	5.5
2000.....	2.8	7.9	2.1	0.6	0.8	3.1	4.7	4.1	4.0	4.4
1999.....	2.9	7.8	2.4	0.9	0.9	3.5	5.1	4.5	4.3	4.3
White, both sexes										
2002.....	3.7	5.4	2.0	0.5	0.7	5.2	8.2	6.8	5.1	5.0
2001 4/.....	4.9	6.2	1.9	0.7	0.7	5.1	9.4	8.9	7.8	7.8
2000.....	3.6	6.6	1.7	0.6	0.9	5.2	7.7	6.2	5.3	4.9
1999.....	3.8	6.7	1.7	0.7	1.0	5.6	8.0	6.5	5.5	5.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)										
White, male										
2002.....	5.4	6.2	2.1	0.4	0.7	8.2	12.9	10.5	7.3	7.0
2001 4/.....	7.2	7.3	2.2	0.6	0.8	7.9	14.5	13.5	11.3	11.6
2000.....	5.2	8.2	1.9	0.7	1.1	8.0	12.0	9.2	7.6	6.6
1999.....	5.5	7.9	1.8	0.7	1.1	8.6	12.1	9.9	7.8	7.0
White, female										
2002.....	2.0	4.6	1.8	0.6	0.6	1.9	3.1	2.8	2.8	3.1
2001 4/.....	2.6	5.1	1.6	0.7	0.6	2.0	3.9	4.1	4.2	3.9
2000.....	2.1	5.0	1.5	0.5	0.7	2.1	3.2	3.0	2.9	3.2
1999.....	2.1	5.4	1.7	0.7	0.8	2.3	3.7	2.9	3.2	3.1
Black, both sexes										
2002.....	22.0	17.3	6.3	1.4	2.6	31.0	63.4	54.2	37.0	26.9
2001 4/.....	22.1	18.9	6.8	0.9	1.9	32.7	62.8	52.6	35.3	28.2
2000.....	21.5	22.8	5.8	1.2	2.5	33.1	64.1	48.2	33.5	25.9
1999.....	21.1	19.5	6.1	1.9	2.5	36.5	60.4	47.0	31.8	25.1
Black, male										
2002.....	38.4	16.3	7.1	1.4	3.5	53.3	115.3	99.9	65.4	45.2
2001 4/.....	38.3	21.0	8.0	1.2	2.7	58.2	116.3	96.0	62.3	42.8
2000.....	37.2	23.3	5.9	*	3.3	57.2	117.8	89.4	58.6	42.5
1999.....	36.1	19.5	5.9	2.0	3.3	62.4	110.9	84.0	54.7	40.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)										
Black, female										
2002.....	7.0	18.5	5.5	1.3	1.6	8.0	12.7	12.3	11.5	10.6
2001 4/.....	7.4	16.7	5.6	*	*	6.4	11.4	13.1	11.1	15.3
2000.....	7.2	22.2	5.7	1.4	1.7	8.4	13.2	10.9	10.9	11.1
1999.....	7.6	19.5	6.3	1.8	1.6	10.0	12.5	13.6	11.2	11.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by hanging, strangulation and suffocation (X91)

All races 3/, both sexes

2002.....	0.2	0.8	0.2	*	*	0.2	0.3	0.3	0.3	0.4
2001.....	0.2	1.0	*	*	*	0.2	0.4	0.3	0.3	0.3
2000.....	0.2	0.7	0.2	*	*	0.2	0.4	0.3	0.4	0.3
1999.....	0.3	0.9	0.1	*	*	0.2	0.4	0.4	0.4	0.4

All races 3/, male

2002.....	0.2	*	*	*	*	*	0.2	0.2	*	0.2
2001.....	0.2	1.2	*	*	*	*	0.2	*	*	0.2
2000.....	0.2	*	*	*	*	*	*	*	0.3	*
1999.....	0.2	1.1	*	*	*	*	*	0.2	0.3	0.2

All races 3/, female

2002.....	0.3	*	*	*	*	0.3	0.5	0.5	0.4	0.6
2001.....	0.3	*	*	*	*	0.3	0.6	0.5	0.5	0.4
2000.....	0.3	*	*	*	*	0.3	0.6	0.5	0.4	0.4
1999.....	0.3	*	*	*	*	0.4	0.6	0.6	0.6	0.5

White, both sexes

2002.....	0.2	0.6	*	*	*	0.1	0.3	0.3	0.2	0.3
2001.....	0.2	0.7	*	*	*	0.2	0.3	0.3	0.3	0.3
2000.....	0.2	*	0.2	*	*	0.2	0.3	0.3	0.2	0.2
1999.....	0.2	0.7	*	*	*	0.2	0.3	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by hanging, strangulation and suffocation (X91)

Black, female

2002.....	0.7	*	*	*	*	*	*	*	*	1.6
2001.....	0.7	*	*	*	*	*	*	*	*	1.4
2000.....	0.8	*	*	*	*	*	1.4	1.4	1.4	*
1999.....	0.8	*	*	*	*	*	*	1.6	1.6	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
All races 3/, both sexes										
2002.....	4.1	*	0.3	0.3	0.7	7.7	13.6	10.6	6.9	5.2
2001.....	4.0	*	0.4	0.3	0.6	7.5	13.6	10.3	6.6	5.1
2000.....	3.8	*	0.2	0.2	0.7	7.7	12.7	9.2	6.3	4.7
1999.....	3.9	*	0.3	0.3	0.8	8.5	12.5	9.2	6.1	4.6
All races 3/, male										
2002.....	7.0	*	0.4	0.3	0.9	13.2	23.8	18.6	11.6	8.4
2001.....	6.8	*	0.3	0.3	0.8	13.2	24.0	17.8	11.1	8.0
2000.....	6.5	*	*	0.3	1.0	13.3	22.4	16.1	10.4	7.4
1999.....	6.5	*	0.3	0.4	1.1	14.5	21.8	15.9	10.1	7.1
All races 3/, female										
2002.....	1.3	*	*	0.3	0.5	1.8	2.9	2.4	2.1	2.0
2001.....	1.3	*	0.4	0.2	0.3	1.5	2.7	2.5	2.1	2.2
2000.....	1.3	*	*	0.2	0.3	1.7	2.7	2.2	2.1	1.9
1999.....	1.3	*	0.3	0.2	0.5	2.2	2.8	2.3	2.1	2.2
White, both sexes										
2002.....	2.2	*	0.3	0.2	0.4	4.1	6.0	4.9	3.4	3.0
2001.....	2.2	*	0.3	0.3	0.4	3.7	6.3	5.0	3.5	3.2
2000.....	2.1	*	*	0.2	0.5	3.9	5.5	4.2	3.3	2.8
1999.....	2.2	*	0.2	0.2	0.6	4.2	5.6	4.3	3.4	3.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
White, male										
2002.....	3.5	*	0.3	*	0.5	6.7	10.0	8.1	5.3	4.5
2001.....	3.5	*	*	0.3	0.6	6.2	10.5	8.2	5.4	4.9
2000.....	3.2	*	*	0.3	0.7	6.5	9.0	6.7	5.1	4.1
1999.....	3.4	*	*	0.2	0.8	6.9	9.1	7.1	5.2	4.3
White, female										
2002.....	1.0	*	*	*	0.3	1.2	1.6	1.6	1.4	1.5
2001.....	1.0	*	*	*	0.3	1.1	1.9	1.6	1.6	1.6
2000.....	1.0	*	*	*	*	1.1	1.7	1.6	1.5	1.5
1999.....	0.9	*	*	*	0.4	1.2	1.9	1.4	1.5	1.6
Black, both sexes										
2002.....	16.7	*	*	*	1.9	27.3	57.0	46.9	30.1	19.5
2001.....	15.8	*	*	*	1.3	28.7	55.7	43.6	26.9	18.4
2000.....	15.6	*	*	*	1.7	28.6	55.3	40.7	26.0	17.3
1999.....	15.4	*	*	0.7	1.8	32.0	53.4	39.3	23.7	15.9
Black, male										
2002.....	31.0	*	*	*	2.7	48.4	105.9	90.2	56.3	35.7
2001.....	29.8	*	*	*	2.1	52.8	106.5	83.2	51.3	31.5
2000.....	29.2	*	*	*	2.8	51.7	105.2	78.7	48.1	31.7
1999.....	28.4	*	*	*	2.8	56.3	101.0	74.4	44.0	27.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
Black, female										
2002.....	3.6	*	*	*	1.2	5.4	9.2	7.3	6.5	5.0
2001.....	3.1	*	*	*	*	3.9	7.0	7.6	5.0	6.8
2000.....	3.2	*	*	*	*	4.9	8.0	6.2	6.0	4.5
1999.....	3.7	*	*	*	*	7.0	8.2	7.5	5.5	5.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by sharp object (X99)										
All races 3/, both sexes										
2002.....	0.7	*	*	0.1	*	0.7	1.4	1.3	1.2	1.1
2001.....	0.7	*	*	*	*	0.8	1.6	1.4	1.0	1.1
2000.....	0.6	*	*	*	0.1	0.8	1.3	1.1	1.0	1.1
1999.....	0.7	*	*	*	0.1	0.8	1.4	1.1	1.2	1.1
All races 3/, male										
2002.....	1.0	*	*	*	*	1.1	2.2	1.9	1.5	1.6
2001.....	1.0	*	*	*	*	1.3	2.4	2.0	1.4	1.4
2000.....	0.9	*	*	*	*	1.1	2.1	1.6	1.5	1.4
1999.....	1.0	*	*	*	*	1.2	2.0	1.6	1.7	1.6
All races 3/, female										
2002.....	0.4	*	*	*	*	0.3	0.5	0.8	0.9	0.7
2001.....	0.4	*	*	*	*	0.3	0.8	0.7	0.6	0.8
2000.....	0.4	*	*	*	*	0.4	0.5	0.5	0.6	0.7
1999.....	0.4	*	*	*	*	0.4	0.7	0.6	0.7	0.6
White, both sexes										
2002.....	0.5	*	*	*	*	0.4	1.0	0.9	0.7	0.8
2001.....	0.5	*	*	*	*	0.5	1.2	1.0	0.7	0.7
2000.....	0.4	*	*	*	*	0.6	0.9	0.8	0.8	0.7
1999.....	0.5	*	*	*	*	0.5	1.0	0.9	0.8	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by sharp object (X99)										
White, male										
2002.....	0.7	*	*	*	*	0.7	1.6	1.3	0.9	1.2
2001.....	0.7	*	*	*	*	0.8	1.9	1.4	0.9	0.9
2000.....	0.6	*	*	*	*	0.9	1.5	1.2	1.0	1.0
1999.....	0.7	*	*	*	*	0.8	1.4	1.3	1.1	0.9
White, female										
2002.....	0.3	*	*	*	*	*	0.4	0.5	0.6	0.4
2001.....	0.3	*	*	*	*	*	0.5	0.5	0.4	0.5
2000.....	0.3	*	*	*	*	0.3	0.3	0.4	0.5	0.5
1999.....	0.3	*	*	*	*	0.3	0.5	0.4	0.5	0.4
Black, both sexes										
2002.....	2.2	*	*	*	*	2.0	3.3	4.2	3.4	3.2
2001.....	2.0	*	*	*	*	2.2	3.7	4.1	2.8	3.5
2000.....	1.9	*	*	*	*	1.6	3.5	2.9	2.7	3.4
1999.....	2.0	*	*	*	*	2.0	3.7	2.6	3.6	3.9
Black, male										
2002.....	3.2	*	*	*	*	3.0	5.4	6.3	4.9	4.1
2001.....	3.0	*	*	*	*	3.3	5.4	6.3	4.2	4.7
2000.....	2.8	*	*	*	*	2.0	5.7	4.5	4.4	4.6
1999.....	3.1	*	*	*	*	3.1	5.3	3.6	5.5	6.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by sharp object (X99)										
Black, female										
2002.....	1.2	*	*	*	*	*	1.3	2.2	2.0	2.3
2001.....	1.1	*	*	*	*	*	2.0	2.1	1.6	2.5
2000.....	1.1	*	*	*	*	*	1.5	1.4	*	2.3
1999.....	1.0	*	*	*	*	*	2.2	1.7	1.9	1.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Neglect, abandonment and other maltreatment syndromes (Y06-Y07)										
All races 3/, both sexes										
2002.....	0.1	2.4	0.6	*	*	*	*	*	*	*
2001.....	0.1	2.9	0.5	*	*	*	*	*	*	*
2000.....	0.1	3.4	0.5	*	*	*	*	*	*	*
1999.....	0.1	3.3	0.4	*	*	*	*	*	*	*
All races 3/, male										
2002.....	0.1	2.5	0.6	*	*	*	*	*	*	*
2001.....	0.1	3.2	0.7	*	*	*	*	*	*	*
2000.....	0.1	4.0	0.5	*	*	*	*	*	*	*
1999.....	0.1	3.7	0.4	*	*	*	*	*	*	*
All races 3/, female										
2002.....	0.1	2.4	0.6	*	*	*	*	*	*	*
2001.....	0.1	2.5	0.4	*	*	*	*	*	*	*
2000.....	0.1	2.7	0.4	*	*	*	*	*	*	*
1999.....	0.1	2.9	0.4	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.1	1.9	0.5	*	*	*	*	*	*	*
2001.....	0.1	2.2	0.4	*	*	*	*	*	*	*
2000.....	0.1	2.6	0.4	*	*	*	*	*	*	*
1999.....	0.1	2.5	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

All races 3/, both sexes

2002.....	0.9	3.7	1.4	0.2	0.1	0.6	0.9	0.8	0.9	1.0
2001 4/.....	2.0	3.5	1.6	0.2	0.2	0.7	1.5	2.8	3.4	3.6
2000.....	1.0	4.3	1.3	0.3	0.2	0.7	1.3	1.1	1.2	1.4
1999.....	1.0	3.8	1.3	0.4	0.2	0.8	1.1	1.1	1.1	1.3

All races 3/, male

2002.....	1.2	4.0	1.6	*	*	0.7	1.2	1.1	1.2	1.3
2001 4/.....	2.9	4.0	1.7	0.3	*	0.9	2.0	3.8	4.9	5.5
2000.....	1.4	4.7	1.5	0.3	0.2	0.9	1.8	1.6	1.5	1.7
1999.....	1.3	4.0	1.4	0.3	0.2	1.0	1.4	1.4	1.4	1.8

All races 3/, female

2002.....	0.5	3.3	1.2	*	*	0.4	0.6	0.5	0.6	0.7
2001 4/.....	1.1	3.1	1.4	*	0.2	0.5	1.0	1.7	1.9	1.8
2000.....	0.7	3.9	1.1	0.3	0.3	0.6	0.8	0.7	0.8	1.1
1999.....	0.7	3.6	1.2	0.4	*	0.6	0.7	0.8	0.9	0.9

White, both sexes

2002.....	0.7	2.5	0.9	*	*	0.4	0.7	0.6	0.7	0.8
2001 4/.....	1.8	2.7	1.1	0.2	0.2	0.6	1.4	2.6	3.3	3.4
2000.....	0.7	3.0	0.8	0.2	0.2	0.5	0.9	0.8	0.8	1.0
1999.....	0.8	2.9	1.0	0.2	0.2	0.6	0.9	0.8	0.8	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

White, male

2002.....	0.9	2.9	1.1	*	*	0.5	0.9	0.9	0.9	1.0
2001 4/.....	2.6	3.1	1.2	*	*	0.7	1.8	3.5	4.7	5.5
2000.....	0.9	3.5	1.0	*	*	0.5	1.2	1.0	1.0	1.2
1999.....	1.0	3.1	1.1	*	*	0.7	1.2	1.0	1.0	1.3

White, female

2002.....	0.4	2.2	0.8	*	*	0.4	0.5	0.4	0.4	0.6
2001 4/.....	1.0	2.3	0.9	*	*	0.4	0.9	1.6	1.7	1.3
2000.....	0.5	2.4	0.7	*	*	0.5	0.6	0.6	0.6	0.7
1999.....	0.5	2.5	0.9	*	*	0.5	0.7	0.6	0.7	0.7

Black, both sexes

2002.....	2.1	9.2	3.8	*	*	1.0	2.0	1.9	2.3	2.6
2001 4/.....	3.1	8.0	4.1	*	*	1.3	2.2	3.8	4.5	5.1
2000.....	2.9	11.2	3.8	0.7	*	2.2	3.9	3.3	3.3	4.0
1999.....	2.5	8.9	3.2	0.9	*	1.7	2.0	3.1	3.0	3.6

Black, male

2002.....	3.1	9.3	4.3	*	*	1.5	2.9	2.5	3.2	3.7
2001 4/.....	4.3	8.4	4.8	*	*	1.6	3.4	5.7	5.9	6.1
2000.....	4.2	11.2	3.9	*	*	3.0	5.7	5.2	4.9	5.3
1999.....	3.4	9.1	3.6	*	*	2.5	3.3	4.2	3.9	4.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

Black, female

2002.....	1.2	9.1	3.3	*	*	*	*	*	1.6	1.6
2001 4/.....	2.1	7.6	3.3	*	*	*	*	2.0	3.1	4.3
2000.....	1.7	11.2	3.7	*	*	1.3	2.2	1.7	1.9	2.8
1999.....	1.6	8.7	2.9	*	*	*	*	2.1	2.1	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2002.....	1.7	2.5	0.4	0.1	0.2	0.7	1.7	1.9	2.3	3.3
2001.....	1.5	2.1	0.3	*	0.2	0.7	1.4	1.6	2.0	3.0
2000.....	1.4	1.7	0.3	*	0.2	0.6	1.3	1.6	1.8	2.6
1999.....	1.4	1.4	0.3	*	0.2	0.8	1.5	1.6	2.1	2.8
All races 3/, male										
2002.....	2.2	2.5	0.4	*	0.2	1.0	2.4	2.7	3.1	4.5
2001.....	2.0	2.4	*	*	0.3	1.0	2.1	2.4	2.7	3.9
2000.....	1.9	1.4	0.3	*	0.4	0.9	2.0	2.3	2.6	3.7
1999.....	2.0	1.7	0.3	*	0.3	1.3	2.2	2.3	2.8	3.9
All races 3/, female										
2002.....	1.2	2.5	0.4	*	*	0.4	1.0	1.0	1.4	2.1
2001.....	0.9	1.7	0.3	*	*	0.3	0.7	0.9	1.3	2.1
2000.....	0.8	2.0	*	*	*	0.3	0.6	0.9	1.1	1.6
1999.....	0.9	1.2	0.3	*	*	0.3	0.6	0.9	1.3	1.7
White, both sexes										
2002.....	1.7	1.8	0.4	*	0.2	0.7	1.9	1.9	2.3	3.4
2001.....	1.5	1.4	0.3	*	0.2	0.7	1.5	1.7	2.1	3.0
2000.....	1.3	1.4	0.2	*	0.2	0.6	1.3	1.6	1.8	2.7
1999.....	1.4	1.2	0.2	*	0.2	0.8	1.5	1.6	2.1	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
White, male										
2002.....	2.2	1.8	0.4	*	0.2	1.1	2.6	2.7	3.1	4.5
2001.....	2.0	1.7	*	*	0.3	1.1	2.2	2.5	2.9	3.8
2000.....	1.8	*	*	*	0.3	1.0	1.9	2.3	2.5	3.7
1999.....	1.9	1.4	*	*	0.3	1.3	2.3	2.3	2.8	3.7
White, female										
2002.....	1.2	1.8	0.4	*	*	0.4	1.1	1.1	1.5	2.3
2001.....	1.0	*	*	*	*	0.3	0.8	0.9	1.4	2.3
2000.....	0.8	1.6	*	*	*	0.3	0.7	0.9	1.1	1.6
1999.....	0.9	*	*	*	*	*	0.6	1.0	1.4	1.8
Black, both sexes										
2002.....	2.0	5.5	*	*	*	*	1.3	2.0	2.5	4.1
2001.....	1.7	5.5	*	*	*	*	1.1	1.5	1.9	3.8
2000.....	1.8	*	*	*	*	*	1.4	1.7	2.2	3.3
1999.....	1.9	*	*	*	*	0.8	1.7	1.9	2.5	3.9
Black, male										
2002.....	2.9	*	*	*	*	*	1.9	3.2	3.7	6.5
2001.....	2.7	6.0	*	*	*	*	1.9	2.5	2.7	5.9
2000.....	2.8	*	*	*	*	*	2.5	2.7	3.4	4.8
1999.....	3.0	*	*	*	*	1.3	2.7	3.4	3.7	6.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
Black, female										
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.9
2001.....	0.9	*	*	*	*	*	*	*	*	2.0
2000.....	0.9	*	*	*	*	*	*	*	*	2.0
1999.....	0.9	*	*	*	*	*	*	*	1.4	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)										
All races 3/, both sexes										
2002.....	1.1	*	*	*	*	0.4	1.0	1.4	1.7	2.5
2001.....	1.0	*	*	*	*	0.3	0.8	1.1	1.5	2.3
2000.....	0.9	*	*	*	*	0.3	0.7	1.0	1.2	2.1
1999.....	0.9	*	*	*	*	0.3	0.8	1.1	1.5	2.1
All races 3/, male										
2002.....	1.4	*	*	*	*	0.5	1.4	1.9	2.1	3.3
2001.....	1.3	*	*	*	*	0.5	1.2	1.5	1.9	2.9
2000.....	1.2	*	*	*	*	0.4	1.0	1.4	1.7	2.8
1999.....	1.2	*	*	*	*	0.4	1.1	1.5	2.0	2.9
All races 3/, female										
2002.....	0.8	*	*	*	*	0.2	0.6	0.8	1.2	1.7
2001.....	0.7	*	*	*	*	*	0.4	0.7	1.1	1.7
2000.....	0.5	*	*	*	*	*	0.4	0.6	0.7	1.3
1999.....	0.6	*	*	*	*	*	0.4	0.6	1.0	1.4
White, both sexes										
2002.....	1.2	*	*	*	*	0.4	1.2	1.5	1.8	2.6
2001.....	1.0	*	*	*	*	0.4	1.0	1.2	1.6	2.4
2000.....	0.9	*	*	*	*	0.3	0.8	1.1	1.3	2.1
1999.....	0.9	*	*	*	*	0.3	0.9	1.1	1.7	2.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)										
White, male										
2002.....	1.4	*	*	*	*	0.5	1.6	2.1	2.2	3.3
2001.....	1.3	*	*	*	*	0.6	1.4	1.6	2.1	2.9
2000.....	1.2	*	*	*	*	0.4	1.2	1.5	1.8	2.9
1999.....	1.1	*	*	*	*	0.5	1.2	1.5	2.2	2.8
White, female										
2002.....	0.9	*	*	*	*	0.3	0.8	0.9	1.2	1.9
2001.....	0.7	*	*	*	*	*	0.6	0.8	1.1	1.8
2000.....	0.6	*	*	*	*	*	0.5	0.7	0.8	1.3
1999.....	0.6	*	*	*	*	*	0.5	0.7	1.1	1.4
Black, both sexes										
2002.....	1.2	*	*	*	*	*	*	1.1	1.7	2.9
2001.....	1.1	*	*	*	*	*	*	0.8	1.6	2.8
2000.....	1.0	*	*	*	*	*	*	0.9	1.3	2.4
1999.....	1.2	*	*	*	*	*	*	1.2	1.5	2.9
Black, male										
2002.....	1.6	*	*	*	*	*	*	1.6	2.2	4.4
2001.....	1.7	*	*	*	*	*	*	*	2.0	4.0
2000.....	1.5	*	*	*	*	*	*	*	2.0	3.3
1999.....	1.8	*	*	*	*	*	*	2.1	2.2	4.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Discharge of firearms, undetermined intent (Y22-Y24)										
All races 3/, both sexes										
2002.....	0.1	*	*	*	*	0.2	0.2	0.1	*	0.1
2001.....	0.1	*	*	*	*	0.1	0.2	0.2	*	0.1
2000.....	0.1	*	*	*	*	0.1	0.2	0.2	0.1	*
1999.....	0.1	*	*	*	*	0.3	0.3	0.2	*	0.1
All races 3/, male										
2002.....	0.1	*	*	*	*	0.3	0.4	0.2	*	0.2
2001.....	0.1	*	*	*	*	0.2	0.4	0.3	*	0.2
2000.....	0.1	*	*	*	*	0.2	0.4	0.3	*	*
1999.....	0.2	*	*	*	*	0.6	0.5	0.3	*	0.2
All races 3/, female										
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.1	*	*	*	*	0.1	0.2	0.1	*	0.1
2001.....	0.1	*	*	*	*	*	0.2	0.2	*	0.2
2000.....	0.1	*	*	*	*	0.1	0.2	0.1	*	*
1999.....	0.1	*	*	*	*	0.3	0.3	0.1	*	0.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2002.....	0.4	2.1	0.3	0.1	0.1	0.1	0.4	0.3	0.4	0.5
2001.....	0.3	1.7	0.2	*	0.1	0.1	0.3	0.3	0.3	0.5
2000.....	0.3	1.6	0.2	*	0.1	0.2	0.3	0.3	0.4	0.4
1999.....	0.3	1.2	0.2	*	0.1	0.2	0.3	0.3	0.4	0.5
All races 3/, male										
2002.....	0.5	2.0	0.3	*	*	*	0.5	0.4	0.6	0.9
2001.....	0.5	2.0	*	*	*	0.2	0.5	0.5	0.4	0.7
2000.....	0.5	1.2	0.3	*	0.2	0.2	0.5	0.4	0.6	0.5
1999.....	0.5	1.4	*	*	*	0.3	0.5	0.5	0.5	0.7
All races 3/, female										
2002.....	0.3	2.1	0.3	*	*	*	0.2	*	*	0.2
2001.....	0.2	1.5	0.3	*	*	*	*	*	*	0.2
2000.....	0.2	1.9	*	*	*	*	*	*	0.2	0.2
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.3
White, both sexes										
2002.....	0.4	1.6	0.3	*	*	0.1	0.4	0.2	0.4	0.5
2001.....	0.3	1.1	0.2	*	*	0.2	0.3	0.3	0.3	0.4
2000.....	0.3	1.2	0.2	*	*	0.2	0.3	0.3	0.4	0.4
1999.....	0.3	1.1	0.2	*	*	0.2	0.3	0.3	0.3	0.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Legal intervention (Y35,Y89.0)										
All races 3/, both sexes										
2002.....	0.1	*	*	*	*	0.1	0.3	0.3	0.2	0.3
2001.....	0.1	*	*	*	*	0.1	0.3	0.2	0.3	0.3
2000.....	0.1	*	*	*	*	0.1	0.2	0.3	0.2	0.3
1999.....	0.1	*	*	*	*	0.1	0.3	0.3	0.3	0.3
All races 3/, male										
2002.....	0.3	*	*	*	*	0.2	0.6	0.5	0.5	0.5
2001.....	0.3	*	*	*	*	0.2	0.6	0.5	0.6	0.6
2000.....	0.3	*	*	*	*	0.3	0.5	0.6	0.5	0.5
1999.....	0.3	*	*	*	*	0.2	0.5	0.7	0.7	0.6
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.3
2001.....	0.1	*	*	*	*	*	0.3	0.2	0.3	0.3
2000.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
1999.....	0.1	*	*	*	*	*	0.1	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Legal intervention (Y35,Y89.0)										
White, male										
2002.....	0.2	*	*	*	*	*	0.4	0.4	0.5	0.5
2001.....	0.2	*	*	*	*	*	0.5	0.4	0.5	0.5
2000.....	0.2	*	*	*	*	*	0.4	0.4	0.4	0.4
1999.....	0.2	*	*	*	*	*	0.3	0.5	0.6	0.5
White, female										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002.....	0.3	*	*	*	*	*	0.9	0.8	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	0.7
1999.....	0.3	*	*	*	*	*	0.8	0.9	*	0.7
Black, male										
2002.....	0.6	*	*	*	*	*	1.7	1.6	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.6	*	*	*	*	*	*	*	*	1.4
1999.....	0.7	*	*	*	*	*	1.7	1.9	*	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)

All races 3/, both sexes

2002.....	0.1	*	*	*	*	0.1	0.3	0.2	0.2	0.2
2001.....	0.1	*	*	*	*	0.1	0.3	0.2	0.3	0.2
2000.....	0.1	*	*	*	*	0.1	0.2	0.3	0.2	0.2
1999.....	0.1	*	*	*	*	0.1	0.2	0.3	0.3	0.2

All races 3/, male

2002.....	0.2	*	*	*	*	0.2	0.6	0.5	0.4	0.4
2001.....	0.2	*	*	*	*	0.2	0.6	0.4	0.5	0.4
2000.....	0.2	*	*	*	*	0.3	0.4	0.5	0.3	0.4
1999.....	0.2	*	*	*	*	0.2	0.5	0.6	0.6	0.4

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
2001.....	0.1	*	*	*	*	*	0.3	0.2	0.2	0.2
2000.....	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
1999.....	0.1	*	*	*	*	*	0.1	0.2	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)

White, male

2002.....	0.2	*	*	*	*	*	0.4	0.3	0.4	0.3
2001.....	0.2	*	*	*	*	*	0.5	0.4	0.5	0.4
2000.....	0.2	*	*	*	*	*	0.4	0.4	0.3	0.3
1999.....	0.2	*	*	*	*	*	0.3	0.4	0.5	0.3

White, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, both sexes

2002.....	0.2	*	*	*	*	*	0.9	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	0.8	*	*	*

Black, male

2002.....	0.5	*	*	*	*	*	1.7	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	1.7	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Complications of medical and surgical care (Y40-Y84,Y88)										
All races 3/, both sexes										
2002.....	1.0	*	*	*	0.1	*	*	0.1	0.2	0.3
2001.....	1.1	0.6	0.2	*	*	*	0.1	0.1	0.3	0.3
2000.....	1.1	*	0.1	*	*	*	0.1	0.2	0.2	0.3
1999.....	1.0	1.4	0.2	*	*	*	*	0.1	0.2	0.3
All races 3/, male										
2002.....	0.9	*	*	*	*	*	*	*	0.3	0.3
2001.....	1.0	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.0	*	*	*	*	*	*	0.2	0.2	0.3
1999.....	1.0	1.6	*	*	*	*	*	*	0.3	0.3
All races 3/, female										
2002.....	1.1	*	*	*	*	*	*	*	0.2	0.3
2001.....	1.1	*	*	*	*	*	*	*	0.2	0.3
2000.....	1.1	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.1	1.2	*	*	1.1	*	*	0.2	*	0.3
White, both sexes										
2002.....	1.0	*	*	*	*	*	*	0.1	0.2	0.3
2001.....	1.1	*	*	*	*	*	0.1	*	0.2	0.3
2000.....	1.1	*	*	*	*	*	*	0.1	0.2	0.3
1999.....	1.0	1.2	0.2	*	*	*	*	*	0.2	0.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

All races 3/, both sexes

2002.....	0.8	*	*	*	*	*	*	*	0.2	0.3
2001.....	0.9	*	*	*	*	*	*	*	0.2	0.2
2000.....	0.9	*	*	*	*	*	*	0.1	0.2	0.2
1999.....	0.9	1.2	0.1	*	*	*	*	*	0.2	0.2

All races 3/, male

2002.....	0.7	*	*	*	*	*	*	*	*	0.2
2001.....	0.8	*	*	*	*	*	*	*	*	0.2
2000.....	0.9	*	*	*	*	*	*	*	*	*
1999.....	0.8	1.4	*	*	*	*	*	*	0.2	0.2

All races 3/, female

2002.....	0.9	*	*	*	*	*	*	*	*	0.3
2001.....	1.0	*	*	*	*	*	*	*	*	0.2
2000.....	1.0	*	*	*	*	*	*	*	0.2	0.2
1999.....	0.9	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.9	*	*	*	*	*	*	*	0.2	0.2
2001.....	0.9	*	*	*	*	*	*	*	0.2	0.2
2000.....	0.9	*	*	*	*	*	*	*	0.2	0.2
1999.....	0.9	1.0	*	*	*	*	*	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
All races 3/, both sexes										
2002.....	242.6	357.1	512.8	770.1	1,187.7	1,832.7	2,845.9	4,449.1	7,103.8	14,828.3
2001.....	240.5	354.8	512.4	771.8	1,210.7	1,869.7	2,878.3	4,494.0	7,151.9	15,112.8
2000.....	239.1	352.5	509.0	792.6	1,241.0	1,906.3	2,929.4	4,553.8	7,335.0	15,524.4
1999.....	236.3	346.6	501.4	801.6	1,254.0	1,956.7	2,995.3	4,608.0	7,378.5	15,554.6
All races 3/, male										
2002.....	306.4	454.7	653.5	961.8	1,476.0	2,265.9	3,533.0	5,522.7	8,652.3	16,254.5
2001.....	304.8	455.4	646.8	958.4	1,496.8	2,315.6	3,587.4	5,603.2	8,781.9	16,744.8
2000.....	305.9	456.3	642.2	989.8	1,535.9	2,362.0	3,676.1	5,693.2	9,095.4	17,501.4
1999.....	304.0	448.0	632.4	1,003.8	1,560.8	2,444.1	3,785.1	5,806.4	9,216.8	17,597.2
All races 3/, female										
2002.....	179.7	262.4	378.2	590.0	926.0	1,458.7	2,296.7	3,690.9	6,162.2	14,209.6
2001.....	177.2	257.2	383.7	596.9	951.3	1,485.9	2,314.8	3,717.6	6,175.1	14,429.9
2000.....	173.4	252.0	381.3	608.1	973.7	1,515.7	2,341.2	3,760.2	6,296.6	14,719.2
1999.....	169.8	248.5	375.9	612.6	976.4	1,539.6	2,378.2	3,774.8	6,302.8	14,731.3
White, both sexes										
2002.....	224.0	327.4	471.3	723.5	1,133.0	1,777.6	2,796.3	4,407.8	7,111.3	15,015.9
2001.....	218.8	322.8	471.4	723.2	1,156.1	1,805.8	2,823.2	4,446.8	7,149.4	15,287.5
2000.....	215.9	317.9	467.4	743.9	1,184.9	1,844.5	2,866.8	4,496.6	7,333.2	15,707.0
1999.....	210.8	310.5	460.0	747.2	1,193.5	1,895.5	2,923.9	4,542.8	7,370.9	15,718.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
White, male										
2002.....	287.2	420.4	601.2	902.0	1,402.7	2,196.1	3,469.7	5,469.3	8,660.8	16,473.2
2001.....	281.4	417.1	595.5	896.0	1,427.9	2,237.6	3,513.9	5,543.5	8,771.1	16,935.4
2000.....	279.4	416.1	588.3	925.1	1,465.1	2,285.1	3,593.1	5,620.3	9,092.5	17,716.4
1999.....	274.0	402.1	579.3	934.6	1,484.0	2,367.8	3,698.3	5,727.2	9,211.2	17,752.9
White, female										
2002.....	160.5	235.1	344.3	552.6	883.7	1,409.1	2,247.4	3,648.0	6,161.6	14,382.8
2001.....	155.9	229.4	350.1	558.4	905.3	1,426.7	2,263.9	3,669.9	6,169.3	14,597.6
2000.....	152.2	220.7	349.3	571.5	926.5	1,459.3	2,284.0	3,706.2	6,286.6	14,890.7
1999.....	147.4	219.9	343.6	569.0	925.9	1,483.5	2,308.5	3,711.2	6,284.5	14,900.6
Black, both sexes										
2002.....	412.9	634.1	925.9	1,321.6	1,880.7	2,632.0	3,824.9	5,571.3	7,962.2	14,260.8
2001.....	428.4	644.1	928.5	1,328.9	1,891.7	2,738.8	3,904.2	5,660.2	8,028.9	14,550.8
2000.....	434.5	666.6	938.7	1,347.0	1,929.0	2,769.0	4,019.7	5,782.6	8,161.0	14,714.3
1999.....	448.6	668.3	929.9	1,399.3	1,969.2	2,813.7	4,148.0	5,878.3	8,169.4	14,757.9
Black, male										
2002.....	502.9	801.4	1,206.6	1,719.3	2,464.7	3,357.1	4,918.1	7,088.6	9,968.4	15,635.5
2001.....	527.8	830.2	1,201.4	1,718.2	2,424.6	3,497.3	5,053.2	7,255.7	10,256.3	16,439.9
2000.....	549.1	853.0	1,227.4	1,776.2	2,462.1	3,513.8	5,203.9	7,410.0	10,405.7	16,791.0
1999.....	572.7	885.6	1,220.1	1,822.6	2,549.7	3,599.1	5,336.3	7,538.1	10,433.8	17,304.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
Black, female										
2002.....	333.6	489.0	688.0	995.2	1,423.7	2,093.9	3,091.2	4,664.6	6,917.8	13,734.2
2001.....	340.6	482.8	696.2	1,009.5	1,472.5	2,178.4	3,138.8	4,707.8	6,894.3	13,849.9
2000.....	333.3	505.0	692.6	995.2	1,508.3	2,219.3	3,233.3	4,801.6	7,030.5	13,941.3
1999.....	339.3	480.9	683.9	1,053.5	1,514.0	2,231.6	3,359.6	4,878.6	7,047.2	13,805.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
I. Certain infectious and parasitic diseases (A00-B99)										
All races 3/, both sexes										
2002.....	19.6	24.7	25.1	24.0	28.8	41.5	60.8	94.0	140.5	252.8
2001.....	19.7	23.2	22.6	22.9	28.7	39.4	57.2	87.5	130.0	246.3
2000.....	19.5	23.5	22.2	21.9	27.5	37.5	54.7	85.1	131.5	256.0
1999.....	20.9	22.9	21.1	22.2	26.5	40.3	56.4	86.9	132.5	267.9
All races 3/, male										
2002.....	27.5	35.6	36.6	31.2	35.3	47.9	69.6	109.1	164.0	270.9
2001.....	28.1	33.9	32.3	30.4	34.0	44.8	65.5	102.4	153.8	269.7
2000.....	27.8	34.9	32.3	28.8	33.4	43.8	62.7	99.8	154.8	273.0
1999.....	30.6	33.9	30.2	29.1	32.0	47.4	65.4	102.6	157.8	290.1
All races 3/, female										
2002.....	11.8	14.0	14.1	17.2	22.9	36.0	53.8	83.4	126.2	245.0
2001.....	11.4	12.8	13.2	15.8	23.9	34.8	50.6	77.1	115.8	236.5
2000.....	11.3	12.5	12.6	15.4	22.1	32.1	48.4	74.9	117.8	249.1
1999.....	11.3	12.4	12.4	15.8	21.5	34.2	49.3	75.9	117.8	258.9
White, both sexes										
2002.....	12.8	17.8	18.4	18.7	23.5	35.8	55.1	87.3	133.9	242.3
2001.....	12.8	16.3	16.3	17.9	23.2	34.0	51.5	81.4	123.5	235.4
2000.....	12.8	16.2	15.9	16.9	22.0	31.9	49.0	77.9	123.0	242.9
1999.....	13.2	15.3	15.3	16.6	21.6	34.7	49.5	78.9	124.1	251.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
I. Certain infectious and parasitic diseases (A00-B99)										
White, male										
2002.....	18.9	26.5	26.6	24.1	28.5	40.7	63.0	100.0	155.3	260.3
2001.....	19.5	24.4	23.5	23.3	27.2	38.3	58.3	95.3	146.1	256.7
2000.....	18.9	24.6	22.9	21.9	26.3	36.8	56.4	90.4	145.0	258.5
1999.....	20.3	22.6	21.8	21.2	25.5	40.2	57.2	94.2	146.7	270.4
White, female										
2002.....	6.6	9.3	10.4	13.5	18.9	31.5	48.6	78.2	120.8	234.5
2001.....	6.1	8.4	9.2	12.6	19.5	30.3	46.0	71.6	109.8	226.4
2000.....	6.6	7.9	9.0	12.1	18.0	27.7	43.1	69.1	110.0	236.5
1999.....	6.2	8.1	8.9	12.3	18.1	29.8	43.4	68.2	110.7	244.5
Black, both sexes										
2002.....	68.7	78.3	81.7	73.6	76.1	94.7	122.7	181.9	238.1	413.1
2001.....	69.7	77.3	75.6	69.7	79.2	89.6	119.4	166.2	226.7	404.7
2000.....	68.7	80.8	77.5	66.3	74.5	89.9	119.3	177.5	253.3	436.7
1999.....	76.9	83.1	71.9	72.0	69.0	93.0	131.6	181.7	249.4	469.0
Black, male										
2002.....	93.7	112.7	126.8	104.1	101.6	123.4	151.2	232.9	297.3	448.1
2001.....	96.3	114.3	111.6	101.8	102.5	111.6	151.1	204.0	278.5	470.9
2000.....	97.4	121.5	120.5	96.7	100.2	116.0	144.1	229.2	311.1	486.7
1999.....	112.3	128.3	108.8	103.4	95.7	120.0	166.2	213.7	325.2	537.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
I. Certain infectious and parasitic diseases (A00-B99)										
Black, female										
2002.....	46.7	48.4	43.5	48.5	56.2	73.4	103.5	151.5	207.3	399.6
2001.....	46.1	45.2	44.9	43.4	60.8	73.4	98.2	143.7	200.3	380.2
2000.....	43.5	45.6	40.9	41.4	54.3	70.7	102.8	146.4	224.2	418.1
1999.....	45.7	44.0	40.7	46.4	48.1	73.1	108.7	162.5	211.7	443.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intestinal infectious diseases (A00-A09)										
All races 3/, both sexes										
2002.....	*	0.1	0.2	0.4	0.7	1.4	2.7	5.1	9.8	21.0
2001.....	*	*	0.2	0.2	0.5	0.8	1.9	3.3	6.6	13.1
2000.....	*	*	0.1	0.3	0.4	0.9	1.6	2.8	5.4	11.9
1999.....	*	*	*	*	0.3	0.8	1.3	2.4	4.6	10.6
All races 3/, male										
2002.....	*	*	*	0.4	0.8	1.6	3.1	5.1	10.3	20.4
2001.....	*	*	*	*	0.4	0.8	2.0	3.6	6.5	12.2
2000.....	*	*	*	0.4	0.4	1.0	1.7	3.0	5.9	11.5
1999.....	*	*	*	*	*	0.8	1.3	2.7	5.1	9.7
All races 3/, female										
2002.....	*	*	*	0.5	0.6	1.3	2.3	5.1	9.5	21.3
2001.....	*	*	0.2	*	0.6	0.8	1.8	3.1	6.6	13.4
2000.....	*	*	*	*	*	0.8	1.6	2.7	5.1	12.0
1999.....	*	*	*	*	*	0.8	1.3	2.2	4.4	10.9
White, both sexes										
2002.....	*	0.1	0.2	0.4	0.6	1.6	2.7	5.3	10.4	22.1
2001.....	*	*	0.2	0.2	0.5	0.8	2.0	3.5	6.7	13.8
2000.....	*	*	*	0.3	0.4	0.9	1.7	3.0	5.8	12.3
1999.....	*	*	*	*	0.3	0.8	1.3	2.5	4.8	11.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other intestinal infectious diseases (A01-A08)										
All races 3/, both sexes										
2002.....	*	0.1	0.2	0.4	0.7	1.4	2.6	5.0	9.4	20.2
2001.....	*	*	0.2	0.2	0.5	0.8	1.8	3.2	6.4	12.6
2000.....	*	*	0.1	0.3	0.3	0.8	1.5	2.7	5.3	11.3
1999.....	*	*	*	*	0.3	0.7	1.1	2.2	4.3	9.9
All races 3/, male										
2002.....	*	*	*	0.4	0.7	1.5	3.0	5.0	9.9	19.7
2001.....	*	*	*	*	0.4	0.8	2.0	3.4	6.4	11.9
2000.....	*	*	*	0.4	*	1.0	1.6	2.9	5.8	10.6
1999.....	*	*	*	*	*	0.7	1.1	2.6	4.9	9.3
All races 3/, female										
2002.....	*	*	*	0.4	0.6	1.3	2.2	5.0	9.1	20.4
2001.....	*	*	*	*	0.6	0.8	1.7	3.0	6.4	13.0
2000.....	*	*	*	*	*	0.7	1.5	2.6	5.0	11.5
1999.....	*	*	*	*	*	0.7	1.2	1.9	4.0	10.2
White, both sexes										
2002.....	*	0.1	0.2	0.4	0.6	1.5	2.6	5.2	10.1	21.1
2001.....	*	*	0.2	0.2	0.5	0.7	1.9	3.3	6.5	13.3
2000.....	*	*	*	0.3	0.3	0.8	1.6	2.8	5.6	11.7
1999.....	*	*	*	*	0.3	0.7	1.2	2.2	4.5	10.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intestinal infections due to other specified organisms (A04,A07-A08)

All races 3/, both sexes

2002.....	*	0.1	0.2	0.4	0.7	1.3	2.5	4.9	9.3	20.2
2001.....	*	*	0.2	0.2	0.5	0.7	1.8	3.1	6.2	12.4
2000.....	*	*	*	0.2	0.3	0.8	1.5	2.6	5.1	11.1
1999.....	*	*	*	*	0.3	0.6	1.1	2.0	4.2	9.7

All races 3/, male

2002.....	*	*	*	0.4	0.7	1.4	2.9	5.0	9.7	19.6
2001.....	*	*	*	*	0.4	0.7	1.9	3.3	6.1	11.8
2000.....	*	*	*	0.3	*	0.9	1.5	2.8	5.7	10.5
1999.....	*	*	*	*	*	0.6	1.0	2.5	4.7	9.1

All races 3/, female

2002.....	*	*	*	0.4	0.6	1.2	2.2	4.9	9.1	20.4
2001.....	*	*	*	*	0.5	0.8	1.7	2.9	6.2	12.7
2000.....	*	*	*	*	*	0.7	1.5	2.5	4.8	11.4
1999.....	*	*	*	*	*	0.7	1.1	1.7	4.0	9.9

White, both sexes

2002.....	*	*	0.2	0.4	0.6	1.4	2.6	5.2	9.9	21.1
2001.....	*	*	0.2	0.2	0.4	0.7	1.9	3.2	6.3	13.1
2000.....	*	*	*	0.3	0.3	0.8	1.5	2.7	5.4	11.5
1999.....	*	*	*	*	0.3	0.7	1.1	2.1	4.4	10.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Tuberculosis (A16-A19)										
All races 3/, both sexes										
2002.....	0.1	0.2	0.2	0.3	0.5	0.9	0.9	1.5	2.3	2.6
2001.....	0.1	0.2	0.3	0.4	0.5	0.6	1.0	1.4	2.0	2.8
2000.....	0.1	0.2	0.3	0.2	0.5	0.7	1.0	1.8	2.4	2.7
1999.....	0.1	0.3	0.3	0.5	0.6	0.7	1.2	1.8	2.4	4.2
All races 3/, male										
2002.....	0.2	0.2	0.4	0.5	0.7	1.2	1.2	1.8	3.2	4.0
2001.....	*	0.3	0.4	0.7	0.7	0.9	1.5	1.7	2.4	4.4
2000.....	*	0.4	0.5	0.4	0.7	1.0	1.1	2.3	3.8	4.6
1999.....	0.2	0.4	0.5	0.8	0.9	1.1	1.7	2.4	3.7	7.1
All races 3/, female										
2002.....	*	*	*	*	0.4	0.5	0.6	1.3	1.7	2.1
2001.....	*	*	*	*	*	0.4	0.7	1.1	1.8	2.2
2000.....	*	*	*	*	*	0.5	0.9	1.4	1.6	2.0
1999.....	*	*	*	*	*	0.4	0.7	1.5	1.7	3.0
White, both sexes										
2002.....	*	*	0.1	0.2	0.4	0.5	0.6	1.2	1.8	2.1
2001.....	*	0.1	0.1	0.3	0.3	0.5	0.7	1.1	1.4	2.2
2000.....	*	*	0.2	*	0.3	0.5	0.6	1.2	1.7	2.2
1999.....	*	0.2	0.2	0.3	0.4	0.6	0.8	1.3	2.0	3.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Respiratory tuberculosis (A16)										
All races 3/, both sexes										
2002.....	0.1	0.1	0.2	0.2	0.4	0.6	0.7	1.2	2.0	2.3
2001.....	*	0.1	0.2	0.4	0.4	0.5	0.8	1.1	1.6	2.4
2000.....	0.1	0.1	0.2	0.2	0.3	0.5	0.8	1.4	2.1	2.3
1999.....	0.1	0.2	0.2	0.4	0.5	0.6	0.9	1.5	2.1	3.7
All races 3/, male										
2002.....	0.2	*	0.3	0.4	0.5	1.0	0.9	1.3	2.8	3.5
2001.....	*	0.2	0.3	0.6	0.7	0.7	1.2	1.5	2.0	3.8
2000.....	*	0.3	0.4	0.3	0.6	0.9	0.9	1.9	3.2	4.1
1999.....	*	0.3	0.4	0.6	0.9	0.9	1.4	2.0	3.3	6.4
All races 3/, female										
2002.....	*	*	*	*	*	*	0.5	1.1	1.5	1.8
2001.....	*	*	*	*	*	*	0.5	0.8	1.4	1.8
2000.....	*	*	*	*	*	*	0.6	1.1	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	1.2	1.4	2.6
White, both sexes										
2002.....	*	*	*	*	0.3	0.4	0.5	0.9	1.5	1.8
2001.....	*	*	*	0.2	0.3	0.4	0.5	0.9	1.2	1.8
2000.....	*	*	0.2	*	*	0.3	0.5	1.0	1.5	1.9
1999.....	*	0.1	0.1	0.2	0.4	0.5	0.6	1.1	1.7	3.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Zoonotic and other bacterial diseases (A20-A49)										
All races 3/, both sexes										
2002.....	2.5	4.5	6.7	10.6	16.8	28.1	45.1	73.3	113.9	210.5
2001.....	2.4	4.3	6.6	10.4	16.3	26.6	42.6	70.3	108.3	213.7
2000.....	2.6	4.2	6.5	9.9	15.9	25.2	40.0	66.9	109.2	222.5
1999.....	2.5	4.1	6.1	9.6	15.2	26.5	40.5	67.5	108.5	231.2
All races 3/, male										
2002.....	2.7	5.1	7.4	11.3	18.6	30.2	50.7	84.9	132.3	222.5
2001.....	2.6	4.7	7.2	11.3	17.6	29.4	48.7	82.3	127.8	233.5
2000.....	2.7	4.8	7.6	10.7	17.4	28.2	45.0	77.3	127.0	233.1
1999.....	2.9	4.6	6.7	10.3	16.9	30.0	46.3	79.3	127.7	246.7
All races 3/, female										
2002.....	2.3	4.0	6.0	10.0	15.2	26.3	40.7	65.1	102.7	205.3
2001.....	2.2	3.9	6.0	9.5	15.1	24.1	37.8	62.0	96.5	205.4
2000.....	2.5	3.6	5.5	9.1	14.6	22.7	36.1	59.6	98.7	218.2
1999.....	2.1	3.6	5.5	9.0	13.7	23.6	35.9	59.3	97.2	225.0
White, both sexes										
2002.....	2.1	3.8	5.3	8.9	14.1	24.7	41.0	67.6	107.5	199.4
2001.....	1.9	3.4	5.3	8.5	14.0	23.4	38.4	65.0	102.8	202.9
2000.....	2.2	3.2	5.3	8.2	13.3	21.8	36.0	60.8	101.6	209.6
1999.....	2.0	3.1	4.9	7.8	13.0	23.1	35.6	60.9	100.8	216.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Zoonotic and other bacterial diseases (A20-A49)										
White, male										
2002.....	2.2	4.3	5.9	9.6	15.9	26.3	46.3	77.5	124.3	211.9
2001.....	2.2	3.7	5.8	9.1	15.3	25.7	43.5	76.2	121.5	221.0
2000.....	2.3	3.7	6.0	9.0	14.8	24.4	41.1	70.0	119.3	219.5
1999.....	2.3	3.5	5.4	8.2	13.9	25.9	40.6	72.4	117.9	230.1
White, female										
2002.....	1.9	3.3	4.8	8.3	12.4	23.3	36.6	60.5	97.3	194.0
2001.....	1.7	3.2	4.8	8.0	12.7	21.4	34.4	57.1	91.5	195.4
2000.....	2.0	2.7	4.6	7.5	12.0	19.5	31.9	54.3	91.0	205.6
1999.....	1.7	2.8	4.5	7.4	12.1	20.6	31.7	52.8	90.6	210.5
Black, both sexes										
2002.....	6.1	10.1	17.8	28.1	42.6	63.2	94.6	151.5	212.7	376.3
2001.....	5.9	11.0	17.9	28.6	40.5	59.8	92.3	143.8	198.0	373.1
2000.....	6.1	11.5	17.9	25.4	41.3	61.3	89.6	150.3	221.7	402.3
1999.....	6.6	11.7	16.6	26.7	37.7	61.3	98.2	151.0	219.6	428.2
Black, male										
2002.....	6.8	10.9	21.3	30.2	47.1	73.1	110.5	191.1	265.2	402.0
2001.....	6.1	12.2	20.1	35.2	43.7	69.9	114.1	175.6	243.2	435.7
2000.....	6.1	13.1	23.0	29.0	46.2	71.3	101.0	188.1	256.3	441.7
1999.....	8.1	13.7	19.5	31.2	48.4	75.6	120.1	176.7	286.0	472.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Zoonotic and other bacterial diseases (A20-A49)										
Black, female										
2002.....	5.5	9.4	14.9	26.3	39.0	55.9	84.0	127.9	185.4	366.4
2001.....	5.7	9.9	16.0	23.2	37.9	52.4	77.8	124.8	175.0	349.8
2000.....	6.1	10.1	13.6	22.4	37.4	54.0	82.1	127.6	204.2	387.6
1999.....	5.3	9.9	14.1	23.0	29.3	50.8	83.7	135.6	186.7	411.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
All races 3/, both sexes										
2002.....	2.3	4.2	6.2	10.0	16.0	27.0	43.1	70.0	109.4	203.0
2001.....	2.3	4.0	6.2	9.8	15.5	25.3	40.9	67.3	103.9	205.9
2000.....	2.5	3.9	6.1	9.4	15.2	24.2	38.4	64.0	105.0	215.7
1999.....	2.2	3.7	5.6	9.0	14.4	24.7	38.3	63.8	102.9	220.7
All races 3/, male										
2002.....	2.5	4.7	6.8	10.6	17.8	29.1	48.2	80.6	127.0	213.9
2001.....	2.5	4.3	6.7	10.6	16.6	27.8	46.7	78.6	122.3	224.3
2000.....	2.6	4.4	7.0	10.1	16.7	26.9	43.2	73.9	121.7	223.0
1999.....	2.6	4.2	6.2	9.7	15.7	27.7	43.7	74.8	120.6	233.5
All races 3/, female										
2002.....	2.2	3.8	5.6	9.5	14.5	25.2	39.0	62.5	98.7	198.3
2001.....	2.1	3.8	5.7	9.1	14.4	23.1	36.2	59.5	92.9	198.3
2000.....	2.3	3.4	5.2	8.7	13.8	21.8	34.6	57.1	95.2	212.7
1999.....	1.9	3.3	5.1	8.4	13.1	22.1	34.1	56.2	92.5	215.5
White, both sexes										
2002.....	1.9	3.6	5.0	8.4	13.5	23.6	39.2	64.4	103.1	191.9
2001.....	1.8	3.2	4.9	8.1	13.2	22.2	36.8	62.1	98.5	195.3
2000.....	2.0	3.0	4.9	7.8	12.6	20.7	34.5	58.0	97.4	203.0
1999.....	1.7	2.8	4.5	7.2	12.1	21.3	33.6	57.6	95.5	205.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
White, male										
2002.....	2.0	4.0	5.4	8.9	15.3	25.2	44.0	73.4	119.0	202.9
2001.....	2.1	3.3	5.3	8.5	14.4	24.4	41.6	72.5	116.0	211.8
2000.....	2.2	3.5	5.5	8.4	14.0	23.1	39.4	66.7	114.0	209.8
1999.....	2.0	3.1	4.9	7.6	12.8	23.7	38.2	68.3	111.6	217.4
White, female										
2002.....	1.9	3.1	4.5	7.8	11.8	22.3	35.2	58.0	93.4	187.1
2001.....	1.6	3.0	4.5	7.7	12.2	20.4	32.9	54.8	87.9	188.3
2000.....	1.9	2.5	4.3	7.2	11.3	18.7	30.6	51.9	87.6	200.2
1999.....	1.4	2.4	4.1	6.9	11.5	19.2	29.9	50.1	86.1	201.4
Black, both sexes										
2002.....	5.5	9.5	17.0	27.2	40.9	62.0	90.6	146.5	208.8	367.0
2001.....	5.6	10.6	16.9	27.1	38.4	57.4	89.6	139.1	193.1	364.8
2000.....	5.8	11.0	16.9	24.3	40.2	59.7	86.6	146.4	216.3	392.6
1999.....	6.2	11.0	15.5	25.4	36.3	57.9	94.6	142.8	209.5	412.3
Black, male										
2002.....	6.1	10.3	20.0	29.0	45.1	71.6	105.5	183.7	259.8	393.5
2001.....	5.7	11.6	19.1	33.3	41.5	66.3	111.1	171.0	236.8	424.4
2000.....	5.8	12.4	21.3	27.8	45.1	68.9	97.0	182.9	250.4	427.8
1999.....	7.7	12.8	18.1	29.7	46.5	71.4	115.0	166.2	266.8	453.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
Black, female										
2002.....	5.0	8.9	14.4	25.7	37.6	54.9	80.7	124.3	182.2	356.9
2001.....	5.4	9.7	15.0	22.1	36.0	50.9	75.3	120.1	170.9	342.7
2000.....	5.8	9.7	13.2	21.5	36.3	52.8	79.7	124.5	199.1	379.5
1999.....	5.0	9.4	13.3	21.9	28.3	47.9	81.0	128.7	181.1	396.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)										
All races 3/, both sexes										
2002.....	0.2	0.2	0.4	0.5	0.7	1.0	2.0	3.2	4.3	7.4
2001.....	0.1	0.2	0.4	0.5	0.8	1.2	1.6	2.9	4.3	7.5
2000.....	0.1	0.2	0.4	0.5	0.7	1.0	1.5	2.8	4.0	6.7
1999.....	0.2	0.3	0.4	0.6	0.8	1.8	2.2	3.5	5.4	10.3
All races 3/, male										
2002.....	0.2	0.3	0.5	0.7	0.8	1.1	2.4	4.2	5.2	8.5
2001.....	*	0.3	0.5	0.6	1.0	1.5	1.9	3.6	5.4	8.9
2000.....	*	0.3	0.6	0.6	0.8	1.4	1.8	3.3	5.2	9.9
1999.....	0.2	0.4	0.4	0.6	1.1	2.2	2.6	4.4	7.0	13.1
All races 3/, female										
2002.....	*	0.2	0.3	0.4	0.7	1.0	1.6	2.5	3.8	6.9
2001.....	*	*	0.3	0.3	0.7	1.0	1.5	2.4	3.6	6.9
2000.....	*	*	*	0.3	0.7	0.8	1.3	2.4	3.4	5.3
1999.....	0.2	0.3	0.4	0.6	0.6	1.4	1.8	2.9	4.5	9.2
White, both sexes										
2002.....	0.1	0.2	0.3	0.5	0.6	1.0	1.8	3.1	4.3	7.3
2001.....	*	0.2	0.3	0.4	0.7	1.1	1.5	2.8	4.2	7.4
2000.....	*	0.2	0.3	0.4	0.7	1.0	1.4	2.7	4.0	6.4
1999.....	0.2	0.3	0.4	0.5	0.8	1.6	2.0	3.1	5.1	9.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)										
White, male										
2002.....	*	0.3	0.4	0.6	0.6	1.1	2.2	4.1	5.1	8.7
2001.....	*	0.3	0.4	0.5	0.9	1.3	1.8	3.6	5.4	8.9
2000.....	*	0.2	0.5	0.6	0.7	1.2	1.6	3.1	5.2	9.5
1999.....	*	0.3	0.4	0.5	1.1	2.0	2.4	4.0	6.2	12.6
White, female										
2002.....	*	*	0.3	0.4	0.6	1.0	1.4	2.4	3.7	6.7
2001.....	*	*	*	*	0.6	0.9	1.4	2.1	3.5	6.8
2000.....	*	*	*	*	0.6	0.7	1.2	2.4	3.2	5.1
1999.....	*	*	0.3	0.5	0.6	1.3	1.8	2.5	4.5	8.8
Black, both sexes										
2002.....	*	*	*	*	*	*	4.0	5.0	*	9.2
2001.....	*	*	1.0	1.4	*	2.3	2.7	4.5	*	8.0
2000.....	*	*	*	*	*	*	2.9	3.9	*	9.7
1999.....	*	*	*	*	*	3.4	3.5	8.0	10.1	15.9
Black, male										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	10.0	19.2	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

Black, female

2002.....	*	*	*	*	*	*	*	*	*	9.4
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	6.9	*	14.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral diseases (A80-B34)										
All races 3/, both sexes										
2002.....	15.9	18.4	15.9	10.6	8.3	7.9	7.8	7.8	7.7	10.1
2001.....	16.5	17.5	14.2	10.3	9.1	8.1	7.6	7.3	7.0	8.3
2000.....	16.1	18.1	14.2	10.1	8.7	7.9	8.0	8.1	8.1	9.6
1999.....	17.2	17.1	12.9	9.9	7.5	8.2	8.1	8.0	8.7	11.6
All races 3/, male										
2002.....	23.2	28.2	25.5	16.3	12.1	10.7	9.2	10.0	9.1	11.7
2001.....	24.4	27.3	22.9	16.1	12.5	10.0	9.1	8.3	8.9	9.3
2000.....	24.0	28.4	22.7	15.7	12.4	10.2	9.5	10.7	9.9	10.8
1999.....	26.2	27.0	20.8	15.5	10.5	10.7	9.8	9.3	10.9	13.3
All races 3/, female										
2002.....	8.8	9.0	6.7	5.2	4.8	5.4	6.7	6.3	6.9	9.3
2001.....	8.7	8.1	5.9	4.8	6.0	6.4	6.5	6.7	5.9	7.9
2000.....	8.3	8.1	6.0	4.8	5.3	6.0	6.9	6.3	7.0	9.1
1999.....	8.3	7.5	5.3	4.7	4.7	6.0	6.7	7.1	7.4	10.9
White, both sexes										
2002.....	9.8	12.5	10.8	7.5	6.0	5.9	6.5	7.0	7.4	10.3
2001.....	10.2	11.7	9.5	7.4	6.4	6.3	6.5	6.8	6.6	8.4
2000.....	10.0	12.0	9.3	6.9	6.2	6.2	6.8	7.6	7.8	9.6
1999.....	10.4	10.9	8.7	6.6	5.5	6.4	6.8	7.4	8.6	11.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral diseases (A80-B34)										
White, male										
2002.....	15.4	19.9	17.4	11.6	8.4	7.8	7.5	8.9	8.6	11.7
2001.....	16.5	19.0	15.9	11.7	8.5	7.8	7.9	7.7	8.3	9.7
2000.....	15.8	19.6	15.3	10.8	8.6	7.6	7.6	9.9	9.3	10.9
1999.....	16.8	17.4	14.3	10.2	7.9	8.3	8.1	8.6	10.9	12.7
White, female										
2002.....	4.1	5.1	4.3	3.6	3.8	4.1	5.8	5.7	6.7	9.7
2001.....	4.0	4.5	3.2	3.3	4.3	5.0	5.4	6.1	5.6	7.9
2000.....	4.3	4.5	3.5	3.2	4.0	5.0	6.1	5.9	6.8	9.1
1999.....	4.0	4.4	3.1	3.2	3.4	4.9	5.7	6.6	7.2	11.1
Black, both sexes										
2002.....	60.5	65.2	59.0	38.4	27.5	24.2	18.4	16.3	9.4	7.4
2001.....	62.3	63.5	54.4	36.3	32.7	21.8	15.9	12.9	10.0	6.1
2000.....	60.6	66.4	55.8	36.7	27.8	21.0	19.3	13.7	11.3	8.8
1999.....	66.5	66.5	49.7	39.3	23.0	22.5	19.6	12.7	9.0	11.5
Black, male										
2002.....	83.9	97.6	98.4	63.1	45.7	39.6	27.7	24.3	*	*
2001.....	87.8	97.7	86.4	60.2	51.5	31.5	21.9	17.2	*	*
2000.....	88.6	104.1	91.6	61.9	46.0	33.8	30.8	19.8	18.5	*
1999.....	100.2	108.7	82.0	66.3	35.9	33.2	28.8	15.7	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral diseases (A80-B34)										
Black, female										
2002.....	39.9	37.1	25.6	18.1	13.2	12.8	12.1	11.5	*	*
2001.....	39.7	33.9	27.1	16.7	17.9	14.6	11.9	10.3	8.2	*
2000.....	35.9	33.7	25.3	16.1	13.4	11.5	11.8	10.0	*	8.6
1999.....	36.8	30.1	22.2	17.3	13.0	14.6	13.5	10.9	8.6	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Herpesviral (herpes simplex) infections (B00)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	0.4	*	0.5	0.5
2001.....	*	*	*	*	*	*	0.3	0.3	0.4	0.5
2000.....	*	*	*	*	*	*	*	0.3	0.5	*
1999.....	*	*	*	*	*	*	0.2	0.3	*	0.5

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	0.5	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	*	0.4	*	0.5	0.6
2001.....	*	*	*	*	*	*	0.3	0.3	0.4	0.5
2000.....	*	*	*	*	*	*	*	0.4	0.5	*
1999.....	*	*	*	*	*	*	0.3	0.3	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Zoster (herpes zoster)(B02)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	1.6
2000.....	*	*	*	*	*	*	*	*	*	1.7
1999.....	*	*	*	*	*	*	*	*	0.6	2.0
All races 3/, male										
2002.....	*	*	*	*	*	*	*	*	*	1.5
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	1.8
2000.....	*	*	*	*	*	*	*	*	*	1.9
1999.....	*	*	*	*	*	*	*	*	*	2.4
White, both sexes										
2002.....	*	*	*	*	*	*	*	*	*	1.8
2001.....	*	*	*	*	*	*	*	*	*	1.8
2000.....	*	*	*	*	*	*	*	*	*	1.8
1999.....	*	*	*	*	*	*	*	*	0.6	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)										
All races 3/, both sexes										
2002.....	2.3	5.6	6.1	4.3	3.7	3.9	4.5	4.7	4.4	3.9
2001.....	2.6	5.3	5.2	4.1	3.8	4.3	4.9	5.1	4.6	3.8
2000.....	2.5	5.2	4.8	3.7	3.8	4.3	4.9	5.4	5.2	4.0
1999.....	2.3	4.4	3.8	3.3	3.3	4.5	5.1	5.3	5.4	4.7
All races 3/, male										
2002.....	3.2	8.5	9.5	6.2	4.6	4.3	4.7	5.7	4.8	5.0
2001.....	3.7	8.1	8.2	5.9	4.5	4.7	4.9	5.1	5.5	4.4
2000.....	3.5	8.0	7.4	5.2	4.7	4.6	5.0	6.5	6.2	5.7
1999.....	3.5	6.7	5.9	4.7	3.8	4.9	5.4	6.1	7.3	6.9
All races 3/, female										
2002.....	1.4	2.8	2.8	2.5	2.9	3.4	4.4	4.1	4.2	3.3
2001.....	1.5	2.5	2.4	2.5	3.3	3.9	4.9	5.0	4.1	3.5
2000.....	1.5	2.5	2.4	2.2	3.1	4.1	4.9	4.6	4.7	3.3
1999.....	1.2	2.2	1.7	2.1	2.8	4.1	4.9	4.7	4.2	3.9
White, both sexes										
2002.....	2.2	5.5	5.8	3.8	3.2	3.4	4.1	4.5	4.3	3.7
2001.....	2.6	5.2	4.7	3.7	3.5	3.8	4.4	4.9	4.4	3.6
2000.....	2.4	4.9	4.3	3.0	3.4	3.9	4.6	5.3	5.0	3.9
1999.....	2.2	4.2	3.3	2.9	2.9	4.0	4.6	4.9	5.3	4.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)										
White, male										
2002.....	3.1	8.3	9.1	5.5	3.9	4.0	4.3	5.5	4.8	4.9
2001.....	3.7	8.1	7.6	5.4	4.1	4.4	4.6	5.2	5.6	4.5
2000.....	3.4	7.6	6.6	4.4	4.2	4.0	4.6	6.4	5.9	5.7
1999.....	3.3	6.1	5.2	4.1	3.6	4.6	5.1	5.8	7.7	6.5
White, female										
2002.....	1.3	2.6	2.7	2.2	2.6	2.8	3.9	3.8	4.0	3.2
2001.....	1.4	2.4	2.0	2.1	2.9	3.3	4.2	4.6	3.7	3.2
2000.....	1.4	2.3	2.0	1.8	2.7	3.7	4.5	4.5	4.4	3.2
1999.....	1.1	2.2	1.5	1.7	2.3	3.5	4.3	4.3	3.9	3.7
Black, both sexes										
2002.....	2.8	7.2	9.0	7.7	6.3	6.1	6.2	6.0	*	*
2001.....	3.3	6.8	9.4	7.3	6.0	5.1	6.2	6.1	*	*
2000.....	3.1	7.3	8.8	6.9	4.6	5.2	6.3	5.7	7.1	*
1999.....	3.5	7.0	6.7	5.7	4.2	6.6	7.2	6.8	*	6.4
Black, male										
2002.....	3.8	10.5	14.6	12.4	8.1	8.0	*	*	*	*
2001.....	4.7	10.1	13.9	10.2	7.3	*	*	*	*	*
2000.....	4.2	11.0	13.9	10.4	5.2	6.8	7.4	*	*	*
1999.....	5.1	11.9	11.2	7.6	4.5	5.5	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)										
Black, female										
2002.....	1.9	4.4	4.2	3.9	4.8	4.6	6.3	5.8	*	*
2001.....	2.0	3.9	5.5	4.9	5.0	5.4	6.7	7.2	*	*
2000.....	2.1	4.0	4.4	4.0	4.1	4.0	5.5	*	*	*
1999.....	2.0	2.8	2.8	4.2	4.0	7.5	7.7	6.6	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All races 3/, both sexes

2002.....	13.4	12.6	9.6	5.9	4.0	2.9	1.4	1.0	0.5	*
2001.....	13.7	12.1	8.7	5.8	4.5	2.7	1.4	0.9	0.4	*
2000.....	13.5	12.7	9.0	5.9	4.1	2.6	1.7	0.9	*	*
1999.....	14.6	12.4	8.7	5.9	3.5	2.7	1.5	0.7	*	*

All races 3/, male

2002.....	19.8	19.4	15.7	9.8	6.8	5.0	2.6	1.7	*	*
2001.....	20.5	19.0	14.4	9.8	7.1	4.2	2.7	1.8	*	*
2000.....	20.4	20.3	15.0	10.1	7.0	4.5	2.9	1.8	*	*
1999.....	22.6	20.0	14.5	10.2	5.9	4.8	2.7	1.3	*	*

All races 3/, female

2002.....	7.2	5.9	3.6	2.3	1.5	1.0	0.5	*	*	*
2001.....	7.0	5.4	3.3	2.0	2.0	1.4	0.4	*	*	*
2000.....	6.7	5.3	3.4	2.0	1.5	1.0	0.7	*	*	*
1999.....	6.9	5.0	3.1	1.9	1.2	1.0	0.6	*	*	*

White, both sexes

2002.....	7.3	6.8	4.6	3.3	2.2	1.3	0.6	0.5	*	*
2001.....	7.5	6.4	4.5	3.3	2.0	1.2	0.8	0.5	*	*
2000.....	7.5	6.9	4.7	3.4	2.1	1.3	0.7	0.5	*	*
1999.....	8.1	6.5	4.9	3.1	1.9	1.4	0.7	0.4	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

White, male

2002.....	12.0	11.4	8.0	5.8	3.8	2.4	1.2	0.8	*	*
2001.....	12.6	10.8	8.0	5.9	3.4	2.1	1.7	1.0	*	*
2000.....	12.3	11.7	8.3	6.0	3.7	2.5	1.3	1.0	*	*
1999.....	13.4	11.0	8.6	5.6	3.5	2.7	1.2	0.8	*	*

White, female

2002.....	2.6	2.2	1.3	0.9	0.7	*	*	*	*	*
2001.....	2.4	2.0	1.0	0.9	0.7	0.5	*	*	*	*
2000.....	2.7	2.1	1.2	0.9	0.7	*	*	*	*	*
1999.....	2.8	2.0	1.2	0.7	*	*	*	*	*	*

Black, both sexes

2002.....	57.3	57.4	49.7	30.2	20.7	17.5	10.5	7.0	*	*
2001.....	58.8	56.5	44.7	28.9	26.2	16.3	8.5	6.4	*	*
2000.....	57.3	58.7	46.6	29.3	22.0	15.1	11.6	6.2	*	*
1999.....	62.8	59.0	42.6	32.7	18.3	15.4	11.2	4.8	*	*

Black, male

2002.....	79.6	86.3	83.5	50.0	37.0	31.4	19.4	14.7	*	*
2001.....	82.8	87.5	72.2	49.9	43.5	26.4	15.5	13.0	*	*
2000.....	84.1	92.7	77.4	51.0	39.5	26.3	22.4	11.8	*	*
1999.....	94.9	96.1	70.6	57.5	30.6	27.3	21.4	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)										
Black, female										
2002.....	37.6	32.4	21.0	13.9	7.9	7.2	4.5	*	*	*
2001.....	37.5	29.7	21.2	11.7	12.6	8.8	*	*	*	*
2000.....	33.7	29.2	20.4	11.5	8.2	6.9	4.4	*	*	*
1999.....	34.6	26.9	18.8	12.5	8.7	6.5	4.5	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

All races 3/, both sexes

2002.....	5.3	5.1	3.9	2.3	1.4	1.0	0.4	0.4	*	*
2001.....	5.3	4.9	3.4	2.2	1.5	0.9	0.6	0.3	*	*
2000.....	5.4	5.2	3.5	2.0	1.4	0.9	0.6	0.4	*	*
1999.....	5.8	5.2	3.6	1.9	1.1	0.9	0.6	0.4	*	*

All races 3/, male

2002.....	7.9	7.8	6.5	3.7	2.2	1.7	0.9	0.7	*	*
2001.....	8.0	7.6	5.6	3.7	2.3	1.2	1.1	*	*	*
2000.....	7.9	8.2	5.8	3.3	2.4	1.7	1.1	*	*	*
1999.....	8.9	8.2	5.9	3.4	1.9	1.6	1.0	0.7	*	*

All races 3/, female

2002.....	2.7	2.4	1.4	0.9	0.7	0.4	*	*	*	*
2001.....	2.7	2.3	1.3	0.8	0.8	0.5	*	*	*	*
2000.....	2.9	2.2	1.2	0.9	0.5	*	*	*	*	*
1999.....	2.7	2.2	1.3	0.6	0.5	*	*	*	*	*

White, both sexes

2002.....	3.0	2.7	1.9	1.3	0.9	0.5	*	*	*	*
2001.....	3.1	2.7	1.7	1.3	0.7	0.4	0.3	*	*	*
2000.....	3.0	2.9	1.9	1.2	0.8	0.5	*	*	*	*
1999.....	3.2	2.6	2.1	1.1	0.7	0.5	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)										
White, male										
2002.....	4.8	4.6	3.2	2.3	1.3	0.9	*	*	*	*
2001.....	5.0	4.6	3.0	2.3	1.2	0.6	0.6	*	*	*
2000.....	4.9	4.8	3.4	1.9	1.3	1.0	*	*	*	*
1999.....	5.2	4.4	3.6	1.9	1.2	0.9	*	*	*	*
White, female										
2002.....	1.2	0.9	0.6	0.4	0.5	*	*	*	*	*
2001.....	1.1	0.9	0.5	0.4	*	*	*	*	*	*
2000.....	1.2	0.9	0.5	0.4	*	*	*	*	*	*
1999.....	1.2	0.9	0.6	*	*	*	*	*	*	*
Black, both sexes										
2002.....	21.6	23.0	20.3	10.9	6.6	6.1	2.7	*	*	*
2001.....	21.9	22.2	17.5	10.7	8.5	5.2	3.6	*	*	*
2000.....	22.4	23.5	16.5	9.9	7.4	5.1	4.4	*	*	*
1999.....	24.5	25.1	16.8	10.1	5.8	4.6	4.3	*	*	*
Black, male										
2002.....	31.3	34.8	35.1	17.7	11.1	9.5	*	*	*	*
2001.....	31.5	33.3	28.3	18.3	13.8	8.1	6.7	*	*	*
2000.....	31.4	36.9	28.0	16.3	13.3	9.4	9.0	*	*	*
1999.....	37.7	40.3	28.2	18.3	8.9	8.8	7.8	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

Black, female

2002.....	13.1	12.7	7.7	5.3	3.1	*	*	*	*	*
2001.....	13.5	12.6	8.3	4.5	4.3	*	*	*	*	*
2000.....	14.4	11.9	6.7	4.6	*	*	*	*	*	*
1999.....	12.8	12.0	7.2	3.3	3.3	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)

All races 3/, both sexes

2002.....	0.9	0.9	0.7	0.4	0.3	*	*	*	*	*
2001.....	1.0	0.8	0.6	0.6	0.4	0.2	*	*	*	*
2000.....	1.0	0.9	0.7	0.5	0.4	0.2	*	*	*	*
1999.....	1.1	0.9	0.7	0.6	0.3	*	*	*	*	*

All races 3/, male

2002.....	1.5	1.5	1.2	0.7	0.6	*	*	*	*	*
2001.....	1.7	1.2	1.1	0.9	0.8	*	*	*	*	*
2000.....	1.7	1.6	1.2	1.0	0.6	*	*	*	*	*
1999.....	1.8	1.5	1.2	1.1	0.5	*	*	*	*	*

All races 3/, female

2002.....	0.3	0.3	0.2	*	*	*	*	*	*	*
2001.....	0.3	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	0.2	0.2	*	*	*	*	*	*	*

White, both sexes

2002.....	0.6	0.6	0.4	0.3	0.2	*	*	*	*	*
2001.....	0.7	0.5	0.4	0.3	0.3	*	*	*	*	*
2000.....	0.7	0.7	0.5	0.3	0.2	*	*	*	*	*
1999.....	0.7	0.6	0.5	0.5	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)										
White, male										
2002.....	1.0	1.2	0.7	0.6	*	*	*	*	*	*
2001.....	1.3	0.9	0.8	0.6	0.5	*	*	*	*	*
2000.....	1.2	1.2	0.9	0.7	0.5	*	*	*	*	*
1999.....	1.3	1.1	0.9	0.9	*	*	*	*	*	*
White, female										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002.....	3.1	3.1	3.0	1.3	*	*	*	*	*	*
2001.....	3.2	2.7	2.2	2.7	2.1	*	*	*	*	*
2000.....	3.5	3.1	2.7	1.9	*	*	*	*	*	*
1999.....	3.4	2.8	2.7	2.4	*	*	*	*	*	*
Black, male										
2002.....	4.7	4.9	5.0	*	*	*	*	*	*	*
2001.....	5.3	4.0	3.8	4.5	4.3	*	*	*	*	*
2000.....	5.7	5.1	4.5	3.8	*	*	*	*	*	*
1999.....	6.0	5.1	4.4	3.6	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)

All races 3/, both sexes

2002.....	1.4	1.1	0.8	0.6	0.4	0.3	*	*	*	*
2001.....	1.2	1.1	0.8	0.6	0.5	0.4	*	*	*	*
2000.....	1.5	1.3	1.0	0.7	0.4	0.3	0.2	*	*	*
1999.....	1.6	1.2	0.9	0.7	0.4	0.3	*	*	*	*

All races 3/, male

2002.....	2.1	1.8	1.4	1.1	0.9	0.6	*	*	*	*
2001.....	1.9	1.8	1.4	1.1	0.8	0.6	*	*	*	*
2000.....	2.5	2.2	1.7	1.3	0.8	*	*	*	*	*
1999.....	2.6	2.1	1.6	1.2	0.7	0.5	*	*	*	*

All races 3/, female

2002.....	0.6	0.5	0.2	*	*	*	*	*	*	*
2001.....	0.5	0.4	0.3	*	*	*	*	*	*	*
2000.....	0.6	0.5	0.3	*	*	*	*	*	*	*
1999.....	0.6	0.4	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.9	0.8	0.5	0.4	0.3	*	*	*	*	*
2001.....	0.7	0.7	0.5	0.4	0.3	*	*	*	*	*
2000.....	1.0	0.7	0.6	0.5	*	*	*	*	*	*
1999.....	1.0	0.9	0.6	0.3	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)										
White, male										
2002.....	1.5	1.4	0.9	0.7	0.5	*	*	*	*	*
2001.....	1.2	1.2	0.9	0.8	0.4	*	*	*	*	*
2000.....	1.7	1.3	1.0	0.8	*	*	*	*	*	*
1999.....	1.7	1.6	1.1	0.6	*	*	*	*	*	*
White, female										
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002.....	5.0	3.8	3.5	3.1	2.2	*	*	*	*	*
2001.....	5.0	4.4	3.5	2.8	3.0	2.2	*	*	*	*
2000.....	5.6	5.7	4.7	3.5	2.3	*	*	*	*	*
1999.....	6.5	4.2	3.7	4.0	2.0	*	*	*	*	*
Black, male										
2002.....	7.2	5.6	6.0	5.5	4.4	*	*	*	*	*
2001.....	7.6	7.0	5.5	4.7	4.3	*	*	*	*	*
2000.....	9.0	9.2	8.2	6.4	4.3	*	*	*	*	*
1999.....	10.1	7.0	6.2	7.1	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

All races 3/, both sexes

2002.....	3.3	3.1	2.4	1.6	1.1	0.8	0.4	0.3	*	*
2001.....	3.3	3.0	2.2	1.4	1.3	0.7	0.4	0.3	*	*
2000.....	3.0	2.9	2.3	1.7	1.2	0.7	0.4	0.3	*	*
1999.....	3.3	2.9	2.2	1.7	1.1	0.9	0.4	*	*	*

All races 3/, male

2002.....	4.4	4.6	3.8	2.6	2.0	1.5	0.5	*	*	*
2001.....	4.6	4.5	3.6	2.5	2.0	1.1	0.7	*	*	*
2000.....	4.4	4.4	3.6	2.8	1.8	1.1	0.7	0.7	*	*
1999.....	4.7	4.5	3.5	2.7	1.9	1.6	0.7	*	*	*

All races 3/, female

2002.....	2.1	1.7	1.1	0.7	0.4	*	*	*	*	*
2001.....	2.1	1.6	0.9	0.5	0.6	*	*	*	*	*
2000.....	1.7	1.4	1.0	0.6	0.6	*	*	*	*	*
1999.....	1.9	1.3	0.9	0.7	0.4	*	*	*	*	*

White, both sexes

2002.....	1.5	1.3	1.0	0.7	0.5	0.3	*	*	*	*
2001.....	1.6	1.3	0.9	0.7	0.5	0.3	*	*	*	*
2000.....	1.5	1.3	1.0	0.9	0.5	0.3	*	*	*	*
1999.....	1.5	1.2	1.0	0.7	0.6	0.4	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

White, male

2002.....	2.4	2.0	1.8	1.3	0.9	0.5	*	*	*	*
2001.....	2.5	2.1	1.7	1.3	1.0	0.5	*	*	*	*
2000.....	2.3	2.2	1.7	1.5	0.8	*	*	*	*	*
1999.....	2.4	2.0	1.7	1.2	1.1	0.8	*	*	*	*

White, female

2002.....	0.7	0.6	0.3	*	*	*	*	*	*	*
2001.....	0.7	0.6	0.3	*	*	*	*	*	*	*
2000.....	0.7	0.5	0.4	*	*	*	*	*	*	*
1999.....	0.7	0.5	*	*	*	*	*	*	*	*

Black, both sexes

2002.....	15.7	16.8	13.7	9.6	7.0	5.8	3.2	*	*	*
2001.....	15.7	15.5	12.9	7.8	8.0	4.1	*	*	*	*
2000.....	14.0	14.7	13.4	9.0	7.2	4.6	2.9	*	*	*
1999.....	16.2	15.5	13.1	10.4	5.7	5.6	3.2	*	*	*

Black, male

2002.....	20.1	25.2	21.6	15.5	12.5	11.2	*	*	*	*
2001.....	20.6	23.5	20.9	14.2	12.4	7.4	*	*	*	*
2000.....	20.1	22.9	21.4	15.1	11.6	8.1	*	*	*	*
1999.....	22.8	25.3	21.1	18.0	9.1	10.4	6.8	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

Black, female

2002.....	11.9	9.4	7.0	4.8	*	*	*	*	*	*
2001.....	11.4	8.6	6.1	2.6	4.5	*	*	*	*	*
2000.....	8.5	7.7	6.5	4.0	3.8	*	*	*	*	*
1999.....	10.4	7.2	6.3	4.3	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified human immunodeficiency virus (HIV) disease (B24)										
All races 3/, both sexes										
2002.....	2.6	2.4	1.8	1.0	0.7	0.5	0.3	*	*	*
2001.....	2.9	2.3	1.7	0.9	0.7	0.6	0.2	*	*	*
2000.....	2.6	2.4	1.6	1.0	0.7	0.5	0.3	*	*	*
1999.....	2.9	2.3	1.3	1.0	0.6	0.5	0.3	*	*	*
All races 3/, male										
2002.....	3.9	3.7	2.9	1.7	1.2	0.8	0.6	*	*	*
2001.....	4.3	3.8	2.8	1.6	1.2	0.9	0.5	*	*	*
2000.....	4.0	3.9	2.6	1.7	1.4	0.9	0.5	*	*	*
1999.....	4.5	3.7	2.2	1.8	1.0	0.8	0.5	*	*	*
All races 3/, female										
2002.....	1.4	1.2	0.7	0.4	*	*	*	*	*	*
2001.....	1.5	0.8	0.6	0.3	*	*	*	*	*	*
2000.....	1.3	1.0	0.7	*	*	*	*	*	*	*
1999.....	1.4	0.9	0.4	0.3	*	*	*	*	*	*
White, both sexes										
2002.....	1.3	1.3	0.8	0.6	0.4	*	*	*	*	*
2001.....	1.5	1.1	0.9	0.5	0.2	*	*	*	*	*
2000.....	1.3	1.3	0.7	0.6	0.4	*	*	*	*	*
1999.....	1.6	1.1	0.8	0.5	0.3	0.3	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified human immunodeficiency virus (HIV) disease (B24)										
White, male										
2002.....	2.2	2.2	1.4	0.9	0.7	*	*	*	*	*
2001.....	2.6	1.9	1.6	0.9	*	*	*	*	*	*
2000.....	2.2	2.2	1.3	1.0	0.7	*	*	*	*	*
1999.....	2.7	1.9	1.4	0.9	0.6	0.5	*	*	*	*
White, female										
2002.....	0.4	0.4	0.3	*	*	*	*	*	*	*
2001.....	0.4	0.3	*	*	*	*	*	*	*	*
2000.....	0.5	0.4	*	*	*	*	*	*	*	*
1999.....	0.5	0.3	*	*	*	*	*	*	*	*
Black, both sexes										
2002.....	11.8	10.7	9.2	5.3	3.5	3.0	*	*	*	*
2001.....	12.8	11.7	8.6	4.8	4.7	3.7	*	*	*	*
2000.....	11.9	11.7	9.4	5.0	3.8	3.1	*	*	*	*
1999.....	12.2	11.3	6.2	5.9	3.5	2.2	*	*	*	*
Black, male										
2002.....	16.3	15.8	15.8	9.1	6.5	5.2	*	*	*	*
2001.....	17.8	19.7	13.7	8.1	8.7	5.8	*	*	*	*
2000.....	17.8	18.5	15.2	9.4	7.9	5.2	*	*	*	*
1999.....	18.2	18.5	10.7	10.5	5.7	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified human immunodeficiency virus (HIV) disease (B24)										
Black, female										
2002.....	7.8	6.3	3.6	*	*	*	*	*	*	*
2001.....	8.4	4.8	4.1	*	*	*	*	*	*	*
2000.....	6.6	5.7	4.5	*	*	*	*	*	*	*
1999.....	6.9	5.1	2.4	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)										
All races 3/, both sexes										
2002.....	0.2	0.2	0.3	0.3	0.5	0.8	1.0	1.2	1.5	2.9
2001.....	0.1	0.1	0.2	0.3	0.6	0.9	0.9	0.9	1.2	2.2
2000.....	0.1	0.2	0.3	0.4	0.6	0.9	1.1	1.2	1.7	3.3
1999.....	0.2	0.2	0.4	0.6	0.6	0.8	1.1	1.5	2.1	4.0
All races 3/, male										
2002.....	0.2	0.2	0.3	0.3	0.5	0.9	1.0	1.4	1.8	2.8
2001.....	0.2	*	0.2	0.3	0.7	0.9	1.0	1.0	1.5	2.4
2000.....	*	*	0.3	0.4	0.7	1.0	1.4	1.7	2.0	3.1
1999.....	*	0.2	0.4	0.5	0.7	0.8	1.4	1.4	2.0	3.9
All races 3/, female										
2002.....	0.2	0.2	0.2	0.3	0.5	0.8	1.0	1.1	1.3	2.9
2001.....	*	*	0.2	0.3	0.5	0.9	0.8	0.8	1.0	2.1
2000.....	*	0.2	0.3	0.4	0.6	0.8	1.0	0.8	1.6	3.4
1999.....	*	*	0.4	0.6	0.6	0.8	0.9	1.5	2.1	4.0
White, both sexes										
2002.....	0.2	0.2	0.3	0.3	0.5	0.9	1.1	1.2	1.6	3.1
2001.....	0.1	*	0.2	0.3	0.7	1.0	0.9	0.9	1.2	2.3
2000.....	0.1	0.2	0.3	0.5	0.6	1.0	1.2	1.2	1.8	3.4
1999.....	0.2	0.2	0.4	0.6	0.7	0.8	1.1	1.5	2.1	4.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

All races 3/, both sexes

2002.....	0.9	1.4	2.1	2.0	2.5	3.2	4.3	6.2	6.7	8.3
2001.....	0.6	1.1	1.2	1.6	2.3	3.3	4.0	5.1	5.9	8.2
2000.....	0.6	0.9	1.0	1.4	2.1	2.7	4.0	5.4	6.1	9.0
1999.....	1.0	1.4	1.7	2.0	2.8	4.1	5.3	7.0	8.2	10.0

All races 3/, male

2002.....	1.2	2.1	3.0	2.7	3.2	4.1	5.3	7.1	9.0	11.9
2001.....	0.9	1.6	1.6	2.1	2.6	3.6	4.1	6.5	8.0	10.1
2000.....	0.8	1.2	1.3	1.6	2.5	3.3	5.4	6.3	7.7	12.6
1999.....	1.3	1.7	2.1	2.2	3.2	4.8	6.2	8.9	10.3	13.1

All races 3/, female

2002.....	0.5	0.7	1.1	1.4	1.8	2.4	3.5	5.6	5.3	6.8
2001.....	0.4	0.6	0.9	1.1	2.0	3.1	3.9	4.2	4.7	7.4
2000.....	0.4	0.7	0.8	1.2	1.7	2.2	2.9	4.7	5.2	7.6
1999.....	0.8	1.1	1.4	1.8	2.4	3.5	4.6	5.6	6.9	8.8

White, both sexes

2002.....	0.8	1.3	1.9	1.7	2.3	3.1	4.2	6.2	6.6	8.3
2001.....	0.5	1.0	1.1	1.4	2.0	3.1	3.8	5.1	5.8	7.9
2000.....	0.5	0.9	0.9	1.3	1.8	2.5	3.8	5.3	6.0	8.9
1999.....	0.7	1.1	1.4	1.8	2.3	3.7	5.0	6.8	7.8	9.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)										
White, male										
2002.....	1.1	2.0	2.8	2.3	3.0	4.0	5.2	6.9	8.9	11.8
2001.....	0.7	1.5	1.4	1.8	2.4	3.4	4.0	6.3	8.0	9.9
2000.....	0.7	1.1	1.2	1.6	2.1	3.1	5.2	5.9	7.5	12.0
1999.....	1.0	1.4	1.7	2.1	2.7	4.4	5.9	8.7	9.4	12.0
White, female										
2002.....	0.4	0.7	1.0	1.2	1.7	2.3	3.4	5.6	5.2	6.7
2001.....	0.4	0.6	0.9	1.0	1.8	2.9	3.7	4.2	4.4	7.1
2000.....	0.3	0.6	0.7	1.1	1.5	2.1	2.8	4.9	5.0	7.6
1999.....	0.4	0.8	1.1	1.4	2.1	3.2	4.3	5.4	6.9	8.6
Black, both sexes										
2002.....	1.5	2.2	3.8	4.8	3.3	3.5	4.0	7.0	5.5	9.5
2001.....	1.3	2.0	2.0	2.9	3.9	4.4	5.9	4.9	7.0	11.4
2000.....	1.3	1.9	2.2	2.5	3.7	3.9	5.3	5.9	9.0	10.9
1999.....	3.0	3.9	4.4	4.1	6.2	6.9	8.4	9.5	11.0	14.3
Black, male										
2002.....	2.1	3.2	5.4	7.3	4.9	*	*	*	*	*
2001.....	2.1	3.0	2.9	3.6	4.7	5.6	*	*	*	*
2000.....	1.7	2.3	3.0	*	5.0	5.5	7.4	9.9	*	*
1999.....	3.0	4.2	5.1	*	7.6	7.8	9.8	10.9	18.3	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

Black, female

2002.....	*	*	2.5	2.7	*	*	*	6.6	*	*
2001.....	*	*	*	*	3.2	*	5.6	*	*	11.8
2000.....	*	*	*	*	*	*	*	*	*	9.0
1999.....	3.1	3.7	3.9	5.0	5.2	6.2	7.4	8.6	*	11.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mycoses (B35-B49)										
All races 3/, both sexes										
2002.....	0.2	0.3	0.5	0.6	0.8	1.1	1.7	2.3	2.7	2.5
2001.....	0.2	0.3	0.3	0.5	0.8	1.2	1.4	2.3	2.0	3.0
2000.....	0.2	0.3	0.5	0.7	1.0	1.2	1.7	2.3	2.1	3.2
1999.....	0.4	0.6	0.8	1.0	1.4	2.1	2.6	3.2	3.3	3.9
All races 3/, male										
2002.....	0.3	0.4	0.7	0.9	1.1	1.7	2.3	3.1	4.1	4.2
2001.....	0.3	0.5	0.4	0.7	1.0	1.4	1.5	3.2	2.9	3.8
2000.....	0.3	0.3	0.6	0.7	1.2	1.6	2.4	3.0	2.8	4.9
1999.....	0.5	0.6	0.8	1.2	1.7	2.4	3.4	4.4	4.9	6.6
All races 3/, female										
2002.....	*	*	0.3	0.4	0.6	0.7	1.2	1.6	1.9	1.7
2001.....	*	0.2	0.3	0.4	0.7	1.0	1.3	1.7	1.5	2.6
2000.....	0.2	0.3	0.4	0.6	0.8	1.0	1.2	1.8	1.7	2.6
1999.....	0.3	0.5	0.7	0.9	1.2	1.9	1.9	2.4	2.3	2.9
White, both sexes										
2002.....	0.2	0.2	0.3	0.5	0.7	1.0	1.6	2.1	2.7	2.4
2001.....	0.2	0.3	0.3	0.5	0.7	1.0	1.2	2.2	1.8	2.6
2000.....	0.2	0.3	0.4	0.6	0.9	1.2	1.6	2.2	2.1	3.0
1999.....	0.3	0.5	0.7	0.9	1.2	1.9	2.4	3.1	3.0	3.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Protozoal diseases (B50-B64)

All races 3/, both sexes

2002.....	*	*	0.1	0.1	*	*	*	*	*	*
2001.....	0.1	0.1	0.1	0.2	0.3	0.3	0.3	*	*	*
2000.....	0.1	0.1	*	0.1	0.2	*	0.3	*	*	*
1999.....	0.2	0.1	0.2	0.2	*	0.2	0.3	0.3	*	*

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	0.6	*	*	*
1999.....	0.2	*	0.2	*	*	*	*	*	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	0.4	0.3	*	*	*
2000.....	*	*	*	*	*	*	0.3	*	*	*
1999.....	0.1	*	*	*	*	*	0.3	0.3	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumocystosis (B59)										
All races 3/, both sexes										
2002.....	*	*	0.1	*	*	*	*	*	*	*
2001.....	*	0.1	*	0.2	0.2	0.3	0.3	*	*	*
2000.....	*	0.1	*	*	0.2	*	0.3	*	*	*
1999.....	0.1	*	0.2	*	*	*	0.3	0.3	*	*
All races 3/, male										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	0.5	*	*	*
1999.....	*	*	0.2	*	*	*	*	*	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	0.3	0.3	*	*	*
2000.....	*	*	*	*	*	*	0.3	*	*	*
1999.....	0.1	*	*	*	*	*	0.3	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Sequelae of tuberculosis (B90)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	*	*	*	*	0.8
2001.....	*	*	*	*	*	*	*	*	0.5	0.6
2000.....	*	*	*	*	*	*	*	0.3	0.6	0.7
1999.....	*	*	*	*	*	*	0.3	0.4	1.0	0.8
All races 3/, male										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	0.9
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	1.0	0.8
White, both sexes										
2002.....	*	*	*	*	*	*	*	*	*	0.9
2001.....	*	*	*	*	*	*	*	*	0.5	0.6
2000.....	*	*	*	*	*	*	*	*	0.5	0.7
1999.....	*	*	*	*	*	*	*	0.4	0.9	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Sequelae of poliomyelitis (B91)										
All races 3/, both sexes										
2002.....	*	*	*	0.1	0.2	0.4	0.6	0.9	0.9	1.3
2001.....	*	*	*	*	0.3	0.4	0.6	0.8	0.9	1.2
2000.....	*	*	*	0.2	0.2	0.3	0.5	0.8	0.8	1.2
1999.....	*	*	*	*	0.2	0.3	0.5	0.6	0.7	1.3
All races 3/, male										
2002.....	*	*	*	*	*	0.5	0.5	0.7	1.1	1.9
2001.....	*	*	*	*	*	*	0.7	1.2	*	1.8
2000.....	*	*	*	*	*	*	0.6	0.8	*	*
1999.....	*	*	*	*	*	*	*	0.8	*	*
All races 3/, female										
2002.....	*	*	*	*	*	0.4	0.6	1.0	0.8	1.0
2001.....	*	*	*	*	*	0.4	0.6	0.6	0.9	1.0
2000.....	*	*	*	*	*	*	0.5	0.8	0.8	1.1
1999.....	*	*	*	*	*	*	0.5	0.5	0.7	1.2
White, both sexes										
2002.....	*	*	*	0.2	0.2	0.5	0.6	1.0	1.0	1.4
2001.....	*	*	*	*	0.4	0.5	0.7	0.9	1.0	1.4
2000.....	*	*	*	0.2	0.2	0.4	0.6	0.9	0.9	1.4
1999.....	*	*	*	*	0.2	0.3	0.5	0.7	0.8	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)										
All races 3/, both sexes										
2002.....	0.5	1.0	1.4	1.1	1.3	1.4	1.7	2.6	2.6	3.6
2001.....	0.3	0.6	0.7	0.7	0.8	1.3	1.4	1.5	2.3	3.2
2000.....	0.2	0.4	0.4	0.4	0.6	0.8	1.2	1.7	2.3	3.7
1999.....	0.5	0.7	0.7	0.7	1.0	1.4	1.6	2.4	3.0	3.9
All races 3/, male										
2002.....	0.8	1.4	2.1	1.5	1.6	1.7	2.0	2.8	3.2	4.7
2001.....	0.5	0.9	1.0	1.0	0.9	1.3	1.3	1.5	3.2	3.4
2000.....	0.4	0.6	0.5	0.5	0.7	1.0	1.5	2.0	2.9	4.3
1999.....	0.6	0.9	0.9	0.7	1.0	1.7	1.6	2.9	3.7	3.8
All races 3/, female										
2002.....	0.3	0.5	0.6	0.7	0.9	1.2	1.5	2.5	2.2	3.1
2001.....	0.2	0.3	0.4	0.4	0.7	1.2	1.5	1.5	1.8	3.2
2000.....	*	0.3	0.3	0.3	0.5	0.6	0.9	1.5	2.0	3.4
1999.....	0.3	0.5	0.5	0.7	0.9	1.1	1.6	2.1	2.6	3.9
White, both sexes										
2002.....	0.5	1.0	1.4	0.9	1.2	1.4	1.8	2.6	2.6	3.5
2001.....	0.3	0.6	0.6	0.6	0.7	1.2	1.3	1.5	2.3	3.3
2000.....	0.2	0.5	0.4	0.3	0.5	0.7	1.2	1.8	2.2	3.6
1999.....	0.3	0.5	0.5	0.6	0.7	1.2	1.5	2.3	2.9	3.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

Black, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	1.4	2.1	2.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
II. Neoplasms (C00-D48)										
All races 3/, both sexes										
2002.....	48.5	92.1	163.6	280.5	452.5	673.5	949.9	1,235.8	1,504.6	1,802.6
2001.....	49.7	93.9	167.1	284.6	459.6	683.7	961.4	1,241.2	1,513.7	1,848.3
2000.....	49.8	95.0	168.5	292.8	470.0	701.2	969.0	1,251.3	1,555.9	1,904.3
1999.....	50.3	95.1	169.3	298.8	477.9	711.0	982.0	1,253.3	1,540.6	1,888.7
All races 3/, male										
2002.....	43.9	91.4	173.9	313.2	526.0	809.1	1,177.8	1,591.4	2,010.6	2,592.2
2001.....	45.0	93.4	176.6	315.6	534.1	826.9	1,200.6	1,607.6	2,041.1	2,721.3
2000.....	45.3	95.1	176.6	325.4	543.2	848.0	1,211.6	1,622.8	2,115.8	2,819.6
1999.....	46.6	96.1	177.6	334.6	558.0	861.6	1,236.8	1,637.2	2,107.0	2,831.4
All races 3/, female										
2002.....	53.1	92.8	153.8	249.8	385.7	556.4	767.8	984.7	1,196.9	1,460.0
2001.....	54.3	94.4	158.0	255.5	391.9	560.4	771.5	984.7	1,197.7	1,483.0
2000.....	54.1	95.0	160.7	262.4	403.7	575.3	777.8	992.7	1,225.7	1,531.6
1999.....	54.0	94.1	161.4	265.4	405.4	582.2	783.0	986.4	1,209.2	1,508.8
White, both sexes										
2002.....	46.5	87.7	156.7	272.3	445.0	669.1	948.1	1,234.8	1,504.1	1,806.4
2001.....	47.3	89.3	160.7	276.2	453.3	676.3	956.5	1,237.9	1,507.6	1,849.1
2000.....	46.7	89.6	161.8	285.0	463.7	695.8	961.4	1,243.6	1,555.2	1,903.1
1999.....	47.7	89.8	162.1	289.3	468.4	703.7	974.3	1,242.7	1,537.1	1,884.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
II. Neoplasms (C00-D48)										
White, male										
2002.....	42.3	87.0	164.9	300.9	509.8	798.0	1,165.9	1,575.7	1,994.7	2,590.9
2001.....	42.9	88.3	167.6	302.3	521.3	810.7	1,183.4	1,588.2	2,014.4	2,707.4
2000.....	42.3	89.2	166.6	311.3	529.3	833.1	1,187.6	1,596.8	2,094.9	2,806.3
1999.....	44.2	89.4	167.0	320.0	540.8	844.6	1,216.4	1,608.8	2,087.0	2,810.7
White, female										
2002.....	50.6	88.4	148.7	245.0	385.2	555.6	770.6	990.7	1,203.4	1,465.6
2001.....	51.8	90.3	154.0	251.2	390.6	558.3	772.7	989.8	1,201.4	1,489.8
2000.....	51.1	89.9	157.1	260.0	403.2	575.6	779.9	995.3	1,234.1	1,536.2
1999.....	51.2	90.3	157.3	260.1	401.8	580.9	781.9	985.6	1,212.5	1,511.8
Black, both sexes										
2002.....	70.0	137.1	241.3	404.8	602.9	843.8	1,141.1	1,450.2	1,707.0	1,949.3
2001.....	71.6	139.2	244.2	409.2	597.0	869.3	1,175.3	1,457.5	1,759.5	2,005.1
2000.....	75.8	148.0	255.0	412.1	611.8	872.0	1,199.5	1,508.3	1,738.5	2,063.0
1999.....	74.6	146.6	258.7	434.4	641.8	892.8	1,209.9	1,531.6	1,742.9	2,070.3
Black, male										
2002.....	61.7	138.8	274.3	491.8	784.4	1,092.0	1,552.3	2,077.8	2,567.7	3,055.5
2001.....	64.1	145.0	283.1	499.5	755.4	1,148.3	1,623.1	2,115.9	2,703.9	3,298.4
2000.....	71.7	154.0	297.3	517.2	779.4	1,152.9	1,678.6	2,200.5	2,722.0	3,340.9
1999.....	69.5	162.3	304.1	539.8	824.0	1,176.7	1,665.8	2,241.5	2,684.7	3,404.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
II. Neoplasms (C00-D48)										
Black, female										
2002.....	77.3	135.7	213.2	333.3	460.9	659.6	865.2	1,075.3	1,258.9	1,525.5
2001.....	78.3	134.3	211.1	335.1	472.5	663.3	877.1	1,064.5	1,278.4	1,525.3
2000.....	79.5	142.8	218.9	325.9	479.5	664.7	881.3	1,091.1	1,243.1	1,587.4
1999.....	79.1	133.2	220.2	348.4	498.8	682.4	907.4	1,103.9	1,276.1	1,571.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)										
All races 3/, both sexes										
2002.....	47.5	90.6	161.5	276.8	447.0	663.8	933.6	1,207.0	1,458.3	1,723.9
2001.....	48.7	92.4	164.8	280.7	453.3	673.2	943.5	1,211.7	1,465.9	1,765.6
2000.....	48.8	93.6	166.2	289.0	463.4	690.6	951.6	1,222.5	1,505.3	1,819.4
1999.....	49.4	93.7	167.0	295.3	471.6	700.2	963.4	1,224.0	1,493.1	1,805.8
All races 3/, male										
2002.....	42.8	89.8	171.7	308.6	519.7	796.8	1,157.8	1,554.1	1,951.5	2,491.1
2001.....	44.0	91.7	174.2	311.6	526.7	814.1	1,178.0	1,568.4	1,978.7	2,613.6
2000.....	44.4	93.5	173.9	321.3	535.7	834.7	1,190.4	1,586.0	2,050.4	2,710.7
1999.....	45.5	94.7	174.9	330.6	550.1	848.2	1,212.9	1,600.1	2,046.0	2,722.2
All races 3/, female										
2002.....	52.2	91.4	151.8	246.8	381.1	548.9	754.4	961.9	1,158.4	1,391.1
2001.....	53.5	93.2	155.9	251.7	386.7	551.9	757.2	962.0	1,158.6	1,410.7
2000.....	53.2	93.7	158.8	258.9	398.0	567.2	763.5	969.3	1,183.8	1,456.4
1999.....	53.3	92.7	159.5	262.3	400.6	573.7	768.4	962.5	1,169.5	1,436.5
White, both sexes										
2002.....	45.5	86.2	154.7	268.7	439.5	659.2	931.4	1,205.1	1,456.9	1,725.7
2001.....	46.4	87.8	158.4	272.2	447.1	665.6	938.4	1,207.5	1,458.8	1,765.2
2000.....	45.8	88.2	159.5	281.1	457.1	685.0	944.1	1,214.2	1,503.0	1,815.7
1999.....	46.8	88.5	159.9	285.8	462.2	693.1	955.3	1,213.0	1,488.4	1,798.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)										
White, male										
2002.....	41.2	85.4	162.8	296.5	503.4	785.6	1,145.2	1,537.3	1,934.6	2,486.8
2001.....	41.9	86.5	165.1	298.2	513.9	797.6	1,160.4	1,547.8	1,950.7	2,597.6
2000.....	41.4	87.7	164.0	307.0	522.0	819.7	1,166.6	1,559.2	2,026.9	2,693.7
1999.....	43.1	88.0	164.3	316.0	532.7	831.3	1,191.9	1,571.1	2,024.1	2,696.3
White, female										
2002.....	49.9	86.9	146.8	242.0	380.4	548.0	757.1	967.4	1,164.2	1,395.1
2001.....	51.0	89.0	152.0	247.4	385.4	549.7	758.6	966.4	1,161.4	1,416.7
2000.....	50.1	88.7	155.2	256.5	397.2	567.2	765.6	971.5	1,191.3	1,459.1
1999.....	50.5	89.1	155.5	257.1	397.2	572.5	767.3	961.5	1,172.1	1,438.2
Black, both sexes										
2002.....	68.3	135.1	238.2	399.6	597.1	833.2	1,126.1	1,425.8	1,665.1	1,885.7
2001.....	70.4	137.8	241.5	404.7	589.4	858.4	1,156.8	1,433.0	1,716.9	1,931.7
2000.....	74.4	145.6	251.9	407.9	604.6	861.4	1,179.2	1,481.7	1,700.6	1,998.9
1999.....	73.3	144.5	255.6	429.5	633.9	879.1	1,191.7	1,502.1	1,704.2	2,008.4
Black, male										
2002.....	60.4	137.1	271.5	484.4	777.8	1,079.3	1,536.2	2,047.9	2,514.3	2,976.1
2001.....	62.8	143.6	280.7	494.9	746.2	1,135.3	1,600.3	2,084.3	2,645.3	3,209.9
2000.....	70.3	151.8	293.7	512.8	769.2	1,137.3	1,654.9	2,166.0	2,683.2	3,264.8
1999.....	68.3	160.5	300.8	533.7	815.7	1,159.8	1,644.1	2,205.4	2,642.0	3,340.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)										
Black, female										
2002.....	75.2	133.5	210.0	330.0	455.6	650.5	850.9	1,054.1	1,223.1	1,468.0
2001.....	77.1	132.8	208.1	330.6	466.1	653.9	861.4	1,044.2	1,244.0	1,457.5
2000.....	78.1	140.2	216.2	321.9	474.8	657.8	863.3	1,069.2	1,205.8	1,527.7
1999.....	77.7	130.6	217.2	344.4	491.4	671.1	891.6	1,078.5	1,239.4	1,510.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
All races 3/, both sexes										
2002.....	0.8	2.0	3.6	5.8	8.1	8.8	12.1	13.1	14.1	18.7
2001.....	0.8	2.1	3.5	6.1	7.9	9.4	11.6	13.5	15.2	18.5
2000.....	0.9	2.0	3.8	5.7	8.1	9.9	11.5	12.6	14.0	19.4
1999.....	0.9	2.3	3.6	5.8	8.1	9.3	11.3	13.5	15.4	19.3
All races 3/, male										
2002.....	1.2	3.2	5.8	9.4	12.9	13.9	18.1	19.2	20.3	26.9
2001.....	1.2	3.2	5.8	10.0	13.0	14.3	17.9	19.6	22.1	25.8
2000.....	1.4	3.1	6.0	9.5	12.7	15.2	16.9	18.7	20.2	27.1
1999.....	1.3	3.6	5.7	9.0	13.1	14.3	17.4	20.1	22.3	27.1
All races 3/, female										
2002.....	0.4	0.9	1.4	2.5	3.7	4.4	7.3	8.8	10.3	15.1
2001.....	0.4	1.0	1.4	2.4	3.3	5.1	6.6	9.3	11.1	15.4
2000.....	0.4	1.0	1.6	2.2	3.9	5.4	7.2	8.3	10.4	16.3
1999.....	0.4	1.0	1.6	2.7	3.6	5.1	6.6	9.0	11.3	16.1
White, both sexes										
2002.....	0.7	1.7	3.1	5.3	7.5	8.4	12.3	13.3	14.2	18.9
2001.....	0.7	1.7	3.2	5.5	7.3	8.7	11.4	13.6	15.5	19.0
2000.....	0.7	1.7	3.2	5.0	7.3	9.4	11.1	12.5	14.0	19.7
1999.....	0.7	1.8	3.0	5.0	7.5	8.9	11.0	13.2	15.5	19.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
White, male										
2002.....	1.0	2.7	5.0	8.6	11.7	12.9	18.0	19.4	20.2	27.0
2001.....	1.0	2.6	5.2	8.9	11.8	13.2	17.1	19.2	22.3	26.4
2000.....	1.1	2.7	4.9	8.3	11.2	14.0	15.9	18.0	19.9	27.6
1999.....	1.1	2.7	4.7	7.6	12.0	13.5	16.8	19.3	22.0	27.3
White, female										
2002.....	0.4	0.7	1.3	2.2	3.6	4.3	7.6	8.8	10.5	15.3
2001.....	0.3	0.8	1.2	2.2	3.2	4.8	6.7	9.6	11.3	15.9
2000.....	0.4	0.8	1.6	1.9	3.6	5.3	7.2	8.6	10.6	16.5
1999.....	0.4	0.8	1.3	2.5	3.3	4.9	6.4	9.0	11.7	16.7
Black, both sexes										
2002.....	1.5	4.2	7.1	10.1	14.7	13.3	12.5	11.3	12.6	16.0
2001.....	1.7	4.3	6.2	12.2	13.1	15.3	15.7	14.2	13.3	12.9
2000.....	1.6	4.1	8.3	12.2	15.2	14.9	16.1	14.6	15.8	15.3
1999.....	1.5	5.9	8.8	12.6	14.6	14.4	16.2	17.7	15.6	14.3
Black, male										
2002.....	2.4	6.7	12.9	18.2	26.9	24.2	21.4	16.5	21.4	26.8
2001.....	2.4	7.0	10.8	22.2	24.8	25.4	29.3	25.1	20.9	*
2000.....	2.7	6.3	15.9	21.9	27.3	27.6	29.1	28.8	28.7	*
1999.....	2.3	10.3	14.5	21.5	25.2	24.1	26.5	33.2	28.8	24.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Black, female

2002.....	*	1.9	2.2	3.4	5.1	5.2	6.5	8.2	8.0	11.9
2001.....	*	2.0	2.2	4.0	3.8	7.9	6.7	7.8	9.4	10.5
2000.....	*	2.1	*	4.2	5.6	5.6	7.5	6.0	9.3	14.2
1999.....	*	2.1	3.9	5.3	6.3	7.1	9.5	8.3	9.1	10.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of tongue (C01-C02)										
All races 3/, both sexes										
2002.....	0.2	0.5	0.9	1.4	2.1	2.1	2.8	3.2	3.4	4.0
2001.....	0.2	0.5	0.9	1.6	1.8	2.0	2.6	3.1	3.5	3.8
2000.....	0.3	0.5	0.9	1.4	2.1	2.3	2.4	2.8	3.2	4.0
1999.....	0.3	0.6	0.9	1.3	1.8	2.2	2.6	3.0	3.4	4.0
All races 3/, male										
2002.....	0.3	0.7	1.4	2.3	3.1	3.5	3.8	4.5	4.3	5.2
2001.....	0.3	0.8	1.5	2.6	3.0	3.1	4.0	4.1	4.9	4.1
2000.....	0.3	0.8	1.5	2.2	3.4	3.5	3.3	3.6	3.8	5.1
1999.....	0.4	0.9	1.4	1.9	3.0	3.5	4.1	4.7	4.3	5.4
All races 3/, female										
2002.....	*	0.2	0.4	0.6	1.2	0.9	2.0	2.3	2.8	3.5
2001.....	*	0.3	0.4	0.6	0.7	1.1	1.4	2.4	2.7	3.6
2000.....	0.2	0.3	0.4	0.6	1.0	1.2	1.7	2.3	2.8	3.6
1999.....	0.2	0.3	0.4	0.7	0.8	1.1	1.4	1.8	2.8	3.5
White, both sexes										
2002.....	0.2	0.4	0.9	1.4	2.0	2.0	2.9	3.3	3.5	4.1
2001.....	0.2	0.5	0.9	1.6	1.8	1.9	2.6	3.2	3.5	3.9
2000.....	0.2	0.5	0.8	1.3	2.0	2.3	2.4	2.9	3.3	4.1
1999.....	0.3	0.5	0.8	1.1	1.8	2.2	2.6	3.0	3.5	4.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of tongue (C01-C02)										
White, male										
2002.....	0.3	0.6	1.4	2.2	3.0	3.2	3.9	4.6	4.4	5.2
2001.....	0.3	0.7	1.5	2.5	2.9	3.0	4.1	4.2	4.8	4.3
2000.....	0.3	0.7	1.3	2.2	3.3	3.5	3.3	3.5	3.8	5.1
1999.....	0.4	0.7	1.3	1.6	2.9	3.5	4.0	4.7	4.5	5.5
White, female										
2002.....	*	*	0.4	0.6	1.2	0.9	2.1	2.4	3.0	3.6
2001.....	*	0.3	0.4	0.7	0.7	1.0	1.4	2.5	2.8	3.8
2000.....	*	0.3	0.5	0.5	0.9	1.3	1.7	2.4	3.0	3.7
1999.....	*	0.3	0.4	0.7	0.8	1.0	1.4	1.9	3.0	3.5
Black, both sexes										
2002.....	*	1.0	1.2	1.9	3.4	3.8	*	*	*	*
2001.....	*	1.0	1.2	2.0	2.9	2.6	*	*	*	*
2000.....	*	*	2.0	2.1	3.1	2.5	3.1	*	*	*
1999.....	*	1.3	2.0	2.8	2.5	2.8	3.0	*	*	*
Black, male										
2002.....	*	*	*	3.1	6.1	7.2	*	*	*	*
2001.....	*	1.7	2.3	4.1	5.5	*	*	*	*	*
2000.....	*	*	4.0	3.6	5.4	5.2	*	*	*	*
1999.....	*	2.3	3.2	5.1	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pharynx (C10-C13,C14.0)										
All races 3/, both sexes										
2002.....	0.3	0.8	1.6	2.5	3.5	3.8	5.2	5.2	4.9	5.4
2001.....	0.4	0.9	1.5	2.6	3.7	4.1	4.9	5.5	5.7	5.9
2000.....	0.4	0.8	1.6	2.7	3.7	4.3	5.1	5.2	5.3	5.6
1999.....	0.4	1.0	1.6	2.7	3.6	4.1	5.1	5.5	5.6	5.2
All races 3/, male										
2002.....	0.5	1.4	2.6	4.0	5.9	6.0	8.2	8.2	8.2	9.8
2001.....	0.6	1.5	2.6	4.4	6.1	6.5	7.9	8.8	9.0	10.0
2000.....	0.6	1.3	2.6	4.6	5.9	6.8	8.2	8.6	8.6	9.9
1999.....	0.6	1.6	2.6	4.5	5.9	6.5	8.0	8.7	9.6	9.9
All races 3/, female										
2002.....	*	0.3	0.6	1.2	1.4	1.8	2.9	3.1	2.8	3.5
2001.....	*	0.4	0.6	0.9	1.6	2.1	2.5	3.1	3.6	4.2
2000.....	*	0.3	0.6	1.0	1.6	2.2	2.8	2.9	3.3	3.9
1999.....	*	0.4	0.7	1.1	1.6	2.1	2.8	3.3	3.2	3.4
White, both sexes										
2002.....	0.2	0.6	1.2	2.2	3.0	3.5	5.2	5.0	4.6	5.2
2001.....	0.2	0.7	1.2	2.2	3.3	3.7	4.6	5.2	5.5	6.0
2000.....	0.2	0.6	1.3	2.2	3.1	3.9	4.7	5.0	5.2	5.3
1999.....	0.2	0.6	1.2	2.3	3.2	3.8	4.7	5.1	5.3	5.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pharynx (C10-C13,C14.0)										
White, male										
2002.....	0.3	1.0	2.0	3.4	4.9	5.4	8.1	8.0	7.6	9.5
2001.....	0.4	1.1	2.0	3.6	5.3	5.6	7.4	8.3	8.8	9.9
2000.....	0.3	0.9	2.0	3.7	4.8	6.1	7.3	7.8	8.4	9.5
1999.....	0.3	1.0	1.9	3.7	5.1	5.8	7.4	7.9	8.9	9.9
White, female										
2002.....	*	0.2	0.5	1.0	1.3	1.8	2.8	2.9	2.7	3.4
2001.....	*	*	0.4	0.8	1.5	2.1	2.4	3.1	3.5	4.3
2000.....	*	0.2	0.6	0.8	1.5	1.9	2.7	2.9	3.3	3.7
1999.....	*	0.3	0.5	0.9	1.4	1.9	2.6	3.2	3.1	3.5
Black, both sexes										
2002.....	0.9	2.0	3.8	5.1	8.1	5.7	6.4	5.3	7.6	7.4
2001.....	0.9	2.2	3.7	6.5	6.4	7.4	7.8	7.3	7.6	*
2000.....	0.8	2.1	3.8	7.1	8.7	8.2	9.1	8.4	7.3	7.5
1999.....	0.8	3.0	4.6	6.5	7.5	7.5	8.9	10.2	9.0	*
Black, male										
2002.....	*	3.4	7.2	9.2	15.2	10.7	10.3	*	*	*
2001.....	*	3.4	6.5	11.7	11.6	14.5	14.5	13.5	*	*
2000.....	*	3.5	7.1	12.7	15.9	14.1	17.7	17.0	*	*
1999.....	*	5.4	8.1	11.2	12.9	13.0	14.6	19.9	17.4	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of pharynx (C10-C13,C14.0)

Black, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	3.8	*	*	*	*
1999.....	*	*	*	*	*	*	5.2	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)

All races 3/, both sexes

2002.....	0.2	0.7	1.1	1.9	2.4	2.9	4.0	4.5	5.7	8.7
2001.....	0.2	0.6	1.0	1.8	2.3	3.1	4.1	4.8	5.9	8.3
2000.....	0.3	0.7	1.2	1.6	2.2	3.3	3.8	4.4	5.2	9.5
1999.....	0.2	0.7	1.1	1.7	2.6	3.0	3.6	4.9	6.3	9.7

All races 3/, male

2002.....	0.3	1.1	1.8	3.1	3.8	4.4	5.9	6.2	7.5	11.1
2001.....	0.3	0.9	1.7	2.9	3.7	4.6	5.8	6.4	8.0	10.6
2000.....	0.4	1.1	1.9	2.6	3.3	4.8	5.2	6.3	7.4	11.4
1999.....	0.3	1.1	1.7	2.5	4.2	4.3	5.2	6.5	8.2	11.4

All races 3/, female

2002.....	*	0.3	0.4	0.7	1.1	1.6	2.4	3.4	4.6	7.7
2001.....	*	0.3	0.4	0.8	1.0	1.9	2.7	3.7	4.7	7.3
2000.....	*	0.4	0.5	0.6	1.3	1.9	2.8	3.1	4.0	8.8
1999.....	*	0.4	0.5	0.9	1.2	1.8	2.3	3.7	5.2	9.0

White, both sexes

2002.....	0.2	0.7	1.0	1.8	2.4	2.9	4.1	4.7	5.9	9.1
2001.....	0.2	0.5	1.0	1.7	2.2	3.0	4.0	5.0	6.3	8.5
2000.....	0.3	0.6	1.1	1.4	2.1	3.2	3.9	4.6	5.2	9.9
1999.....	0.2	0.6	1.0	1.6	2.4	2.9	3.6	4.9	6.5	10.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)

White, male

2002.....	0.3	1.0	1.7	2.9	3.8	4.4	5.9	6.4	7.8	11.5
2001.....	0.3	0.8	1.6	2.6	3.5	4.5	5.5	6.5	8.4	11.0
2000.....	0.4	1.0	1.6	2.3	3.1	4.4	5.2	6.4	7.2	12.2
1999.....	0.3	0.9	1.6	2.3	3.9	4.0	5.2	6.5	8.3	11.5

White, female

2002.....	*	0.3	0.4	0.7	1.1	1.6	2.6	3.5	4.7	8.0
2001.....	*	0.3	0.4	0.8	1.0	1.7	2.8	3.9	5.0	7.5
2000.....	*	0.3	0.5	0.6	1.3	2.1	2.8	3.2	4.1	8.9
1999.....	*	0.3	0.4	0.9	1.1	1.9	2.3	3.8	5.5	9.5

Black, both sexes

2002.....	*	1.2	2.0	3.0	3.1	3.6	3.9	4.1	*	6.2
2001.....	*	1.2	1.2	3.7	3.7	5.3	5.5	4.5	*	*
2000.....	*	1.4	2.6	3.0	3.4	4.2	4.0	3.7	6.2	*
1999.....	*	1.6	2.2	3.3	4.7	4.1	4.3	4.3	*	6.4

Black, male

2002.....	*	1.9	3.7	5.8	5.5	6.2	7.3	*	*	*
2001.....	*	1.8	*	6.4	7.5	7.1	10.1	*	*	*
2000.....	*	*	4.9	5.6	6.0	8.3	6.7	*	*	*
1999.....	*	2.6	3.2	5.3	8.3	7.3	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of digestive organs (C15-C26)										
All races 3/, both sexes										
2002.....	10.1	21.1	39.5	64.8	101.0	152.4	212.5	281.9	354.0	467.4
2001.....	9.8	21.6	39.7	63.4	102.1	154.5	215.5	282.2	362.2	487.9
2000.....	10.0	21.7	38.8	65.0	105.6	153.7	217.5	290.1	374.6	504.9
1999.....	10.2	21.8	38.1	65.5	106.6	157.7	218.4	288.9	367.2	507.4
All races 3/, male										
2002.....	12.7	27.5	53.5	86.9	135.8	202.1	278.3	364.4	449.0	557.9
2001.....	11.6	28.1	52.7	84.4	136.1	203.6	278.0	366.3	458.9	593.2
2000.....	12.3	28.4	51.4	86.2	139.3	202.8	281.4	377.0	474.6	614.1
1999.....	12.5	29.0	50.4	86.9	138.8	205.8	285.0	374.7	472.3	620.8
All races 3/, female										
2002.....	7.5	14.8	26.1	44.1	69.4	109.5	159.9	223.5	296.1	428.1
2001.....	7.9	15.3	27.2	43.8	71.3	112.3	165.8	223.4	304.3	443.9
2000.....	7.7	15.2	26.6	45.1	75.1	111.6	167.2	229.6	315.5	460.4
1999.....	8.0	14.9	26.2	45.5	77.5	116.7	166.3	229.3	305.7	461.7
White, both sexes										
2002.....	9.1	19.2	36.3	60.5	95.8	145.8	205.6	274.7	346.3	460.4
2001.....	8.8	19.6	36.6	58.8	97.3	147.3	208.2	274.5	354.1	481.6
2000.....	8.8	19.6	35.6	60.3	100.2	147.0	210.3	281.8	365.9	497.0
1999.....	9.1	19.9	34.8	60.2	100.9	151.7	211.0	280.4	358.9	499.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of digestive organs (C15-C26)										
White, male										
2002.....	11.6	25.3	49.1	81.2	129.0	194.6	269.7	356.9	439.1	550.5
2001.....	10.5	25.5	48.6	78.4	130.0	194.4	269.8	358.3	449.9	588.1
2000.....	10.9	25.6	47.1	79.3	132.3	195.1	271.9	368.1	464.1	605.3
1999.....	11.3	26.3	45.8	79.5	131.0	198.2	277.1	363.3	463.6	614.4
White, female										
2002.....	6.7	13.1	23.9	40.6	65.2	102.9	153.3	215.9	289.5	421.3
2001.....	7.2	13.7	24.8	40.1	67.2	105.9	158.4	215.0	296.1	437.0
2000.....	6.8	13.6	24.4	42.3	70.6	105.0	160.8	221.1	307.4	453.1
1999.....	7.0	13.6	24.1	41.8	73.1	111.2	158.5	222.3	297.0	453.7
Black, both sexes										
2002.....	16.2	34.7	64.6	107.2	151.8	217.4	295.9	380.0	450.0	566.6
2001.....	15.7	36.7	65.8	107.4	150.7	224.7	297.8	380.2	475.1	585.6
2000.....	16.8	37.8	68.1	105.8	158.3	221.2	302.3	395.9	483.7	606.9
1999.....	17.0	36.1	66.2	112.9	160.5	222.0	306.6	395.8	484.5	613.1
Black, male										
2002.....	20.1	44.6	90.5	147.5	206.9	288.2	394.9	489.1	600.7	671.1
2001.....	19.1	48.4	89.0	146.3	199.9	305.9	385.4	489.7	602.7	676.3
2000.....	21.6	50.2	93.2	152.0	212.2	289.6	403.5	508.4	629.7	728.8
1999.....	21.1	50.3	92.5	159.0	216.8	292.9	397.5	543.8	615.6	715.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of digestive organs (C15-C26)										
Black, female										
2002.....	12.7	26.1	42.7	74.1	108.7	164.8	229.4	314.8	371.6	526.6
2001.....	12.7	26.5	46.1	75.6	112.0	164.6	239.4	314.8	410.0	551.9
2000.....	12.5	27.0	46.8	67.9	115.7	170.7	235.1	328.1	410.1	561.5
1999.....	13.4	23.9	44.0	75.2	116.3	169.5	246.2	306.7	419.6	574.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of esophagus (C15)										
All races 3/, both sexes										
2002.....	1.0	2.4	4.7	8.6	12.7	18.1	22.5	25.7	27.8	24.6
2001.....	1.1	2.4	5.0	7.9	13.0	18.5	22.9	26.1	26.5	25.8
2000.....	1.0	2.4	4.9	8.8	13.2	17.9	22.1	24.5	27.2	25.9
1999.....	1.0	2.4	4.7	8.7	13.0	18.2	22.0	24.3	24.6	26.8
All races 3/, male										
2002.....	1.6	4.1	8.3	15.1	22.1	31.4	40.3	46.8	50.6	43.4
2001.....	1.8	4.1	8.7	13.7	22.6	32.6	39.9	46.5	48.8	48.1
2000.....	1.8	4.2	8.7	15.2	23.0	30.7	38.0	43.9	48.8	47.8
1999.....	1.7	4.2	8.3	15.3	21.9	31.7	39.3	44.1	44.8	48.0
All races 3/, female										
2002.....	0.4	0.7	1.3	2.4	4.2	6.7	8.2	10.9	14.0	16.5
2001.....	0.3	0.7	1.4	2.4	4.2	6.4	9.4	11.9	13.2	16.5
2000.....	0.2	0.6	1.4	2.8	4.4	7.1	9.6	11.0	14.5	17.1
1999.....	0.3	0.6	1.3	2.5	5.0	6.7	8.5	10.5	12.8	18.3
White, both sexes										
2002.....	0.9	2.3	4.6	8.1	12.2	17.5	22.4	25.7	28.2	25.1
2001.....	1.0	2.3	4.6	7.1	12.3	17.7	22.1	26.2	26.9	26.3
2000.....	0.9	2.3	4.5	8.1	12.4	17.4	21.8	24.0	27.2	26.6
1999.....	0.9	2.2	4.3	8.0	12.1	17.5	21.5	23.7	24.9	27.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of esophagus (C15)										
White, male										
2002.....	1.6	4.0	8.1	14.5	21.5	30.7	40.2	47.2	51.3	44.6
2001.....	1.8	3.9	8.1	12.5	21.6	31.2	39.0	46.9	49.5	48.8
2000.....	1.8	4.1	8.0	14.1	22.0	30.0	37.5	43.2	49.7	49.3
1999.....	1.6	4.0	7.6	14.1	20.5	30.8	38.4	43.0	45.6	48.0
White, female										
2002.....	0.3	0.6	1.2	2.1	3.7	5.9	7.8	10.3	14.1	16.7
2001.....	0.3	0.6	1.2	1.9	3.7	5.8	8.5	11.6	13.2	16.8
2000.....	*	0.4	1.1	2.4	3.6	6.4	9.1	10.6	13.8	17.3
1999.....	*	0.5	1.0	2.1	4.4	5.8	8.0	10.1	12.6	18.9
Black, both sexes										
2002.....	1.6	3.6	6.9	14.9	20.4	27.5	29.8	31.8	26.2	20.8
2001.....	1.6	3.7	8.6	15.9	21.9	29.6	35.8	30.7	27.6	23.6
2000.....	1.5	4.1	9.8	16.9	23.3	27.0	29.6	36.1	29.9	22.2
1999.....	1.7	4.3	9.4	17.1	23.5	28.7	31.3	35.7	23.7	23.9
Black, male										
2002.....	2.5	5.7	11.5	26.8	34.4	45.4	54.4	54.2	48.1	37.5
2001.....	2.6	6.0	15.3	27.5	38.8	51.8	59.9	55.3	53.8	46.5
2000.....	2.4	6.3	17.4	30.5	38.5	44.8	49.8	65.7	45.5	38.1
1999.....	3.2	7.4	16.4	30.3	40.3	46.5	56.0	67.9	36.6	51.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of esophagus (C15)										
Black, female										
2002.....	*	1.8	3.0	5.1	9.4	14.2	13.2	18.4	14.7	14.4
2001.....	*	1.7	2.9	6.3	8.6	13.1	19.7	16.1	14.3	15.2
2000.....	*	2.1	3.4	5.8	11.3	13.8	16.2	18.2	22.1	16.3
1999.....	*	1.6	3.4	6.4	10.3	15.6	14.9	16.3	17.3	13.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of stomach (C16)										
All races 3/, both sexes										
2002.....	1.2	2.1	3.5	5.6	8.5	12.3	18.8	25.6	33.0	47.6
2001.....	1.3	2.2	3.7	5.3	9.2	13.7	19.5	27.0	34.7	45.4
2000.....	1.4	2.3	3.6	5.6	9.7	13.8	19.6	27.1	36.9	52.7
1999.....	1.4	2.3	3.8	6.2	9.8	14.2	20.2	28.1	36.8	50.1
All races 3/, male										
2002.....	1.4	2.7	4.8	7.7	11.9	18.1	27.5	36.4	47.1	67.2
2001.....	1.4	2.8	4.9	7.5	13.7	19.5	27.5	39.1	49.4	62.6
2000.....	1.7	2.9	4.8	7.8	14.1	19.5	28.3	39.4	51.4	75.6
1999.....	1.7	3.1	5.4	8.8	13.7	20.3	28.6	41.5	54.9	71.3
All races 3/, female										
2002.....	1.0	1.5	2.3	3.7	5.4	7.3	11.9	18.1	24.4	39.1
2001.....	1.1	1.6	2.5	3.2	5.1	8.8	13.1	18.5	25.9	38.3
2000.....	1.2	1.7	2.4	3.5	5.8	8.9	12.7	18.5	28.3	43.4
1999.....	1.2	1.5	2.4	3.7	6.4	9.1	13.7	18.8	26.2	41.6
White, both sexes										
2002.....	1.0	1.8	2.9	4.9	7.3	10.5	16.5	22.6	29.3	42.2
2001.....	1.1	1.9	3.1	4.7	7.9	11.7	17.4	24.0	31.0	41.1
2000.....	1.1	1.9	3.1	4.9	8.5	11.9	17.2	23.6	32.7	47.3
1999.....	1.1	2.0	3.1	5.2	8.3	12.8	17.7	24.8	33.0	44.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of stomach (C16)										
White, male										
2002.....	1.1	2.4	3.9	6.8	10.3	15.7	24.3	32.8	42.4	60.5
2001.....	1.2	2.4	4.1	6.5	11.7	16.3	24.5	35.4	44.0	57.7
2000.....	1.2	2.3	4.2	6.7	12.5	17.1	25.1	34.6	46.1	68.9
1999.....	1.2	2.7	4.4	7.6	11.5	18.4	25.2	36.9	49.0	63.4
White, female										
2002.....	0.8	1.2	1.8	3.0	4.6	5.9	10.1	15.4	21.3	34.2
2001.....	1.0	1.3	2.1	2.9	4.3	7.7	11.7	16.0	23.1	34.1
2000.....	1.0	1.5	2.0	3.1	4.9	7.4	10.9	15.8	24.7	38.6
1999.....	0.9	1.2	1.9	2.9	5.4	8.0	11.7	16.3	23.5	37.3
Black, both sexes										
2002.....	2.2	3.6	7.5	10.4	16.1	24.6	38.9	52.7	63.1	100.4
2001.....	2.5	3.7	7.5	9.2	17.7	27.1	34.7	51.1	69.9	88.1
2000.....	2.9	4.7	7.0	11.1	17.3	26.8	38.2	56.9	75.1	101.7
1999.....	3.2	3.9	8.5	13.2	19.1	23.6	40.3	58.6	74.3	102.4
Black, male										
2002.....	2.4	5.1	11.0	15.2	24.0	36.9	59.8	73.5	90.2	137.2
2001.....	2.8	5.0	11.5	13.8	27.5	43.7	54.2	75.7	106.7	104.4
2000.....	3.9	6.6	9.6	16.5	25.9	38.5	57.2	83.2	105.4	136.1
1999.....	4.0	5.7	12.2	19.0	28.2	35.1	60.4	89.3	121.2	147.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of stomach (C16)										
Black, female										
2002.....	2.1	2.3	4.6	6.5	9.9	15.5	24.9	40.3	48.9	86.3
2001.....	2.2	2.5	4.1	5.4	10.1	14.8	21.7	36.3	51.1	82.1
2000.....	1.9	3.0	4.7	6.6	10.5	18.2	25.5	41.0	59.9	88.9
1999.....	2.4	2.4	5.4	8.4	12.0	15.2	27.0	40.0	51.0	85.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of small intestine (C17)										
All races 3/, both sexes										
2002.....	0.1	0.2	0.4	0.6	0.7	1.2	1.6	2.1	2.6	2.8
2001.....	0.1	0.2	0.4	0.5	1.0	1.0	1.9	2.4	2.6	3.2
2000.....	0.1	0.3	0.4	0.6	1.0	1.2	1.7	2.2	2.8	3.0
1999.....	0.1	0.4	0.5	0.4	0.9	1.2	1.7	2.2	2.6	3.2
All races 3/, male										
2002.....	0.2	0.3	0.4	0.6	1.0	1.4	1.9	2.7	2.9	3.7
2001.....	*	0.3	0.5	0.6	1.3	1.3	2.5	2.7	3.3	4.1
2000.....	*	0.3	0.4	0.6	1.2	1.4	2.2	2.5	3.7	4.2
1999.....	0.2	0.4	0.5	0.6	1.4	1.5	2.0	3.0	3.6	4.4
All races 3/, female										
2002.....	*	*	0.3	0.5	0.5	1.1	1.3	1.7	2.5	2.4
2001.....	*	0.2	0.2	0.5	0.8	0.8	1.4	2.2	2.1	2.8
2000.....	*	0.2	0.3	0.5	0.8	1.1	1.4	1.9	2.3	2.5
1999.....	*	0.3	0.4	0.3	0.4	0.9	1.4	1.7	2.0	2.7
White, both sexes										
2002.....	0.1	0.2	0.4	0.5	0.6	1.2	1.6	2.0	2.5	2.6
2001.....	0.1	0.2	0.3	0.5	1.0	1.0	1.8	2.4	2.5	3.0
2000.....	0.1	0.2	0.3	0.6	1.0	1.2	1.7	2.1	2.7	3.1
1999.....	0.1	0.3	0.4	0.4	0.8	1.2	1.6	2.1	2.4	3.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of colon, rectum and anus (C18-C21)										
All races 3/, both sexes										
2002.....	4.3	8.2	15.0	24.6	40.7	61.6	86.2	118.9	163.2	241.4
2001.....	4.1	8.4	15.2	25.4	39.3	62.9	88.8	121.1	165.8	255.7
2000.....	4.0	8.2	14.7	25.2	42.0	62.0	92.4	126.9	174.8	270.4
1999.....	4.0	8.3	14.4	25.3	43.6	63.9	91.9	129.0	173.2	271.7
All races 3/, male										
2002.....	4.9	9.1	17.5	30.1	51.6	77.8	106.8	147.7	197.4	275.1
2001.....	4.3	9.5	17.1	30.8	48.0	78.9	110.4	149.4	201.3	298.3
2000.....	4.4	9.0	17.0	30.6	51.0	77.4	115.7	159.4	213.0	315.6
1999.....	4.1	9.1	16.6	30.6	52.9	78.0	114.3	161.3	216.2	324.0
All races 3/, female										
2002.....	3.6	7.4	12.6	19.5	30.8	47.6	69.7	98.6	142.3	226.8
2001.....	3.8	7.4	13.4	20.4	31.5	49.2	71.5	101.2	144.6	237.8
2000.....	3.5	7.5	12.4	20.2	33.9	48.8	74.1	104.3	152.2	251.9
1999.....	3.9	7.5	12.3	20.4	35.1	51.8	74.3	106.6	148.0	250.6
White, both sexes										
2002.....	3.9	7.5	14.2	23.3	38.9	59.9	84.6	116.9	160.3	240.7
2001.....	3.8	7.7	14.3	23.9	37.9	60.8	86.8	118.0	163.7	255.3
2000.....	3.6	7.5	13.8	23.7	40.5	60.3	90.6	125.0	172.1	269.2
1999.....	3.7	7.7	13.5	23.8	42.0	62.0	90.0	126.9	170.8	270.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of colon, rectum and anus (C18-C21)										
White, male										
2002.....	4.5	8.4	16.5	28.7	49.4	76.0	105.4	144.9	193.3	273.6
2001.....	4.0	8.7	16.4	29.3	46.7	76.7	108.7	146.7	198.9	297.3
2000.....	3.9	8.3	16.1	28.6	49.1	75.3	113.5	157.6	208.6	313.3
1999.....	3.9	8.5	15.8	28.8	51.1	76.1	113.3	158.1	214.4	324.5
White, female										
2002.....	3.3	6.5	12.0	18.2	29.3	45.7	67.6	96.9	140.1	226.4
2001.....	3.6	6.7	12.2	18.8	29.8	46.9	69.1	97.7	142.5	237.7
2000.....	3.3	6.6	11.5	19.1	32.6	47.2	72.3	102.1	150.4	251.3
1999.....	3.5	6.9	11.2	19.2	33.6	49.7	71.5	104.9	145.1	249.3
Black, both sexes										
2002.....	7.2	14.5	23.5	40.0	61.9	86.8	114.9	160.5	218.7	274.7
2001.....	6.5	14.5	24.6	43.1	58.3	90.9	120.3	172.6	211.0	286.5
2000.....	6.6	15.2	24.8	39.9	62.7	86.9	126.1	167.4	226.2	310.0
1999.....	6.6	13.3	23.4	42.0	65.7	90.5	125.4	172.3	226.8	306.1
Black, male										
2002.....	8.4	15.7	27.9	47.5	79.4	109.2	138.2	209.0	285.1	328.1
2001.....	7.6	16.4	26.4	51.3	67.7	114.1	146.8	208.2	263.2	347.2
2000.....	8.0	16.4	29.2	51.4	76.4	109.3	160.1	208.9	298.4	377.1
1999.....	6.3	15.0	26.9	54.1	80.4	110.6	147.6	228.0	274.7	350.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of colon, rectum and anus (C18-C21)										
Black, female										
2002.....	6.1	13.5	19.7	33.9	48.2	70.1	99.2	131.5	184.2	254.2
2001.....	5.6	12.8	23.1	36.3	50.9	73.8	102.7	151.4	184.4	264.0
2000.....	5.3	14.2	21.1	30.5	52.0	70.5	103.4	142.4	189.8	285.0
1999.....	6.7	11.9	20.5	32.1	54.2	75.6	110.7	138.7	203.1	289.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Colon (C18)										
All races 3/, both sexes										
2002.....	3.3	6.3	11.9	19.8	32.5	50.5	72.7	101.5	142.3	211.3
2001.....	3.1	6.4	11.8	20.4	32.4	52.4	75.6	103.9	143.5	221.6
2000.....	3.1	6.3	11.8	20.1	34.4	51.4	78.2	108.3	153.7	236.6
1999.....	3.1	6.4	11.6	20.4	35.9	53.3	77.7	110.7	151.6	238.1
All races 3/, male										
2002.....	3.7	6.8	13.5	23.8	40.2	63.2	89.8	124.7	169.8	240.2
2001.....	3.2	7.1	12.9	24.5	38.5	64.5	92.6	126.3	173.0	256.7
2000.....	3.4	6.7	13.6	24.0	40.9	62.7	96.6	135.2	185.0	275.8
1999.....	3.2	7.0	13.2	24.2	43.3	64.3	96.0	137.4	187.2	282.2
All races 3/, female										
2002.....	2.9	5.8	10.3	16.1	25.4	39.5	59.1	85.1	125.6	198.7
2001.....	3.0	5.7	10.7	16.7	26.9	42.0	62.1	88.3	125.8	206.9
2000.....	2.8	5.9	10.1	16.6	28.5	41.7	63.7	89.6	135.2	220.6
1999.....	3.0	5.9	10.1	16.8	29.3	43.8	63.4	92.1	130.8	220.4
White, both sexes										
2002.....	3.0	5.5	11.1	18.6	30.8	48.6	71.0	99.4	140.0	210.5
2001.....	2.8	5.8	10.9	19.1	30.9	50.1	73.7	100.7	141.6	221.2
2000.....	2.8	5.6	11.0	18.8	33.0	49.9	76.4	106.4	150.8	235.0
1999.....	2.8	5.9	10.8	19.0	34.5	51.6	75.8	108.5	149.3	237.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Colon (C18)										
White, male										
2002.....	3.4	6.1	12.6	22.5	38.1	61.2	88.2	121.7	166.9	238.7
2001.....	2.9	6.4	12.1	23.2	37.1	62.2	91.1	123.1	170.6	255.5
2000.....	3.0	6.1	12.7	22.3	39.0	61.2	94.6	133.3	180.4	272.5
1999.....	3.0	6.5	12.6	22.6	41.6	62.7	95.0	134.3	185.6	282.9
White, female										
2002.....	2.6	4.9	9.7	14.8	24.0	37.6	57.0	83.4	123.5	198.2
2001.....	2.7	5.1	9.6	15.1	25.1	39.5	59.7	84.8	124.1	206.9
2000.....	2.6	5.1	9.3	15.5	27.3	40.1	61.8	87.5	133.2	219.8
1999.....	2.6	5.3	9.2	15.5	27.9	41.8	60.6	90.5	127.8	218.7
Black, both sexes										
2002.....	5.7	12.1	19.7	33.6	52.1	74.8	100.3	142.8	190.2	244.1
2001.....	5.5	11.8	20.5	36.2	51.3	80.1	105.9	154.9	182.6	249.9
2000.....	5.4	12.5	21.0	33.5	53.6	73.0	109.7	148.6	206.7	279.7
1999.....	5.5	11.4	19.6	35.7	56.4	78.1	109.4	151.6	204.0	273.9
Black, male										
2002.....	6.3	12.1	23.0	39.3	66.3	94.2	119.2	186.9	238.4	292.7
2001.....	5.9	13.5	22.0	41.6	57.2	97.8	124.9	184.9	231.1	307.5
2000.....	6.6	12.9	24.3	41.5	64.4	86.9	136.1	184.3	271.4	346.0
1999.....	5.1	12.3	21.6	44.4	67.7	94.8	126.5	200.9	238.9	304.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Colon (C18)										
Black, female										
2002.....	5.1	12.0	16.9	29.0	41.0	60.3	87.6	116.4	165.1	225.5
2001.....	5.1	10.3	19.2	31.8	46.7	67.0	93.3	137.0	157.8	228.6
2000.....	4.4	12.3	18.1	27.0	45.1	62.8	92.1	127.0	174.1	255.0
1999.....	5.9	10.7	18.0	28.5	47.6	65.8	98.1	121.9	186.7	262.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Rectosigmoid junction and rectum (C19-C20)										
All races 3/, both sexes										
2002.....	0.9	1.8	2.8	4.5	7.7	10.6	12.9	16.7	19.9	29.0
2001.....	0.8	1.8	3.2	4.7	6.6	10.0	12.5	16.3	21.4	32.6
2000.....	0.7	1.7	2.6	4.7	7.2	10.1	13.6	17.7	20.3	32.6
1999.....	0.8	1.6	2.5	4.6	7.3	10.1	13.5	17.6	20.6	32.1
All races 3/, male										
2002.....	1.1	2.2	3.6	6.1	10.8	14.2	16.6	22.4	26.9	33.6
2001.....	1.0	2.2	4.0	6.0	9.2	13.9	17.2	22.3	27.7	40.7
2000.....	0.9	2.2	3.2	6.3	9.7	14.2	18.7	23.5	27.0	38.9
1999.....	0.9	1.9	3.2	6.2	9.4	13.3	17.8	23.3	28.2	40.7
All races 3/, female										
2002.....	0.6	1.4	2.0	3.0	4.8	7.5	9.9	12.6	15.6	27.0
2001.....	0.7	1.5	2.4	3.3	4.2	6.6	8.8	12.0	17.6	29.2
2000.....	0.6	1.3	2.0	3.3	4.9	6.6	9.6	13.7	16.4	30.1
1999.....	0.8	1.4	1.9	3.1	5.3	7.4	10.1	13.6	16.2	28.7
White, both sexes										
2002.....	0.8	1.7	2.8	4.4	7.6	10.7	13.0	16.8	19.3	29.0
2001.....	0.8	1.8	3.1	4.5	6.6	10.1	12.4	16.4	21.1	32.6
2000.....	0.7	1.7	2.5	4.6	7.1	9.8	13.5	17.8	20.5	33.0
1999.....	0.8	1.6	2.4	4.5	7.1	10.0	13.5	17.5	20.7	32.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Rectosigmoid junction and rectum (C19-C20)										
White, male										
2002.....	1.0	2.0	3.6	5.9	10.8	14.3	16.7	22.7	25.6	33.7
2001.....	0.9	2.1	4.0	5.8	9.2	14.0	16.9	22.8	27.8	41.0
2000.....	0.9	2.1	3.1	5.9	9.7	13.6	18.4	23.7	27.2	39.8
1999.....	0.8	1.8	3.0	5.9	9.3	13.0	17.8	23.2	28.0	40.6
White, female										
2002.....	0.6	1.4	1.9	2.9	4.7	7.4	9.9	12.5	15.4	27.0
2001.....	0.7	1.4	2.2	3.3	4.2	6.8	8.7	11.9	17.1	29.1
2000.....	0.6	1.3	1.9	3.3	4.6	6.5	9.7	13.7	16.5	30.3
1999.....	0.8	1.4	1.8	3.2	5.2	7.4	10.1	13.5	16.4	28.9
Black, both sexes										
2002.....	1.3	2.3	3.5	6.0	9.0	11.6	13.9	17.2	27.7	29.7
2001.....	0.9	2.6	3.7	6.5	6.9	10.3	14.1	17.2	26.8	35.6
2000.....	1.0	2.5	3.4	6.1	8.8	13.6	15.6	17.6	19.2	28.8
1999.....	0.9	1.7	3.3	6.1	9.1	11.9	14.7	20.4	21.4	31.9
Black, male										
2002.....	1.7	3.5	4.4	7.9	12.5	14.5	18.7	21.1	45.9	34.3
2001.....	1.6	2.8	4.2	9.2	10.4	15.8	21.5	22.8	30.5	38.6
2000.....	*	3.5	4.2	9.6	11.4	21.9	23.4	22.7	27.0	30.0
1999.....	*	2.5	4.7	9.5	12.5	15.6	20.4	27.1	34.9	45.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Rectosigmoid junction and rectum (C19-C20)										
Black, female										
2002.....	*	*	2.7	4.5	6.3	9.4	10.8	14.8	18.3	27.9
2001.....	*	2.3	3.3	4.2	4.2	6.2	9.2	13.9	24.9	34.5
2000.....	*	1.7	2.8	3.2	6.7	7.5	10.4	14.5	15.3	28.3
1999.....	*	*	2.1	3.3	6.5	9.2	11.0	16.3	14.7	26.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anus and anus canal (C21)										
All races 3/, both sexes										
2002.....	0.1	0.2	0.3	0.3	0.6	0.5	0.6	0.8	1.0	1.2
2001.....	0.1	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.0	1.4
2000.....	0.1	0.2	0.3	0.3	0.5	0.5	0.7	0.9	0.8	1.2
1999.....	0.1	0.2	0.2	0.3	0.4	0.5	0.7	0.7	0.9	1.4
All races 3/, male										
2002.....	*	0.2	0.3	*	0.5	0.5	*	*	*	*
2001.....	*	*	0.2	0.3	*	0.5	0.6	0.8	*	*
2000.....	*	*	0.3	0.4	0.4	0.5	0.5	0.7	*	*
1999.....	*	0.2	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	*	0.2	0.3	0.4	0.6	0.6	0.7	0.9	1.1	1.2
2001.....	*	*	0.4	0.4	0.4	0.6	0.7	0.9	1.2	1.7
2000.....	*	0.3	0.3	0.3	0.5	0.5	0.8	1.0	0.7	1.3
1999.....	*	*	0.3	0.4	0.4	0.5	0.8	0.9	1.0	1.6
White, both sexes										
2002.....	0.1	0.2	0.4	0.4	0.6	0.6	0.6	0.8	1.0	1.2
2001.....	0.1	0.2	0.3	0.4	0.4	0.5	0.7	0.9	1.0	1.5
2000.....	0.1	0.2	0.3	0.3	0.5	0.5	0.7	0.8	0.8	1.1
1999.....	*	0.2	0.2	0.4	0.4	0.5	0.6	0.8	0.9	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of liver and intrahepatic bile ducts (C22)										
All races 3/, both sexes										
2002.....	1.3	3.4	6.6	8.1	11.6	16.9	22.6	27.6	31.6	35.0
2001.....	1.2	3.8	5.8	7.5	11.6	16.3	21.5	27.0	32.0	34.1
2000.....	1.4	3.7	5.4	7.7	11.3	16.0	21.6	27.1	31.3	32.7
1999.....	1.4	3.4	4.9	7.1	10.9	16.3	21.8	26.1	30.9	30.6
All races 3/, male										
2002.....	2.0	5.7	11.1	12.4	17.4	24.8	32.8	39.5	44.8	46.2
2001.....	1.8	6.0	9.7	11.4	17.2	23.5	30.6	39.2	43.9	50.3
2000.....	2.0	6.0	8.5	11.7	16.5	23.6	30.9	39.3	46.7	44.7
1999.....	2.1	5.5	7.6	10.6	15.6	23.7	31.9	36.6	42.9	40.1
All races 3/, female										
2002.....	0.6	1.2	2.4	4.0	6.4	10.1	14.4	19.2	23.6	30.2
2001.....	0.5	1.7	2.1	3.9	6.5	10.2	14.2	18.4	24.8	27.2
2000.....	0.8	1.4	2.4	4.0	6.5	9.5	14.2	18.6	22.2	27.8
1999.....	0.8	1.3	2.3	3.8	6.6	10.1	13.8	18.7	23.9	26.8
White, both sexes										
2002.....	1.1	3.0	5.5	6.8	10.6	15.8	21.1	26.7	30.7	33.8
2001.....	0.8	3.2	4.9	6.4	10.5	15.1	20.2	26.1	30.4	33.3
2000.....	1.1	3.1	4.4	6.6	10.0	14.6	20.2	25.8	30.7	32.0
1999.....	1.1	2.7	4.2	5.9	9.7	15.2	20.5	25.2	30.2	29.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of liver and intrahepatic bile ducts (C22)										
White, male										
2002.....	1.6	4.9	9.0	10.4	15.6	23.0	30.4	38.0	42.9	45.1
2001.....	1.3	5.0	8.1	9.5	15.7	22.1	29.1	37.7	42.6	49.7
2000.....	1.5	4.8	6.8	9.6	14.4	21.4	28.9	38.1	45.8	43.1
1999.....	1.7	4.3	6.3	8.8	13.9	21.9	30.2	35.3	42.4	40.6
White, female										
2002.....	0.5	1.1	2.0	3.4	5.9	9.3	13.5	18.6	23.2	29.0
2001.....	0.4	1.4	1.9	3.4	5.7	8.9	13.0	17.8	23.1	26.5
2000.....	0.6	1.3	2.0	3.6	5.9	8.5	13.2	17.1	21.7	27.4
1999.....	0.6	1.2	2.1	3.1	5.8	9.3	12.8	18.1	23.0	25.6
Black, both sexes										
2002.....	2.1	5.8	12.3	15.5	14.0	19.5	27.3	30.2	31.9	38.0
2001.....	2.3	7.1	10.7	15.2	14.3	20.0	24.0	26.9	40.9	35.0
2000.....	2.7	7.0	10.6	12.7	15.8	20.9	26.9	28.8	28.5	31.6
1999.....	2.2	6.6	8.1	12.0	14.7	21.1	28.0	26.6	33.8	34.4
Black, male										
2002.....	3.3	9.8	21.7	25.3	22.8	32.2	45.1	44.6	51.2	41.8
2001.....	3.6	11.6	18.6	25.3	21.4	28.0	32.7	39.5	49.0	42.0
2000.....	3.8	12.6	17.7	22.1	25.3	34.1	41.5	36.4	45.5	48.4
1999.....	3.4	11.8	14.2	18.6	21.6	32.2	42.1	40.4	42.7	30.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of liver and intrahepatic bile ducts (C22)										
Black, female										
2002.....	*	2.3	4.3	7.5	7.1	10.2	15.5	21.7	21.9	36.6
2001.....	*	3.2	3.9	6.8	8.8	14.1	18.2	19.4	36.8	32.4
2000.....	1.8	2.2	4.7	5.0	8.3	11.1	17.3	24.2	20.0	25.3
1999.....	*	2.0	2.9	6.5	9.3	12.9	18.7	18.3	29.4	35.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Liver (C22.0,C22.2-C22.9)										
All races 3/, both sexes										
2002.....	1.0	3.0	5.7	6.4	9.1	13.5	18.1	21.2	23.6	24.7
2001.....	0.9	3.4	5.0	6.2	9.2	12.8	16.7	20.6	23.8	24.2
2000.....	1.2	3.2	4.5	6.4	8.9	12.7	17.1	20.7	24.1	23.4
1999.....	1.2	3.0	4.1	5.7	8.7	13.0	17.5	20.4	23.3	22.5
All races 3/, male										
2002.....	1.7	5.1	10.0	10.6	14.5	20.9	27.6	31.4	34.8	34.6
2001.....	1.5	5.6	8.7	10.0	14.5	19.5	24.9	31.3	33.6	38.5
2000.....	1.8	5.4	7.6	10.2	14.0	19.7	25.5	31.4	37.6	33.9
1999.....	1.9	5.1	6.9	8.9	13.4	19.9	26.7	29.9	34.2	31.2
All races 3/, female										
2002.....	0.4	0.9	1.6	2.5	4.2	7.1	10.5	14.0	16.8	20.4
2001.....	0.4	1.2	1.5	2.5	4.5	7.1	10.1	13.1	17.8	18.2
2000.....	0.6	1.1	1.5	2.7	4.4	6.6	10.4	13.3	16.1	19.1
1999.....	0.6	0.9	1.5	2.7	4.5	7.1	10.3	13.7	16.9	18.9
White, both sexes										
2002.....	0.8	2.6	4.6	5.3	8.1	12.4	16.6	20.4	22.8	23.7
2001.....	0.6	2.8	4.2	5.1	8.2	11.6	15.5	19.6	22.5	23.5
2000.....	0.8	2.6	3.5	5.3	7.7	11.2	15.6	19.5	23.4	22.5
1999.....	0.9	2.3	3.4	4.6	7.6	11.8	16.1	19.4	22.5	21.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Liver (C22.0,C22.2-C22.9)										
White, male										
2002.....	1.4	4.3	8.0	8.7	12.8	19.3	25.1	30.2	33.2	33.5
2001.....	1.0	4.6	7.1	8.2	13.2	18.1	23.6	29.9	32.4	37.8
2000.....	1.3	4.2	5.9	8.4	12.0	17.5	23.4	30.2	36.6	31.8
1999.....	1.5	3.8	5.6	7.2	11.6	18.2	24.9	28.7	33.8	31.4
White, female										
2002.....	0.3	0.8	1.3	2.0	3.8	6.2	9.7	13.3	16.5	19.4
2001.....	0.2	1.0	1.3	2.1	3.6	5.8	8.9	12.4	16.5	17.5
2000.....	0.4	0.9	1.2	2.3	3.8	5.7	9.3	11.9	15.5	18.7
1999.....	0.4	0.8	1.3	2.1	3.8	6.3	9.1	13.0	15.9	17.8
Black, both sexes										
2002.....	1.9	5.3	11.5	13.8	12.0	16.5	23.9	24.2	24.9	28.8
2001.....	2.1	6.5	9.9	13.4	12.1	16.7	20.5	22.8	31.4	27.0
2000.....	2.5	6.5	9.8	11.3	13.1	18.1	23.1	23.8	24.3	24.7
1999.....	1.9	6.3	7.6	10.4	13.1	17.9	24.6	22.7	26.9	28.4
Black, male										
2002.....	3.2	9.2	21.0	23.8	20.4	27.4	40.1	36.3	41.3	35.4
2001.....	3.4	11.2	17.8	23.2	17.9	23.9	27.9	32.1	40.9	34.0
2000.....	3.4	12.0	17.0	20.3	23.0	30.7	36.4	30.7	40.5	42.7
1999.....	3.1	11.5	13.8	17.5	20.4	28.6	37.3	34.7	34.9	24.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Liver (C22.0,C22.2-C22.9)										
Black, female										
2002.....	*	1.9	3.5	5.5	5.4	8.3	13.0	17.0	16.3	26.3
2001.....	*	2.3	3.1	5.4	7.5	11.5	15.5	17.2	26.6	24.4
2000.....	1.6	1.8	3.7	4.0	5.2	8.8	14.2	19.7	16.1	18.0
1999.....	*	1.9	2.3	4.7	7.3	10.0	16.2	15.4	22.9	29.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intrahepatic bile duct carcinoma (C22.1)										
All races 3/, both sexes										
2002.....	0.2	0.4	0.9	1.6	2.5	3.4	4.5	6.4	8.0	10.3
2001.....	0.2	0.4	0.8	1.3	2.3	3.5	4.8	6.4	8.2	9.9
2000.....	0.2	0.5	0.9	1.4	2.3	3.3	4.5	6.4	7.2	9.3
1999.....	0.2	0.4	0.7	1.4	2.2	3.3	4.3	5.7	7.7	8.1
All races 3/, male										
2002.....	0.3	0.5	1.0	1.8	2.8	3.9	5.3	8.0	10.0	11.6
2001.....	0.3	0.4	1.0	1.3	2.7	4.0	5.7	7.9	10.3	11.9
2000.....	0.3	0.6	0.9	1.4	2.5	3.9	5.5	7.9	9.0	10.8
1999.....	0.2	0.4	0.7	1.7	2.3	3.8	5.2	6.7	8.8	8.9
All races 3/, female										
2002.....	0.2	0.3	0.8	1.5	2.3	3.0	3.9	5.3	6.8	9.8
2001.....	0.2	0.4	0.6	1.3	2.0	3.0	4.1	5.3	7.0	9.0
2000.....	0.2	0.4	0.9	1.3	2.2	2.8	3.8	5.4	6.1	8.8
1999.....	0.2	0.4	0.8	1.1	2.1	3.0	3.6	5.0	7.0	7.8
White, both sexes										
2002.....	0.2	0.4	0.8	1.5	2.5	3.4	4.5	6.4	7.9	10.2
2001.....	0.2	0.4	0.8	1.3	2.3	3.5	4.7	6.4	7.9	9.8
2000.....	0.2	0.5	0.9	1.3	2.2	3.4	4.6	6.3	7.3	9.5
1999.....	0.2	0.4	0.7	1.3	2.2	3.3	4.4	5.7	7.7	8.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of gallbladder and extrahepatic bile ducts (C23-C24)										
All races 3/, both sexes										
2002.....	0.2	0.4	0.8	1.4	2.0	3.7	5.7	8.3	9.9	14.5
2001.....	0.2	0.5	0.8	1.4	2.3	3.7	6.0	8.3	12.1	15.8
2000.....	0.2	0.4	0.9	1.5	2.6	4.2	6.4	9.1	11.4	16.0
1999.....	0.2	0.4	0.8	1.6	2.5	4.2	6.0	8.8	11.9	16.2
All races 3/, male										
2002.....	*	0.2	0.6	1.2	1.8	3.3	5.5	8.2	9.1	15.0
2001.....	*	0.3	0.7	1.2	1.7	2.8	5.3	8.1	12.8	15.8
2000.....	*	0.3	0.6	1.1	2.2	3.9	6.2	8.6	11.0	15.8
1999.....	0.2	0.3	0.6	1.2	2.1	3.8	5.4	8.3	11.0	15.2
All races 3/, female										
2002.....	0.3	0.6	1.0	1.6	2.2	4.1	5.8	8.3	10.3	14.3
2001.....	0.3	0.6	0.9	1.5	2.8	4.5	6.5	8.5	11.7	15.8
2000.....	0.2	0.5	1.1	1.8	2.9	4.5	6.6	9.4	11.7	16.0
1999.....	0.3	0.4	1.0	2.0	2.9	4.6	6.5	9.1	12.4	16.7
White, both sexes										
2002.....	0.2	0.4	0.8	1.3	2.0	3.7	5.6	8.1	9.8	14.5
2001.....	0.2	0.5	0.8	1.4	2.2	3.5	5.9	8.3	12.2	15.8
2000.....	0.2	0.4	0.8	1.5	2.5	4.2	6.3	9.2	11.6	16.3
1999.....	0.2	0.3	0.8	1.6	2.5	4.2	6.0	8.7	11.5	16.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of gallbladder and extrahepatic bile ducts (C23-C24)

Black, female

2002.....	*	*	*	2.6	*	5.6	6.0	10.7	9.9	16.4
2001.....	*	*	*	*	4.2	6.4	6.5	7.5	10.2	16.0
2000.....	*	*	2.2	*	3.6	6.9	6.0	7.7	9.3	12.0
1999.....	*	*	*	2.9	*	4.0	6.5	8.3	13.8	14.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pancreas (C25)										
All races 3/, both sexes										
2002.....	1.9	4.2	8.2	15.5	24.0	37.2	53.2	70.2	81.8	94.5
2001.....	1.8	4.0	8.6	14.9	24.8	37.0	52.8	67.2	84.2	100.1
2000.....	1.8	4.2	8.7	15.1	24.8	37.2	51.4	70.4	85.7	96.4
1999.....	1.9	4.5	8.7	15.6	25.0	38.1	52.5	67.2	82.0	99.7
All races 3/, male										
2002.....	2.4	5.3	10.6	19.2	29.3	43.3	60.8	78.9	91.9	100.9
2001.....	2.0	4.9	10.7	18.5	30.3	43.6	59.0	77.2	94.9	105.1
2000.....	2.1	5.4	11.1	18.5	30.0	44.7	57.0	80.4	94.9	102.9
1999.....	2.4	5.9	11.0	19.1	29.9	44.9	60.9	75.7	92.0	107.5
All races 3/, female										
2002.....	1.4	3.1	6.0	12.0	19.2	31.9	47.1	64.1	75.6	91.7
2001.....	1.6	3.1	6.5	11.5	19.8	31.3	47.8	60.1	77.8	98.1
2000.....	1.5	3.0	6.4	11.8	20.0	30.8	47.1	63.5	80.2	93.8
1999.....	1.4	3.1	6.5	12.3	20.5	32.2	46.0	61.2	76.1	96.6
White, both sexes										
2002.....	1.9	4.1	7.8	15.0	23.3	36.1	52.0	69.2	81.4	94.3
2001.....	1.7	3.8	8.3	14.4	24.7	36.3	51.7	66.3	83.3	99.0
2000.....	1.7	4.1	8.4	14.5	24.3	36.2	50.4	69.2	84.5	94.7
1999.....	1.8	4.4	8.2	14.8	24.3	37.2	51.5	65.8	80.6	98.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pancreas (C25)										
White, male										
2002.....	2.4	5.1	10.1	18.6	28.8	42.6	59.7	78.9	92.2	101.6
2001.....	1.9	4.7	10.5	18.2	30.3	42.8	58.0	77.0	94.5	106.5
2000.....	2.0	5.3	10.7	17.8	29.6	44.3	55.9	79.9	94.0	102.4
1999.....	2.3	5.8	10.3	17.9	29.2	44.0	60.1	74.6	90.9	107.4
White, female										
2002.....	1.4	3.0	5.5	11.6	18.2	30.3	45.7	62.2	74.9	91.1
2001.....	1.6	2.9	6.3	10.8	19.6	30.6	46.5	58.7	76.5	95.9
2000.....	1.4	2.9	6.1	11.4	19.4	29.2	46.0	61.6	78.9	91.5
1999.....	1.3	3.1	6.3	11.9	19.9	31.3	44.6	59.5	74.5	95.0
Black, both sexes										
2002.....	2.5	5.6	12.3	22.4	34.4	50.4	73.7	88.6	91.8	106.6
2001.....	2.2	6.4	12.1	20.7	31.6	48.5	70.8	84.9	104.8	121.3
2000.....	2.6	5.8	13.2	22.8	33.3	50.3	69.8	93.8	104.2	119.8
1999.....	2.8	6.2	14.1	25.0	32.2	50.9	70.1	87.5	104.6	118.6
Black, male										
2002.....	2.9	7.2	16.5	29.0	40.8	56.6	84.1	90.5	105.5	100.8
2001.....	2.3	7.9	15.2	24.2	38.2	60.0	78.8	95.2	109.2	99.8
2000.....	3.0	7.3	17.5	29.3	40.4	55.7	78.9	101.6	116.3	109.6
1999.....	3.4	8.3	19.5	34.1	39.5	60.0	78.4	100.7	117.7	114.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pancreas (C25)										
Black, female										
2002.....	2.1	4.3	8.8	17.0	29.4	45.9	66.8	87.5	84.7	108.8
2001.....	2.2	5.1	9.6	17.8	26.4	40.0	65.5	78.8	102.6	129.2
2000.....	2.3	4.4	9.4	17.4	27.8	46.3	63.7	89.1	98.1	123.6
1999.....	2.2	4.3	9.5	17.6	26.5	44.3	64.6	79.5	98.1	120.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of other and ill-defined digestive organs (C26)										
All races 3/, both sexes										
2002.....	*	0.1	0.2	0.5	0.7	1.3	2.0	3.4	4.1	6.9
2001.....	*	0.2	0.3	0.5	0.9	1.3	2.2	3.2	4.3	7.8
2000.....	*	0.2	0.3	0.5	1.0	1.3	2.1	2.8	4.5	7.8
1999.....	0.1	0.3	0.3	0.6	1.0	1.6	2.3	3.3	5.3	9.1
All races 3/, male										
2002.....	*	*	0.3	0.7	0.8	2.0	2.6	4.3	5.3	6.4
2001.....	*	0.2	0.4	0.7	1.3	1.4	2.7	4.1	4.5	8.9
2000.....	*	*	0.3	0.7	1.3	1.6	3.1	3.4	5.2	7.5
1999.....	0.2	0.3	0.5	0.7	1.3	1.9	2.6	4.2	6.9	10.4
All races 3/, female										
2002.....	*	*	*	0.3	0.7	0.8	1.5	2.7	3.4	7.2
2001.....	*	*	*	0.3	0.7	1.2	1.8	2.6	4.2	7.3
2000.....	*	*	0.2	0.4	0.7	1.0	1.4	2.3	4.1	8.0
1999.....	*	*	*	0.4	0.7	1.4	2.1	2.7	4.4	8.5
White, both sexes										
2002.....	*	*	0.2	0.5	0.7	1.3	1.9	3.5	4.0	7.0
2001.....	*	0.1	0.2	0.5	0.8	1.3	2.2	3.2	4.1	7.7
2000.....	*	0.2	0.2	0.6	1.0	1.2	2.0	2.8	4.3	7.9
1999.....	0.1	0.2	0.3	0.5	1.0	1.6	2.3	3.3	5.5	9.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)										
All races 3/, both sexes										
2002.....	9.6	21.5	43.4	89.5	157.8	239.7	328.9	386.5	387.5	305.1
2001.....	10.0	21.6	44.1	92.3	159.7	246.4	331.7	382.6	377.8	312.2
2000.....	9.8	22.0	45.6	95.9	165.1	252.9	333.8	377.8	385.1	310.8
1999.....	9.7	20.9	46.4	96.7	166.0	250.7	331.0	370.6	372.2	306.6
All races 3/, male										
2002.....	10.1	25.3	54.2	112.3	201.0	307.8	436.0	532.5	563.8	498.8
2001.....	11.2	25.9	55.2	115.5	204.0	323.8	449.3	538.5	558.1	534.1
2000.....	11.2	27.2	56.2	121.6	211.2	333.9	455.0	533.6	578.3	541.7
1999.....	11.2	25.9	57.7	124.3	216.4	334.1	453.2	529.0	569.3	550.6
All races 3/, female										
2002.....	9.0	17.8	32.9	68.0	118.6	180.9	243.4	283.4	280.3	221.1
2001.....	8.7	17.3	33.6	70.6	119.5	179.8	238.3	273.4	269.7	219.3
2000.....	8.4	17.0	35.5	71.8	123.3	183.5	238.3	269.3	271.1	216.8
1999.....	8.4	16.1	35.5	70.9	120.4	179.4	235.5	260.5	256.8	208.3
White, both sexes										
2002.....	9.1	20.4	41.6	87.8	157.2	242.4	334.8	391.2	393.1	309.0
2001.....	9.5	20.2	42.6	91.1	160.7	248.3	335.4	385.9	380.2	314.8
2000.....	9.0	20.4	43.8	94.4	165.8	254.9	335.8	379.8	391.0	313.4
1999.....	9.2	19.3	44.3	95.3	165.4	251.2	333.1	371.7	374.7	308.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)										
White, male										
2002.....	9.4	23.3	50.6	107.5	194.8	305.5	436.9	530.6	564.7	503.6
2001.....	10.4	23.6	51.5	110.8	201.1	319.8	446.3	534.5	554.0	535.5
2000.....	9.8	24.8	52.0	116.2	208.2	329.8	450.1	527.8	580.1	543.3
1999.....	10.2	23.2	53.4	119.7	210.9	328.7	448.4	524.2	565.4	551.9
White, female										
2002.....	8.9	17.5	32.7	68.9	122.4	186.8	251.5	291.5	287.9	224.5
2001.....	8.6	16.9	33.8	72.2	123.4	185.6	245.6	280.6	275.2	222.3
2000.....	8.2	16.0	35.7	73.7	126.7	189.5	244.1	275.7	278.5	220.1
1999.....	8.2	15.4	35.5	72.1	123.5	183.7	241.5	264.7	262.1	210.1
Black, both sexes										
2002.....	14.9	34.8	68.4	127.3	203.6	278.0	343.7	407.8	378.3	278.0
2001.....	15.4	35.0	67.8	127.4	191.9	284.8	363.2	410.1	381.9	290.2
2000.....	16.8	37.8	74.0	131.6	200.8	291.8	375.8	412.1	357.5	295.9
1999.....	15.7	37.5	75.3	133.2	209.9	294.7	364.5	414.0	375.3	300.4
Black, male										
2002.....	18.1	46.6	98.0	184.4	311.8	417.4	531.1	666.4	642.7	498.5
2001.....	19.6	48.9	100.0	189.0	285.8	440.1	582.0	691.4	685.4	571.9
2000.....	23.1	51.1	108.6	200.9	294.2	453.8	593.7	688.0	645.7	584.7
1999.....	20.7	53.3	111.5	201.4	321.8	451.3	581.4	680.1	694.9	599.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)										
Black, female										
2002.....	12.0	24.6	43.2	80.4	118.9	174.6	218.0	253.3	240.7	193.5
2001.....	11.7	23.0	40.3	77.0	118.1	170.1	217.5	242.1	227.3	185.7
2000.....	11.3	26.3	44.4	74.8	127.0	172.2	231.1	245.8	212.3	188.4
1999.....	11.3	23.9	44.7	77.5	122.1	178.7	220.5	253.7	216.9	188.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of nasal cavity, middle ear and accessory sinuses (C30-C31)										
All races 3/, both sexes										
2002.....	*	0.2	0.2	0.2	0.4	0.4	0.7	0.7	1.0	1.2
2001.....	0.1	0.1	0.2	0.3	0.4	0.4	0.6	0.8	1.4	1.7
2000.....	0.1	*	0.2	0.3	0.4	0.5	0.6	0.6	0.9	1.2
1999.....	0.1	0.1	0.3	0.3	0.3	0.5	0.6	0.8	1.0	1.5
All races 3/, male										
2002.....	*	0.3	0.3	0.3	0.6	0.7	1.0	1.0	1.1	1.7
2001.....	*	0.2	0.3	0.4	0.5	0.5	0.8	0.9	1.9	1.8
2000.....	*	*	0.2	0.4	0.6	0.7	0.9	1.0	1.1	*
1999.....	*	*	0.4	0.5	0.4	0.7	0.6	1.2	1.1	1.8
All races 3/, female										
2002.....	*	*	*	*	*	*	0.5	0.6	0.8	1.0
2001.....	*	*	*	*	*	*	0.4	0.8	1.1	1.6
2000.....	*	*	*	*	*	*	0.4	*	0.8	1.1
1999.....	*	*	*	*	*	*	0.6	0.6	0.9	1.4
White, both sexes										
2002.....	*	0.2	0.2	0.3	0.4	0.4	0.7	0.8	1.0	1.3
2001.....	*	0.2	0.2	0.3	0.4	0.4	0.5	0.8	1.4	1.7
2000.....	*	*	0.2	0.3	0.4	0.5	0.6	0.7	0.9	1.3
1999.....	0.1	*	0.3	0.3	0.3	0.5	0.7	0.9	1.0	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of larynx (C32)										
All races 3/, both sexes										
2002.....	0.2	0.7	1.4	2.7	4.2	5.3	6.6	7.9	7.2	5.5
2001.....	0.3	0.7	1.5	2.9	4.5	5.6	7.3	7.4	6.9	6.7
2000.....	0.2	0.8	1.5	2.9	4.5	6.0	7.3	8.0	8.1	6.5
1999.....	0.3	0.8	1.5	3.1	4.6	6.1	7.4	7.2	7.4	7.0
All races 3/, male										
2002.....	0.4	1.2	2.4	4.7	7.2	8.9	11.3	14.8	14.9	13.2
2001.....	0.5	1.2	2.5	4.9	7.5	9.5	12.9	14.5	13.9	18.3
2000.....	0.4	1.2	2.5	4.9	7.9	10.0	12.9	14.8	17.1	17.4
1999.....	0.4	1.3	2.6	5.4	7.8	10.4	12.6	13.2	15.8	17.8
All races 3/, female										
2002.....	*	0.2	0.5	0.9	1.5	2.2	2.9	3.1	2.5	2.2
2001.....	*	0.2	0.5	1.0	1.7	2.2	2.8	2.5	2.8	1.9
2000.....	*	0.3	0.6	0.9	1.4	2.6	2.8	3.3	2.8	2.1
1999.....	*	0.2	0.4	1.0	1.7	2.5	3.3	3.0	2.6	2.6
White, both sexes										
2002.....	0.2	0.6	1.2	2.5	3.9	5.0	6.4	7.5	6.9	5.6
2001.....	0.3	0.6	1.2	2.5	4.1	5.2	6.7	7.0	6.8	6.6
2000.....	0.2	0.6	1.3	2.4	3.9	5.6	6.7	7.5	8.0	6.6
1999.....	0.2	0.6	1.3	2.7	4.2	5.7	7.1	7.1	7.3	6.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of larynx (C32)										
White, male										
2002.....	0.3	1.0	2.0	4.1	6.3	8.3	10.7	13.8	14.6	13.3
2001.....	0.4	0.9	2.0	4.2	6.9	8.8	11.9	13.3	13.2	18.3
2000.....	0.3	1.0	2.1	4.1	6.8	9.2	11.7	13.9	16.8	17.6
1999.....	0.3	1.1	2.2	4.6	7.0	9.6	12.0	12.9	15.1	16.8
White, female										
2002.....	*	*	0.4	0.9	1.5	2.1	2.8	2.9	2.3	2.3
2001.....	*	*	0.4	0.9	1.5	2.1	2.6	2.5	2.9	1.7
2000.....	*	0.3	0.5	0.9	1.3	2.5	2.6	3.0	2.7	2.1
1999.....	*	*	0.3	1.0	1.5	2.3	3.3	3.0	2.7	2.7
Black, both sexes										
2002.....	*	1.9	3.6	6.1	9.3	9.1	11.8	15.3	11.3	*
2001.....	*	1.9	3.8	7.1	8.9	9.5	13.9	13.4	9.2	8.3
2000.....	*	2.0	4.1	7.5	10.8	10.6	15.3	15.6	11.6	6.9
1999.....	0.9	2.1	3.9	7.5	10.0	11.4	11.8	10.2	9.8	8.3
Black, male										
2002.....	*	3.2	6.0	11.5	18.8	17.0	22.4	33.1	22.9	*
2001.....	*	3.6	7.0	13.0	15.9	17.5	26.3	30.7	22.5	*
2000.....	*	3.2	7.6	14.3	20.7	19.8	30.1	29.3	27.0	*
1999.....	1.6	3.9	6.9	14.4	17.8	21.0	22.7	19.9	27.9	24.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of larynx (C32)										
Black, female										
2002.....	*	*	*	*	*	*	4.7	*	*	*
2001.....	*	*	*	*	3.4	*	5.6	*	*	*
2000.....	*	*	*	*	*	3.8	5.5	7.4	*	*
1999.....	*	*	*	*	3.8	4.2	4.5	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of trachea, bronchus and lung (C33-C34)										
All races 3/, both sexes										
2002.....	9.2	20.4	41.6	86.3	152.9	233.4	320.9	376.9	378.5	297.2
2001.....	9.5	20.5	42.3	88.8	154.5	239.7	323.2	373.4	368.6	302.7
2000.....	9.3	21.0	43.8	92.5	159.8	245.8	325.4	368.2	374.7	302.1
1999.....	9.3	19.9	44.4	93.0	160.7	243.6	322.3	361.7	362.6	297.0
All races 3/, male										
2002.....	9.6	23.6	51.4	107.0	192.7	297.5	423.0	515.5	546.8	482.2
2001.....	10.5	24.3	52.2	109.6	195.7	312.9	434.7	522.3	541.0	512.4
2000.....	10.5	25.7	53.2	115.9	202.3	322.5	440.5	516.5	558.1	521.2
1999.....	10.5	24.3	54.4	118.1	207.7	322.4	439.2	513.4	551.3	529.2
All races 3/, female										
2002.....	8.8	17.3	32.2	66.8	116.8	177.9	239.3	279.1	276.1	217.0
2001.....	8.4	16.8	32.8	69.2	117.2	176.7	234.5	269.1	265.2	215.0
2000.....	8.1	16.5	34.7	70.5	121.3	180.2	234.7	264.9	266.6	212.8
1999.....	8.1	15.7	34.9	69.5	118.1	176.1	230.9	256.2	252.2	203.5
White, both sexes										
2002.....	8.8	19.4	40.0	84.9	152.6	236.4	327.0	382.1	384.1	300.9
2001.....	9.1	19.3	40.9	87.9	155.9	242.0	327.4	377.2	371.2	305.4
2000.....	8.6	19.5	42.2	91.5	161.2	248.2	328.0	370.7	380.7	304.5
1999.....	8.8	18.4	42.6	92.0	160.6	244.5	324.6	362.9	365.2	298.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of trachea, bronchus and lung (C33-C34)										
White, male										
2002.....	8.9	21.8	48.1	102.8	187.3	295.9	424.5	514.5	547.8	486.6
2001.....	9.7	22.2	49.0	105.6	193.4	309.8	432.8	519.4	537.4	513.9
2000.....	9.3	23.5	49.5	111.5	200.5	319.1	436.8	511.6	560.0	522.4
1999.....	9.7	21.9	50.5	114.3	203.1	317.9	435.0	508.8	548.0	531.5
White, female										
2002.....	8.7	17.1	32.0	67.7	120.6	184.0	247.5	287.3	283.8	220.2
2001.....	8.4	16.4	33.1	71.0	121.3	182.5	242.1	276.4	270.7	218.1
2000.....	8.0	15.6	35.0	72.5	125.0	186.2	240.7	271.6	274.1	215.9
1999.....	7.9	15.0	35.0	70.8	121.6	180.5	236.9	260.4	257.3	205.3
Black, both sexes										
2002.....	14.1	32.4	64.4	120.6	193.7	267.2	330.7	391.2	366.3	272.3
2001.....	14.5	32.6	63.5	119.3	182.2	273.6	347.6	395.1	370.3	280.7
2000.....	15.7	35.5	69.3	123.1	188.7	279.9	359.1	394.9	343.3	288.1
1999.....	14.5	35.1	70.8	125.2	198.4	282.1	351.1	402.9	363.5	290.5
Black, male										
2002.....	17.0	42.6	91.6	172.1	292.0	398.4	506.4	630.6	619.0	484.6
2001.....	18.3	45.0	92.4	174.1	268.7	418.8	552.7	659.8	659.7	549.2
2000.....	21.4	47.4	100.4	185.2	271.7	432.2	561.6	656.8	617.1	562.8
1999.....	18.6	48.8	103.6	186.3	301.7	429.2	556.6	658.7	665.3	574.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of trachea, bronchus and lung (C33-C34)										
Black, female										
2002.....	11.5	23.6	41.5	78.3	116.8	169.8	212.8	248.1	234.7	191.0
2001.....	11.1	21.9	38.8	74.3	114.1	166.3	211.0	237.1	222.8	181.0
2000.....	10.7	25.1	42.7	72.2	123.2	167.6	224.7	237.0	205.5	185.9
1999.....	10.8	23.3	42.9	75.3	117.3	173.2	214.7	248.9	213.9	184.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of thymus, heart, mediastinum and pleura (C37-C38)										
All races 3/, both sexes										
2002.....	*	0.2	0.1	0.2	0.3	0.5	0.6	0.8	0.8	1.1
2001.....	0.1	0.2	0.2	0.3	0.3	0.6	0.6	0.8	0.9	1.0
2000.....	0.1	0.1	0.1	0.2	0.3	0.5	0.5	0.8	1.3	0.9
1999.....	*	0.1	0.2	0.3	0.3	0.4	0.7	0.8	1.0	1.0
All races 3/, male										
2002.....	*	0.2	*	*	0.5	0.6	0.6	1.0	*	1.7
2001.....	*	0.2	0.2	0.5	*	0.7	0.8	0.8	1.2	1.5
2000.....	*	*	*	*	0.4	0.6	0.6	1.1	1.8	*
1999.....	*	*	0.3	0.3	0.4	*	0.7	1.0	*	*
All races 3/, female										
2002.....	*	*	*	*	*	0.4	0.6	0.6	0.7	0.9
2001.....	*	*	*	*	*	0.4	0.5	0.8	0.7	0.8
2000.....	*	*	*	*	*	0.4	*	0.6	0.9	0.7
1999.....	*	*	*	*	*	0.4	0.6	0.7	1.0	0.8
White, both sexes										
2002.....	*	0.2	0.1	*	0.3	0.5	0.7	0.8	0.8	1.3
2001.....	0.1	0.2	0.2	0.3	0.3	0.5	0.6	0.7	0.9	1.0
2000.....	*	*	*	0.2	0.3	0.5	0.5	0.8	1.3	0.9
1999.....	*	*	0.2	0.3	0.3	0.4	0.6	0.8	1.1	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of bone and articular cartilage (C40-C41)										
All races 3/, both sexes										
2002.....	0.2	0.3	0.3	0.5	0.7	0.6	1.0	1.3	2.0	2.9
2001.....	0.2	0.3	0.3	0.5	0.6	0.9	1.3	1.7	2.2	3.5
2000.....	0.2	0.3	0.3	0.5	0.6	0.9	1.0	1.8	2.1	3.4
1999.....	0.2	0.3	0.4	0.3	0.6	0.8	1.1	1.6	2.1	3.0
All races 3/, male										
2002.....	0.3	0.4	0.4	0.6	0.7	0.7	1.1	1.8	2.4	3.7
2001.....	0.3	0.4	0.4	0.5	0.7	1.1	1.7	2.3	2.9	3.6
2000.....	0.2	0.4	0.5	0.7	0.7	1.2	1.2	2.2	2.7	4.2
1999.....	0.3	0.4	0.5	0.4	0.8	0.9	1.6	1.8	2.2	4.5
All races 3/, female										
2002.....	*	*	*	0.4	0.6	0.6	1.0	1.0	1.8	2.5
2001.....	*	*	0.2	0.5	0.5	0.7	1.0	1.3	1.8	3.4
2000.....	*	*	*	0.4	0.5	0.7	0.8	1.5	1.8	3.1
1999.....	*	0.3	0.3	*	0.4	0.8	0.8	1.5	2.0	2.4
White, both sexes										
2002.....	0.2	0.3	0.3	0.5	0.6	0.6	1.0	1.3	2.1	3.0
2001.....	0.2	0.2	0.3	0.5	0.6	0.8	1.3	1.7	2.3	3.6
2000.....	0.2	0.3	0.3	0.5	0.6	1.0	1.0	1.8	2.2	3.5
1999.....	0.2	0.3	0.4	0.3	0.5	0.9	1.2	1.7	2.2	3.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Melanoma and other malignant neoplasms of skin (C43-C44)										
All races 3/, both sexes										
2002.....	1.8	2.6	4.1	5.4	7.7	10.0	13.2	17.8	21.8	29.8
2001.....	1.9	2.9	4.4	5.9	8.0	10.2	14.2	17.1	21.0	31.2
2000.....	1.9	3.0	4.5	5.8	7.8	10.5	13.3	16.6	20.3	31.8
1999.....	1.9	3.1	4.2	5.7	8.0	10.6	13.1	17.2	21.3	29.2
All races 3/, male										
2002.....	2.4	3.8	5.6	7.7	11.5	14.9	20.2	28.3	35.3	50.2
2001.....	2.4	3.8	5.9	8.3	11.9	15.6	22.2	27.1	34.5	54.0
2000.....	2.4	3.7	6.3	8.3	11.0	15.5	20.4	26.5	34.7	55.4
1999.....	2.5	4.2	5.7	8.4	11.6	15.5	20.5	26.9	35.2	50.6
All races 3/, female										
2002.....	1.3	1.5	2.7	3.2	4.3	5.7	7.6	10.5	13.6	21.0
2001.....	1.5	2.1	2.9	3.6	4.5	5.5	7.8	10.1	12.9	21.7
2000.....	1.4	2.2	2.7	3.4	4.9	6.2	7.8	9.7	11.8	22.2
1999.....	1.3	2.0	2.8	3.2	4.7	6.3	7.2	10.4	13.1	20.6
White, both sexes										
2002.....	2.2	3.1	4.7	6.0	8.6	11.1	14.5	19.6	23.8	32.2
2001.....	2.3	3.4	5.0	6.6	8.9	11.3	15.5	18.7	22.6	33.7
2000.....	2.2	3.4	5.1	6.4	8.7	11.6	14.5	17.9	21.8	34.3
1999.....	2.3	3.5	4.8	6.4	8.9	11.8	14.3	18.5	22.9	31.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Melanoma and other malignant neoplasms of skin (C43-C44)										
White, male										
2002.....	2.8	4.4	6.2	8.4	12.8	16.4	21.9	30.9	38.5	54.5
2001.....	2.8	4.3	6.7	9.3	13.1	17.1	24.1	29.6	37.4	58.7
2000.....	2.7	4.3	7.0	9.1	12.2	17.0	22.1	28.6	37.2	60.6
1999.....	3.0	4.7	6.3	9.3	12.8	17.2	22.3	28.7	37.9	55.0
White, female										
2002.....	1.6	1.8	3.2	3.7	4.8	6.4	8.4	11.5	14.7	22.5
2001.....	1.8	2.5	3.4	4.1	5.0	6.2	8.5	11.0	13.7	23.2
2000.....	1.7	2.6	3.2	3.7	5.4	6.9	8.4	10.4	12.6	23.6
1999.....	1.5	2.3	3.3	3.7	5.3	7.0	7.9	11.3	14.1	22.1
Black, both sexes										
2002.....	*	*	1.7	2.5	2.4	3.6	3.9	3.8	*	7.7
2001.....	*	*	1.2	1.6	3.0	4.0	5.1	3.8	6.0	8.0
2000.....	*	*	1.3	2.8	2.9	3.6	4.9	4.8	6.8	8.1
1999.....	*	1.0	1.5	1.7	2.9	2.8	3.7	5.4	*	6.7
Black, male										
2002.....	*	*	2.8	4.5	*	6.2	*	*	*	*
2001.....	*	*	*	*	4.1	6.9	7.7	*	*	*
2000.....	*	*	*	4.1	4.3	5.2	*	*	*	*
1999.....	*	*	2.6	*	5.1	*	*	10.4	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant melanoma of skin (C43)										
All races 3/, both sexes										
2002.....	1.7	2.3	3.4	4.4	6.0	7.8	10.0	12.9	14.9	17.9
2001.....	1.8	2.6	3.6	4.6	6.1	7.8	10.7	12.4	14.7	17.8
2000.....	1.7	2.7	3.8	4.7	6.3	7.9	10.0	12.0	14.9	18.9
1999.....	1.7	2.6	3.6	4.7	6.1	8.3	9.7	12.9	14.2	16.4
All races 3/, male										
2002.....	2.1	3.3	4.4	6.1	8.5	11.3	14.9	19.8	23.3	29.2
2001.....	2.2	3.2	4.7	6.3	8.8	11.9	16.3	19.6	24.0	29.9
2000.....	2.1	3.2	5.2	6.5	8.7	11.2	14.7	18.4	24.7	31.8
1999.....	2.2	3.5	4.6	6.7	8.4	12.0	14.6	19.8	22.7	27.5
All races 3/, female										
2002.....	1.2	1.4	2.5	2.8	3.6	4.8	6.1	7.9	9.8	12.9
2001.....	1.4	1.9	2.6	3.0	3.7	4.3	6.4	7.3	9.2	12.8
2000.....	1.4	2.1	2.4	3.0	4.1	5.0	6.3	7.5	9.1	13.7
1999.....	1.3	1.8	2.6	2.8	4.0	5.2	5.8	8.1	9.2	11.9
White, both sexes										
2002.....	2.0	2.8	4.0	5.0	6.8	8.9	11.0	14.2	16.3	19.2
2001.....	2.2	3.1	4.3	5.3	7.0	8.8	11.9	13.6	15.9	19.3
2000.....	2.0	3.1	4.4	5.3	7.2	8.9	11.0	13.1	16.0	20.3
1999.....	2.1	3.1	4.2	5.4	6.9	9.4	10.7	14.0	15.3	17.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other malignant neoplasms of skin (C44)										
All races 3/, both sexes										
2002.....	0.2	0.3	0.7	1.0	1.7	2.2	3.2	5.0	6.9	12.0
2001.....	0.1	0.3	0.7	1.3	1.9	2.4	3.5	4.8	6.3	13.4
2000.....	0.1	0.3	0.7	1.1	1.5	2.6	3.3	4.6	5.4	12.9
1999.....	0.2	0.4	0.7	1.0	1.9	2.3	3.4	4.3	7.0	12.8
All races 3/, male										
2002.....	0.2	0.5	1.2	1.6	3.0	3.7	5.3	8.5	12.0	21.0
2001.....	0.2	0.5	1.2	2.1	3.1	3.7	5.9	7.5	10.5	24.1
2000.....	0.2	0.5	1.1	1.9	2.4	4.2	5.7	8.1	9.9	23.6
1999.....	0.3	0.7	1.1	1.7	3.2	3.6	5.9	7.1	12.4	23.1
All races 3/, female										
2002.....	*	*	0.3	0.5	0.7	0.9	1.6	2.5	3.8	8.1
2001.....	*	*	0.2	0.5	0.8	1.2	1.5	2.8	3.7	8.9
2000.....	*	*	0.3	0.4	0.8	1.2	1.5	2.2	2.7	8.5
1999.....	*	*	0.3	0.4	0.7	1.1	1.4	2.4	3.9	8.6
White, both sexes										
2002.....	0.2	0.3	0.7	1.0	1.8	2.2	3.4	5.4	7.4	13.0
2001.....	0.1	0.3	0.7	1.3	1.9	2.5	3.6	5.1	6.7	14.3
2000.....	0.1	0.3	0.7	1.1	1.5	2.7	3.5	4.9	5.8	14.0
1999.....	0.2	0.4	0.6	1.0	2.0	2.4	3.6	4.5	7.6	13.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of mesothelial and soft tissue (C45-C49)										
All races 3/, both sexes										
2002.....	0.9	1.2	2.0	3.4	5.5	7.4	10.9	13.8	16.3	15.7
2001.....	0.9	1.4	2.3	3.4	5.3	7.7	10.6	14.7	15.7	14.9
2000.....	0.9	1.4	2.2	3.7	5.3	8.1	11.2	14.4	16.2	15.7
1999.....	1.0	1.4	2.3	3.5	5.6	7.7	11.0	14.0	15.9	14.8
All races 3/, male										
2002.....	1.1	1.3	2.0	3.7	6.7	9.4	15.6	20.8	25.1	28.1
2001.....	1.0	1.5	2.6	3.6	6.3	10.0	15.3	21.3	24.9	25.9
2000.....	0.9	1.5	2.2	4.2	6.6	10.8	15.2	21.9	26.2	28.4
1999.....	1.3	1.4	2.5	4.3	6.9	10.4	15.6	20.8	25.7	24.6
All races 3/, female										
2002.....	0.8	1.1	1.9	3.1	4.4	5.7	7.1	8.9	11.0	10.4
2001.....	0.9	1.2	2.0	3.2	4.3	5.7	6.8	10.1	10.2	10.3
2000.....	0.9	1.3	2.2	3.2	4.1	5.8	8.0	9.1	10.3	10.5
1999.....	0.7	1.5	2.0	2.7	4.4	5.3	7.4	9.3	10.1	10.8
White, both sexes										
2002.....	0.9	1.2	2.0	3.4	5.6	7.7	11.4	14.6	17.0	16.5
2001.....	0.9	1.3	2.3	3.4	5.6	8.3	11.1	15.4	16.5	15.4
2000.....	0.9	1.4	2.2	3.8	5.5	8.3	11.7	15.1	17.1	16.2
1999.....	0.9	1.5	2.3	3.6	5.7	8.1	11.4	14.4	16.3	15.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of mesothelial and soft tissue (C45-C49)										
White, male										
2002.....	1.0	1.3	2.1	3.8	7.0	10.0	16.4	22.0	26.5	29.8
2001.....	0.9	1.4	2.7	3.6	6.8	10.8	16.3	22.4	26.2	27.0
2000.....	0.9	1.6	2.3	4.5	6.9	11.1	16.0	23.1	27.6	29.5
1999.....	1.2	1.4	2.6	4.6	7.1	11.0	16.2	21.6	26.9	26.2
White, female										
2002.....	0.8	1.1	1.9	3.1	4.4	5.8	7.3	9.3	11.2	10.7
2001.....	0.8	1.2	1.9	3.2	4.5	6.0	6.9	10.4	10.7	10.5
2000.....	0.9	1.3	2.0	3.2	4.2	5.9	8.3	9.5	10.9	10.7
1999.....	0.6	1.5	2.0	2.7	4.4	5.5	7.7	9.4	10.1	10.9
Black, both sexes										
2002.....	1.3	1.7	2.0	3.8	4.7	6.6	7.5	9.1	11.8	7.7
2001.....	1.3	1.7	2.9	3.9	3.4	4.5	7.5	8.7	8.1	11.1
2000.....	1.4	1.9	2.6	3.8	4.8	7.7	8.8	8.5	7.9	10.9
1999.....	1.5	1.5	2.4	2.9	6.1	5.1	8.4	11.4	11.6	10.5
Black, male										
2002.....	1.7	2.3	*	4.0	4.8	6.7	9.7	11.9	*	*
2001.....	1.5	*	2.9	4.4	*	5.1	8.1	10.7	*	*
2000.....	*	1.8	*	4.0	6.2	9.4	11.0	12.3	*	*
1999.....	1.9	*	2.5	3.6	7.2	6.0	10.5	14.7	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of mesothelial and soft tissue (C45-C49)

Black, female

2002.....	*	*	2.1	3.5	4.6	6.5	6.0	7.4	10.3	*
2001.....	*	1.8	2.9	3.6	3.5	4.1	7.2	7.5	*	8.8
2000.....	1.3	1.9	3.7	3.6	3.8	6.5	7.3	6.3	*	8.6
1999.....	*	1.6	2.3	*	5.2	4.4	7.0	9.4	10.4	10.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mesothelioma (C45)										
All races 3/, both sexes										
2002.....	0.1	0.1	0.4	1.0	1.9	3.0	5.1	7.0	7.4	5.9
2001.....	0.1	0.2	0.5	0.9	1.9	3.1	4.8	7.2	6.9	5.5
2000.....	0.1	0.2	0.5	1.1	1.9	3.3	5.2	6.6	7.2	5.7
1999.....	*	0.2	0.5	1.2	2.0	3.4	5.1	6.6	7.1	4.8
All races 3/, male										
2002.....	*	*	0.6	1.6	3.4	5.6	9.8	14.0	15.6	15.9
2001.....	*	0.3	0.8	1.5	3.2	5.6	9.1	14.1	14.7	14.5
2000.....	*	0.2	0.7	1.9	3.4	5.9	9.8	13.4	15.5	15.5
1999.....	*	0.4	0.8	1.9	3.3	6.1	9.6	12.9	15.8	12.7
All races 3/, female										
2002.....	*	*	*	0.4	0.5	0.7	1.3	2.0	2.3	1.6
2001.....	*	*	0.3	0.4	0.7	0.9	1.4	2.4	2.1	1.7
2000.....	*	*	0.2	0.4	0.6	1.1	1.5	1.9	2.3	1.8
1999.....	*	*	0.3	0.5	0.7	1.0	1.6	2.2	2.1	1.6
White, both sexes										
2002.....	0.1	0.2	0.4	1.1	2.1	3.3	5.5	7.5	7.9	6.4
2001.....	*	0.2	0.5	1.0	2.2	3.5	5.3	7.9	7.3	5.9
2000.....	*	0.2	0.5	1.2	2.1	3.5	5.6	7.1	7.8	6.1
1999.....	*	0.3	0.5	1.3	2.1	3.7	5.5	7.0	7.5	5.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of other mesothelial and soft tissue (C47-C49)										
All races 3/, both sexes										
2002.....	0.8	1.1	1.6	2.4	3.6	4.4	5.8	6.8	8.8	9.3
2001.....	0.8	1.1	1.8	2.4	3.4	4.6	5.8	7.4	8.7	9.1
2000.....	0.8	1.2	1.7	2.6	3.4	4.8	5.9	7.7	8.8	9.6
1999.....	0.9	1.2	1.8	2.3	3.6	4.3	5.9	7.4	8.6	9.6
All races 3/, male										
2002.....	0.9	1.1	1.4	2.1	3.3	3.8	5.7	6.7	9.3	11.7
2001.....	0.9	1.1	1.8	2.1	3.1	4.4	6.1	7.1	10.0	10.7
2000.....	0.8	1.3	1.5	2.3	3.3	4.9	5.2	8.4	10.4	12.5
1999.....	1.1	1.0	1.7	2.4	3.6	4.4	5.9	7.8	9.6	11.3
All races 3/, female										
2002.....	0.8	1.0	1.7	2.7	3.8	4.9	5.8	6.8	8.6	8.3
2001.....	0.8	1.1	1.8	2.8	3.6	4.8	5.4	7.6	8.0	8.4
2000.....	0.8	1.1	1.9	2.8	3.5	4.7	6.5	7.1	7.8	8.4
1999.....	0.7	1.3	1.8	2.2	3.7	4.3	5.8	7.1	8.0	8.9
White, both sexes										
2002.....	0.8	1.0	1.6	2.4	3.6	4.4	5.9	7.1	9.0	9.6
2001.....	0.8	1.1	1.7	2.4	3.4	4.8	5.8	7.4	9.1	9.1
2000.....	0.8	1.2	1.7	2.6	3.4	4.8	6.1	7.9	9.1	9.7
1999.....	0.8	1.2	1.8	2.3	3.6	4.4	5.9	7.4	8.7	9.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of other mesothelial and soft tissue (C47-C49)										
White, male										
2002.....	0.8	1.1	1.5	2.1	3.3	3.8	5.9	7.1	9.6	12.0
2001.....	0.8	1.0	1.8	2.0	3.2	4.5	6.4	7.1	10.4	10.6
2000.....	0.8	1.3	1.6	2.4	3.2	4.9	5.4	8.6	10.8	12.4
1999.....	1.1	1.0	1.8	2.4	3.6	4.3	5.9	7.9	9.8	12.0
White, female										
2002.....	0.7	1.0	1.8	2.6	3.8	4.9	5.9	7.1	8.6	8.5
2001.....	0.7	1.1	1.7	2.8	3.7	5.1	5.4	7.7	8.3	8.4
2000.....	0.8	1.1	1.7	2.8	3.6	4.7	6.6	7.4	8.2	8.5
1999.....	0.5	1.3	1.7	2.1	3.6	4.4	6.0	7.1	8.0	8.9
Black, both sexes										
2002.....	1.2	1.6	1.7	3.2	4.1	5.3	5.5	5.7	8.9	5.9
2001.....	1.2	1.4	2.5	3.5	3.0	3.8	6.1	7.3	6.0	9.8
2000.....	1.2	1.6	2.4	2.9	3.7	5.2	5.9	6.0	*	9.1
1999.....	1.3	1.3	1.9	2.4	4.8	4.0	6.6	8.0	7.5	8.3
Black, male										
2002.....	1.5	2.1	*	*	*	*	*	*	*	*
2001.....	*	*	2.3	3.4	*	*	*	*	*	*
2000.....	*	*	*	*	4.1	*	*	*	*	*
1999.....	1.5	*	*	*	4.5	*	7.1	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of other mesothelial and soft tissue (C47-C49)										
Black, female										
2002.....	*	*	2.0	3.4	4.3	6.3	5.4	6.3	9.5	*
2001.....	*	1.7	2.7	3.5	3.5	3.8	6.7	7.2	*	8.8
2000.....	*	1.7	3.7	3.4	3.4	5.6	6.7	*	*	*
1999.....	*	*	2.2	*	5.0	*	6.3	8.0	*	9.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)										
All races 3/, both sexes										
2002.....	7.9	12.9	19.7	26.2	33.7	41.3	52.4	67.1	89.8	134.6
2001.....	8.2	13.4	20.6	27.1	34.1	41.7	54.0	69.1	88.0	134.2
2000.....	8.3	13.6	20.8	28.1	35.0	43.4	55.2	70.9	90.9	147.4
1999.....	7.9	13.3	21.0	28.1	34.8	43.5	55.9	70.0	92.4	144.4
All races 3/, male										
2002.....	*	*	0.3	0.4	0.7	0.8	1.3	1.9	2.7	3.4
2001.....	*	*	0.4	0.5	0.8	1.2	1.7	2.3	2.4	3.5
2000.....	*	*	0.3	0.7	0.8	1.1	1.7	2.1	2.4	4.2
1999.....	*	*	0.4	0.6	0.7	1.1	1.4	1.9	2.3	4.5
All races 3/, female										
2002.....	15.6	25.3	38.3	50.4	63.7	76.2	93.2	113.2	142.7	191.5
2001.....	16.2	26.3	40.1	52.0	64.3	76.5	95.6	115.9	139.4	188.9
2000.....	16.4	26.6	40.4	53.8	66.0	79.6	97.3	118.8	143.1	205.7
1999.....	15.6	26.1	40.8	53.9	65.6	79.7	98.4	117.4	145.1	200.8
White, both sexes										
2002.....	7.0	11.7	18.5	25.0	33.1	40.8	52.0	67.5	89.0	135.3
2001.....	7.3	12.4	19.4	25.8	33.3	40.9	53.6	69.4	88.2	134.8
2000.....	7.5	12.4	19.7	26.9	34.4	43.3	55.5	71.1	91.7	147.3
1999.....	7.1	12.1	19.8	26.8	33.8	42.8	55.5	69.7	92.8	146.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)										
Black, female										
2002.....	28.7	42.4	58.7	73.1	81.2	92.9	111.1	123.2	176.7	209.1
2001.....	27.5	41.9	60.5	76.7	83.7	99.5	115.0	130.9	157.8	205.9
2000.....	27.7	44.3	60.4	78.8	83.3	91.0	107.4	133.6	149.0	238.7
1999.....	27.5	44.9	63.3	79.6	85.2	99.5	114.1	139.3	154.7	199.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of female genital organs (C51-C58)										
All races 3/, both sexes										
2002.....	3.7	6.4	10.3	15.1	22.5	30.3	39.9	52.0	63.6	77.5
2001.....	3.8	6.4	10.6	14.7	22.8	30.1	39.6	52.1	66.9	79.9
2000.....	3.8	6.5	10.4	15.5	22.2	31.0	38.9	52.5	65.4	82.3
1999.....	3.9	6.6	10.3	15.1	22.7	30.8	39.4	51.7	64.6	79.1
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	7.3	12.6	20.2	29.2	42.9	56.4	71.8	88.8	102.3	111.2
2001.....	7.5	12.6	20.7	28.4	43.4	56.0	71.0	88.6	107.0	113.4
2000.....	7.5	12.7	20.3	30.1	42.4	57.5	69.6	89.1	103.9	115.8
1999.....	7.8	13.0	20.3	29.3	43.2	57.2	70.3	87.6	102.4	110.9
White, both sexes										
2002.....	3.6	6.2	10.3	14.9	22.1	29.2	38.9	51.3	63.3	76.9
2001.....	3.7	6.3	10.4	14.5	21.9	29.0	38.9	51.6	66.9	79.7
2000.....	3.6	6.4	10.2	15.4	21.7	29.8	37.7	52.0	65.0	82.1
1999.....	3.8	6.7	10.3	14.9	22.2	29.7	38.3	50.7	64.5	78.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of female genital organs (C51-C58)										
Black, female										
2002.....	9.2	14.9	22.4	34.3	52.3	76.6	95.7	106.8	117.8	136.0
2001.....	9.8	13.1	23.0	32.6	56.8	77.5	91.7	107.1	117.7	130.1
2000.....	10.5	13.7	23.0	33.7	54.1	81.6	96.3	110.5	119.7	132.6
1999.....	9.6	12.0	21.9	36.4	55.9	80.0	101.3	113.3	110.6	125.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of vulva and vagina (C51-C52)										
All races 3/, both sexes										
2002.....	0.1	0.1	0.3	0.3	0.6	0.7	1.5	2.5	3.5	7.7
2001.....	0.1	0.2	0.3	0.3	0.5	0.9	1.4	2.5	3.9	7.3
2000.....	0.1	0.1	0.3	0.3	0.6	1.0	1.5	2.5	3.6	7.7
1999.....	0.1	0.1	0.3	0.3	0.6	0.9	1.5	2.5	3.9	8.6
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	0.2	0.3	0.5	0.7	1.2	1.3	2.7	4.3	5.7	11.0
2001.....	0.2	0.3	0.6	0.6	1.0	1.6	2.6	4.2	6.3	10.3
2000.....	0.2	0.3	0.6	0.6	1.1	1.9	2.8	4.3	5.7	10.9
1999.....	0.2	0.3	0.6	0.7	1.2	1.6	2.6	4.2	6.1	12.0
White, both sexes										
2002.....	0.1	0.1	0.3	0.4	0.7	0.7	1.5	2.6	3.6	8.0
2001.....	*	0.1	0.3	0.3	0.4	0.9	1.4	2.5	4.1	7.4
2000.....	*	0.1	0.3	0.3	0.6	1.1	1.6	2.5	3.7	7.9
1999.....	*	0.1	0.3	0.3	0.6	0.9	1.5	2.5	3.9	9.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of cervix uteri (C53)										
All races 3/, both sexes										
2002.....	1.7	2.1	2.4	2.7	2.8	3.3	3.6	4.1	4.2	6.2
2001.....	1.8	2.3	2.6	2.6	3.2	2.9	3.5	4.0	5.5	7.2
2000.....	1.8	2.2	2.7	3.1	3.1	3.6	3.9	4.3	5.6	7.3
1999.....	1.8	2.4	2.7	3.0	3.5	3.7	3.9	4.7	4.6	7.9
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	3.4	4.1	4.7	5.2	5.3	6.1	6.5	7.0	6.7	8.9
2001.....	3.7	4.5	5.1	5.1	6.0	5.4	6.3	6.8	8.9	10.1
2000.....	3.5	4.4	5.2	6.0	5.9	6.7	6.9	7.3	9.0	10.3
1999.....	3.6	4.6	5.2	5.8	6.7	6.8	6.9	8.0	7.2	11.1
White, both sexes										
2002.....	1.6	1.8	2.2	2.5	2.4	2.7	3.0	3.6	3.5	5.1
2001.....	1.7	2.2	2.3	2.3	2.9	2.4	3.0	3.4	4.7	6.2
2000.....	1.6	2.0	2.4	2.7	2.6	2.9	3.3	3.8	5.0	6.3
1999.....	1.7	2.3	2.5	2.7	3.0	3.1	3.1	4.0	4.1	6.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of cervix uteri (C53)										
Black, female										
2002.....	5.6	7.0	8.4	7.6	10.7	14.2	16.8	15.4	19.1	24.2
2001.....	5.6	6.1	8.2	8.3	10.6	13.7	12.8	15.5	22.9	26.1
2000.....	5.9	7.0	9.1	11.2	11.4	16.3	16.2	17.4	20.4	27.0
1999.....	5.7	5.5	7.5	11.1	14.8	15.2	21.8	19.7	14.3	29.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of corpus uteri and uterus, part unspecified (C54-C55)										
All races 3/, both sexes										
2002.....	0.4	0.9	2.0	3.5	5.8	8.4	11.8	14.2	17.0	24.6
2001.....	0.4	0.9	2.0	3.2	5.8	8.2	11.4	14.5	19.2	24.7
2000.....	0.4	0.9	1.8	3.5	6.0	8.4	11.1	14.0	18.6	24.7
1999.....	0.5	0.9	1.7	3.3	5.5	8.8	10.9	14.4	19.4	24.0
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	0.9	1.7	3.9	6.9	11.0	15.7	21.2	24.2	27.4	35.2
2001.....	0.8	1.8	4.0	6.2	11.1	15.3	20.4	24.6	30.7	35.0
2000.....	0.9	1.9	3.5	6.9	11.4	15.6	19.8	23.8	29.5	34.8
1999.....	0.9	1.8	3.3	6.3	10.5	16.2	19.5	24.5	30.7	33.7
White, both sexes										
2002.....	0.4	0.8	1.9	3.2	5.2	7.2	10.8	13.1	16.3	23.8
2001.....	0.4	0.9	1.9	3.1	5.2	7.2	10.5	13.7	18.6	24.5
2000.....	0.4	0.9	1.8	3.4	5.4	7.3	9.9	13.0	17.5	24.4
1999.....	0.4	0.9	1.6	3.0	5.2	7.6	10.2	13.7	18.8	23.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of corpus uteri and uterus, part unspecified (C54-C55)

Black, female

2002.....	*	2.0	5.4	12.1	19.2	34.8	42.1	44.5	44.2	53.8
2001.....	*	2.6	5.4	8.3	20.8	32.7	40.4	42.4	47.0	44.2
2000.....	*	1.9	4.4	9.6	21.4	34.0	44.0	44.4	52.6	45.5
1999.....	1.6	*	4.1	10.5	18.8	36.4	36.2	41.2	44.5	48.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of ovary (C56)										
All races 3/, both sexes										
2002.....	1.4	3.2	5.5	8.3	12.9	17.3	22.3	30.3	37.8	38.1
2001.....	1.4	2.9	5.5	8.3	13.0	17.6	22.6	30.2	37.4	39.6
2000.....	1.5	3.1	5.5	8.3	12.2	17.5	21.8	30.9	36.5	40.9
1999.....	1.5	3.1	5.5	8.3	12.6	17.0	22.4	29.2	35.5	37.6
All races 3/, male										
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2002.....	2.8	6.4	10.7	16.2	24.6	32.3	40.1	51.7	60.8	54.6
2001.....	2.8	5.8	10.8	16.0	24.7	32.8	40.5	51.4	59.8	56.1
2000.....	2.9	6.0	10.8	16.1	23.3	32.5	39.0	52.5	58.0	57.6
1999.....	3.0	6.2	10.8	16.0	24.0	31.6	39.9	49.5	56.2	52.8
White, both sexes										
2002.....	1.4	3.3	5.8	8.6	13.5	18.0	22.9	31.1	38.8	39.0
2001.....	1.4	3.1	5.7	8.5	13.2	18.1	23.3	31.1	38.6	40.4
2000.....	1.5	3.2	5.6	8.7	12.7	18.0	22.4	32.0	37.6	42.0
1999.....	1.6	3.3	5.7	8.6	13.0	17.5	22.9	29.6	36.4	38.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of ovary (C56)										
Black, female										
2002.....	2.3	5.3	7.7	13.9	19.5	25.9	33.4	42.3	49.3	48.1
2001.....	2.7	3.6	8.1	14.4	22.6	28.3	34.5	41.9	42.1	50.1
2000.....	2.9	4.0	8.8	11.6	19.4	29.2	32.6	41.9	41.2	47.2
1999.....	2.1	4.5	9.3	13.2	20.3	26.2	37.8	46.3	42.3	39.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other and unspecified female genital organs (C57)

All races 3/, both sexes

2002.....	*	*	0.2	0.2	0.4	0.5	0.6	1.0	1.1	1.0
2001.....	*	*	0.1	0.3	0.3	0.4	0.7	0.9	0.9	1.2
2000.....	*	*	0.1	0.2	0.4	0.5	0.6	0.7	1.1	1.6
1999.....	*	*	0.2	0.3	0.4	0.5	0.7	0.8	1.3	0.9

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	*	*	0.3	0.4	0.8	1.0	1.2	1.6	1.7	1.5
2001.....	*	*	0.3	0.5	0.6	0.8	1.2	1.5	1.4	1.7
2000.....	*	*	0.2	0.5	0.8	0.8	1.1	1.3	1.7	2.2
1999.....	*	*	0.4	0.5	0.8	0.9	1.3	1.4	2.1	1.3

White, both sexes

2002.....	*	*	0.2	0.2	0.3	0.5	0.7	1.0	1.1	1.0
2001.....	*	*	0.1	0.3	0.3	0.4	0.7	0.9	0.9	1.3
2000.....	*	*	*	0.2	0.4	0.5	0.6	0.6	1.1	1.6
1999.....	*	*	0.2	0.3	0.4	0.5	0.7	0.8	1.2	0.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of male genital organs (C60-C63)										
All races 3/, both sexes										
2002.....	0.4	0.6	1.8	4.9	11.8	24.6	46.4	80.3	129.4	195.9
2001.....	0.4	0.7	1.9	4.8	11.6	25.3	47.9	83.0	133.8	201.9
2000.....	0.3	0.7	1.7	4.8	12.4	26.3	49.4	85.7	137.7	210.1
1999.....	0.4	0.8	1.8	5.1	12.6	28.9	52.3	89.5	142.3	208.7
All races 3/, male										
2002.....	0.8	1.3	3.7	10.1	24.7	53.1	104.4	193.9	342.2	647.3
2001.....	0.7	1.5	3.9	9.9	24.4	54.8	108.2	201.5	357.2	684.2
2000.....	0.7	1.5	3.5	10.0	26.2	56.9	112.1	208.7	371.2	726.1
1999.....	0.9	1.5	3.7	10.6	26.4	62.6	119.3	218.2	385.6	726.5
All races 3/, female										
2002.....
2001.....
2000.....
1999.....
White, both sexes										
2002.....	0.4	0.5	1.5	4.2	10.0	21.6	41.3	73.2	121.0	187.6
2001.....	0.4	0.7	1.6	4.1	9.9	22.5	42.7	75.8	123.9	190.3
2000.....	0.3	0.6	1.5	4.1	10.8	22.8	43.6	78.1	127.0	199.5
1999.....	0.4	0.7	1.5	4.5	10.4	25.3	46.8	81.6	133.9	196.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of male genital organs (C60-C63)										
White, male										
2002.....	0.8	1.1	3.1	8.6	20.8	46.2	91.9	175.4	318.4	619.3
2001.....	0.8	1.4	3.2	8.5	20.7	48.1	95.5	182.8	328.9	645.0
2000.....	0.6	1.3	3.1	8.4	22.4	48.9	98.0	189.2	340.4	690.4
1999.....	0.9	1.4	3.1	9.2	21.7	54.4	105.7	197.7	360.7	685.1
White, female										
2002.....
2001.....
2000.....
1999.....
Black, both sexes										
2002.....	*	1.5	4.4	11.9	29.8	58.2	111.3	177.3	262.7	328.7
2001.....	*	1.4	5.1	12.1	29.0	58.1	115.3	181.7	281.4	363.3
2000.....	*	1.7	3.7	12.9	29.7	65.1	120.7	191.2	294.2	357.9
1999.....	*	1.4	4.4	12.1	34.1	69.0	124.1	197.3	273.6	367.6
Black, male										
2002.....	*	3.2	9.7	26.5	67.9	136.6	277.1	474.0	767.3	1,186.8
2001.....	*	3.1	11.0	26.7	65.9	136.7	288.5	486.0	833.9	1,342.3
2000.....	*	3.6	8.1	28.7	67.3	153.3	302.6	508.4	878.4	1,319.3
1999.....	*	3.0	9.6	27.0	77.6	162.0	311.0	524.8	825.7	1,351.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of prostate (C61)										
All races 3/, both sexes										
2002.....	0.1	0.4	1.6	4.6	11.4	24.2	45.8	79.7	128.7	195.0
2001.....	0.1	0.5	1.7	4.6	11.3	25.0	47.4	82.4	133.2	200.9
2000.....	*	0.5	1.5	4.6	12.1	25.9	49.0	85.1	136.6	209.3
1999.....	0.1	0.5	1.6	4.9	12.3	28.4	51.9	89.0	141.4	207.5
All races 3/, male										
2002.....	0.2	0.8	3.3	9.6	24.0	52.2	103.2	192.6	340.3	644.5
2001.....	0.2	1.0	3.5	9.4	23.8	53.9	107.1	200.1	355.4	681.2
2000.....	*	1.1	3.1	9.6	25.4	56.0	111.3	207.4	368.2	723.1
1999.....	0.3	1.0	3.3	10.1	25.8	61.6	118.3	216.9	383.0	722.4
All races 3/, female										
2002.....
2001.....
2000.....
1999.....
White, both sexes										
2002.....	*	0.3	1.3	3.9	9.6	21.2	40.7	72.6	120.3	186.7
2001.....	*	0.4	1.4	3.9	9.7	22.1	42.2	75.3	123.3	189.4
2000.....	*	0.4	1.3	3.8	10.4	22.4	43.2	77.6	125.8	198.6
1999.....	0.1	0.4	1.4	4.2	10.2	24.9	46.4	81.1	132.9	195.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of prostate (C61)										
White, male										
2002.....	*	0.5	2.7	8.1	20.1	45.2	90.7	174.1	316.4	616.3
2001.....	*	0.8	2.8	8.0	20.2	47.3	94.3	181.5	327.2	641.8
2000.....	*	0.8	2.6	7.9	21.6	48.0	97.0	188.0	337.4	687.5
1999.....	0.2	0.8	2.8	8.6	21.2	53.4	104.7	196.5	358.0	681.0
White, female										
2002.....
2001.....
2000.....
1999.....
Black, both sexes										
2002.....	*	1.2	4.2	11.7	29.4	57.9	110.6	176.8	261.9	328.4
2001.....	*	1.1	4.8	11.8	28.7	57.4	114.7	180.8	280.3	362.9
2000.....	*	1.5	3.6	12.9	29.5	64.7	120.7	190.1	292.8	356.6
1999.....	*	1.4	4.2	12.0	33.4	68.7	123.4	196.6	273.0	366.4
Black, male										
2002.....	*	2.6	9.1	25.9	66.9	135.9	275.4	472.6	765.0	1,185.7
2001.....	*	2.4	10.4	26.1	65.1	135.2	286.8	483.7	830.7	1,341.2
2000.....	*	3.2	7.8	28.7	66.9	152.5	302.6	505.6	874.2	1,314.7
1999.....	*	2.9	9.2	26.6	76.0	161.5	309.3	522.9	824.0	1,346.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of penis and other and unspecified male genital organs (C60,C63)

All races 3/, both sexes

2002.....	*	*	*	0.1	0.2	0.3	0.4	0.5	0.5	0.6
2001.....	*	*	*	*	0.2	0.3	0.4	0.4	0.6	0.8
2000.....	*	*	*	*	0.2	0.3	0.3	0.4	1.0	0.7
1999.....	*	*	*	*	0.2	0.3	0.3	0.5	0.7	1.1

All races 3/, male

2002.....	*	*	*	0.3	0.5	0.6	0.9	1.2	1.3	2.0
2001.....	*	*	*	*	0.4	0.6	0.8	1.0	1.5	2.8
2000.....	*	*	*	*	0.5	0.7	0.7	1.1	2.7	2.4
1999.....	*	*	*	*	0.4	0.7	0.7	1.2	1.8	3.7

All races 3/, female

2002.....
2001.....
2000.....
1999.....

White, both sexes

2002.....	*	*	*	*	*	0.3	0.4	0.5	0.5	0.7
2001.....	*	*	*	*	*	0.3	0.4	0.4	0.5	0.9
2000.....	*	*	*	*	0.3	0.4	0.3	0.4	1.0	0.7
1999.....	*	*	*	*	*	0.3	0.3	0.5	0.7	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of urinary tract (C64-C68)										
All races 3/, both sexes										
2002.....	1.4	3.3	6.1	11.2	17.6	28.7	41.0	58.2	74.7	102.0
2001.....	1.7	3.2	6.3	10.7	19.3	27.3	40.9	55.2	76.1	103.9
2000.....	1.5	3.1	6.2	11.4	17.9	27.5	40.1	55.2	77.4	106.0
1999.....	1.4	2.9	6.3	11.3	18.2	26.9	40.6	54.4	73.1	103.0
All races 3/, male										
2002.....	1.9	4.7	9.0	16.8	26.8	42.7	61.3	92.6	119.6	183.6
2001.....	2.1	4.7	9.2	15.9	28.9	41.4	62.9	87.5	126.4	192.4
2000.....	2.0	4.3	9.2	17.1	26.7	41.5	61.0	87.4	129.7	198.9
1999.....	1.9	4.3	9.2	16.9	27.2	40.6	61.8	88.2	121.7	188.4
All races 3/, female										
2002.....	0.9	2.0	3.4	6.0	9.4	16.5	24.8	33.9	47.4	66.6
2001.....	1.2	1.8	3.4	5.8	10.5	15.2	23.3	32.7	46.0	66.9
2000.....	1.0	1.9	3.3	6.2	10.0	15.4	23.7	32.7	46.6	68.1
1999.....	0.9	1.6	3.5	6.1	10.0	15.1	23.9	30.8	44.6	68.6
White, both sexes										
2002.....	1.4	3.4	6.3	11.5	18.3	29.8	42.4	59.6	76.4	103.9
2001.....	1.6	3.3	6.3	10.9	19.7	28.6	42.2	56.6	77.4	106.4
2000.....	1.5	3.2	6.4	11.8	18.7	28.4	41.1	56.2	78.9	109.2
1999.....	1.4	2.8	6.4	11.5	18.8	27.6	41.4	55.5	74.7	104.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of urinary tract (C64-C68)										
White, male										
2002.....	1.9	4.9	9.3	17.4	27.8	44.6	63.7	95.5	122.7	190.7
2001.....	2.1	4.9	9.3	16.4	29.5	43.5	65.2	89.8	129.9	200.5
2000.....	2.1	4.4	9.4	17.5	28.0	42.9	62.8	89.8	132.5	206.8
1999.....	1.9	4.2	9.1	17.2	28.2	41.8	63.7	90.7	125.5	194.9
White, female										
2002.....	0.9	1.9	3.3	5.9	9.5	16.8	25.0	33.9	48.1	66.2
2001.....	1.1	1.7	3.5	5.7	10.6	15.5	23.6	33.0	45.7	67.0
2000.....	1.0	1.9	3.5	6.4	10.1	15.8	23.7	32.5	47.0	69.6
1999.....	0.9	1.5	3.7	6.1	10.1	15.2	23.8	30.8	44.7	68.4
Black, both sexes										
2002.....	1.3	3.5	6.4	11.7	16.5	25.7	36.1	54.1	64.9	93.2
2001.....	2.4	3.5	6.9	11.1	19.8	22.5	34.5	50.4	71.2	86.6
2000.....	1.5	3.4	6.0	10.9	15.9	25.1	37.5	53.5	74.0	80.4
1999.....	1.7	4.1	7.4	12.8	17.0	26.0	39.2	49.1	61.0	93.1
Black, male										
2002.....	1.7	4.4	8.3	16.5	24.5	34.4	47.4	77.2	98.6	135.1
2001.....	2.7	4.4	10.7	14.9	30.3	31.0	50.2	76.2	101.9	127.1
2000.....	1.7	4.2	9.1	16.3	22.6	38.0	53.5	76.5	122.2	140.7
1999.....	2.3	5.8	11.9	18.6	24.0	36.1	52.2	73.1	89.8	138.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of urinary tract (C64-C68)										
Black, female										
2002.....	*	2.8	4.8	7.8	10.2	19.2	28.5	40.3	47.3	77.2
2001.....	2.2	2.7	3.7	8.1	11.5	16.1	24.0	34.9	55.6	71.6
2000.....	*	2.7	3.5	6.5	10.6	15.6	26.9	39.6	49.7	57.9
1999.....	*	2.5	3.6	8.0	11.5	18.5	30.6	34.6	46.7	76.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of kidney and renal pelvis (C64-C65)										
All races 3/, both sexes										
2002.....	0.9	2.4	4.3	7.2	10.9	16.0	19.6	24.6	28.7	32.5
2001.....	1.2	2.4	4.4	7.1	11.9	14.6	20.8	24.4	29.1	32.3
2000.....	1.1	2.2	4.5	7.6	10.8	15.0	19.4	24.5	29.6	33.8
1999.....	1.1	2.2	4.3	7.3	10.8	14.4	19.2	23.1	26.9	31.4
All races 3/, male										
2002.....	1.3	3.4	6.4	10.6	16.1	23.0	27.6	36.6	39.9	50.1
2001.....	1.6	3.5	6.5	10.5	17.2	21.0	30.0	34.4	42.2	52.9
2000.....	1.6	3.1	6.6	11.2	15.6	21.4	27.4	34.6	44.2	55.1
1999.....	1.5	3.2	6.1	10.6	15.8	20.3	27.3	33.7	38.9	49.4
All races 3/, female										
2002.....	0.5	1.5	2.3	3.9	6.2	10.0	13.2	16.2	21.9	24.9
2001.....	0.9	1.4	2.4	3.9	7.0	9.1	13.5	17.4	21.2	23.7
2000.....	0.7	1.4	2.4	4.3	6.5	9.5	13.0	17.5	21.0	25.1
1999.....	0.6	1.2	2.5	4.3	6.2	9.3	12.8	15.7	19.9	24.1
White, both sexes										
2002.....	1.0	2.4	4.4	7.4	11.2	16.6	20.2	25.0	29.2	33.2
2001.....	1.2	2.5	4.4	7.2	12.1	15.3	21.3	24.7	29.5	32.8
2000.....	1.1	2.3	4.6	7.8	11.1	15.4	19.7	24.9	30.0	34.5
1999.....	1.1	2.1	4.3	7.5	11.2	14.8	19.4	23.4	27.3	31.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of kidney and renal pelvis (C64-C65)										
White, male										
2002.....	1.4	3.5	6.6	10.9	16.4	23.8	28.3	37.5	40.2	51.9
2001.....	1.6	3.6	6.5	10.8	17.2	21.9	30.6	34.4	42.8	54.4
2000.....	1.6	3.2	6.7	11.3	16.1	21.5	27.7	35.1	44.5	56.6
1999.....	1.5	3.1	6.0	10.7	16.4	20.6	27.6	34.1	40.1	49.8
White, female										
2002.....	0.5	1.4	2.2	4.1	6.4	10.3	13.6	16.1	22.5	25.1
2001.....	0.8	1.3	2.4	3.8	7.4	9.5	13.8	17.8	21.4	23.8
2000.....	0.7	1.4	2.5	4.5	6.5	10.0	13.3	17.7	21.4	25.5
1999.....	0.6	1.1	2.6	4.4	6.3	9.7	13.0	15.8	19.8	24.2
Black, both sexes										
2002.....	0.9	2.8	4.4	7.0	10.6	14.3	16.8	25.8	26.7	28.2
2001.....	1.8	2.6	5.0	7.4	12.2	11.7	19.2	24.0	28.2	29.8
2000.....	1.1	2.5	4.7	7.4	10.4	14.8	18.5	23.5	29.1	29.7
1999.....	1.2	2.9	4.9	7.8	9.7	13.5	19.3	23.0	24.3	31.2
Black, male										
2002.....	*	3.5	6.4	11.8	16.8	19.9	23.7	34.4	39.0	39.7
2001.....	1.9	3.5	7.8	10.5	20.5	17.0	29.3	39.0	42.5	46.5
2000.....	*	3.2	7.1	11.4	15.3	25.0	28.1	32.1	48.9	49.6
1999.....	1.8	4.1	8.0	12.5	14.0	20.0	28.2	34.2	27.9	49.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of kidney and renal pelvis (C64-C65)										
Black, female										
2002.....	*	2.3	2.6	3.1	5.7	10.2	12.1	20.6	20.3	23.8
2001.....	1.7	1.8	2.7	4.9	5.6	7.7	12.6	15.0	20.8	23.6
2000.....	*	1.8	2.6	4.2	6.5	7.3	12.2	18.2	19.1	22.3
1999.....	*	1.9	2.3	3.9	6.3	8.7	13.5	16.3	22.5	24.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of bladder (C67)										
All races 3/, both sexes										
2002.....	0.4	0.9	1.7	3.8	6.4	12.0	20.2	31.9	44.1	66.6
2001.....	0.4	0.8	1.7	3.4	7.0	12.0	19.0	29.3	44.8	68.9
2000.....	0.3	0.8	1.6	3.6	6.8	11.7	19.6	29.1	45.5	69.6
1999.....	0.3	0.7	1.9	3.8	7.0	11.8	20.2	29.7	43.8	68.5
All races 3/, male										
2002.....	0.5	1.2	2.4	5.9	10.1	19.0	32.2	54.0	76.6	128.9
2001.....	0.5	1.2	2.6	5.2	11.3	19.5	31.6	50.8	81.0	134.8
2000.....	0.4	1.1	2.5	5.7	10.6	19.0	32.2	50.6	82.4	139.4
1999.....	0.4	1.1	2.9	5.9	11.0	19.3	32.9	52.3	79.5	134.8
All races 3/, female										
2002.....	0.3	0.5	1.0	1.9	2.9	6.0	10.6	16.3	24.2	39.6
2001.....	0.3	0.4	0.9	1.7	3.2	5.6	9.0	14.3	23.2	41.4
2000.....	0.3	0.5	0.8	1.7	3.4	5.5	9.7	14.0	23.8	41.2
1999.....	0.2	0.3	0.9	1.7	3.4	5.4	10.3	13.9	22.9	41.8
White, both sexes										
2002.....	0.4	0.9	1.8	3.9	6.7	12.5	20.9	32.9	45.2	67.7
2001.....	0.4	0.8	1.8	3.5	7.3	12.6	19.8	30.3	45.7	70.8
2000.....	0.3	0.8	1.7	3.8	7.2	12.3	20.2	29.7	46.4	72.0
1999.....	0.3	0.7	2.0	3.8	7.2	12.1	20.8	30.5	45.0	69.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of bladder (C67)										
White, male										
2002.....	0.5	1.3	2.6	6.2	10.8	20.1	33.7	56.0	79.4	134.1
2001.....	0.5	1.2	2.7	5.4	12.0	20.7	33.1	53.1	83.7	141.1
2000.....	0.4	1.2	2.7	6.0	11.4	20.2	33.6	52.7	84.6	145.7
1999.....	0.4	1.1	3.0	6.2	11.4	20.1	34.4	54.3	82.1	140.7
White, female										
2002.....	0.3	0.5	1.0	1.7	2.9	5.9	10.4	16.4	24.3	38.9
2001.....	0.3	0.3	0.9	1.7	3.0	5.5	9.0	14.2	22.7	41.3
2000.....	0.3	0.5	0.8	1.8	3.4	5.5	9.6	13.6	23.7	42.1
1999.....	0.2	0.3	1.0	1.5	3.4	5.1	10.0	13.8	23.2	41.4
Black, both sexes										
2002.....	*	*	1.6	4.0	5.2	10.9	18.6	27.0	37.2	63.3
2001.....	*	0.9	1.7	3.4	6.9	10.0	14.1	24.8	41.2	55.0
2000.....	*	0.8	1.3	3.0	5.4	9.5	17.9	28.4	44.6	49.7
1999.....	*	1.0	2.2	4.5	6.9	11.8	18.9	25.4	35.3	60.3
Black, male										
2002.....	*	*	*	4.5	7.1	14.2	23.7	41.8	57.3	91.1
2001.....	*	*	2.8	4.2	9.2	12.7	20.5	34.4	57.8	78.3
2000.....	*	*	*	4.6	7.2	12.5	24.1	41.1	73.3	88.8
1999.....	*	*	3.9	5.4	9.8	15.6	23.7	38.9	59.3	86.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of bladder (C67)										
Black, female										
2002.....	*	*	*	3.7	3.7	8.5	15.2	18.1	26.7	52.6
2001.....	*	*	*	2.7	5.1	8.1	9.9	19.1	32.7	46.3
2000.....	*	*	*	*	3.9	7.3	13.8	20.8	30.1	35.2
1999.....	*	*	*	3.7	4.7	9.0	15.8	17.2	23.3	50.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other and unspecified sites within the urinary tract (C66,C68)

All races 3/, both sexes

2002.....	*	*	0.1	0.2	0.4	0.7	1.2	1.6	1.9	2.9
2001.....	*	*	0.1	0.2	0.4	0.7	1.1	1.5	2.2	2.7
2000.....	*	*	*	0.2	0.3	0.7	1.2	1.6	2.3	2.6
1999.....	*	*	0.1	0.2	0.4	0.7	1.1	1.6	2.3	3.1

All races 3/, male

2002.....	*	*	*	0.3	0.6	0.7	1.5	2.0	3.1	4.7
2001.....	*	*	*	*	0.4	0.9	1.3	2.2	3.3	4.7
2000.....	*	*	*	*	0.4	1.1	1.5	2.1	3.1	4.4
1999.....	*	*	*	0.3	*	1.0	1.6	2.2	3.3	4.2

All races 3/, female

2002.....	*	*	*	*	*	0.6	1.0	1.4	1.2	2.2
2001.....	*	*	*	*	*	0.4	0.9	1.0	1.6	1.8
2000.....	*	*	*	*	*	*	0.9	1.2	1.8	1.8
1999.....	*	*	*	*	0.4	0.4	0.8	1.2	1.8	2.7

White, both sexes

2002.....	*	*	*	0.2	0.4	0.7	1.3	1.7	2.0	3.0
2001.....	*	*	0.1	*	0.3	0.7	1.1	1.6	2.3	2.8
2000.....	*	*	*	0.2	0.3	0.7	1.2	1.6	2.5	2.7
1999.....	*	*	*	0.2	0.4	0.7	1.2	1.7	2.4	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of eye and adnexa (C69)										
All races 3/, both sexes										
2002.....	*	*	*	0.2	0.2	0.2	0.3	0.4	0.5	0.7
2001.....	*	*	*	*	*	0.3	0.3	0.4	0.6	0.6
2000.....	*	*	*	0.2	0.2	0.3	0.3	0.4	0.5	0.8
1999.....	*	*	*	*	*	0.2	0.3	0.4	0.6	0.8
All races 3/, male										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	0.5	*	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	0.7
2001.....	*	*	*	*	*	*	*	*	*	0.6
2000.....	*	*	*	*	*	*	*	*	*	0.7
1999.....	*	*	*	*	*	*	*	*	0.8	0.8
White, both sexes										
2002.....	*	*	*	0.2	0.2	*	0.3	0.5	0.6	0.8
2001.....	*	*	*	*	*	0.3	0.3	0.4	0.7	0.7
2000.....	*	*	*	*	0.2	0.3	0.4	0.5	0.5	0.8
1999.....	*	*	*	*	*	*	0.4	0.4	0.7	0.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
All races 3/, both sexes										
2002.....	2.8	4.2	6.2	8.5	11.1	14.7	17.8	19.6	18.5	16.4
2001.....	2.8	4.0	6.1	8.3	11.6	14.2	16.7	20.3	20.6	16.4
2000.....	2.9	4.1	6.3	8.2	11.4	14.8	18.3	19.4	21.1	16.9
1999.....	2.8	4.3	6.1	9.2	11.7	15.7	18.3	21.0	19.4	16.0
All races 3/, male										
2002.....	3.5	5.3	7.2	10.6	13.1	18.4	22.2	24.4	22.9	20.1
2001.....	3.5	4.9	7.5	10.7	13.9	17.3	21.2	25.4	26.5	20.1
2000.....	3.7	4.9	7.7	10.1	14.0	17.9	22.8	23.2	26.3	22.4
1999.....	3.4	5.2	7.4	10.6	14.7	19.5	22.3	26.1	24.5	20.7
All races 3/, female										
2002.....	2.1	3.1	5.1	6.5	9.2	11.4	14.2	16.2	15.8	14.8
2001.....	2.1	3.1	4.8	6.0	9.5	11.5	13.1	16.7	17.1	14.8
2000.....	2.0	3.3	4.9	6.5	9.0	12.1	14.8	16.7	18.0	14.7
1999.....	2.3	3.4	4.9	7.9	8.9	12.4	15.1	17.5	16.5	14.1
White, both sexes										
2002.....	3.1	4.7	6.7	9.1	12.0	15.9	19.0	20.7	19.3	17.1
2001.....	3.0	4.4	6.6	8.8	12.6	15.2	17.8	21.6	21.6	17.0
2000.....	3.1	4.5	6.9	8.8	12.3	16.0	19.7	20.5	21.8	17.5
1999.....	3.0	4.7	6.6	10.0	12.5	16.9	19.6	22.2	20.3	16.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
White, male										
2002.....	3.9	5.9	7.9	11.2	14.0	19.9	23.6	25.6	23.7	21.2
2001.....	3.8	5.4	8.1	11.4	14.9	18.6	22.6	26.7	27.8	20.5
2000.....	4.0	5.4	8.3	10.9	15.2	19.4	24.3	24.3	27.5	23.5
1999.....	3.6	5.7	7.9	11.4	15.7	20.9	23.9	27.4	25.8	21.8
White, female										
2002.....	2.3	3.5	5.5	7.1	10.2	12.4	15.3	17.2	16.7	15.4
2001.....	2.3	3.4	5.2	6.4	10.5	12.2	13.9	18.0	17.9	15.6
2000.....	2.1	3.5	5.5	6.9	9.7	13.0	16.0	17.7	18.5	15.0
1999.....	2.5	3.8	5.4	8.7	9.7	13.4	16.2	18.5	17.1	14.5
Black, both sexes										
2002.....	1.3	1.9	3.7	5.7	5.6	8.0	9.5	12.2	11.3	9.5
2001.....	1.6	2.6	3.4	5.4	6.4	9.0	9.4	9.4	10.8	11.7
2000.....	2.0	2.2	3.7	5.4	6.0	7.5	8.8	10.3	14.4	10.6
1999.....	1.8	2.6	4.0	5.3	6.7	8.5	7.8	12.0	11.6	9.9
Black, male										
2002.....	*	2.1	3.3	8.2	7.9	9.5	11.7	15.6	17.6	*
2001.....	1.9	3.0	4.4	6.9	9.2	9.1	11.4	12.5	*	*
2000.....	2.7	2.2	5.7	6.6	7.7	7.8	12.4	11.8	*	*
1999.....	2.6	3.7	5.3	5.9	8.7	10.9	8.8	14.2	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Black, female

2002.....	*	1.8	4.0	3.7	3.7	6.8	8.1	10.2	8.0	10.3
2001.....	1.3	2.3	2.6	4.1	4.3	8.8	8.1	7.5	10.2	9.7
2000.....	1.5	2.2	2.1	4.5	4.7	7.3	6.4	9.4	14.0	10.7
1999.....	*	1.6	2.8	4.8	5.2	6.7	7.2	10.6	12.5	9.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of brain (C71)										
All races 3/, both sexes										
2002.....	2.7	4.2	6.0	8.4	10.9	14.4	17.3	19.2	18.0	16.0
2001.....	2.7	3.9	6.0	8.2	11.4	14.0	16.4	19.8	20.1	16.0
2000.....	2.8	4.0	6.2	8.1	11.2	14.5	18.0	19.1	20.6	16.3
1999.....	2.7	4.2	6.0	9.1	11.4	15.5	17.9	20.5	19.0	15.5
All races 3/, male										
2002.....	3.4	5.3	7.1	10.4	12.9	18.0	21.6	23.9	21.9	19.7
2001.....	3.4	4.8	7.4	10.5	13.7	17.0	20.8	24.9	26.0	19.7
2000.....	3.6	4.8	7.6	10.0	13.9	17.7	22.5	22.8	25.8	21.8
1999.....	3.3	5.2	7.3	10.5	14.4	19.2	21.9	25.5	24.0	20.1
All races 3/, female										
2002.....	2.0	3.1	5.0	6.4	9.1	11.2	13.9	15.9	15.5	14.4
2001.....	2.0	3.1	4.7	6.0	9.3	11.4	12.9	16.3	16.6	14.4
2000.....	2.0	3.2	4.8	6.3	8.8	11.9	14.5	16.5	17.6	14.1
1999.....	2.2	3.3	4.7	7.8	8.7	12.3	14.8	17.1	16.0	13.6
White, both sexes										
2002.....	3.0	4.6	6.6	9.0	11.8	15.6	18.6	20.3	18.8	16.7
2001.....	3.0	4.3	6.6	8.7	12.4	14.9	17.4	21.1	21.2	16.6
2000.....	3.0	4.4	6.8	8.7	12.2	15.7	19.4	20.2	21.4	16.9
1999.....	3.0	4.6	6.5	9.9	12.3	16.6	19.2	21.7	19.8	16.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of brain (C71)										
White, male										
2002.....	3.8	5.8	7.9	11.0	13.9	19.4	23.0	25.1	22.7	20.8
2001.....	3.8	5.3	8.0	11.2	14.6	18.3	22.1	26.1	27.4	20.1
2000.....	3.9	5.3	8.1	10.7	15.0	19.1	24.0	24.1	27.0	23.0
1999.....	3.5	5.6	7.8	11.3	15.4	20.6	23.5	26.8	25.3	21.2
White, female										
2002.....	2.2	3.5	5.4	7.0	10.0	12.2	15.0	16.9	16.4	14.9
2001.....	2.3	3.3	5.2	6.3	10.3	12.0	13.6	17.5	17.4	15.1
2000.....	2.1	3.4	5.4	6.7	9.5	12.8	15.7	17.5	18.1	14.5
1999.....	2.5	3.7	5.2	8.5	9.5	13.2	15.9	18.1	16.6	14.1
Black, both sexes										
2002.....	1.3	1.8	3.6	5.3	5.3	7.5	9.0	11.9	10.5	9.2
2001.....	1.5	2.6	3.3	5.3	6.2	8.9	9.4	9.2	10.6	11.7
2000.....	1.9	2.2	3.6	5.3	5.8	7.2	8.8	9.8	14.1	10.0
1999.....	1.8	2.6	3.7	5.0	6.4	8.5	7.4	11.4	10.7	9.2
Black, male										
2002.....	*	2.0	3.1	7.7	7.5	9.0	10.7	15.2	16.0	*
2001.....	1.8	2.9	4.2	6.7	8.7	9.1	11.4	12.5	*	*
2000.....	2.6	2.2	5.6	6.4	7.5	7.5	12.4	10.9	*	*
1999.....	2.5	3.7	4.9	5.8	8.5	10.9	7.8	13.8	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of brain (C71)										
Black, female										
2002.....	*	1.7	4.0	3.3	3.6	6.5	7.8	9.9	*	10.3
2001.....	*	2.3	2.6	4.1	4.2	8.6	8.1	7.2	9.8	9.7
2000.....	*	2.1	2.0	4.3	4.6	6.9	6.4	9.1	14.0	10.3
1999.....	*	1.6	2.7	4.4	4.8	6.7	7.2	10.0	11.7	8.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other parts of central nervous system (C70,C72)

All races 3/, both sexes

2002.....	*	*	0.1	0.2	0.2	0.3	0.5	0.4	0.5	*
2001.....	*	*	*	*	0.2	0.2	0.4	0.5	0.4	*
2000.....	*	*	0.1	*	*	0.2	0.3	0.3	0.4	0.6
1999.....	*	*	0.2	*	0.2	0.2	0.4	0.5	0.5	*

All races 3/, male

2002.....	*	*	*	*	*	*	0.6	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	0.3	0.5	0.4	0.5	*
2001.....	*	*	*	*	*	0.3	0.4	0.5	0.5	*
2000.....	*	*	*	*	*	0.2	0.3	*	0.4	0.5
1999.....	*	*	0.2	*	*	0.2	0.4	0.5	0.5	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)										
All races 3/, both sexes										
2002.....	0.2	0.5	0.6	1.0	1.4	1.9	2.7	3.9	4.7	5.3
2001.....	0.2	0.5	0.7	1.0	1.7	2.2	3.1	3.8	4.7	5.3
2000.....	0.2	0.4	0.5	1.0	1.6	2.3	2.9	3.9	4.5	6.2
1999.....	0.2	0.4	0.6	1.1	1.6	2.0	2.6	3.9	4.4	5.8
All races 3/, male										
2002.....	0.2	0.5	0.7	1.0	1.6	1.9	2.6	4.1	4.6	4.2
2001.....	0.3	0.5	0.7	1.0	1.8	2.5	3.3	3.4	4.7	4.5
2000.....	0.2	0.5	0.5	1.0	1.8	2.5	3.0	3.9	5.0	5.3
1999.....	0.3	0.4	0.7	1.1	1.7	2.1	3.1	4.0	4.5	5.1
All races 3/, female										
2002.....	0.2	0.4	0.6	0.9	1.3	1.9	2.8	3.8	4.8	5.8
2001.....	0.2	0.5	0.6	0.9	1.6	1.9	3.0	4.0	4.7	5.7
2000.....	0.3	0.3	0.6	1.1	1.5	2.1	2.8	3.8	4.3	6.5
1999.....	0.2	0.4	0.5	1.1	1.6	1.9	2.2	3.8	4.4	6.1
White, both sexes										
2002.....	0.2	0.5	0.6	1.0	1.5	2.0	2.6	4.0	4.7	5.3
2001.....	0.3	0.5	0.7	1.0	1.7	2.3	3.3	3.7	4.5	5.3
2000.....	0.2	0.4	0.5	1.1	1.6	2.2	2.8	3.9	4.6	6.2
1999.....	0.2	0.5	0.6	1.1	1.6	2.0	2.6	3.8	4.5	5.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)										
All races 3/, both sexes										
2002.....	3.2	6.5	11.3	18.8	31.0	44.0	64.4	85.4	112.6	154.5
2001.....	3.3	6.3	11.3	19.9	30.4	44.2	64.4	86.9	112.4	155.2
2000.....	3.4	6.5	11.8	19.6	32.0	46.9	66.7	90.0	119.6	159.0
1999.....	3.9	7.0	12.2	22.1	35.9	52.6	74.2	96.8	128.2	168.4
All races 3/, male										
2002.....	3.3	7.2	14.0	21.9	38.3	54.3	79.5	105.3	138.8	196.9
2001.....	3.5	7.2	13.8	23.5	37.1	52.9	79.1	107.0	138.3	196.5
2000.....	3.6	7.5	13.9	23.3	38.6	56.6	81.7	110.4	148.1	200.1
1999.....	4.0	8.3	14.3	26.5	43.1	62.9	92.2	118.6	158.2	218.9
All races 3/, female										
2002.....	3.2	5.8	8.6	15.9	24.4	35.1	52.3	71.4	96.6	136.0
2001.....	3.0	5.4	8.9	16.4	24.3	36.8	52.8	72.9	96.9	137.9
2000.....	3.1	5.5	9.7	16.2	26.0	38.6	54.8	75.8	102.8	142.3
1999.....	3.8	5.8	10.3	18.0	29.3	43.8	60.1	81.5	110.7	148.1
White, both sexes										
2002.....	3.0	6.1	10.7	18.2	30.0	43.3	63.6	84.8	112.5	154.9
2001.....	3.0	5.8	10.6	18.8	29.3	43.2	62.9	86.8	111.6	157.1
2000.....	3.1	5.9	11.0	19.0	30.9	46.5	65.6	88.5	120.1	160.3
1999.....	3.6	6.5	11.5	20.9	34.6	52.1	72.9	96.0	128.0	168.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)										
White, male										
2002.....	2.9	6.7	13.2	21.1	36.5	53.9	78.5	103.6	139.1	198.4
2001.....	3.2	6.6	12.5	21.8	35.2	51.0	76.9	106.2	136.9	199.0
2000.....	3.2	6.7	12.8	22.1	36.1	55.9	80.1	106.9	149.0	201.3
1999.....	3.7	7.3	13.3	24.8	41.4	61.7	90.2	117.7	158.1	218.7
White, female										
2002.....	3.2	5.4	8.2	15.5	23.9	33.9	51.4	71.3	96.2	136.0
2001.....	2.9	5.0	8.7	16.0	23.9	36.4	51.5	73.0	96.3	139.6
2000.....	3.0	5.2	9.2	16.1	26.1	38.3	53.9	75.6	102.9	143.7
1999.....	3.5	5.6	9.8	17.1	28.3	43.8	59.2	80.8	110.2	147.7
Black, both sexes										
2002.....	4.8	10.3	17.2	28.5	46.1	58.8	84.7	108.1	127.9	168.4
2001.....	4.8	11.1	18.8	31.3	45.0	61.9	88.7	101.9	134.1	152.9
2000.....	5.9	11.5	19.9	29.1	46.8	59.5	86.2	117.5	128.8	155.2
1999.....	6.3	12.4	20.1	36.2	53.4	65.5	95.9	117.3	143.9	184.3
Black, male										
2002.....	5.9	11.7	22.7	35.6	63.4	71.6	108.2	145.6	162.0	212.3
2001.....	5.6	13.4	26.6	41.9	62.2	82.1	117.5	138.5	170.1	208.8
2000.....	7.5	15.1	26.0	38.7	67.5	76.0	110.7	169.6	160.2	204.1
1999.....	6.5	16.7	25.7	49.0	69.0	87.8	126.2	145.3	183.1	241.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)										
Black, female										
2002.....	3.8	9.1	12.6	22.7	32.5	49.4	69.0	85.6	110.2	151.6
2001.....	4.1	9.2	12.1	22.6	31.5	46.9	69.5	80.2	115.7	132.2
2000.....	4.5	8.4	14.7	21.2	30.4	47.4	69.9	86.0	112.9	136.9
1999.....	6.1	8.7	15.2	25.8	41.3	49.1	75.8	100.4	124.5	163.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
All races 3/, both sexes										
2002.....	4.6	7.5	12.6	21.5	36.9	59.2	90.1	125.6	168.8	197.5
2001.....	4.8	8.1	13.0	22.7	38.1	58.8	91.8	129.0	168.5	199.9
2000.....	4.8	8.3	13.4	23.4	38.2	62.2	91.6	131.4	176.0	204.6
1999.....	4.8	8.4	13.6	25.6	39.4	62.8	93.9	130.7	174.0	199.4
All races 3/, male										
2002.....	5.5	9.1	15.1	26.8	45.7	76.4	116.8	164.3	224.2	269.1
2001.....	5.9	9.8	16.0	27.7	47.5	75.2	116.9	165.7	220.9	275.3
2000.....	5.7	10.3	16.1	28.4	45.8	78.3	117.7	169.8	230.6	281.8
1999.....	5.9	10.2	16.7	31.0	48.6	78.1	118.9	169.4	221.8	279.0
All races 3/, female										
2002.....	3.7	6.0	10.1	16.4	28.9	44.4	68.8	98.2	135.1	166.5
2001.....	3.7	6.4	10.1	18.1	29.5	44.6	71.8	103.4	137.1	168.4
2000.....	3.9	6.5	10.8	18.8	31.3	48.4	71.0	104.7	143.8	173.1
1999.....	3.8	6.6	10.6	20.5	31.0	49.8	74.3	103.8	145.9	167.3
White, both sexes										
2002.....	4.6	7.2	12.1	21.0	36.9	60.3	91.7	128.7	173.6	204.0
2001.....	4.6	7.9	12.8	22.3	38.0	59.0	93.7	131.9	172.8	205.9
2000.....	4.7	8.0	13.2	23.3	38.5	63.5	93.4	134.4	181.5	208.7
1999.....	4.7	8.3	13.4	25.3	39.2	63.8	95.6	133.1	178.4	204.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
White, male										
2002.....	5.5	8.8	14.8	26.5	45.7	77.9	119.2	169.0	231.0	279.4
2001.....	5.7	9.7	15.8	27.1	47.3	75.6	119.2	169.8	226.4	284.8
2000.....	5.8	9.9	15.6	28.3	46.1	80.4	119.0	174.1	238.0	290.2
1999.....	5.6	10.0	16.5	30.6	48.6	79.4	121.0	172.1	228.5	285.8
White, female										
2002.....	3.7	5.7	9.6	15.8	28.8	44.8	69.4	99.9	138.5	171.2
2001.....	3.5	6.1	9.8	17.7	29.3	44.4	73.0	105.0	140.4	172.8
2000.....	3.6	6.1	10.7	18.6	31.4	48.7	72.8	106.5	147.9	175.5
1999.....	3.8	6.6	10.3	20.2	30.5	50.3	75.5	105.7	148.8	171.3
Black, both sexes										
2002.....	5.7	10.5	17.3	29.8	43.9	62.3	93.2	111.8	138.2	152.3
2001.....	6.6	10.4	16.5	30.2	44.9	67.8	90.7	116.7	142.7	155.7
2000.....	6.3	12.5	17.5	29.1	44.0	62.2	90.5	114.3	131.6	175.5
1999.....	6.5	10.4	18.2	33.2	45.9	64.6	90.4	119.1	139.3	162.0
Black, male										
2002.....	7.2	13.1	19.7	36.0	55.6	78.8	123.2	137.8	177.3	202.6
2001.....	8.2	11.9	21.0	37.5	56.9	87.9	114.1	141.2	191.0	206.5
2000.....	6.6	14.9	21.8	36.7	54.2	73.1	125.4	147.9	167.8	224.9
1999.....	8.4	13.3	22.5	41.2	55.2	80.7	114.6	159.1	164.8	220.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
Black, female										
2002.....	4.3	8.3	15.2	24.7	34.7	50.1	73.0	96.3	117.8	133.1
2001.....	5.3	9.2	12.7	24.2	35.4	52.9	75.1	102.1	118.1	136.8
2000.....	6.0	10.4	13.9	22.9	36.0	54.1	67.3	94.0	113.4	157.1
1999.....	4.8	8.0	14.5	26.6	38.6	52.7	74.3	95.0	126.6	140.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hodgkin's disease (C81)										
All races 3/, both sexes										
2002.....	0.3	0.4	0.5	0.5	0.8	1.0	1.6	1.7	2.3	2.7
2001.....	0.4	0.4	0.5	0.4	0.7	1.1	1.2	2.0	2.5	2.2
2000.....	0.3	0.4	0.5	0.5	0.9	1.0	1.6	1.8	2.4	2.5
1999.....	0.3	0.4	0.5	0.7	0.9	1.3	1.8	2.0	2.3	2.4
All races 3/, male										
2002.....	0.4	0.5	0.6	0.7	1.0	1.4	2.2	2.0	3.2	3.6
2001.....	0.5	0.5	0.6	0.6	0.9	1.5	1.6	2.5	2.9	2.6
2000.....	0.4	0.5	0.6	0.5	1.1	1.3	1.9	2.3	2.7	3.7
1999.....	0.4	0.5	0.6	0.9	1.2	1.6	2.3	2.3	2.8	3.8
All races 3/, female										
2002.....	0.3	0.3	0.4	0.3	0.7	0.6	1.1	1.5	1.7	2.3
2001.....	0.3	0.4	0.4	0.3	0.6	0.7	0.9	1.7	2.2	2.0
2000.....	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.4	2.3	2.0
1999.....	0.3	0.3	0.4	0.5	0.5	0.9	1.5	1.8	2.0	1.8
White, both sexes										
2002.....	0.4	0.4	0.5	0.5	0.8	1.0	1.7	1.8	2.4	2.9
2001.....	0.4	0.4	0.5	0.4	0.8	1.2	1.3	2.1	2.6	2.4
2000.....	0.3	0.4	0.5	0.5	0.9	1.0	1.6	2.0	2.5	2.6
1999.....	0.4	0.4	0.5	0.7	0.9	1.3	2.0	2.2	2.5	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)										
All races 3/, both sexes										
2002.....	1.9	3.0	5.0	8.8	14.9	23.3	35.1	50.3	67.3	77.6
2001.....	2.0	3.4	5.6	9.6	15.6	23.1	36.2	51.0	70.1	80.1
2000.....	2.1	3.7	5.8	10.7	16.1	25.1	36.4	53.9	72.1	80.7
1999.....	2.1	3.6	6.2	10.9	16.3	25.7	38.5	54.8	72.8	78.0
All races 3/, male										
2002.....	2.4	3.9	6.3	11.3	18.4	30.2	43.0	61.9	84.8	100.2
2001.....	2.7	4.4	7.4	11.9	19.1	29.3	45.0	61.4	87.8	107.1
2000.....	2.6	4.8	7.5	13.4	19.2	30.7	45.3	66.6	88.5	106.6
1999.....	2.8	4.8	7.9	13.5	19.6	30.9	46.8	68.4	85.9	104.7
All races 3/, female										
2002.....	1.3	2.2	3.9	6.4	11.7	17.4	28.7	42.1	56.6	67.8
2001.....	1.3	2.5	3.8	7.5	12.4	17.8	29.2	43.7	59.5	68.8
2000.....	1.5	2.6	4.1	8.2	13.3	20.3	29.4	45.0	62.5	70.2
1999.....	1.4	2.4	4.6	8.3	13.3	21.2	31.9	45.3	65.2	67.3
White, both sexes										
2002.....	1.8	3.0	5.0	9.0	15.6	24.5	36.6	53.0	70.6	82.4
2001.....	2.0	3.4	5.7	9.8	16.3	24.0	38.1	53.7	73.3	83.8
2000.....	2.1	3.6	5.9	11.2	17.0	26.4	38.3	56.7	76.2	84.8
1999.....	2.0	3.6	6.5	11.2	17.1	27.2	40.3	57.2	76.9	81.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)										
White, male										
2002.....	2.3	4.0	6.2	11.6	19.1	31.7	44.8	65.2	88.9	106.1
2001.....	2.7	4.4	7.5	12.0	19.8	30.5	46.9	64.3	91.7	111.9
2000.....	2.7	4.7	7.6	13.8	20.1	32.4	46.9	70.1	93.1	112.5
1999.....	2.7	4.7	8.1	13.8	20.3	32.4	49.0	70.5	91.1	108.3
White, female										
2002.....	1.3	2.1	3.7	6.5	12.3	18.2	29.9	44.3	59.5	72.1
2001.....	1.3	2.4	3.9	7.7	13.1	18.3	30.9	46.2	62.2	72.1
2000.....	1.4	2.6	4.3	8.8	14.1	21.2	31.3	47.2	66.1	73.5
1999.....	1.4	2.4	4.8	8.7	14.2	22.7	33.4	47.8	68.5	70.7
Black, both sexes										
2002.....	2.7	3.5	6.1	9.3	12.8	17.7	25.7	27.1	35.6	31.2
2001.....	2.4	4.3	5.9	9.9	12.4	19.2	23.5	25.9	38.7	39.6
2000.....	2.3	4.6	5.5	8.9	12.6	16.6	22.7	26.3	28.5	36.9
1999.....	2.8	4.1	5.7	10.6	11.7	17.0	23.9	32.1	31.5	40.8
Black, male										
2002.....	3.8	4.0	7.4	11.5	16.6	23.2	34.4	31.7	42.0	42.9
2001.....	3.4	5.1	8.7	12.4	15.5	24.9	31.6	31.1	48.2	55.6
2000.....	2.6	6.3	7.9	13.7	15.7	19.3	32.1	31.2	32.0	46.1
1999.....	3.9	5.6	8.0	13.9	15.7	23.4	28.5	47.5	28.8	59.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)										
Black, female										
2002.....	1.7	3.1	5.0	7.6	9.8	13.7	19.9	24.4	32.2	26.7
2001.....	1.6	3.6	3.6	8.0	9.9	15.0	18.2	22.7	33.9	33.7
2000.....	2.1	3.2	3.5	5.0	10.1	14.6	16.4	23.4	26.7	33.5
1999.....	1.8	2.8	3.7	7.9	8.7	12.3	20.9	22.9	32.8	33.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)										
All races 3/, both sexes										
2002.....	1.9	3.0	5.0	8.8	14.9	23.2	35.0	50.2	67.0	77.3
2001.....	2.0	3.4	5.5	9.6	15.6	23.0	36.0	50.9	69.8	79.6
2000.....	2.1	3.6	5.8	10.6	16.1	25.0	36.3	53.7	71.8	80.2
1999.....	2.1	3.5	6.2	10.8	16.3	25.6	38.4	54.6	72.6	77.7
All races 3/, male										
2002.....	2.4	3.9	6.2	11.3	18.4	30.0	42.8	61.8	84.4	99.7
2001.....	2.7	4.4	7.4	11.9	19.1	29.2	44.7	61.3	87.3	106.6
2000.....	2.6	4.7	7.5	13.3	19.2	30.5	45.2	66.3	88.1	105.9
1999.....	2.8	4.7	7.9	13.5	19.6	30.8	46.8	68.1	85.7	104.4
All races 3/, female										
2002.....	1.3	2.2	3.8	6.4	11.7	17.3	28.7	41.9	56.4	67.6
2001.....	1.3	2.5	3.8	7.5	12.4	17.7	29.1	43.6	59.3	68.3
2000.....	1.5	2.6	4.1	8.2	13.3	20.3	29.2	45.0	62.2	69.8
1999.....	1.4	2.4	4.5	8.3	13.3	21.1	31.8	45.2	65.0	66.9
White, both sexes										
2002.....	1.8	3.0	4.9	9.0	15.6	24.5	36.5	52.9	70.4	82.1
2001.....	2.0	3.4	5.7	9.8	16.2	23.9	37.9	53.6	73.0	83.4
2000.....	2.1	3.6	5.9	11.2	16.9	26.3	38.1	56.5	75.8	84.3
1999.....	2.0	3.6	6.4	11.1	17.0	27.1	40.2	57.0	76.7	81.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)										
White, male										
2002.....	2.3	4.0	6.2	11.6	19.1	31.6	44.5	65.0	88.6	105.6
2001.....	2.6	4.4	7.4	12.0	19.7	30.3	46.6	64.2	91.2	111.5
2000.....	2.7	4.7	7.6	13.7	20.0	32.2	46.8	69.8	92.6	111.8
1999.....	2.7	4.7	8.1	13.7	20.2	32.3	48.9	70.2	90.8	108.0
White, female										
2002.....	1.3	2.1	3.7	6.5	12.3	18.2	29.9	44.2	59.3	71.8
2001.....	1.3	2.4	3.9	7.7	13.0	18.3	30.9	46.0	62.0	71.6
2000.....	1.4	2.6	4.3	8.8	14.1	21.2	31.1	47.2	65.8	73.1
1999.....	1.4	2.4	4.8	8.7	14.1	22.6	33.3	47.7	68.3	70.4
Black, both sexes										
2002.....	2.7	3.5	6.1	9.3	12.8	17.6	25.7	26.8	35.3	30.6
2001.....	2.4	4.3	5.9	9.8	12.3	19.1	23.4	25.7	37.9	39.3
2000.....	2.3	4.5	5.5	8.8	12.6	16.6	22.7	25.9	28.2	36.3
1999.....	2.8	4.1	5.6	10.5	11.7	17.0	23.9	32.0	31.5	40.5
Black, male										
2002.....	3.8	4.0	7.4	11.5	16.6	22.9	34.4	31.7	41.3	42.9
2001.....	3.4	5.1	8.7	12.2	15.5	24.6	31.6	30.7	47.4	54.5
2000.....	2.6	6.1	7.9	13.5	15.7	19.3	32.1	30.7	32.0	43.8
1999.....	3.9	5.6	7.9	13.7	15.7	23.4	28.5	47.0	28.8	59.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)										
Black, female										
2002.....	1.7	3.1	5.0	7.6	9.8	13.7	19.9	23.9	32.2	25.9
2001.....	1.6	3.6	3.6	7.8	9.8	15.0	17.9	22.7	33.1	33.7
2000.....	2.1	3.2	3.5	5.0	10.1	14.6	16.4	23.1	26.3	33.5
1999.....	1.7	2.8	3.7	7.9	8.7	12.3	20.9	22.9	32.8	33.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant immunoproliferative diseases (C88)

All races 3/, both sexes

2002.....	*	*	*	*	0.2	0.3	0.6	1.2	1.7	2.0
2001.....	*	*	*	*	0.2	0.4	0.6	0.8	1.6	2.0
2000.....	*	*	*	*	*	0.3	0.6	0.9	1.7	2.1
1999.....	*	*	*	*	0.4	0.3	0.7	1.2	1.6	2.7

All races 3/, male

2002.....	*	*	*	*	0.4	0.5	0.9	1.7	2.9	2.8
2001.....	*	*	*	*	*	0.6	0.7	1.1	2.2	2.9
2000.....	*	*	*	*	*	*	0.8	1.3	2.4	3.4
1999.....	*	*	*	*	0.5	0.5	1.1	1.7	2.3	3.9

All races 3/, female

2002.....	*	*	*	*	*	*	*	0.8	1.0	1.6
2001.....	*	*	*	*	*	*	0.5	0.6	1.2	1.7
2000.....	*	*	*	*	*	*	0.4	0.5	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	0.9	1.2	2.2

White, both sexes

2002.....	*	*	*	*	0.3	0.3	0.6	1.3	1.8	2.1
2001.....	*	*	*	*	*	0.4	0.6	0.9	1.7	2.2
2000.....	*	*	*	*	*	0.3	0.6	0.9	1.8	2.2
1999.....	*	*	*	*	0.4	0.4	0.8	1.2	1.8	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and malignant plasma cell neoplasms (C90)										
All races 3/, both sexes										
2002.....	0.4	1.2	2.4	4.7	8.0	13.8	20.5	26.7	34.1	34.7
2001.....	0.5	1.2	2.3	5.0	8.3	13.6	20.5	27.4	33.1	34.2
2000.....	0.5	1.2	2.6	4.4	8.3	14.1	19.8	27.4	34.6	35.7
1999.....	0.5	1.2	2.4	5.4	8.7	13.8	19.8	26.3	34.5	33.9
All races 3/, male										
2002.....	0.5	1.5	2.9	5.5	9.8	16.4	25.0	33.8	43.3	48.0
2001.....	0.6	1.5	2.8	5.8	9.9	16.2	24.6	34.5	41.2	45.1
2000.....	0.6	1.4	2.8	4.8	9.2	16.6	23.4	34.4	43.7	48.3
1999.....	0.6	1.3	2.9	6.4	10.5	16.0	23.1	32.4	43.2	47.4
All races 3/, female										
2002.....	0.4	1.0	1.9	3.9	6.4	11.6	16.9	21.7	28.6	29.0
2001.....	0.4	0.8	1.9	4.2	7.0	11.3	17.2	22.5	28.3	29.7
2000.....	0.4	1.0	2.3	4.1	7.5	12.0	16.9	22.6	29.3	30.6
1999.....	0.4	1.1	2.0	4.6	7.1	11.8	17.3	22.1	29.4	28.4
White, both sexes										
2002.....	0.4	1.1	2.1	4.1	7.2	12.7	19.2	25.6	33.1	33.6
2001.....	0.4	1.0	2.0	4.3	7.3	12.5	19.2	26.1	31.3	32.8
2000.....	0.4	1.0	2.2	3.7	7.6	13.1	18.5	25.6	33.4	33.0
1999.....	0.4	1.1	2.0	4.6	7.5	12.8	18.4	24.5	33.2	31.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and malignant plasma cell neoplasms (C90)										
White, male										
2002.....	0.4	1.3	2.6	4.9	9.0	15.4	23.9	32.9	42.3	47.3
2001.....	0.5	1.3	2.5	5.0	8.7	15.0	23.4	33.4	39.3	44.0
2000.....	0.5	1.2	2.4	4.1	8.6	15.6	21.9	32.5	42.9	46.7
1999.....	0.4	1.2	2.6	5.5	9.5	15.2	21.4	30.8	41.9	45.4
White, female										
2002.....	0.3	0.8	1.5	3.4	5.6	10.4	15.4	20.4	27.4	27.7
2001.....	*	0.7	1.6	3.6	6.0	10.2	15.9	21.0	26.5	28.1
2000.....	0.2	0.8	2.0	3.3	6.7	10.9	15.8	20.8	27.8	27.4
1999.....	0.3	0.9	1.5	3.8	5.7	10.7	16.0	20.1	28.1	26.5
Black, both sexes										
2002.....	1.0	2.9	5.6	10.5	16.2	26.3	37.7	44.1	54.2	54.3
2001.....	1.5	2.6	4.8	11.8	19.0	25.5	38.1	47.6	60.9	57.1
2000.....	1.1	3.1	6.1	11.6	16.5	27.2	37.6	52.8	57.0	70.1
1999.....	1.4	2.4	6.4	13.6	19.6	24.1	37.5	50.2	56.1	62.5
Black, male										
2002.....	*	3.5	5.9	12.2	20.0	30.7	44.7	51.4	69.5	70.8
2001.....	1.6	3.4	5.3	14.5	23.0	29.7	45.1	57.1	79.5	69.2
2000.....	*	3.3	7.0	12.2	18.4	31.0	48.1	66.2	67.4	76.1
1999.....	1.9	2.6	6.4	15.9	20.8	26.0	46.5	58.4	73.2	80.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and malignant plasma cell neoplasms (C90)										
Black, female										
2002.....	*	2.3	5.3	9.2	13.3	23.1	32.9	39.8	46.2	48.1
2001.....	1.5	1.9	4.3	9.5	15.8	22.3	33.4	41.9	51.5	52.6
2000.....	*	2.8	5.4	11.2	15.0	24.4	30.6	44.7	51.8	67.8
1999.....	*	2.3	6.4	11.8	18.6	22.7	31.5	45.2	47.5	55.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
All races 3/, both sexes										
2002.....	1.9	2.8	4.5	7.4	12.9	20.9	32.4	45.7	63.4	80.5
2001.....	1.9	3.0	4.6	7.6	13.1	20.6	33.3	47.7	61.3	81.4
2000.....	1.9	3.1	4.5	7.7	12.7	21.7	33.2	47.4	65.1	83.6
1999.....	1.9	3.2	4.4	8.5	13.1	21.8	33.1	46.3	62.7	82.4
All races 3/, male										
2002.....	2.2	3.2	5.3	9.2	16.1	28.0	45.8	65.0	90.0	114.5
2001.....	2.1	3.4	5.2	9.2	17.4	27.6	44.9	66.2	86.9	117.5
2000.....	2.1	3.6	5.1	9.7	16.0	29.2	46.2	65.1	93.4	119.8
1999.....	2.1	3.6	5.2	10.1	16.8	29.0	45.7	64.6	87.6	119.2
All races 3/, female										
2002.....	1.6	2.5	3.8	5.7	10.0	14.7	21.8	32.1	47.3	65.7
2001.....	1.7	2.7	4.0	6.0	9.3	14.6	24.1	34.8	45.9	66.3
2000.....	1.7	2.6	3.9	5.9	9.7	15.3	22.9	35.1	48.4	68.8
1999.....	1.7	2.8	3.6	7.0	9.8	15.6	23.2	33.6	48.1	67.6
White, both sexes										
2002.....	2.0	2.7	4.5	7.4	13.1	21.8	33.6	47.0	65.7	83.0
2001.....	1.9	3.1	4.5	7.6	13.5	20.9	34.5	49.1	63.8	84.7
2000.....	1.9	3.0	4.5	7.8	12.8	22.6	34.4	49.2	67.6	86.0
1999.....	1.9	3.2	4.3	8.7	13.3	22.2	34.2	48.0	64.1	85.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
White, male										
2002.....	2.3	3.0	5.3	9.3	16.3	29.0	47.2	67.0	93.3	119.1
2001.....	2.0	3.5	5.1	9.4	17.8	27.8	46.5	68.3	90.1	123.1
2000.....	2.2	3.5	5.0	9.8	16.1	30.5	47.4	67.5	96.7	123.6
1999.....	2.1	3.5	5.1	10.3	17.0	29.5	47.1	66.7	90.2	124.1
White, female										
2002.....	1.7	2.5	3.8	5.6	10.1	15.4	22.6	32.8	48.9	67.3
2001.....	1.7	2.7	3.9	6.0	9.5	14.8	24.7	35.4	47.9	68.6
2000.....	1.6	2.5	4.0	5.9	9.8	15.7	23.9	36.3	50.2	70.8
1999.....	1.8	2.9	3.6	7.2	9.8	15.8	23.9	34.9	48.7	69.9
Black, both sexes										
2002.....	1.7	3.6	5.1	9.1	13.8	17.3	28.6	39.2	46.6	64.7
2001.....	2.3	3.0	5.6	7.9	12.3	22.3	28.0	41.3	42.0	57.1
2000.....	2.4	4.5	5.2	7.9	14.1	17.9	28.4	34.7	43.8	66.0
1999.....	2.0	3.5	5.4	8.1	13.6	22.2	27.9	35.0	51.1	55.8
Black, male										
2002.....	1.7	4.8	5.7	11.0	17.2	23.4	42.4	52.4	64.2	86.8
2001.....	2.7	2.8	6.6	9.9	17.1	32.3	36.4	51.6	61.0	77.2
2000.....	2.5	4.9	6.2	10.0	18.6	22.4	43.5	49.6	64.1	98.0
1999.....	2.2	4.4	6.9	10.3	17.2	29.1	38.0	51.3	61.0	77.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
Black, female										
2002.....	1.6	2.6	4.7	7.6	11.2	12.8	19.3	31.3	37.4	56.3
2001.....	2.0	3.3	4.7	6.3	8.5	15.0	22.4	35.2	32.3	49.7
2000.....	2.3	4.2	4.4	6.2	10.5	14.6	18.4	25.6	33.5	54.1
1999.....	1.7	2.6	4.2	6.2	10.8	17.1	21.2	25.2	46.2	47.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lymphoid leukemia (C91)										
All races 3/, both sexes										
2002.....	0.5	0.7	1.1	1.8	3.3	5.3	8.2	12.2	18.6	30.1
2001.....	0.4	0.6	1.1	1.9	3.3	5.4	8.7	12.9	18.2	31.2
2000.....	0.4	0.6	1.0	1.8	3.2	5.5	8.9	13.1	19.5	32.0
1999.....	0.3	0.6	1.1	2.2	3.5	6.0	9.2	13.4	19.7	30.8
All races 3/, male										
2002.....	0.6	0.8	1.5	2.4	4.4	7.8	12.5	17.7	26.9	43.0
2001.....	0.5	0.7	1.4	2.6	4.9	8.1	12.2	18.4	25.7	43.8
2000.....	0.5	0.8	1.2	2.7	4.6	8.2	12.8	18.9	28.4	43.8
1999.....	0.4	0.7	1.5	3.0	5.1	8.6	13.1	19.1	28.7	45.6
All races 3/, female										
2002.....	0.3	0.5	0.7	1.3	2.3	3.2	4.8	8.3	13.6	24.5
2001.....	0.3	0.5	0.7	1.1	1.9	3.1	5.8	9.0	13.8	25.9
2000.....	0.2	0.5	0.7	1.0	1.9	3.2	5.8	9.1	14.2	27.3
1999.....	0.3	0.5	0.6	1.4	2.1	3.7	6.2	9.4	14.4	24.8
White, both sexes										
2002.....	0.5	0.7	1.2	1.8	3.3	5.5	8.5	12.3	19.5	31.6
2001.....	0.4	0.6	1.0	1.9	3.4	5.3	9.0	13.2	19.1	32.8
2000.....	0.4	0.6	1.0	1.9	3.2	5.7	9.1	13.6	20.3	33.1
1999.....	0.4	0.6	1.1	2.1	3.5	6.1	9.6	13.9	20.3	32.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lymphoid leukemia (C91)										
White, male										
2002.....	0.7	0.8	1.6	2.4	4.5	8.2	12.8	17.8	28.3	45.1
2001.....	0.5	0.7	1.4	2.7	4.9	8.0	12.7	19.0	26.6	46.2
2000.....	0.5	0.8	1.2	2.8	4.6	8.4	13.1	19.5	29.5	45.0
1999.....	0.4	0.7	1.6	3.0	5.2	8.7	13.5	19.8	29.8	47.7
White, female										
2002.....	0.3	0.5	0.7	1.2	2.2	3.2	5.0	8.4	14.1	25.8
2001.....	0.3	0.4	0.6	1.1	2.0	2.9	5.9	9.0	14.5	27.3
2000.....	0.2	0.5	0.8	1.0	2.0	3.3	6.0	9.5	14.8	28.3
1999.....	0.3	0.5	0.6	1.4	2.1	3.8	6.5	9.8	14.7	25.7
Black, both sexes										
2002.....	*	*	1.0	2.6	4.3	5.2	7.9	14.3	13.1	18.4
2001.....	*	0.9	1.7	2.4	4.0	7.4	7.9	13.0	13.5	19.0
2000.....	*	1.0	1.1	1.5	3.5	4.5	9.1	10.5	13.6	25.0
1999.....	*	*	1.2	2.8	4.0	7.0	8.0	10.0	16.2	22.0
Black, male										
2002.....	*	*	*	3.3	5.7	7.0	11.7	21.6	18.3	31.1
2001.....	*	*	2.2	3.1	6.5	11.2	10.8	16.7	22.5	28.4
2000.....	*	*	*	*	6.0	7.3	13.7	16.5	21.1	40.4
1999.....	*	*	*	3.9	5.5	10.9	12.2	15.2	21.8	32.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lymphoid leukemia (C91)										
Black, female										
2002.....	*	*	*	*	3.1	3.9	5.4	9.9	10.3	13.6
2001.....	*	*	*	*	*	4.7	6.1	10.8	9.0	15.6
2000.....	*	*	*	*	*	*	6.0	6.8	9.8	19.3
1999.....	*	*	*	*	*	4.0	5.2	6.9	13.4	18.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Myeloid leukemia (C92)										
All races 3/, both sexes										
2002.....	1.1	1.6	2.5	4.0	7.1	10.9	16.1	21.6	26.9	25.7
2001.....	1.2	1.9	2.7	4.2	6.9	10.7	16.8	22.2	25.5	25.1
2000.....	1.2	1.8	2.7	4.3	6.8	11.1	16.0	22.0	26.7	25.3
1999.....	1.2	2.0	2.5	4.6	6.5	10.8	15.2	19.5	24.5	25.6
All races 3/, male										
2002.....	1.2	1.8	2.8	5.0	8.6	14.1	22.1	30.5	38.6	37.6
2001.....	1.3	2.0	2.8	4.9	9.0	13.4	22.2	29.9	37.0	37.2
2000.....	1.2	2.0	3.1	5.0	8.2	14.1	21.9	29.7	39.1	39.1
1999.....	1.2	2.1	2.8	5.2	8.0	13.8	20.7	26.3	34.6	37.7
All races 3/, female										
2002.....	1.0	1.5	2.3	3.1	5.7	8.2	11.3	15.3	19.9	20.5
2001.....	1.2	1.8	2.6	3.6	5.1	8.4	12.4	16.9	18.6	20.1
2000.....	1.2	1.6	2.3	3.7	5.6	8.5	11.3	16.6	19.4	19.7
1999.....	1.2	1.9	2.2	4.1	5.2	8.3	10.9	14.7	18.6	20.7
White, both sexes										
2002.....	1.2	1.6	2.5	4.0	7.3	11.4	16.9	22.5	27.9	26.5
2001.....	1.2	1.9	2.7	4.3	7.3	11.0	17.4	23.0	26.5	26.0
2000.....	1.1	1.7	2.7	4.3	6.8	11.6	16.8	23.0	27.7	25.9
1999.....	1.2	2.0	2.5	4.8	6.6	11.1	15.7	20.2	25.2	26.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Myeloid leukemia (C92)										
White, male										
2002.....	1.2	1.7	2.8	5.0	8.8	14.5	23.1	31.8	39.9	38.7
2001.....	1.2	2.0	2.8	4.9	9.4	13.7	22.9	30.9	38.7	39.1
2000.....	1.2	2.0	3.1	5.0	8.2	14.8	22.6	31.4	40.5	41.0
1999.....	1.2	2.1	2.7	5.4	8.0	14.0	21.4	27.2	35.8	39.5
White, female										
2002.....	1.1	1.5	2.2	3.0	5.9	8.6	11.8	15.9	20.6	21.2
2001.....	1.2	1.8	2.6	3.7	5.3	8.7	13.0	17.4	19.1	20.5
2000.....	1.1	1.5	2.4	3.7	5.5	8.8	12.1	17.2	20.1	19.8
1999.....	1.2	1.9	2.2	4.2	5.4	8.5	11.1	15.3	18.9	21.6
Black, both sexes										
2002.....	0.8	1.9	2.4	4.8	6.6	8.4	12.5	15.1	18.1	19.3
2001.....	1.6	1.6	2.8	4.2	4.6	10.4	12.8	17.2	15.7	17.2
2000.....	1.7	2.5	3.2	4.5	7.9	8.6	11.2	13.3	16.1	20.3
1999.....	1.4	2.1	2.8	3.4	6.7	9.6	12.4	14.1	17.9	14.3
Black, male										
2002.....	*	2.5	2.5	5.8	8.1	11.0	17.7	19.7	22.9	32.2
2001.....	1.7	*	3.1	5.3	6.3	14.0	17.8	20.9	19.3	*
2000.....	1.8	2.5	3.8	5.1	9.3	9.6	17.1	15.6	22.8	24.2
1999.....	*	2.5	3.7	4.4	8.7	12.5	16.6	19.9	20.9	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Myeloid leukemia (C92)

Black, female

2002.....	*	*	2.4	3.9	5.4	6.5	9.0	12.3	15.5	14.4
2001.....	1.4	1.8	2.5	3.3	3.2	7.7	9.4	15.0	13.9	16.4
2000.....	1.7	2.6	2.6	3.9	6.9	7.9	7.3	12.0	12.7	18.9
1999.....	1.4	1.6	*	*	5.2	7.5	9.7	10.6	16.4	12.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Monocytic leukemia (C93)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	*	*	0.4	0.4	0.6
2001.....	*	*	*	*	*	*	*	0.4	0.5	0.7
2000.....	*	*	*	*	*	*	0.2	0.3	0.4	0.7
1999.....	*	*	*	*	*	0.2	0.3	0.5	0.6	0.9
All races 3/, male										
2002.....	*	*	*	*	*	*	*	0.7	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	0.7	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	0.7
White, both sexes										
2002.....	*	*	*	*	*	*	*	0.4	0.4	0.6
2001.....	*	*	*	*	*	*	*	0.4	0.6	0.8
2000.....	*	*	*	*	*	*	*	0.4	*	0.7
1999.....	*	*	*	*	*	*	0.3	0.5	0.6	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified leukemia (C94-C95)										
All races 3/, both sexes										
2002.....	0.4	0.5	0.9	1.5	2.4	4.5	7.9	11.6	17.5	24.1
2001.....	0.3	0.5	0.8	1.4	2.8	4.4	7.7	12.2	17.0	24.4
2000.....	0.4	0.6	0.8	1.6	2.6	5.0	8.0	12.0	18.5	25.6
1999.....	0.4	0.6	0.8	1.7	3.0	4.8	8.3	13.0	17.8	25.2
All races 3/, male										
2002.....	0.4	0.6	1.0	1.7	3.0	5.9	10.9	16.1	24.3	32.6
2001.....	0.4	0.7	0.9	1.7	3.4	6.0	10.2	17.3	23.6	35.3
2000.....	0.4	0.8	0.7	2.0	3.1	6.8	11.1	16.1	25.1	35.7
1999.....	0.4	0.7	0.8	1.8	3.7	6.3	11.5	18.4	23.4	34.6
All races 3/, female										
2002.....	0.3	0.5	0.8	1.3	2.0	3.2	5.5	8.4	13.4	20.4
2001.....	0.2	0.4	0.7	1.2	2.2	3.0	5.6	8.6	13.1	19.8
2000.....	0.3	0.5	0.9	1.2	2.1	3.5	5.6	9.2	14.6	21.4
1999.....	0.3	0.5	0.8	1.5	2.4	3.4	5.9	9.2	14.6	21.3
White, both sexes										
2002.....	0.4	0.5	0.8	1.5	2.4	4.7	8.1	11.8	18.0	24.2
2001.....	0.3	0.6	0.8	1.5	2.7	4.4	7.8	12.5	17.6	25.1
2000.....	0.4	0.6	0.8	1.6	2.7	5.1	8.2	12.2	19.2	26.3
1999.....	0.4	0.6	0.7	1.7	3.1	4.8	8.6	13.4	18.0	25.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified leukemia (C94-C95)										
White, male										
2002.....	0.4	0.5	0.9	1.8	3.0	6.0	10.9	16.6	24.9	34.0
2001.....	0.3	0.7	0.9	1.7	3.3	6.0	10.6	17.8	24.1	36.4
2000.....	0.4	0.7	0.7	1.9	3.2	7.0	11.3	16.1	26.0	36.4
1999.....	0.5	0.7	0.8	1.8	3.9	6.5	11.9	19.0	23.6	35.4
White, female										
2002.....	0.3	0.4	0.8	1.3	1.9	3.5	5.7	8.4	13.7	20.0
2001.....	0.2	0.4	0.6	1.2	2.1	3.0	5.6	8.8	13.7	20.3
2000.....	0.3	0.5	0.9	1.2	2.2	3.5	5.7	9.4	15.1	22.2
1999.....	0.3	0.5	0.7	1.5	2.4	3.2	6.0	9.5	14.7	21.8
Black, both sexes										
2002.....	*	1.0	1.6	1.7	3.0	3.7	8.0	9.4	15.4	26.1
2001.....	*	*	1.0	*	3.6	4.5	7.3	10.9	12.5	20.6
2000.....	*	0.9	*	2.0	2.6	4.6	8.0	10.8	13.8	20.0
1999.....	*	*	1.5	1.8	2.9	5.4	7.2	10.5	16.5	19.4
Black, male										
2002.....	*	*	*	*	*	5.5	13.0	10.6	22.9	22.5
2001.....	*	*	*	*	4.3	7.1	7.7	13.9	18.5	29.5
2000.....	*	*	*	*	*	5.5	12.7	17.5	19.4	32.3
1999.....	*	*	*	*	*	5.7	8.8	15.2	18.3	24.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified leukemia (C94-C95)										
Black, female										
2002.....	*	*	*	*	*	*	4.7	8.8	11.5	27.5
2001.....	*	*	*	*	*	*	7.0	9.2	9.4	17.3
2000.....	*	*	*	*	*	4.0	4.9	6.8	11.0	15.5
1999.....	*	*	*	*	*	5.2	6.1	7.7	15.6	17.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Benign neoplasms (D10-D36)										
All races 3/, both sexes										
2002.....	0.2	0.4	0.4	0.5	0.8	1.4	1.8	3.1	4.3	7.8
2001.....	0.2	0.2	0.4	0.6	0.9	1.3	1.7	2.9	4.6	8.7
2000.....	0.2	0.2	0.4	0.4	0.7	1.2	1.9	3.2	5.5	9.0
1999.....	0.1	0.2	0.3	0.6	0.9	1.3	1.8	3.4	4.8	8.4
All races 3/, male										
2002.....	*	0.3	0.3	0.5	0.8	1.4	1.9	3.1	3.8	6.3
2001.....	*	*	0.4	0.6	1.0	1.3	1.6	3.1	5.1	8.9
2000.....	*	0.3	0.4	0.4	0.5	1.3	1.7	3.2	4.9	8.7
1999.....	0.2	*	0.3	0.6	0.9	1.1	2.1	3.1	4.6	9.2
All races 3/, female										
2002.....	0.3	0.4	0.4	0.5	0.7	1.5	1.8	3.2	4.6	8.4
2001.....	0.2	0.3	0.4	0.6	0.8	1.2	1.7	2.7	4.3	8.6
2000.....	0.3	0.2	0.4	0.5	0.9	1.1	2.0	3.3	5.9	9.1
1999.....	*	0.2	0.3	0.6	0.9	1.5	1.6	3.7	4.9	8.1
White, both sexes										
2002.....	0.2	0.3	0.4	0.5	0.7	1.3	1.8	3.1	4.1	7.4
2001.....	0.2	0.2	0.4	0.5	0.7	1.1	1.6	2.8	4.3	8.2
2000.....	0.2	0.2	0.4	0.4	0.7	1.1	1.7	3.2	5.5	8.9
1999.....	0.1	0.2	0.3	0.5	0.8	1.2	1.7	3.2	4.7	8.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of eye, brain and other parts of central nervous system (D31-D33)

All races 3/, both sexes

2002.....	0.1	0.2	0.2	0.3	0.4	0.9	1.2	2.1	2.9	5.1
2001.....	*	0.1	0.2	0.3	0.5	0.8	1.2	1.8	3.0	5.9
2000.....	*	0.1	0.2	0.2	0.5	0.7	1.2	2.3	3.7	6.2
1999.....	*	*	0.2	0.4	0.6	0.9	1.1	2.2	3.5	5.5

All races 3/, male

2002.....	*	*	*	*	0.4	0.8	1.2	1.7	2.4	3.2
2001.....	*	*	*	0.3	0.4	0.8	1.0	1.8	2.7	5.4
2000.....	*	*	*	*	*	0.8	1.1	2.0	3.3	5.9
1999.....	*	*	*	0.3	0.4	0.8	1.2	1.9	3.2	5.5

All races 3/, female

2002.....	*	*	0.3	0.3	0.4	1.0	1.2	2.3	3.2	5.9
2001.....	*	*	0.2	0.3	0.5	0.8	1.3	1.9	3.1	6.2
2000.....	*	*	*	0.3	0.6	0.7	1.3	2.4	4.0	6.4
1999.....	*	*	*	0.4	0.7	1.1	1.1	2.5	3.7	5.5

White, both sexes

2002.....	*	0.2	0.2	0.3	0.4	0.8	1.2	2.0	2.8	4.9
2001.....	*	*	0.2	0.2	0.4	0.7	1.1	1.8	2.8	5.5
2000.....	*	*	0.2	0.2	0.5	0.7	1.1	2.2	3.7	6.3
1999.....	*	*	0.1	0.3	0.5	0.9	1.1	2.1	3.4	5.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Of other and unspecified sites (D10-D24,D29-D30,D34-D36)

All races 3/, both sexes

2002.....	0.1	0.2	0.1	0.2	0.3	0.5	0.6	1.0	1.3	2.6
2001.....	0.1	0.1	0.2	0.3	0.4	0.5	0.5	1.0	1.6	2.5
2000.....	*	0.1	0.2	0.2	0.2	0.4	0.6	1.0	1.7	2.6
1999.....	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.3	2.8

All races 3/, male

2002.....	*	*	*	0.3	0.4	0.6	0.7	1.4	1.4	3.1
2001.....	*	*	0.2	0.3	0.5	0.5	0.5	1.3	2.4	3.5
2000.....	*	*	0.2	*	*	0.5	0.6	1.1	1.6	2.9
1999.....	*	*	*	*	0.5	*	0.9	1.2	1.4	3.7

All races 3/, female

2002.....	*	0.2	*	*	*	0.4	0.5	0.8	1.3	2.4
2001.....	*	*	*	*	*	0.4	*	0.8	1.1	2.1
2000.....	*	*	*	*	*	0.4	0.6	0.8	1.8	2.6
1999.....	*	*	*	*	*	0.4	0.5	1.2	1.2	2.4

White, both sexes

2002.....	*	0.1	0.1	0.2	0.3	0.4	0.5	1.1	1.3	2.3
2001.....	*	*	0.2	0.3	0.4	0.4	0.5	1.0	1.4	2.5
2000.....	*	*	0.2	0.2	0.2	0.4	0.5	0.9	1.8	2.5
1999.....	*	*	*	0.2	0.3	0.3	0.6	1.1	1.3	2.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)										
All races 3/, both sexes										
2002.....	0.7	1.0	1.5	3.0	4.4	7.9	14.0	24.7	40.0	67.0
2001.....	0.7	1.1	1.7	3.0	5.1	8.7	15.6	25.6	41.5	70.1
2000.....	0.7	1.1	1.8	3.2	5.5	9.0	14.9	24.6	43.2	72.0
1999.....	0.7	1.1	1.8	2.8	5.1	9.1	16.2	24.9	41.1	70.8
All races 3/, male										
2002.....	0.8	1.2	1.7	3.9	5.3	10.5	17.5	33.2	53.7	91.9
2001.....	0.8	1.4	1.9	3.2	6.3	10.9	20.3	35.1	55.7	96.6
2000.....	0.8	1.2	2.1	3.6	6.7	11.6	18.9	32.6	58.8	97.6
1999.....	0.9	1.2	2.2	3.2	6.7	11.8	21.2	33.2	55.0	97.8
All races 3/, female										
2002.....	0.6	0.9	1.3	2.3	3.5	5.6	11.1	18.7	31.7	56.2
2001.....	0.6	0.9	1.5	2.9	4.1	6.9	11.8	19.0	33.0	59.0
2000.....	0.6	1.0	1.4	2.8	4.4	6.8	11.7	19.0	33.9	61.5
1999.....	0.6	1.0	1.4	2.3	3.6	6.8	12.2	19.2	33.0	60.0
White, both sexes										
2002.....	0.6	1.1	1.5	3.0	4.5	8.2	14.4	25.6	41.1	69.3
2001.....	0.7	1.2	1.7	3.2	5.3	9.2	15.9	26.5	42.8	71.6
2000.....	0.7	1.1	1.7	3.3	5.5	9.3	15.0	25.3	44.9	74.3
1999.....	0.7	1.1	1.8	2.8	5.1	9.2	16.7	25.6	42.5	73.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)										
White, male										
2002.....	0.8	1.3	1.6	3.7	5.5	10.8	18.2	34.4	54.9	95.4
2001.....	0.8	1.4	2.0	3.4	6.4	11.4	21.0	36.1	57.3	99.2
2000.....	0.7	1.2	2.1	3.7	6.6	11.9	18.9	33.4	61.3	101.1
1999.....	0.9	1.2	2.2	3.3	6.9	11.8	21.9	34.2	57.1	102.7
White, female										
2002.....	0.5	0.9	1.3	2.2	3.6	5.9	11.4	19.3	32.6	57.9
2001.....	0.6	0.9	1.5	2.9	4.3	7.2	11.8	19.7	34.1	60.1
2000.....	0.6	1.0	1.4	2.8	4.5	7.0	11.8	19.6	35.0	63.4
1999.....	0.6	1.0	1.4	2.3	3.5	6.9	12.5	19.6	33.9	61.6
Black, both sexes										
2002.....	1.1	1.1	2.0	4.2	4.3	7.3	11.9	19.2	33.2	49.9
2001.....	1.0	0.9	1.9	2.8	5.1	6.9	14.8	19.8	33.3	55.6
2000.....	*	1.4	2.5	3.0	5.7	8.0	15.6	20.1	28.5	51.6
1999.....	1.0	1.4	2.1	2.9	5.6	10.5	14.2	21.6	30.3	48.8
Black, male										
2002.....	*	*	2.4	6.3	4.2	9.5	13.4	24.8	45.1	66.5
2001.....	*	*	*	*	6.3	9.1	17.8	27.4	46.6	71.5
2000.....	*	*	2.9	3.5	7.9	11.4	19.4	27.9	32.9	64.6
1999.....	*	*	*	3.6	5.9	14.5	17.6	26.1	34.9	52.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

Black, female

2002.....	*	*	*	2.6	4.3	5.7	11.0	15.9	27.1	43.5
2001.....	*	*	2.1	3.0	4.2	5.3	12.8	15.3	26.6	49.7
2000.....	*	*	2.2	2.7	3.9	5.4	13.1	15.4	26.3	46.8
1999.....	*	*	*	*	5.3	7.5	11.9	18.9	28.1	47.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Neoplasms of uncertain or unknown behavior of unspecified sites (D48)

All races 3/, both sexes

2002.....	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.9	1.9	3.8
2001.....	*	0.1	0.2	0.2	0.3	0.5	0.6	0.9	1.6	3.8
2000.....	*	0.1	*	0.2	0.4	0.3	0.6	1.1	1.9	3.9
1999.....	*	*	0.2	*	0.3	0.3	0.6	0.9	1.5	3.5

All races 3/, male

2002.....	*	*	*	*	*	*	0.6	0.9	1.5	3.0
2001.....	*	*	*	*	*	0.5	0.6	0.9	1.4	1.8
2000.....	*	*	*	*	*	*	*	1.1	1.6	2.4
1999.....	*	*	*	*	*	0.5	0.6	0.9	1.2	2.1

All races 3/, female

2002.....	*	*	*	*	0.4	*	0.4	0.9	2.2	4.2
2001.....	*	*	*	0.3	*	0.4	0.7	0.9	1.7	4.6
2000.....	*	*	*	*	0.4	*	0.6	1.1	2.0	4.4
1999.....	*	*	*	*	*	*	0.6	0.9	1.7	4.1

White, both sexes

2002.....	*	0.1	*	0.2	0.3	0.4	0.5	1.0	2.0	3.9
2001.....	*	0.1	0.2	0.2	0.3	0.4	0.5	0.9	1.6	4.0
2000.....	*	0.1	*	0.2	0.4	0.3	0.5	1.0	1.9	4.1
1999.....	*	*	0.1	*	0.3	0.3	0.6	0.8	1.5	3.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

All races 3/, both sexes

2002.....	1.7	2.2	2.6	3.0	4.3	6.0	9.4	13.9	24.7	56.8
2001.....	1.8	2.2	2.4	3.0	4.3	6.1	8.7	15.0	24.5	58.5
2000.....	1.6	2.1	2.5	3.1	4.3	6.4	8.8	14.1	25.9	58.9
1999.....	1.6	2.1	2.1	2.9	3.9	5.4	9.3	14.5	23.7	60.3

All races 3/, male

2002.....	1.6	2.1	2.5	3.1	4.5	5.5	10.2	14.8	25.1	57.2
2001.....	1.6	2.1	2.4	2.7	4.1	5.9	8.6	15.3	27.3	58.1
2000.....	1.5	2.2	2.4	2.9	4.2	6.5	9.8	14.2	27.3	60.4
1999.....	1.6	2.2	2.1	2.7	3.6	5.4	9.9	15.7	25.4	63.2

All races 3/, female

2002.....	1.8	2.3	2.6	3.0	4.1	6.4	8.8	13.3	24.4	56.6
2001.....	2.0	2.2	2.5	3.4	4.4	6.2	8.8	14.8	22.8	58.7
2000.....	1.6	1.9	2.6	3.3	4.4	6.3	8.1	14.1	25.0	58.3
1999.....	1.7	2.0	2.2	3.0	4.1	5.4	8.8	13.6	22.8	59.2

White, both sexes

2002.....	0.8	1.3	1.6	2.3	3.5	5.3	8.8	13.1	24.3	56.2
2001.....	0.9	1.1	1.5	2.3	3.6	5.5	8.1	14.5	23.8	58.3
2000.....	0.8	1.1	1.6	2.4	3.5	5.7	8.3	13.7	25.5	58.8
1999.....	0.8	1.2	1.4	2.0	3.2	4.7	8.7	14.0	23.5	59.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)										
White, male										
2002.....	0.8	1.4	1.6	2.4	3.8	5.0	9.5	14.2	24.5	56.3
2001.....	0.9	1.2	1.6	2.1	3.6	5.4	7.9	14.9	27.0	58.9
2000.....	0.9	1.3	1.6	2.4	3.5	6.2	9.2	13.8	26.8	59.7
1999.....	0.9	1.4	1.4	1.9	3.0	4.9	9.2	15.2	24.8	63.6
White, female										
2002.....	0.8	1.3	1.6	2.2	3.2	5.6	8.2	12.3	24.2	56.2
2001.....	0.9	1.0	1.5	2.5	3.7	5.5	8.3	14.3	21.9	58.0
2000.....	0.7	1.0	1.6	2.3	3.6	5.3	7.6	13.6	24.6	58.4
1999.....	0.7	1.0	1.3	2.0	3.5	4.6	8.2	13.1	22.7	58.0
Black, both sexes										
2002.....	7.6	9.0	10.7	10.5	12.1	13.4	16.9	25.6	31.9	71.9
2001.....	8.5	10.2	10.0	9.9	10.1	12.3	16.8	22.9	35.8	64.8
2000.....	7.0	9.2	10.5	10.3	11.8	13.0	14.7	21.0	35.9	66.6
1999.....	7.4	8.3	8.7	11.3	10.1	12.2	15.6	22.3	29.2	71.1
Black, male										
2002.....	7.0	8.1	11.1	10.4	12.1	12.2	19.4	24.8	35.9	78.3
2001.....	7.6	9.6	9.7	8.6	9.6	11.2	18.2	24.6	36.1	52.2
2000.....	6.1	9.4	10.6	8.1	11.8	10.9	15.7	19.8	36.2	73.8
1999.....	6.8	8.2	8.3	10.2	10.4	11.7	16.6	23.3	34.9	63.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

Black, female

2002.....	8.0	9.8	10.4	10.5	12.1	14.2	15.2	26.1	29.8	69.4
2001.....	9.3	10.8	10.1	11.0	10.6	13.1	15.9	21.9	35.6	69.5
2000.....	7.7	9.0	10.5	12.1	11.8	14.6	14.0	21.6	35.7	64.0
1999.....	7.9	8.4	9.1	12.2	9.8	12.5	14.9	21.7	26.4	74.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)										
All races 3/, both sexes										
2002.....	0.4	0.6	0.6	0.8	1.2	1.9	3.7	6.4	12.9	40.3
2001.....	0.6	0.5	0.7	0.8	1.2	2.0	3.5	7.1	13.6	42.0
2000.....	0.5	0.5	0.7	0.9	1.2	2.2	3.4	6.5	13.9	42.4
1999.....	0.5	0.7	0.6	0.8	1.2	1.8	3.5	6.8	13.7	43.8
All races 3/, male										
2002.....	0.4	0.5	0.6	0.8	1.3	1.7	4.3	6.6	12.4	38.3
2001.....	0.4	0.6	0.6	0.8	1.1	2.1	3.8	7.2	14.3	40.0
2000.....	0.4	0.5	0.7	1.0	1.2	2.2	3.6	6.5	14.6	40.1
1999.....	0.5	0.7	0.6	0.7	1.2	2.1	3.4	7.2	14.8	43.4
All races 3/, female										
2002.....	0.5	0.7	0.7	0.8	1.1	2.1	3.2	6.2	13.2	41.2
2001.....	0.7	0.5	0.7	0.9	1.2	1.9	3.3	7.0	13.2	42.9
2000.....	0.5	0.6	0.7	0.9	1.2	2.2	3.3	6.5	13.6	43.3
1999.....	0.4	0.7	0.6	0.9	1.1	1.6	3.6	6.5	13.0	44.0
White, both sexes										
2002.....	0.1	0.3	0.3	0.6	0.9	1.7	3.4	5.9	12.6	39.7
2001.....	0.2	0.2	0.4	0.6	1.0	1.8	3.2	6.7	13.2	41.7
2000.....	0.1	0.2	0.4	0.7	0.9	1.9	3.2	6.2	13.8	42.0
1999.....	0.1	0.3	0.3	0.4	0.8	1.6	3.2	6.5	13.6	43.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)										
White, male										
2002.....	*	0.3	0.3	0.6	1.0	1.6	3.7	6.3	11.9	37.2
2001.....	*	*	0.4	0.5	1.0	1.9	3.3	7.0	14.1	40.4
2000.....	*	*	0.4	0.8	0.9	2.1	3.4	6.1	14.2	39.1
1999.....	*	0.4	0.3	*	0.7	1.9	3.1	7.0	14.8	43.4
White, female										
2002.....	*	0.3	0.3	0.5	0.8	1.8	3.1	5.6	13.0	40.8
2001.....	0.3	*	0.4	0.6	1.1	1.6	3.1	6.5	12.6	42.3
2000.....	*	0.3	0.4	0.7	0.9	1.8	3.0	6.3	13.5	43.2
1999.....	*	0.3	0.3	0.5	0.9	1.4	3.4	6.1	12.9	42.9
Black, both sexes										
2002.....	2.4	3.1	3.1	3.2	4.1	4.4	7.9	13.7	19.4	52.3
2001.....	3.2	3.2	2.9	3.2	2.6	3.9	7.4	12.5	21.4	48.5
2000.....	2.7	2.9	3.2	3.0	3.9	4.8	6.1	10.3	18.1	51.0
1999.....	2.8	3.2	2.9	4.2	4.3	3.6	6.6	11.4	16.5	52.9
Black, male										
2002.....	2.3	2.6	2.6	3.0	5.1	*	11.7	12.9	23.7	56.8
2001.....	2.7	3.9	3.2	3.3	*	*	10.4	12.1	20.1	42.0
2000.....	2.2	3.2	3.4	*	*	*	*	10.9	19.4	55.4
1999.....	2.9	3.3	3.2	4.2	6.2	*	7.8	10.9	*	44.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)										
Black, female										
2002.....	2.6	3.6	3.5	3.3	3.3	4.6	5.4	14.3	17.1	50.5
2001.....	3.7	2.6	2.6	3.2	*	3.9	5.4	12.8	22.1	50.9
2000.....	3.2	2.7	3.0	3.0	3.9	5.8	6.2	10.0	17.4	49.4
1999.....	2.6	3.2	2.7	4.2	*	*	5.9	11.7	16.4	56.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional anemias (D50-D53)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	*	*	0.3	0.5	3.5
2001.....	*	*	*	*	*	*	*	0.5	0.8	3.6
2000.....	*	*	*	*	*	*	*	0.4	0.9	4.3
1999.....	*	*	*	*	*	*	0.2	0.4	1.1	4.0
All races 3/, male										
2002.....	*	*	*	*	*	*	*	*	*	2.8
2001.....	*	*	*	*	*	*	*	*	1.0	3.2
2000.....	*	*	*	*	*	*	*	*	*	5.1
1999.....	*	*	*	*	*	*	*	*	*	3.9
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	0.7	3.8
2001.....	*	*	*	*	*	*	*	0.5	0.7	3.8
2000.....	*	*	*	*	*	*	*	*	1.2	4.0
1999.....	*	*	*	*	*	*	*	*	1.2	4.0
White, both sexes										
2002.....	*	*	*	*	*	*	*	0.3	0.5	3.6
2001.....	*	*	*	*	*	*	*	0.5	0.8	3.7
2000.....	*	*	*	*	*	*	*	0.3	0.8	4.4
1999.....	*	*	*	*	*	*	*	0.3	1.1	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hemolytic anemias (D55-D59)										
All races 3/, both sexes										
2002.....	0.3	0.4	0.3	0.2	0.2	0.4	0.4	0.7	1.0	2.2
2001.....	0.4	0.3	0.3	0.3	0.2	0.3	0.4	0.7	1.2	2.2
2000.....	0.3	0.3	0.4	0.3	0.2	0.3	0.4	0.6	1.2	2.4
1999.....	0.3	0.4	0.2	0.3	0.2	0.3	0.4	0.7	1.2	2.2
All races 3/, male										
2002.....	0.3	0.3	0.3	*	*	*	*	*	*	2.2
2001.....	0.3	0.4	0.3	*	*	*	*	0.8	1.4	2.2
2000.....	0.3	0.3	0.3	*	*	*	*	*	1.2	2.5
1999.....	0.4	0.4	0.2	*	*	*	*	0.8	1.4	1.8
All races 3/, female										
2002.....	0.3	0.4	0.4	*	*	0.4	0.4	0.8	1.1	2.2
2001.....	0.5	0.3	0.3	0.3	*	*	*	0.7	1.1	2.2
2000.....	0.4	0.3	0.4	0.3	*	*	0.4	0.7	1.2	2.3
1999.....	0.3	0.4	*	0.4	*	*	0.5	0.7	1.0	2.4
White, both sexes										
2002.....	*	*	*	*	*	0.3	0.3	0.6	1.1	2.3
2001.....	*	*	*	*	*	0.3	0.3	0.7	1.2	2.2
2000.....	*	*	*	*	*	*	0.3	0.7	1.3	2.5
1999.....	*	*	*	*	*	0.2	0.4	0.8	1.2	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hemolytic anemias (D55-D59)										
Black, female										
2002.....	2.3	3.0	2.6	*	*	*	*	*	*	*
2001.....	3.1	2.3	1.9	*	*	*	*	*	*	*
2000.....	2.6	2.1	2.5	*	*	*	*	*	*	*
1999.....	2.0	2.6	*	2.8	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aplastic anemias (D60-D61)										
All races 3/, both sexes										
2002.....	*	0.1	0.1	0.2	0.4	0.6	1.1	1.8	3.2	5.3
2001.....	0.1	0.1	0.2	0.3	0.5	0.6	1.2	2.1	3.5	6.8
2000.....	*	0.1	0.2	0.3	0.5	0.8	1.1	1.8	3.6	7.4
1999.....	*	0.2	0.1	0.2	0.5	0.7	1.3	2.0	3.5	7.7
All races 3/, male										
2002.....	*	*	*	*	0.5	*	1.3	2.3	3.7	7.0
2001.....	*	*	0.2	*	0.5	0.6	1.1	2.3	3.9	7.8
2000.....	*	*	*	0.3	0.5	1.0	1.2	1.8	4.4	7.6
1999.....	*	*	*	*	0.5	1.0	1.2	2.4	4.3	10.1
All races 3/, female										
2002.....	*	*	*	0.3	*	0.8	0.9	1.5	2.8	4.5
2001.....	*	*	*	0.4	0.5	0.6	1.3	1.9	3.3	6.3
2000.....	*	*	*	*	0.5	0.7	1.0	1.8	3.1	7.3
1999.....	*	*	*	*	0.4	0.4	1.3	1.8	3.0	6.8
White, both sexes										
2002.....	*	0.1	*	0.2	0.3	0.6	1.0	1.9	3.2	5.2
2001.....	*	0.1	0.2	0.3	0.5	0.6	1.2	2.2	3.6	7.1
2000.....	*	0.1	0.1	0.3	0.5	0.8	1.1	1.8	3.7	7.4
1999.....	*	0.1	0.1	*	0.4	0.7	1.2	2.0	3.6	7.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute posthemorrhagic and other anemias (D62,D64)										
All races 3/, both sexes										
2002.....	*	0.1	0.1	0.3	0.6	0.8	2.0	3.6	8.3	29.4
2001.....	*	*	0.1	0.2	0.4	0.9	1.7	3.8	8.1	29.5
2000.....	*	*	0.1	0.3	0.5	1.0	1.8	3.6	8.3	28.2
1999.....	*	0.1	0.2	0.2	0.4	0.8	1.6	3.6	7.9	29.9
All races 3/, male										
2002.....	*	*	*	0.4	0.6	0.9	2.4	3.6	7.8	26.3
2001.....	*	*	*	*	*	1.0	2.1	3.7	8.0	26.9
2000.....	*	*	*	0.4	0.6	0.9	1.9	3.6	8.6	24.9
1999.....	*	*	*	*	0.4	0.8	1.7	3.5	8.0	27.7
All races 3/, female										
2002.....	*	*	*	0.3	0.5	0.7	1.8	3.6	8.5	30.7
2001.....	*	*	*	*	0.4	0.9	1.4	3.9	8.1	30.5
2000.....	*	*	*	0.3	0.4	1.1	1.7	3.6	8.1	29.6
1999.....	*	*	*	*	0.5	0.8	1.6	3.7	7.8	30.8
White, both sexes										
2002.....	*	*	*	0.3	0.4	0.7	1.9	3.0	7.8	28.7
2001.....	*	*	*	0.2	0.3	0.8	1.5	3.4	7.6	28.8
2000.....	*	*	*	0.3	0.3	0.8	1.6	3.4	8.0	27.8
1999.....	*	*	*	*	0.3	0.7	1.4	3.4	7.8	28.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute posthemorrhagic and other anemias (D62,D64)

Black, female

2002.....	*	*	*	*	*	*	*	11.3	13.9	40.3
2001.....	*	*	*	*	*	*	*	9.7	16.4	42.5
2000.....	*	*	*	*	*	*	*	7.7	12.7	38.6
1999.....	*	*	*	*	*	*	*	7.7	10.8	42.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)										
All races 3/, both sexes										
2002.....	0.4	0.7	0.7	0.9	1.4	2.2	3.5	4.8	7.7	11.3
2001.....	0.5	0.6	0.7	1.1	1.5	2.3	3.1	4.9	7.6	11.1
2000.....	0.4	0.5	0.8	1.1	1.6	2.6	3.3	4.8	7.9	11.6
1999.....	0.4	0.6	0.7	1.0	1.5	2.2	3.7	5.4	6.9	11.5
All races 3/, male										
2002.....	0.4	0.6	0.7	0.9	1.6	2.1	3.5	5.6	8.4	13.9
2001.....	0.4	0.6	0.7	1.0	1.6	2.3	3.3	4.8	9.6	12.3
2000.....	0.4	0.7	0.8	1.0	1.5	2.7	3.8	4.9	8.3	14.6
1999.....	0.4	0.6	0.8	1.0	1.3	2.2	4.1	5.8	6.8	14.3
All races 3/, female										
2002.....	0.4	0.8	0.7	0.9	1.2	2.3	3.6	4.3	7.2	10.1
2001.....	0.5	0.6	0.7	1.1	1.4	2.2	3.0	5.0	6.4	10.5
2000.....	0.3	0.4	0.8	1.3	1.7	2.5	2.9	4.7	7.7	10.4
1999.....	0.5	0.5	0.6	1.0	1.8	2.2	3.3	5.1	6.9	10.3
White, both sexes										
2002.....	0.4	0.6	0.6	0.8	1.3	2.0	3.5	4.6	7.8	11.3
2001.....	0.4	0.5	0.6	1.0	1.3	2.2	3.1	4.8	7.4	11.0
2000.....	0.3	0.5	0.6	1.0	1.5	2.5	3.1	4.7	7.9	11.8
1999.....	0.4	0.5	0.6	0.8	1.5	1.9	3.5	5.2	6.7	11.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)										
White, male										
2002.....	0.4	0.5	0.6	0.8	1.5	1.9	3.5	5.4	8.2	14.1
2001.....	0.4	0.4	0.6	0.9	1.4	2.2	3.1	4.6	9.6	12.5
2000.....	0.3	0.6	0.6	0.9	1.4	2.5	3.6	4.9	8.3	14.8
1999.....	0.4	0.5	0.7	0.8	1.2	1.9	3.9	5.4	6.3	14.6
White, female										
2002.....	0.4	0.6	0.6	0.8	1.1	2.1	3.4	4.0	7.5	10.1
2001.....	0.4	0.5	0.6	1.1	1.2	2.2	3.0	5.0	6.1	10.4
2000.....	0.2	0.4	0.7	1.1	1.6	2.4	2.8	4.5	7.6	10.5
1999.....	0.3	0.4	0.5	0.7	1.8	2.0	3.1	5.0	7.0	10.2
Black, both sexes										
2002.....	*	1.4	1.8	1.9	2.3	4.5	4.8	7.9	7.3	13.1
2001.....	1.0	1.9	1.8	1.6	2.9	3.0	4.2	6.4	10.3	12.3
2000.....	0.8	1.1	2.5	2.4	2.8	3.6	5.3	6.9	10.2	11.3
1999.....	1.1	1.1	1.8	2.8	2.3	4.6	5.4	8.2	8.7	13.7
Black, male										
2002.....	*	*	2.3	*	*	*	*	*	*	*
2001.....	*	1.9	*	*	*	*	*	*	*	*
2000.....	*	*	2.6	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	5.5	*	10.0	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)

Black, female

2002.....	*	1.8	*	*	*	4.6	5.4	8.0	*	12.3
2001.....	*	1.8	*	*	*	*	*	5.8	9.8	13.9
2000.....	*	*	2.4	3.0	*	*	4.7	7.7	11.0	10.3
1999.....	*	*	*	2.9	*	4.0	5.4	7.2	*	13.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of blood and blood-forming organs (D70-D76)										
All races 3/, both sexes										
2002.....	0.1	0.1	0.3	0.4	0.5	0.7	1.1	1.6	2.6	3.4
2001.....	0.1	0.1	0.2	0.3	0.5	0.8	0.9	1.8	2.0	3.9
2000.....	0.1	0.2	0.3	0.3	0.5	0.8	1.0	1.6	2.2	3.4
1999.....	0.1	0.1	0.2	0.4	0.4	0.5	1.2	1.3	2.2	3.6
All races 3/, male										
2002.....	*	*	0.3	0.4	0.6	0.8	1.4	1.8	2.7	3.5
2001.....	*	*	0.3	0.3	0.6	0.8	0.8	2.2	2.2	3.9
2000.....	*	0.2	*	*	0.5	1.0	1.2	1.8	2.7	4.1
1999.....	*	*	*	0.4	0.5	0.5	1.5	1.8	2.7	4.0
All races 3/, female										
2002.....	*	*	0.2	0.3	0.5	0.6	0.8	1.4	2.5	3.3
2001.....	*	*	0.2	0.3	0.5	0.7	0.9	1.5	1.9	3.8
2000.....	*	*	0.3	*	0.6	0.7	0.8	1.4	1.9	3.1
1999.....	*	*	0.3	0.4	0.4	0.6	0.9	1.0	1.8	3.4
White, both sexes										
2002.....	0.1	0.1	0.3	0.4	0.6	0.7	1.1	1.6	2.6	3.4
2001.....	0.1	0.1	0.2	0.3	0.5	0.7	0.9	1.9	2.0	3.9
2000.....	*	0.1	0.2	0.3	0.5	0.8	1.0	1.6	2.2	3.5
1999.....	0.1	*	0.2	0.4	0.4	0.5	1.2	1.4	2.2	3.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Certain disorders involving the immune mechanism (D80-D89)										
All races 3/, both sexes										
2002.....	0.7	0.8	1.0	1.0	1.1	1.1	1.0	1.1	1.5	1.8
2001.....	0.7	0.9	0.8	0.8	1.1	1.1	1.2	1.2	1.3	1.5
2000.....	0.6	0.8	0.8	0.8	0.9	0.8	1.1	1.3	1.8	1.6
1999.....	0.6	0.7	0.6	0.7	0.7	0.9	0.9	1.0	1.0	1.4
All races 3/, male										
2002.....	0.7	0.9	0.9	1.0	1.0	0.9	1.0	0.8	1.5	1.5
2001.....	0.7	0.8	0.8	0.6	0.9	0.7	0.8	1.1	1.2	1.8
2000.....	0.6	0.8	0.7	0.7	0.9	0.7	1.1	1.0	1.7	1.6
1999.....	0.5	0.8	0.6	0.6	0.7	0.7	0.8	1.0	*	*
All races 3/, female										
2002.....	0.8	0.7	1.0	1.0	1.2	1.4	1.1	1.3	1.5	1.9
2001.....	0.7	1.0	0.9	1.0	1.3	1.4	1.6	1.3	1.4	1.4
2000.....	0.6	0.8	0.8	0.9	0.9	0.9	1.0	1.5	1.9	1.5
1999.....	0.6	0.7	0.6	0.8	0.8	1.0	1.0	1.1	1.0	1.5
White, both sexes										
2002.....	0.2	0.3	0.4	0.6	0.7	0.8	0.9	1.1	1.4	1.8
2001.....	0.2	0.3	0.4	0.4	0.8	0.8	0.9	1.1	1.2	1.6
2000.....	0.3	0.3	0.3	0.4	0.6	0.5	1.0	1.2	1.6	1.5
1999.....	0.2	0.3	0.3	0.4	0.5	0.7	0.7	1.0	0.9	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Certain disorders involving the immune mechanism (D80-D89)

White, male

2002.....	0.2	0.4	0.4	0.6	0.8	0.6	1.0	0.8	1.6	*
2001.....	0.2	0.4	0.4	*	0.7	0.6	0.6	0.9	1.1	2.0
2000.....	0.3	0.3	0.4	0.5	0.7	0.6	1.1	1.0	1.5	*
1999.....	0.3	0.4	0.3	0.4	0.6	0.6	0.6	1.0	*	*

White, female

2002.....	*	0.2	0.4	0.5	0.6	1.0	0.8	1.2	1.3	1.9
2001.....	*	0.3	0.3	0.5	0.9	1.0	1.1	1.2	1.3	1.5
2000.....	0.2	*	0.3	*	0.5	0.5	0.9	1.3	1.6	1.5
1999.....	*	*	*	0.4	0.5	0.7	0.8	1.0	0.9	1.4

Black, both sexes

2002.....	4.4	4.3	5.5	5.0	5.0	4.1	3.1	*	*	*
2001.....	4.0	5.0	4.8	4.8	4.0	4.0	4.6	*	*	*
2000.....	3.3	4.9	4.5	4.5	4.2	3.5	*	*	*	*
1999.....	3.4	3.7	3.6	3.3	2.7	3.1	*	*	*	*

Black, male

2002.....	4.0	4.3	5.8	4.8	4.0	*	*	*	*	*
2001.....	3.9	3.7	3.9	3.4	*	*	*	*	*	*
2000.....	3.1	4.7	4.3	*	*	*	*	*	*	*
1999.....	2.8	3.5	3.0	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Certain disorders involving the immune mechanism (D80-D89)

Black, female

2002.....	4.9	4.3	5.3	5.1	5.9	4.8	*	*	*	*
2001.....	4.2	6.2	5.6	5.9	4.6	5.3	5.8	*	*	*
2000.....	3.4	5.1	4.8	5.9	4.4	4.6	*	*	*	*
1999.....	4.0	3.9	4.1	4.0	3.5	4.2	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
IV. Endocrine, nutritional and metabolic diseases (E00-E88)										
All races 3/, both sexes										
2002.....	9.2	14.0	24.2	39.1	59.2	91.3	133.0	197.1	288.8	504.8
2001.....	8.0	14.0	23.8	37.3	59.1	90.6	132.7	194.8	286.9	519.2
2000.....	8.4	13.3	22.9	37.3	58.9	89.4	131.9	195.7	286.5	521.0
1999.....	8.1	13.3	21.7	36.5	58.8	90.3	132.3	191.6	285.9	529.8
All races 3/, male										
2002.....	11.4	16.3	28.7	46.4	68.9	102.1	147.7	217.5	313.8	504.5
2001.....	9.5	16.8	27.9	42.9	67.4	102.8	148.3	215.1	310.2	507.7
2000.....	9.8	16.0	26.4	42.5	66.6	98.5	146.0	215.1	311.8	517.4
1999.....	9.4	16.1	25.2	41.9	65.3	98.8	146.0	213.4	312.5	521.0
All races 3/, female										
2002.....	7.0	11.7	19.8	32.2	50.4	81.9	121.2	182.7	273.5	504.9
2001.....	6.5	11.3	19.9	31.9	51.6	80.0	120.3	180.6	272.9	524.0
2000.....	7.0	10.7	19.4	32.4	52.0	81.6	120.7	182.3	271.5	522.5
1999.....	6.8	10.6	18.3	31.4	52.9	83.0	121.6	176.4	270.3	533.3
White, both sexes										
2002.....	8.3	12.3	21.2	34.5	52.5	81.0	121.2	183.8	274.3	487.1
2001.....	7.2	12.2	21.0	32.8	52.6	81.3	121.4	181.3	272.1	502.4
2000.....	7.7	11.6	19.9	32.9	52.9	80.1	120.6	181.8	270.5	503.5
1999.....	7.2	11.4	18.7	31.1	51.4	80.8	119.6	178.3	269.9	511.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
IV. Endocrine, nutritional and metabolic diseases (E00-E88)										
White, male										
2002.....	10.5	14.6	25.9	42.1	62.2	93.2	138.7	208.7	303.9	498.9
2001.....	8.9	14.8	25.1	38.7	61.2	95.1	139.9	204.3	297.8	500.5
2000.....	9.2	14.4	23.8	38.3	60.9	91.0	137.2	204.2	302.4	510.5
1999.....	8.5	14.2	22.5	36.8	58.9	90.4	136.6	203.3	301.4	504.7
White, female										
2002.....	6.1	10.1	16.5	27.3	43.6	70.2	107.0	166.0	256.1	482.1
2001.....	5.4	9.6	17.0	27.1	44.6	69.2	106.5	165.0	256.6	503.2
2000.....	6.2	8.9	16.1	27.8	45.5	70.5	107.4	166.0	251.5	500.6
1999.....	5.8	8.7	15.0	25.6	44.5	72.4	106.1	160.7	251.2	514.0
Black, both sexes										
2002.....	17.3	28.3	51.2	83.3	125.4	186.0	261.4	356.3	491.9	757.0
2001.....	15.1	28.7	50.2	81.1	121.3	180.4	263.0	363.7	482.9	763.1
2000.....	15.2	28.3	49.9	79.6	117.6	183.3	258.5	375.1	503.2	766.1
1999.....	16.2	28.7	49.8	84.7	127.7	183.9	268.1	349.9	497.3	763.6
Black, male										
2002.....	20.6	32.0	56.7	92.7	143.3	192.0	264.0	352.7	492.9	610.0
2001.....	16.2	33.8	55.1	88.9	133.2	189.6	265.6	375.4	511.2	658.1
2000.....	16.5	31.6	53.2	87.3	126.7	180.1	265.1	376.6	480.5	655.0
1999.....	18.0	33.5	53.0	92.9	130.5	193.2	261.9	347.6	485.7	746.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
IV. Endocrine, nutritional and metabolic diseases (E00-E88)										
Black, female										
2002.....	14.4	25.0	46.5	75.5	111.4	181.5	259.7	358.4	491.4	813.3
2001.....	14.2	24.2	46.0	74.8	112.0	173.7	261.2	356.7	468.5	802.0
2000.....	14.0	25.4	47.1	73.3	110.5	185.7	254.2	374.2	514.6	807.4
1999.....	14.7	24.6	47.2	78.0	125.5	177.0	272.3	351.3	503.0	770.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Endocrine diseases (E00-E34)										
All races 3/, both sexes										
2002.....	6.3	10.2	18.5	30.7	48.4	76.9	111.3	161.9	226.5	347.6
2001.....	5.6	10.2	18.3	30.1	49.3	76.6	110.8	160.4	227.1	350.3
2000.....	5.8	9.8	17.7	30.3	48.9	75.2	110.7	160.9	222.4	345.2
1999.....	5.7	10.3	17.0	30.2	49.7	77.0	111.6	157.7	223.2	343.6
All races 3/, male										
2002.....	7.9	12.1	21.9	36.6	56.4	85.4	122.9	178.9	248.9	365.2
2001.....	6.8	12.3	21.8	34.9	55.3	85.7	123.0	175.6	247.7	362.2
2000.....	7.0	11.6	20.1	34.0	54.5	81.7	121.4	177.0	247.4	360.9
1999.....	6.6	12.5	19.8	34.1	54.6	83.6	121.3	175.9	248.6	361.8
All races 3/, female										
2002.....	4.7	8.3	15.3	25.2	41.2	69.6	102.0	149.9	212.9	339.9
2001.....	4.5	8.1	15.0	25.6	44.0	68.7	101.2	149.7	214.7	345.3
2000.....	4.7	8.1	15.4	26.9	43.8	69.5	102.3	149.7	207.7	338.8
1999.....	4.8	8.1	14.3	26.5	45.3	71.3	104.1	145.0	208.4	336.3
White, both sexes										
2002.....	5.5	8.7	15.7	26.5	42.2	67.2	100.5	149.5	213.6	331.6
2001.....	5.0	8.8	15.7	25.8	43.1	67.9	100.1	147.7	214.1	335.2
2000.....	5.2	8.3	15.0	26.4	43.1	66.2	100.1	147.8	207.5	329.3
1999.....	4.9	8.6	14.3	25.0	42.6	68.1	99.9	145.2	208.9	325.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Endocrine diseases (E00-E34)										
White, male										
2002.....	7.0	10.5	19.1	32.4	50.2	76.9	114.5	170.9	240.8	361.5
2001.....	6.3	10.7	19.1	30.8	49.6	78.5	115.3	165.5	237.4	356.8
2000.....	6.4	10.0	17.6	30.1	49.3	74.6	113.2	167.2	238.9	353.0
1999.....	5.7	10.8	17.2	29.3	48.4	75.9	113.0	167.1	240.0	349.4
White, female										
2002.....	4.0	6.9	12.5	20.8	34.7	58.7	89.1	134.2	196.9	318.5
2001.....	3.6	6.9	12.5	21.0	37.2	58.5	87.8	135.1	200.0	326.1
2000.....	4.0	6.7	12.4	22.9	37.4	58.8	89.6	134.2	188.8	319.7
1999.....	4.1	6.5	11.4	21.0	37.4	61.3	89.4	129.9	190.5	316.4
Black, both sexes										
2002.....	12.6	22.0	42.1	69.3	107.6	163.3	226.6	305.3	400.6	559.8
2001.....	11.4	21.6	41.1	70.7	106.2	159.8	230.8	312.3	394.1	555.5
2000.....	10.8	22.4	41.4	67.6	103.3	162.5	228.0	325.0	415.6	550.6
1999.....	11.8	23.7	41.5	74.8	113.5	162.3	233.8	301.0	408.8	552.6
Black, male										
2002.....	15.4	25.9	47.9	77.6	121.2	167.3	228.0	299.5	391.3	427.8
2001.....	12.3	26.2	46.3	77.7	113.5	164.4	226.5	317.3	409.3	469.7
2000.....	12.1	25.8	44.2	73.8	108.9	156.9	228.7	317.5	389.5	474.0
1999.....	13.6	28.3	45.1	80.5	115.2	168.2	220.8	288.7	388.0	523.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Endocrine diseases (E00-E34)										
Black, female										
2002.....	10.1	18.7	37.2	62.4	97.0	160.4	225.6	308.7	405.4	610.4
2001.....	10.6	17.6	36.7	64.9	100.5	156.4	233.6	309.2	386.3	587.3
2000.....	9.7	19.4	39.0	62.5	98.9	166.7	227.5	329.5	428.8	579.1
1999.....	10.3	19.7	38.5	70.1	112.1	158.0	242.4	308.4	419.2	563.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of thyroid gland (E00-E07)										
All races 3/, both sexes										
2002.....	0.1	0.1	0.2	0.3	0.4	0.5	1.4	2.5	6.0	22.7
2001.....	0.1	0.1	0.2	0.3	0.4	0.6	1.1	3.0	5.8	23.2
2000.....	0.1	*	0.2	0.3	0.4	0.5	1.3	2.7	5.9	21.1
1999.....	0.1	0.2	0.2	0.3	0.3	0.6	1.3	2.5	6.2	22.4
All races 3/, male										
2002.....	*	*	*	*	*	*	0.8	1.6	3.0	12.7
2001.....	*	*	*	*	*	*	0.7	2.3	2.8	14.1
2000.....	*	*	*	*	*	*	0.9	1.5	2.6	12.1
1999.....	*	*	*	*	*	0.5	0.8	1.5	4.4	12.9
All races 3/, female										
2002.....	*	0.2	0.3	0.4	0.6	0.6	1.8	3.1	7.8	27.0
2001.....	*	*	0.3	0.4	0.5	0.8	1.4	3.6	7.6	27.1
2000.....	0.2	*	0.3	0.4	0.6	0.7	1.5	3.5	7.8	24.8
1999.....	*	0.3	*	0.4	0.4	0.8	1.7	3.2	7.3	26.3
White, both sexes										
2002.....	*	*	0.1	0.2	0.4	0.6	1.3	2.5	6.2	23.5
2001.....	*	*	0.2	0.3	0.4	0.6	1.1	3.0	5.5	23.9
2000.....	*	*	0.2	0.3	0.5	0.4	1.2	2.6	5.8	21.6
1999.....	*	0.1	0.2	0.2	0.3	0.6	1.3	2.5	6.1	23.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)										
All races 3/, both sexes										
2002.....	6.0	9.9	18.0	30.1	47.5	75.7	108.8	157.7	217.9	320.6
2001.....	5.4	9.9	17.8	29.5	48.4	75.5	108.6	155.5	218.7	321.8
2000.....	5.5	9.5	17.3	29.6	48.1	74.0	108.6	156.5	213.9	319.7
1999.....	5.5	9.9	16.5	29.5	48.9	75.6	109.2	153.5	214.8	317.2
All races 3/, male										
2002.....	7.7	11.9	21.5	36.0	55.5	84.3	121.1	175.9	243.5	348.5
2001.....	6.6	12.1	21.3	34.4	54.4	84.7	121.0	171.5	242.3	343.1
2000.....	6.8	11.4	19.7	33.4	53.9	80.8	119.7	173.7	242.5	345.1
1999.....	6.4	12.3	19.4	33.5	53.9	82.5	119.4	172.3	241.3	345.7
All races 3/, female										
2002.....	4.4	7.9	14.7	24.5	40.2	68.2	99.0	144.9	202.3	308.4
2001.....	4.2	7.9	14.5	24.9	42.9	67.6	98.8	144.3	204.5	312.9
2000.....	4.3	7.8	14.9	26.1	42.8	68.1	99.9	144.6	197.0	309.4
1999.....	4.6	7.5	13.8	25.8	44.5	69.7	101.2	140.4	199.3	305.8
White, both sexes										
2002.....	5.3	8.5	15.4	26.0	41.3	66.0	98.2	145.4	204.9	303.7
2001.....	4.8	8.6	15.3	25.3	42.3	66.8	98.0	143.0	205.9	306.0
2000.....	5.0	8.0	14.6	25.7	42.3	65.2	98.1	143.7	199.1	303.5
1999.....	4.7	8.3	13.8	24.5	41.9	66.9	97.6	141.2	200.8	298.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)										
White, male										
2002.....	6.9	10.3	18.8	32.0	49.5	75.9	112.9	167.9	235.5	344.3
2001.....	6.1	10.5	18.6	30.4	48.8	77.6	113.5	161.8	232.1	337.3
2000.....	6.3	9.7	17.3	29.5	48.7	73.8	111.6	164.0	234.1	337.4
1999.....	5.6	10.6	16.8	28.9	47.7	75.2	111.3	163.7	233.2	333.4
White, female										
2002.....	3.8	6.6	12.0	20.2	33.7	57.3	86.3	129.3	186.1	286.1
2001.....	3.4	6.7	12.0	20.4	36.3	57.4	85.4	129.7	190.1	293.0
2000.....	3.7	6.4	12.0	22.1	36.3	57.6	87.3	129.3	178.3	289.7
1999.....	3.9	6.1	11.0	20.4	36.6	59.6	86.7	125.4	181.7	285.1
Black, both sexes										
2002.....	12.0	21.4	40.7	67.8	105.5	161.6	221.6	300.3	392.7	537.8
2001.....	10.7	21.0	40.2	69.6	104.4	157.6	226.8	305.2	382.2	529.7
2000.....	10.1	21.9	40.7	66.3	102.4	159.2	223.7	316.5	404.9	524.9
1999.....	11.3	22.5	40.5	72.7	111.6	159.6	229.0	294.9	395.8	529.0
Black, male										
2002.....	14.9	25.4	46.4	76.5	118.8	165.3	223.3	295.8	386.7	417.0
2001.....	11.7	25.5	45.6	76.7	112.5	162.1	221.5	308.1	400.5	450.5
2000.....	11.8	25.5	43.5	72.1	108.5	154.3	224.3	311.4	382.7	452.1
1999.....	13.2	27.5	44.1	78.2	113.3	163.8	216.1	282.6	374.1	506.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)										
Black, female										
2002.....	9.4	17.9	35.8	60.6	95.1	158.9	220.4	303.0	395.9	584.1
2001.....	9.8	17.0	35.6	63.9	98.1	154.3	230.3	303.4	372.8	559.1
2000.....	8.7	18.7	38.4	61.5	97.6	162.8	223.3	319.6	416.1	552.0
1999.....	9.7	18.1	37.4	68.3	110.3	156.4	237.6	302.4	406.6	537.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Disorders of pituitary gland (E22-E23)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	0.2	0.4	0.5	1.1
2001.....	*	*	*	*	*	*	*	0.4	0.6	1.0
2000.....	*	*	*	*	*	0.2	0.3	0.3	0.7	1.1
1999.....	*	*	*	*	*	0.2	0.2	0.5	0.5	1.1

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	0.7	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	*	0.5	*	1.1
2001.....	*	*	*	*	*	*	*	*	0.6	1.1
2000.....	*	*	*	*	*	*	*	*	1.0	1.1
1999.....	*	*	*	*	*	*	*	*	*	1.2

White, both sexes

2002.....	*	*	*	*	*	*	*	0.4	0.4	1.1
2001.....	*	*	*	*	*	*	*	0.4	0.7	1.0
2000.....	*	*	*	*	*	*	0.3	0.3	0.8	1.1
1999.....	*	*	*	*	*	*	*	0.4	0.5	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of adrenal glands (E24-E27)										
All races 3/, both sexes										
2002.....	*	*	*	*	0.2	0.2	0.3	0.5	0.9	1.1
2001.....	*	*	*	*	*	*	0.5	0.7	1.0	1.8
2000.....	*	*	*	0.2	*	0.2	0.2	0.5	1.0	1.0
1999.....	*	*	0.1	*	*	0.3	0.3	0.5	0.4	1.0
All races 3/, male										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	0.7	1.2	2.1
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	0.4	0.6	0.9	1.1
2001.....	*	*	*	*	*	*	0.5	0.7	0.8	1.7
2000.....	*	*	*	*	*	*	*	0.5	1.0	1.1
1999.....	*	*	*	*	*	*	0.4	0.6	*	1.0
White, both sexes										
2002.....	*	*	*	*	*	*	0.3	0.5	0.9	1.1
2001.....	*	*	*	*	*	*	0.5	0.7	1.0	1.8
2000.....	*	*	*	0.2	*	*	*	0.5	1.0	1.1
1999.....	*	*	0.1	*	*	0.3	0.3	0.6	0.5	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other endocrine diseases (E15-E21,E28-E31,E34)										
All races 3/, both sexes										
2002.....	*	0.1	0.2	0.2	0.2	0.3	0.5	0.8	1.3	2.2
2001.....	*	*	0.1	0.1	0.3	0.2	0.5	0.7	1.0	2.5
2000.....	*	0.1	*	*	*	0.2	0.4	0.9	0.9	2.2
1999.....	*	*	0.2	0.2	0.2	0.2	0.6	0.7	1.2	1.9
All races 3/, male										
2002.....	*	*	*	*	*	*	*	0.7	1.1	1.8
2001.....	*	*	*	*	*	*	0.6	0.8	*	2.2
2000.....	*	*	*	*	*	*	*	0.9	*	1.7
1999.....	*	*	*	*	*	*	0.7	0.8	1.4	*
All races 3/, female										
2002.....	*	*	*	*	*	*	0.6	0.8	1.3	2.3
2001.....	*	*	*	*	*	*	*	0.7	1.2	2.6
2000.....	*	*	*	*	*	*	0.4	0.9	0.9	2.4
1999.....	*	*	*	*	*	*	0.5	0.6	1.1	2.0
White, both sexes										
2002.....	*	*	0.1	*	0.2	0.3	0.4	0.7	1.2	2.1
2001.....	*	*	*	*	0.2	*	0.3	0.6	0.9	2.4
2000.....	*	*	*	*	*	*	0.3	0.7	0.7	2.0
1999.....	*	*	*	*	*	*	0.5	0.6	1.1	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional diseases (E40-E68)										
All races 3/, both sexes										
2002.....	1.9	2.2	3.0	3.8	3.9	4.7	5.0	8.7	15.0	42.9
2001.....	1.4	2.1	2.8	3.1	3.5	3.9	5.4	8.1	14.5	44.5
2000.....	1.6	2.0	2.7	2.9	3.6	4.5	5.4	9.1	17.9	51.7
1999.....	1.5	1.7	2.3	2.8	2.9	3.8	5.6	8.6	17.9	55.3
All races 3/, male										
2002.....	2.2	2.2	3.1	3.7	3.6	4.3	4.9	8.2	13.5	37.8
2001.....	1.5	2.2	2.6	2.9	3.4	3.4	5.3	6.9	14.4	35.8
2000.....	1.7	2.2	2.8	3.3	3.2	4.0	5.3	8.5	15.8	44.6
1999.....	1.8	1.8	2.3	3.1	3.0	3.5	5.8	8.4	17.1	48.7
All races 3/, female										
2002.....	1.6	2.2	2.9	3.9	4.2	5.1	5.2	9.0	15.9	45.1
2001.....	1.3	2.0	3.0	3.3	3.7	4.3	5.6	9.0	14.6	48.1
2000.....	1.6	1.8	2.7	2.6	3.9	4.9	5.5	9.6	19.2	54.6
1999.....	1.3	1.6	2.4	2.4	2.9	4.0	5.5	8.8	18.4	58.0
White, both sexes										
2002.....	1.8	2.1	2.9	3.6	3.7	4.4	4.8	8.2	14.4	41.6
2001.....	1.2	1.9	2.6	3.0	3.4	3.6	5.0	7.6	13.6	43.8
2000.....	1.5	1.8	2.6	2.7	3.5	4.3	5.0	8.5	17.4	50.6
1999.....	1.4	1.5	2.2	2.6	2.8	3.6	4.9	8.0	16.8	54.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional diseases (E40-E68)										
White, male										
2002.....	2.1	2.3	3.2	3.6	3.5	4.1	4.6	8.0	12.5	37.1
2001.....	1.4	2.0	2.5	2.9	3.2	3.1	4.9	6.6	13.5	34.5
2000.....	1.6	2.2	2.8	3.1	3.0	3.8	4.9	7.9	15.3	43.7
1999.....	1.8	1.7	2.3	2.9	2.8	3.3	5.0	7.6	15.8	46.9
White, female										
2002.....	1.4	2.0	2.6	3.5	3.8	4.6	5.0	8.3	15.5	43.5
2001.....	1.1	1.7	2.7	3.1	3.6	4.1	5.1	8.3	13.8	47.7
2000.....	1.4	1.4	2.4	2.3	3.9	4.7	5.2	9.0	18.7	53.4
1999.....	1.1	1.4	2.2	2.2	2.7	3.8	4.8	8.3	17.4	57.2
Black, both sexes										
2002.....	3.4	3.7	5.2	7.5	7.0	8.7	8.8	16.3	24.9	62.1
2001.....	2.9	4.3	5.5	5.4	5.7	6.8	10.6	15.5	26.5	56.2
2000.....	3.0	3.8	4.9	6.2	5.3	7.5	10.4	16.9	26.5	68.5
1999.....	2.6	3.2	4.0	5.3	5.5	6.6	13.4	16.2	34.4	72.4
Black, male										
2002.....	3.7	2.6	3.8	6.7	5.7	7.0	8.7	12.4	29.8	48.2
2001.....	2.9	3.9	4.4	4.2	6.1	6.4	10.4	13.5	26.5	57.9
2000.....	2.8	2.8	4.0	6.4	5.6	6.5	12.0	18.0	25.3	65.7
1999.....	2.4	2.6	3.1	6.1	5.3	6.2	14.9	18.0	35.8	75.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional diseases (E40-E68)										
Black, female										
2002.....	3.2	4.6	6.4	8.2	8.1	10.0	9.0	18.7	22.3	67.4
2001.....	2.8	4.5	6.4	6.3	5.4	7.1	10.8	16.6	26.6	55.6
2000.....	3.1	4.7	5.6	6.1	5.1	8.3	9.3	16.2	27.2	69.5
1999.....	2.9	3.7	4.8	4.6	5.7	6.9	12.4	15.2	33.7	71.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)										
All races 3/, both sexes										
2002.....	0.1	0.2	0.3	0.5	0.7	1.4	2.5	5.6	12.2	38.4
2001.....	0.1	0.2	0.3	0.4	0.8	1.3	2.6	5.5	12.0	40.2
2000.....	0.1	0.2	0.3	0.5	0.9	1.9	2.8	6.4	14.6	46.3
1999.....	0.2	0.2	0.3	0.5	0.8	1.5	3.0	6.0	14.7	50.0
All races 3/, male										
2002.....	*	*	0.3	0.5	0.8	1.4	2.8	5.7	11.4	35.1
2001.....	*	0.2	0.3	0.5	0.8	1.3	2.6	5.1	12.6	33.6
2000.....	*	0.2	*	0.6	0.9	1.8	3.3	6.4	13.1	40.7
1999.....	*	0.2	0.3	0.6	1.1	1.5	3.6	6.2	14.8	44.4
All races 3/, female										
2002.....	*	*	0.3	0.4	0.6	1.4	2.2	5.6	12.6	39.8
2001.....	*	0.2	0.3	0.4	0.7	1.3	2.5	5.7	11.6	43.0
2000.....	*	0.2	0.3	0.3	0.9	1.9	2.4	6.5	15.5	48.6
1999.....	0.2	0.2	0.3	0.4	0.6	1.4	2.5	5.8	14.7	52.3
White, both sexes										
2002.....	0.1	0.2	0.3	0.4	0.6	1.3	2.4	5.4	11.7	37.0
2001.....	*	0.2	0.2	0.4	0.7	1.1	2.4	5.1	11.2	39.5
2000.....	*	0.2	0.2	0.4	0.8	1.8	2.4	6.1	14.2	45.2
1999.....	0.2	0.2	0.3	0.4	0.7	1.3	2.5	5.5	13.8	48.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)										
White, male										
2002.....	*	*	*	0.4	0.6	1.3	2.7	5.5	10.5	34.3
2001.....	*	*	*	0.4	0.7	1.0	2.3	4.7	11.7	32.2
2000.....	*	*	*	0.5	0.7	1.6	2.8	5.9	12.5	39.7
1999.....	*	*	0.3	0.4	0.9	1.2	2.8	5.6	13.4	42.2
White, female										
2002.....	*	*	0.3	0.4	0.6	1.2	2.1	5.3	12.4	38.2
2001.....	*	*	0.3	0.4	0.8	1.2	2.5	5.4	10.9	42.6
2000.....	*	*	0.3	*	0.9	1.9	2.1	6.2	15.1	47.4
1999.....	*	*	0.3	0.4	0.5	1.3	2.3	5.5	14.0	51.6
Black, both sexes										
2002.....	*	*	*	1.5	1.9	3.0	4.4	10.1	19.9	57.6
2001.....	*	*	*	*	*	2.8	4.0	10.1	22.2	51.6
2000.....	*	*	*	*	1.9	3.1	6.5	11.2	20.9	63.8
1999.....	*	*	*	1.6	2.0	3.2	8.0	11.1	28.9	66.0
Black, male										
2002.....	*	*	*	*	*	*	*	9.2	25.2	46.1
2001.....	*	*	*	*	*	*	*	11.2	24.1	55.6
2000.....	*	*	*	*	*	*	9.7	14.2	22.8	61.1
1999.....	*	*	*	*	*	*	12.2	13.3	33.1	73.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)										
Black, female										
2002.....	*	*	*	*	*	*	*	10.7	17.1	62.0
2001.....	*	*	*	*	*	*	*	9.4	21.3	50.1
2000.....	*	*	*	*	*	*	4.4	9.4	20.0	64.8
1999.....	*	*	*	*	*	*	5.2	9.7	26.8	63.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nutritional marasmus (E41)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*	*	0.6	2.4
2001.....	*	*	*	*	*	*	*	*	0.6	2.5
2000.....	*	*	*	*	*	*	*	*	0.8	2.7
1999.....	*	*	*	*	*	*	*	0.3	0.8	2.5

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	2.2
2000.....	*	*	*	*	*	*	*	*	*	2.9
1999.....	*	*	*	*	*	*	*	*	*	3.0

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	0.6	2.7
2001.....	*	*	*	*	*	*	*	*	*	2.6
2000.....	*	*	*	*	*	*	*	*	0.9	2.7
1999.....	*	*	*	*	*	*	*	*	0.6	2.2

White, both sexes

2002.....	*	*	*	*	*	*	*	*	0.6	2.4
2001.....	*	*	*	*	*	*	*	*	0.6	2.5
2000.....	*	*	*	*	*	*	*	*	0.9	2.7
1999.....	*	*	*	*	*	*	*	*	0.8	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified malnutrition (E40,E42-E46)										
All races 3/, both sexes										
2002.....	0.1	0.2	0.3	0.5	0.7	1.4	2.4	5.4	11.6	36.0
2001.....	0.1	0.2	0.3	0.4	0.7	1.2	2.4	5.2	11.4	37.7
2000.....	0.1	0.2	0.2	0.4	0.9	1.8	2.7	6.2	13.8	43.6
1999.....	0.2	0.2	0.3	0.5	0.8	1.4	2.8	5.7	14.0	47.5
All races 3/, male										
2002.....	*	*	0.2	0.5	0.8	1.4	2.8	5.6	10.8	33.5
2001.....	*	0.2	0.3	0.5	0.7	1.2	2.5	4.8	11.9	31.3
2000.....	*	0.2	*	0.6	0.9	1.7	3.2	6.1	12.4	37.8
1999.....	*	0.2	0.3	0.6	1.0	1.5	3.3	6.0	13.8	41.4
All races 3/, female										
2002.....	*	*	0.3	0.4	0.6	1.3	2.1	5.3	12.0	37.1
2001.....	*	0.2	0.3	0.4	0.7	1.2	2.4	5.5	11.1	40.4
2000.....	*	0.2	0.3	0.3	0.8	1.9	2.3	6.3	14.6	45.9
1999.....	0.2	*	0.3	0.4	0.6	1.4	2.4	5.5	14.1	50.0
White, both sexes										
2002.....	0.1	0.2	0.2	0.4	0.6	1.2	2.3	5.2	11.1	34.7
2001.....	*	0.2	0.2	0.4	0.7	1.1	2.3	4.8	10.6	37.0
2000.....	*	0.2	0.2	0.4	0.7	1.7	2.3	5.9	13.3	42.5
1999.....	0.2	0.2	0.3	0.3	0.7	1.2	2.3	5.4	12.9	46.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified malnutrition (E40,E42-E46)

White, male

2002.....	*	*	*	0.4	0.6	1.3	2.6	5.4	10.0	32.5
2001.....	*	*	*	0.4	0.6	1.0	2.3	4.4	11.0	30.1
2000.....	*	*	*	0.4	0.7	1.6	2.6	5.7	11.7	37.0
1999.....	*	*	0.3	*	0.9	1.2	2.6	5.4	12.5	39.4

White, female

2002.....	*	*	0.3	0.4	0.6	1.2	2.0	5.0	11.8	35.6
2001.....	*	*	0.3	0.3	0.8	1.2	2.4	5.1	10.4	39.9
2000.....	*	*	*	*	0.8	1.8	2.0	6.0	14.2	44.7
1999.....	*	*	0.3	0.3	0.5	1.3	2.1	5.3	13.2	49.4

Black, both sexes

2002.....	*	*	*	1.4	1.8	2.9	4.3	9.3	18.8	55.8
2001.....	*	*	*	*	*	2.7	3.9	10.1	20.9	49.1
2000.....	*	*	*	*	1.9	3.0	6.3	10.7	20.9	59.7
1999.....	*	*	*	1.5	1.9	3.1	7.4	9.6	28.6	62.8

Black, male

2002.....	*	*	*	*	*	*	*	*	22.9	45.0
2001.....	*	*	*	*	*	*	*	11.2	23.3	53.3
2000.....	*	*	*	*	*	*	9.4	12.8	22.8	55.4
1999.....	*	*	*	*	*	*	11.5	12.8	32.3	69.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified malnutrition (E40,E42-E46)										
Black, female										
2002.....	*	*	*	*	*	*	*	9.6	16.7	60.0
2001.....	*	*	*	*	*	*	*	9.4	19.6	47.6
2000.....	*	*	*	*	*	*	*	9.4	20.0	61.4
1999.....	*	*	*	*	*	*	4.7	7.7	26.8	60.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other nutritional deficiencies (E50-E64)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	*	*	0.4	0.6	3.1
2001.....	*	*	*	*	*	*	*	0.4	0.8	2.7
2000.....	*	*	*	*	*	*	0.3	0.4	1.2	3.7
1999.....	*	*	*	*	*	*	*	0.4	1.1	3.9
All races 3/, male										
2002.....	*	*	*	*	*	*	*	*	*	1.9
2001.....	*	*	*	*	*	*	*	*	*	1.6
2000.....	*	*	*	*	*	*	*	*	1.3	3.0
1999.....	*	*	*	*	*	*	*	*	1.3	3.9
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	0.6	3.6
2001.....	*	*	*	*	*	*	*	*	1.0	3.1
2000.....	*	*	*	*	*	*	*	*	1.1	4.0
1999.....	*	*	*	*	*	*	*	*	1.1	3.9
White, both sexes										
2002.....	*	*	*	*	*	*	*	0.5	0.6	3.1
2001.....	*	*	*	*	*	*	*	0.4	0.8	2.7
2000.....	*	*	*	*	*	*	0.3	0.3	1.2	3.7
1999.....	*	*	*	*	*	*	*	0.4	1.1	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Obesity and other hyperalimentation (E65-E68)										
All races 3/, both sexes										
2002.....	1.8	2.0	2.7	3.3	3.1	3.1	2.4	2.6	2.2	1.4
2001.....	1.3	1.9	2.5	2.7	2.7	2.5	2.7	2.2	1.7	1.6
2000.....	1.5	1.7	2.4	2.4	2.6	2.5	2.4	2.3	2.2	1.7
1999.....	1.3	1.5	2.0	2.2	2.1	2.2	2.4	2.2	2.0	1.4
All races 3/, male										
2002.....	2.1	2.0	2.8	3.2	2.7	2.7	1.9	1.9	1.5	*
2001.....	1.4	1.9	2.3	2.4	2.5	1.9	2.4	1.3	1.2	*
2000.....	1.6	1.9	2.5	2.6	2.2	2.0	1.9	1.7	1.4	*
1999.....	1.6	1.5	1.9	2.5	1.9	1.9	1.9	1.6	*	*
All races 3/, female										
2002.....	1.5	2.1	2.6	3.4	3.5	3.5	2.8	3.1	2.6	1.6
2001.....	1.1	1.8	2.6	2.9	3.0	3.0	2.9	2.9	2.0	2.0
2000.....	1.4	1.6	2.3	2.2	3.0	2.9	2.8	2.7	2.6	2.0
1999.....	1.1	1.4	2.0	1.9	2.2	2.5	2.8	2.6	2.6	1.9
White, both sexes										
2002.....	1.6	2.0	2.6	3.1	3.0	3.0	2.3	2.3	2.1	1.4
2001.....	1.1	1.7	2.3	2.6	2.6	2.4	2.4	2.1	1.7	1.6
2000.....	1.4	1.6	2.3	2.2	2.6	2.3	2.3	2.2	2.1	1.7
1999.....	1.2	1.4	1.9	2.2	2.0	2.2	2.2	2.0	2.0	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Obesity and other hyperalimentation (E65-E68)										
Black, female										
2002.....	3.0	4.5	6.0	7.3	7.1	6.5	*	7.7	*	*
2001.....	2.6	4.5	5.7	5.6	5.0	5.3	7.8	6.4	*	*
2000.....	2.8	4.4	4.9	5.7	3.8	5.6	4.9	6.0	*	*
1999.....	2.5	2.8	4.3	3.5	4.2	4.6	7.2	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Metabolic disorders (E70-E88)										
All races 3/, both sexes										
2002.....	1.0	1.5	2.6	4.6	6.9	9.7	16.6	26.5	47.3	114.3
2001.....	1.0	1.7	2.7	4.1	6.3	10.1	16.4	26.3	45.2	124.4
2000.....	1.0	1.5	2.5	4.0	6.5	9.8	15.7	25.7	46.1	124.1
1999.....	0.9	1.3	2.3	3.5	6.2	9.5	15.1	25.3	44.7	130.8
All races 3/, male										
2002.....	1.3	2.0	3.7	6.1	8.9	12.5	20.0	30.4	51.5	101.5
2001.....	1.2	2.3	3.5	5.2	8.8	13.7	20.0	32.6	48.0	109.8
2000.....	1.2	2.2	3.6	5.3	8.9	12.8	19.2	29.6	48.6	112.0
1999.....	1.1	1.8	3.1	4.7	7.8	11.6	19.0	29.2	46.9	110.5
All races 3/, female										
2002.....	0.8	1.1	1.6	3.2	5.0	7.2	13.9	23.8	44.7	119.9
2001.....	0.7	1.1	1.9	3.0	3.9	7.0	13.6	22.0	43.5	130.6
2000.....	0.8	0.8	1.4	2.8	4.3	7.2	12.9	22.9	44.7	129.1
1999.....	0.7	0.9	1.7	2.4	4.7	7.7	12.1	22.7	43.4	139.0
White, both sexes										
2002.....	1.0	1.5	2.6	4.5	6.6	9.4	16.0	26.1	46.3	114.0
2001.....	1.0	1.5	2.7	4.0	6.1	9.9	16.3	26.0	44.4	123.4
2000.....	0.9	1.5	2.4	3.9	6.3	9.7	15.5	25.4	45.6	123.6
1999.....	0.8	1.3	2.2	3.5	6.0	9.1	14.8	25.0	44.2	131.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Metabolic disorders (E70-E88)										
White, male										
2002.....	1.3	1.8	3.7	6.0	8.4	12.1	19.6	29.8	50.6	100.2
2001.....	1.2	2.1	3.5	5.1	8.4	13.5	19.7	32.2	46.9	109.2
2000.....	1.1	2.2	3.5	5.2	8.6	12.6	19.1	29.1	48.1	113.9
1999.....	1.0	1.7	3.0	4.7	7.8	11.2	18.6	28.6	45.7	108.4
White, female										
2002.....	0.7	1.1	1.5	3.0	5.0	6.9	13.0	23.4	43.7	120.0
2001.....	0.8	1.0	1.9	3.0	3.8	6.7	13.6	21.6	42.9	129.4
2000.....	0.7	0.8	1.2	2.7	4.2	7.1	12.6	22.8	44.1	127.6
1999.....	0.6	0.9	1.4	2.4	4.4	7.4	11.9	22.5	43.3	140.4
Black, both sexes										
2002.....	1.3	2.5	3.8	6.5	10.8	13.9	26.0	34.7	66.5	135.1
2001.....	0.9	2.8	3.6	5.1	9.4	13.8	21.5	36.0	62.3	151.4
2000.....	1.4	2.1	3.6	5.8	9.0	13.3	20.1	33.2	61.0	147.0
1999.....	1.8	1.9	4.3	4.6	8.7	14.9	21.0	32.7	54.0	138.7
Black, male										
2002.....	1.5	3.5	5.0	8.5	16.4	17.7	27.4	40.9	71.8	134.0
2001.....	*	3.7	4.3	6.9	13.6	18.8	28.6	44.6	75.4	130.5
2000.....	1.7	2.9	5.1	7.1	12.2	16.7	24.4	41.1	65.8	115.3
1999.....	2.0	2.5	4.8	6.3	10.0	18.7	26.1	40.8	61.9	147.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Metabolic disorders (E70-E88)										
Black, female										
2002.....	*	1.8	2.8	4.9	6.3	11.1	25.1	31.0	63.7	135.5
2001.....	*	2.1	2.9	3.6	6.1	10.1	16.8	30.8	55.6	159.1
2000.....	*	*	2.4	4.7	6.5	10.8	17.3	28.5	58.6	158.8
1999.....	1.6	*	3.9	3.3	7.7	12.1	17.6	27.7	50.1	135.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)										
All races 3/, both sexes										
2002.....	0.3	0.5	0.6	1.0	1.6	2.5	5.2	9.8	24.7	83.2
2001.....	0.3	0.5	0.6	0.8	1.7	2.6	5.6	11.0	25.1	93.9
2000.....	0.4	0.4	0.6	1.0	1.7	3.0	5.2	11.3	26.6	96.8
1999.....	0.3	0.4	0.7	0.8	1.4	3.0	5.5	11.9	27.7	106.4
All races 3/, male										
2002.....	0.3	0.5	0.6	1.0	1.7	2.6	5.2	9.6	24.1	67.9
2001.....	0.3	0.5	0.6	0.8	2.0	2.8	5.5	12.2	24.0	76.5
2000.....	0.4	0.5	0.7	1.2	1.9	3.4	5.8	12.0	27.2	84.3
1999.....	0.4	0.5	0.8	1.1	1.4	3.1	6.2	13.2	27.2	85.0
All races 3/, female										
2002.....	0.2	0.4	0.5	1.0	1.5	2.3	5.2	10.0	25.0	89.8
2001.....	0.3	0.4	0.6	0.8	1.4	2.5	5.7	10.2	25.7	101.2
2000.....	0.3	0.3	0.5	0.9	1.5	2.6	4.7	10.8	26.2	101.9
1999.....	0.2	0.3	0.5	0.7	1.5	2.9	5.0	11.1	27.9	115.1
White, both sexes										
2002.....	0.2	0.4	0.5	0.9	1.4	2.2	4.7	9.2	23.5	82.4
2001.....	0.3	0.3	0.5	0.7	1.4	2.4	5.1	10.4	24.0	93.0
2000.....	0.3	0.3	0.5	0.9	1.5	2.7	4.8	10.9	25.9	96.0
1999.....	0.2	0.4	0.5	0.7	1.2	2.5	5.2	11.4	27.2	106.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)										
White, male										
2002.....	0.2	0.4	0.5	0.8	1.5	2.2	4.8	8.9	22.7	66.1
2001.....	0.2	0.4	0.5	0.6	1.6	2.6	4.7	11.5	22.3	75.8
2000.....	0.3	0.4	0.6	1.0	1.6	3.0	5.3	11.2	26.3	84.9
1999.....	0.2	0.5	0.7	0.9	1.1	2.3	5.6	12.3	26.0	82.4
White, female										
2002.....	*	0.4	0.5	1.0	1.4	2.2	4.7	9.4	24.0	89.5
2001.....	0.3	0.3	0.5	0.7	1.3	2.1	5.4	9.7	25.0	100.2
2000.....	0.3	0.3	0.4	0.8	1.4	2.3	4.4	10.7	25.7	100.5
1999.....	*	0.3	0.4	0.6	1.2	2.6	4.8	10.8	27.9	116.4
Black, both sexes										
2002.....	0.8	1.2	1.3	2.0	3.1	5.4	11.4	18.9	42.9	104.2
2001.....	*	1.4	1.6	1.8	4.2	5.5	12.1	20.7	42.8	119.4
2000.....	0.8	1.3	1.6	2.2	3.4	6.6	10.1	18.5	40.9	119.8
1999.....	1.1	1.0	1.9	2.0	4.0	7.8	11.0	19.8	36.7	115.1
Black, male										
2002.....	*	1.8	*	*	4.4	7.0	11.3	20.7	46.6	99.7
2001.....	*	*	*	*	5.9	5.8	15.8	24.2	54.6	97.6
2000.....	*	*	*	*	4.6	8.1	12.0	26.0	47.2	98.0
1999.....	1.5	*	*	*	*	11.4	14.9	26.6	44.5	125.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

Black, female

2002.....	*	*	*	*	*	4.3	11.4	17.8	41.0	106.0
2001.....	*	*	*	*	*	5.3	9.6	18.6	36.8	127.6
2000.....	*	*	*	*	*	5.6	8.9	14.0	37.8	127.9
1999.....	*	*	*	*	4.0	5.2	8.3	15.7	32.8	111.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other metabolic disorders (E70-E83,E85,E88)										
All races 3/, both sexes										
2002.....	0.6	1.0	2.0	3.6	5.2	7.2	11.4	16.7	22.6	31.1
2001.....	0.5	1.1	2.0	3.2	4.6	7.4	10.8	15.3	20.0	30.5
2000.....	0.4	1.0	1.9	2.9	4.8	6.8	10.5	14.3	19.6	27.3
1999.....	0.5	0.9	1.6	2.6	4.7	6.5	9.5	13.4	17.0	24.3
All races 3/, male										
2002.....	0.9	1.4	3.1	5.1	7.1	9.8	14.7	20.8	27.3	33.6
2001.....	0.7	1.6	2.8	4.4	6.8	10.9	14.5	20.3	23.8	33.3
2000.....	0.6	1.5	2.9	4.0	6.9	9.3	13.4	17.6	21.4	27.7
1999.....	0.6	1.3	2.2	3.6	6.4	8.6	12.7	15.9	19.6	25.5
All races 3/, female										
2002.....	0.4	0.7	1.0	2.2	3.5	4.9	8.7	13.8	19.7	30.0
2001.....	0.4	0.6	1.2	2.1	2.6	4.5	7.8	11.7	17.8	29.3
2000.....	0.3	0.5	0.9	1.9	2.8	4.6	8.2	12.1	18.5	27.1
1999.....	0.4	0.5	1.0	1.7	3.2	4.8	7.1	11.6	15.5	23.9
White, both sexes										
2002.....	0.6	1.0	2.0	3.6	5.2	7.2	11.2	16.9	22.7	31.6
2001.....	0.6	1.1	2.1	3.3	4.6	7.5	11.2	15.5	20.3	30.4
2000.....	0.4	1.1	1.9	3.0	4.8	7.0	10.7	14.5	19.7	27.5
1999.....	0.5	0.9	1.6	2.7	4.8	6.7	9.7	13.6	16.9	24.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other metabolic disorders (E70-E83,E85,E88)										
White, male										
2002.....	0.9	1.4	3.2	5.2	6.9	9.9	14.8	20.9	27.8	34.1
2001.....	0.8	1.6	2.9	4.4	6.8	10.9	15.0	20.7	24.4	33.3
2000.....	0.6	1.7	2.9	4.1	6.9	9.5	13.8	18.0	21.8	29.0
1999.....	0.6	1.2	2.2	3.7	6.6	8.9	13.0	16.2	19.6	25.9
White, female										
2002.....	0.4	0.7	0.9	2.1	3.6	4.7	8.3	14.0	19.6	30.4
2001.....	0.4	0.6	1.3	2.2	2.6	4.5	8.1	11.8	17.9	29.1
2000.....	0.2	0.5	0.9	1.8	2.8	4.7	8.2	12.1	18.4	26.9
1999.....	0.3	0.5	0.9	1.7	3.2	4.7	7.0	11.8	15.4	24.0
Black, both sexes										
2002.....	*	1.3	2.4	4.4	7.6	8.5	14.5	15.8	23.5	30.9
2001.....	*	1.5	1.9	3.2	5.2	8.3	9.4	15.3	19.5	31.9
2000.....	*	*	2.0	3.6	5.7	6.6	10.0	14.7	20.0	27.2
1999.....	0.7	0.9	2.3	2.7	4.7	7.1	10.0	12.9	17.3	23.6
Black, male										
2002.....	*	1.7	3.3	5.8	12.1	10.7	15.7	20.2	25.2	34.3
2001.....	*	2.1	2.5	4.7	7.7	13.0	12.8	20.4	20.9	32.9
2000.....	*	*	3.1	4.0	7.7	8.6	12.4	15.1	18.5	*
1999.....	*	*	2.5	*	5.9	7.3	11.2	14.2	17.4	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other metabolic disorders (E70-E83,E85,E88)										
Black, female										
2002.....	*	*	*	3.3	4.0	6.8	13.7	13.2	22.7	29.6
2001.....	*	*	*	*	3.2	4.9	7.2	12.2	18.8	31.6
2000.....	*	*	*	3.2	4.1	5.2	8.4	14.5	20.8	30.9
1999.....	*	*	2.2	*	3.7	6.9	9.2	12.0	17.3	24.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
V. Mental and behavioral disorders (F01-F99)										
All races 3/, both sexes										
2002.....	6.5	7.9	8.3	8.3	9.5	13.0	25.6	64.2	168.2	646.7
2001.....	6.0	7.9	7.7	7.7	9.4	13.0	23.8	60.6	154.3	604.8
2000.....	5.8	7.5	7.1	7.8	8.6	11.7	22.5	54.1	143.7	561.6
1999.....	6.1	7.4	6.8	7.3	8.9	11.2	21.2	49.9	128.6	515.3
All races 3/, male										
2002.....	9.3	12.0	13.4	12.9	14.4	18.4	30.2	68.2	166.1	506.0
2001.....	8.7	12.2	12.3	11.7	14.3	18.1	29.3	63.3	151.3	488.0
2000.....	8.5	11.6	11.4	11.9	13.0	16.5	28.0	59.4	140.5	451.0
1999.....	9.4	11.5	11.1	11.7	13.8	16.1	26.9	56.9	133.4	415.8
All races 3/, female										
2002.....	3.6	4.0	3.3	4.0	5.1	8.4	21.9	61.3	169.4	707.8
2001.....	3.3	3.8	3.3	3.9	4.9	8.6	19.5	58.8	156.0	653.7
2000.....	3.2	3.5	3.1	4.0	4.6	7.7	18.3	50.4	145.6	606.6
1999.....	2.8	3.4	2.8	3.3	4.5	7.0	16.7	45.1	125.8	555.4
White, both sexes										
2002.....	6.2	7.5	7.6	7.7	8.8	12.5	25.3	64.1	170.5	663.8
2001.....	5.5	7.1	6.8	6.9	8.9	12.5	23.5	61.1	157.4	621.6
2000.....	5.4	6.7	6.6	7.2	7.9	11.2	22.0	54.0	145.3	577.6
1999.....	5.5	6.5	6.1	6.5	8.3	10.7	20.7	50.0	130.3	534.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
V. Mental and behavioral disorders (F01-F99)										
White, male										
2002.....	9.2	11.3	12.3	11.6	13.0	16.9	29.0	66.5	167.4	514.9
2001.....	8.0	10.9	10.8	10.2	13.2	16.8	28.3	63.1	152.8	498.4
2000.....	7.9	10.3	10.4	10.8	11.6	15.2	26.7	58.9	140.3	459.4
1999.....	8.6	10.2	9.8	10.1	12.8	14.9	25.7	55.8	132.9	429.2
White, female										
2002.....	3.3	3.7	3.1	4.0	4.9	8.7	22.3	62.4	172.4	728.4
2001.....	2.9	3.4	2.9	3.8	4.9	8.7	19.6	59.7	160.2	673.1
2000.....	2.9	3.1	2.8	3.8	4.4	7.7	18.3	50.6	148.3	625.6
1999.....	2.3	2.8	2.4	3.1	4.2	6.9	16.7	45.9	128.8	576.9
Black, both sexes										
2002.....	9.5	12.6	15.0	15.9	16.8	19.8	34.5	78.3	173.2	540.8
2001.....	10.3	15.3	16.2	15.4	15.9	20.7	33.5	70.5	147.9	488.5
2000.....	9.4	15.2	13.7	14.2	16.2	18.7	33.4	68.4	145.4	442.6
1999.....	10.5	15.1	13.9	16.5	15.4	18.8	30.8	57.8	128.3	349.1
Black, male										
2002.....	12.8	19.5	25.7	28.0	28.9	35.4	51.1	103.3	189.5	457.8
2001.....	14.9	24.5	27.3	27.4	29.1	34.6	47.8	84.6	171.8	423.2
2000.....	13.6	24.4	23.6	24.4	27.7	32.5	48.8	82.2	169.4	415.2
1999.....	15.4	23.1	23.2	29.5	26.5	31.4	46.5	79.8	168.3	309.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
V. Mental and behavioral disorders (F01-F99)										
Black, female										
2002.....	6.7	6.7	5.8	6.0	7.3	8.1	23.3	63.4	164.7	572.6
2001.....	6.2	7.4	6.7	5.5	5.6	10.5	24.0	62.1	135.7	512.8
2000.....	5.6	7.2	5.4	5.9	7.0	8.4	23.1	60.1	133.3	452.9
1999.....	6.2	8.1	6.1	6.0	6.7	9.4	20.5	44.6	108.5	364.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Organic dementia (F01,F03)										
All races 3/, both sexes										
2002.....	*	0.1	0.2	0.5	1.8	5.4	17.0	53.8	150.9	598.3
2001.....	*	*	0.2	0.6	1.8	5.5	15.9	49.5	135.8	554.0
2000.....	*	0.1	0.1	0.6	1.4	4.5	14.4	43.0	125.3	505.6
1999.....	*	*	0.2	0.5	1.4	4.0	13.0	37.8	108.1	454.4
All races 3/, male										
2002.....	*	*	*	0.5	1.9	6.2	18.7	55.1	147.0	465.7
2001.....	*	*	*	0.5	2.0	6.2	17.5	49.7	130.8	442.4
2000.....	*	*	*	0.5	1.7	5.2	16.3	45.4	120.5	398.9
1999.....	*	*	*	0.6	1.7	4.9	15.3	40.5	109.7	361.7
All races 3/, female										
2002.....	*	*	*	0.6	1.6	4.6	15.7	52.9	153.3	655.9
2001.....	*	*	*	0.6	1.5	4.9	14.5	49.4	138.8	600.8
2000.....	*	*	*	0.7	1.2	4.0	12.8	41.4	128.2	549.1
1999.....	*	*	*	*	1.1	3.2	11.3	35.9	107.1	491.7
White, both sexes										
2002.....	*	*	0.2	0.5	1.8	5.2	16.9	53.6	153.0	613.9
2001.....	*	*	0.2	0.6	1.8	5.4	15.6	49.9	138.3	569.1
2000.....	*	*	0.1	0.6	1.3	4.4	13.9	42.9	126.6	519.6
1999.....	*	*	0.2	0.5	1.4	3.8	12.9	37.9	109.6	471.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Organic dementia (F01,F03)										
White, male										
2002.....	*	*	*	0.5	1.9	5.8	18.2	53.7	148.1	473.7
2001.....	*	*	*	0.5	2.0	5.9	16.9	49.8	132.0	452.2
2000.....	*	*	*	0.5	1.6	4.9	15.4	45.2	120.2	406.1
1999.....	*	*	*	0.6	1.7	4.8	14.9	40.2	109.4	373.7
White, female										
2002.....	*	*	*	0.6	1.7	4.8	15.9	53.6	155.9	674.7
2001.....	*	*	*	0.6	1.6	5.0	14.5	50.0	142.2	618.0
2000.....	*	*	*	0.7	1.1	3.9	12.6	41.3	130.4	565.7
1999.....	*	*	*	*	1.1	3.0	11.2	36.3	109.7	510.4
Black, both sexes										
2002.....	*	*	*	*	2.2	7.8	22.4	66.8	157.3	503.1
2001.....	*	*	*	*	2.2	8.2	23.0	57.5	132.7	452.3
2000.....	*	*	*	*	2.5	6.4	22.5	55.3	127.6	403.2
1999.....	*	*	*	*	*	6.0	17.7	42.8	107.5	309.3
Black, male										
2002.....	*	*	*	*	*	11.0	28.7	84.0	169.7	425.6
2001.....	*	*	*	*	*	11.2	29.3	62.7	150.9	382.4
2000.....	*	*	*	*	*	9.1	29.4	61.4	145.8	372.5
1999.....	*	*	*	*	*	7.5	23.1	51.3	136.0	266.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Organic dementia (F01,F03)										
Black, female										
2002.....	*	*	*	*	*	5.4	18.1	56.5	150.8	532.7
2001.....	*	*	*	*	*	6.0	18.8	54.4	123.5	478.2
2000.....	*	*	*	*	*	4.4	18.0	51.6	118.5	414.7
1999.....	*	*	*	*	*	4.8	14.2	37.8	93.3	325.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified organic mental disorders (F04-F09)

All races 3/, both sexes

2002.....	*	*	*	0.1	0.2	0.5	1.2	2.4	6.5	26.7
2001.....	*	*	*	*	0.2	0.5	1.1	2.9	7.8	29.3
2000.....	*	*	*	0.2	0.3	0.5	1.2	3.2	7.5	33.7
1999.....	*	*	*	*	0.2	0.4	1.2	3.3	8.7	37.7

All races 3/, male

2002.....	*	*	*	*	*	0.7	1.3	2.4	6.9	20.1
2001.....	*	*	*	*	*	0.7	1.3	3.0	7.7	23.8
2000.....	*	*	*	*	*	0.6	1.4	3.4	6.4	28.4
1999.....	*	*	*	*	*	0.5	1.3	3.6	9.0	29.8

All races 3/, female

2002.....	*	*	*	*	*	*	1.1	2.4	6.4	29.6
2001.....	*	*	*	*	*	*	1.0	2.9	7.9	31.6
2000.....	*	*	*	*	*	*	1.1	3.1	8.2	35.8
1999.....	*	*	*	*	*	*	1.1	3.1	8.6	40.9

White, both sexes

2002.....	*	*	*	*	*	0.4	1.1	2.5	6.7	27.5
2001.....	*	*	*	*	*	0.5	1.1	2.9	8.0	30.2
2000.....	*	*	*	0.2	0.2	0.5	1.3	3.3	7.5	34.7
1999.....	*	*	*	*	0.2	0.4	1.1	3.2	8.8	38.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified organic mental disorders (F04-F09)

Black, female

2002.....	*	*	*	*	*	*	*	*	*	25.1
2001.....	*	*	*	*	*	*	*	*	*	24.8
2000.....	*	*	*	*	*	*	*	*	9.8	25.3
1999.....	*	*	*	*	*	*	*	*	8.6	28.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mental and behavioral disorders due to psychoactive substance use (F10-F19)										
All races 3/, both sexes										
2002.....	6.0	7.3	7.6	6.9	6.7	6.1	5.3	4.7	3.8	3.7
2001.....	5.5	7.4	7.0	6.4	6.6	5.7	5.1	4.6	4.5	4.0
2000.....	5.5	7.0	6.5	6.4	6.0	5.8	5.1	4.7	4.5	4.1
1999.....	5.7	6.9	6.2	6.2	6.6	5.8	5.0	5.3	4.4	3.8
All races 3/, male										
2002.....	8.8	11.2	12.6	11.5	11.2	10.4	8.1	7.8	6.2	7.0
2001.....	8.2	11.6	11.5	10.4	11.1	9.7	8.6	7.2	7.3	7.5
2000.....	8.1	11.1	10.7	10.4	10.1	9.7	8.3	8.0	7.9	7.3
1999.....	9.0	11.0	10.2	10.2	11.1	9.5	8.1	9.0	8.4	7.4
All races 3/, female										
2002.....	3.2	3.4	2.8	2.6	2.5	2.4	3.1	2.5	2.4	2.2
2001.....	2.9	3.4	2.8	2.7	2.6	2.3	2.3	2.9	2.9	2.5
2000.....	2.9	3.0	2.5	2.7	2.3	2.4	2.6	2.5	2.6	2.8
1999.....	2.5	3.1	2.3	2.5	2.5	2.6	2.5	2.7	2.0	2.4
White, both sexes										
2002.....	5.8	6.8	6.9	6.3	5.9	5.8	5.1	4.7	3.8	3.6
2001.....	5.0	6.7	6.2	5.7	6.1	5.3	5.0	4.6	4.6	4.0
2000.....	5.0	6.2	5.9	5.8	5.4	5.3	5.0	4.7	4.6	4.1
1999.....	5.1	6.0	5.4	5.4	6.0	5.4	4.7	5.1	4.3	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mental and behavioral disorders due to psychoactive substance use (F10-F19)										
White, male										
2002.....	8.7	10.6	11.5	10.2	9.8	9.4	7.4	7.7	5.9	6.8
2001.....	7.5	10.4	10.1	9.0	10.0	8.7	8.1	6.9	7.2	7.3
2000.....	7.6	9.8	9.6	9.3	9.0	8.7	7.9	7.6	7.8	7.2
1999.....	8.2	9.7	9.0	8.6	10.1	8.4	7.4	8.4	8.3	7.4
White, female										
2002.....	2.8	3.1	2.5	2.5	2.3	2.5	3.2	2.6	2.5	2.2
2001.....	2.6	3.0	2.3	2.6	2.5	2.4	2.4	3.0	2.9	2.6
2000.....	2.5	2.5	2.2	2.4	2.2	2.4	2.6	2.6	2.7	2.9
1999.....	2.0	2.5	1.8	2.3	2.3	2.7	2.5	2.7	2.0	2.5
Black, both sexes										
2002.....	8.6	11.9	14.2	14.3	13.5	9.6	8.3	5.8	*	*
2001.....	9.6	14.5	15.0	13.2	12.7	10.0	7.4	6.4	*	*
2000.....	9.0	14.6	13.1	12.9	12.4	11.0	7.5	6.9	*	*
1999.....	10.2	14.3	13.1	14.9	12.4	10.8	9.5	8.0	6.1	*
Black, male										
2002.....	11.5	18.5	24.8	26.5	25.3	20.7	16.7	11.5	*	*
2001.....	14.0	23.1	25.3	24.2	24.8	20.6	15.1	13.9	*	*
2000.....	12.7	23.6	22.9	23.1	22.8	21.6	15.0	15.6	*	*
1999.....	15.0	22.3	21.6	27.6	22.9	21.6	19.0	16.6	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

Black, female

2002.....	6.1	6.1	5.3	4.3	4.2	*	*	*	*	*
2001.....	5.7	7.0	6.3	4.1	3.2	*	*	*	*	*
2000.....	5.6	6.8	4.8	4.6	4.2	*	*	*	*	*
1999.....	5.9	7.5	5.8	4.6	4.2	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mental and behavioral disorders due to use of alcohol (F10)										
All races 3/, both sexes										
2002.....	3.9	5.3	6.0	5.9	5.8	5.1	4.1	3.5	2.4	2.0
2001.....	3.7	5.4	5.6	5.5	5.9	5.1	4.1	3.3	3.0	2.1
2000.....	3.6	5.1	5.2	5.5	5.3	5.0	4.1	3.4	2.9	2.5
1999.....	3.6	5.1	5.0	5.3	5.7	4.9	3.8	3.8	2.9	2.3
All races 3/, male										
2002.....	6.1	8.3	10.1	9.9	10.0	9.0	6.8	6.4	4.5	4.4
2001.....	5.7	8.6	9.2	9.0	10.0	8.8	7.3	5.8	5.4	4.7
2000.....	5.7	8.3	8.6	9.2	9.1	8.8	7.2	6.1	5.9	5.2
1999.....	5.9	8.1	8.4	8.9	9.7	8.4	6.7	7.2	6.2	4.9
All races 3/, female										
2002.....	1.9	2.3	2.1	2.0	1.9	1.6	1.9	1.4	1.1	1.0
2001.....	1.8	2.3	2.1	2.2	2.1	1.9	1.5	1.6	1.6	1.1
2000.....	1.6	2.0	1.9	2.1	1.8	1.8	1.6	1.5	1.2	1.5
1999.....	1.5	2.1	1.7	2.0	2.1	2.0	1.6	1.5	1.0	1.3
White, both sexes										
2002.....	4.0	5.3	5.8	5.4	5.1	4.7	3.8	3.4	2.3	2.0
2001.....	3.5	5.2	5.1	4.9	5.5	4.6	3.9	3.2	3.0	2.1
2000.....	3.5	4.7	4.9	5.1	4.7	4.5	3.9	3.2	2.9	2.5
1999.....	3.5	4.7	4.5	4.7	5.2	4.5	3.5	3.6	2.8	2.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mental and behavioral disorders due to use of alcohol (F10)										
White, male										
2002.....	6.2	8.2	9.8	9.0	8.9	8.2	6.0	6.3	4.2	4.1
2001.....	5.3	8.2	8.4	7.9	9.1	7.7	6.8	5.5	5.4	4.4
2000.....	5.5	7.7	8.0	8.3	8.0	7.8	6.7	5.6	5.7	5.1
1999.....	5.6	7.6	7.7	7.7	8.7	7.3	5.9	6.7	5.9	4.7
White, female										
2002.....	1.8	2.4	2.0	2.0	1.7	1.7	1.9	1.4	1.1	1.1
2001.....	1.7	2.2	1.9	2.1	2.0	1.9	1.5	1.6	1.6	1.1
2000.....	1.5	1.8	1.8	1.9	1.6	1.7	1.6	1.6	1.2	1.5
1999.....	1.3	1.9	1.4	1.9	1.9	2.0	1.6	1.4	0.9	1.3
Black, both sexes										
2002.....	4.1	6.2	8.4	10.9	11.2	8.3	7.9	5.0	*	*
2001.....	5.2	7.9	10.3	11.0	11.0	9.6	6.3	5.2	*	*
2000.....	4.7	8.8	8.7	10.3	11.1	10.2	6.8	6.0	*	*
1999.....	4.9	7.9	8.9	11.6	11.3	9.6	8.4	7.1	*	*
Black, male										
2002.....	5.9	10.4	14.6	20.7	21.0	18.0	16.4	9.6	*	*
2001.....	8.4	13.4	17.4	20.6	21.6	19.6	13.5	11.6	*	*
2000.....	7.3	14.5	15.7	18.9	21.3	20.3	14.4	14.6	*	*
1999.....	7.5	12.9	15.2	21.9	21.0	19.5	17.3	14.2	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to use of alcohol (F10)

Black, female

2002.....	2.4	2.5	3.2	2.8	3.6	*	*	*	*	*
2001.....	2.4	3.1	4.2	3.1	*	*	*	*	*	*
2000.....	2.3	3.8	2.8	3.2	*	*	*	*	*	*
1999.....	2.5	3.7	3.6	3.2	3.7	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)										
All races 3/, both sexes										
2002.....	2.0	2.0	1.6	1.1	0.9	1.0	1.2	1.2	1.4	1.7
2001.....	1.8	2.1	1.4	0.9	0.7	0.7	1.0	1.3	1.5	1.8
2000.....	1.8	1.9	1.3	0.9	0.8	0.8	1.0	1.4	1.6	1.6
1999.....	2.1	1.9	1.2	0.9	0.9	0.9	1.2	1.4	1.5	1.5
All races 3/, male										
2002.....	2.8	2.9	2.6	1.6	1.2	1.3	1.3	1.4	1.7	2.6
2001.....	2.5	3.0	2.3	1.3	1.0	0.9	1.3	1.4	1.9	2.8
2000.....	2.4	2.8	2.1	1.2	1.0	0.9	1.1	1.9	2.0	2.1
1999.....	3.1	2.8	1.8	1.3	1.4	1.1	1.5	1.8	2.3	2.5
All races 3/, female										
2002.....	1.3	1.1	0.7	0.6	0.6	0.7	1.2	1.1	1.3	1.2
2001.....	1.2	1.1	0.6	0.5	0.5	0.4	0.9	1.3	1.3	1.4
2000.....	1.3	1.0	0.6	0.6	0.6	0.6	1.0	1.0	1.4	1.3
1999.....	1.0	1.0	0.6	0.5	0.4	0.7	0.9	1.2	1.0	1.1
White, both sexes										
2002.....	1.8	1.6	1.1	0.9	0.8	1.0	1.3	1.3	1.5	1.6
2001.....	1.5	1.5	1.1	0.8	0.7	0.7	1.1	1.4	1.5	1.9
2000.....	1.6	1.4	1.0	0.7	0.8	0.8	1.1	1.4	1.7	1.6
1999.....	1.6	1.3	0.9	0.6	0.9	0.9	1.2	1.5	1.6	1.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)										
White, male										
2002.....	2.5	2.4	1.7	1.2	0.9	1.2	1.4	1.4	1.8	2.7
2001.....	2.2	2.2	1.7	1.1	0.9	0.9	1.3	1.3	1.9	2.9
2000.....	2.1	2.1	1.6	0.9	1.0	0.9	1.1	2.0	2.1	2.1
1999.....	2.6	2.1	1.3	0.9	1.4	1.1	1.5	1.7	2.4	2.7
White, female										
2002.....	1.0	0.7	0.5	0.5	0.6	0.8	1.3	1.2	1.4	1.1
2001.....	0.9	0.8	0.4	0.4	0.5	0.5	0.9	1.4	1.3	1.5
2000.....	1.0	0.8	0.4	0.5	0.5	0.7	1.0	1.0	1.5	1.4
1999.....	0.7	0.6	0.4	0.4	*	0.7	1.0	1.3	1.1	1.1
Black, both sexes										
2002.....	4.6	5.7	5.8	3.4	2.3	*	*	*	*	*
2001.....	4.4	6.6	4.7	2.2	*	*	*	*	*	*
2000.....	4.3	5.8	4.4	2.6	*	*	*	*	*	*
1999.....	5.3	6.4	4.2	3.4	*	*	*	*	*	*
Black, male										
2002.....	5.6	8.1	10.2	5.8	4.4	*	*	*	*	*
2001.....	5.6	9.7	7.9	3.6	*	*	*	*	*	*
2000.....	5.5	9.1	7.2	4.1	*	*	*	*	*	*
1999.....	7.5	9.3	6.4	5.8	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

Black, female

2002.....	3.7	3.7	2.1	*	*	*	*	*	*	*
2001.....	3.3	3.9	2.0	*	*	*	*	*	*	*
2000.....	3.3	3.0	2.0	*	*	*	*	*	*	*
1999.....	3.4	3.8	2.3	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Schizophrenia, schizotypal and delusional disorders (F20-F29)										
All races 3/, both sexes										
2002.....	0.1	0.1	*	0.2	*	0.3	0.7	1.0	1.3	3.3
2001.....	*	0.1	*	0.1	0.2	0.3	0.4	1.1	1.3	2.7
2000.....	*	*	*	*	0.2	0.3	0.7	0.9	1.7	3.3
1999.....	*	*	*	*	0.2	0.3	0.6	1.0	2.1	4.9
All races 3/, male										
2002.....	*	*	*	*	*	*	0.7	0.9	*	1.9
2001.....	*	*	*	*	*	0.5	0.5	0.9	1.1	1.5
2000.....	*	*	*	*	*	*	0.7	0.7	1.6	2.4
1999.....	*	*	*	*	*	*	0.7	1.0	1.4	4.6
All races 3/, female										
2002.....	*	*	*	*	*	*	0.6	1.1	1.6	3.9
2001.....	*	*	*	*	*	*	*	1.2	1.4	3.1
2000.....	*	*	*	*	*	*	0.7	1.0	1.8	3.8
1999.....	*	*	*	*	*	*	0.5	1.0	2.5	5.0
White, both sexes										
2002.....	*	*	*	0.2	*	0.3	0.6	1.0	1.3	3.4
2001.....	*	*	*	*	0.2	0.3	0.4	1.1	1.3	2.8
2000.....	*	*	*	*	*	0.4	0.7	0.9	1.8	3.4
1999.....	*	*	*	*	*	0.3	0.6	1.0	2.1	5.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mood (affective) disorders (F30-F39)										
All races 3/, both sexes										
2002.....	*	*	*	*	0.2	0.2	0.7	1.3	3.5	9.1
2001.....	0.1	*	*	*	*	0.4	0.6	1.4	3.2	9.6
2000.....	*	*	*	*	*	0.3	0.5	1.2	2.9	9.8
1999.....	*	*	*	*	*	0.3	0.6	1.6	3.4	9.4
All races 3/, male										
2002.....	*	*	*	*	*	*	0.5	0.9	2.6	6.6
2001.....	*	*	*	*	*	*	0.6	1.3	2.4	8.0
2000.....	*	*	*	*	*	*	0.5	0.9	2.3	9.1
1999.....	*	*	*	*	*	*	*	1.6	2.7	8.8
All races 3/, female										
2002.....	*	*	*	*	*	*	0.7	1.6	4.0	10.1
2001.....	*	*	*	*	*	*	0.6	1.5	3.7	10.3
2000.....	*	*	*	*	*	*	0.5	1.5	3.2	10.1
1999.....	*	*	*	*	*	*	0.6	1.7	3.9	9.7
White, both sexes										
2002.....	*	*	*	*	0.2	0.2	0.7	1.4	3.6	9.8
2001.....	*	*	*	*	*	0.4	0.6	1.6	3.4	10.1
2000.....	*	*	*	*	*	0.3	0.6	1.3	3.0	10.6
1999.....	*	*	*	*	*	0.3	0.6	1.8	3.7	10.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mental retardation (F70-F79)										
All races 3/, both sexes										
2002.....	0.2	0.2	0.2	0.4	0.4	0.4	0.4	0.3	0.5	0.5
2001.....	0.1	0.2	0.2	0.3	0.3	0.3	0.4	0.4	0.5	*
2000.....	0.1	0.2	0.2	0.3	0.3	0.2	0.3	0.4	*	0.5
1999.....	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.5	*
All races 3/, male										
2002.....	0.2	0.3	0.3	0.5	0.4	*	*	*	*	*
2001.....	*	0.2	0.2	*	0.5	*	*	*	*	*
2000.....	*	0.2	0.2	0.3	*	*	*	*	*	*
1999.....	0.2	0.2	0.3	0.3	*	0.5	*	*	*	*
All races 3/, female										
2002.....	*	0.2	*	0.3	0.3	*	*	*	*	*
2001.....	*	*	0.2	0.3	*	*	*	*	*	*
2000.....	*	*	*	0.3	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.2	0.2	0.2	0.4	0.4	0.4	0.4	*	0.6	0.6
2001.....	0.1	0.2	0.2	0.3	0.3	0.3	0.4	0.3	0.6	*
2000.....	0.1	0.2	0.2	0.4	0.3	0.2	0.3	0.4	*	0.5
1999.....	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.6	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	*	0.3	0.4	1.2	4.4
2001.....	*	*	*	*	*	*	*	0.5	1.0	4.1
2000.....	*	*	*	*	*	*	*	0.4	1.2	3.9
1999.....	*	*	*	*	*	*	0.4	0.4	1.0	4.0
All races 3/, male										
2002.....	*	*	*	*	*	*	*	*	1.6	3.6
2001.....	*	*	*	*	*	*	*	*	*	3.5
2000.....	*	*	*	*	*	*	*	*	1.3	3.3
1999.....	*	*	*	*	*	*	*	*	*	2.5
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	1.0	4.7
2001.....	*	*	*	*	*	*	*	0.5	1.0	4.4
2000.....	*	*	*	*	*	*	*	0.5	1.1	4.1
1999.....	*	*	*	*	*	*	*	*	1.1	4.7
White, both sexes										
2002.....	*	*	*	*	*	*	*	0.4	1.3	4.4
2001.....	*	*	*	*	*	*	*	0.5	1.0	4.4
2000.....	*	*	*	*	*	*	*	0.4	1.2	4.0
1999.....	*	*	*	*	*	*	0.4	0.5	1.1	4.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
VI. Diseases of the nervous system (G00-G98)										
All races 3/, both sexes										
2002.....	5.1	7.8	10.5	15.7	24.9	42.7	83.2	185.2	388.8	951.0
2001.....	5.0	7.3	10.3	15.7	24.0	41.0	81.1	178.4	375.0	906.2
2000.....	5.0	7.3	11.0	15.6	23.8	40.2	79.5	172.9	361.9	864.9
1999.....	4.6	7.0	10.3	14.4	22.1	39.3	76.3	163.6	340.4	790.8
All races 3/, male										
2002.....	5.7	8.5	11.2	16.5	27.4	48.2	95.7	210.0	422.2	867.4
2001.....	5.5	7.7	10.9	16.2	25.7	47.4	92.9	206.0	421.1	861.6
2000.....	5.7	8.3	11.8	16.5	25.9	46.5	92.8	198.3	407.5	826.2
1999.....	5.1	7.2	11.0	15.9	24.4	44.9	89.1	193.7	394.2	772.7
All races 3/, female										
2002.....	4.4	7.0	9.9	14.9	22.6	37.8	73.2	167.7	368.5	987.3
2001.....	4.5	6.9	9.7	15.2	22.5	35.5	71.7	159.1	347.4	924.9
2000.....	4.4	6.4	10.3	14.8	21.8	34.9	69.0	155.1	335.0	880.7
1999.....	4.1	6.7	9.6	13.0	20.1	34.6	66.3	142.7	308.9	798.2
White, both sexes										
2002.....	4.9	7.5	10.2	15.7	25.4	43.8	85.9	192.1	405.4	989.7
2001.....	4.8	7.0	10.2	15.5	24.3	42.0	84.1	184.4	389.0	942.3
2000.....	4.8	7.0	10.9	15.6	24.1	41.2	81.7	179.1	375.6	900.8
1999.....	4.4	6.6	10.1	14.3	22.2	40.1	78.1	169.0	353.7	825.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
VI. Diseases of the nervous system (G00-G98)										
White, male										
2002.....	5.5	8.2	10.7	16.4	28.0	49.0	98.4	217.7	438.9	904.3
2001.....	5.3	7.5	10.7	16.1	26.0	48.4	95.9	213.3	436.5	896.4
2000.....	5.3	8.1	11.9	16.3	26.0	47.8	95.3	205.0	422.6	859.9
1999.....	4.7	6.9	10.7	15.9	24.4	45.7	91.7	200.0	411.2	806.2
White, female										
2002.....	4.3	6.9	9.7	15.1	22.9	39.2	75.8	173.7	384.8	1,026.8
2001.....	4.4	6.6	9.7	15.0	22.7	36.3	74.5	163.9	360.3	961.5
2000.....	4.3	5.9	10.0	14.9	22.3	35.4	70.7	160.9	347.7	917.4
1999.....	4.1	6.3	9.4	12.8	20.1	35.3	67.3	147.1	319.8	832.7
Black, both sexes										
2002.....	7.7	11.7	16.0	18.8	28.2	43.4	76.3	150.1	269.2	655.4
2001.....	7.4	10.7	14.9	21.3	27.3	42.2	69.6	152.0	273.6	624.9
2000.....	7.9	12.1	15.4	20.0	26.2	41.6	73.1	136.1	259.8	561.2
1999.....	7.1	11.7	15.6	18.6	26.3	40.1	73.7	131.6	225.9	485.3
Black, male										
2002.....	8.5	13.5	17.1	20.5	30.7	52.1	93.8	168.1	303.4	570.4
2001.....	8.5	11.5	16.5	22.1	29.5	49.3	81.8	170.5	302.6	593.4
2000.....	9.6	12.2	15.3	23.9	30.6	46.6	86.9	159.7	285.8	537.4
1999.....	8.9	12.3	18.1	19.3	29.7	46.2	81.1	152.4	231.1	481.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

VI. Diseases of the nervous system (G00-G98)

Black, female

2002.....	7.0	10.2	15.0	17.4	26.2	37.0	64.5	139.4	251.5	688.0
2001.....	6.4	10.0	13.6	20.7	25.6	37.0	61.4	140.9	258.8	636.5
2000.....	6.4	11.9	15.5	16.9	22.7	38.0	63.9	121.9	246.7	570.1
1999.....	5.6	11.2	13.5	18.0	23.6	35.6	68.9	119.0	223.4	486.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningitis (G00,G03)										
All races 3/, both sexes										
2002.....	0.2	0.3	0.3	0.3	0.3	0.5	0.7	0.6	0.7	1.2
2001.....	0.2	0.3	0.3	0.4	0.5	0.5	0.5	0.6	1.0	0.9
2000.....	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1	1.2
1999.....	0.2	0.3	0.3	0.3	0.5	0.5	0.7	0.9	1.2	1.5
All races 3/, male										
2002.....	0.2	0.4	0.3	0.4	*	0.6	0.9	0.7	1.0	*
2001.....	*	0.3	0.3	0.4	0.6	0.5	0.7	0.9	1.1	*
2000.....	0.2	0.3	0.4	0.4	0.5	0.6	0.9	1.3	1.1	*
1999.....	0.2	0.4	0.4	0.4	0.4	0.6	0.8	0.9	1.3	1.7
All races 3/, female										
2002.....	*	0.2	0.2	*	0.3	0.4	0.6	0.6	*	1.1
2001.....	*	0.2	0.2	0.4	0.4	0.4	0.4	0.5	0.9	0.8
2000.....	*	0.2	0.3	*	*	0.4	0.5	0.6	1.1	1.0
1999.....	*	0.2	*	*	0.5	0.4	0.7	0.9	1.1	1.4
White, both sexes										
2002.....	0.1	0.2	0.2	0.3	0.3	0.5	0.7	0.6	0.6	1.2
2001.....	0.1	0.2	0.2	0.3	0.5	0.4	0.6	0.6	0.9	0.8
2000.....	0.1	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1	1.0
1999.....	0.1	0.2	0.2	0.3	0.4	0.5	0.6	0.7	1.1	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other inflammatory diseases of central nervous system (G04-G09)										
All races 3/, both sexes										
2002.....	0.2	0.2	0.2	0.3	0.4	0.5	0.8	1.1	1.4	1.2
2001.....	0.1	0.2	0.2	0.3	0.4	0.5	0.8	0.9	1.1	1.0
2000.....	0.2	0.1	0.2	0.3	0.3	0.5	0.8	1.1	1.1	1.0
1999.....	0.1	0.2	0.2	0.3	0.3	0.5	0.7	0.7	1.0	1.1
All races 3/, male										
2002.....	0.2	0.3	0.3	0.4	0.5	0.8	1.0	1.3	2.0	*
2001.....	*	0.2	0.3	0.3	0.4	0.7	0.8	1.1	1.3	1.8
2000.....	0.2	*	0.3	0.3	0.4	0.6	1.0	1.6	1.4	*
1999.....	0.2	*	*	0.5	*	0.6	0.6	0.9	1.6	*
All races 3/, female										
2002.....	*	0.2	0.2	0.3	0.3	*	0.6	0.9	1.1	1.1
2001.....	*	0.2	*	0.3	0.4	0.4	0.8	0.8	0.9	0.7
2000.....	*	*	*	0.3	*	0.5	0.6	0.8	0.9	0.9
1999.....	*	*	*	*	0.4	0.4	0.8	0.6	0.7	0.9
White, both sexes										
2002.....	0.1	0.2	0.2	0.3	0.4	0.5	0.7	1.0	1.4	1.3
2001.....	0.1	0.2	0.2	0.3	0.4	0.5	0.7	0.9	1.0	1.0
2000.....	0.2	*	0.2	0.3	0.4	0.5	0.7	1.1	1.0	1.0
1999.....	0.1	0.2	0.2	0.3	0.3	0.5	0.6	0.7	1.1	1.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Parkinson's disease (G20-G21)										
All races 3/, both sexes										
2002.....	*	*	0.2	0.6	1.9	6.5	18.4	45.4	89.8	135.2
2001.....	*	*	0.2	0.7	1.8	5.8	18.1	46.4	90.9	134.2
2000.....	*	*	0.2	0.6	1.7	6.6	16.7	44.8	87.6	131.9
1999.....	*	*	0.2	0.6	1.6	5.6	16.7	41.8	82.8	124.4
All races 3/, male										
2002.....	*	*	0.3	0.8	2.5	9.8	27.3	67.2	136.6	220.1
2001.....	*	*	0.3	1.0	2.7	8.2	26.5	68.8	141.7	224.4
2000.....	*	*	*	0.8	2.2	9.0	24.1	67.4	135.0	222.3
1999.....	*	*	0.3	0.7	2.3	7.7	24.8	64.4	131.3	205.6
All races 3/, female										
2002.....	*	*	*	0.5	1.4	3.8	11.3	30.0	61.3	98.4
2001.....	*	*	*	0.4	1.0	3.7	11.5	30.7	60.4	96.4
2000.....	*	*	*	0.4	1.3	4.4	10.9	29.1	59.5	95.0
1999.....	*	*	*	0.4	0.9	3.9	10.4	26.0	54.4	91.7
White, both sexes										
2002.....	*	*	0.2	0.6	2.1	6.9	19.6	48.3	95.7	143.3
2001.....	*	*	0.2	0.7	1.9	6.2	19.4	49.6	96.4	141.8
2000.....	*	*	0.2	0.6	1.9	7.1	17.8	47.5	92.5	139.5
1999.....	*	*	0.2	0.6	1.6	6.0	17.7	44.1	87.4	131.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Parkinson's disease (G20-G21)										
White, male										
2002.....	*	*	0.3	0.8	2.7	10.2	28.8	71.2	145.0	232.7
2001.....	*	*	0.3	1.0	3.0	8.8	28.1	73.4	149.6	237.6
2000.....	*	*	*	0.9	2.5	9.6	25.5	70.7	141.5	235.3
1999.....	*	*	0.3	0.7	2.4	8.1	26.4	67.7	138.6	216.7
White, female										
2002.....	*	*	*	0.5	1.5	3.9	12.0	31.9	65.5	104.4
2001.....	*	*	*	0.5	0.9	4.0	12.3	32.7	64.2	101.7
2000.....	*	*	*	0.4	1.4	4.8	11.7	31.2	63.3	100.6
1999.....	*	*	*	0.4	0.9	4.2	10.8	27.6	57.3	96.8
Black, both sexes										
2002.....	*	*	*	*	*	4.6	9.5	20.6	32.7	51.7
2001.....	*	*	*	*	*	2.7	8.9	18.9	34.9	59.0
2000.....	*	*	*	*	*	3.1	8.8	21.0	37.6	51.6
1999.....	*	*	*	*	*	3.6	8.7	19.5	32.4	52.0
Black, male										
2002.....	*	*	*	*	*	7.5	16.0	30.3	55.0	89.0
2001.....	*	*	*	*	*	*	14.8	25.6	57.8	99.8
2000.....	*	*	*	*	*	*	13.7	37.8	65.8	86.5
1999.....	*	*	*	*	*	5.7	12.6	31.8	48.8	86.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Parkinson's disease (G20-G21)										
Black, female										
2002.....	*	*	*	*	*	*	5.2	14.8	21.1	37.4
2001.....	*	*	*	*	*	*	4.9	15.0	23.3	43.8
2000.....	*	*	*	*	*	*	5.5	10.8	23.4	38.6
1999.....	*	*	*	*	*	*	6.1	12.0	24.2	39.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)										
All races 3/, both sexes										
2002.....	*	*	0.2	0.9	3.2	9.8	30.6	95.4	245.7	752.3
2001.....	*	*	0.3	1.2	3.3	9.0	29.2	88.5	232.5	710.3
2000.....	*	*	0.3	1.1	3.2	9.1	28.9	85.2	221.1	667.7
1999.....	*	0.1	0.2	1.0	3.0	8.7	26.7	78.6	206.2	601.3
All races 3/, male										
2002.....	*	*	0.2	0.8	3.1	9.2	29.5	90.9	220.6	572.0
2001.....	*	*	0.3	0.9	3.2	9.0	27.7	85.6	216.6	556.1
2000.....	*	*	0.3	0.9	2.9	9.5	29.1	79.9	206.8	522.1
1999.....	*	*	*	1.1	3.0	8.8	26.9	77.4	198.2	485.7
All races 3/, female										
2002.....	*	*	*	1.0	3.3	10.3	31.5	98.5	260.9	830.5
2001.....	*	*	0.3	1.4	3.4	9.0	30.4	90.5	242.1	774.9
2000.....	*	*	0.3	1.2	3.5	8.7	28.8	88.9	229.5	727.0
1999.....	*	*	0.3	0.8	3.0	8.6	26.6	79.4	210.9	648.0
White, both sexes										
2002.....	*	*	0.2	1.0	3.4	10.3	31.9	99.1	255.3	781.8
2001.....	*	*	0.3	1.3	3.6	9.3	30.6	90.9	240.5	738.1
2000.....	*	*	0.3	1.1	3.4	9.6	29.9	88.1	229.1	695.3
1999.....	*	*	0.2	1.0	3.3	9.1	27.6	81.3	213.6	627.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)										
White, male										
2002.....	*	*	*	0.8	3.3	9.4	30.5	94.3	227.6	594.8
2001.....	*	*	0.3	0.9	3.4	9.2	28.8	88.1	222.7	577.2
2000.....	*	*	0.3	1.0	3.1	10.1	29.9	82.6	214.4	542.1
1999.....	*	*	*	1.1	3.3	9.2	27.6	80.1	205.6	506.9
White, female										
2002.....	*	*	*	1.2	3.5	11.0	33.1	102.6	272.2	863.0
2001.....	*	*	0.4	1.6	3.8	9.4	32.1	92.9	251.3	805.4
2000.....	*	*	0.3	1.3	3.7	9.2	29.9	92.0	237.7	757.6
1999.....	*	*	*	0.9	3.3	9.1	27.5	82.1	218.3	676.5
Black, both sexes										
2002.....	*	*	*	*	2.5	8.5	27.2	81.3	186.8	543.7
2001.....	*	*	*	*	1.9	8.7	23.3	86.8	183.6	506.7
2000.....	*	*	*	*	2.1	7.4	25.9	71.1	171.7	445.1
1999.....	*	*	*	*	*	6.9	25.3	65.0	152.0	372.4
Black, male										
2002.....	*	*	*	*	*	9.5	29.4	82.2	189.5	412.8
2001.....	*	*	*	*	*	9.9	24.2	84.6	185.4	414.1
2000.....	*	*	*	*	*	7.8	27.7	64.7	156.0	371.3
1999.....	*	*	*	*	*	8.0	27.5	62.7	139.5	317.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)										
Black, female										
2002.....	*	*	*	*	*	7.8	25.8	80.7	185.4	593.9
2001.....	*	*	*	*	*	7.9	22.6	88.2	182.7	541.0
2000.....	*	*	*	*	*	7.1	24.6	74.9	179.6	472.6
1999.....	*	*	*	*	*	6.0	23.9	66.4	158.2	392.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple sclerosis (G35)										
All races 3/, both sexes										
2002.....	0.8	1.7	2.3	3.0	3.5	3.4	3.5	3.4	3.3	2.2
2001.....	0.9	1.6	2.3	3.0	3.3	3.4	3.3	3.6	3.4	2.1
2000.....	0.9	1.4	2.5	3.0	3.1	3.1	3.1	3.0	3.1	1.7
1999.....	0.9	1.4	2.4	2.4	2.7	3.0	2.8	3.2	2.8	2.1
All races 3/, male										
2002.....	0.5	1.1	1.6	2.2	3.1	2.4	2.6	2.7	2.4	1.9
2001.....	0.6	1.2	1.7	2.1	2.3	3.1	2.3	3.1	2.1	1.5
2000.....	0.7	1.1	1.8	2.2	2.3	2.3	2.4	2.4	2.2	1.8
1999.....	0.7	0.8	1.8	1.9	1.8	2.4	2.4	2.3	2.2	*
All races 3/, female										
2002.....	1.1	2.3	3.0	3.8	3.8	4.2	4.2	3.9	3.8	2.4
2001.....	1.2	2.0	3.0	3.7	4.1	3.6	4.1	4.0	4.1	2.3
2000.....	1.1	1.8	3.1	3.8	3.8	3.7	3.6	3.5	3.7	1.7
1999.....	1.2	2.0	3.0	2.9	3.5	3.5	3.1	3.8	3.1	2.4
White, both sexes										
2002.....	0.8	1.7	2.3	3.2	3.6	3.6	3.8	3.6	3.5	2.4
2001.....	0.9	1.6	2.4	3.2	3.5	3.6	3.6	3.9	3.6	2.2
2000.....	0.9	1.4	2.6	3.1	3.2	3.3	3.3	3.3	3.3	1.9
1999.....	0.9	1.5	2.5	2.7	2.9	3.1	2.9	3.5	3.0	2.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple sclerosis (G35)										
Black, female										
2002.....	2.1	2.9	4.1	3.5	4.0	4.3	*	*	*	*
2001.....	1.6	2.5	3.3	3.5	4.0	*	*	*	*	*
2000.....	1.7	3.1	3.1	3.4	3.9	*	*	*	*	*
1999.....	1.6	1.8	2.8	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Epilepsy (G40-G41)										
All races 3/, both sexes										
2002.....	0.6	0.6	0.6	0.5	0.6	0.7	0.9	1.3	1.4	1.8
2001.....	0.6	0.5	0.6	0.6	0.6	0.7	0.8	1.1	1.2	1.6
2000.....	0.6	0.6	0.6	0.7	0.6	0.7	1.0	1.0	1.6	2.0
1999.....	0.6	0.5	0.6	0.5	0.6	0.8	1.0	1.1	1.5	1.8
All races 3/, male										
2002.....	0.8	0.8	0.8	0.4	0.7	0.8	0.9	1.2	1.5	2.3
2001.....	0.8	0.6	0.7	0.7	0.7	0.8	0.8	1.0	*	2.0
2000.....	0.8	0.8	0.7	0.8	0.8	0.7	1.0	1.0	*	2.4
1999.....	0.7	0.7	0.6	0.6	0.7	0.9	1.0	0.9	1.3	2.3
All races 3/, female										
2002.....	0.4	0.5	0.4	0.5	0.6	0.7	0.9	1.3	1.4	1.5
2001.....	0.5	0.4	0.5	0.5	0.6	0.6	0.7	1.2	1.5	1.4
2000.....	0.4	0.4	0.5	0.5	0.5	0.6	1.0	1.0	2.0	1.9
1999.....	0.4	0.4	0.5	0.5	0.5	0.7	0.9	1.2	1.6	1.7
White, both sexes										
2002.....	0.6	0.6	0.5	0.4	0.6	0.6	0.7	1.1	1.3	1.6
2001.....	0.6	0.5	0.5	0.5	0.6	0.6	0.6	0.9	1.0	1.6
2000.....	0.6	0.5	0.6	0.6	0.5	0.6	0.8	0.8	1.5	1.7
1999.....	0.5	0.5	0.5	0.4	0.4	0.6	0.8	0.9	1.4	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transient cerebral ischemic attacks and related syndromes (G45)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	*	*	0.4	0.8	2.9
2001.....	*	*	*	*	*	*	*	0.4	0.7	3.8
2000.....	*	*	*	*	*	*	*	0.5	1.0	3.9
1999.....	*	*	*	*	*	*	*	0.4	1.0	4.6
All races 3/, male										
2002.....	*	*	*	*	*	*	*	*	1.0	1.8
2001.....	*	*	*	*	*	*	*	*	*	4.2
2000.....	*	*	*	*	*	*	*	0.7	*	2.9
1999.....	*	*	*	*	*	*	*	0.7	1.2	3.8
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	0.6	3.4
2001.....	*	*	*	*	*	*	*	*	*	3.6
2000.....	*	*	*	*	*	*	*	*	1.1	4.2
1999.....	*	*	*	*	*	*	*	*	0.9	4.9
White, both sexes										
2002.....	*	*	*	*	*	*	*	0.4	0.8	3.0
2001.....	*	*	*	*	*	*	*	0.5	0.8	3.9
2000.....	*	*	*	*	*	*	*	0.5	1.1	4.0
1999.....	*	*	*	*	*	*	*	0.4	1.1	4.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Infantile cerebral palsy (G80)										
All races 3/, both sexes										
2002.....	0.2	0.3	0.3	0.3	0.5	0.4	0.5	0.5	0.6	*
2001.....	0.2	0.2	0.3	0.4	0.4	0.3	0.6	0.6	0.4	*
2000.....	0.2	0.2	0.3	0.3	0.4	0.5	0.4	0.4	0.7	*
1999.....	0.2	0.2	0.2	0.2	0.3	0.3	0.4	0.4	0.5	0.5
All races 3/, male										
2002.....	0.2	*	0.4	0.4	0.6	0.5	0.5	*	*	*
2001.....	0.3	0.2	0.4	0.4	0.4	*	0.9	0.8	*	*
2000.....	0.2	0.3	0.3	0.3	*	0.7	0.7	*	*	*
1999.....	0.2	0.3	0.3	0.3	0.4	0.5	*	*	*	*
All races 3/, female										
2002.....	0.2	0.3	0.3	0.3	0.4	*	0.4	*	*	*
2001.....	*	0.2	0.3	0.4	*	*	*	*	*	*
2000.....	0.2	0.2	*	0.3	0.4	*	*	*	0.7	*
1999.....	0.2	0.2	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.2	0.3	0.3	0.4	0.5	0.4	0.5	0.5	0.6	*
2001.....	0.2	0.2	0.3	0.4	0.4	0.3	0.6	0.6	0.4	*
2000.....	0.2	0.3	0.3	0.3	0.4	0.5	0.5	0.4	0.7	*
1999.....	0.2	0.3	0.2	0.3	0.3	0.4	0.4	0.4	0.5	0.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)										
All races 3/, both sexes										
2002.....	3.1	4.5	6.4	9.7	14.4	20.7	27.7	37.2	45.2	53.8
2001.....	2.9	4.3	6.1	9.2	13.8	20.7	27.6	36.3	43.9	51.9
2000.....	2.9	4.5	6.7	9.3	13.9	19.2	27.7	35.9	44.6	55.4
1999.....	2.5	4.1	6.2	9.1	13.1	19.8	27.1	36.6	43.4	53.6
All races 3/, male										
2002.....	3.7	5.6	7.3	11.1	16.5	24.2	32.9	44.8	56.2	66.1
2001.....	3.4	5.0	7.0	10.4	15.3	24.6	33.0	44.4	56.2	69.7
2000.....	3.4	5.5	7.7	10.7	16.3	22.9	33.5	43.6	58.4	72.0
1999.....	3.0	4.6	7.2	10.5	15.3	23.3	32.0	45.6	56.5	70.3
All races 3/, female										
2002.....	2.4	3.4	5.5	8.3	12.4	17.7	23.6	31.8	38.4	48.5
2001.....	2.3	3.7	5.2	8.0	12.4	17.3	23.4	30.6	36.6	44.5
2000.....	2.4	3.6	5.7	8.0	11.8	16.0	23.2	30.6	36.5	48.7
1999.....	2.0	3.6	5.1	7.8	11.0	16.8	23.2	30.4	35.7	46.8
White, both sexes										
2002.....	3.0	4.4	6.2	9.5	14.4	20.9	28.0	37.5	46.1	54.7
2001.....	2.8	4.1	5.9	8.8	13.6	20.8	27.9	36.6	44.4	52.5
2000.....	2.7	4.3	6.5	9.2	13.9	19.1	27.8	36.6	45.4	56.2
1999.....	2.5	3.7	6.0	8.8	12.9	19.8	27.2	36.9	44.5	54.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)										
White, male										
2002.....	3.7	5.5	7.0	10.9	16.6	24.1	33.0	45.5	57.7	67.5
2001.....	3.3	4.8	6.7	10.1	15.2	24.8	33.3	44.7	57.4	70.1
2000.....	3.2	5.4	7.7	10.5	16.1	23.0	33.8	44.5	59.3	73.1
1999.....	3.0	4.5	7.1	10.4	15.2	23.4	32.3	46.3	58.5	71.5
White, female										
2002.....	2.4	3.3	5.3	8.1	12.4	18.1	23.9	31.8	39.0	49.2
2001.....	2.3	3.5	5.1	7.5	12.1	17.3	23.5	30.8	36.5	45.1
2000.....	2.3	3.3	5.4	8.0	11.8	15.7	23.1	31.1	37.1	49.3
1999.....	2.0	3.0	4.9	7.3	10.8	16.7	23.1	30.4	36.2	47.3
Black, both sexes										
2002.....	4.2	6.4	9.2	13.0	18.1	23.5	31.6	37.6	42.6	50.5
2001.....	4.0	6.5	9.7	14.6	17.7	24.3	30.7	38.4	45.8	48.8
2000.....	4.5	7.5	9.9	12.5	16.7	24.2	30.4	35.2	40.7	51.3
1999.....	3.5	7.6	9.4	13.1	16.5	22.5	31.4	38.0	34.7	50.7
Black, male										
2002.....	5.1	7.9	10.5	15.6	20.4	29.2	40.1	42.3	49.7	57.9
2001.....	5.0	7.5	11.4	15.2	19.3	28.2	35.7	49.7	53.0	62.4
2000.....	5.6	8.3	10.2	16.0	20.1	27.6	37.4	44.9	56.5	63.4
1999.....	4.2	7.5	11.1	13.1	18.9	24.7	33.9	47.0	37.5	65.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

Black, female

2002.....	3.5	5.1	8.2	10.9	16.3	19.2	26.0	34.9	39.0	47.6
2001.....	3.1	5.5	8.2	14.1	16.5	21.4	27.4	31.6	42.1	43.8
2000.....	3.5	6.7	9.6	9.7	14.1	21.7	25.8	29.3	32.7	46.8
1999.....	2.9	7.6	7.9	13.2	14.6	21.0	29.7	32.6	33.3	45.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
IX. Diseases of the circulatory system (I00-I99)										
All races 3/, both sexes										
2002.....	50.4	89.0	145.7	235.5	376.5	605.9	997.5	1,707.6	3,047.6	7,414.0
2001.....	49.6	86.6	146.1	236.8	388.4	623.5	1,022.9	1,767.1	3,145.7	7,699.8
2000.....	50.5	89.8	148.2	250.1	410.1	651.3	1,071.3	1,832.2	3,265.6	8,055.8
1999.....	50.8	89.3	149.7	256.4	418.3	682.1	1,115.6	1,895.5	3,369.1	8,220.5
All races 3/, male										
2002.....	68.7	125.2	207.7	329.0	515.2	805.6	1,291.7	2,147.8	3,646.7	7,659.3
2001.....	67.3	122.7	206.2	329.7	529.2	825.3	1,331.5	2,231.5	3,786.4	7,958.0
2000.....	69.1	128.3	210.8	352.5	559.6	861.1	1,409.4	2,313.7	3,969.1	8,470.3
1999.....	69.8	126.7	213.7	360.1	570.7	912.4	1,466.7	2,400.1	4,092.1	8,624.0
All races 3/, female										
2002.....	32.4	53.9	86.3	147.7	250.7	433.5	762.4	1,396.8	2,683.2	7,307.6
2001.....	32.2	51.6	88.6	149.8	260.6	449.7	777.7	1,441.9	2,761.9	7,591.7
2000.....	32.1	52.4	88.3	154.3	274.6	471.5	804.9	1,496.7	2,850.6	7,887.0
1999.....	32.0	53.1	88.4	159.5	280.3	485.1	841.2	1,544.6	2,946.1	8,057.9
White, both sexes										
2002.....	44.3	77.7	128.7	213.7	346.9	571.0	961.2	1,670.2	3,036.0	7,507.0
2001.....	42.3	75.2	128.8	214.1	357.9	583.9	985.3	1,728.3	3,134.6	7,796.7
2000.....	43.4	77.7	132.0	227.1	378.7	615.7	1,032.2	1,790.7	3,255.4	8,159.4
1999.....	43.1	77.5	133.3	231.7	387.3	647.6	1,072.5	1,854.7	3,357.7	8,322.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
IX. Diseases of the circulatory system (I00-I99)										
White, male										
2002.....	62.6	113.0	188.2	303.5	481.4	768.5	1,257.1	2,119.7	3,654.4	7,786.5
2001.....	59.8	110.2	187.0	303.7	496.2	785.5	1,291.8	2,200.7	3,789.8	8,082.3
2000.....	61.7	115.8	192.1	325.2	526.1	826.1	1,371.9	2,281.0	3,975.9	8,607.0
1999.....	61.6	113.9	194.5	331.6	535.7	877.3	1,424.9	2,368.9	4,101.9	8,738.2
White, female										
2002.....	25.9	42.6	70.5	127.7	222.6	397.2	720.1	1,348.4	2,656.9	7,385.6
2001.....	24.6	40.5	72.0	128.6	230.3	406.9	737.1	1,393.7	2,738.6	7,677.1
2000.....	25.0	39.9	73.1	133.8	242.8	431.7	759.7	1,445.8	2,826.7	7,977.5
1999.....	24.5	41.5	73.6	136.8	250.6	447.2	792.4	1,493.6	2,918.3	8,155.3
Black, both sexes										
2002.....	100.9	186.4	305.2	472.5	703.6	1,007.6	1,517.4	2,356.8	3,577.5	7,111.7
2001.....	107.5	184.1	309.6	475.0	719.8	1,064.4	1,557.8	2,436.9	3,638.1	7,346.8
2000.....	106.7	194.4	306.7	493.2	742.0	1,070.6	1,632.2	2,510.7	3,733.9	7,548.9
1999.....	111.7	190.9	313.3	511.4	747.2	1,100.0	1,711.0	2,564.7	3,805.3	7,610.9
Black, male										
2002.....	124.8	240.2	409.4	638.6	940.4	1,300.5	1,916.4	2,815.9	4,121.4	6,961.1
2001.....	132.1	239.0	408.8	632.4	942.9	1,350.6	1,998.5	2,960.2	4,299.5	7,393.4
2000.....	133.7	250.3	410.7	675.6	960.7	1,337.4	2,049.3	3,012.7	4,411.3	7,717.4
1999.....	140.9	249.2	426.4	688.3	996.7	1,409.3	2,153.2	3,061.2	4,428.7	7,943.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
IX. Diseases of the circulatory system (I00-I99)										
Black, female										
2002.....	79.8	139.8	216.8	336.2	518.4	790.4	1,249.6	2,082.4	3,294.4	7,169.4
2001.....	85.8	136.5	225.1	345.8	544.3	853.0	1,264.3	2,124.5	3,301.2	7,329.5
2000.....	82.8	145.9	218.0	343.7	569.4	873.8	1,355.3	2,208.1	3,392.7	7,486.2
1999.....	86.0	140.6	217.4	367.0	551.6	870.8	1,417.5	2,265.6	3,496.4	7,486.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
All races 3/, both sexes										
2002.....	0.2	0.5	0.7	1.3	2.1	2.8	5.0	8.4	11.4	17.8
2001.....	0.4	0.5	0.7	1.2	1.7	3.2	5.0	8.1	11.4	17.7
2000.....	0.4	0.6	0.8	1.4	2.2	3.4	5.6	8.1	11.6	16.8
1999.....	0.3	0.5	0.9	1.4	2.1	3.5	5.8	8.4	12.6	18.0
All races 3/, male										
2002.....	0.2	0.4	0.6	1.0	1.5	1.9	4.2	5.8	8.2	11.8
2001.....	0.3	0.4	0.5	1.1	1.2	2.4	3.2	6.0	7.7	12.8
2000.....	0.3	0.5	0.6	1.1	1.8	2.2	3.6	5.8	7.8	12.2
1999.....	0.3	0.4	0.6	1.0	1.4	2.4	4.7	5.4	8.9	10.3
All races 3/, female										
2002.....	0.3	0.5	0.9	1.5	2.6	3.7	5.6	10.2	13.4	20.4
2001.....	0.4	0.6	0.9	1.3	2.2	3.9	6.4	9.5	13.7	19.8
2000.....	0.5	0.7	0.9	1.6	2.5	4.5	7.1	9.7	13.9	18.7
1999.....	0.4	0.6	1.1	1.8	2.7	4.5	6.7	10.4	14.7	21.1
White, both sexes										
2002.....	0.2	0.4	0.7	1.2	2.0	2.8	5.2	8.6	12.1	18.9
2001.....	0.3	0.4	0.6	1.0	1.7	3.2	5.0	8.4	12.2	18.7
2000.....	0.3	0.5	0.7	1.3	2.2	3.5	5.6	8.6	12.4	17.6
1999.....	0.3	0.5	0.8	1.4	2.0	3.6	6.0	8.8	13.1	18.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Black, female

2002.....	*	*	*	*	3.6	*	*	5.5	*	*
2001.....	*	*	2.2	2.7	*	3.8	7.2	6.4	*	*
2000.....	*	*	*	*	*	4.4	*	*	*	*
1999.....	*	*	*	2.8	*	*	4.7	*	*	11.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic rheumatic heart diseases (I05-I09)										
All races 3/, both sexes										
2002.....	0.2	0.4	0.7	1.3	2.1	2.8	4.9	8.2	11.3	17.5
2001.....	0.4	0.5	0.7	1.2	1.7	3.2	4.9	8.0	11.3	17.2
2000.....	0.4	0.6	0.7	1.4	2.1	3.4	5.5	8.0	11.6	16.7
1999.....	0.3	0.5	0.8	1.4	2.1	3.5	5.8	8.3	12.5	17.9
All races 3/, male										
2002.....	0.2	0.4	0.6	1.0	1.5	1.9	4.1	5.7	8.1	11.4
2001.....	0.3	0.4	0.4	1.0	1.2	2.4	3.2	5.9	7.5	12.3
2000.....	0.3	0.5	0.6	1.1	1.8	2.2	3.5	5.7	7.8	12.1
1999.....	0.3	0.4	0.6	1.0	1.3	2.4	4.6	5.4	8.8	10.3
All races 3/, female										
2002.....	0.3	0.5	0.9	1.5	2.6	3.6	5.5	10.0	13.3	20.2
2001.....	0.4	0.6	0.9	1.3	2.2	3.8	6.3	9.4	13.5	19.3
2000.....	0.5	0.7	0.8	1.6	2.5	4.4	7.1	9.7	13.8	18.5
1999.....	0.4	0.6	1.1	1.8	2.7	4.5	6.6	10.3	14.6	21.0
White, both sexes										
2002.....	0.2	0.4	0.7	1.1	2.0	2.8	5.2	8.5	12.0	18.6
2001.....	0.3	0.4	0.6	1.0	1.7	3.1	4.9	8.3	12.0	18.3
2000.....	0.3	0.5	0.7	1.3	2.1	3.5	5.6	8.6	12.3	17.5
1999.....	0.3	0.5	0.7	1.4	2.0	3.6	6.0	8.7	13.1	18.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic rheumatic heart diseases (I05-I09)

Black, female

2002.....	*	*	*	*	3.6	*	*	*	*	*
2001.....	*	*	2.2	2.7	*	*	7.2	6.4	*	*
2000.....	*	*	*	*	*	4.2	*	*	*	*
1999.....	*	*	*	*	*	*	4.7	*	*	10.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Rheumatic mitral valve diseases (I05)										
All races 3/, both sexes										
2002.....	0.2	0.2	0.5	0.8	1.3	1.4	2.6	4.4	5.7	7.4
2001.....	0.2	0.3	0.5	0.7	1.0	1.8	2.8	4.5	6.0	8.5
2000.....	0.2	0.4	0.5	0.8	1.3	2.0	3.4	4.4	6.8	8.5
1999.....	0.2	0.3	0.5	0.8	1.2	2.1	3.2	4.9	6.8	9.0
All races 3/, male										
2002.....	*	*	0.4	0.6	0.9	0.9	2.3	2.6	3.7	4.2
2001.....	0.2	0.3	0.3	0.6	0.5	1.4	1.8	2.9	3.7	6.2
2000.....	0.2	0.3	0.3	0.7	1.1	1.2	2.2	3.2	4.7	6.1
1999.....	*	0.3	0.4	0.6	0.7	1.5	2.4	3.2	4.4	4.9
All races 3/, female										
2002.....	*	0.3	0.6	0.9	1.6	1.8	2.9	5.7	7.0	8.7
2001.....	0.2	0.4	0.6	0.8	1.4	2.1	3.6	5.5	7.4	9.4
2000.....	0.3	0.4	0.6	0.9	1.5	2.7	4.3	5.2	8.0	9.5
1999.....	0.2	0.4	0.7	1.1	1.6	2.7	3.8	6.1	8.3	10.7
White, both sexes										
2002.....	0.1	0.2	0.5	0.7	1.2	1.3	2.8	4.6	6.2	7.9
2001.....	0.2	0.3	0.4	0.6	1.0	1.7	2.8	4.7	6.5	9.1
2000.....	0.2	0.4	0.5	0.8	1.3	2.2	3.4	4.6	7.3	9.1
1999.....	0.1	0.3	0.4	0.8	1.2	2.2	3.3	5.2	7.2	9.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Disorders of both mitral and aortic valves (I08.0)

All races 3/, both sexes

2002.....	*	*	*	*	*	0.3	0.7	1.3	1.8	3.5
2001.....	*	*	*	*	*	0.3	0.6	0.7	1.6	2.5
2000.....	*	*	*	*	0.2	0.2	0.5	0.8	1.0	2.0
1999.....	*	*	*	*	0.2	0.2	0.6	0.5	1.1	2.0

All races 3/, male

2002.....	*	*	*	*	*	*	0.7	1.4	1.5	2.4
2001.....	*	*	*	*	*	*	0.6	*	1.6	2.3
2000.....	*	*	*	*	*	*	*	*	*	1.9
1999.....	*	*	*	*	*	*	0.6	*	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	0.6	1.2	2.1	4.0
2001.....	*	*	*	*	*	0.4	0.6	0.7	1.6	2.5
2000.....	*	*	*	*	*	*	0.5	0.9	1.2	2.1
1999.....	*	*	*	*	*	*	0.6	0.6	1.2	2.2

White, both sexes

2002.....	*	*	*	*	*	0.4	0.7	1.4	1.9	3.8
2001.....	*	*	*	*	*	0.4	0.6	0.7	1.7	2.6
2000.....	*	*	*	*	*	*	0.5	0.8	1.1	2.0
1999.....	*	*	*	*	*	*	0.6	0.6	1.2	2.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)										
All races 3/, both sexes										
2002.....	*	0.1	0.2	0.4	0.6	1.0	1.5	2.2	3.2	5.6
2001.....	0.1	0.1	0.2	0.4	0.5	1.0	1.4	2.7	3.4	5.9
2000.....	0.1	0.2	0.2	0.4	0.7	1.1	1.6	2.9	3.7	5.8
1999.....	0.1	0.1	0.2	0.4	0.7	1.1	1.8	2.7	4.3	6.6
All races 3/, male										
2002.....	*	*	*	0.4	0.4	0.6	1.0	1.5	2.4	3.8
2001.....	*	*	*	0.4	0.4	0.7	0.8	2.2	1.8	3.6
2000.....	*	*	*	*	0.5	0.7	0.9	1.9	2.2	4.0
1999.....	*	*	*	*	0.4	0.7	1.6	1.6	3.2	3.7
All races 3/, female										
2002.....	*	*	*	0.4	0.8	1.3	1.9	2.7	3.8	6.3
2001.....	*	*	*	0.4	0.6	1.2	2.0	3.1	4.4	6.8
2000.....	*	0.2	0.3	0.5	0.8	1.4	2.2	3.5	4.6	6.6
1999.....	*	*	0.3	0.6	0.9	1.5	2.0	3.5	5.0	7.8
White, both sexes										
2002.....	*	*	0.1	0.4	0.6	1.0	1.6	2.3	3.4	5.9
2001.....	*	*	*	0.3	0.5	1.0	1.5	2.8	3.6	6.3
2000.....	*	0.1	0.2	0.3	0.7	1.1	1.7	3.1	3.8	6.0
1999.....	*	*	0.2	0.4	0.6	1.2	1.9	2.8	4.6	6.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive diseases (I10-I13)										
All races 3/, both sexes										
2002.....	4.7	8.1	12.0	16.5	22.8	32.2	47.5	77.8	142.1	390.8
2001.....	4.5	7.2	11.1	14.9	21.2	30.3	45.3	77.2	140.5	384.7
2000.....	4.3	6.8	10.0	14.5	21.9	29.5	45.1	75.2	138.9	380.3
1999.....	4.0	6.6	9.4	14.6	20.6	30.4	43.8	75.7	136.9	366.0
All races 3/, male										
2002.....	6.4	10.8	16.0	21.9	28.6	37.8	52.5	82.0	136.9	311.1
2001.....	6.2	9.8	14.6	18.8	25.8	36.4	50.0	80.0	134.2	309.4
2000.....	5.6	9.0	13.8	18.4	27.5	33.8	51.2	76.7	135.5	309.3
1999.....	5.4	8.6	12.2	18.7	24.7	35.1	47.9	79.8	131.9	297.3
All races 3/, female										
2002.....	3.1	5.5	8.1	11.5	17.5	27.5	43.5	74.8	145.3	425.4
2001.....	2.9	4.7	7.7	11.3	17.0	25.2	41.5	75.2	144.3	416.3
2000.....	3.0	4.7	6.4	10.8	16.9	25.8	40.3	74.1	140.9	409.3
1999.....	2.7	4.7	6.7	10.8	16.9	26.3	40.6	72.9	139.8	393.7
White, both sexes										
2002.....	3.1	5.4	8.1	11.5	16.5	24.3	38.0	65.5	126.8	368.7
2001.....	2.7	4.5	7.3	10.2	15.2	22.3	35.8	65.7	126.0	364.3
2000.....	2.8	4.3	6.4	9.6	15.3	21.9	35.2	63.1	123.9	357.1
1999.....	2.3	4.0	6.0	9.6	14.4	22.3	33.9	62.7	121.1	343.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive diseases (I10-I13)										
White, male										
2002.....	4.6	7.7	11.4	16.0	21.1	28.7	42.7	69.5	121.3	290.6
2001.....	4.0	6.3	10.0	13.5	19.2	27.5	40.3	68.4	119.3	288.6
2000.....	3.8	6.2	9.3	12.8	19.7	25.8	40.8	65.4	120.1	286.2
1999.....	3.5	5.6	8.1	12.9	17.9	26.2	37.7	66.4	116.4	273.5
White, female										
2002.....	1.6	3.2	4.8	7.1	12.2	20.4	34.2	62.6	130.2	402.6
2001.....	1.4	2.7	4.6	7.1	11.6	17.8	32.2	63.7	130.1	396.0
2000.....	1.7	2.4	3.5	6.6	11.2	18.5	30.6	61.5	126.1	385.8
1999.....	1.2	2.5	3.9	6.5	11.2	19.0	30.9	60.1	123.9	371.5
Black, both sexes										
2002.....	16.3	28.9	44.5	64.1	81.0	105.7	150.4	224.0	341.7	685.4
2001.....	17.2	28.3	43.3	58.1	76.4	104.6	145.9	213.3	325.6	664.2
2000.....	15.3	26.9	40.6	59.6	82.1	103.2	152.4	221.1	326.4	673.8
1999.....	16.3	26.7	39.8	60.1	75.9	105.6	151.7	230.5	335.2	639.9
Black, male										
2002.....	20.3	36.6	57.8	82.5	104.1	126.9	170.9	247.6	354.6	602.5
2001.....	22.3	38.1	56.4	72.4	92.5	129.1	167.0	230.5	349.1	604.8
2000.....	19.5	33.2	54.3	75.6	103.7	119.4	173.5	224.0	349.8	606.6
1999.....	20.3	33.9	51.4	75.3	92.3	126.7	169.6	251.7	351.4	589.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive diseases (I10-I13)										
Black, female										
2002.....	12.8	22.2	33.2	49.0	62.9	89.9	136.7	209.9	335.0	717.1
2001.....	12.7	19.7	32.2	46.3	63.7	86.5	131.8	203.0	313.6	686.2
2000.....	11.5	21.5	28.9	46.6	65.0	91.2	138.3	219.3	314.6	698.8
1999.....	12.7	20.5	30.1	47.6	63.1	90.0	139.7	217.7	327.1	658.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Essential (primary) hypertension (I10)										
All races 3/, both sexes										
2002.....	0.5	1.0	1.6	2.5	4.2	6.6	10.9	18.5	37.1	112.0
2001.....	0.4	1.0	1.6	2.7	4.3	6.9	10.1	19.1	35.6	107.4
2000.....	0.5	1.1	1.7	2.6	4.3	6.1	10.4	18.2	33.1	101.1
1999.....	0.4	0.9	1.6	2.5	4.3	6.7	9.7	18.0	33.0	98.1
All races 3/, male										
2002.....	0.7	1.3	2.1	3.3	5.0	7.3	11.9	17.5	32.0	74.2
2001.....	0.5	1.2	2.2	3.4	5.2	7.9	10.9	18.8	30.1	74.8
2000.....	0.6	1.5	2.3	3.3	5.4	6.5	11.7	17.4	29.5	70.3
1999.....	0.5	1.1	1.9	3.2	5.1	7.8	10.7	17.2	29.7	68.1
All races 3/, female										
2002.....	0.4	0.7	1.2	1.8	3.5	6.0	10.2	19.1	40.2	128.3
2001.....	0.4	0.7	1.1	2.0	3.5	6.1	9.4	19.3	38.9	121.1
2000.....	0.4	0.6	1.2	1.9	3.4	5.7	9.4	18.7	35.2	113.6
1999.....	0.3	0.7	1.3	1.9	3.6	5.7	8.9	18.6	34.9	110.2
White, both sexes										
2002.....	0.4	0.7	1.3	2.1	3.7	5.7	9.5	16.4	35.1	108.6
2001.....	0.3	0.7	1.2	2.1	3.6	5.7	8.6	17.6	33.1	104.6
2000.....	0.4	0.8	1.2	2.1	3.5	5.3	9.2	16.2	31.2	97.5
1999.....	0.3	0.7	1.1	2.0	3.6	5.6	8.7	16.0	30.9	94.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Essential (primary) hypertension (I10)										
White, male										
2002.....	0.5	1.0	1.7	2.8	4.4	6.3	10.4	15.6	29.1	71.0
2001.....	0.4	0.9	1.6	2.6	4.4	6.8	9.2	16.9	26.9	71.1
2000.....	0.5	1.1	1.7	2.7	4.3	5.8	10.3	15.6	26.8	66.4
1999.....	0.4	0.9	1.4	2.6	4.3	6.7	9.5	15.3	27.0	62.7
White, female										
2002.....	0.3	0.4	0.9	1.4	3.0	5.1	8.7	17.1	38.7	124.9
2001.....	*	0.4	0.8	1.6	2.7	4.8	8.1	18.1	36.9	118.7
2000.....	0.3	0.4	0.8	1.5	2.8	4.8	8.4	16.7	33.8	110.2
1999.....	*	0.5	0.8	1.4	2.9	4.7	8.0	16.5	33.2	107.0
Black, both sexes										
2002.....	1.5	3.4	4.7	6.9	9.1	15.4	27.5	43.6	67.0	158.6
2001.....	1.6	3.4	4.9	8.3	11.9	18.2	25.8	39.4	66.6	150.8
2000.....	1.4	3.2	5.9	7.9	11.9	14.7	23.6	41.4	57.0	149.2
1999.....	1.4	2.4	6.0	8.0	11.2	16.1	22.2	42.3	58.9	143.2
Black, male										
2002.....	1.8	4.0	5.7	8.8	10.3	17.2	30.4	44.1	73.4	117.9
2001.....	1.7	3.8	6.9	11.3	13.6	20.3	31.3	46.0	72.2	132.8
2000.....	*	4.5	7.7	10.7	15.9	14.6	28.4	39.7	64.9	123.4
1999.....	1.7	2.6	6.3	9.5	13.6	19.0	26.5	41.3	66.3	124.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Essential (primary) hypertension (I10)										
Black, female										
2002.....	*	2.9	3.9	5.4	8.2	14.1	25.5	43.4	63.7	174.2
2001.....	1.5	3.0	3.1	5.9	10.6	16.7	22.2	35.5	63.8	157.5
2000.....	1.4	2.1	4.5	5.5	8.7	14.8	20.4	42.4	53.1	158.8
1999.....	*	2.1	5.7	6.8	9.3	14.0	19.4	42.9	55.3	150.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)										
All races 3/, both sexes										
2002.....	3.5	5.9	8.6	11.1	14.3	18.9	24.0	37.9	66.0	187.0
2001.....	3.2	5.1	7.7	9.7	12.5	16.3	23.5	36.6	66.0	189.5
2000.....	2.9	4.7	6.6	9.3	13.2	16.5	23.0	36.3	66.1	193.1
1999.....	2.8	4.5	6.2	9.6	11.9	16.3	22.1	36.4	64.7	184.9
All races 3/, male										
2002.....	4.9	8.0	11.9	15.2	19.0	23.6	27.0	41.0	62.2	134.6
2001.....	4.5	7.0	10.3	12.8	16.5	20.5	27.1	38.9	63.5	134.4
2000.....	3.9	6.3	9.4	12.1	17.4	20.7	27.2	37.3	61.8	139.1
1999.....	3.9	6.1	8.1	12.7	15.0	19.5	24.6	39.4	59.2	133.6
All races 3/, female										
2002.....	2.2	3.9	5.4	7.2	10.0	14.8	21.6	35.8	68.2	209.7
2001.....	2.0	3.2	5.1	6.8	8.9	12.7	20.6	35.1	67.5	212.6
2000.....	2.0	3.2	3.8	6.6	9.3	12.9	19.6	35.7	68.6	215.1
1999.....	1.7	3.0	4.3	6.8	9.2	13.5	20.2	34.3	67.9	205.5
White, both sexes										
2002.....	2.4	4.3	6.0	8.1	10.5	14.4	19.3	31.9	58.1	177.3
2001.....	2.1	3.3	5.2	6.9	9.1	12.1	18.5	30.9	59.2	179.9
2000.....	2.0	3.1	4.3	6.2	9.4	12.5	17.6	30.2	58.4	181.8
1999.....	1.8	2.9	4.1	6.5	8.7	12.0	16.9	30.0	57.3	175.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)										
White, male										
2002.....	3.7	6.1	8.9	11.6	14.0	18.1	22.0	34.5	54.8	124.8
2001.....	3.1	4.8	7.5	9.6	12.5	15.9	21.9	32.9	56.1	123.7
2000.....	2.8	4.6	6.5	8.6	12.7	16.0	21.5	31.1	53.4	129.3
1999.....	2.7	4.2	5.8	8.9	11.3	14.4	19.1	32.4	51.8	123.7
White, female										
2002.....	1.2	2.4	3.3	4.7	7.1	11.1	17.1	30.0	60.1	200.1
2001.....	1.0	1.8	3.0	4.4	6.0	8.8	15.8	29.5	61.1	203.5
2000.....	1.2	1.6	2.2	4.0	6.3	9.4	14.5	29.6	61.3	203.1
1999.....	0.8	1.6	2.5	4.2	6.3	10.0	15.1	28.2	60.6	196.0
Black, both sexes										
2002.....	11.4	19.3	30.3	39.7	50.5	61.2	76.5	110.3	167.7	320.4
2001.....	11.5	18.6	28.4	35.7	43.6	55.8	77.8	105.1	154.7	326.1
2000.....	9.9	17.8	25.4	37.6	47.9	54.6	81.6	111.1	164.9	340.0
1999.....	10.2	17.9	24.8	38.0	41.6	57.6	79.0	113.7	161.2	305.1
Black, male										
2002.....	14.6	24.4	39.9	53.1	71.1	79.0	89.8	125.4	167.4	267.0
2001.....	14.8	25.1	36.6	45.1	55.9	69.6	91.6	117.1	171.8	284.8
2000.....	12.7	21.4	35.8	47.9	63.8	70.0	95.9	117.2	178.7	273.3
1999.....	13.1	22.5	31.3	49.5	52.0	73.0	90.2	129.2	166.5	263.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)										
Black, female										
2002.....	8.6	14.9	22.3	28.7	34.4	47.9	67.7	101.3	167.9	340.9
2001.....	8.6	13.0	21.5	28.0	33.9	45.6	68.6	97.9	145.9	341.4
2000.....	7.5	14.6	16.5	29.2	35.3	43.2	72.1	107.4	157.9	364.9
1999.....	7.6	13.8	19.3	28.7	33.4	46.2	71.6	104.4	158.6	320.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive renal disease (I12)										
All races 3/, both sexes										
2002.....	0.5	0.8	1.2	2.0	3.1	5.2	9.7	16.5	29.6	68.4
2001.....	0.6	0.9	1.3	1.7	3.2	5.3	8.9	16.4	29.6	64.5
2000.....	0.5	0.7	1.3	1.8	3.5	5.4	8.7	15.7	29.9	61.8
1999.....	0.5	0.8	1.2	1.6	3.0	5.3	8.9	14.7	26.9	53.9
All races 3/, male										
2002.....	0.6	0.9	1.4	2.2	3.2	5.4	10.3	18.4	32.7	79.2
2001.....	0.8	1.1	1.5	1.8	2.9	5.8	9.0	17.2	31.2	76.0
2000.....	0.7	0.8	1.6	2.1	3.7	5.0	8.9	17.0	33.4	72.0
1999.....	0.6	0.9	1.6	1.8	3.1	5.6	9.4	15.9	29.7	64.3
All races 3/, female										
2002.....	0.4	0.6	1.1	1.9	2.9	5.0	9.3	15.1	27.7	63.7
2001.....	0.4	0.7	1.2	1.7	3.4	5.0	8.8	15.9	28.6	59.7
2000.....	0.4	0.6	1.0	1.6	3.3	5.7	8.5	14.7	27.8	57.7
1999.....	0.4	0.7	0.8	1.4	2.9	5.1	8.5	13.8	25.3	49.8
White, both sexes										
2002.....	0.2	0.3	0.6	1.0	1.7	3.3	7.3	13.3	25.3	61.9
2001.....	0.2	0.4	0.6	0.8	1.8	3.4	6.7	13.3	25.5	58.7
2000.....	0.2	0.2	0.6	0.9	1.9	3.3	6.1	12.7	26.3	55.7
1999.....	0.3	0.3	0.5	0.7	1.4	3.5	6.5	11.5	22.4	47.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive renal disease (I12)										
White, male										
2002.....	0.3	0.4	0.7	1.1	1.9	3.5	7.9	15.2	28.6	74.2
2001.....	0.3	0.5	0.6	0.9	1.5	3.5	7.2	14.5	27.6	71.1
2000.....	0.3	0.3	0.8	1.1	2.2	3.0	6.5	14.3	30.6	66.1
1999.....	0.3	0.4	0.6	0.9	1.5	3.7	7.1	12.8	25.5	58.3
White, female										
2002.....	*	0.3	0.5	0.9	1.5	3.2	6.9	12.0	23.3	56.5
2001.....	*	0.3	0.5	0.8	2.0	3.4	6.4	12.4	24.2	53.5
2000.....	*	*	0.3	0.7	1.7	3.5	5.8	11.5	23.6	51.4
1999.....	*	0.3	0.4	0.6	1.4	3.2	6.0	10.6	20.7	43.5
Black, both sexes										
2002.....	2.3	4.0	6.1	11.5	15.2	21.1	34.1	52.7	84.0	150.3
2001.....	2.8	4.4	7.7	9.7	15.1	22.1	30.4	50.4	81.5	137.3
2000.....	2.8	4.0	6.8	10.1	17.1	25.6	35.2	51.0	73.4	134.2
1999.....	2.6	4.0	6.8	9.0	15.9	21.9	35.0	52.7	79.7	126.3
Black, male										
2002.....	2.9	5.1	7.6	13.0	15.4	23.4	36.7	61.1	90.9	156.5
2001.....	3.8	6.2	9.4	10.5	16.7	28.5	30.6	49.3	85.9	143.0
2000.....	3.9	4.6	7.9	11.9	18.4	25.5	35.4	53.4	72.5	146.5
1999.....	2.9	5.2	10.3	10.0	17.4	24.1	35.6	57.5	88.9	141.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive renal disease (I12)										
Black, female										
2002.....	1.8	2.9	4.9	10.3	15.0	19.4	32.3	47.8	80.4	147.9
2001.....	1.9	2.8	6.2	9.1	13.9	17.5	30.3	51.0	79.3	135.1
2000.....	1.7	3.6	5.8	8.6	16.0	25.7	35.1	49.6	73.9	129.6
1999.....	2.2	2.9	3.8	8.2	14.6	20.2	34.7	49.8	75.2	120.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)										
All races 3/, both sexes										
2002.....	0.2	0.4	0.5	0.9	1.2	1.6	2.8	4.9	9.5	23.5
2001.....	0.3	0.3	0.4	0.8	1.2	1.8	2.8	5.0	9.3	23.3
2000.....	0.3	0.3	0.4	0.8	1.0	1.6	3.0	5.1	9.8	24.3
1999.....	0.3	0.4	0.4	0.9	1.4	2.1	3.0	6.6	12.3	29.1
All races 3/, male										
2002.....	0.2	0.5	0.6	1.1	1.4	1.5	3.4	5.1	9.9	23.1
2001.....	0.4	0.4	0.6	0.9	1.2	2.1	2.9	5.2	9.4	24.2
2000.....	0.3	0.4	0.5	0.9	1.0	1.7	3.4	5.0	10.8	27.8
1999.....	0.4	0.5	0.6	1.0	1.6	2.3	3.2	7.3	13.3	31.3
All races 3/, female										
2002.....	0.2	0.2	0.4	0.7	1.0	1.7	2.4	4.7	9.2	23.7
2001.....	*	*	0.3	0.7	1.3	1.5	2.8	4.9	9.3	22.9
2000.....	0.2	0.2	0.4	0.7	1.0	1.5	2.7	5.1	9.3	22.9
1999.....	0.3	0.3	0.3	0.7	1.2	2.0	2.9	6.1	11.7	28.2
White, both sexes										
2002.....	*	*	0.1	0.3	0.6	0.9	1.9	3.8	8.4	21.0
2001.....	*	*	0.2	0.4	0.7	1.1	2.0	3.9	8.3	21.0
2000.....	0.2	0.1	0.2	0.4	0.5	0.9	2.2	4.0	8.1	22.0
1999.....	*	0.1	0.2	0.4	0.7	1.2	1.9	5.2	10.4	26.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)										
White, male										
2002.....	*	*	*	0.5	0.7	0.8	2.4	4.2	8.8	20.6
2001.....	*	*	0.3	0.4	0.7	1.3	2.0	4.1	8.7	22.7
2000.....	*	*	*	0.4	0.5	1.0	2.5	4.4	9.2	24.4
1999.....	*	*	*	0.5	0.8	1.4	2.0	5.8	12.1	28.8
White, female										
2002.....	*	*	*	*	0.5	0.9	1.5	3.6	8.1	21.1
2001.....	*	*	*	0.3	0.8	0.8	2.0	3.7	8.0	20.3
2000.....	*	*	*	0.4	0.5	0.8	2.0	3.7	7.4	21.1
1999.....	*	*	*	0.4	0.6	1.1	1.8	4.7	9.5	25.0
Black, both sexes										
2002.....	1.1	2.2	3.3	6.0	6.2	8.0	12.3	17.4	23.0	56.1
2001.....	1.4	1.9	2.3	4.4	5.7	8.4	11.8	18.4	22.8	50.1
2000.....	1.2	1.9	2.5	4.1	5.3	8.3	11.9	17.6	31.1	50.4
1999.....	2.1	2.5	2.3	5.0	7.3	10.1	15.4	21.8	35.3	65.4
Black, male										
2002.....	*	3.0	4.7	7.6	7.3	7.2	14.0	17.0	22.9	61.1
2001.....	2.1	3.0	3.5	5.6	6.3	10.7	13.5	18.1	19.3	44.3
2000.....	1.5	2.7	2.9	5.1	5.6	9.4	13.7	13.7	33.7	63.4
1999.....	2.6	3.5	3.4	6.3	9.3	10.6	17.3	23.7	29.6	59.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)										
Black, female										
2002.....	*	1.5	2.1	4.6	5.3	8.5	11.2	17.6	23.1	54.2
2001.....	*	*	*	3.3	5.3	6.8	10.8	18.6	24.5	52.2
2000.....	*	*	2.1	3.2	5.1	7.5	10.7	19.9	29.7	45.5
1999.....	1.6	1.6	*	4.0	5.7	9.6	14.2	20.6	38.0	67.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
All races 3/, both sexes										
2002.....	23.8	46.4	83.5	140.1	226.2	358.4	569.8	940.2	1,611.2	3,775.0
2001.....	23.4	45.6	84.8	142.8	236.6	372.4	591.8	984.3	1,685.2	3,958.2
2000.....	23.6	48.0	86.7	153.9	250.0	390.8	628.0	1,032.5	1,761.7	4,182.5
1999.....	24.2	48.5	90.1	157.2	260.0	420.1	661.9	1,081.1	1,841.7	4,318.1
All races 3/, male										
2002.....	36.5	72.3	130.3	211.4	331.4	508.5	791.5	1,274.8	2,094.2	4,205.3
2001.....	35.8	71.6	131.1	215.5	344.4	525.7	820.3	1,337.5	2,180.0	4,395.8
2000.....	37.0	76.9	134.2	232.8	362.9	550.5	879.6	1,394.4	2,305.0	4,701.7
1999.....	37.9	77.0	140.3	237.7	376.1	595.2	924.6	1,462.5	2,400.4	4,837.5
All races 3/, female										
2002.....	11.2	21.4	38.7	73.1	130.7	228.8	392.6	704.0	1,317.5	3,588.3
2001.....	11.1	20.3	40.3	74.8	138.8	240.5	410.2	737.0	1,388.7	3,775.1
2000.....	10.3	20.0	41.3	80.0	147.7	253.9	429.9	780.5	1,441.2	3,971.0
1999.....	10.7	20.8	42.1	82.1	155.0	270.2	456.7	815.9	1,514.7	4,108.8
White, both sexes										
2002.....	22.9	44.2	79.1	133.3	216.1	346.6	559.2	928.9	1,616.4	3,836.4
2001.....	22.1	43.1	80.3	135.4	225.8	357.9	580.5	972.9	1,689.9	4,022.7
2000.....	22.2	45.3	82.4	146.2	238.5	378.4	615.0	1,019.7	1,768.5	4,257.7
1999.....	22.6	46.0	85.4	149.1	249.5	408.5	647.3	1,070.6	1,849.5	4,389.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
White, male										
2002.....	35.9	70.3	125.6	203.9	320.5	498.1	784.6	1,273.1	2,115.1	4,299.7
2001.....	34.6	69.5	126.4	207.2	333.9	512.8	810.3	1,335.1	2,201.7	4,494.0
2000.....	35.9	74.5	130.0	223.7	352.1	541.2	870.1	1,390.0	2,328.3	4,813.9
1999.....	36.2	74.7	135.0	228.2	364.4	586.0	913.4	1,463.9	2,429.7	4,933.9
White, female										
2002.....	9.8	18.2	33.6	65.7	119.6	213.2	375.5	682.5	1,310.6	3,635.2
2001.....	9.5	17.0	35.2	66.8	126.1	222.0	394.4	716.3	1,380.6	3,825.5
2000.....	8.5	16.4	36.0	72.5	133.8	236.0	410.4	759.2	1,435.5	4,031.8
1999.....	8.9	17.6	37.0	73.8	143.6	253.7	435.9	794.5	1,507.0	4,170.3
Black, both sexes										
2002.....	35.1	72.8	136.5	230.9	362.0	522.5	779.6	1,196.0	1,768.0	3,462.9
2001.....	36.5	72.5	138.8	236.1	376.4	562.5	810.8	1,249.3	1,837.2	3,615.1
2000.....	37.1	78.2	140.3	250.9	392.9	566.1	866.5	1,296.9	1,882.8	3,684.7
1999.....	39.7	76.9	149.3	258.5	395.4	596.2	906.3	1,321.5	1,926.9	3,805.8
Black, male										
2002.....	48.4	101.6	196.9	330.3	509.6	703.3	1,024.8	1,495.0	2,171.1	3,556.2
2001.....	50.8	101.9	199.9	333.7	515.8	743.8	1,093.0	1,582.5	2,244.8	3,775.0
2000.....	51.8	111.6	197.6	366.7	530.3	727.5	1,139.7	1,630.6	2,316.5	3,885.2
1999.....	56.8	110.4	217.6	372.5	553.5	788.8	1,188.9	1,630.3	2,312.4	4,080.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
Black, female										
2002.....	23.5	47.9	85.4	149.3	246.6	388.3	615.0	1,017.3	1,558.1	3,427.2
2001.....	23.9	47.1	86.8	156.0	266.7	428.6	622.8	1,050.3	1,629.6	3,555.7
2000.....	24.1	49.2	91.4	156.0	284.5	447.0	685.1	1,095.7	1,664.3	3,610.0
1999.....	24.7	48.0	91.4	165.5	271.4	453.5	718.8	1,135.4	1,735.9	3,703.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)										
All races 3/, both sexes										
2002.....	9.9	19.0	34.9	58.4	96.2	148.5	227.5	352.5	572.7	1,191.3
2001.....	9.9	19.7	36.5	61.3	100.1	154.1	235.6	374.0	611.5	1,265.4
2000.....	9.9	21.1	37.8	67.0	110.9	166.8	254.7	405.3	650.1	1,338.5
1999.....	10.4	21.5	39.0	69.7	114.9	181.7	271.6	422.9	689.4	1,379.3
All races 3/, male										
2002.....	14.8	29.2	53.7	86.6	139.0	207.6	312.0	466.1	726.6	1,366.4
2001.....	15.0	30.8	55.6	91.3	143.5	214.2	322.0	495.6	776.1	1,461.3
2000.....	15.5	33.6	57.5	99.7	158.6	231.3	349.6	535.2	835.8	1,568.3
1999.....	16.0	33.7	60.9	105.3	164.9	253.5	373.8	564.2	885.1	1,622.1
All races 3/, female										
2002.....	5.1	9.1	17.0	32.0	57.4	97.6	159.8	272.2	479.1	1,115.3
2001.....	4.9	9.0	18.2	33.2	60.7	102.4	167.0	288.8	512.9	1,183.5
2000.....	4.4	8.9	18.9	36.5	67.7	111.5	179.9	314.9	540.6	1,244.8
1999.....	4.9	9.7	18.0	36.5	69.6	120.2	191.7	324.7	574.9	1,281.5
White, both sexes										
2002.....	9.7	18.4	34.2	57.3	94.3	146.4	225.0	349.5	572.7	1,201.2
2001.....	9.5	19.3	35.4	59.8	98.4	150.8	233.3	371.9	612.7	1,277.8
2000.....	9.5	20.6	36.7	65.6	108.9	164.1	251.8	401.7	652.6	1,353.5
1999.....	10.0	21.1	38.3	68.2	113.4	180.6	268.3	421.1	692.5	1,393.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)										
White, male										
2002.....	14.7	29.0	53.3	86.2	138.7	207.7	313.1	467.1	730.8	1,386.1
2001.....	14.9	30.9	55.1	90.6	143.7	213.3	321.9	497.6	784.0	1,481.1
2000.....	15.3	33.6	57.2	98.8	158.3	231.1	349.3	536.3	843.5	1,592.9
1999.....	15.8	33.8	60.8	104.0	165.1	254.3	374.0	568.8	896.6	1,644.0
White, female										
2002.....	4.6	8.0	15.4	29.7	53.3	92.5	153.2	265.3	475.8	1,120.8
2001.....	4.2	7.7	16.1	30.3	56.7	96.0	161.5	282.8	509.1	1,192.7
2000.....	3.7	7.7	16.6	34.1	63.5	105.5	173.6	307.0	539.1	1,256.2
1999.....	4.3	8.4	16.2	34.2	65.8	116.3	184.3	317.3	572.1	1,292.4
Black, both sexes										
2002.....	13.9	27.7	50.4	84.2	136.3	193.6	295.9	434.1	643.9	1,201.2
2001.....	14.3	27.8	53.2	89.2	135.9	211.1	304.3	452.3	671.2	1,239.9
2000.....	15.1	29.6	55.5	94.7	151.4	219.0	328.8	490.6	698.8	1,275.7
1999.....	15.4	30.0	54.9	96.9	150.2	224.0	347.5	495.1	729.6	1,318.8
Black, male										
2002.....	19.0	37.1	70.4	114.5	177.6	247.8	370.2	531.4	776.4	1,280.1
2001.....	18.5	36.4	71.2	120.1	174.5	264.8	386.4	548.7	792.2	1,382.0
2000.....	20.6	41.0	72.7	131.4	196.7	270.9	419.2	592.5	856.5	1,418.5
1999.....	21.1	41.1	76.9	138.8	198.8	290.8	435.5	591.7	858.0	1,498.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)										
Black, female										
2002.....	9.3	19.5	33.4	59.4	104.1	153.4	246.0	376.0	574.9	1,171.0
2001.....	10.5	20.3	37.8	63.8	105.6	171.4	249.5	394.7	609.5	1,187.2
2000.....	10.1	19.8	40.8	64.6	115.7	180.7	268.8	429.2	619.4	1,222.5
1999.....	10.4	20.5	36.2	62.6	112.1	174.5	289.2	436.8	665.9	1,251.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)										
All races 3/, both sexes										
2002.....	0.3	0.6	1.2	1.4	2.2	3.0	3.9	5.7	9.0	21.1
2001.....	0.3	0.6	1.0	1.4	2.4	2.9	4.1	5.3	9.2	21.5
2000.....	0.3	0.5	1.1	1.6	2.5	3.1	4.0	5.9	9.4	21.9
1999.....	0.3	0.5	0.9	1.6	2.2	3.1	4.3	5.5	9.2	21.9
All races 3/, male										
2002.....	0.4	0.8	1.8	2.1	3.0	4.3	5.1	7.0	11.3	21.4
2001.....	0.4	0.9	1.6	2.0	3.5	4.1	5.5	6.7	11.3	20.7
2000.....	0.4	0.8	1.6	2.5	3.5	4.4	5.2	7.6	11.7	20.7
1999.....	0.4	0.7	1.3	2.5	3.1	4.3	5.6	7.6	10.8	22.3
All races 3/, female										
2002.....	*	0.3	0.7	0.7	1.4	1.9	3.0	4.7	7.6	21.0
2001.....	*	0.3	0.5	0.9	1.3	2.0	3.1	4.4	8.0	21.8
2000.....	*	*	0.5	0.8	1.5	1.9	2.9	4.8	8.1	22.4
1999.....	*	0.2	0.6	0.7	1.5	2.0	3.2	4.1	8.3	21.7
White, both sexes										
2002.....	0.2	0.5	1.1	1.3	2.0	2.9	3.8	5.6	8.9	21.4
2001.....	0.2	0.5	0.9	1.3	2.1	2.7	4.0	5.2	9.2	21.5
2000.....	0.2	0.4	1.0	1.4	2.2	2.9	3.8	5.6	9.3	22.1
1999.....	0.2	0.3	0.8	1.4	2.1	3.0	4.0	5.4	8.9	22.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)										
White, male										
2002.....	0.4	0.7	1.6	2.0	2.8	4.1	4.8	7.0	11.1	21.9
2001.....	0.3	0.8	1.3	1.8	3.1	3.5	5.4	6.5	11.6	20.7
2000.....	0.4	0.7	1.5	2.2	3.2	4.0	4.9	7.2	11.6	21.0
1999.....	0.3	0.5	1.2	2.2	2.8	4.1	5.2	7.6	10.3	22.9
White, female										
2002.....	*	0.3	0.6	0.7	1.2	1.8	2.9	4.6	7.5	21.1
2001.....	*	0.2	0.4	0.8	1.2	1.9	2.9	4.3	7.7	21.7
2000.....	*	*	0.5	0.6	1.3	1.9	2.8	4.5	8.0	22.6
1999.....	*	*	0.5	0.7	1.5	2.0	3.0	3.8	8.1	22.1
Black, both sexes										
2002.....	*	1.1	2.4	2.1	4.1	5.0	7.1	7.7	11.8	21.1
2001.....	0.7	1.1	2.1	2.8	5.2	5.3	6.2	6.9	10.8	24.0
2000.....	*	1.0	2.1	3.8	4.7	5.2	6.1	10.0	12.1	21.9
1999.....	*	1.5	2.2	3.3	3.9	4.3	7.3	8.7	13.9	17.2
Black, male										
2002.....	*	1.9	3.6	3.1	5.7	8.0	10.7	9.2	16.0	21.4
2001.....	*	*	3.6	3.4	8.1	9.1	7.7	11.2	*	23.8
2000.....	*	*	3.6	5.8	6.6	8.3	9.0	13.2	*	*
1999.....	*	2.5	2.8	5.8	6.6	6.8	10.2	10.0	17.4	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute ischemic heart diseases (I24)

Black, female

2002.....	*	*	*	*	*	*	4.7	6.9	9.5	20.9
2001.....	*	*	*	*	*	*	5.2	*	11.9	24.0
2000.....	*	*	*	*	3.3	*	*	8.0	10.6	22.3
1999.....	*	*	*	*	*	*	5.4	8.0	12.1	16.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)										
All races 3/, both sexes										
2002.....	13.6	26.8	47.3	80.2	127.8	206.8	338.4	582.1	1,029.5	2,562.5
2001.....	13.2	25.2	47.3	80.1	134.1	215.4	352.0	605.0	1,064.5	2,671.3
2000.....	13.4	26.5	47.9	85.2	136.6	221.0	369.4	621.2	1,102.2	2,822.1
1999.....	13.5	26.5	50.2	85.9	142.9	235.3	386.1	652.6	1,143.1	2,917.0
All races 3/, male										
2002.....	21.4	42.2	74.9	122.7	189.4	296.6	474.3	801.6	1,356.2	2,817.4
2001.....	20.4	40.0	74.0	122.2	197.4	307.4	492.9	835.3	1,392.6	2,913.8
2000.....	21.1	42.5	75.1	130.6	200.8	314.8	524.7	851.5	1,457.6	3,112.7
1999.....	21.5	42.6	78.1	129.9	208.1	337.3	545.2	890.7	1,504.6	3,193.1
All races 3/, female										
2002.....	6.0	11.9	21.0	40.3	72.0	129.4	229.7	427.1	830.8	2,451.9
2001.....	6.1	11.0	21.7	40.6	76.8	136.2	240.1	443.8	867.9	2,569.8
2000.....	5.8	10.9	21.8	42.7	78.5	140.5	247.0	460.8	892.5	2,703.8
1999.....	5.6	10.9	23.4	44.9	83.9	148.1	261.7	487.1	931.6	2,805.7
White, both sexes										
2002.....	13.0	25.2	43.9	74.7	119.8	197.3	330.5	573.8	1,034.7	2,613.9
2001.....	12.3	23.3	44.0	74.3	125.3	204.5	343.2	595.8	1,068.1	2,723.5
2000.....	12.5	24.3	44.8	79.2	127.3	211.4	359.4	612.4	1,106.6	2,882.1
1999.....	12.3	24.6	46.3	79.4	134.0	225.0	375.0	644.2	1,148.1	2,973.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)										
White, male										
2002.....	20.8	40.6	70.7	115.8	179.1	286.2	466.8	799.0	1,373.2	2,891.6
2001.....	19.4	37.7	69.9	114.8	187.1	296.0	483.0	831.0	1,406.1	2,992.2
2000.....	20.3	40.1	71.3	122.7	190.6	306.1	515.9	846.5	1,473.2	3,200.0
1999.....	20.1	40.4	72.9	122.0	196.5	327.6	534.1	887.5	1,522.9	3,267.1
White, female										
2002.....	5.1	10.0	17.6	35.3	65.1	118.9	219.3	412.6	827.2	2,493.2
2001.....	5.2	9.0	18.7	35.8	68.2	124.1	230.0	429.2	863.8	2,611.0
2000.....	4.7	8.6	18.8	37.8	69.0	128.7	233.9	447.7	888.4	2,753.0
1999.....	4.5	9.0	20.3	38.9	76.3	135.5	248.5	473.4	926.8	2,855.7
Black, both sexes										
2002.....	20.7	44.0	83.7	144.6	221.6	323.9	476.6	754.2	1,112.3	2,240.6
2001.....	21.5	43.7	83.6	144.1	235.3	346.1	500.3	790.1	1,155.2	2,351.2
2000.....	21.5	47.6	82.7	152.4	236.7	341.9	531.5	796.3	1,171.8	2,387.1
1999.....	23.8	45.4	92.2	158.4	241.2	367.9	551.4	817.7	1,183.5	2,469.8
Black, male										
2002.....	29.0	62.5	122.8	212.7	326.3	447.5	643.9	954.4	1,378.6	2,254.6
2001.....	31.2	64.0	125.1	210.1	333.1	469.8	698.8	1,022.7	1,443.9	2,369.1
2000.....	30.3	69.2	121.3	229.5	326.9	448.3	711.4	1,024.9	1,444.9	2,446.0
1999.....	34.9	66.9	138.0	227.8	348.1	491.2	743.2	1,028.6	1,437.0	2,561.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)										
Black, female										
2002.....	13.3	28.0	50.5	88.6	139.7	232.2	364.3	634.5	973.6	2,235.2
2001.....	13.0	26.1	48.2	89.9	158.2	254.8	368.1	651.2	1,008.1	2,344.5
2000.....	13.7	28.8	49.7	89.2	165.6	263.4	412.0	658.5	1,034.3	2,365.2
1999.....	14.1	26.8	53.5	101.6	157.4	276.5	424.2	690.6	1,057.9	2,435.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)										
All races 3/, both sexes										
2002.....	4.8	9.8	17.5	27.9	39.7	53.9	76.5	116.6	187.6	474.5
2001.....	4.6	8.8	17.4	27.7	40.6	55.3	76.7	121.8	197.2	493.6
2000.....	4.4	9.4	17.5	28.5	40.7	55.7	81.9	125.0	207.8	534.6
1999.....	4.7	9.1	17.4	28.6	43.4	59.6	86.8	131.6	218.2	563.4
All races 3/, male										
2002.....	7.8	15.6	28.4	44.8	59.7	78.8	106.7	154.5	232.4	491.8
2001.....	7.2	14.6	28.3	43.4	61.8	79.0	107.6	160.2	247.4	502.0
2000.....	7.1	15.5	28.4	45.8	61.3	80.8	115.6	167.1	252.8	550.8
1999.....	7.6	14.9	28.0	45.0	64.6	87.3	120.8	172.7	264.3	578.6
All races 3/, female										
2002.....	1.8	4.1	7.1	12.1	21.4	32.4	52.5	89.7	160.4	467.1
2001.....	2.1	3.3	7.0	12.9	21.4	34.9	52.1	94.9	167.1	490.1
2000.....	1.8	3.5	7.2	12.4	22.1	34.1	55.4	95.7	181.2	528.0
1999.....	1.8	3.5	7.3	13.4	24.1	36.0	60.2	103.1	191.3	557.2
White, both sexes										
2002.....	4.4	9.1	15.6	24.7	35.3	47.3	69.7	107.8	180.5	470.6
2001.....	4.2	7.9	15.6	23.9	35.0	47.9	69.4	112.2	190.6	489.7
2000.....	4.0	8.3	15.7	24.8	35.2	49.5	73.7	116.0	198.5	529.5
1999.....	4.2	8.0	15.2	24.6	37.5	52.3	77.9	121.5	209.8	557.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)										
White, male										
2002.....	7.4	14.7	25.7	40.0	53.6	69.4	98.0	143.3	222.2	488.0
2001.....	6.6	13.3	25.7	38.3	54.0	69.3	97.6	148.7	239.9	498.6
2000.....	6.6	14.0	25.9	40.0	54.2	72.8	104.4	154.6	241.6	545.6
1999.....	7.0	13.2	24.8	39.4	56.2	77.6	109.0	159.8	253.9	571.2
White, female										
2002.....	1.5	3.5	5.6	10.2	18.4	27.8	46.6	82.5	154.9	463.0
2001.....	1.8	2.6	5.7	10.1	17.6	29.1	46.7	86.4	160.8	486.0
2000.....	1.4	2.7	5.8	10.2	17.6	29.0	49.0	88.8	172.9	522.9
1999.....	1.3	2.8	5.8	10.6	20.3	30.3	53.1	94.6	183.8	552.0
Black, both sexes										
2002.....	8.1	17.0	35.9	60.5	85.5	119.5	157.5	231.9	300.4	575.5
2001.....	8.2	16.8	35.7	64.2	94.4	128.0	162.6	244.0	301.5	594.5
2000.....	7.4	19.4	36.4	65.4	94.0	119.6	176.8	239.7	340.0	646.6
1999.....	9.1	19.1	38.4	67.8	98.2	133.9	188.9	261.7	338.9	668.6
Black, male										
2002.....	11.8	25.5	55.0	97.1	130.5	180.0	220.3	321.0	417.3	599.3
2001.....	12.4	25.7	56.0	97.0	144.0	183.5	241.0	330.4	388.5	609.3
2000.....	11.0	30.3	56.8	106.4	137.1	168.6	257.1	338.3	425.7	701.2
1999.....	13.7	29.9	59.7	103.8	147.9	197.3	271.0	350.0	435.1	715.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)										
Black, female										
2002.....	4.8	9.7	19.7	30.4	50.3	74.6	115.4	178.7	239.5	566.4
2001.....	4.5	9.0	18.4	37.3	55.4	87.1	110.3	192.5	257.1	589.0
2000.....	4.2	10.0	19.0	31.8	60.0	83.5	123.4	180.3	296.8	626.3
1999.....	5.0	9.8	20.4	38.4	59.2	87.0	134.3	208.5	291.3	650.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
All races 3/, both sexes										
2002.....	8.8	17.1	29.9	52.3	88.2	152.9	261.8	465.6	841.9	2,088.0
2001.....	8.6	16.4	29.8	52.4	93.5	160.1	275.3	483.2	867.3	2,177.7
2000.....	9.0	17.0	30.3	56.7	95.9	165.3	287.4	496.2	894.4	2,287.5
1999.....	8.8	17.4	32.8	57.3	99.5	175.7	299.3	521.0	924.9	2,353.6
All races 3/, male										
2002.....	13.6	26.6	46.5	78.0	129.6	217.8	367.6	647.1	1,123.8	2,325.6
2001.....	13.2	25.4	45.7	78.8	135.6	228.4	385.2	675.1	1,145.1	2,411.8
2000.....	14.0	27.0	46.7	84.9	139.5	234.0	409.1	684.5	1,204.8	2,561.9
1999.....	13.9	27.7	50.2	84.9	143.5	250.0	424.4	718.0	1,240.3	2,614.4
All races 3/, female										
2002.....	4.2	7.8	13.9	28.2	50.5	97.0	177.3	337.3	670.4	1,984.9
2001.....	4.0	7.6	14.7	27.7	55.4	101.2	188.0	348.9	700.7	2,079.8
2000.....	4.0	7.3	14.6	30.3	56.4	106.3	191.6	365.1	711.3	2,175.8
1999.....	3.9	7.4	16.1	31.5	59.7	112.1	201.5	384.0	740.3	2,248.4
White, both sexes										
2002.....	8.6	16.1	28.3	49.9	84.5	150.0	260.7	466.0	854.2	2,143.3
2001.....	8.1	15.4	28.4	50.5	90.2	156.6	273.8	483.6	877.5	2,233.8
2000.....	8.5	16.0	29.1	54.4	92.2	162.0	285.8	496.4	908.0	2,352.7
1999.....	8.2	16.6	31.1	54.8	96.5	172.6	297.1	522.7	938.2	2,416.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
White, male										
2002.....	13.4	25.9	45.0	75.8	125.4	216.8	368.7	655.7	1,151.0	2,403.6
2001.....	12.8	24.4	44.2	76.5	133.1	226.8	385.4	682.4	1,166.2	2,493.6
2000.....	13.7	26.1	45.5	82.7	136.4	233.3	411.5	691.9	1,231.6	2,654.4
1999.....	13.1	27.2	48.1	82.6	140.3	249.9	425.1	727.7	1,269.0	2,695.9
White, female										
2002.....	3.6	6.5	12.0	25.2	46.7	91.1	172.7	330.1	672.3	2,030.2
2001.....	3.5	6.4	13.0	25.7	50.7	95.0	183.3	342.8	703.0	2,125.0
2000.....	3.3	5.9	13.0	27.5	51.4	99.6	184.9	358.9	715.6	2,230.1
1999.....	3.2	6.2	14.5	28.3	56.1	105.2	195.4	378.8	743.0	2,303.8
Black, both sexes										
2002.....	12.6	27.0	47.8	84.1	136.1	204.4	319.1	522.2	811.9	1,665.1
2001.....	13.3	26.9	47.9	79.9	140.9	218.1	337.8	546.0	853.7	1,756.7
2000.....	14.1	28.1	46.3	87.0	142.8	222.3	354.7	556.6	831.8	1,740.5
1999.....	14.7	26.3	53.8	90.6	143.1	233.9	362.6	555.9	844.6	1,801.2
Black, male										
2002.....	17.2	37.0	67.8	115.7	195.8	267.5	423.6	633.4	961.4	1,655.3
2001.....	18.8	38.3	69.2	113.1	189.2	286.4	457.8	692.3	1,055.4	1,759.8
2000.....	19.3	39.0	64.5	123.2	189.9	279.7	454.3	686.6	1,019.2	1,744.8
1999.....	21.2	37.0	78.2	124.1	200.3	293.9	472.2	678.6	1,001.9	1,845.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
Black, female										
2002.....	8.5	18.4	30.8	58.2	89.4	157.6	248.9	455.8	734.1	1,668.8
2001.....	8.5	17.0	29.8	52.6	102.9	167.6	257.8	458.7	751.0	1,755.5
2000.....	9.5	18.7	30.7	57.4	105.6	179.9	288.6	478.2	737.5	1,738.9
1999.....	9.1	17.1	33.1	63.3	98.2	189.5	289.9	482.0	766.6	1,784.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)										
All races 3/, both sexes										
2002.....	12.4	18.1	25.9	39.4	60.6	97.9	164.5	286.7	536.3	1,443.5
2001.....	11.9	17.9	25.9	39.7	62.0	100.7	164.7	287.2	537.0	1,475.5
2000.....	12.3	17.6	25.9	40.2	64.3	103.6	167.1	290.2	542.8	1,509.4
1999.....	12.5	17.8	25.9	41.8	66.5	105.2	175.0	295.4	553.5	1,512.9
All races 3/, male										
2002.....	15.9	23.5	33.6	49.4	75.7	118.8	199.8	340.7	612.4	1,448.6
2001.....	14.9	23.1	32.7	49.4	77.7	121.5	200.9	344.7	622.1	1,473.4
2000.....	15.4	23.0	33.4	52.0	80.8	126.9	205.9	345.1	626.8	1,549.1
1999.....	15.6	23.0	33.3	54.1	84.3	133.4	217.0	350.9	641.4	1,548.9
All races 3/, female										
2002.....	9.0	12.8	18.5	30.1	46.8	79.8	136.3	248.6	490.0	1,441.3
2001.....	8.9	12.8	19.5	30.6	47.8	82.8	135.9	247.0	486.0	1,476.4
2000.....	9.2	12.3	18.7	29.2	49.3	83.6	136.6	252.1	493.2	1,493.2
1999.....	9.4	12.8	18.9	30.2	50.5	81.0	142.2	256.8	502.0	1,498.5
White, both sexes										
2002.....	10.7	15.0	22.1	35.0	55.3	92.4	158.5	282.4	539.1	1,475.5
2001.....	9.7	14.9	22.3	34.8	57.2	94.1	159.0	281.6	538.8	1,509.6
2000.....	10.2	14.6	22.1	35.9	59.0	97.3	160.6	284.5	545.7	1,541.6
1999.....	10.2	14.6	22.2	36.3	60.4	98.6	167.6	288.5	553.7	1,543.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)										
White, male										
2002.....	14.1	19.9	29.2	44.1	69.9	112.5	193.0	336.5	617.5	1,482.1
2001.....	12.7	19.7	28.9	43.5	72.1	114.9	193.7	338.2	621.9	1,503.4
2000.....	13.0	19.7	28.6	46.9	74.8	119.7	199.2	339.4	630.8	1,580.5
1999.....	13.3	19.4	28.7	47.5	76.9	125.6	209.0	344.0	641.4	1,571.5
White, female										
2002.....	7.3	10.2	15.2	26.2	41.8	74.7	130.3	243.7	491.0	1,472.6
2001.....	6.6	10.1	15.8	26.5	43.5	75.9	130.9	241.5	488.5	1,512.2
2000.....	7.4	9.6	15.8	25.4	44.3	77.7	129.5	245.9	495.2	1,525.8
1999.....	7.2	9.8	15.8	25.7	45.1	75.0	134.8	249.5	501.9	1,531.9
Black, both sexes										
2002.....	26.6	43.6	60.8	87.1	121.2	169.4	255.8	386.7	594.7	1,270.4
2001.....	29.2	43.1	60.9	91.4	118.8	179.7	254.5	402.9	601.8	1,277.7
2000.....	28.5	43.3	62.0	86.2	122.8	181.5	266.8	406.9	590.7	1,306.0
1999.....	30.2	44.9	62.2	96.8	133.5	185.5	284.2	417.4	621.2	1,302.8
Black, male										
2002.....	32.0	56.7	77.1	110.7	150.7	208.7	317.5	454.2	675.6	1,238.3
2001.....	33.7	53.5	73.7	117.2	151.5	212.9	318.8	494.8	743.2	1,349.1
2000.....	34.8	54.3	82.1	112.1	153.6	224.8	318.9	480.5	696.3	1,402.3
1999.....	36.2	56.1	81.9	127.0	172.7	237.8	353.1	489.1	747.3	1,454.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)										
Black, female										
2002.....	21.8	32.3	47.0	67.7	98.2	140.2	214.4	346.3	552.6	1,282.7
2001.....	25.3	34.0	50.1	70.3	93.1	155.1	211.6	348.1	529.8	1,251.2
2000.....	22.9	33.9	44.8	64.9	98.4	149.6	232.2	362.6	537.5	1,270.2
1999.....	24.9	35.2	45.4	72.1	102.8	146.8	238.5	374.2	558.7	1,246.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)										
All races 3/, both sexes										
2002.....	1.9	2.9	3.8	5.9	8.1	12.7	17.5	25.7	35.5	56.3
2001.....	2.1	2.9	3.9	5.9	8.2	12.0	17.3	24.2	34.8	57.5
2000.....	2.1	2.8	4.2	5.2	8.5	12.3	16.8	24.1	34.7	57.4
1999.....	2.1	2.8	3.9	5.6	8.0	11.9	17.9	25.7	36.5	59.6
All races 3/, male										
2002.....	1.6	2.8	3.5	5.3	8.0	12.3	16.3	23.8	33.8	52.1
2001.....	1.9	2.6	3.4	5.1	8.2	11.7	17.6	23.1	35.5	55.6
2000.....	1.7	2.7	4.0	4.9	8.3	11.5	16.6	25.1	33.5	54.8
1999.....	1.8	2.5	3.6	5.3	8.6	12.0	18.5	25.5	35.2	60.3
All races 3/, female										
2002.....	2.2	3.1	4.1	6.4	8.3	13.0	18.4	27.1	36.6	58.1
2001.....	2.2	3.1	4.4	6.6	8.2	12.3	17.1	24.9	34.4	58.2
2000.....	2.5	3.0	4.4	5.5	8.8	12.9	17.1	23.5	35.3	58.5
1999.....	2.3	3.1	4.1	5.9	7.5	11.8	17.4	25.8	37.2	59.2
White, both sexes										
2002.....	1.7	2.6	3.5	5.3	7.7	12.0	16.7	24.8	35.1	56.1
2001.....	1.8	2.5	3.6	5.3	7.7	11.2	16.7	23.2	34.6	56.8
2000.....	1.8	2.5	3.7	4.9	7.9	11.7	16.1	23.3	34.2	56.4
1999.....	1.7	2.4	3.5	4.9	7.2	11.0	16.7	24.7	35.3	59.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)										
White, male										
2002.....	1.5	2.5	3.5	4.8	7.9	11.8	15.9	23.2	33.1	51.9
2001.....	1.7	2.3	3.3	4.5	7.7	11.2	17.1	22.2	34.8	53.6
2000.....	1.4	2.5	3.6	4.7	7.8	11.3	16.2	24.6	33.0	52.9
1999.....	1.6	2.3	3.3	4.9	7.8	11.2	17.4	24.7	33.8	60.3
White, female										
2002.....	1.9	2.7	3.5	5.8	7.4	12.1	17.4	25.9	36.3	58.0
2001.....	1.9	2.7	3.9	6.1	7.7	11.2	16.3	23.9	34.4	58.1
2000.....	2.1	2.6	3.8	5.2	8.1	12.1	16.0	22.5	35.0	57.8
1999.....	1.9	2.6	3.6	5.0	6.7	10.9	16.1	24.7	36.2	58.8
Black, both sexes										
2002.....	3.9	6.4	7.3	12.8	15.0	22.6	30.3	43.6	49.5	68.6
2001.....	4.5	6.3	7.4	12.4	15.3	23.8	28.5	39.8	45.2	77.7
2000.....	4.9	5.8	9.3	8.9	16.3	20.9	29.2	39.3	46.9	76.6
1999.....	4.9	6.2	8.6	12.6	17.4	23.0	35.9	43.7	57.8	72.1
Black, male										
2002.....	2.9	6.2	4.9	11.9	11.7	21.7	25.0	37.7	51.2	65.4
2001.....	3.8	5.5	5.2	11.9	16.3	21.1	28.6	38.1	54.6	95.3
2000.....	3.9	4.8	8.5	9.2	15.9	17.2	26.4	38.7	50.6	92.3
1999.....	3.9	5.0	7.9	10.3	18.9	23.9	38.3	43.2	62.8	71.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)										
Black, female										
2002.....	4.9	6.7	9.3	13.6	17.7	23.3	33.8	47.2	48.5	69.8
2001.....	5.2	7.0	9.3	12.8	14.6	25.7	28.5	40.8	40.5	71.1
2000.....	5.7	6.7	9.9	8.6	16.7	23.6	31.1	39.6	45.0	70.8
1999.....	5.7	7.3	9.2	14.4	16.3	22.3	34.2	44.1	55.3	72.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pulmonary embolism (I26)										
All races 3/, both sexes										
2002.....	1.3	2.0	2.5	3.9	5.3	8.1	11.0	15.8	22.2	35.4
2001.....	1.4	1.9	2.7	4.1	5.4	7.8	11.4	15.7	22.3	37.8
2000.....	1.4	1.9	2.8	3.5	5.8	7.9	10.8	15.7	22.5	38.4
1999.....	1.4	2.0	2.7	3.9	5.6	7.9	12.6	17.9	25.8	42.0
All races 3/, male										
2002.....	1.1	2.0	2.4	3.8	5.5	8.2	10.8	15.0	21.3	34.1
2001.....	1.3	1.8	2.4	3.6	5.6	8.4	11.8	16.1	23.5	38.0
2000.....	1.1	2.0	3.0	3.5	6.0	8.0	11.2	17.4	22.4	37.5
1999.....	1.2	1.9	2.6	4.1	6.3	8.3	13.7	18.1	24.2	42.0
All races 3/, female										
2002.....	1.4	1.9	2.6	4.0	5.1	8.0	11.1	16.4	22.8	36.0
2001.....	1.5	2.1	2.9	4.6	5.2	7.3	11.1	15.4	21.5	37.7
2000.....	1.6	1.9	2.6	3.5	5.6	7.8	10.6	14.5	22.5	38.8
1999.....	1.6	2.1	2.8	3.8	5.0	7.5	11.7	17.8	26.7	42.1
White, both sexes										
2002.....	1.2	1.8	2.3	3.5	5.1	7.7	10.4	15.1	21.7	34.8
2001.....	1.2	1.7	2.5	3.8	5.1	7.4	10.9	15.0	22.1	37.0
2000.....	1.1	1.7	2.5	3.3	5.4	7.4	10.4	15.0	22.0	37.4
1999.....	1.2	1.7	2.4	3.5	5.1	7.3	11.6	17.0	25.1	41.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pulmonary embolism (I26)										
White, male										
2002.....	1.1	1.8	2.4	3.4	5.4	7.8	10.6	14.6	20.6	33.3
2001.....	1.1	1.6	2.4	3.3	5.2	8.1	11.5	15.4	22.9	36.1
2000.....	0.9	1.8	2.7	3.2	5.7	7.8	11.0	16.8	21.6	35.8
1999.....	1.1	1.7	2.3	3.7	5.7	7.6	12.8	17.2	23.0	41.3
White, female										
2002.....	1.3	1.8	2.3	3.6	4.7	7.6	10.3	15.5	22.4	35.5
2001.....	1.3	1.8	2.5	4.4	5.0	6.8	10.5	14.7	21.6	37.4
2000.....	1.4	1.6	2.3	3.3	5.2	7.1	9.9	13.6	22.2	38.1
1999.....	1.3	1.7	2.5	3.4	4.4	7.1	10.6	16.8	26.4	41.8
Black, both sexes										
2002.....	2.4	4.1	5.0	8.5	9.1	14.3	19.4	28.7	33.5	49.0
2001.....	3.1	4.4	5.0	7.9	9.9	14.3	19.2	27.3	30.3	55.9
2000.....	3.3	4.3	6.1	6.0	11.1	14.7	18.5	28.8	33.0	56.6
1999.....	3.2	4.4	6.0	8.8	12.1	15.7	26.4	33.4	40.7	53.2
Black, male										
2002.....	1.8	4.6	3.8	8.6	8.5	14.5	16.0	24.8	35.9	51.5
2001.....	2.9	4.0	3.4	8.4	12.0	14.2	18.2	28.8	40.1	72.6
2000.....	3.0	3.9	7.0	7.4	12.0	12.7	17.1	30.2	38.8	71.5
1999.....	2.6	3.8	5.9	8.8	13.4	18.7	29.2	35.6	49.7	60.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pulmonary embolism (I26)										
Black, female										
2002.....	3.0	3.7	6.1	8.4	9.6	14.2	21.7	31.0	32.2	48.1
2001.....	3.2	4.8	6.4	7.4	8.3	14.3	20.0	26.3	25.3	49.7
2000.....	3.6	4.7	5.4	4.9	10.3	16.1	19.5	27.9	30.1	51.1
1999.....	3.7	5.0	6.1	8.7	11.1	13.5	24.5	32.0	36.3	50.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

All races 3/, both sexes

2002.....	0.6	1.0	1.3	2.0	2.9	4.6	6.5	9.9	13.3	20.9
2001.....	0.7	1.0	1.2	1.8	2.8	4.2	6.0	8.5	12.5	19.6
2000.....	0.7	0.9	1.4	1.7	2.7	4.4	6.0	8.4	12.2	19.0
1999.....	0.7	0.9	1.2	1.7	2.4	4.0	5.4	7.8	10.7	17.5

All races 3/, male

2002.....	0.4	0.7	1.1	1.5	2.5	4.1	5.5	8.8	12.5	18.0
2001.....	0.6	0.9	1.0	1.5	2.6	3.3	5.9	7.0	11.9	17.6
2000.....	0.5	0.7	1.0	1.4	2.3	3.5	5.4	7.7	11.1	17.3
1999.....	0.6	0.7	1.1	1.3	2.3	3.7	4.9	7.4	10.9	18.4

All races 3/, female

2002.....	0.8	1.2	1.5	2.4	3.2	4.9	7.3	10.7	13.8	22.1
2001.....	0.8	1.1	1.5	2.0	3.0	5.0	6.0	9.5	12.9	20.5
2000.....	0.9	1.1	1.8	2.0	3.2	5.2	6.5	8.9	12.8	19.7
1999.....	0.8	1.1	1.4	2.0	2.5	4.3	5.7	8.0	10.5	17.2

White, both sexes

2002.....	0.5	0.8	1.2	1.7	2.6	4.3	6.3	9.7	13.3	21.3
2001.....	0.6	0.8	1.1	1.5	2.6	3.8	5.8	8.2	12.5	19.7
2000.....	0.6	0.9	1.2	1.7	2.5	4.3	5.7	8.4	12.3	19.0
1999.....	0.5	0.7	1.1	1.4	2.1	3.7	5.1	7.7	10.2	17.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)										
White, male										
2002.....	0.4	0.6	1.1	1.3	2.5	3.9	5.3	8.7	12.5	18.6
2001.....	0.5	0.8	0.9	1.3	2.5	3.1	5.6	6.8	11.9	17.4
2000.....	0.5	0.7	1.0	1.5	2.2	3.4	5.2	7.8	11.4	17.1
1999.....	0.4	0.6	1.0	1.3	2.0	3.5	4.6	7.5	10.8	19.0
White, female										
2002.....	0.7	1.0	1.3	2.1	2.7	4.6	7.1	10.4	13.9	22.5
2001.....	0.6	0.9	1.3	1.7	2.7	4.4	5.9	9.2	12.8	20.7
2000.....	0.7	1.0	1.5	1.9	2.8	5.0	6.1	8.8	12.8	19.8
1999.....	0.6	0.8	1.2	1.6	2.2	3.8	5.4	7.8	9.8	17.0
Black, both sexes										
2002.....	1.5	2.3	2.2	4.3	5.9	8.3	10.9	14.9	16.0	19.6
2001.....	1.5	1.9	2.4	4.5	5.4	9.5	9.3	12.5	14.9	21.8
2000.....	1.6	1.5	3.1	2.9	5.3	6.2	10.7	10.5	13.8	20.0
1999.....	1.7	1.8	2.6	3.8	5.3	7.3	9.5	10.4	17.0	18.8
Black, male										
2002.....	*	*	*	3.3	*	7.2	9.0	12.9	15.3	*
2001.....	*	*	*	3.4	4.3	6.9	10.4	9.3	*	22.7
2000.....	*	*	*	*	*	*	9.4	*	*	*
1999.....	*	*	*	*	5.5	5.2	9.2	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)										
Black, female										
2002.....	1.8	3.0	3.3	5.1	8.1	9.1	12.1	16.2	16.3	21.8
2001.....	2.0	2.3	2.9	5.4	6.2	11.5	8.5	14.4	15.1	21.5
2000.....	2.1	2.1	4.5	3.8	6.4	7.5	11.5	11.7	14.9	19.7
1999.....	1.9	2.4	3.1	5.7	5.2	8.9	9.7	12.0	19.0	21.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other non-pulmonary forms of heart disease (I30-I51)										
All races 3/, both sexes										
2002.....	10.5	15.1	22.1	33.6	52.4	85.2	147.1	261.0	500.8	1,387.2
2001.....	9.8	15.0	22.0	33.8	53.8	88.6	147.4	263.0	502.3	1,418.1
2000.....	10.2	14.8	21.7	35.0	55.7	91.3	150.3	266.1	508.1	1,452.0
1999.....	10.4	15.0	22.0	36.2	58.5	93.3	157.1	269.7	517.0	1,453.4
All races 3/, male										
2002.....	14.3	20.7	30.0	44.1	67.8	106.5	183.6	316.9	578.7	1,396.5
2001.....	13.0	20.5	29.3	44.3	69.5	109.7	183.3	321.6	586.6	1,417.8
2000.....	13.7	20.3	29.4	47.1	72.5	115.4	189.3	320.0	593.3	1,494.4
1999.....	13.8	20.5	29.7	48.8	75.7	121.4	198.5	325.4	606.3	1,488.5
All races 3/, female										
2002.....	6.8	9.7	14.4	23.6	38.5	66.8	117.9	221.5	453.4	1,383.1
2001.....	6.7	9.6	15.1	24.0	39.6	70.4	118.8	222.0	451.7	1,418.2
2000.....	6.7	9.3	14.3	23.7	40.6	70.7	119.5	228.6	457.9	1,434.8
1999.....	7.0	9.7	14.7	24.4	42.9	69.2	124.8	231.0	464.8	1,439.2
White, both sexes										
2002.....	9.0	12.4	18.6	29.7	47.6	80.4	141.8	257.7	504.0	1,419.3
2001.....	7.9	12.3	18.7	29.5	49.5	83.0	142.3	258.4	504.2	1,452.9
2000.....	8.5	12.1	18.4	30.9	51.0	85.6	144.5	261.2	511.5	1,485.2
1999.....	8.5	12.1	18.7	31.4	53.2	87.6	151.0	263.8	518.4	1,484.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other non-pulmonary forms of heart disease (I30-I51)										
White, male										
2002.....	12.6	17.4	25.7	39.3	62.0	100.7	177.1	313.3	584.4	1,430.2
2001.....	11.0	17.3	25.6	38.9	64.5	103.7	176.6	316.0	587.2	1,449.9
2000.....	11.6	17.2	25.0	42.2	67.0	108.4	183.0	314.8	597.8	1,527.6
1999.....	11.7	17.1	25.5	42.6	69.2	114.4	191.6	319.3	607.6	1,511.1
White, female										
2002.....	5.4	7.5	11.7	20.5	34.3	62.5	112.9	217.8	454.8	1,414.6
2001.....	4.8	7.4	12.0	20.4	35.8	64.8	114.5	217.6	454.1	1,454.1
2000.....	5.3	7.0	11.9	20.2	36.3	65.6	113.5	223.4	460.2	1,468.0
1999.....	5.4	7.3	12.2	20.8	38.4	64.2	118.7	224.8	465.7	1,473.2
Black, both sexes										
2002.....	22.7	37.2	53.5	74.3	106.2	146.8	225.5	343.1	545.3	1,201.8
2001.....	24.7	36.7	53.5	79.0	103.5	155.9	225.9	363.2	556.6	1,200.0
2000.....	23.6	37.5	52.7	77.3	106.4	160.6	237.6	367.7	543.8	1,229.4
1999.....	25.3	38.6	53.6	84.2	116.1	162.6	248.4	373.7	563.4	1,230.8
Black, male										
2002.....	29.1	50.5	72.3	98.8	139.0	187.0	292.4	416.6	624.4	1,172.9
2001.....	29.9	48.0	68.5	105.3	135.3	191.8	290.2	456.7	688.6	1,253.8
2000.....	30.9	49.4	73.7	102.9	137.7	207.6	292.5	441.8	645.7	1,310.1
1999.....	32.3	51.1	74.0	116.6	153.8	213.9	314.8	445.9	684.5	1,382.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other non-pulmonary forms of heart disease (I30-I51)										
Black, female										
2002.....	17.0	25.6	37.7	54.1	80.5	116.9	180.6	299.1	504.1	1,212.9
2001.....	20.1	27.0	40.7	57.5	78.6	129.3	183.2	307.3	489.4	1,180.0
2000.....	17.2	27.1	34.9	56.3	81.7	126.0	201.1	323.0	492.5	1,199.3
1999.....	19.2	27.9	36.2	57.7	86.5	124.5	204.3	330.1	503.4	1,174.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and subacute endocarditis (I33)										
All races 3/, both sexes										
2002.....	0.3	0.4	0.5	0.5	0.8	1.1	1.5	2.2	2.7	2.6
2001.....	0.3	0.4	0.5	0.5	0.8	1.1	1.7	2.2	3.0	2.5
2000.....	0.3	0.4	0.6	0.7	0.7	1.1	1.3	2.2	2.8	2.8
1999.....	0.3	0.3	0.4	0.7	0.7	1.1	1.5	2.1	2.7	2.9
All races 3/, male										
2002.....	0.4	0.5	0.7	0.6	0.9	1.5	1.7	2.5	3.7	3.5
2001.....	0.3	0.7	0.7	0.6	1.1	1.1	2.1	3.1	4.3	3.8
2000.....	0.4	0.6	0.7	0.8	0.9	1.3	1.6	3.1	4.0	4.2
1999.....	0.4	0.5	0.7	1.0	0.8	1.2	1.7	2.8	3.9	4.7
All races 3/, female										
2002.....	0.2	0.3	0.3	0.4	0.8	0.8	1.3	1.9	2.1	2.2
2001.....	0.3	0.2	0.3	0.5	0.5	1.0	1.3	1.6	2.2	2.0
2000.....	0.2	0.3	0.4	0.5	0.6	0.9	1.1	1.6	2.0	2.3
1999.....	*	*	0.2	0.5	0.7	1.0	1.4	1.7	2.0	2.1
White, both sexes										
2002.....	0.2	0.3	0.4	0.4	0.8	1.0	1.4	2.2	2.9	2.7
2001.....	0.2	0.4	0.4	0.4	0.7	0.9	1.5	2.0	3.0	2.6
2000.....	0.3	0.3	0.4	0.5	0.7	1.1	1.3	2.2	2.7	2.7
1999.....	0.2	0.3	0.3	0.5	0.6	1.0	1.4	2.2	2.8	3.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
All races 3/, both sexes										
2002.....	0.2	0.3	0.3	0.4	0.4	0.6	1.0	1.1	1.4	2.1
2001.....	0.2	0.3	0.3	0.4	0.5	0.5	0.8	1.2	1.3	1.9
2000.....	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.1	1.3	1.8
1999.....	0.3	0.3	0.3	0.5	0.6	0.7	0.9	1.2	1.5	2.0
All races 3/, male										
2002.....	0.3	0.4	0.3	0.5	0.6	0.7	1.1	1.1	1.3	2.4
2001.....	*	0.3	0.3	0.4	0.6	0.6	0.8	1.3	1.1	1.8
2000.....	0.3	0.3	*	0.5	0.4	0.5	1.1	1.4	1.7	2.1
1999.....	0.3	0.3	0.3	0.6	0.6	0.8	0.9	1.3	1.5	2.5
All races 3/, female										
2002.....	*	*	0.2	0.4	*	0.5	1.0	1.1	1.4	2.0
2001.....	0.3	0.3	0.3	0.4	*	*	0.9	1.1	1.4	2.0
2000.....	*	*	0.2	*	0.4	0.5	0.7	0.9	1.0	1.7
1999.....	0.3	0.3	0.2	0.3	0.6	0.7	0.8	1.2	1.5	1.9
White, both sexes										
2002.....	0.2	0.2	0.2	0.4	0.4	0.6	1.0	1.0	1.3	2.2
2001.....	0.2	0.2	0.2	0.3	0.5	0.4	0.9	1.2	1.3	1.9
2000.....	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.0	1.4	1.8
1999.....	0.2	0.3	0.2	0.5	0.6	0.7	0.8	1.2	1.5	2.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of endocardium (I34-I38)										
All races 3/, both sexes										
2002.....	1.0	1.3	1.9	2.9	4.8	8.4	15.9	33.0	72.0	203.9
2001.....	0.8	1.2	2.1	2.8	4.7	9.0	17.3	33.9	70.3	204.2
2000.....	0.9	1.4	1.9	3.0	5.1	8.9	17.8	34.1	73.3	213.2
1999.....	1.1	1.3	2.0	2.9	5.1	9.2	17.6	35.2	76.6	209.8
All races 3/, male										
2002.....	1.1	1.7	2.3	3.3	5.2	8.7	17.3	36.4	74.9	198.5
2001.....	0.9	1.6	2.7	3.3	5.3	9.7	18.9	38.6	75.4	200.8
2000.....	1.0	1.9	2.2	3.8	5.9	9.3	19.2	35.1	78.9	215.8
1999.....	1.2	1.5	2.4	3.5	5.8	10.4	19.6	38.5	86.3	206.4
All races 3/, female										
2002.....	0.8	1.0	1.5	2.5	4.5	8.2	14.8	30.6	70.2	206.3
2001.....	0.6	0.9	1.5	2.4	4.2	8.4	16.1	30.6	67.2	205.6
2000.....	0.7	1.0	1.5	2.4	4.4	8.6	16.8	33.5	69.9	212.1
1999.....	0.9	1.2	1.7	2.4	4.4	8.2	16.0	32.9	70.9	211.2
White, both sexes										
2002.....	0.9	1.1	1.7	2.7	4.8	8.2	16.4	34.7	75.8	216.2
2001.....	0.7	1.1	1.9	2.6	4.6	8.8	17.8	35.2	74.1	216.5
2000.....	0.8	1.3	1.7	2.7	5.0	8.9	18.1	35.1	77.0	225.8
1999.....	1.0	1.1	1.9	2.7	4.9	9.1	17.4	36.4	80.2	221.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of endocardium (I34-I38)										
White, male										
2002.....	1.1	1.4	2.1	3.1	5.1	8.7	17.8	38.2	78.9	211.7
2001.....	0.8	1.4	2.5	3.1	5.2	9.5	19.4	39.9	79.8	212.0
2000.....	1.0	1.7	2.0	3.5	5.9	9.4	19.5	36.2	82.7	229.0
1999.....	1.1	1.1	2.3	3.3	5.6	10.5	19.4	39.9	90.2	218.2
White, female										
2002.....	0.6	0.7	1.3	2.3	4.4	7.8	15.2	32.1	73.9	218.2
2001.....	0.5	0.7	1.4	2.1	4.2	8.2	16.5	31.8	70.6	218.4
2000.....	0.6	0.8	1.4	2.0	4.2	8.4	17.0	34.3	73.6	224.5
1999.....	0.8	1.0	1.4	2.2	4.2	7.9	15.8	33.9	74.4	222.7
Black, both sexes										
2002.....	1.8	3.6	3.8	4.8	6.7	12.1	13.8	22.2	37.2	86.4
2001.....	1.5	2.9	3.9	5.9	6.4	12.2	15.1	25.7	38.2	85.4
2000.....	1.9	3.0	3.6	5.9	7.2	10.9	18.9	29.9	39.5	92.9
1999.....	2.2	3.2	3.7	5.3	7.8	11.4	21.8	25.9	41.9	91.5
Black, male										
2002.....	1.5	4.5	4.0	6.1	7.1	10.7	14.0	24.3	35.2	67.5
2001.....	1.9	3.8	5.1	6.7	7.9	13.0	15.8	29.7	32.9	86.2
2000.....	1.8	4.1	4.5	6.6	7.5	10.4	19.7	29.8	43.0	86.5
1999.....	2.3	3.8	4.1	5.4	9.3	11.2	25.1	27.1	47.1	77.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of endocardium (I34-I38)										
Black, female										
2002.....	2.0	2.8	3.6	3.8	6.3	13.1	13.7	20.9	38.2	93.6
2001.....	*	2.1	2.9	5.1	5.1	11.6	14.6	23.3	40.9	85.0
2000.....	1.9	2.0	2.8	5.3	7.0	11.3	18.4	29.9	37.8	95.3
1999.....	2.2	2.7	3.4	5.1	6.7	11.5	19.6	25.2	39.3	96.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nonrheumatic mitral valve disorders (I34)										
All races 3/, both sexes										
2002.....	0.2	0.3	0.3	0.5	1.0	1.5	2.8	5.0	9.8	22.1
2001.....	0.2	0.3	0.4	0.5	0.7	1.7	2.7	5.7	9.6	22.7
2000.....	0.3	0.3	0.3	0.5	1.0	1.6	3.0	5.8	10.6	24.3
1999.....	0.3	0.3	0.4	0.5	0.9	1.6	3.1	5.8	10.5	25.1
All races 3/, male										
2002.....	0.2	0.2	0.3	0.4	0.9	1.3	2.4	4.2	8.2	21.0
2001.....	0.2	0.3	0.4	0.5	0.7	1.7	2.8	4.7	8.1	19.0
2000.....	0.2	0.4	0.3	0.5	0.8	1.6	2.6	5.6	9.8	22.9
1999.....	0.3	0.3	0.3	0.4	0.9	1.3	3.1	5.6	10.6	22.2
All races 3/, female										
2002.....	0.2	0.3	0.3	0.5	1.1	1.8	3.1	5.6	10.9	22.6
2001.....	0.3	0.3	0.4	0.5	0.8	1.7	2.7	6.3	10.5	24.3
2000.....	0.3	0.2	0.4	0.6	1.1	1.5	3.2	5.9	11.1	24.9
1999.....	0.4	0.4	0.5	0.6	0.9	1.8	3.1	5.9	10.5	26.2
White, both sexes										
2002.....	0.2	0.2	0.3	0.4	1.0	1.5	2.9	5.2	10.4	23.8
2001.....	0.2	0.3	0.4	0.5	0.7	1.6	2.8	6.0	10.1	24.2
2000.....	0.3	0.3	0.3	0.5	1.0	1.6	3.0	6.0	11.3	25.5
1999.....	0.3	0.3	0.3	0.5	0.9	1.5	3.1	5.9	11.1	26.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nonrheumatic aortic valve disorders (I35)										
All races 3/, both sexes										
2002.....	0.3	0.5	0.8	1.2	2.2	4.0	8.8	19.2	45.4	144.8
2001.....	0.2	0.5	0.9	1.2	2.3	4.5	9.2	19.9	44.6	144.1
2000.....	0.3	0.5	0.8	1.4	2.3	4.5	9.9	20.0	47.1	151.3
1999.....	0.3	0.5	0.8	1.3	2.3	4.8	9.3	20.7	49.9	147.8
All races 3/, male										
2002.....	0.4	0.7	1.0	1.5	2.6	4.5	10.8	23.2	51.4	144.9
2001.....	0.3	0.8	1.3	1.5	2.9	5.3	11.0	24.7	50.7	145.9
2000.....	0.4	0.8	1.2	2.1	3.1	5.2	11.7	21.5	52.9	158.2
1999.....	0.4	0.7	1.2	1.7	3.0	5.9	11.7	24.3	58.9	150.3
All races 3/, female										
2002.....	0.2	0.2	0.6	0.8	1.9	3.5	7.1	16.4	41.7	144.8
2001.....	*	0.3	0.5	0.9	1.8	3.8	7.8	16.6	40.9	143.4
2000.....	*	0.2	0.4	0.8	1.6	3.8	8.5	19.0	43.7	148.5
1999.....	0.2	0.4	0.5	0.9	1.7	3.8	7.4	18.1	44.7	146.8
White, both sexes										
2002.....	0.3	0.4	0.7	1.2	2.3	3.9	9.1	20.3	47.9	153.2
2001.....	0.2	0.5	0.9	1.2	2.4	4.6	9.6	20.5	47.3	152.8
2000.....	0.3	0.5	0.8	1.4	2.3	4.4	10.2	20.9	49.5	160.7
1999.....	0.3	0.4	0.8	1.3	2.3	4.8	9.2	21.7	52.3	156.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nonrheumatic aortic valve disorders (I35)										
White, male										
2002.....	0.4	0.6	1.0	1.5	2.7	4.6	11.3	24.5	54.3	154.4
2001.....	0.3	0.7	1.2	1.5	3.0	5.4	11.4	25.2	53.8	153.5
2000.....	0.4	0.7	1.2	2.0	3.2	5.2	12.0	22.6	55.4	168.4
1999.....	0.4	0.6	1.1	1.7	3.0	6.1	11.5	25.6	61.2	159.0
White, female										
2002.....	*	*	0.5	0.9	1.9	3.4	7.4	17.3	44.0	152.7
2001.....	*	0.3	0.5	0.8	2.0	3.9	8.2	17.2	43.3	152.5
2000.....	*	*	0.5	0.7	1.5	3.8	8.7	19.7	46.1	157.6
1999.....	0.2	0.3	0.4	0.8	1.8	3.7	7.4	19.0	47.0	155.0
Black, both sexes										
2002.....	*	0.9	1.1	*	2.2	5.1	6.8	11.5	23.3	63.0
2001.....	*	*	1.3	1.7	2.0	4.4	6.9	16.3	21.4	59.6
2000.....	*	0.8	*	2.1	2.6	5.5	9.1	13.9	25.4	62.3
1999.....	*	1.0	1.4	1.7	2.8	5.0	10.7	12.0	26.9	64.4
Black, male										
2002.....	*	*	*	*	*	5.2	7.7	13.3	22.2	52.5
2001.....	*	*	2.3	*	*	*	8.1	22.8	19.3	68.1
2000.....	*	*	*	*	*	6.2	10.7	13.2	30.3	57.7
1999.....	*	*	*	*	*	5.7	13.9	13.3	36.6	58.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nonrheumatic aortic valve disorders (I35)

Black, female

2002.....	*	*	*	*	*	5.0	6.3	10.4	23.9	66.9
2001.....	*	*	*	*	*	4.1	6.1	12.5	22.5	56.4
2000.....	*	*	*	*	*	5.0	8.0	14.2	22.9	64.0
1999.....	*	*	*	*	*	4.4	8.6	11.2	22.0	66.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of endocardium (I36-I38)										
All races 3/, both sexes										
2002.....	0.4	0.6	0.8	1.2	1.6	2.9	4.4	8.7	16.7	37.0
2001.....	0.3	0.4	0.8	1.1	1.6	2.8	5.3	8.3	16.1	37.4
2000.....	0.3	0.6	0.8	1.1	1.8	2.9	5.0	8.3	15.6	37.6
1999.....	0.4	0.5	0.8	1.1	1.8	2.9	5.2	8.7	16.1	37.0
All races 3/, male										
2002.....	0.5	0.8	1.0	1.3	1.7	2.9	4.1	9.1	15.3	32.6
2001.....	0.4	0.5	1.0	1.3	1.7	2.7	5.1	9.2	16.5	35.9
2000.....	0.4	0.7	0.8	1.2	2.0	2.5	4.9	8.0	16.2	34.7
1999.....	0.5	0.5	0.9	1.3	1.9	3.1	4.9	8.6	16.8	33.9
All races 3/, female										
2002.....	0.4	0.5	0.6	1.1	1.5	3.0	4.6	8.5	17.6	38.9
2001.....	0.2	0.3	0.6	1.0	1.6	2.8	5.5	7.7	15.8	38.0
2000.....	0.2	0.5	0.7	0.9	1.7	3.2	5.0	8.6	15.2	38.7
1999.....	0.3	0.5	0.7	0.9	1.7	2.7	5.5	8.8	15.7	38.2
White, both sexes										
2002.....	0.3	0.4	0.7	1.0	1.4	2.8	4.3	9.2	17.5	39.2
2001.....	0.3	0.3	0.7	0.9	1.5	2.6	5.4	8.6	16.7	39.5
2000.....	0.2	0.5	0.6	0.8	1.7	2.9	4.9	8.2	16.2	39.6
1999.....	0.3	0.3	0.7	0.9	1.6	2.8	5.1	8.8	16.8	38.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of endocardium (I36-I38)										
White, male										
2002.....	0.4	0.5	0.9	1.1	1.5	2.9	4.1	9.4	15.9	34.8
2001.....	0.3	0.3	0.9	1.0	1.5	2.5	5.2	9.7	17.3	38.0
2000.....	0.4	0.6	0.6	1.0	1.8	2.5	4.8	7.8	16.9	36.4
1999.....	0.4	0.3	0.8	1.1	1.7	3.0	4.8	8.6	17.6	35.5
White, female										
2002.....	0.2	0.3	0.4	1.0	1.3	2.7	4.6	9.0	18.5	41.1
2001.....	*	*	0.5	0.8	1.4	2.7	5.5	7.9	16.3	40.2
2000.....	*	0.4	0.6	0.7	1.6	3.2	4.9	8.4	15.7	40.9
1999.....	0.2	0.3	0.6	0.8	1.6	2.5	5.3	8.9	16.4	40.2
Black, both sexes										
2002.....	1.3	2.2	2.1	2.8	3.6	4.9	4.8	6.9	9.7	16.6
2001.....	*	1.6	1.9	3.3	3.5	5.3	5.4	6.9	11.6	18.1
2000.....	1.0	1.5	2.1	2.9	3.6	3.8	7.2	11.0	10.4	18.5
1999.....	1.3	1.6	1.7	2.8	3.7	4.5	7.7	9.5	10.7	16.9
Black, male										
2002.....	*	3.0	*	3.7	*	*	*	*	*	*
2001.....	*	1.9	2.6	4.4	4.3	5.8	*	*	*	*
2000.....	*	1.9	2.4	*	*	*	7.0	12.8	*	*
1999.....	*	*	*	*	4.9	*	*	10.0	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of endocardium (I36-I38)										
Black, female										
2002.....	1.4	1.6	2.2	*	3.6	5.4	5.4	6.3	9.5	19.3
2001.....	*	*	*	*	*	4.9	5.6	7.8	11.9	18.9
2000.....	*	*	*	2.7	3.3	4.0	7.3	10.0	11.0	17.6
1999.....	*	*	*	*	*	4.6	8.6	9.2	11.2	18.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cardiomyopathy (I42)										
All races 3/, both sexes										
2002.....	4.1	5.9	7.2	10.5	13.8	21.1	33.6	49.5	77.3	142.0
2001.....	4.0	5.5	7.3	10.6	14.5	22.9	31.9	48.9	76.8	138.3
2000.....	4.1	5.6	8.0	10.8	15.8	24.1	33.7	51.5	76.2	142.3
1999.....	4.2	5.7	7.9	11.7	16.8	25.6	35.5	50.0	76.1	139.2
All races 3/, male										
2002.....	5.9	8.6	10.4	14.7	19.4	28.9	47.9	71.8	110.8	189.7
2001.....	5.8	7.9	10.0	14.8	20.5	31.0	44.7	69.9	108.8	184.7
2000.....	5.9	7.9	11.5	15.5	21.6	33.1	47.9	74.4	109.5	195.3
1999.....	6.1	8.2	11.3	16.8	23.0	36.1	50.7	70.9	106.8	188.2
All races 3/, female										
2002.....	2.4	3.2	4.1	6.5	8.7	14.3	22.1	33.7	56.9	121.3
2001.....	2.3	3.2	4.7	6.7	9.0	15.9	21.8	34.3	57.6	118.9
2000.....	2.2	3.4	4.7	6.4	10.5	16.4	22.5	35.6	56.5	120.7
1999.....	2.3	3.4	4.7	6.9	11.2	16.6	23.6	35.5	58.2	119.4
White, both sexes										
2002.....	3.3	4.7	5.8	8.9	12.2	19.5	31.7	47.7	76.6	141.7
2001.....	3.0	4.2	5.9	8.8	12.7	21.0	30.3	46.9	75.3	139.2
2000.....	3.2	4.2	6.4	9.0	14.0	21.6	31.7	49.1	74.7	142.4
1999.....	3.1	4.4	6.3	9.6	14.2	23.4	33.1	48.0	74.3	138.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cardiomyopathy (I42)										
White, male										
2002.....	4.8	7.0	8.5	12.6	17.4	26.7	45.3	70.3	110.8	190.9
2001.....	4.5	6.2	8.3	12.4	18.2	28.5	42.7	67.7	107.1	187.0
2000.....	4.6	6.3	9.2	13.0	19.4	29.5	45.4	71.2	108.2	197.5
1999.....	4.7	6.4	9.1	13.8	19.4	33.2	47.2	68.7	105.1	188.0
White, female										
2002.....	1.8	2.4	3.2	5.3	7.4	13.1	20.7	31.6	55.6	120.4
2001.....	1.5	2.2	3.5	5.4	7.6	14.3	20.2	32.2	56.1	119.2
2000.....	1.7	2.3	3.6	5.2	9.0	14.6	20.6	33.5	54.8	120.0
1999.....	1.5	2.4	3.6	5.6	9.4	14.9	21.9	33.4	56.1	118.8
Black, both sexes										
2002.....	10.4	15.2	18.4	27.0	30.4	39.5	57.5	77.3	96.3	161.0
2001.....	11.8	15.5	19.8	27.8	33.3	44.7	54.5	80.9	104.5	141.2
2000.....	10.7	16.6	22.7	28.6	32.2	51.2	59.6	85.5	104.8	151.1
1999.....	12.1	16.6	22.7	30.9	40.5	49.4	64.3	79.1	103.7	153.0
Black, male										
2002.....	14.8	22.4	27.4	38.6	42.0	55.8	85.5	103.8	130.7	204.8
2001.....	16.3	21.4	26.6	40.0	47.6	62.5	73.4	112.9	153.3	180.4
2000.....	16.2	22.0	33.1	42.8	44.5	73.6	82.2	123.3	149.2	193.7
1999.....	16.8	23.5	33.6	47.1	58.1	70.6	96.3	105.0	140.4	215.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cardiomyopathy (I42)										
Black, female										
2002.....	6.5	9.1	10.7	17.5	21.4	27.4	38.8	61.5	78.4	144.2
2001.....	7.8	10.3	14.0	17.8	22.1	31.5	41.9	61.8	79.7	126.7
2000.....	5.8	12.0	13.9	17.0	22.6	34.6	44.6	62.7	82.4	135.2
1999.....	7.9	10.7	13.4	17.7	26.6	33.7	43.0	63.5	85.6	129.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Conduction disorders and cardiac dysrhythmias (I44-I49)										
All races 3/, both sexes										
2002.....	2.6	3.6	6.4	10.1	15.8	24.7	39.0	70.1	126.0	323.2
2001.....	2.2	3.9	6.3	10.0	15.9	25.3	40.5	69.3	125.7	328.2
2000.....	2.5	3.8	5.9	10.5	16.2	26.1	41.0	68.8	127.3	340.2
1999.....	2.5	4.1	6.5	10.8	19.1	27.3	44.8	72.4	134.3	353.7
All races 3/, male										
2002.....	3.4	4.8	8.7	13.5	21.2	31.1	48.1	81.3	138.2	310.5
2001.....	2.9	5.1	8.5	13.1	20.9	31.0	49.7	82.2	140.8	311.6
2000.....	3.0	5.1	8.2	14.3	21.7	34.5	52.3	80.6	137.4	333.2
1999.....	3.1	5.7	8.7	14.7	24.9	36.8	55.1	84.9	154.2	353.4
All races 3/, female										
2002.....	1.8	2.4	4.3	6.9	10.9	19.1	31.8	62.2	118.6	328.8
2001.....	1.5	2.7	4.3	7.0	11.4	20.4	33.3	60.3	116.7	335.2
2000.....	1.9	2.5	3.7	7.0	11.2	18.8	32.2	60.6	121.4	343.0
1999.....	1.8	2.5	4.4	7.2	13.8	19.2	36.7	63.7	122.7	353.8
White, both sexes										
2002.....	2.4	3.2	5.7	9.1	14.1	23.4	37.1	68.0	124.4	324.4
2001.....	1.9	3.5	5.6	8.9	14.7	23.7	38.6	66.9	123.8	329.7
2000.....	2.2	3.3	5.4	9.8	14.7	24.7	39.4	66.6	127.1	341.6
1999.....	2.3	3.6	5.9	9.8	17.9	26.0	43.0	70.0	133.1	355.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Conduction disorders and cardiac dysrhythmias (I44-I49)										
White, male										
2002.....	3.4	4.4	7.8	12.4	19.3	29.9	45.7	78.2	136.9	308.9
2001.....	2.7	4.7	7.7	11.6	19.4	29.7	46.9	78.8	137.5	310.2
2000.....	2.8	4.6	7.4	13.3	20.0	32.7	50.4	78.4	138.4	332.8
1999.....	3.0	5.3	8.0	13.3	23.6	35.1	53.3	81.3	152.3	350.9
White, female										
2002.....	1.5	2.0	3.6	6.1	9.4	17.7	30.0	60.6	116.8	331.1
2001.....	1.2	2.2	3.6	6.2	10.4	18.4	31.8	58.5	115.5	337.9
2000.....	1.6	2.0	3.3	6.4	9.9	17.6	30.5	58.4	120.4	345.2
1999.....	1.6	2.0	3.9	6.5	12.6	18.1	34.9	62.1	121.8	357.9
Black, both sexes										
2002.....	4.4	7.2	14.0	20.3	33.5	41.6	63.8	99.1	160.7	337.6
2001.....	4.0	8.0	13.6	21.1	29.8	44.7	65.7	105.2	162.5	340.2
2000.....	4.8	8.3	11.8	19.4	32.3	43.4	64.4	105.6	147.1	346.3
1999.....	4.2	8.9	12.8	21.7	33.6	44.6	70.6	109.4	170.2	359.0
Black, male										
2002.....	5.1	9.3	18.4	26.5	43.9	51.4	82.5	120.3	173.5	336.6
2001.....	4.4	9.4	17.6	28.5	40.5	50.6	86.2	135.7	198.2	357.4
2000.....	5.1	10.8	17.1	26.2	42.9	58.8	79.6	121.4	152.6	361.0
1999.....	4.7	11.1	16.5	31.0	43.5	61.0	85.8	135.3	215.4	411.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Conduction disorders and cardiac dysrhythmias (I44-I49)										
Black, female										
2002.....	3.8	5.4	10.2	15.3	25.4	34.4	51.3	86.4	154.0	338.0
2001.....	3.7	6.7	10.2	15.0	21.4	40.4	52.0	87.1	144.3	333.8
2000.....	4.6	6.2	7.3	13.8	24.0	32.1	54.4	96.0	144.4	340.8
1999.....	3.7	6.9	9.6	14.1	25.8	32.5	60.5	93.8	147.8	339.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
All races 3/, both sexes										
2002.....	0.8	1.7	3.4	6.0	12.4	22.9	45.7	89.0	194.0	645.0
2001.....	0.8	1.7	3.3	6.5	13.0	23.8	45.6	92.6	199.2	674.6
2000.....	0.8	1.4	3.0	6.1	12.8	24.0	46.5	93.5	199.9	684.9
1999.....	0.8	1.5	2.9	6.5	11.7	23.5	48.3	94.3	201.6	681.9
All races 3/, male										
2002.....	1.0	2.0	4.4	7.2	14.8	27.0	53.2	103.9	218.0	621.4
2001.....	0.9	2.2	4.0	8.1	15.0	28.0	54.2	107.9	226.6	646.4
2000.....	1.1	1.9	3.5	7.4	15.3	28.2	55.6	107.9	229.5	672.2
1999.....	1.0	1.9	3.6	8.0	14.2	28.0	58.9	108.4	224.9	667.0
All races 3/, female										
2002.....	0.5	1.3	2.4	4.9	10.1	19.3	39.8	78.4	179.3	655.3
2001.....	0.7	1.2	2.6	5.1	11.3	20.2	38.7	81.9	182.7	686.4
2000.....	0.6	0.9	2.5	5.0	10.5	20.3	39.3	83.5	182.4	690.0
1999.....	0.6	1.1	2.1	5.1	9.5	19.6	39.9	84.5	188.0	687.9
White, both sexes										
2002.....	0.6	1.2	2.7	5.3	11.4	21.7	44.1	88.1	195.6	664.5
2001.....	0.6	1.4	2.7	5.7	12.1	22.5	44.3	92.1	201.1	695.1
2000.....	0.6	1.1	2.5	5.4	11.8	22.6	44.9	92.8	201.8	703.7
1999.....	0.6	1.1	2.3	5.6	10.9	22.0	46.8	92.5	202.5	699.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
White, male										
2002.....	0.8	1.5	3.6	6.4	13.6	25.5	51.4	103.4	221.4	641.9
2001.....	0.7	1.9	3.4	7.2	14.2	26.5	52.8	107.6	228.2	666.5
2000.....	0.9	1.5	3.0	6.8	14.1	27.2	54.3	107.7	231.7	690.0
1999.....	0.8	1.4	2.9	7.1	13.3	26.2	58.1	107.2	226.3	680.0
White, female										
2002.....	0.4	0.9	1.9	4.2	9.3	18.4	38.2	77.1	179.8	674.3
2001.....	0.5	0.8	2.0	4.2	10.1	18.9	37.4	81.0	184.7	707.1
2000.....	0.4	0.6	2.0	4.1	9.6	18.6	37.3	82.4	183.9	709.3
1999.....	0.4	0.8	1.7	4.2	8.8	18.4	37.8	82.1	188.5	707.0
Black, both sexes										
2002.....	2.2	5.5	9.3	13.9	24.6	38.6	71.7	118.5	215.3	528.0
2001.....	2.7	4.5	8.7	15.7	24.6	41.2	71.1	121.1	213.7	543.8
2000.....	2.2	4.2	7.6	14.0	24.3	41.2	72.6	119.8	209.5	558.4
1999.....	2.4	4.8	8.4	16.4	22.0	42.0	74.0	130.5	216.7	552.9
Black, male										
2002.....	2.7	6.6	12.0	16.5	31.3	47.6	87.5	136.9	236.9	479.2
2001.....	2.6	5.0	10.4	18.9	26.4	49.6	86.2	140.8	256.8	541.2
2000.....	2.5	5.5	9.1	14.8	29.6	45.5	82.9	134.7	242.8	581.2
1999.....	2.7	6.4	11.0	19.3	27.4	51.4	83.1	146.7	238.9	603.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
Black, female										
2002.....	1.7	4.5	7.0	11.7	19.4	31.8	61.2	107.6	204.1	546.7
2001.....	2.8	4.1	7.2	13.1	23.2	35.1	61.0	109.3	191.7	544.8
2000.....	1.9	3.0	6.4	13.4	20.1	38.0	65.7	110.8	192.8	549.9
1999.....	2.1	3.5	6.3	14.0	17.8	35.0	68.0	120.7	205.7	534.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congestive heart failure (I50.0)										
All races 3/, both sexes										
2002.....	0.6	1.3	2.8	5.1	10.7	20.3	41.4	82.3	181.6	609.0
2001.....	0.7	1.4	2.8	5.4	11.3	21.5	41.5	85.8	185.4	636.1
2000.....	0.7	1.1	2.5	5.4	11.1	20.9	41.8	86.1	186.9	644.4
1999.....	0.7	1.2	2.3	5.3	10.0	20.9	43.6	86.6	188.4	642.7
All races 3/, male										
2002.....	0.8	1.6	3.5	6.1	12.5	23.8	48.2	95.6	203.2	584.6
2001.....	0.8	1.9	3.3	6.6	12.8	24.9	49.1	98.7	208.5	609.3
2000.....	0.8	1.6	2.9	6.4	13.0	24.7	49.6	99.1	214.1	632.2
1999.....	0.8	1.5	2.9	6.5	11.9	24.5	53.2	99.2	209.4	628.2
All races 3/, female										
2002.....	0.5	1.0	2.1	4.2	9.0	17.2	36.0	72.8	168.4	619.6
2001.....	0.6	1.0	2.3	4.2	10.0	18.5	35.5	76.8	171.5	647.3
2000.....	0.5	0.7	2.1	4.4	9.4	17.7	35.7	77.0	170.9	649.4
1999.....	0.5	0.9	1.8	4.2	8.4	17.9	36.1	77.9	176.1	648.5
White, both sexes										
2002.....	0.5	0.9	2.2	4.5	9.8	19.3	40.1	81.7	183.6	628.8
2001.....	0.5	1.1	2.3	4.7	10.5	20.2	40.5	85.7	187.6	656.5
2000.....	0.5	0.8	2.1	4.8	10.2	19.7	40.6	85.6	189.3	663.6
1999.....	0.5	0.8	1.8	4.5	9.4	19.6	42.3	85.3	189.4	660.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congestive heart failure (I50.0)										
White, male										
2002.....	0.6	1.2	2.8	5.4	11.4	22.5	46.5	95.3	207.2	605.8
2001.....	0.6	1.6	2.8	6.0	12.1	23.5	47.9	99.1	210.8	629.0
2000.....	0.7	1.2	2.4	6.1	11.8	23.9	48.6	99.2	216.9	650.5
1999.....	0.7	1.1	2.3	5.7	11.0	22.9	52.5	98.4	210.7	641.7
White, female										
2002.....	0.3	0.7	1.6	3.6	8.2	16.5	35.0	71.9	169.1	638.8
2001.....	0.4	0.7	1.9	3.5	9.1	17.2	34.4	76.1	173.6	667.9
2000.....	0.4	0.5	1.7	3.5	8.7	16.0	34.1	76.1	172.8	668.8
1999.....	0.4	0.6	1.3	3.4	7.8	16.8	34.2	76.2	176.9	667.4
Black, both sexes										
2002.....	1.9	4.6	7.7	11.9	21.9	34.4	63.8	106.7	196.0	482.6
2001.....	2.2	4.0	7.3	13.3	21.0	37.6	63.1	108.5	194.2	502.4
2000.....	1.8	3.6	6.7	11.9	21.3	36.5	63.0	108.4	190.3	510.2
1999.....	1.8	4.2	7.3	13.6	18.7	37.2	66.6	116.4	200.2	509.5
Black, male										
2002.....	2.2	5.6	9.6	14.3	27.3	42.9	81.5	122.6	209.4	428.8
2001.....	2.3	4.5	8.8	15.2	22.6	44.2	75.4	123.1	224.7	500.4
2000.....	2.0	4.7	8.0	11.7	26.1	38.2	71.2	120.0	216.6	528.2
1999.....	2.1	5.6	9.2	15.4	23.1	44.9	74.6	131.5	224.1	554.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congestive heart failure (I50.0)										
Black, female										
2002.....	1.6	3.7	6.1	9.9	17.7	28.1	52.0	97.2	189.0	503.2
2001.....	2.2	3.6	6.0	11.7	19.7	32.7	54.9	99.8	178.7	503.1
2000.....	1.5	2.5	5.5	12.0	17.5	35.1	57.5	101.4	177.0	503.5
1999.....	1.6	3.0	5.7	12.0	15.3	31.6	61.2	107.3	188.4	492.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified heart failure (I50.1-I50.9)										
All races 3/, both sexes										
2002.....	0.1	0.4	0.6	0.9	1.7	2.6	4.3	6.7	12.4	36.1
2001.....	0.2	0.3	0.5	1.1	1.7	2.4	4.1	6.8	13.8	38.5
2000.....	0.2	0.3	0.5	0.8	1.7	3.0	4.7	7.4	12.9	40.5
1999.....	0.2	0.3	0.5	1.2	1.7	2.6	4.6	7.7	13.3	39.2
All races 3/, male										
2002.....	0.2	0.4	1.0	1.1	2.3	3.2	5.0	8.3	14.8	36.8
2001.....	0.2	0.3	0.7	1.5	2.1	3.1	5.2	9.2	18.1	37.2
2000.....	0.3	0.4	0.7	0.9	2.3	3.6	6.1	8.8	15.4	40.0
1999.....	0.2	0.4	0.7	1.6	2.4	3.5	5.7	9.2	15.5	38.8
All races 3/, female										
2002.....	*	0.3	0.3	0.7	1.1	2.1	3.8	5.6	10.9	35.7
2001.....	*	*	0.3	0.8	1.3	1.7	3.2	5.2	11.2	39.1
2000.....	*	*	0.4	0.6	1.1	2.5	3.6	6.5	11.5	40.6
1999.....	*	0.3	0.4	0.9	1.1	1.7	3.8	6.6	11.9	39.4
White, both sexes										
2002.....	0.1	0.3	0.5	0.8	1.6	2.4	4.0	6.4	12.0	35.6
2001.....	0.1	0.2	0.4	1.0	1.5	2.3	3.8	6.4	13.5	38.7
2000.....	0.1	0.2	0.5	0.7	1.6	2.9	4.3	7.2	12.5	40.2
1999.....	0.1	0.3	0.5	1.1	1.6	2.4	4.5	7.2	13.1	39.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified heart failure (I50.1-I50.9)										
White, male										
2002.....	*	0.3	0.8	1.0	2.2	3.0	4.9	8.0	14.2	36.0
2001.....	*	0.3	0.6	1.3	2.0	3.0	4.9	8.5	17.4	37.5
2000.....	0.2	0.3	0.6	0.7	2.2	3.2	5.7	8.4	14.8	39.5
1999.....	*	0.4	0.6	1.4	2.2	3.3	5.6	8.8	15.6	38.3
White, female										
2002.....	*	0.2	*	0.6	1.1	1.9	3.3	5.2	10.6	35.5
2001.....	*	*	*	0.8	1.1	1.7	3.0	4.9	11.1	39.2
2000.....	*	*	0.3	0.6	1.0	2.6	3.2	6.3	11.1	40.5
1999.....	*	*	0.3	0.8	1.0	1.6	3.6	6.0	11.6	39.5
Black, both sexes										
2002.....	*	0.9	1.6	2.0	2.7	4.1	7.9	11.9	19.4	45.4
2001.....	*	*	1.4	2.5	3.7	3.7	7.9	12.5	19.5	41.5
2000.....	*	*	*	2.2	3.0	4.8	9.6	11.4	19.2	48.2
1999.....	*	*	*	2.8	3.3	4.8	7.4	14.1	16.5	43.4
Black, male										
2002.....	*	*	2.5	*	4.0	*	*	14.2	27.5	50.4
2001.....	*	*	*	3.8	*	5.3	10.8	17.7	32.1	40.8
2000.....	*	*	*	*	*	7.3	11.7	14.6	26.1	53.0
1999.....	*	*	*	3.9	4.2	6.5	8.5	15.2	*	49.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified heart failure (I50.1-I50.9)

Black, female

2002.....	*	*	*	*	*	3.7	9.2	10.4	15.1	43.5
2001.....	*	*	*	*	3.5	*	6.1	9.4	13.1	41.7
2000.....	*	*	*	*	*	*	8.2	9.4	15.7	46.4
1999.....	*	*	*	*	*	*	6.8	13.4	17.3	41.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)										
All races 3/, both sexes										
2002.....	0.3	0.3	0.3	0.4	0.2	0.4	0.6	0.8	1.1	3.2
2001.....	0.3	0.3	0.3	0.2	0.3	0.4	0.4	0.5	0.6	2.0
2000.....	0.3	0.3	0.1	0.3	0.3	*	0.4	0.4	0.7	2.4
1999.....	0.3	0.3	0.3	0.2	0.2	0.3	0.3	0.4	0.7	2.4
All races 3/, male										
2002.....	0.4	0.3	0.3	0.3	*	*	0.8	0.8	1.0	3.8
2001.....	0.3	0.3	0.3	*	*	0.5	0.5	0.8	*	2.2
2000.....	0.3	0.2	*	0.4	*	*	*	*	*	2.6
1999.....	0.3	0.3	0.3	*	*	*	*	*	*	2.1
All races 3/, female										
2002.....	0.3	0.3	0.3	0.4	*	*	0.5	0.8	1.1	3.0
2001.....	0.3	0.3	*	*	*	*	*	*	*	1.9
2000.....	0.2	0.3	*	0.3	*	*	*	*	*	2.4
1999.....	0.3	0.3	0.2	*	*	*	*	*	*	2.5
White, both sexes										
2002.....	0.3	0.3	0.2	0.4	0.2	0.3	0.6	0.8	1.1	3.2
2001.....	0.2	0.2	0.2	0.2	0.3	0.3	0.4	0.5	0.6	2.1
2000.....	0.2	0.2	*	0.4	0.2	*	0.4	0.4	0.7	2.6
1999.....	0.3	0.2	0.2	0.2	*	0.3	0.3	0.3	0.8	2.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)										
All races 3/, both sexes										
2002.....	1.3	1.7	2.1	2.8	4.2	6.1	9.7	15.3	26.4	65.0
2001.....	1.2	1.7	2.0	2.8	4.1	5.7	9.0	14.4	25.4	66.3
2000.....	1.1	1.6	2.0	3.1	4.5	6.4	8.7	14.3	26.8	64.6
1999.....	1.0	1.4	1.8	2.8	4.3	5.6	8.3	14.1	23.4	61.5
All races 3/, male										
2002.....	1.8	2.4	3.0	4.0	5.5	8.2	13.5	18.9	30.7	66.7
2001.....	1.7	2.4	2.8	3.9	5.8	7.8	12.3	17.8	28.8	66.4
2000.....	1.7	2.3	2.8	4.4	6.5	8.1	11.3	17.0	31.2	69.0
1999.....	1.5	2.1	2.5	4.0	6.2	7.8	11.2	18.3	27.7	64.3
All races 3/, female										
2002.....	0.8	1.1	1.3	1.6	3.0	4.3	6.6	12.8	23.8	64.3
2001.....	0.7	0.9	1.2	1.9	2.6	3.9	6.5	12.0	23.3	66.3
2000.....	0.6	0.8	1.2	1.9	2.7	4.9	6.6	12.5	24.1	62.7
1999.....	0.6	0.8	1.1	1.7	2.7	3.7	6.0	11.2	20.8	60.4
White, both sexes										
2002.....	1.1	1.5	1.8	2.6	3.7	5.7	9.4	15.2	26.3	64.5
2001.....	1.0	1.4	1.7	2.5	4.0	5.4	8.6	13.7	25.1	65.7
2000.....	1.0	1.5	1.7	2.8	4.3	6.1	7.9	13.9	26.1	64.7
1999.....	0.9	1.2	1.6	2.4	3.9	5.1	8.1	13.3	23.2	61.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)										
White, male										
2002.....	1.6	2.1	2.6	3.7	4.9	7.6	13.3	18.7	30.2	66.9
2001.....	1.6	2.1	2.6	3.5	5.7	7.5	11.6	17.0	28.4	66.2
2000.....	1.5	2.2	2.4	3.9	6.3	7.7	10.3	16.4	30.0	69.2
1999.....	1.2	2.0	2.2	3.5	5.7	7.2	10.8	17.5	27.2	64.5
White, female										
2002.....	0.7	0.8	1.0	1.5	2.7	4.0	6.3	12.8	23.9	63.4
2001.....	0.5	0.7	0.9	1.6	2.5	3.5	6.2	11.3	23.1	65.5
2000.....	0.5	0.7	0.9	1.7	2.4	4.6	6.0	12.1	23.8	62.8
1999.....	0.5	0.5	0.9	1.4	2.2	3.2	5.9	10.3	20.8	60.2
Black, both sexes										
2002.....	2.3	3.6	5.4	5.1	8.7	11.2	14.5	19.9	32.2	80.5
2001.....	2.6	3.7	4.5	6.1	6.4	9.4	15.2	23.8	32.2	83.8
2000.....	2.4	2.9	4.8	6.5	8.0	11.0	18.7	22.2	37.8	73.8
1999.....	2.6	3.1	3.8	6.7	9.1	11.3	13.3	24.8	26.9	68.9
Black, male										
2002.....	3.2	4.6	7.2	7.3	12.1	16.0	18.7	26.6	42.8	77.2
2001.....	3.0	5.1	5.3	8.4	8.3	10.9	21.9	28.8	39.3	84.0
2000.....	3.3	4.1	7.1	9.7	10.6	14.1	25.4	26.9	52.3	80.7
1999.....	3.6	3.8	5.3	9.7	11.7	16.1	19.0	29.9	36.6	72.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

Black, female

2002.....	1.5	2.6	4.0	3.3	6.0	7.6	11.6	15.9	26.7	81.7
2001.....	2.3	2.5	3.9	4.2	4.8	8.3	10.8	20.8	28.6	83.8
2000.....	1.7	1.8	2.8	3.9	6.0	8.8	14.2	19.4	30.6	71.3
1999.....	1.6	2.6	2.6	4.3	7.2	7.7	9.5	21.7	22.0	67.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)										
All races 3/, both sexes										
2002.....	7.1	12.3	18.3	28.6	48.3	83.8	160.7	308.8	601.7	1,445.9
2001.....	7.2	12.1	18.5	28.8	49.6	85.7	164.3	320.2	622.4	1,500.2
2000.....	7.7	13.1	19.3	30.8	53.8	91.0	169.1	332.8	654.0	1,589.2
1999.....	7.5	12.5	18.4	31.7	51.4	91.1	173.5	340.6	664.0	1,614.8
All races 3/, male										
2002.....	6.9	13.4	20.4	32.7	55.8	97.0	178.7	334.8	615.6	1,317.9
2001.....	7.4	13.2	20.5	32.8	57.2	98.5	187.9	349.0	654.6	1,380.2
2000.....	7.8	14.2	21.3	35.2	62.3	102.6	192.9	367.6	695.1	1,484.3
1999.....	7.8	13.3	20.7	35.7	60.0	102.8	197.7	376.9	706.8	1,512.8
All races 3/, female										
2002.....	7.2	11.2	16.3	24.8	41.4	72.4	146.2	290.4	593.3	1,501.5
2001.....	7.1	10.9	16.5	25.1	42.7	74.7	145.6	300.0	603.2	1,550.4
2000.....	7.5	12.0	17.3	26.6	46.0	81.0	150.3	308.5	629.7	1,632.0
1999.....	7.1	11.7	16.1	28.0	43.7	81.1	154.5	315.3	639.0	1,656.0
White, both sexes										
2002.....	5.5	9.5	14.0	23.8	41.2	74.2	150.1	298.6	595.7	1,459.5
2001.....	5.6	9.3	13.8	23.9	41.7	75.2	153.2	309.7	617.1	1,511.2
2000.....	5.9	9.7	15.4	25.4	46.5	81.9	159.0	321.3	647.0	1,601.6
1999.....	5.7	9.4	14.4	26.2	43.5	82.9	161.3	329.3	658.4	1,631.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)										
White, male										
2002.....	5.5	10.6	15.5	26.9	47.2	85.1	167.3	323.4	610.1	1,327.4
2001.....	5.9	10.2	15.4	26.9	48.1	86.7	174.8	337.7	648.8	1,386.2
2000.....	6.1	10.8	16.9	28.6	53.8	91.7	180.5	354.6	686.1	1,490.7
1999.....	6.1	9.8	16.2	29.5	50.6	93.2	183.6	363.2	700.5	1,522.7
White, female										
2002.....	5.5	8.4	12.6	20.9	35.6	64.5	136.0	280.8	586.8	1,516.9
2001.....	5.4	8.3	12.2	21.1	35.8	65.1	135.7	289.8	598.0	1,563.5
2000.....	5.7	8.7	13.9	22.4	39.8	73.3	141.9	297.9	623.8	1,646.7
1999.....	5.3	8.9	12.7	23.0	37.0	73.9	143.7	305.5	633.6	1,675.4
Black, both sexes										
2002.....	17.4	32.1	51.2	71.1	109.9	169.6	272.5	447.2	712.5	1,383.9
2001.....	18.4	32.6	55.0	70.6	117.1	178.3	277.6	460.2	716.4	1,452.7
2000.....	19.7	37.2	51.2	77.4	114.0	174.8	281.1	475.2	769.4	1,530.0
1999.....	19.9	34.4	50.6	78.4	118.2	170.0	307.8	487.9	756.4	1,492.5
Black, male										
2002.....	16.9	34.3	61.6	88.9	138.8	213.7	328.8	502.5	729.8	1,252.2
2001.....	17.8	36.5	63.9	87.6	143.4	218.5	336.3	524.6	768.1	1,349.1
2000.....	20.5	41.0	60.4	97.8	137.5	212.8	340.0	544.8	863.2	1,458.8
1999.....	20.8	38.8	61.3	92.7	149.4	204.3	366.7	559.0	818.8	1,448.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)										
Black, female										
2002.....	17.9	30.1	42.4	56.5	87.3	136.9	234.8	414.1	703.4	1,434.3
2001.....	19.0	29.1	47.4	56.7	96.5	148.5	238.6	421.9	690.1	1,491.2
2000.....	19.0	33.9	43.4	60.6	95.4	146.7	242.0	433.2	722.2	1,556.5
1999.....	19.0	30.5	41.6	66.6	93.7	144.5	268.7	445.1	725.6	1,509.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Subarachnoid hemorrhage (I60)										
All races 3/, both sexes										
2002.....	2.3	3.0	3.1	3.8	4.8	5.5	6.2	7.9	10.3	12.2
2001.....	2.2	3.0	3.4	4.2	4.3	5.3	6.3	8.0	9.8	13.2
2000.....	2.5	3.3	3.8	4.7	5.0	5.2	6.4	8.6	10.7	14.1
1999.....	2.6	3.3	3.7	4.4	5.1	5.7	6.7	8.4	10.9	13.1
All races 3/, male										
2002.....	1.8	2.4	2.2	3.0	3.8	4.2	4.6	5.6	6.1	9.1
2001.....	1.9	2.4	2.8	3.2	3.1	3.5	5.2	5.7	7.5	9.5
2000.....	2.0	2.6	2.9	3.4	3.6	3.4	5.2	6.3	7.7	11.2
1999.....	2.2	2.7	2.9	3.3	3.6	4.2	4.4	6.1	7.7	9.0
All races 3/, female										
2002.....	2.7	3.6	4.0	4.6	5.6	6.6	7.4	9.6	12.9	13.6
2001.....	2.6	3.6	3.9	5.2	5.3	6.8	7.2	9.7	11.2	14.8
2000.....	2.9	4.0	4.7	6.0	6.4	6.7	7.4	10.3	12.4	15.3
1999.....	3.0	3.9	4.4	5.5	6.4	6.9	8.5	10.0	12.8	14.8
White, both sexes										
2002.....	2.0	2.8	2.8	3.5	4.6	5.3	5.7	7.7	10.1	12.0
2001.....	1.9	2.7	2.9	3.8	4.2	4.8	6.1	7.7	9.9	12.7
2000.....	2.1	2.9	3.5	4.6	4.8	4.8	6.0	8.4	10.4	13.6
1999.....	2.3	2.9	3.4	4.2	4.7	5.6	6.5	8.3	10.7	12.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Subarachnoid hemorrhage (I60)										
White, male										
2002.....	1.7	2.4	2.0	2.7	3.6	4.1	4.3	5.6	5.9	9.1
2001.....	1.7	2.1	2.5	2.8	3.1	3.3	5.0	5.5	7.9	9.2
2000.....	1.8	2.4	2.7	3.3	3.3	3.2	4.8	6.3	7.8	11.8
1999.....	2.1	2.5	2.7	3.2	3.4	4.0	4.4	6.3	7.6	9.0
White, female										
2002.....	2.4	3.1	3.6	4.3	5.5	6.4	6.8	9.3	12.7	13.3
2001.....	2.2	3.2	3.4	4.8	5.3	6.1	6.9	9.3	11.1	14.2
2000.....	2.5	3.4	4.3	5.9	6.2	6.1	7.0	10.0	12.0	14.3
1999.....	2.5	3.3	4.0	5.1	5.9	7.0	8.3	9.7	12.5	14.1
Black, both sexes										
2002.....	4.1	5.3	5.5	6.0	5.6	6.2	6.7	8.2	9.2	12.8
2001.....	4.4	5.1	6.5	7.2	4.8	8.2	6.3	9.7	6.8	13.8
2000.....	4.7	6.1	6.2	6.2	6.2	7.4	8.7	11.0	10.4	17.2
1999.....	5.0	6.0	6.3	6.7	6.7	5.5	8.7	7.5	12.4	14.7
Black, male										
2002.....	2.7	3.1	3.4	5.2	5.1	5.7	6.7	*	*	*
2001.....	3.4	3.9	5.3	5.8	4.1	*	*	*	*	*
2000.....	3.6	4.2	4.6	3.8	4.8	*	8.4	*	*	*
1999.....	3.5	4.5	5.3	5.1	4.9	5.7	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Subarachnoid hemorrhage (I60)										
Black, female										
2002.....	5.2	7.2	7.3	6.6	6.0	6.5	6.7	9.9	10.7	15.6
2001.....	5.3	6.1	7.6	8.3	5.4	10.9	7.2	10.8	8.6	16.4
2000.....	5.6	7.8	7.5	8.1	7.4	9.4	8.9	12.8	13.2	22.8
1999.....	6.4	7.3	7.2	8.0	8.2	5.4	10.8	10.3	13.0	18.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intracerebral and other intracranial hemorrhage (I61-I62)										
All races 3/, both sexes										
2002.....	2.8	5.0	7.2	9.7	14.3	20.9	36.0	59.1	90.6	127.2
2001.....	3.1	5.2	7.4	9.5	14.8	21.2	35.2	58.2	91.1	127.9
2000.....	3.2	5.6	7.6	10.6	15.9	23.5	35.5	60.1	95.1	137.9
1999.....	2.8	4.9	7.1	10.7	14.3	20.7	36.4	59.5	90.5	122.3
All races 3/, male										
2002.....	3.2	6.0	8.8	11.0	17.3	24.7	41.0	63.4	98.4	135.0
2001.....	3.4	6.3	8.8	11.2	17.3	23.7	41.5	65.5	99.5	134.2
2000.....	3.6	6.8	8.9	13.4	19.0	27.0	40.0	65.9	100.6	149.6
1999.....	3.4	5.6	8.6	12.7	17.3	23.5	41.7	65.1	94.2	130.7
All races 3/, female										
2002.....	2.4	3.9	5.7	8.4	11.7	17.6	32.1	56.1	85.9	123.8
2001.....	2.7	4.1	6.2	7.8	12.6	19.1	30.2	53.1	86.2	125.2
2000.....	2.8	4.4	6.4	7.9	13.0	20.4	32.0	56.1	91.8	133.1
1999.....	2.3	4.2	5.6	8.8	11.6	18.2	32.3	55.6	88.4	118.9
White, both sexes										
2002.....	1.8	3.4	5.0	7.8	12.5	18.8	34.9	58.7	91.3	129.2
2001.....	2.2	3.5	5.1	7.8	12.5	19.1	33.7	57.8	91.7	128.9
2000.....	2.2	3.7	5.6	8.5	13.7	21.7	34.7	59.4	95.6	139.4
1999.....	1.9	3.3	5.1	8.5	12.2	19.4	35.4	58.9	90.7	123.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intracerebral and other intracranial hemorrhage (I61-I62)										
White, male										
2002.....	2.1	4.2	6.1	8.7	15.1	22.2	40.0	63.5	99.5	136.9
2001.....	2.5	4.5	6.0	9.2	14.9	21.2	39.7	65.1	100.4	134.9
2000.....	2.5	4.7	6.6	10.7	16.1	24.5	38.9	65.5	101.3	150.5
1999.....	2.3	3.7	6.2	10.1	14.8	22.2	40.7	64.0	93.7	131.7
White, female										
2002.....	1.5	2.6	4.0	7.0	10.1	15.8	30.7	55.2	86.2	125.9
2001.....	1.9	2.6	4.1	6.5	10.2	17.3	28.8	52.7	86.4	126.3
2000.....	1.8	2.7	4.6	6.4	11.4	19.2	31.3	55.1	92.2	134.9
1999.....	1.5	2.8	4.1	7.1	9.7	16.9	31.2	55.3	89.0	119.7
Black, both sexes										
2002.....	8.8	15.7	22.9	24.1	28.1	34.8	44.6	61.7	81.1	98.3
2001.....	8.7	16.6	25.5	22.6	32.0	37.0	43.7	59.2	83.2	106.6
2000.....	10.0	19.0	23.4	27.5	32.3	36.1	42.4	63.3	78.2	117.0
1999.....	9.4	16.1	22.3	26.6	29.6	30.5	44.4	62.5	77.4	96.9
Black, male										
2002.....	10.4	19.3	30.0	30.8	36.0	43.4	52.4	61.5	81.0	101.8
2001.....	9.3	20.3	31.3	29.1	36.4	46.0	54.2	62.7	85.9	112.3
2000.....	11.6	23.4	28.5	38.4	43.5	46.3	47.1	69.9	78.4	130.3
1999.....	11.6	19.2	28.6	33.2	37.8	34.0	48.2	70.8	81.1	109.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intracerebral and other intracranial hemorrhage (I61-I62)										
Black, female										
2002.....	7.4	12.5	16.9	18.6	21.8	28.5	39.4	61.7	81.2	96.9
2001.....	8.1	13.4	20.6	17.2	28.5	30.4	36.8	57.1	81.8	104.4
2000.....	8.5	15.2	19.1	18.6	23.5	28.6	39.3	59.2	78.1	112.0
1999.....	7.4	13.3	16.9	21.2	23.1	27.9	41.9	57.5	75.6	92.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebral infarction (I63)										
All races 3/, both sexes										
2002.....	0.3	0.7	1.3	1.7	3.5	5.8	11.6	22.6	45.6	118.3
2001.....	0.4	0.8	1.1	1.9	3.1	6.1	13.0	24.3	47.6	124.8
2000.....	0.5	0.7	1.2	2.0	4.0	6.6	12.8	27.2	54.5	141.0
1999.....	0.5	0.7	1.2	2.4	4.0	8.1	14.9	29.7	58.5	153.1
All races 3/, male										
2002.....	0.3	0.9	1.6	2.3	4.2	7.2	12.4	25.8	50.1	108.7
2001.....	0.4	0.9	1.2	2.6	3.8	7.7	14.9	28.0	53.3	111.2
2000.....	0.5	0.9	1.4	2.5	5.2	7.8	15.4	31.2	60.8	133.3
1999.....	0.5	0.9	1.5	2.9	4.8	10.0	17.4	34.0	63.3	144.6
All races 3/, female										
2002.....	0.3	0.5	0.9	1.1	2.9	4.6	11.0	20.4	42.8	122.4
2001.....	0.3	0.7	0.9	1.3	2.5	4.6	11.4	21.8	44.2	130.4
2000.....	0.4	0.5	1.1	1.6	3.0	5.6	10.7	24.4	50.7	144.2
1999.....	0.4	0.5	0.9	1.8	3.3	6.4	12.9	26.7	55.7	156.6
White, both sexes										
2002.....	0.3	0.6	1.0	1.5	3.2	5.3	10.9	22.1	44.8	119.1
2001.....	0.3	0.6	0.9	1.6	2.7	5.4	12.3	23.6	47.5	126.0
2000.....	0.3	0.6	1.0	1.6	3.7	6.0	12.0	26.3	54.0	141.7
1999.....	0.4	0.6	1.0	2.0	3.5	7.5	13.8	28.5	58.6	154.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebral infarction (I63)										
White, male										
2002.....	0.3	0.9	1.3	2.0	3.7	6.5	11.7	25.3	49.3	108.6
2001.....	0.4	0.7	1.1	2.2	3.4	7.1	14.3	26.8	52.9	110.5
2000.....	0.4	0.8	1.0	2.0	4.7	7.0	14.6	29.8	59.8	133.1
1999.....	0.4	0.7	1.3	2.6	4.2	9.5	15.9	32.3	63.5	144.0
White, female										
2002.....	0.3	0.4	0.8	1.0	2.7	4.2	10.3	19.8	42.1	123.7
2001.....	0.2	0.5	0.8	1.1	2.1	4.0	10.7	21.3	44.2	132.5
2000.....	0.3	0.4	0.9	1.2	2.7	5.2	10.0	23.8	50.5	145.2
1999.....	0.3	0.5	0.8	1.5	2.9	5.7	12.1	25.9	55.7	158.5
Black, both sexes										
2002.....	*	1.3	3.1	3.6	6.7	11.0	20.0	32.3	56.8	113.1
2001.....	0.9	2.0	2.2	4.4	7.3	12.3	21.7	34.9	53.4	121.3
2000.....	1.2	1.3	3.1	5.4	7.8	13.0	21.1	37.7	67.5	142.6
1999.....	1.1	1.8	2.7	5.0	8.7	14.4	27.7	47.0	62.7	150.8
Black, male										
2002.....	*	*	4.4	4.9	8.9	14.7	22.7	35.8	63.4	114.7
2001.....	*	2.4	2.6	6.6	8.3	15.8	23.2	47.4	65.0	127.1
2000.....	*	*	4.5	7.1	10.1	17.7	24.7	46.8	83.5	137.2
1999.....	*	2.7	3.1	6.3	10.8	17.1	37.7	60.3	68.9	162.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebral infarction (I63)										
Black, female										
2002.....	*	*	2.0	2.4	5.0	8.3	18.1	30.2	53.3	112.5
2001.....	*	1.7	1.9	2.7	6.4	9.8	20.6	27.5	47.4	119.1
2000.....	1.3	*	*	4.0	5.9	9.6	18.6	32.2	59.4	144.7
1999.....	*	*	2.4	3.9	7.0	12.3	21.2	38.9	59.6	146.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Stroke, not specified as hemorrhage or infarction (I64)										
All races 3/, both sexes										
2002.....	1.2	2.8	5.3	10.7	20.5	40.3	82.7	165.7	338.0	859.4
2001.....	1.2	2.5	5.2	10.5	22.0	41.7	85.7	176.0	355.4	906.2
2000.....	1.2	2.6	5.2	10.7	23.0	44.6	89.8	181.1	372.2	954.7
1999.....	1.1	2.7	5.0	11.3	22.4	45.1	89.6	186.1	380.9	982.4
All races 3/, male										
2002.....	1.2	3.2	6.1	13.2	24.3	47.5	93.1	179.9	334.5	762.8
2001.....	1.3	3.0	6.2	12.6	26.5	49.4	97.9	188.5	363.9	819.7
2000.....	1.3	3.1	6.5	12.8	27.8	51.0	102.3	198.1	390.9	863.2
1999.....	1.3	3.2	6.2	13.1	27.2	51.7	103.2	206.0	404.7	896.4
All races 3/, female										
2002.....	1.2	2.4	4.4	8.4	16.9	34.1	74.3	155.8	340.2	901.3
2001.....	1.1	2.0	4.2	8.6	17.8	35.1	76.1	167.2	350.3	942.4
2000.....	1.1	2.2	3.9	8.6	18.7	39.2	80.0	169.2	361.2	992.0
1999.....	0.9	2.2	3.9	9.6	18.2	39.4	78.9	172.4	367.0	1,017.1
White, both sexes										
2002.....	1.0	2.1	4.0	8.6	16.6	34.9	76.1	159.4	333.5	867.3
2001.....	0.9	1.9	3.9	8.5	17.8	35.8	78.8	169.6	350.8	913.6
2000.....	1.0	1.9	4.1	8.4	19.4	39.5	83.1	174.3	366.9	963.5
1999.....	0.8	1.9	3.8	9.0	18.4	39.9	82.3	178.9	377.1	993.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Stroke, not specified as hemorrhage or infarction (I64)										
White, male										
2002.....	1.1	2.5	4.7	10.6	19.6	40.6	85.5	172.6	329.7	766.7
2001.....	1.1	2.3	4.7	10.0	21.2	42.3	89.6	181.7	357.5	823.5
2000.....	1.1	2.2	5.3	10.1	23.6	45.2	94.2	189.8	384.8	865.2
1999.....	1.0	2.3	4.8	10.7	22.3	45.4	95.3	197.4	401.4	903.1
White, female										
2002.....	0.9	1.6	3.3	6.7	13.8	30.0	68.4	149.9	335.9	910.9
2001.....	0.7	1.4	3.1	7.0	14.6	30.1	70.1	161.0	346.7	951.3
2000.....	0.8	1.5	2.9	6.8	15.5	34.6	74.2	163.4	356.3	1,003.5
1999.....	0.6	1.5	2.9	7.4	14.9	35.1	72.0	165.9	362.8	1,029.7
Black, both sexes										
2002.....	3.2	8.0	15.7	30.9	55.6	91.8	156.7	259.7	425.2	840.7
2001.....	3.6	7.1	16.3	29.3	59.5	96.7	163.1	268.9	435.0	883.4
2000.....	2.9	8.6	14.7	31.0	55.2	97.3	167.8	275.6	473.0	919.4
1999.....	3.4	8.5	16.0	32.5	59.9	97.2	174.4	287.1	454.8	917.6
Black, male										
2002.....	2.5	8.9	18.8	40.0	71.1	118.4	194.6	293.9	428.7	761.2
2001.....	3.6	8.2	19.8	37.9	77.9	122.7	199.6	302.9	470.3	816.9
2000.....	3.0	10.0	18.6	39.0	66.7	117.1	203.6	321.8	519.3	897.2
1999.....	3.8	10.1	19.8	37.8	77.2	120.2	202.2	328.6	489.2	877.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Stroke, not specified as hemorrhage or infarction (I64)										
Black, female										
2002.....	3.8	7.2	13.0	23.5	43.5	72.0	131.3	239.3	423.3	871.2
2001.....	3.7	6.1	13.3	22.2	45.0	77.5	138.8	248.5	417.0	908.1
2000.....	2.8	7.4	11.4	24.4	46.1	82.8	144.1	247.8	449.6	927.6
1999.....	3.1	7.2	12.7	28.1	46.3	80.2	156.0	262.0	437.8	932.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other cerebrovascular diseases and their sequelae (I67,I69)										
All races 3/, both sexes										
2002.....	0.5	0.8	1.5	2.7	5.2	11.3	24.2	53.4	117.2	328.9
2001.....	0.4	0.7	1.4	2.7	5.4	11.4	24.1	53.6	118.5	328.2
2000.....	0.4	0.8	1.4	2.8	5.8	11.0	24.6	55.7	121.5	341.4
1999.....	0.4	0.8	1.4	2.9	5.6	11.7	25.8	56.9	123.1	343.9
All races 3/, male										
2002.....	0.4	0.8	1.7	3.2	6.2	13.3	27.6	60.2	126.5	302.3
2001.....	0.4	0.7	1.5	3.2	6.5	14.2	28.3	61.4	130.6	305.5
2000.....	0.4	0.8	1.6	3.1	6.8	13.3	30.1	66.1	135.0	326.9
1999.....	0.4	0.8	1.5	3.6	7.1	13.4	31.0	65.8	136.9	332.1
All races 3/, female										
2002.....	0.6	0.8	1.2	2.4	4.3	9.5	21.5	48.6	111.5	340.4
2001.....	0.4	0.7	1.2	2.3	4.4	9.0	20.7	48.2	111.3	337.6
2000.....	0.4	0.9	1.3	2.4	5.0	9.0	20.3	48.5	113.6	347.4
1999.....	0.5	0.9	1.2	2.3	4.2	10.2	21.8	50.7	115.1	348.7
White, both sexes										
2002.....	0.4	0.6	1.2	2.4	4.3	9.8	22.4	50.7	115.9	332.0
2001.....	0.3	0.6	1.0	2.2	4.5	10.0	22.3	51.0	117.3	329.9
2000.....	0.3	0.7	1.2	2.3	5.0	9.9	23.1	52.9	120.2	343.4
1999.....	0.4	0.7	1.1	2.4	4.7	10.5	23.3	54.7	121.3	348.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other cerebrovascular diseases and their sequelae (I67,I69)										
White, male										
2002.....	0.3	0.7	1.4	2.7	5.1	11.7	25.8	56.5	125.7	306.3
2001.....	0.3	0.6	1.0	2.7	5.5	12.8	26.2	58.6	130.0	308.1
2000.....	0.3	0.7	1.3	2.5	6.2	11.8	27.9	63.3	132.5	330.0
1999.....	0.3	0.6	1.1	2.9	5.9	12.1	27.4	63.2	134.3	334.9
White, female										
2002.....	0.5	0.6	1.0	2.0	3.5	8.1	19.7	46.6	109.9	343.2
2001.....	0.3	0.6	0.9	1.8	3.6	7.6	19.2	45.6	109.6	339.1
2000.....	0.3	0.6	1.0	2.1	4.0	8.1	19.3	45.7	112.9	348.8
1999.....	0.4	0.7	1.1	1.9	3.6	9.2	20.2	48.7	113.6	353.4
Black, both sexes										
2002.....	0.8	1.8	4.0	6.5	13.9	25.8	44.5	85.2	140.2	318.9
2001.....	0.9	1.7	4.4	7.2	13.6	24.0	42.8	87.5	138.1	327.6
2000.....	0.9	2.2	3.8	7.3	12.5	20.9	41.1	87.6	140.3	333.8
1999.....	0.9	2.0	3.3	7.6	13.3	22.3	52.6	83.9	149.1	312.5
Black, male										
2002.....	*	*	5.0	7.9	17.6	31.4	52.4	105.6	150.5	269.1
2001.....	*	*	4.9	8.3	16.7	29.5	54.2	103.6	143.7	285.9
2000.....	*	*	4.2	9.6	12.4	27.1	56.2	98.3	177.0	291.8
1999.....	*	2.3	4.6	10.3	18.7	27.3	73.3	96.4	168.3	294.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other cerebrovascular diseases and their sequelae (I67,I69)

Black, female

2002.....	*	1.9	3.2	5.4	11.0	21.6	39.2	73.0	134.9	338.0
2001.....	*	1.8	4.1	6.3	11.2	19.9	35.2	77.9	135.3	343.1
2000.....	*	2.6	3.5	5.4	12.6	16.3	31.1	81.2	121.9	349.4
1999.....	*	1.7	2.3	5.4	9.2	18.7	38.9	76.4	139.6	319.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of arteries, arterioles and capillaries (I70-I78)										
All races 3/, both sexes										
2002.....	1.4	2.5	3.8	7.6	14.1	27.5	45.8	78.3	131.6	316.5
2001.....	1.4	2.3	3.9	7.4	14.6	27.3	46.5	82.5	137.2	337.3
2000.....	1.4	2.5	4.1	7.7	15.3	29.6	51.1	85.6	144.6	352.0
1999.....	1.5	2.4	4.0	8.0	15.4	28.6	51.3	87.6	147.8	367.2
All races 3/, male										
2002.....	2.0	3.6	5.2	10.3	19.6	38.1	60.7	102.7	166.1	344.0
2001.....	2.0	3.3	5.2	10.0	20.0	37.0	63.3	106.0	175.7	364.5
2000.....	1.9	3.3	5.8	11.0	21.6	41.2	70.6	116.1	186.4	391.4
1999.....	1.8	3.4	5.5	11.3	22.0	39.8	70.0	117.2	189.4	396.5
All races 3/, female										
2002.....	0.8	1.5	2.5	5.0	9.2	18.3	33.8	61.1	110.6	304.6
2001.....	0.9	1.4	2.7	5.0	9.7	18.9	33.1	66.0	114.2	325.9
2000.....	1.0	1.7	2.5	4.6	9.6	19.7	35.6	64.4	120.0	336.0
1999.....	1.2	1.5	2.7	5.0	9.5	19.1	36.7	67.0	123.5	355.4
White, both sexes										
2002.....	1.2	2.1	3.3	7.1	13.7	27.7	46.1	79.0	133.2	323.7
2001.....	1.2	2.1	3.5	7.0	14.0	27.5	46.6	82.8	138.7	344.1
2000.....	1.2	2.1	3.5	7.2	14.8	29.7	51.8	86.1	146.1	358.3
1999.....	1.3	2.1	3.7	7.6	15.4	28.7	52.1	88.3	149.4	372.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of arteries, arterioles and capillaries (I70-I78)										
White, male										
2002.....	1.8	3.1	4.4	9.6	19.0	39.0	61.2	104.5	169.2	354.1
2001.....	1.6	2.9	4.6	9.6	19.3	37.6	63.8	107.1	178.0	375.1
2000.....	1.5	2.9	5.0	10.5	21.3	42.0	72.3	117.6	190.0	401.0
1999.....	1.6	3.1	5.0	10.8	22.3	40.5	71.7	118.9	191.8	404.8
White, female										
2002.....	0.7	1.2	2.2	4.7	8.8	17.7	33.9	60.8	111.2	310.4
2001.....	0.8	1.3	2.4	4.5	9.0	18.6	32.7	65.6	115.0	331.0
2000.....	0.8	1.3	2.1	4.1	8.7	19.0	35.3	63.9	119.9	340.9
1999.....	1.0	1.2	2.4	4.5	9.2	18.4	36.5	66.8	124.4	360.0
Black, both sexes										
2002.....	3.1	5.9	8.1	13.2	20.7	30.9	49.3	85.2	134.7	273.8
2001.....	3.3	4.4	7.5	12.7	23.0	29.2	54.9	92.2	138.9	300.3
2000.....	3.5	5.3	8.3	13.3	22.9	32.9	52.8	92.4	146.3	315.9
1999.....	3.0	4.8	7.4	13.4	18.2	33.0	50.5	91.8	143.3	333.2
Black, male										
2002.....	4.4	7.4	12.0	19.1	29.3	36.9	64.1	100.6	159.7	276.6
2001.....	4.7	6.2	10.8	15.3	29.7	35.6	71.0	109.2	176.6	279.1
2000.....	4.6	6.5	12.0	17.5	28.6	41.4	65.9	116.2	171.1	325.2
1999.....	3.5	6.5	10.0	17.5	23.5	40.8	64.8	111.6	174.4	348.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of arteries, arterioles and capillaries (I70-I78)										
Black, female										
2002.....	1.9	4.5	4.8	8.3	13.9	26.5	39.4	76.0	121.7	272.7
2001.....	2.2	2.9	4.7	10.5	17.8	24.4	44.2	82.1	119.8	308.2
2000.....	2.5	4.4	5.2	9.9	18.5	26.7	44.2	78.0	133.7	312.5
1999.....	2.7	3.3	5.2	10.1	14.0	27.3	41.0	79.8	127.9	327.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)										
All races 3/, both sexes										
2002.....	0.1	0.3	0.6	1.4	2.4	5.2	9.3	20.5	47.8	168.1
2001.....	0.1	0.3	0.7	1.4	2.5	4.9	9.8	21.9	49.2	179.1
2000.....	0.1	0.4	0.6	1.3	2.6	5.2	10.5	22.7	52.4	188.8
1999.....	0.1	0.3	0.8	1.3	2.9	5.1	11.2	24.1	53.6	202.0
All races 3/, male										
2002.....	0.2	0.4	0.8	1.7	3.2	6.4	11.5	24.4	52.5	156.6
2001.....	*	0.4	0.8	1.7	3.2	6.2	12.1	25.0	53.2	167.2
2000.....	0.2	0.5	0.8	1.7	3.6	6.6	13.1	27.5	59.8	176.5
1999.....	0.2	0.4	1.0	1.8	3.8	6.3	13.3	28.1	60.0	189.0
All races 3/, female										
2002.....	*	*	0.5	1.0	1.7	4.1	7.5	17.8	45.0	173.1
2001.....	*	0.2	0.5	1.1	1.8	3.9	8.1	19.7	46.8	184.1
2000.....	*	0.2	0.4	0.9	1.7	4.0	8.4	19.4	48.1	193.8
1999.....	*	*	0.5	0.9	2.2	4.0	9.5	21.4	49.8	207.2
White, both sexes										
2002.....	0.1	0.2	0.5	1.3	2.3	5.3	9.1	20.4	48.4	174.7
2001.....	0.1	0.3	0.6	1.4	2.3	4.9	9.8	22.2	50.3	185.9
2000.....	0.1	0.3	0.5	1.2	2.4	5.2	10.4	23.1	53.0	195.2
1999.....	*	0.3	0.7	1.2	2.9	5.0	11.3	24.2	54.7	207.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)										
White, male										
2002.....	*	0.4	0.6	1.7	3.0	6.5	11.3	24.0	53.1	163.2
2001.....	*	0.4	0.8	1.7	3.0	6.2	12.1	25.2	54.0	173.9
2000.....	*	0.4	0.7	1.7	3.5	6.5	13.2	28.2	60.3	181.4
1999.....	*	0.4	1.0	1.6	3.7	6.2	13.3	27.7	60.8	193.7
White, female										
2002.....	*	*	0.4	1.0	1.6	4.2	7.3	17.7	45.5	179.7
2001.....	*	*	0.5	1.0	1.6	3.9	8.0	20.2	48.0	190.9
2000.....	*	*	0.3	0.8	1.5	4.1	8.2	19.5	48.7	200.8
1999.....	*	*	0.5	0.9	2.0	3.9	9.7	21.7	51.0	213.3
Black, both sexes										
2002.....	*	*	1.5	1.9	4.2	5.2	11.5	26.8	51.3	117.0
2001.....	*	*	1.0	2.0	5.0	6.3	12.1	22.9	42.8	125.6
2000.....	*	*	1.5	2.1	4.7	5.6	12.9	23.5	53.9	138.3
1999.....	*	*	1.3	2.1	4.7	7.0	11.9	27.5	46.8	154.3
Black, male										
2002.....	*	*	2.3	*	6.3	6.2	15.4	34.9	59.6	110.4
2001.....	*	*	*	*	5.9	7.6	15.1	30.2	49.0	113.5
2000.....	*	*	*	*	6.0	8.6	14.0	25.5	66.6	147.6
1999.....	*	*	*	3.4	5.7	8.8	15.6	37.0	56.7	164.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)										
Black, female										
2002.....	*	*	*	*	*	4.4	9.0	22.0	46.9	119.5
2001.....	*	*	*	*	4.3	5.3	10.1	18.6	39.7	130.1
2000.....	*	*	*	*	3.6	*	12.2	22.2	47.6	134.8
1999.....	*	*	*	*	3.8	5.6	9.5	21.7	41.9	150.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)										
All races 3/, both sexes										
2002.....	0.9	1.5	2.0	4.0	7.4	14.7	23.9	37.5	51.5	69.4
2001.....	0.9	1.4	2.0	3.7	7.8	15.0	23.8	39.0	55.4	77.5
2000.....	0.9	1.3	2.2	4.3	8.4	16.3	27.8	41.0	56.5	79.2
1999.....	0.9	1.4	2.1	4.4	8.1	16.5	27.3	42.2	58.2	80.3
All races 3/, male										
2002.....	1.4	2.4	3.2	6.3	11.5	23.0	34.9	55.0	74.5	106.6
2001.....	1.4	2.1	3.2	5.8	12.0	22.2	36.0	56.3	82.5	114.8
2000.....	1.3	1.9	3.5	6.7	13.1	24.9	41.8	61.9	84.9	127.3
1999.....	1.2	2.2	3.3	6.8	13.0	25.3	41.0	63.3	88.6	125.3
All races 3/, female										
2002.....	0.3	0.7	1.0	1.8	3.7	7.5	15.1	25.1	37.6	53.3
2001.....	0.3	0.6	0.9	1.8	3.9	8.8	14.0	26.9	39.2	61.9
2000.....	0.5	0.8	1.0	2.1	4.1	8.9	16.7	26.5	39.7	59.6
1999.....	0.5	0.7	1.0	2.1	3.7	9.0	16.6	27.5	40.4	62.2
White, both sexes										
2002.....	0.7	1.2	1.7	3.8	7.5	15.3	24.8	39.0	53.1	71.2
2001.....	0.7	1.2	1.7	3.6	7.7	15.5	24.6	39.8	57.0	79.7
2000.....	0.7	1.2	2.0	4.1	8.4	16.7	29.0	42.0	58.1	80.8
1999.....	0.7	1.2	1.9	4.2	8.5	17.1	28.4	43.5	60.0	82.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)										
White, male										
2002.....	1.2	2.0	2.7	5.9	11.6	24.3	36.1	57.5	77.7	111.8
2001.....	1.1	1.9	2.8	5.6	11.9	23.2	37.3	57.7	85.5	120.3
2000.....	1.0	1.8	3.2	6.5	13.3	26.1	43.8	63.6	88.6	132.8
1999.....	1.0	1.9	2.9	6.7	13.6	26.5	43.0	66.3	91.5	129.8
White, female										
2002.....	0.3	0.5	0.8	1.7	3.7	7.4	15.5	25.8	38.1	53.5
2001.....	0.3	0.5	0.7	1.6	3.9	8.8	14.3	27.1	39.8	62.6
2000.....	0.3	0.5	0.9	1.9	3.9	8.5	17.1	26.8	40.0	59.6
1999.....	0.4	0.5	0.9	1.8	3.7	8.9	16.7	27.5	41.3	63.1
Black, both sexes										
2002.....	2.1	3.8	4.7	6.8	8.3	11.6	18.4	24.9	37.4	50.5
2001.....	2.1	2.7	4.4	6.2	9.3	11.6	18.4	34.4	42.5	55.3
2000.....	2.3	2.9	3.7	7.3	9.3	14.3	20.3	33.8	40.4	62.3
1999.....	2.0	3.5	4.0	6.5	6.5	14.1	19.8	29.6	40.5	60.3
Black, male										
2002.....	3.5	5.7	7.7	11.3	13.1	14.7	24.7	31.2	41.3	49.3
2001.....	3.4	3.9	6.9	8.6	14.6	15.0	26.6	43.2	54.6	48.8
2000.....	3.5	3.6	6.0	10.4	12.8	17.7	27.7	47.7	45.5	69.2
1999.....	2.7	5.4	6.4	9.0	9.3	19.2	25.4	29.9	54.1	76.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)										
Black, female										
2002.....	*	2.1	2.2	3.2	4.5	9.3	14.1	21.1	35.4	50.9
2001.....	*	1.7	2.2	4.2	5.1	9.0	13.0	29.1	36.4	57.7
2000.....	*	2.3	*	4.7	6.5	11.7	15.3	25.3	37.8	59.7
1999.....	1.4	2.0	*	4.4	4.3	10.4	16.0	29.5	33.7	54.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)										
All races 3/, both sexes										
2002.....	0.4	0.7	1.1	2.2	4.3	7.7	12.6	20.4	32.3	79.0
2001.....	0.5	0.6	1.2	2.2	4.3	7.3	12.9	21.5	32.7	80.7
2000.....	0.4	0.8	1.3	2.1	4.3	8.2	12.8	21.8	35.8	84.0
1999.....	0.5	0.7	1.2	2.3	4.4	7.0	12.9	21.3	36.1	84.9
All races 3/, male										
2002.....	0.4	0.8	1.2	2.3	4.9	8.7	14.2	23.4	39.2	80.8
2001.....	0.4	0.7	1.2	2.4	4.8	8.5	15.2	24.7	40.0	82.5
2000.....	0.3	0.8	1.4	2.6	4.9	9.7	15.7	26.6	41.7	87.6
1999.....	0.4	0.8	1.2	2.6	5.2	8.1	15.7	25.9	40.8	82.2
All races 3/, female										
2002.....	0.4	0.6	1.0	2.1	3.7	6.7	11.3	18.3	28.1	78.2
2001.....	0.5	0.6	1.3	2.0	3.9	6.3	11.1	19.3	28.3	79.9
2000.....	0.5	0.7	1.1	1.6	3.7	6.9	10.5	18.5	32.2	82.6
1999.....	0.5	0.6	1.2	2.1	3.6	6.0	10.6	18.2	33.3	86.0
White, both sexes										
2002.....	0.4	0.6	1.0	2.0	3.9	7.1	12.2	19.7	31.7	77.8
2001.....	0.4	0.6	1.2	2.1	4.0	7.1	12.2	20.8	31.4	78.5
2000.....	0.3	0.7	1.0	1.9	3.9	7.8	12.4	21.0	34.9	82.3
1999.....	0.5	0.7	1.1	2.1	4.1	6.6	12.5	20.6	34.8	82.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)										
White, male										
2002.....	0.4	0.8	1.1	2.0	4.4	8.2	13.7	23.0	38.4	79.1
2001.....	0.3	0.6	1.1	2.3	4.4	8.3	14.4	24.2	38.5	80.9
2000.....	0.3	0.7	1.1	2.4	4.5	9.4	15.3	25.9	41.1	86.9
1999.....	0.4	0.8	1.1	2.4	4.9	7.9	15.3	24.8	39.6	81.3
White, female										
2002.....	0.3	0.5	0.9	2.0	3.4	6.1	11.0	17.3	27.5	77.2
2001.....	0.5	0.6	1.2	1.8	3.5	6.0	10.4	18.4	27.1	77.5
2000.....	0.4	0.6	0.9	1.4	3.3	6.5	10.0	17.6	31.2	80.4
1999.....	0.5	0.5	1.0	1.8	3.4	5.6	10.2	17.6	32.0	83.5
Black, both sexes										
2002.....	0.7	1.5	2.0	4.4	8.3	14.1	19.4	33.5	46.1	106.3
2001.....	1.1	1.3	2.1	4.5	8.7	11.3	24.4	34.9	53.6	119.4
2000.....	0.9	1.7	3.1	3.9	9.0	13.0	19.6	35.2	52.0	115.4
1999.....	0.8	1.0	2.1	4.9	7.0	11.9	18.8	34.6	56.1	118.6
Black, male										
2002.....	*	*	2.1	5.2	9.9	16.0	24.0	34.4	58.8	116.9
2001.....	*	*	2.5	4.5	9.2	13.0	29.3	35.8	73.0	116.9
2000.....	*	1.9	4.0	4.8	9.7	15.1	24.1	43.0	59.0	108.4
1999.....	*	*	*	5.1	8.5	12.7	23.7	44.6	63.7	107.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Black, female

2002.....	*	1.9	1.8	3.7	7.0	12.8	16.4	32.9	39.4	102.3
2001.....	*	*	1.8	4.5	8.3	10.1	21.1	34.4	43.7	120.4
2000.....	*	*	2.4	3.2	8.3	11.5	16.6	30.5	48.4	118.0
1999.....	*	*	*	4.7	5.8	11.4	15.5	28.6	52.3	122.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of circulatory system (I80-I99)										
All races 3/, both sexes										
2002.....	0.8	1.2	1.5	2.0	2.5	3.4	4.3	7.4	13.2	24.4
2001.....	0.8	1.1	1.3	1.9	2.6	3.8	5.4	7.7	11.9	26.1
2000.....	0.9	1.2	1.5	1.7	2.6	3.4	5.3	7.7	11.9	25.6
1999.....	0.8	1.0	1.0	1.6	2.2	3.3	4.3	6.8	12.7	23.4
All races 3/, male										
2002.....	0.8	1.3	1.6	2.3	2.6	3.6	4.2	7.0	13.3	20.5
2001.....	0.9	1.2	1.5	2.2	2.9	3.9	5.9	8.3	12.1	22.0
2000.....	1.1	1.5	1.7	1.9	2.7	3.9	5.6	8.0	12.4	22.2
1999.....	0.9	1.1	1.2	1.7	2.3	3.7	4.8	7.5	13.2	20.8
All races 3/, female										
2002.....	0.8	1.1	1.4	1.7	2.5	3.2	4.4	7.7	13.1	26.2
2001.....	0.8	0.9	1.0	1.7	2.3	3.7	5.0	7.3	11.7	27.8
2000.....	0.7	1.0	1.3	1.5	2.6	3.0	5.0	7.5	11.6	26.9
1999.....	0.6	1.0	0.9	1.5	2.1	2.9	3.9	6.3	12.4	24.5
White, both sexes										
2002.....	0.6	1.0	1.4	1.9	2.2	3.2	4.1	7.1	12.7	24.3
2001.....	0.7	0.9	1.1	1.8	2.3	3.6	5.1	7.2	11.8	26.0
2000.....	0.8	1.1	1.4	1.5	2.4	2.9	5.0	7.3	11.8	25.5
1999.....	0.6	0.9	0.9	1.6	2.1	2.9	4.1	6.5	12.4	23.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of circulatory system (I80-I99)										
White, male										
2002.....	0.6	1.1	1.5	2.1	2.3	3.3	4.0	6.7	12.4	19.9
2001.....	0.7	1.2	1.3	2.0	2.5	3.6	5.6	7.8	12.0	21.7
2000.....	1.0	1.3	1.6	1.7	2.4	3.4	5.5	7.8	12.3	22.0
1999.....	0.8	0.9	1.0	1.7	2.3	3.2	4.7	7.0	12.8	20.7
White, female										
2002.....	0.7	1.0	1.2	1.6	2.1	3.0	4.2	7.4	12.9	26.3
2001.....	0.7	0.7	1.0	1.5	2.1	3.5	4.8	6.8	11.7	27.9
2000.....	0.5	0.9	1.1	1.4	2.4	2.5	4.7	7.0	11.5	26.9
1999.....	0.5	0.9	0.9	1.5	1.9	2.7	3.7	6.1	12.2	24.2
Black, both sexes										
2002.....	1.8	2.4	2.9	4.2	5.9	6.4	7.2	12.9	22.2	30.0
2001.....	1.8	2.1	2.4	3.9	6.1	6.9	9.0	14.1	14.9	30.1
2000.....	1.8	2.5	3.2	3.9	5.1	8.7	9.1	14.6	14.1	30.0
1999.....	1.9	2.4	2.3	2.1	3.9	6.9	6.4	11.4	17.6	28.4
Black, male										
2002.....	2.1	3.0	2.9	5.4	5.9	7.7	7.7	12.4	28.3	32.2
2001.....	2.2	1.7	3.1	4.7	8.5	7.9	10.4	16.3	*	29.5
2000.....	2.0	3.1	3.7	4.8	5.6	9.4	8.7	13.2	*	30.0
1999.....	2.6	2.7	3.0	*	*	9.3	6.8	14.7	20.9	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of circulatory system (I80-I99)										
Black, female										
2002.....	1.6	1.8	2.9	3.2	5.9	5.4	6.9	13.2	19.1	29.2
2001.....	1.5	2.4	*	3.2	4.2	6.2	8.1	12.8	14.7	30.3
2000.....	1.7	1.9	2.8	3.2	4.7	8.3	9.3	15.4	14.4	30.0
1999.....	*	2.1	*	*	4.2	5.0	6.1	9.4	16.0	30.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)										
All races 3/, both sexes										
2002.....	0.7	0.9	1.2	1.6	1.9	2.6	3.1	5.4	10.1	18.6
2001.....	0.6	0.8	0.9	1.4	2.0	2.8	4.1	5.9	8.9	19.6
2000.....	0.6	0.9	1.1	1.3	1.8	2.5	3.9	5.8	8.8	18.7
1999.....	0.6	0.7	0.7	1.1	1.4	2.2	2.8	4.3	8.0	14.6
All races 3/, male										
2002.....	0.6	0.9	1.1	1.7	1.7	2.7	2.9	4.7	9.6	14.8
2001.....	0.6	0.8	1.0	1.5	2.1	2.7	4.4	5.9	8.3	15.2
2000.....	0.7	0.9	1.1	1.3	1.6	2.8	3.8	5.7	8.9	15.0
1999.....	0.6	0.6	0.6	1.0	1.2	2.3	3.1	4.4	7.9	11.2
All races 3/, female										
2002.....	0.7	0.9	1.2	1.5	2.0	2.6	3.2	5.9	10.4	20.2
2001.....	0.7	0.8	0.8	1.3	1.9	2.9	3.8	5.9	9.2	21.4
2000.....	0.5	0.9	1.1	1.2	1.9	2.3	4.0	5.9	8.7	20.2
1999.....	0.6	0.7	0.7	1.1	1.5	2.0	2.5	4.2	8.1	15.9
White, both sexes										
2002.....	0.5	0.8	1.0	1.5	1.6	2.5	2.9	5.2	9.6	18.4
2001.....	0.5	0.7	0.8	1.3	1.7	2.6	3.8	5.6	8.9	19.6
2000.....	0.6	0.8	1.0	1.1	1.6	2.1	3.7	5.5	8.8	18.8
1999.....	0.5	0.6	0.6	1.0	1.3	2.0	2.7	4.1	7.8	14.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)										
White, male										
2002.....	0.4	0.8	1.0	1.5	1.5	2.4	2.8	4.7	8.8	14.4
2001.....	0.5	0.7	0.9	1.4	1.7	2.6	4.1	5.7	8.3	15.0
2000.....	0.7	0.8	1.0	1.1	1.4	2.5	3.6	5.6	9.0	14.9
1999.....	0.5	0.6	0.5	1.0	1.2	2.0	3.0	4.2	7.7	11.4
White, female										
2002.....	0.6	0.9	1.0	1.4	1.8	2.5	3.0	5.6	10.1	20.2
2001.....	0.6	0.6	0.8	1.2	1.7	2.6	3.5	5.6	9.2	21.5
2000.....	0.5	0.8	0.9	1.1	1.8	1.8	3.8	5.4	8.7	20.4
1999.....	0.5	0.6	0.7	1.1	1.4	1.9	2.4	4.1	7.8	15.5
Black, both sexes										
2002.....	1.6	1.8	2.5	3.2	4.4	4.9	5.6	9.3	18.3	23.8
2001.....	1.5	1.8	1.7	2.9	5.3	5.3	7.8	10.4	11.6	23.3
2000.....	1.3	1.9	2.5	2.9	3.9	6.8	7.1	11.7	9.3	21.9
1999.....	1.5	1.6	1.5	1.6	2.4	4.8	4.5	7.1	12.7	18.8
Black, male										
2002.....	1.7	2.1	2.4	4.2	4.4	5.7	*	*	22.2	22.5
2001.....	1.7	*	2.4	3.3	6.9	5.1	8.8	11.2	*	*
2000.....	*	2.2	2.4	*	*	6.5	7.0	*	*	*
1999.....	1.7	*	*	*	*	6.8	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)										
Black, female										
2002.....	1.6	1.6	2.6	*	4.5	4.3	5.8	11.0	16.3	24.2
2001.....	1.3	2.2	*	2.6	4.0	5.4	7.2	10.0	12.3	24.0
2000.....	*	1.7	2.6	2.7	3.9	7.1	7.1	13.4	10.6	22.3
1999.....	*	1.8	*	*	*	*	4.5	6.3	12.5	21.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified disorders of circulatory system (I83-I99)										
All races 3/, both sexes										
2002.....	0.1	0.3	0.3	0.4	0.7	0.7	1.2	2.1	3.1	5.9
2001.....	0.2	0.3	0.4	0.5	0.6	1.0	1.4	1.8	3.0	6.5
2000.....	0.2	0.3	0.4	0.4	0.9	0.9	1.4	1.9	3.1	6.8
1999.....	0.2	0.3	0.4	0.5	0.8	1.1	1.5	2.5	4.7	8.8
All races 3/, male										
2002.....	0.2	0.4	0.5	0.6	0.8	0.9	1.3	2.3	3.7	5.7
2001.....	0.3	0.4	0.5	0.7	0.8	1.2	1.5	2.4	3.8	6.8
2000.....	0.3	0.5	0.6	0.6	1.1	1.0	1.8	2.4	3.5	7.3
1999.....	0.3	0.4	0.5	0.7	1.1	1.4	1.7	3.1	5.3	9.6
All races 3/, female										
2002.....	*	*	0.2	0.3	0.5	0.6	1.2	1.9	2.7	5.9
2001.....	*	*	0.2	0.3	0.4	0.8	1.2	1.4	2.5	6.4
2000.....	*	*	0.2	*	0.7	0.8	1.0	1.6	2.9	6.7
1999.....	*	0.3	*	0.4	0.6	0.9	1.3	2.1	4.3	8.5
White, both sexes										
2002.....	0.1	0.2	0.3	0.4	0.6	0.7	1.2	2.0	3.1	5.9
2001.....	0.2	0.3	0.3	0.5	0.6	0.9	1.3	1.6	3.0	6.5
2000.....	0.2	0.3	0.4	0.4	0.8	0.8	1.3	1.9	3.0	6.7
1999.....	0.2	0.3	0.3	0.5	0.8	1.0	1.5	2.4	4.6	8.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
X. Diseases of the respiratory system (J00-J98)										
All races 3/, both sexes										
2002.....	7.6	13.1	22.8	45.7	91.4	172.4	315.3	528.4	849.0	1,663.7
2001.....	7.4	12.7	23.1	46.9	93.4	178.6	319.0	526.3	833.1	1,671.8
2000.....	8.0	13.2	23.3	48.3	94.2	181.4	324.3	529.4	874.7	1,734.0
1999.....	7.4	12.9	22.4	49.4	97.1	185.1	330.2	537.8	873.1	1,733.0
All races 3/, male										
2002.....	8.3	14.2	25.5	50.2	102.9	197.5	379.7	661.7	1,094.2	2,150.0
2001.....	8.0	14.3	25.7	51.9	104.0	204.9	385.2	662.7	1,080.0	2,187.2
2000.....	8.9	14.8	25.7	54.2	106.1	210.7	394.4	666.6	1,158.0	2,344.4
1999.....	7.9	14.6	24.5	54.1	111.1	220.6	411.8	691.1	1,170.2	2,356.4
All races 3/, female										
2002.....	6.8	12.0	20.2	41.5	81.1	150.7	263.8	434.3	699.8	1,452.7
2001.....	6.8	11.0	20.6	42.2	83.9	155.9	266.4	430.8	685.2	1,456.2
2000.....	7.1	11.6	21.0	42.9	83.5	156.3	269.0	433.8	707.6	1,485.4
1999.....	7.0	11.4	20.5	45.0	84.3	154.8	266.4	431.1	699.3	1,481.7
White, both sexes										
2002.....	6.7	12.1	21.6	45.3	93.0	179.4	326.6	545.8	873.3	1,713.3
2001.....	6.5	11.4	21.7	46.7	95.4	184.8	329.8	542.3	856.5	1,713.5
2000.....	7.0	11.6	21.8	47.7	95.6	185.4	333.7	544.9	898.9	1,783.5
1999.....	6.1	11.2	21.2	48.7	98.7	190.9	338.4	551.4	897.0	1,778.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
X. Diseases of the respiratory system (J00-J98)										
White, male										
2002.....	7.3	13.0	23.9	48.8	102.3	202.3	386.7	672.5	1,114.4	2,201.7
2001.....	7.1	13.1	23.6	51.2	103.7	208.2	391.6	672.0	1,098.6	2,230.5
2000.....	7.8	12.9	23.5	52.6	105.9	210.3	398.5	676.6	1,179.6	2,402.2
1999.....	6.3	12.6	22.5	52.3	110.8	223.6	415.4	699.7	1,188.8	2,409.4
White, female										
2002.....	6.0	11.3	19.3	42.0	84.4	159.3	277.5	455.0	725.5	1,501.2
2001.....	5.9	9.6	19.8	42.3	87.8	164.3	279.7	450.5	710.2	1,497.0
2000.....	6.2	10.3	20.2	43.1	86.1	163.6	281.7	452.2	731.9	1,532.2
1999.....	5.9	9.9	19.9	45.3	87.5	162.3	277.1	447.2	724.8	1,525.0
Black, both sexes										
2002.....	15.5	23.1	36.6	61.9	102.5	154.4	271.2	424.1	670.3	1,204.5
2001.....	14.9	24.9	39.3	60.7	99.0	162.5	275.3	430.7	647.1	1,258.0
2000.....	16.0	27.1	40.4	65.3	106.0	183.3	288.6	427.5	662.7	1,239.4
1999.....	17.6	28.0	38.2	68.0	105.5	173.1	302.0	456.5	655.6	1,298.0
Black, male										
2002.....	17.9	26.6	43.7	77.9	135.8	204.2	378.9	634.7	986.6	1,685.3
2001.....	16.2	27.3	47.7	72.0	130.6	218.5	390.5	653.7	981.6	1,775.7
2000.....	18.5	32.3	48.7	82.0	134.0	254.7	419.2	634.6	980.4	1,808.3
1999.....	19.8	33.3	47.4	83.9	141.1	237.6	438.2	682.9	1,034.1	1,900.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
X. Diseases of the respiratory system (J00-J98)										
Black, female										
2002.....	13.4	20.1	30.5	48.8	76.5	117.5	198.9	298.3	505.7	1,020.3
2001.....	13.8	22.8	32.1	51.4	74.1	121.1	198.6	297.6	476.7	1,066.0
2000.....	13.7	22.6	33.4	51.7	84.0	130.6	201.8	302.8	502.7	1,027.6
1999.....	15.6	23.5	30.4	55.0	77.5	125.3	211.5	320.1	468.0	1,072.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute upper respiratory infections (J00-J06)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*	*	*	1.1
2001.....	*	*	*	*	*	*	*	*	*	1.3
2000.....	*	*	*	*	*	*	*	*	0.4	1.7
1999.....	*	*	*	*	*	*	*	*	0.5	1.7

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	1.2
2001.....	*	*	*	*	*	*	*	*	*	1.6
2000.....	*	*	*	*	*	*	*	*	*	2.0
1999.....	*	*	*	*	*	*	*	*	*	2.0

White, both sexes

2002.....	*	*	*	*	*	*	*	*	*	1.1
2001.....	*	*	*	*	*	*	*	*	*	1.4
2000.....	*	*	*	*	*	*	*	*	*	1.8
1999.....	*	*	*	*	*	*	*	*	0.5	1.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*	*	*	1.0
2001.....	*	*	*	*	*	*	*	*	*	1.1
2000.....	*	*	*	*	*	*	*	*	*	1.7
1999.....	*	*	*	*	*	*	*	*	0.5	1.6

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	1.2
2001.....	*	*	*	*	*	*	*	*	*	1.4
2000.....	*	*	*	*	*	*	*	*	*	1.9
1999.....	*	*	*	*	*	*	*	*	*	1.9

White, both sexes

2002.....	*	*	*	*	*	*	*	*	*	1.1
2001.....	*	*	*	*	*	*	*	*	*	1.2
2000.....	*	*	*	*	*	*	*	*	*	1.7
1999.....	*	*	*	*	*	*	*	*	0.5	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of the respiratory system (J10-J98)										
All races 3/, both sexes										
2002.....	7.5	13.1	22.7	45.6	91.4	172.3	315.1	528.3	848.8	1,662.6
2001.....	7.4	12.6	23.1	46.8	93.4	178.5	318.9	526.1	832.8	1,670.6
2000.....	8.0	13.2	23.2	48.3	94.2	181.3	324.1	529.1	874.3	1,732.2
1999.....	7.4	12.9	22.4	49.4	97.0	185.1	330.1	537.5	872.6	1,731.2
All races 3/, male										
2002.....	8.3	14.2	25.4	50.1	102.9	197.4	379.5	661.6	1,094.0	2,149.3
2001.....	7.9	14.3	25.7	51.9	104.0	204.9	385.2	662.6	1,079.8	2,186.6
2000.....	8.9	14.8	25.6	54.2	106.1	210.6	394.3	666.5	1,157.7	2,343.3
1999.....	7.8	14.5	24.5	54.1	111.0	220.5	411.7	690.9	1,169.5	2,355.3
All races 3/, female										
2002.....	6.8	11.9	20.1	41.4	81.0	150.6	263.7	434.1	699.7	1,451.5
2001.....	6.8	11.0	20.6	42.1	83.8	155.9	266.3	430.6	684.9	1,454.6
2000.....	7.1	11.6	20.9	42.8	83.5	156.2	268.8	433.5	707.2	1,483.4
1999.....	7.0	11.3	20.4	45.0	84.3	154.8	266.3	430.9	698.8	1,479.7
White, both sexes										
2002.....	6.6	12.1	21.6	45.3	93.0	179.4	326.4	545.6	873.2	1,712.2
2001.....	6.5	11.3	21.7	46.6	95.4	184.8	329.7	542.2	856.2	1,712.1
2000.....	6.9	11.6	21.8	47.7	95.5	185.3	333.5	544.6	898.5	1,781.7
1999.....	6.1	11.2	21.2	48.7	98.6	190.8	338.3	551.1	896.5	1,776.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of the respiratory system (J10-J98)										
White, male										
2002.....	7.3	12.9	23.9	48.8	102.3	202.2	386.6	672.4	1,114.3	2,201.0
2001.....	7.0	13.1	23.6	51.2	103.6	208.1	391.6	671.9	1,098.3	2,229.8
2000.....	7.8	12.9	23.5	52.6	105.8	210.3	398.3	676.5	1,179.3	2,401.1
1999.....	6.2	12.5	22.5	52.3	110.8	223.5	415.3	699.4	1,188.2	2,408.6
White, female										
2002.....	6.0	11.2	19.3	41.9	84.3	159.2	277.4	454.8	725.4	1,499.8
2001.....	5.9	9.6	19.8	42.3	87.8	164.2	279.6	450.3	709.9	1,495.4
2000.....	6.1	10.3	20.2	43.1	86.1	163.5	281.5	451.8	731.5	1,530.1
1999.....	5.9	9.9	19.8	45.3	87.4	162.3	277.0	447.0	724.3	1,522.9
Black, both sexes										
2002.....	15.4	23.0	36.5	61.8	102.5	154.2	271.2	424.1	669.8	1,204.5
2001.....	14.9	24.8	39.2	60.6	98.8	162.5	275.2	430.5	647.1	1,257.1
2000.....	16.0	27.0	40.4	65.3	106.0	183.1	288.6	427.5	662.4	1,238.8
1999.....	17.6	28.0	38.1	67.9	105.3	173.1	301.6	456.5	655.3	1,296.4
Black, male										
2002.....	17.8	26.6	43.5	77.9	135.8	203.9	378.9	634.7	985.8	1,685.3
2001.....	16.1	27.2	47.7	71.8	130.4	218.5	390.5	653.7	981.6	1,775.7
2000.....	18.5	32.3	48.7	82.0	134.0	254.2	419.2	634.6	979.5	1,806.0
1999.....	19.8	33.2	47.4	83.9	140.7	237.6	437.9	682.9	1,034.1	1,895.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of the respiratory system (J10-J98)										
Black, female										
2002.....	13.4	20.0	30.5	48.6	76.5	117.3	198.9	298.3	505.3	1,020.3
2001.....	13.8	22.8	32.0	51.4	73.9	121.1	198.4	297.3	476.7	1,064.7
2000.....	13.7	22.5	33.3	51.7	84.0	130.6	201.8	302.8	502.7	1,027.6
1999.....	15.6	23.5	30.3	54.8	77.5	125.3	211.1	320.1	467.6	1,072.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza (J10-J11)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	*	0.4	0.8	2.2	8.8
2001.....	*	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	0.1	0.1	0.3	0.5	0.8	1.4	2.5	6.1	20.2
1999.....	*	*	*	0.2	0.4	0.6	1.2	2.1	6.1	21.2
All races 3/, male										
2002.....	*	*	*	*	*	*	*	0.9	2.6	8.9
2001.....	*	*	*	*	*	*	*	*	*	2.2
2000.....	*	*	*	0.3	0.5	1.0	1.6	2.7	6.3	20.7
1999.....	*	*	*	*	0.5	0.6	1.2	2.4	7.3	19.9
All races 3/, female										
2002.....	*	*	*	*	*	*	*	0.7	2.0	8.7
2001.....	*	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	*	*	*	0.5	0.6	1.2	2.5	6.0	19.9
1999.....	*	*	*	*	*	0.6	1.1	1.9	5.4	21.7
White, both sexes										
2002.....	*	*	*	*	*	*	0.4	0.9	2.4	9.4
2001.....	*	*	*	*	*	*	*	*	0.7	2.3
2000.....	*	*	0.1	0.3	0.5	0.8	1.4	2.7	6.5	21.4
1999.....	*	*	*	0.2	0.4	0.7	1.2	2.2	6.4	22.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)										
All races 3/, both sexes										
2002.....	2.6	4.1	5.4	8.4	14.6	25.1	50.5	105.7	225.1	687.8
2001.....	2.6	3.9	5.3	8.1	13.8	24.3	49.2	100.9	216.1	683.4
2000.....	2.9	3.8	5.5	8.6	15.1	25.8	51.2	104.1	234.7	723.9
1999.....	2.7	4.0	5.2	8.2	13.9	24.3	49.3	103.8	227.7	730.6
All races 3/, male										
2002.....	3.0	4.9	6.7	10.1	17.8	30.9	63.4	133.0	285.0	796.9
2001.....	3.0	4.8	6.6	9.9	16.9	30.3	61.4	130.5	273.9	806.2
2000.....	3.7	4.6	6.8	10.6	18.4	31.9	64.1	131.5	302.6	871.3
1999.....	3.1	5.1	6.7	9.5	16.3	31.1	63.0	131.1	294.5	874.7
All races 3/, female										
2002.....	2.2	3.4	4.3	6.8	11.8	20.0	40.2	86.5	188.6	640.5
2001.....	2.2	2.9	4.1	6.3	11.0	19.1	39.4	80.2	181.5	632.0
2000.....	2.1	3.0	4.2	6.7	12.0	20.5	41.0	85.1	194.6	663.9
1999.....	2.3	2.9	3.9	7.0	11.7	18.6	38.6	84.8	188.7	672.5
White, both sexes										
2002.....	2.3	3.8	4.9	7.6	13.7	24.3	49.5	104.5	225.6	699.6
2001.....	2.1	3.3	4.8	7.4	12.9	23.2	48.0	98.3	215.8	692.7
2000.....	2.4	3.2	4.8	7.8	13.8	23.7	48.9	102.9	234.3	737.0
1999.....	2.1	3.3	4.6	7.4	12.6	23.4	47.6	103.3	229.2	741.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)										
White, male										
2002.....	2.7	4.3	6.0	9.0	16.3	29.6	61.9	130.4	283.7	806.2
2001.....	2.5	4.1	5.6	9.0	15.3	28.3	59.2	125.3	271.5	812.1
2000.....	3.1	3.8	5.7	9.2	16.7	28.6	60.6	129.1	299.7	884.4
1999.....	2.2	4.2	5.6	8.4	14.6	29.5	59.7	129.6	294.3	884.0
White, female										
2002.....	1.9	3.2	3.9	6.2	11.3	19.7	39.4	86.0	190.0	653.3
2001.....	1.8	2.4	3.9	5.9	10.6	18.8	38.9	79.3	182.2	642.8
2000.....	1.8	2.7	3.9	6.4	11.1	19.4	39.6	84.5	195.4	677.1
1999.....	1.9	2.5	3.5	6.5	10.8	18.0	37.9	84.8	190.7	684.4
Black, both sexes										
2002.....	5.2	7.5	10.1	17.2	26.1	35.0	68.6	123.2	230.8	572.8
2001.....	5.8	9.0	10.7	15.4	23.1	34.1	65.7	133.7	222.9	579.7
2000.....	6.0	8.1	12.1	17.7	26.9	45.4	76.0	123.5	243.1	591.5
1999.....	6.8	9.7	11.6	16.0	26.4	36.5	73.9	117.8	217.0	629.1
Black, male										
2002.....	6.2	9.9	12.9	23.5	35.6	47.4	89.8	173.2	320.2	687.2
2001.....	7.0	11.4	15.7	20.6	33.2	49.6	94.9	203.5	309.0	712.6
2000.....	8.1	11.0	17.0	25.5	35.0	64.3	106.6	170.6	343.1	742.7
1999.....	8.9	13.3	17.1	21.7	33.9	51.4	108.2	159.6	307.8	800.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)										
Black, female										
2002.....	4.3	5.5	7.7	12.0	18.6	25.9	54.4	93.3	184.2	529.0
2001.....	4.7	6.9	6.5	11.2	15.2	22.7	46.2	92.1	179.1	530.5
2000.....	4.2	5.6	7.9	11.2	20.4	31.5	55.7	95.1	192.8	535.3
1999.....	5.0	6.6	7.0	11.4	20.5	25.4	51.1	92.7	172.0	565.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral pneumonia, not elsewhere classified (J12)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	0.3	0.4	0.7	1.7
2001.....	*	*	*	*	*	*	*	0.3	0.7	1.4
2000.....	*	*	*	*	*	0.2	0.3	0.6	1.0	2.9
1999.....	*	*	*	*	*	0.2	0.2	0.5	1.2	2.4

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	1.0	1.9
2001.....	*	*	*	*	*	*	*	*	1.3	1.9
2000.....	*	*	*	*	*	*	*	0.7	*	2.4
1999.....	*	*	*	*	*	*	*	0.9	1.7	1.8

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	1.6
2001.....	*	*	*	*	*	*	*	*	*	1.2
2000.....	*	*	*	*	*	*	*	0.5	1.0	3.0
1999.....	*	*	*	*	*	*	*	*	0.9	2.6

White, both sexes

2002.....	*	*	*	*	*	*	0.3	0.4	0.8	1.8
2001.....	*	*	*	*	*	*	*	0.3	0.8	1.5
2000.....	*	*	*	*	*	0.3	0.3	0.6	1.0	3.0
1999.....	*	*	*	*	*	0.2	0.3	0.5	1.3	2.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bacterial pneumonia (J13-J15)										
All races 3/, both sexes										
2002.....	0.4	0.6	0.7	1.2	1.6	2.6	4.7	8.3	15.5	35.7
2001.....	0.3	0.5	0.7	1.4	1.6	2.7	4.4	8.4	14.8	34.8
2000.....	0.5	0.6	0.8	1.3	1.9	2.7	4.9	8.1	15.3	35.7
1999.....	0.5	0.5	0.6	1.1	1.5	2.4	4.3	7.2	13.6	30.1
All races 3/, male										
2002.....	0.4	0.6	0.8	1.4	1.8	3.0	6.1	11.3	20.9	46.8
2001.....	0.5	0.6	0.9	1.6	2.0	3.6	5.6	11.7	20.6	46.8
2000.....	0.6	0.6	0.9	1.5	2.3	3.3	6.4	11.9	21.7	52.3
1999.....	0.5	0.6	0.8	1.2	1.6	3.0	5.5	9.0	18.4	40.4
All races 3/, female										
2002.....	0.4	0.5	0.6	0.9	1.4	2.3	3.6	6.2	12.2	31.0
2001.....	0.2	0.5	0.6	1.1	1.3	1.9	3.4	6.2	11.3	29.8
2000.....	0.4	0.5	0.6	1.1	1.6	2.1	3.7	5.4	11.6	29.0
1999.....	0.5	0.4	0.4	0.9	1.5	1.8	3.3	5.9	10.8	26.0
White, both sexes										
2002.....	0.4	0.5	0.7	1.0	1.5	2.6	4.7	8.3	15.3	36.3
2001.....	0.3	0.5	0.7	1.4	1.5	2.6	4.2	8.1	14.8	35.0
2000.....	0.4	0.5	0.7	1.2	1.9	2.5	4.7	7.9	15.0	35.6
1999.....	0.4	0.5	0.5	1.0	1.4	2.3	4.1	7.1	13.6	30.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bacterial pneumonia (J13-J15)										
White, male										
2002.....	0.4	0.5	0.7	1.2	1.7	2.9	5.9	11.1	20.1	47.9
2001.....	0.4	0.5	0.8	1.6	1.8	3.5	5.3	10.9	20.5	46.6
2000.....	0.6	0.6	0.9	1.4	2.1	3.0	6.1	11.4	21.7	51.1
1999.....	0.4	0.5	0.7	1.1	1.5	2.9	5.2	8.9	18.1	40.1
White, female										
2002.....	0.3	0.5	0.6	0.8	1.4	2.3	3.7	6.3	12.3	31.2
2001.....	*	0.5	0.7	1.1	1.2	1.8	3.4	6.1	11.3	30.2
2000.....	0.3	0.4	0.6	1.0	1.6	2.0	3.6	5.5	11.0	29.2
1999.....	0.4	0.4	0.4	0.9	1.3	1.8	3.2	5.9	11.0	26.3
Black, both sexes										
2002.....	0.8	1.1	1.0	2.3	2.2	2.9	5.4	8.4	16.0	28.5
2001.....	0.7	*	1.2	1.6	3.0	3.3	6.1	12.3	14.1	32.5
2000.....	*	1.0	1.3	2.0	2.8	4.6	7.1	8.9	17.5	34.7
1999.....	1.1	1.0	*	2.1	2.6	2.8	6.9	8.2	12.1	25.2
Black, male										
2002.....	*	*	*	3.4	*	*	7.7	13.3	26.0	32.2
2001.....	*	*	*	*	*	5.1	10.1	21.4	18.5	49.9
2000.....	*	*	*	*	4.1	5.5	10.7	16.1	21.1	60.0
1999.....	*	*	*	*	*	*	9.8	10.9	20.9	38.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bacterial pneumonia (J13-J15)										
Black, female										
2002.....	*	*	*	*	*	*	*	5.5	10.7	27.1
2001.....	*	*	*	*	*	*	*	6.9	11.9	26.1
2000.....	*	*	*	*	*	4.0	4.7	*	15.7	25.3
1999.....	*	*	*	*	*	*	5.0	6.6	*	20.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia due to other or unspecified organisms (J16,J18)										
All races 3/, both sexes										
2002.....	2.2	3.5	4.7	7.2	13.0	22.4	45.5	97.0	208.8	650.3
2001.....	2.2	3.3	4.6	6.7	12.1	21.5	44.7	92.1	200.6	647.1
2000.....	2.4	3.2	4.7	7.2	13.0	22.9	46.0	95.5	218.3	685.3
1999.....	2.2	3.5	4.6	7.0	12.2	21.7	44.8	96.1	212.9	698.1
All races 3/, male										
2002.....	2.6	4.2	5.8	8.6	15.9	27.8	56.9	121.3	263.1	748.2
2001.....	2.5	4.2	5.7	8.2	14.9	26.5	55.7	118.3	252.1	757.4
2000.....	3.1	3.9	5.8	9.0	16.0	28.3	57.5	118.9	279.9	816.5
1999.....	2.6	4.5	5.8	8.1	14.5	27.9	57.2	121.2	274.4	832.5
All races 3/, female										
2002.....	1.7	2.8	3.6	5.8	10.3	17.7	36.4	79.8	175.8	607.9
2001.....	1.9	2.4	3.5	5.2	9.6	17.1	36.0	73.8	169.8	601.0
2000.....	1.7	2.5	3.6	5.6	10.3	18.2	37.0	79.2	182.0	631.9
1999.....	1.8	2.5	3.4	6.0	10.1	16.5	35.0	78.7	177.0	643.9
White, both sexes										
2002.....	1.9	3.2	4.2	6.5	12.1	21.7	44.5	95.8	209.5	661.6
2001.....	1.8	2.7	4.0	6.0	11.3	20.5	43.6	89.9	200.3	656.2
2000.....	2.0	2.7	4.0	6.5	11.8	21.0	43.9	94.4	218.3	698.3
1999.....	1.7	2.9	4.0	6.4	11.1	20.8	43.2	95.8	214.3	708.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia due to other or unspecified organisms (J16,J18)										
White, male										
2002.....	2.3	3.8	5.2	7.7	14.5	26.6	55.6	118.8	262.5	756.4
2001.....	2.1	3.5	4.8	7.3	13.5	24.6	53.8	113.8	249.6	763.5
2000.....	2.5	3.3	4.8	7.7	14.5	25.3	54.2	117.0	276.9	830.6
1999.....	1.9	3.7	4.9	7.2	12.9	26.4	54.2	119.9	274.5	841.9
White, female										
2002.....	1.6	2.7	3.2	5.3	9.8	17.3	35.5	79.3	177.0	620.4
2001.....	1.6	2.0	3.2	4.7	9.3	16.9	35.4	73.0	170.5	611.3
2000.....	1.5	2.2	3.2	5.3	9.4	17.2	35.7	78.5	183.3	644.6
1999.....	1.5	2.1	3.1	5.6	9.4	15.9	34.5	78.8	178.7	655.4
Black, both sexes										
2002.....	4.4	6.4	9.1	14.9	23.9	32.2	63.1	114.6	214.6	543.4
2001.....	5.1	8.2	9.5	13.8	20.1	30.6	59.5	121.4	208.6	546.9
2000.....	5.4	7.1	10.7	15.6	23.8	40.7	68.7	114.6	225.3	556.5
1999.....	5.7	8.6	10.6	13.9	23.8	33.5	67.0	109.1	204.3	602.6
Black, male										
2002.....	5.5	8.6	11.7	20.1	32.7	44.4	82.1	159.4	293.5	652.9
2001.....	5.9	10.1	13.7	18.6	29.3	44.2	84.5	182.1	290.5	661.5
2000.....	7.5	10.0	15.3	22.6	30.6	58.8	95.6	154.5	322.0	682.7
1999.....	7.7	11.8	15.4	18.8	31.6	47.5	98.4	147.2	285.1	760.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia due to other or unspecified organisms (J16,J18)										
Black, female										
2002.....	3.4	4.5	6.9	10.6	17.0	23.1	50.4	87.8	173.5	501.5
2001.....	4.3	6.4	5.9	9.9	12.8	20.5	42.8	85.1	166.8	504.4
2000.....	3.5	4.6	6.8	9.9	18.5	27.3	50.8	90.6	176.6	509.5
1999.....	4.0	5.9	6.6	9.8	17.6	23.1	46.1	86.1	164.2	543.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute lower respiratory infections (J20-J22)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	0.3	0.3	0.8	3.3
2001.....	*	*	*	*	*	*	*	0.3	0.7	3.3
2000.....	*	*	*	*	*	*	0.3	0.6	0.8	4.5
1999.....	*	*	*	*	*	*	0.2	0.6	1.3	6.0

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	*	2.8
2001.....	*	*	*	*	*	*	*	*	*	2.3
2000.....	*	*	*	*	*	*	*	0.7	*	3.8
1999.....	*	*	*	*	*	*	*	0.7	1.6	4.0

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	0.9	3.5
2001.....	*	*	*	*	*	*	*	*	0.7	3.7
2000.....	*	*	*	*	*	*	*	0.5	0.9	4.8
1999.....	*	*	*	*	*	*	*	0.5	1.1	6.8

White, both sexes

2002.....	*	*	*	*	*	*	*	*	0.8	3.5
2001.....	*	*	*	*	*	*	*	0.3	0.7	3.5
2000.....	*	*	*	*	*	*	0.3	0.6	0.9	4.7
1999.....	*	*	*	*	*	*	*	0.6	1.3	6.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute bronchitis and bronchiolitis (J20-J21)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*	*	0.5	2.0
2001.....	*	*	*	*	*	*	*	*	0.4	2.1
2000.....	*	*	*	*	*	*	*	0.4	0.5	2.6
1999.....	*	*	*	*	*	*	*	0.4	0.8	3.6

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	2.4
1999.....	*	*	*	*	*	*	*	*	*	2.6

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	2.2
2001.....	*	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	*	*	*	*	*	*	2.7
1999.....	*	*	*	*	*	*	*	*	0.7	4.0

White, both sexes

2002.....	*	*	*	*	*	*	*	*	0.5	2.1
2001.....	*	*	*	*	*	*	*	*	0.5	2.1
2000.....	*	*	*	*	*	*	*	0.4	0.5	2.7
1999.....	*	*	*	*	*	*	*	0.4	0.9	3.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Unspecified acute lower respiratory infection (J22)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*	*	0.4	1.3
2001.....	*	*	*	*	*	*	*	*	*	1.2
2000.....	*	*	*	*	*	*	*	*	*	1.9
1999.....	*	*	*	*	*	*	*	*	0.5	2.4

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	1.3
2001.....	*	*	*	*	*	*	*	*	*	1.3
2000.....	*	*	*	*	*	*	*	*	*	2.1
1999.....	*	*	*	*	*	*	*	*	*	2.8

White, both sexes

2002.....	*	*	*	*	*	*	*	*	*	1.4
2001.....	*	*	*	*	*	*	*	*	*	1.3
2000.....	*	*	*	*	*	*	*	*	*	2.0
1999.....	*	*	*	*	*	*	*	*	0.5	2.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of upper respiratory tract (J30-J39)

All races 3/, both sexes

2002.....	*	0.1	*	0.2	0.2	0.3	0.4	0.6	0.7	2.1
2001.....	*	0.1	0.1	*	0.2	0.3	0.4	0.6	1.3	2.4
2000.....	*	0.1	0.1	0.1	0.2	0.3	0.5	0.7	1.2	2.1
1999.....	*	*	*	*	0.3	0.4	0.4	0.9	1.1	2.4

All races 3/, male

2002.....	*	*	*	*	*	*	*	0.7	*	2.5
2001.....	*	*	*	*	*	*	*	0.7	1.9	3.4
2000.....	*	*	*	*	*	*	0.6	0.7	1.9	3.1
1999.....	*	*	*	*	*	0.5	0.5	1.1	1.3	3.5

All races 3/, female

2002.....	*	*	*	*	*	*	*	0.5	0.6	1.9
2001.....	*	*	*	*	*	*	*	0.6	1.0	1.9
2000.....	*	*	*	*	*	*	0.5	0.6	0.8	1.7
1999.....	*	*	*	*	*	*	*	0.7	1.0	2.0

White, both sexes

2002.....	*	*	*	0.2	*	0.2	0.4	0.6	0.6	2.2
2001.....	*	*	*	*	0.2	0.3	0.3	0.6	1.4	2.3
2000.....	*	*	0.1	*	0.2	0.3	0.5	0.6	1.3	2.2
1999.....	*	*	*	*	0.3	0.3	0.4	0.8	1.1	2.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)										
All races 3/, both sexes										
2002.....	2.9	5.8	12.0	28.2	60.8	119.0	211.5	330.1	465.7	637.6
2001.....	2.7	5.4	11.9	29.4	62.9	124.2	215.4	329.2	452.9	644.7
2000.....	2.8	5.8	11.8	29.6	62.5	124.1	218.2	327.8	473.4	648.6
1999.....	2.6	5.5	12.0	31.9	66.6	130.4	227.6	340.2	484.5	646.0
All races 3/, male										
2002.....	3.0	5.9	12.8	29.5	65.9	131.5	246.0	405.6	595.9	894.8
2001.....	2.5	5.8	12.3	31.4	67.0	137.1	252.9	406.7	583.5	916.9
2000.....	2.6	6.2	12.1	31.8	68.0	139.7	257.8	404.2	628.5	967.9
1999.....	2.4	5.7	12.1	34.3	74.6	151.0	278.3	432.4	648.3	996.7
All races 3/, female										
2002.....	2.8	5.6	11.2	26.9	56.1	108.3	183.8	276.8	386.5	526.0
2001.....	2.9	5.1	11.5	27.5	59.1	113.1	185.6	275.0	374.6	530.8
2000.....	2.9	5.5	11.5	27.5	57.5	110.7	187.0	274.6	381.9	518.5
1999.....	2.9	5.3	11.9	29.7	59.3	112.8	188.0	276.1	388.7	504.6
White, both sexes										
2002.....	2.5	5.5	11.7	28.9	63.8	126.8	223.3	346.9	486.5	666.2
2001.....	2.4	5.1	11.7	30.2	66.2	131.3	227.1	346.1	473.5	671.8
2000.....	2.3	5.3	11.4	30.1	65.5	130.5	229.6	343.1	495.5	675.6
1999.....	2.2	4.9	11.8	32.6	69.6	137.2	237.9	353.4	504.9	672.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)										
White, male										
2002.....	2.5	5.6	12.4	29.6	67.5	137.1	254.3	417.4	612.9	923.4
2001.....	2.2	5.6	11.9	31.8	69.1	142.0	260.8	418.9	600.0	943.6
2000.....	2.2	5.7	11.5	31.8	69.8	143.1	265.8	415.5	648.5	997.4
1999.....	2.0	5.0	11.6	34.5	76.3	155.7	285.6	441.3	664.4	1,029.4
White, female										
2002.....	2.5	5.4	11.0	28.3	60.4	117.6	198.0	296.4	408.9	554.5
2001.....	2.6	4.6	11.6	28.6	63.5	122.0	199.7	294.5	397.1	557.9
2000.....	2.4	4.9	11.3	28.4	61.5	119.4	200.5	292.1	404.4	544.8
1999.....	2.5	4.7	12.0	30.8	63.5	121.1	200.0	291.6	410.7	529.7
Black, both sexes										
2002.....	6.5	9.1	17.4	30.9	53.4	86.1	148.1	215.9	298.5	368.2
2001.....	5.1	9.1	16.6	31.6	52.2	94.3	149.6	206.9	276.5	375.8
2000.....	6.1	11.0	18.2	33.0	57.4	97.3	151.2	211.6	271.3	373.8
1999.....	6.1	11.2	17.2	36.1	57.6	100.0	163.3	242.4	295.3	376.2
Black, male										
2002.....	6.9	9.9	19.9	38.1	71.1	116.2	214.0	340.8	477.6	645.4
2001.....	5.1	8.9	18.8	37.1	66.3	127.1	222.2	329.9	468.7	671.7
2000.....	5.9	11.4	19.6	39.5	73.5	138.4	226.3	326.5	448.5	693.1
1999.....	6.2	11.9	19.7	43.2	79.3	139.4	243.5	387.0	521.4	701.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic lower respiratory diseases (J40-J47)

Black, female

2002.....	6.1	8.5	15.3	25.1	39.6	63.8	104.0	141.3	205.3	262.0
2001.....	5.1	9.2	14.7	27.1	41.1	70.0	101.3	133.4	178.7	266.1
2000.....	6.2	10.6	16.9	27.7	44.6	67.0	101.2	142.4	182.1	255.0
1999.....	6.1	10.7	15.0	30.2	40.6	70.8	110.0	155.3	183.2	254.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
All races 3/, both sexes										
2002.....	2.9	5.7	11.9	28.0	60.4	118.4	210.3	327.6	461.0	631.8
2001.....	2.7	5.4	11.8	29.3	62.5	123.4	214.2	326.9	449.0	637.7
2000.....	2.7	5.8	11.8	29.3	62.2	123.4	216.9	325.4	468.8	642.4
1999.....	2.6	5.4	11.9	31.7	66.2	129.6	226.5	337.6	479.9	639.8
All races 3/, male										
2002.....	2.9	5.9	12.7	29.4	65.6	131.1	245.1	404.0	592.8	890.1
2001.....	2.5	5.7	12.3	31.3	66.6	136.5	252.0	405.0	581.3	912.1
2000.....	2.6	6.2	12.1	31.6	67.7	139.3	256.8	402.6	625.4	963.6
1999.....	2.4	5.6	12.1	34.2	74.3	150.6	277.3	429.9	644.2	990.1
All races 3/, female										
2002.....	2.8	5.6	11.1	26.7	55.6	107.4	182.4	273.7	380.9	519.7
2001.....	2.8	5.1	11.4	27.4	58.7	112.2	184.2	272.2	369.7	522.9
2000.....	2.8	5.4	11.4	27.2	57.2	109.9	185.6	271.7	376.5	511.6
1999.....	2.8	5.2	11.7	29.4	58.9	111.6	186.8	273.5	383.7	498.6
White, both sexes										
2002.....	2.5	5.5	11.6	28.8	63.3	126.2	222.0	344.2	481.6	660.2
2001.....	2.4	5.1	11.6	30.1	65.8	130.6	225.8	343.6	469.4	664.3
2000.....	2.3	5.2	11.4	29.8	65.2	129.8	228.4	340.6	490.6	669.0
1999.....	2.2	4.8	11.7	32.3	69.2	136.4	236.8	350.8	500.1	666.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
White, male										
2002.....	2.5	5.5	12.3	29.5	67.1	136.8	253.5	415.7	609.7	918.7
2001.....	2.2	5.5	11.8	31.7	68.6	141.6	260.0	417.3	597.9	938.8
2000.....	2.2	5.6	11.5	31.6	69.6	142.8	264.8	414.0	645.1	992.8
1999.....	2.0	5.0	11.6	34.3	76.0	155.3	284.6	438.9	660.3	1,022.5
White, female										
2002.....	2.4	5.4	11.0	28.1	59.7	116.8	196.4	293.1	403.0	547.9
2001.....	2.6	4.6	11.5	28.5	63.2	121.0	198.2	291.4	391.8	549.4
2000.....	2.4	4.9	11.2	28.1	61.0	118.4	199.2	289.0	398.7	537.5
1999.....	2.5	4.7	11.8	30.4	63.0	119.9	198.8	288.9	405.5	523.1
Black, both sexes										
2002.....	6.5	9.0	17.2	30.9	53.3	85.6	147.6	215.4	297.0	367.0
2001.....	5.1	9.0	16.6	31.5	51.9	94.0	149.2	206.0	275.2	374.6
2000.....	6.1	10.9	18.1	32.6	57.1	96.6	150.4	211.1	270.5	373.2
1999.....	6.1	11.1	17.1	35.8	57.4	99.7	162.5	241.0	293.6	375.3
Black, male										
2002.....	6.9	9.8	19.7	38.0	70.9	115.7	212.6	339.9	476.1	645.4
2001.....	5.1	8.8	18.8	36.9	66.1	126.5	221.5	328.5	465.5	669.4
2000.....	5.9	11.3	19.6	39.5	73.1	137.1	225.0	325.1	448.5	693.1
1999.....	6.1	11.8	19.6	42.9	78.9	139.4	242.9	385.1	520.6	700.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
Black, female										
2002.....	6.1	8.3	15.1	25.1	39.5	63.3	104.0	141.1	203.7	260.4
2001.....	5.1	9.2	14.7	27.1	40.8	70.0	101.1	132.9	178.2	265.2
2000.....	6.2	10.6	16.8	27.0	44.5	66.6	100.8	142.4	180.9	254.1
1999.....	6.1	10.6	14.9	30.0	40.4	70.2	109.1	154.2	181.1	253.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Bronchitis, not specified as acute or chronic (J40)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	0.4	0.7	1.4	5.2
2001.....	*	*	*	*	*	0.2	0.4	1.0	1.3	5.2
2000.....	*	*	*	*	0.2	0.3	0.4	1.1	2.0	6.8
1999.....	*	*	*	*	0.3	0.3	0.7	1.0	2.2	6.2

All races 3/, male

2002.....	*	*	*	*	*	*	*	0.8	1.2	4.7
2001.....	*	*	*	*	*	*	*	1.0	1.2	5.1
2000.....	*	*	*	*	*	*	*	1.3	3.0	6.4
1999.....	*	*	*	*	*	*	0.7	1.2	2.3	5.9

All races 3/, female

2002.....	*	*	*	*	*	*	0.5	0.7	1.5	5.4
2001.....	*	*	*	*	*	*	*	0.9	1.4	5.3
2000.....	*	*	*	*	*	*	0.5	1.0	1.4	7.0
1999.....	*	*	*	*	*	*	0.6	0.8	2.1	6.3

White, both sexes

2002.....	*	*	*	*	*	*	0.5	0.8	1.5	5.5
2001.....	*	*	*	*	*	*	0.4	1.0	1.4	5.5
2000.....	*	*	*	*	0.2	0.3	0.4	1.2	2.1	7.3
1999.....	*	*	*	*	0.3	0.3	0.7	0.9	2.2	6.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic bronchitis (J41-J42)										
All races 3/, both sexes										
2002.....	*	*	*	*	0.2	0.4	0.5	1.0	1.5	2.9
2001.....	*	*	*	*	0.2	0.3	0.4	0.8	1.2	3.9
2000.....	*	*	*	*	0.2	0.4	0.6	1.0	1.6	5.0
1999.....	*	*	*	0.2	0.3	0.3	0.6	0.9	1.9	4.5
All races 3/, male										
2002.....	*	*	*	*	*	0.5	0.7	1.5	1.6	4.0
2001.....	*	*	*	*	*	0.5	0.6	1.1	1.2	4.1
2000.....	*	*	*	*	*	0.6	0.6	1.4	1.9	6.4
1999.....	*	*	*	*	*	0.5	0.7	1.3	2.3	5.9
All races 3/, female										
2002.....	*	*	*	*	*	*	*	0.7	1.4	2.4
2001.....	*	*	*	*	*	*	*	0.5	1.2	3.8
2000.....	*	*	*	*	*	*	0.6	0.7	1.4	4.4
1999.....	*	*	*	*	0.4	*	0.4	0.7	1.7	4.0
White, both sexes										
2002.....	*	*	*	*	*	0.4	0.6	1.1	1.5	3.1
2001.....	*	*	*	*	0.2	0.4	0.4	0.8	1.3	4.1
2000.....	*	*	*	*	0.2	0.4	0.6	1.0	1.6	5.3
1999.....	*	*	*	0.2	0.3	0.4	0.6	1.0	1.9	4.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
All races 3/, both sexes										
2002.....	0.3	0.7	1.6	4.3	9.1	17.4	31.3	43.2	54.7	59.3
2001.....	0.2	0.7	1.7	5.0	10.1	19.8	33.2	46.9	55.5	62.1
2000.....	0.3	0.8	1.8	5.0	10.4	20.7	35.5	47.5	60.4	63.3
1999.....	0.4	0.7	2.0	5.5	12.1	23.5	38.4	51.3	65.3	62.1
All races 3/, male										
2002.....	0.5	0.9	1.9	4.6	10.7	19.7	37.8	56.1	74.9	90.0
2001.....	0.3	0.9	2.0	5.8	10.8	22.4	40.3	59.2	74.9	101.2
2000.....	0.3	1.0	2.1	5.6	11.2	24.5	42.6	60.4	83.6	104.2
1999.....	0.4	0.9	2.2	6.2	13.6	28.0	48.9	67.3	91.5	105.4
All races 3/, female										
2002.....	0.2	0.6	1.3	4.0	7.7	15.4	26.1	34.0	42.4	46.0
2001.....	*	0.5	1.4	4.2	9.5	17.6	27.5	38.3	43.8	45.8
2000.....	0.2	0.6	1.5	4.5	9.7	17.5	29.9	38.5	46.8	46.7
1999.....	0.3	0.5	1.7	4.9	10.8	19.8	30.2	40.2	49.9	44.7
White, both sexes										
2002.....	0.3	0.7	1.7	4.6	9.8	18.6	33.3	45.8	57.4	62.3
2001.....	0.2	0.7	1.7	5.1	10.9	21.5	35.2	49.8	58.7	65.1
2000.....	0.2	0.7	1.8	5.3	11.2	22.0	37.9	50.2	63.6	65.8
1999.....	0.3	0.6	2.0	5.8	13.0	25.4	40.9	53.8	68.8	64.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
White, male										
2002.....	0.4	0.9	1.9	4.8	11.1	20.2	39.5	58.2	77.4	93.0
2001.....	0.3	1.0	1.9	5.8	11.3	23.9	41.6	61.7	78.0	105.1
2000.....	0.2	0.9	2.0	5.8	11.8	25.0	44.6	63.1	87.0	105.5
1999.....	0.3	0.8	2.1	6.4	14.2	29.4	50.6	69.6	94.4	109.2
White, female										
2002.....	*	0.6	1.5	4.4	8.6	17.3	28.3	37.0	45.1	48.9
2001.....	*	0.5	1.4	4.5	10.6	19.4	30.0	41.3	47.0	48.4
2000.....	0.2	0.6	1.5	4.8	10.7	19.3	32.5	41.1	49.8	49.7
1999.....	0.3	0.5	1.9	5.3	11.9	21.9	33.1	42.6	53.7	46.9
Black, both sexes										
2002.....	0.8	1.0	1.5	3.3	5.9	11.9	20.9	23.5	34.8	32.1
2001.....	*	*	2.3	4.9	6.4	11.6	20.1	21.9	26.3	35.3
2000.....	*	1.6	2.5	4.7	7.7	14.9	19.5	24.5	30.8	37.2
1999.....	0.8	1.4	2.4	4.6	8.0	14.0	20.8	31.4	31.2	37.6
Black, male										
2002.....	*	*	2.4	4.5	9.3	20.4	30.7	41.3	61.1	64.3
2001.....	*	*	2.9	6.9	9.6	15.8	31.6	34.4	49.0	68.1
2000.....	*	2.4	3.5	5.9	10.8	25.2	30.1	35.0	53.1	87.6
1999.....	*	2.0	4.1	6.4	11.9	22.8	37.0	48.9	65.4	72.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
Black, female										
2002.....	*	*	*	*	3.3	5.6	14.3	12.9	21.1	19.7
2001.....	*	*	*	3.3	4.0	8.4	12.3	14.4	14.7	23.2
2000.....	*	*	*	3.6	5.2	7.3	12.4	18.2	19.5	18.5
1999.....	*	*	*	3.0	5.0	7.5	10.1	20.9	14.3	24.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic obstructive pulmonary disease (J44)										
All races 3/, both sexes										
2002.....	1.4	3.5	8.6	21.6	48.8	97.8	174.2	277.3	395.7	549.7
2001.....	1.3	3.3	8.3	22.3	49.4	100.0	176.2	273.1	382.9	550.3
2000.....	1.2	3.3	8.2	22.1	48.8	98.6	175.8	270.2	396.1	550.6
1999.....	1.1	3.1	8.0	23.3	51.2	101.7	182.1	278.2	400.8	549.6
All races 3/, male										
2002.....	1.5	3.7	9.4	23.2	52.9	108.7	203.4	341.3	508.8	780.4
2001.....	1.3	3.7	8.9	24.1	53.5	111.0	207.7	339.9	498.3	789.6
2000.....	1.2	3.8	8.7	24.2	54.2	111.1	209.8	335.5	530.2	833.7
1999.....	1.2	3.6	8.5	25.7	58.4	119.2	222.9	355.4	540.9	860.0
All races 3/, female										
2002.....	1.2	3.2	7.8	20.2	45.1	88.4	150.9	232.1	326.9	449.5
2001.....	1.4	2.8	7.7	20.6	45.7	90.5	151.2	226.3	313.7	450.2
2000.....	1.2	2.9	7.7	20.2	43.9	87.8	149.0	224.6	317.1	435.3
1999.....	1.1	2.5	7.6	21.1	44.7	86.7	150.2	224.6	318.9	424.5
White, both sexes										
2002.....	1.4	3.6	8.8	22.7	51.4	104.8	184.5	291.4	413.7	574.7
2001.....	1.3	3.3	8.5	23.3	52.7	105.9	186.4	287.4	400.3	573.9
2000.....	1.1	3.3	8.3	23.0	51.5	104.3	185.5	283.0	414.8	574.7
1999.....	1.1	3.0	8.3	24.2	53.8	107.3	190.7	289.4	417.8	573.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic obstructive pulmonary disease (J44)										
White, male										
2002.....	1.5	3.8	9.4	23.6	54.4	114.4	210.6	351.3	523.6	805.8
2001.....	1.3	3.8	8.9	24.7	55.8	114.9	214.9	350.1	512.4	812.9
2000.....	1.2	3.7	8.5	24.7	56.0	114.7	216.3	344.3	546.6	862.1
1999.....	1.1	3.5	8.6	26.1	60.0	123.0	229.5	362.6	554.3	890.0
White, female										
2002.....	1.3	3.5	8.1	21.9	48.7	96.3	163.2	248.5	346.4	474.3
2001.....	1.4	2.8	8.1	22.0	49.8	98.0	163.3	243.0	332.6	473.8
2000.....	1.1	3.0	8.1	21.4	47.3	95.2	160.8	239.8	336.4	457.9
1999.....	1.1	2.5	8.1	22.4	48.2	93.5	159.9	237.9	337.2	446.6
Black, both sexes										
2002.....	1.7	3.8	10.0	20.3	41.6	67.2	116.4	183.8	250.9	318.9
2001.....	1.6	4.0	9.3	20.6	37.3	75.5	118.7	172.5	237.0	319.7
2000.....	1.8	3.8	10.1	21.3	41.9	72.0	119.9	174.1	227.0	310.0
1999.....	1.9	4.7	8.4	23.6	42.8	76.8	129.1	197.6	247.0	312.8
Black, male										
2002.....	2.0	4.7	12.5	27.4	56.0	88.3	172.9	290.7	405.8	565.0
2001.....	1.6	4.3	11.3	26.4	50.2	105.2	178.7	283.9	405.3	580.9
2000.....	1.7	4.9	12.5	27.3	55.7	102.8	186.5	283.5	383.6	576.6
1999.....	2.4	5.9	10.8	30.9	60.9	109.6	191.6	323.9	440.3	605.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic obstructive pulmonary disease (J44)										
Black, female										
2002.....	1.4	2.9	7.8	14.4	30.4	51.6	78.4	119.9	170.3	224.7
2001.....	1.5	3.7	7.7	15.8	27.2	53.5	78.7	105.9	151.3	222.7
2000.....	1.9	2.8	7.9	16.3	31.1	49.3	75.7	108.2	148.2	210.8
1999.....	1.5	3.7	6.4	17.7	28.6	52.5	87.5	121.6	151.2	203.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)										
All races 3/, both sexes										
2002.....	1.2	1.4	1.6	1.9	2.1	2.7	3.8	5.4	7.7	14.7
2001.....	1.1	1.3	1.8	1.9	2.6	3.0	3.9	5.2	8.1	16.1
2000.....	1.2	1.6	1.7	2.0	2.5	3.4	4.6	5.7	8.7	16.7
1999.....	1.1	1.6	1.7	2.6	2.2	3.7	4.7	6.2	9.6	17.3
All races 3/, male										
2002.....	0.9	1.1	1.3	1.5	1.7	2.0	2.8	4.3	6.2	10.9
2001.....	0.9	1.0	1.3	1.2	1.9	2.4	3.0	3.8	5.6	12.1
2000.....	1.0	1.3	1.3	1.6	1.8	2.7	3.4	3.9	6.7	12.9
1999.....	0.8	1.0	1.2	1.9	1.7	2.6	3.9	4.8	7.2	12.7
All races 3/, female										
2002.....	1.4	1.8	1.9	2.3	2.5	3.3	4.5	6.2	8.7	16.3
2001.....	1.3	1.7	2.2	2.5	3.3	3.6	4.7	6.1	9.6	17.8
2000.....	1.4	1.9	2.1	2.4	3.1	4.0	5.5	6.9	9.8	18.2
1999.....	1.3	2.2	2.2	3.3	2.7	4.6	5.4	7.2	11.1	19.2
White, both sexes										
2002.....	0.8	1.1	1.1	1.4	1.8	2.2	3.2	5.1	7.5	14.7
2001.....	0.8	1.0	1.4	1.4	1.9	2.6	3.4	4.7	7.8	15.7
2000.....	0.9	1.1	1.2	1.4	2.0	2.8	4.0	5.2	8.4	15.9
1999.....	0.8	1.2	1.3	2.0	1.8	3.1	4.0	5.8	9.4	16.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)										
White, male										
2002.....	0.6	0.8	0.9	1.0	1.4	1.5	2.3	3.8	5.8	10.7
2001.....	0.6	0.7	0.9	1.0	1.2	2.0	2.4	3.3	5.1	11.3
2000.....	0.7	1.0	1.0	1.0	1.4	2.0	3.0	3.7	6.3	11.4
1999.....	0.5	0.7	0.9	1.5	1.2	2.0	3.0	4.2	7.0	10.8
White, female										
2002.....	1.0	1.3	1.3	1.7	2.2	2.8	3.9	6.1	8.6	16.4
2001.....	1.0	1.2	1.8	1.9	2.5	3.2	4.2	5.7	9.4	17.6
2000.....	1.0	1.2	1.5	1.8	2.6	3.4	4.8	6.3	9.7	17.7
1999.....	1.1	1.7	1.7	2.6	2.2	4.0	4.8	6.9	10.8	18.9
Black, both sexes										
2002.....	4.0	4.1	5.7	7.1	5.6	6.1	10.1	7.4	8.9	11.9
2001.....	3.1	4.3	4.9	5.9	7.9	6.3	9.4	10.2	11.1	16.3
2000.....	3.6	5.4	5.3	6.6	7.2	8.8	9.7	10.7	9.9	22.2
1999.....	3.3	4.8	5.8	7.3	6.2	8.4	11.1	9.8	11.3	20.4
Black, male										
2002.....	3.7	3.4	4.6	6.0	5.3	6.2	8.7	*	*	*
2001.....	2.9	3.8	4.5	3.6	5.9	*	9.8	*	*	*
2000.....	3.1	4.0	3.5	6.3	6.0	8.6	7.0	*	*	25.4
1999.....	2.9	3.6	4.2	5.1	5.7	6.8	12.2	9.5	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)										
Black, female										
2002.....	4.2	4.7	6.6	8.1	5.7	5.9	11.0	8.0	9.5	12.3
2001.....	3.3	4.8	5.3	7.8	9.4	7.5	9.2	11.4	11.9	16.8
2000.....	4.0	6.6	6.9	6.9	8.2	9.0	11.5	14.2	10.2	21.0
1999.....	3.6	5.9	7.1	9.1	6.5	9.6	10.4	10.0	13.8	21.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchiectasis (J47)										
All races 3/, both sexes										
2002.....	*	*	*	0.1	0.4	0.6	1.2	2.5	4.6	5.8
2001.....	*	*	*	0.2	0.4	0.7	1.2	2.3	3.9	7.0
2000.....	*	*	*	0.3	0.3	0.7	1.2	2.4	4.6	6.1
1999.....	*	*	*	0.2	0.4	0.8	1.1	2.6	4.7	6.2
All races 3/, male										
2002.....	*	*	*	*	*	*	0.9	1.6	3.1	4.7
2001.....	*	*	*	*	0.4	0.6	0.8	1.6	2.2	4.8
2000.....	*	*	*	*	*	0.5	1.0	1.6	3.2	4.3
1999.....	*	*	*	*	*	*	1.0	2.5	4.1	6.6
All races 3/, female										
2002.....	*	*	*	*	0.5	0.9	1.4	3.1	5.5	6.3
2001.....	*	*	*	*	0.4	0.9	1.5	2.8	4.9	7.9
2000.....	*	*	*	0.3	0.4	0.9	1.4	3.0	5.5	6.9
1999.....	*	*	*	0.3	0.4	1.1	1.2	2.6	5.1	6.0
White, both sexes										
2002.....	*	*	*	0.2	0.5	0.6	1.3	2.7	4.9	6.1
2001.....	*	*	*	*	0.4	0.8	1.2	2.4	4.1	7.4
2000.....	*	*	*	0.2	0.3	0.7	1.2	2.5	4.9	6.6
1999.....	*	*	*	0.2	0.4	0.8	1.1	2.6	4.8	6.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lung diseases due to external agents (J60-J70)										
All races 3/, both sexes										
2002.....	0.6	0.7	1.2	2.0	3.9	7.5	15.2	34.9	71.7	192.9
2001.....	0.5	0.7	1.3	2.1	4.0	7.3	16.2	34.0	73.3	195.9
2000.....	0.5	0.8	1.2	2.1	3.5	7.7	16.5	34.0	70.6	195.3
1999.....	0.5	0.7	1.0	2.1	3.5	7.3	15.3	31.9	66.1	181.8
All races 3/, male										
2002.....	0.7	0.9	1.6	2.6	5.1	10.5	22.0	51.6	105.9	290.1
2001.....	0.7	0.8	1.8	2.8	5.0	10.4	23.7	49.1	111.6	291.3
2000.....	0.7	1.1	1.6	2.7	4.6	11.2	25.2	50.6	107.8	301.5
1999.....	0.6	0.9	1.2	2.7	5.1	10.7	22.9	50.0	103.9	279.1
All races 3/, female										
2002.....	0.4	0.5	0.8	1.5	2.9	5.0	9.9	23.2	50.9	150.7
2001.....	0.4	0.6	0.9	1.4	3.0	4.6	10.3	23.5	50.3	156.0
2000.....	0.4	0.6	0.9	1.6	2.5	4.8	9.7	22.4	48.6	152.1
1999.....	0.4	0.5	0.9	1.6	2.1	4.3	9.3	19.3	44.0	142.6
White, both sexes										
2002.....	0.5	0.7	1.1	2.0	3.8	7.3	15.1	34.7	72.8	196.8
2001.....	0.5	0.7	1.2	2.1	3.8	7.2	15.8	34.1	73.7	197.6
2000.....	0.6	0.7	1.2	2.1	3.3	7.5	15.9	33.8	70.8	199.2
1999.....	0.4	0.6	0.9	2.0	3.3	7.0	14.7	31.5	66.1	184.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lung diseases due to external agents (J60-J70)										
White, male										
2002.....	0.7	0.9	1.5	2.6	4.9	10.1	21.6	51.1	107.8	297.5
2001.....	0.7	0.8	1.6	2.8	4.7	10.4	23.2	49.4	112.8	297.5
2000.....	0.7	0.9	1.4	2.7	4.4	10.6	23.9	50.0	109.3	311.0
1999.....	0.5	0.8	1.0	2.5	4.7	10.2	21.9	49.2	103.7	285.0
White, female										
2002.....	0.4	0.5	0.8	1.5	2.7	4.7	9.8	23.0	51.3	153.0
2001.....	0.4	0.5	0.9	1.4	3.0	4.3	9.8	23.3	50.0	155.8
2000.....	0.4	0.5	0.9	1.5	2.3	4.8	9.4	22.5	47.9	153.8
1999.....	0.4	0.4	0.9	1.5	2.1	4.2	9.0	19.0	43.9	144.0
Black, both sexes										
2002.....	1.0	0.9	2.1	2.6	5.9	12.0	19.8	44.0	70.6	169.3
2001.....	0.8	1.1	2.5	2.6	6.4	9.8	23.4	39.9	79.6	189.2
2000.....	*	1.7	2.2	3.0	5.9	11.8	26.8	41.6	74.3	163.6
1999.....	1.0	1.2	2.0	4.0	5.9	10.9	24.8	42.7	73.4	172.2
Black, male										
2002.....	*	*	3.2	3.7	7.3	17.5	30.7	67.5	99.3	247.7
2001.....	*	*	3.5	3.6	9.2	13.5	33.7	59.9	118.8	257.6
2000.....	*	2.4	3.3	3.3	7.7	18.7	45.1	69.0	103.7	223.7
1999.....	*	*	2.7	5.3	10.0	17.4	40.0	69.8	116.0	246.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lung diseases due to external agents (J60-J70)										
Black, female										
2002.....	*	*	*	*	4.8	8.0	12.5	29.9	55.7	139.2
2001.....	*	*	*	*	4.3	7.1	16.6	28.0	59.7	163.8
2000.....	*	*	*	2.7	4.6	6.7	14.7	25.1	59.4	141.2
1999.....	*	*	*	2.9	*	6.2	14.6	26.3	52.3	144.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumoconioses and chemical effects (J60-J66,J68)

All races 3/, both sexes

2002.....	*	*	*	0.2	0.4	0.8	1.6	3.4	4.2	6.8
2001.....	*	*	*	0.2	0.3	1.0	2.0	3.4	5.1	6.2
2000.....	*	*	*	0.2	0.3	0.9	2.2	3.4	4.8	7.5
1999.....	*	*	*	*	0.4	1.1	1.9	3.4	5.2	6.1

All races 3/, male

2002.....	*	*	*	0.3	0.8	1.7	3.5	8.0	10.5	21.6
2001.....	*	*	*	0.4	0.7	2.0	4.4	7.8	13.2	20.2
2000.....	*	*	*	*	0.7	1.8	4.8	7.9	12.3	24.9
1999.....	*	*	*	*	0.7	2.3	4.4	8.1	13.5	20.4

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	0.2	0.4	0.9	1.8	3.6	4.4	7.2
2001.....	*	*	*	0.2	0.3	1.1	2.1	3.7	5.4	6.5
2000.....	*	*	*	0.2	0.4	1.0	2.4	3.5	5.0	7.9
1999.....	*	*	*	*	0.3	1.1	2.0	3.6	5.5	6.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)										
All races 3/, both sexes										
2002.....	0.5	0.7	1.1	1.8	3.4	6.5	13.4	31.3	67.3	186.0
2001.....	0.5	0.7	1.3	1.8	3.6	6.2	14.1	30.4	68.0	189.4
2000.....	0.5	0.8	1.2	1.9	3.2	6.7	14.2	30.4	65.7	187.6
1999.....	0.5	0.7	0.9	2.0	3.1	6.1	13.2	28.2	60.7	175.6
All races 3/, male										
2002.....	0.7	0.9	1.5	2.2	4.2	8.6	18.2	43.4	95.1	268.5
2001.....	0.7	0.8	1.7	2.3	4.2	8.2	19.1	41.0	98.1	270.3
2000.....	0.6	1.0	1.5	2.4	3.9	9.2	20.2	42.4	95.2	276.0
1999.....	0.6	0.9	1.0	2.4	4.2	8.3	18.4	41.6	90.1	258.4
All races 3/, female										
2002.....	0.4	0.5	0.8	1.4	2.8	4.8	9.6	22.8	50.4	150.2
2001.....	0.4	0.6	0.9	1.4	3.0	4.4	10.2	23.0	49.9	155.6
2000.....	0.4	0.6	0.9	1.5	2.5	4.6	9.4	22.1	48.2	151.6
1999.....	0.4	0.5	0.8	1.6	2.0	4.2	9.2	18.8	43.5	142.2
White, both sexes										
2002.....	0.5	0.7	1.0	1.8	3.3	6.1	13.1	30.9	68.1	189.5
2001.....	0.5	0.7	1.2	1.8	3.5	6.0	13.6	30.2	68.1	190.8
2000.....	0.5	0.7	1.1	1.9	2.9	6.4	13.4	30.1	65.7	191.0
1999.....	0.4	0.6	0.8	1.8	2.9	5.8	12.5	27.6	60.3	178.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)										
White, male										
2002.....	0.6	0.8	1.3	2.1	4.1	8.0	17.6	42.5	96.5	274.6
2001.....	0.7	0.8	1.5	2.3	4.0	8.0	18.4	40.6	98.7	275.4
2000.....	0.7	0.9	1.3	2.3	3.6	8.5	18.7	41.4	96.2	284.0
1999.....	0.5	0.8	0.9	2.2	3.9	7.7	17.1	40.4	88.9	263.6
White, female										
2002.....	0.4	0.5	0.7	1.5	2.6	4.5	9.5	22.6	50.7	152.5
2001.....	0.4	0.5	0.8	1.3	3.0	4.2	9.7	22.8	49.6	155.3
2000.....	0.4	0.5	0.9	1.4	2.3	4.6	9.1	22.2	47.5	153.2
1999.....	0.4	0.4	0.8	1.5	2.0	4.1	8.8	18.6	43.4	143.6
Black, both sexes										
2002.....	0.9	0.8	2.1	2.4	5.5	11.7	18.8	41.9	67.8	165.1
2001.....	0.7	1.0	2.4	2.4	6.0	9.3	22.3	38.6	76.1	185.5
2000.....	*	1.7	2.2	2.9	5.8	11.3	25.1	38.9	71.4	159.8
1999.....	1.0	1.2	1.8	3.9	5.0	10.1	23.7	40.5	71.1	167.1
Black, male										
2002.....	*	*	3.1	3.7	6.3	16.7	28.4	62.5	91.7	233.7
2001.....	*	*	3.4	3.1	8.1	12.5	31.0	56.2	108.3	243.9
2000.....	*	2.4	3.1	3.3	7.2	17.4	41.1	62.4	96.1	209.9
1999.....	*	*	*	5.1	8.1	15.6	37.3	65.1	109.9	228.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)										
Black, female										
2002.....	*	*	*	*	4.8	8.0	12.3	29.6	55.3	138.8
2001.....	*	*	*	*	4.3	6.9	16.6	28.0	59.7	163.8
2000.....	*	*	*	*	4.6	6.7	14.4	24.8	59.0	141.2
1999.....	*	*	*	2.9	*	6.0	14.6	25.7	51.9	144.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to food and vomit (J69.0)										
All races 3/, both sexes										
2002.....	0.5	0.7	1.1	1.8	3.4	6.5	13.4	31.3	67.3	186.0
2001.....	0.5	0.7	1.3	1.8	3.6	6.2	14.1	30.4	67.9	189.4
2000.....	0.5	0.8	1.2	1.9	3.1	6.7	14.2	30.4	65.7	187.6
1999.....	0.5	0.7	0.9	2.0	3.0	6.1	13.2	28.1	60.6	175.6
All races 3/, male										
2002.....	0.7	0.9	1.5	2.2	4.2	8.5	18.2	43.4	95.1	268.5
2001.....	0.7	0.8	1.7	2.3	4.2	8.2	19.1	41.0	98.0	270.2
2000.....	0.6	1.0	1.5	2.3	3.9	9.2	20.2	42.4	95.2	276.0
1999.....	0.6	0.9	1.0	2.4	4.2	8.3	18.4	41.6	90.1	258.4
All races 3/, female										
2002.....	0.4	0.5	0.8	1.4	2.8	4.8	9.6	22.8	50.3	150.2
2001.....	0.4	0.6	0.9	1.4	3.0	4.4	10.2	23.0	49.9	155.5
2000.....	0.4	0.6	0.9	1.5	2.5	4.6	9.4	22.1	48.2	151.6
1999.....	0.4	0.5	0.8	1.6	2.0	4.2	9.2	18.8	43.4	142.2
White, both sexes										
2002.....	0.5	0.7	1.0	1.8	3.3	6.1	13.1	30.9	68.1	189.5
2001.....	0.5	0.7	1.2	1.8	3.5	5.9	13.6	30.2	68.1	190.7
2000.....	0.5	0.7	1.1	1.8	2.9	6.4	13.4	30.1	65.7	191.0
1999.....	0.4	0.6	0.8	1.8	2.9	5.7	12.5	27.6	60.2	178.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to food and vomit (J69.0)										
White, male										
2002.....	0.6	0.8	1.3	2.1	4.1	8.0	17.6	42.5	96.5	274.6
2001.....	0.7	0.8	1.5	2.3	4.0	8.0	18.4	40.6	98.6	275.3
2000.....	0.7	0.9	1.3	2.3	3.6	8.5	18.7	41.4	96.2	284.0
1999.....	0.5	0.8	0.9	2.2	3.9	7.7	17.1	40.4	88.9	263.6
White, female										
2002.....	0.4	0.5	0.7	1.5	2.6	4.4	9.5	22.5	50.7	152.5
2001.....	0.4	0.5	0.8	1.3	3.0	4.2	9.7	22.8	49.6	155.3
2000.....	0.4	0.5	0.9	1.4	2.3	4.6	9.1	22.2	47.5	153.2
1999.....	0.4	0.4	0.8	1.5	2.0	4.1	8.8	18.6	43.3	143.6
Black, both sexes										
2002.....	0.9	0.8	2.1	2.4	5.5	11.7	18.8	41.9	67.5	165.1
2001.....	0.7	1.0	2.4	2.4	6.0	9.3	22.3	38.6	76.1	185.5
2000.....	*	1.7	2.2	2.9	5.8	11.3	25.1	38.9	71.4	159.8
1999.....	1.0	1.2	1.8	3.9	5.0	9.9	23.7	40.3	71.1	167.1
Black, male										
2002.....	*	*	3.1	3.7	6.3	16.7	28.4	62.5	91.7	233.7
2001.....	*	*	3.4	3.1	8.1	12.5	31.0	56.2	108.3	243.9
2000.....	*	2.4	3.1	3.3	7.2	17.4	41.1	62.4	96.1	209.9
1999.....	*	*	*	5.1	8.1	15.3	37.3	65.1	109.9	228.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to food and vomit (J69.0)										
Black, female										
2002.....	*	*	*	*	4.8	8.0	12.3	29.6	54.9	138.8
2001.....	*	*	*	*	4.3	6.9	16.6	28.0	59.7	163.8
2000.....	*	*	*	*	4.6	6.7	14.4	24.8	59.0	141.2
1999.....	*	*	*	2.9	*	6.0	14.6	25.5	51.9	144.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Suppurative and necrotic conditions of lower respiratory tract (J85-J86)

All races 3/, both sexes

2002.....	0.1	0.2	0.3	0.3	0.5	0.6	1.0	1.3	1.9	2.3
2001.....	0.1	0.2	0.3	0.4	0.5	0.7	0.9	1.5	1.8	3.0
2000.....	0.2	0.2	0.3	0.3	0.5	0.6	1.1	1.5	1.6	2.9
1999.....	0.2	0.2	0.3	0.4	0.5	1.0	1.0	1.3	1.7	2.8

All races 3/, male

2002.....	0.2	0.4	0.4	0.4	0.8	0.9	1.3	1.9	3.0	3.4
2001.....	0.2	0.2	0.5	0.5	0.7	0.9	1.3	2.2	2.5	4.7
2000.....	0.2	0.3	0.4	0.5	0.6	1.0	1.4	2.2	2.1	5.8
1999.....	0.3	0.3	0.5	0.5	0.7	1.6	1.5	1.7	2.8	4.0

All races 3/, female

2002.....	*	*	*	*	*	0.4	0.6	0.9	1.1	1.8
2001.....	*	*	*	0.3	*	0.5	0.7	1.1	1.4	2.3
2000.....	*	*	*	*	0.4	*	0.8	0.9	1.4	1.7
1999.....	*	*	*	*	*	0.4	0.6	1.1	1.0	2.3

White, both sexes

2002.....	0.1	0.2	0.2	0.3	0.4	0.6	0.9	1.1	1.8	2.2
2001.....	0.1	0.2	0.3	0.4	0.5	0.7	0.9	1.5	1.8	3.0
2000.....	0.1	0.1	0.3	0.3	0.4	0.6	1.0	1.5	1.6	2.7
1999.....	0.2	0.2	0.3	0.3	0.5	0.9	1.0	1.3	1.7	2.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pleural effusion and plaque (J90,J92)										
All races 3/, both sexes										
2002.....	*	*	*	*	0.2	0.5	0.9	1.6	3.7	9.9
2001.....	*	*	*	*	0.3	0.4	0.9	1.6	3.8	10.0
2000.....	*	*	*	*	0.2	0.5	0.8	1.6	3.6	8.9
1999.....	*	*	*	0.2	0.2	0.5	0.8	1.7	3.4	9.0
All races 3/, male										
2002.....	*	*	*	*	*	*	0.9	1.8	4.7	11.0
2001.....	*	*	*	*	*	0.5	1.2	1.6	4.8	12.2
2000.....	*	*	*	*	*	0.5	1.0	2.0	4.2	10.8
1999.....	*	*	*	*	*	0.6	1.1	2.2	4.2	11.5
All races 3/, female										
2002.....	*	*	*	*	*	0.6	0.8	1.4	3.1	9.4
2001.....	*	*	*	*	*	*	0.7	1.7	3.3	9.1
2000.....	*	*	*	*	*	0.4	0.6	1.3	3.2	8.1
1999.....	*	*	*	*	*	*	0.7	1.3	2.9	8.0
White, both sexes										
2002.....	*	*	*	*	0.2	0.5	0.8	1.6	3.8	10.2
2001.....	*	*	*	*	0.3	0.4	0.9	1.6	3.9	10.0
2000.....	*	*	*	*	*	0.4	0.8	1.6	3.5	9.0
1999.....	*	*	*	*	0.3	0.5	0.8	1.7	3.5	9.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of respiratory system (J80-J84,J93-J98)										
All races 3/, both sexes										
2002.....	1.2	2.0	3.5	6.3	10.9	19.0	35.0	53.0	77.0	118.1
2001.....	1.3	2.2	3.8	6.6	11.5	21.2	35.6	57.6	82.1	125.7
2000.....	1.4	2.2	4.0	7.1	11.7	21.3	34.2	56.4	82.3	125.8
1999.....	1.3	2.2	3.5	6.1	11.5	20.5	34.2	55.1	80.6	131.6
All races 3/, male										
2002.....	1.3	2.0	3.6	7.0	12.5	22.7	44.7	65.9	95.2	138.9
2001.....	1.4	2.4	4.2	7.1	13.5	25.3	44.1	71.3	100.2	147.4
2000.....	1.4	2.3	4.3	8.0	13.5	24.6	42.3	71.9	103.5	158.3
1999.....	1.4	2.2	3.6	6.5	13.2	24.3	43.0	69.3	105.7	161.8
All races 3/, female										
2002.....	1.2	2.1	3.5	5.7	9.5	15.8	27.3	43.9	65.9	109.0
2001.....	1.2	2.0	3.5	6.2	9.6	17.7	28.9	48.0	71.2	116.6
2000.....	1.4	2.1	3.7	6.3	10.0	18.4	27.8	45.6	69.8	112.6
1999.....	1.3	2.2	3.4	5.8	10.0	17.3	27.4	45.3	65.9	119.4
White, both sexes										
2002.....	1.1	1.7	3.3	6.0	10.7	19.4	35.8	55.0	78.9	122.1
2001.....	1.1	1.9	3.4	6.3	11.3	21.4	36.5	59.3	84.7	129.0
2000.....	1.3	1.9	3.7	6.9	11.5	21.4	35.1	57.8	84.3	130.0
1999.....	1.1	1.9	3.2	5.9	11.5	20.8	34.5	56.2	82.4	134.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of respiratory system (J80-J84,J93-J98)										
White, male										
2002.....	1.2	1.7	3.3	6.8	12.2	23.6	45.3	68.1	97.9	143.8
2001.....	1.4	2.1	3.8	7.0	13.1	25.8	45.5	73.7	103.5	151.5
2000.....	1.3	2.0	3.9	7.8	13.4	25.0	43.1	73.5	106.2	163.4
1999.....	1.2	1.9	3.3	6.1	13.3	24.9	43.5	71.0	108.0	166.3
White, female										
2002.....	1.1	1.8	3.3	5.3	9.3	15.7	28.0	45.6	67.3	112.7
2001.....	0.9	1.7	3.0	5.7	9.6	17.6	29.2	49.1	73.3	119.6
2000.....	1.3	1.9	3.5	6.1	9.8	18.2	28.7	46.8	71.3	116.4
1999.....	1.0	1.9	3.1	5.6	9.9	17.1	27.3	45.9	67.4	122.2
Black, both sexes										
2002.....	2.4	4.5	6.1	9.7	15.1	18.8	30.3	35.2	59.9	76.9
2001.....	2.6	5.0	7.9	10.5	15.2	22.2	32.0	44.6	59.0	90.0
2000.....	2.4	5.2	7.2	10.5	14.2	24.5	29.2	45.3	64.1	82.3
1999.....	3.0	5.0	6.4	9.1	13.3	22.0	34.8	48.2	58.9	94.1
Black, male										
2002.....	2.8	4.2	6.3	10.4	18.6	20.2	40.4	45.0	73.4	86.8
2001.....	2.3	5.3	7.7	9.9	18.9	25.2	35.3	51.1	71.4	106.7
2000.....	2.5	5.9	8.0	11.7	15.7	27.1	35.4	61.4	74.2	104.9
1999.....	2.8	5.4	6.8	9.7	14.9	23.6	41.0	59.4	78.5	114.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of respiratory system (J80-J84,J93-J98)										
Black, female										
2002.....	2.0	4.8	5.9	9.2	12.4	17.8	23.5	29.4	52.9	73.1
2001.....	3.0	4.6	8.0	11.0	12.3	20.1	29.8	40.8	52.7	83.8
2000.....	2.3	4.5	6.4	9.6	13.1	22.7	25.1	35.6	59.0	73.8
1999.....	3.3	4.6	6.1	8.7	12.1	20.8	30.6	41.5	49.3	86.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XI. Diseases of the digestive system (K00-K92)										
All races 3/, both sexes										
2002.....	15.9	26.3	33.8	40.2	50.5	68.6	96.0	139.2	212.9	415.3
2001.....	15.2	26.8	33.0	39.0	53.2	71.4	97.3	138.2	212.8	431.4
2000.....	15.9	25.8	31.7	39.8	52.9	68.9	96.3	140.7	214.9	440.7
1999.....	15.8	25.3	31.0	39.6	53.5	72.3	99.0	138.5	214.9	436.4
All races 3/, male										
2002.....	20.7	37.0	48.7	54.2	65.3	83.9	111.8	155.6	227.7	404.4
2001.....	20.1	37.5	46.8	53.1	68.5	86.9	113.9	153.0	227.3	428.2
2000.....	21.5	37.2	44.7	53.3	68.7	84.7	112.4	160.6	229.3	441.9
1999.....	22.0	37.0	44.4	54.2	68.6	89.0	117.3	154.4	238.2	438.5
All races 3/, female										
2002.....	11.2	16.0	19.5	27.1	37.0	55.5	83.4	127.6	203.9	420.0
2001.....	10.4	16.4	19.8	25.9	39.3	58.1	84.1	127.8	204.1	432.7
2000.....	10.4	14.9	19.3	27.1	38.7	55.5	83.7	126.8	206.5	440.2
1999.....	9.7	14.0	18.2	25.9	39.8	57.9	84.7	127.4	201.3	435.6
White, both sexes										
2002.....	15.9	25.7	32.8	39.3	49.5	69.0	96.4	141.0	215.3	426.3
2001.....	14.7	26.0	31.4	37.0	52.3	71.1	97.7	138.3	216.0	441.6
2000.....	15.4	24.9	29.9	38.2	51.5	68.0	96.5	141.2	217.1	449.3
1999.....	15.0	23.8	29.7	37.8	52.0	71.2	98.7	138.4	217.3	445.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XI. Diseases of the digestive system (K00-K92)										
White, male										
2002.....	21.0	36.4	47.4	52.9	63.7	83.9	111.4	157.0	228.7	412.5
2001.....	19.4	36.7	44.6	50.1	67.0	85.5	113.6	151.9	230.6	434.7
2000.....	20.9	36.5	42.1	51.4	66.8	83.5	111.8	160.1	230.7	448.1
1999.....	21.1	35.0	42.5	51.8	67.1	87.8	117.1	154.3	240.5	443.1
White, female										
2002.....	10.9	15.1	18.5	26.4	36.4	55.9	84.2	129.6	207.2	432.3
2001.....	10.0	15.5	18.5	24.5	38.7	58.4	84.9	128.7	207.2	444.5
2000.....	9.8	13.4	18.0	25.6	37.4	54.4	84.3	127.9	208.9	449.8
1999.....	8.9	12.7	17.2	24.6	38.1	56.7	84.1	127.3	203.5	446.0
Black, both sexes										
2002.....	17.6	35.1	48.3	54.2	65.2	76.8	105.5	137.1	208.0	333.2
2001.....	20.1	36.6	51.1	62.2	68.6	87.7	108.1	151.8	201.0	351.0
2000.....	21.3	37.2	51.0	58.7	72.2	86.3	109.0	147.5	207.0	382.0
1999.....	22.6	40.1	46.6	59.1	74.3	90.3	112.7	150.0	207.5	364.4
Black, male										
2002.....	20.9	47.7	69.4	76.8	89.3	99.5	133.2	158.0	241.5	355.9
2001.....	26.6	51.3	75.3	90.1	91.9	118.7	138.0	184.0	217.5	386.9
2000.....	28.0	49.6	73.8	81.0	97.3	109.0	138.7	181.4	241.1	392.1
1999.....	30.0	59.2	68.5	84.6	95.0	112.4	135.0	171.0	239.8	400.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XI. Diseases of the digestive system (K00-K92)										
Black, female										
2002.....	14.6	24.1	30.4	35.6	46.3	59.9	86.9	124.6	190.6	324.5
2001.....	14.3	23.9	30.5	39.4	50.2	64.8	88.1	132.6	192.6	337.6
2000.....	15.4	26.3	31.5	40.5	52.5	69.5	89.2	127.0	189.8	378.2
1999.....	16.1	23.6	28.0	38.2	58.1	73.9	97.9	137.3	191.4	350.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of other parts of digestive system (K20-K92)										
All races 3/, both sexes										
2002.....	15.9	26.3	33.8	40.2	50.4	68.5	95.9	139.0	212.5	413.8
2001.....	15.2	26.8	33.0	39.0	53.1	71.4	97.2	137.9	212.2	430.2
2000.....	15.9	25.8	31.7	39.8	52.9	68.9	96.2	140.6	214.5	439.5
1999.....	15.8	25.3	31.0	39.6	53.5	72.2	98.9	138.3	214.6	435.4
All races 3/, male										
2002.....	20.7	37.0	48.7	54.1	65.2	83.8	111.6	155.3	227.4	403.1
2001.....	20.1	37.5	46.8	53.0	68.4	86.9	113.8	152.8	226.7	427.0
2000.....	21.4	37.1	44.7	53.3	68.6	84.6	112.2	160.4	229.0	440.8
1999.....	21.9	37.0	44.4	54.2	68.6	88.9	117.1	154.2	238.0	437.4
All races 3/, female										
2002.....	11.2	16.0	19.5	27.0	37.0	55.4	83.3	127.5	203.4	418.4
2001.....	10.4	16.4	19.8	25.9	39.3	58.1	84.1	127.5	203.6	431.5
2000.....	10.4	14.9	19.2	27.1	38.7	55.4	83.6	126.7	206.0	439.0
1999.....	9.7	13.9	18.2	25.9	39.7	57.8	84.6	127.1	200.9	434.6
White, both sexes										
2002.....	15.9	25.7	32.8	39.3	49.5	68.9	96.3	140.8	214.9	424.8
2001.....	14.7	26.0	31.4	37.0	52.2	71.0	97.7	138.1	215.4	440.4
2000.....	15.3	24.9	29.9	38.2	51.5	67.9	96.4	141.1	216.7	448.1
1999.....	15.0	23.7	29.7	37.8	52.0	71.1	98.6	138.2	217.0	444.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of other parts of digestive system (K20-K92)										
White, male										
2002.....	21.0	36.3	47.4	52.8	63.6	83.8	111.2	156.7	228.3	411.2
2001.....	19.4	36.6	44.6	50.0	67.0	85.5	113.6	151.7	230.0	433.4
2000.....	20.9	36.4	42.0	51.4	66.7	83.4	111.6	160.0	230.4	446.9
1999.....	21.1	34.9	42.5	51.7	67.1	87.7	116.9	154.1	240.4	441.9
White, female										
2002.....	10.9	15.1	18.5	26.4	36.4	55.8	84.1	129.5	206.7	430.6
2001.....	10.0	15.5	18.5	24.5	38.7	58.3	84.8	128.5	206.6	443.3
2000.....	9.8	13.4	18.0	25.6	37.4	54.4	84.2	127.8	208.5	448.6
1999.....	8.8	12.7	17.2	24.6	38.0	56.6	84.1	127.1	203.2	445.0
Black, both sexes										
2002.....	17.6	35.0	48.3	54.1	65.2	76.7	105.5	136.9	207.7	332.3
2001.....	20.1	36.6	51.1	62.2	68.6	87.7	108.1	150.9	200.2	350.1
2000.....	21.3	37.2	51.0	58.7	72.2	86.2	108.9	147.5	206.1	381.6
1999.....	22.5	40.0	46.6	59.1	74.2	90.3	112.6	149.6	206.9	363.8
Black, male										
2002.....	20.9	47.5	69.3	76.8	89.3	99.2	133.2	158.0	240.7	355.9
2001.....	26.6	51.3	75.3	90.1	91.9	118.7	138.0	183.1	215.9	385.8
2000.....	28.0	49.6	73.8	81.0	97.3	109.0	138.4	181.4	241.1	392.1
1999.....	29.9	59.2	68.5	84.6	94.8	112.4	135.0	171.0	238.9	400.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of other parts of digestive system (K20-K92)										
Black, female										
2002.....	14.6	24.1	30.4	35.4	46.3	59.9	86.9	124.3	190.6	323.2
2001.....	14.3	23.9	30.5	39.4	50.2	64.8	88.1	131.7	192.1	336.8
2000.....	15.4	26.3	31.5	40.5	52.5	69.3	89.2	127.0	188.5	377.7
1999.....	16.1	23.5	28.0	38.2	58.1	73.9	97.7	136.7	191.0	350.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of esophagus (K20-K22)										
All races 3/, both sexes										
2002.....	0.2	0.3	0.3	0.3	0.5	0.9	1.7	3.5	6.8	20.7
2001.....	*	0.1	0.2	0.4	0.6	1.0	1.8	3.5	7.4	21.2
2000.....	0.1	0.2	0.3	0.4	0.6	1.1	1.8	3.6	7.7	23.3
1999.....	0.1	0.2	0.3	0.4	0.6	1.1	1.9	3.3	7.7	23.5
All races 3/, male										
2002.....	0.2	0.4	0.4	0.3	0.6	1.4	1.9	4.2	8.3	25.4
2001.....	*	*	0.3	0.5	0.9	1.3	2.1	4.1	8.8	23.6
2000.....	0.2	0.3	0.4	0.5	1.0	1.5	2.1	4.6	9.3	28.6
1999.....	*	0.3	0.3	0.4	0.8	1.5	2.7	4.4	9.9	28.3
All races 3/, female										
2002.....	*	*	*	0.3	0.4	0.6	1.6	3.0	5.9	18.7
2001.....	*	*	*	*	0.4	0.8	1.5	3.0	6.6	20.2
2000.....	*	*	*	*	*	0.7	1.6	2.8	6.8	21.1
1999.....	*	*	0.2	0.4	0.4	0.8	1.3	2.5	6.4	21.5
White, both sexes										
2002.....	0.2	0.3	0.3	0.3	0.5	0.9	1.7	3.4	7.0	21.7
2001.....	*	0.1	0.2	0.4	0.6	1.0	1.7	3.6	7.6	22.2
2000.....	0.1	0.2	0.3	0.4	0.6	1.1	1.8	3.7	8.1	24.4
1999.....	*	0.2	0.2	0.4	0.6	1.1	1.9	3.4	7.9	24.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)										
All races 3/, both sexes										
2002.....	0.3	0.6	0.8	1.4	1.7	2.7	4.7	8.0	13.2	28.3
2001.....	0.4	0.7	0.8	1.3	2.1	3.1	4.8	8.3	16.0	33.4
2000.....	0.3	0.6	1.0	1.5	2.2	3.1	5.1	8.7	15.6	35.6
1999.....	0.4	0.6	0.8	1.4	2.2	3.3	5.9	9.1	17.1	34.6
All races 3/, male										
2002.....	0.4	0.7	1.2	1.8	2.1	3.2	6.1	9.3	16.7	31.6
2001.....	0.5	0.9	1.1	1.8	2.8	4.1	5.6	10.3	17.3	34.9
2000.....	0.4	0.9	1.5	2.1	2.9	3.9	6.9	11.2	17.8	40.7
1999.....	0.4	0.7	1.2	1.8	3.0	4.5	7.9	11.0	21.1	37.5
All races 3/, female										
2002.....	0.3	0.4	0.5	1.0	1.4	2.2	3.6	7.0	11.0	26.9
2001.....	0.2	0.4	0.6	0.8	1.5	2.2	4.2	6.9	15.2	32.7
2000.....	0.2	0.4	0.5	0.9	1.6	2.5	3.6	6.9	14.4	33.5
1999.....	0.3	0.5	0.5	1.1	1.5	2.3	4.3	7.7	14.7	33.5
White, both sexes										
2002.....	0.2	0.5	0.8	1.3	1.6	2.6	4.8	7.9	13.6	29.1
2001.....	0.3	0.6	0.7	1.1	2.0	3.0	4.8	8.3	16.2	34.4
2000.....	0.3	0.6	0.9	1.4	2.2	3.0	5.1	8.5	16.0	36.3
1999.....	0.3	0.5	0.7	1.4	2.2	3.1	5.8	9.1	17.1	35.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)										
White, male										
2002.....	0.3	0.6	1.0	1.6	2.0	3.1	6.0	9.2	17.1	32.5
2001.....	0.5	0.8	0.9	1.5	2.7	3.8	5.3	10.1	17.3	35.7
2000.....	0.4	0.8	1.3	2.0	2.9	3.8	6.7	10.9	17.8	40.7
1999.....	0.3	0.6	1.1	1.6	2.9	4.1	7.8	11.1	21.1	38.4
White, female										
2002.....	*	0.4	0.5	1.0	1.3	2.1	3.8	7.0	11.5	27.6
2001.....	*	0.4	0.6	0.8	1.4	2.2	4.4	7.0	15.6	33.9
2000.....	*	0.3	0.5	0.9	1.6	2.4	3.8	6.9	14.9	34.5
1999.....	0.3	0.4	0.4	1.1	1.5	2.2	4.2	7.7	14.8	34.1
Black, both sexes										
2002.....	0.9	0.9	1.6	2.3	2.9	3.2	5.2	8.8	6.8	19.0
2001.....	*	1.1	1.6	2.5	3.0	3.6	5.5	7.8	11.1	20.9
2000.....	*	1.4	2.0	2.6	2.7	4.1	5.7	9.4	11.0	26.6
1999.....	1.0	1.2	1.8	2.4	3.2	5.0	6.5	7.7	16.2	25.5
Black, male										
2002.....	*	*	2.8	3.6	4.2	*	8.0	11.9	*	*
2001.....	*	1.8	2.7	4.1	4.5	5.3	9.1	10.7	*	23.8
2000.....	*	*	3.3	3.5	*	*	10.0	12.8	*	31.1
1999.....	1.5	1.9	2.8	3.7	5.3	7.5	8.8	*	18.3	28.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)										
Black, female										
2002.....	*	*	*	*	*	*	*	6.9	*	18.5
2001.....	*	*	*	*	*	*	*	6.1	9.0	19.8
2000.....	*	*	*	*	*	*	*	7.4	8.5	24.9
1999.....	*	*	*	*	*	*	5.0	6.9	15.1	24.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Gastric ulcer (K25)										
All races 3/, both sexes										
2002.....	0.1	0.2	0.3	0.5	0.5	0.8	1.4	2.1	3.8	7.7
2001.....	0.1	0.3	0.3	0.4	0.6	1.0	1.5	2.4	4.6	8.0
2000.....	0.1	0.3	0.4	0.5	0.7	1.0	1.5	2.4	4.0	9.5
1999.....	0.2	0.2	0.3	0.5	0.6	0.9	1.8	2.7	4.8	8.8
All races 3/, male										
2002.....	*	0.2	0.3	0.6	0.8	1.0	1.8	2.2	3.9	8.1
2001.....	0.2	0.3	0.3	0.5	0.7	1.2	1.8	2.7	4.6	6.0
2000.....	*	0.4	0.5	0.5	0.8	1.2	2.0	3.0	3.8	9.9
1999.....	0.2	0.3	0.5	0.6	0.7	1.0	2.2	2.7	5.4	8.7
All races 3/, female										
2002.....	*	*	*	0.4	*	0.7	1.1	2.0	3.8	7.5
2001.....	*	*	0.2	*	0.6	0.9	1.4	2.2	4.6	8.8
2000.....	*	*	0.2	0.4	0.7	0.9	1.2	2.0	4.1	9.4
1999.....	*	*	*	0.4	0.5	0.8	1.5	2.7	4.4	8.8
White, both sexes										
2002.....	*	0.2	0.2	0.5	0.5	0.8	1.5	2.0	3.8	7.8
2001.....	*	0.2	0.2	0.3	0.6	0.9	1.5	2.4	4.6	8.0
2000.....	*	0.2	0.3	0.4	0.7	1.0	1.5	2.3	4.0	9.4
1999.....	0.1	0.2	0.3	0.4	0.6	0.7	1.8	2.7	4.8	8.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Duodenal ulcer (K26)										
All races 3/, both sexes										
2002.....	0.1	0.2	0.3	0.5	0.7	0.9	1.7	3.1	4.3	9.0
2001.....	0.2	0.3	0.3	0.4	0.8	1.1	1.6	2.9	5.4	11.0
2000.....	0.1	0.2	0.4	0.6	0.9	1.0	1.7	3.0	5.6	10.1
1999.....	0.1	0.2	0.3	0.4	1.0	1.2	1.9	2.9	4.9	10.0
All races 3/, male										
2002.....	*	0.3	0.4	0.8	0.9	1.2	2.2	3.6	6.0	10.1
2001.....	0.3	0.4	0.5	0.6	1.1	1.4	2.0	3.6	6.4	13.8
2000.....	0.2	0.3	0.7	0.8	1.1	1.4	2.3	3.8	6.4	12.3
1999.....	*	0.3	0.5	0.5	1.4	1.9	2.9	4.1	6.7	11.9
All races 3/, female										
2002.....	*	0.2	*	0.3	0.5	0.6	1.3	2.8	3.3	8.5
2001.....	*	*	*	0.3	0.4	0.8	1.3	2.4	4.8	9.8
2000.....	*	*	*	0.3	0.7	0.7	1.3	2.5	5.1	9.2
1999.....	*	*	*	0.3	0.6	0.6	1.2	2.1	3.9	9.3
White, both sexes										
2002.....	0.1	0.2	0.2	0.5	0.7	0.9	1.8	3.2	4.7	9.5
2001.....	0.2	0.2	0.3	0.4	0.8	1.1	1.6	2.9	5.6	11.6
2000.....	*	0.2	0.4	0.6	0.9	1.0	1.7	3.1	5.8	10.5
1999.....	0.1	0.2	0.3	0.4	1.0	1.2	1.9	3.0	5.1	10.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer, site unspecified (K27)										
All races 3/, both sexes										
2002.....	*	0.1	0.3	0.3	0.5	0.8	1.5	2.7	4.9	11.3
2001.....	*	0.1	0.2	0.5	0.7	1.0	1.7	2.9	5.8	14.2
2000.....	*	0.2	0.2	0.5	0.6	1.0	1.7	3.1	5.9	15.5
1999.....	*	0.1	0.2	0.5	0.6	1.2	2.1	3.4	7.2	15.6
All races 3/, male										
2002.....	*	*	0.4	0.4	0.5	1.0	2.0	3.4	6.5	12.8
2001.....	*	*	0.3	0.7	0.9	1.4	1.8	3.9	6.2	14.9
2000.....	*	0.2	0.3	0.7	1.0	1.3	2.5	4.3	7.2	18.2
1999.....	*	*	0.3	0.6	0.9	1.6	2.7	4.1	8.8	16.4
All races 3/, female										
2002.....	*	*	*	0.3	0.5	0.7	1.1	2.2	3.8	10.7
2001.....	*	*	*	0.3	0.4	0.6	1.5	2.2	5.6	13.9
2000.....	*	*	*	*	*	0.8	1.2	2.3	5.1	14.5
1999.....	*	*	*	0.3	0.4	0.9	1.6	2.9	6.3	15.2
White, both sexes										
2002.....	*	0.1	0.3	0.3	0.5	0.8	1.5	2.6	5.0	11.5
2001.....	*	0.2	0.2	0.4	0.6	1.0	1.7	2.9	5.8	14.6
2000.....	*	0.2	0.2	0.5	0.6	1.0	1.8	3.1	6.0	15.9
1999.....	*	*	0.2	0.5	0.6	1.2	2.0	3.4	7.2	16.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Gastritis and duodenitis (K29)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	*	0.3	0.6	1.1	2.9
2001.....	*	*	*	*	*	0.2	0.3	0.6	1.0	3.8
2000.....	*	*	*	0.2	0.2	0.2	0.3	0.6	1.2	3.5
1999.....	*	*	*	*	*	0.2	0.3	0.6	1.5	3.8
All races 3/, male										
2002.....	*	*	*	*	*	*	*	0.8	1.0	2.9
2001.....	*	*	*	*	*	*	*	*	*	4.8
2000.....	*	*	*	*	*	*	*	*	1.2	3.3
1999.....	*	*	*	*	*	*	*	0.8	1.7	3.7
All races 3/, female										
2002.....	*	*	*	*	*	*	*	0.5	1.1	2.9
2001.....	*	*	*	*	*	*	*	0.5	1.0	3.3
2000.....	*	*	*	*	*	*	*	0.6	1.2	3.6
1999.....	*	*	*	*	*	*	*	*	1.4	3.9
White, both sexes										
2002.....	*	*	*	*	*	*	0.3	0.7	1.0	2.9
2001.....	*	*	*	*	*	*	0.3	0.6	0.9	4.0
2000.....	*	*	*	0.2	*	*	0.3	0.6	1.3	3.5
1999.....	*	*	*	*	*	*	0.3	0.5	1.3	3.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Dyspepsia and other diseases of stomach and duodenum (K30-K31)

All races 3/, both sexes

2002.....	0.1	*	0.1	0.2	0.3	0.4	0.6	1.1	1.7	3.2
2001.....	*	*	0.2	*	0.2	0.4	0.6	1.0	1.7	3.6
2000.....	*	*	0.2	0.2	0.2	0.5	0.7	1.2	1.6	4.0
1999.....	*	0.1	*	*	0.3	0.3	0.6	1.0	1.9	4.2

All races 3/, male

2002.....	*	*	*	*	*	0.6	0.7	1.3	1.6	2.9
2001.....	*	*	*	*	*	0.5	0.6	0.9	2.2	3.3
2000.....	*	*	*	*	*	0.7	0.9	1.2	1.7	4.3
1999.....	*	*	*	*	*	*	0.6	1.2	1.6	4.8

All races 3/, female

2002.....	*	*	*	0.3	*	*	0.5	1.1	1.8	3.4
2001.....	*	*	*	*	*	*	0.6	1.0	1.5	3.7
2000.....	*	*	*	*	*	*	0.6	1.1	1.5	3.9
1999.....	*	*	*	*	0.4	*	0.6	0.9	2.1	4.0

White, both sexes

2002.....	*	*	0.1	0.2	0.3	0.4	0.6	1.2	1.6	3.3
2001.....	*	*	0.2	*	*	0.4	0.6	0.9	1.8	3.7
2000.....	*	*	0.2	0.2	0.2	0.4	0.7	1.1	1.6	4.1
1999.....	*	*	*	*	0.3	0.3	0.6	1.1	1.9	4.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of appendix (K35-K38)										
All races 3/, both sexes										
2002.....	*	0.1	0.2	0.2	0.3	0.3	0.7	0.8	1.1	2.0
2001.....	*	0.1	*	0.2	0.2	0.4	0.6	0.5	1.0	2.1
2000.....	*	*	0.1	0.2	0.3	0.3	0.5	0.7	1.4	1.6
1999.....	*	0.1	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.5
All races 3/, male										
2002.....	*	*	0.2	*	0.5	0.5	0.8	1.1	1.6	2.2
2001.....	*	*	*	*	*	0.5	0.9	0.7	1.6	2.8
2000.....	*	*	*	0.3	0.4	*	0.6	1.0	1.6	2.0
1999.....	*	*	*	*	*	*	0.5	0.9	*	2.9
All races 3/, female										
2002.....	*	*	*	*	*	*	0.5	0.6	0.8	2.0
2001.....	*	*	*	*	*	*	0.4	*	0.7	1.8
2000.....	*	*	*	*	*	*	0.4	0.5	1.2	1.5
1999.....	*	*	*	*	*	*	*	*	1.1	0.9
White, both sexes										
2002.....	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.2	2.1
2001.....	*	*	*	*	*	0.3	0.7	0.5	0.9	2.1
2000.....	*	*	*	0.2	0.3	0.2	0.4	0.6	1.3	1.7
1999.....	*	*	0.2	*	*	0.3	0.4	0.6	1.0	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hernia (K40-K46)										
All races 3/, both sexes										
2002.....	0.1	0.2	0.2	0.4	0.6	0.9	1.6	2.8	5.3	12.8
2001.....	0.1	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.6
2000.....	*	0.1	0.3	0.4	0.6	0.9	1.4	3.0	4.9	13.1
1999.....	0.1	*	0.2	0.3	0.7	0.8	1.6	2.5	5.4	12.7
All races 3/, male										
2002.....	*	*	*	0.4	0.5	0.9	1.6	2.7	5.2	11.7
2001.....	*	*	0.3	0.3	0.8	0.9	1.4	2.3	5.3	12.0
2000.....	*	*	0.4	0.3	0.6	0.9	1.4	3.0	5.7	12.6
1999.....	*	*	0.2	0.3	0.8	0.8	1.5	2.5	6.5	13.2
All races 3/, female										
2002.....	*	*	0.3	0.4	0.7	1.0	1.5	2.9	5.3	13.3
2001.....	*	*	0.3	0.5	0.6	0.8	1.6	2.6	4.8	12.8
2000.....	*	*	0.3	0.4	0.6	0.9	1.5	3.0	4.4	13.3
1999.....	*	*	*	*	0.6	0.8	1.6	2.6	4.8	12.6
White, both sexes										
2002.....	*	0.1	0.2	0.4	0.6	0.9	1.6	3.0	5.4	13.4
2001.....	*	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.8
2000.....	*	0.1	0.3	0.4	0.6	0.8	1.5	3.0	5.0	13.3
1999.....	*	*	0.2	0.3	0.7	0.7	1.6	2.5	5.5	13.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Crohn's disease and ulcerative colitis (K50-K51)										
All races 3/, both sexes										
2002.....	0.2	0.2	0.3	0.3	0.4	0.7	1.0	1.7	2.3	3.1
2001.....	*	0.2	0.2	0.5	0.5	0.7	1.1	1.6	2.4	3.8
2000.....	*	0.1	0.2	0.4	0.4	0.6	1.1	1.6	2.2	3.3
1999.....	0.1	0.2	0.3	0.3	0.4	0.6	1.1	1.6	2.5	3.3
All races 3/, male										
2002.....	0.2	0.2	*	*	*	0.8	1.0	1.7	2.3	2.7
2001.....	*	*	0.2	0.5	0.5	0.7	1.3	1.6	2.3	3.5
2000.....	*	*	*	0.3	*	0.6	1.1	1.6	1.6	2.9
1999.....	*	*	0.3	0.3	*	0.7	1.1	1.6	2.5	2.7
All races 3/, female										
2002.....	0.2	0.2	0.3	0.3	0.5	0.7	1.0	1.7	2.3	3.2
2001.....	*	*	0.2	0.5	0.4	0.6	0.9	1.5	2.5	3.9
2000.....	*	*	0.3	0.4	0.4	0.7	1.2	1.6	2.6	3.4
1999.....	*	0.3	0.3	0.3	0.5	0.5	1.1	1.6	2.6	3.5
White, both sexes										
2002.....	0.2	0.2	0.3	0.3	0.5	0.8	1.0	1.8	2.5	3.4
2001.....	*	0.2	0.2	0.5	0.5	0.7	1.2	1.7	2.5	3.9
2000.....	*	0.2	0.2	0.4	0.4	0.7	1.2	1.7	2.4	3.5
1999.....	0.2	0.2	0.3	0.3	0.4	0.7	1.2	1.6	2.8	3.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

All races 3/, both sexes

2002.....	0.5	0.9	1.4	2.8	4.8	9.0	17.0	29.6	53.0	113.3
2001.....	0.4	0.8	1.6	2.7	5.4	10.1	17.8	29.7	52.1	113.5
2000.....	0.5	0.9	1.7	2.8	5.3	9.5	17.5	31.5	52.9	116.2
1999.....	0.5	0.8	1.5	2.9	5.3	10.1	17.1	29.9	52.5	116.2

All races 3/, male

2002.....	0.6	0.8	1.5	2.5	4.7	8.1	15.8	26.8	49.7	96.7
2001.....	0.5	0.8	1.6	2.3	5.1	9.1	17.6	27.2	47.9	98.5
2000.....	0.4	0.9	1.6	2.7	5.2	8.9	16.6	30.0	48.2	98.5
1999.....	0.6	0.8	1.4	2.9	4.9	8.7	16.7	27.9	48.7	95.6

All races 3/, female

2002.....	0.4	0.9	1.4	3.0	4.9	9.8	18.0	31.6	55.0	120.5
2001.....	0.4	0.9	1.6	3.0	5.6	11.0	18.0	31.5	54.6	119.8
2000.....	0.5	0.9	1.8	2.8	5.4	10.0	18.2	32.5	55.7	123.4
1999.....	0.5	0.7	1.6	3.0	5.7	11.3	17.5	31.3	54.8	124.5

White, both sexes

2002.....	0.5	0.8	1.3	2.6	4.6	8.7	16.4	29.8	53.0	115.9
2001.....	0.4	0.8	1.3	2.4	5.0	9.7	17.7	29.1	52.9	116.2
2000.....	0.4	0.7	1.5	2.6	4.8	9.2	17.3	31.1	52.9	118.0
1999.....	0.4	0.6	1.4	2.9	5.3	9.7	16.5	29.2	52.9	118.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Vascular disorders and obstruction of intestine without hernia (K55-K56)										
White, male										
2002.....	0.5	0.7	1.3	2.3	4.2	7.8	14.9	26.7	48.7	97.5
2001.....	0.4	0.7	1.3	2.1	4.8	8.4	17.3	26.2	48.3	98.9
2000.....	0.3	0.6	1.4	2.4	4.5	8.2	16.2	29.0	47.0	98.9
1999.....	0.5	0.7	1.2	2.8	4.7	8.2	16.0	27.3	47.9	94.8
White, female										
2002.....	0.4	0.8	1.2	2.9	4.9	9.5	17.7	32.0	55.6	123.9
2001.....	0.4	0.8	1.4	2.7	5.2	10.9	18.0	31.1	55.7	123.4
2000.....	0.5	0.7	1.5	2.8	5.1	9.9	18.2	32.5	56.4	125.8
1999.....	0.4	0.6	1.5	2.9	5.8	11.0	16.8	30.6	55.9	127.4
Black, both sexes										
2002.....	1.0	1.8	3.0	4.6	7.2	13.3	25.3	32.3	63.3	99.8
2001.....	0.8	1.8	3.9	5.1	10.2	15.4	24.0	40.1	53.6	96.7
2000.....	0.9	2.4	4.0	5.4	10.4	14.3	21.7	40.7	59.0	112.0
1999.....	1.4	1.8	3.2	4.3	7.4	15.9	26.1	42.3	54.9	108.7
Black, male										
2002.....	*	1.9	3.5	4.8	9.5	12.5	27.4	30.3	71.8	105.1
2001.....	*	1.8	4.0	5.2	9.6	18.8	26.9	42.7	57.0	120.3
2000.....	*	3.0	4.1	6.9	13.0	16.4	23.4	45.8	70.8	110.7
1999.....	1.8	2.3	3.3	4.6	8.3	15.3	25.1	39.9	68.0	110.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

Black, female

2002.....	*	1.8	2.6	4.5	5.4	13.9	24.0	33.5	58.9	97.8
2001.....	*	1.8	3.8	5.1	10.7	13.0	22.0	38.6	51.9	88.0
2000.....	*	1.8	4.0	4.2	8.3	12.7	20.6	37.6	53.1	112.5
1999.....	*	*	3.0	4.2	6.7	16.4	26.8	43.8	48.4	108.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diverticular disease of intestine (K57)										
All races 3/, both sexes										
2002.....	*	0.2	0.3	0.5	0.9	1.9	3.5	7.3	13.4	27.7
2001.....	*	0.2	0.3	0.6	1.3	2.2	3.6	7.0	12.8	31.4
2000.....	0.1	0.2	0.3	0.4	1.1	2.2	3.7	6.5	13.3	30.9
1999.....	*	0.2	0.3	0.7	1.1	2.2	4.1	6.9	12.8	33.4
All races 3/, male										
2002.....	*	0.3	0.4	0.6	0.9	1.7	3.1	6.5	10.4	19.9
2001.....	*	0.3	0.3	0.6	1.1	1.6	3.2	6.6	10.7	23.4
2000.....	*	0.2	0.3	0.6	1.0	2.1	3.4	5.2	10.8	22.3
1999.....	*	*	0.2	0.6	1.2	1.9	3.7	5.5	10.4	25.8
All races 3/, female										
2002.....	*	*	*	0.5	1.0	2.0	3.8	7.9	15.2	31.0
2001.....	*	*	0.2	0.6	1.4	2.7	3.9	7.3	14.1	34.7
2000.....	*	*	0.3	0.3	1.2	2.4	4.0	7.4	14.8	34.4
1999.....	*	*	0.3	0.8	1.0	2.6	4.3	7.9	14.1	36.4
White, both sexes										
2002.....	*	0.2	0.3	0.5	0.9	1.9	3.6	7.4	13.7	28.5
2001.....	*	0.2	0.3	0.6	1.3	2.0	3.7	7.2	13.5	32.4
2000.....	*	0.2	0.3	0.4	1.0	2.2	3.8	6.7	13.7	32.3
1999.....	*	0.2	0.3	0.7	1.1	2.2	4.2	7.2	13.1	34.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diverticular disease of intestine (K57)										
Black, female										
2002.....	*	*	*	*	*	*	*	6.0	13.1	24.6
2001.....	*	*	*	*	*	4.7	4.7	6.1	*	27.8
2000.....	*	*	*	*	*	*	*	6.0	12.3	24.5
1999.....	*	*	*	*	*	4.2	*	*	11.7	27.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of intestines and peritoneum (K52,K58-K63)										
All races 3/, both sexes										
2002.....	0.3	0.4	0.7	1.2	2.1	3.5	5.2	9.9	16.2	32.8
2001.....	0.3	0.4	0.7	1.1	1.9	3.5	5.6	10.1	16.8	34.1
2000.....	0.3	0.5	0.6	1.2	2.0	3.0	5.3	9.5	16.9	33.3
1999.....	0.3	0.4	0.7	1.2	1.9	3.4	5.6	9.7	15.7	32.0
All races 3/, male										
2002.....	0.3	0.5	0.7	1.2	2.0	3.4	5.5	10.5	13.8	29.4
2001.....	0.2	0.4	0.7	1.0	1.9	3.2	5.4	9.9	15.6	29.9
2000.....	0.3	0.4	0.5	1.3	2.1	2.7	5.6	9.1	15.1	28.7
1999.....	0.3	0.6	0.7	1.2	1.8	3.1	5.5	9.8	15.5	28.9
All races 3/, female										
2002.....	0.3	0.4	0.7	1.2	2.3	3.5	5.1	9.4	17.7	34.3
2001.....	0.3	0.5	0.8	1.1	2.0	3.7	5.7	10.2	17.5	35.8
2000.....	0.2	0.5	0.8	1.1	1.8	3.3	5.1	9.7	17.9	35.2
1999.....	0.3	0.3	0.6	1.2	2.1	3.6	5.7	9.7	15.8	33.2
White, both sexes										
2002.....	0.3	0.4	0.7	1.2	2.1	3.6	5.1	10.1	16.7	34.1
2001.....	0.3	0.4	0.7	0.9	1.8	3.5	5.7	10.0	16.8	35.2
2000.....	0.2	0.5	0.6	1.2	1.8	3.0	5.4	9.6	17.2	33.7
1999.....	0.3	0.4	0.6	1.2	1.9	3.3	5.6	9.8	15.9	33.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of intestines and peritoneum (K52,K58-K63)										
White, male										
2002.....	0.3	0.5	0.7	1.2	1.9	3.4	5.1	10.7	13.7	30.3
2001.....	0.2	0.4	0.7	0.9	1.7	3.0	5.1	9.8	15.7	31.2
2000.....	0.3	0.4	0.4	1.2	1.8	2.7	5.7	9.1	15.2	28.1
1999.....	0.3	0.5	0.7	1.2	1.7	3.1	5.6	9.6	15.7	28.3
White, female										
2002.....	0.3	0.4	0.7	1.2	2.2	3.8	5.1	9.7	18.5	35.8
2001.....	0.3	0.4	0.8	1.0	1.9	3.9	6.1	10.2	17.4	36.8
2000.....	*	0.5	0.7	1.1	1.8	3.2	5.1	10.0	18.4	36.1
1999.....	0.2	0.3	0.6	1.2	2.1	3.6	5.7	9.9	16.0	34.9
Black, both sexes										
2002.....	*	*	1.0	1.7	3.3	3.3	7.8	8.9	13.9	23.2
2001.....	*	0.8	1.1	2.3	2.7	4.1	6.1	13.0	19.8	27.0
2000.....	*	*	1.1	1.8	3.5	3.4	6.1	8.0	15.0	29.7
1999.....	*	0.9	1.2	1.5	2.3	4.3	6.4	8.9	16.2	23.6
Black, male										
2002.....	*	*	*	*	*	5.0	10.7	10.6	16.0	24.7
2001.....	*	*	*	*	*	5.8	8.8	15.8	16.9	*
2000.....	*	*	*	*	4.1	*	*	10.9	17.7	30.0
1999.....	*	*	*	*	*	*	*	10.4	17.4	39.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of intestines and peritoneum (K52,K58-K63)										
Black, female										
2002.....	*	*	*	*	3.4	*	5.8	8.0	12.7	22.6
2001.....	*	*	*	*	*	*	*	11.4	21.3	29.5
2000.....	*	*	*	*	*	4.0	6.4	6.3	13.6	29.6
1999.....	*	*	*	*	*	4.2	7.2	8.0	15.6	17.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of peritoneum (K65-K66)										
All races 3/, both sexes										
2002.....	0.2	0.4	0.6	0.7	1.0	1.3	2.3	3.7	6.2	10.3
2001.....	0.2	0.4	0.5	0.8	0.9	1.4	2.3	3.6	5.7	10.4
2000.....	0.3	0.4	0.5	0.9	1.2	1.4	2.7	3.9	5.3	10.6
1999.....	0.3	0.4	0.5	0.8	1.1	1.6	2.8	3.7	5.5	9.7
All races 3/, male										
2002.....	0.2	0.4	0.7	0.8	1.1	1.3	2.0	3.7	5.1	9.1
2001.....	0.3	0.5	0.6	0.9	1.0	1.7	2.2	3.6	6.2	10.1
2000.....	0.3	0.4	0.5	1.0	1.4	1.5	2.7	4.3	5.2	11.0
1999.....	0.3	0.5	0.7	0.8	1.3	1.7	3.1	4.1	6.3	10.1
All races 3/, female										
2002.....	0.2	0.3	0.5	0.6	1.0	1.3	2.5	3.7	6.8	10.8
2001.....	0.2	0.4	0.5	0.6	0.8	1.1	2.3	3.5	5.4	10.5
2000.....	0.3	0.3	0.5	0.9	1.0	1.4	2.6	3.6	5.3	10.5
1999.....	0.2	0.2	0.4	0.8	0.9	1.5	2.5	3.5	5.1	9.6
White, both sexes										
2002.....	0.2	0.3	0.5	0.7	0.9	1.2	2.2	3.8	6.1	10.5
2001.....	0.2	0.4	0.5	0.6	0.7	1.3	2.2	3.5	5.6	10.4
2000.....	0.2	0.3	0.5	0.8	1.1	1.3	2.6	3.8	5.2	10.6
1999.....	0.2	0.4	0.5	0.7	1.0	1.4	2.7	3.5	5.6	9.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of peritoneum (K65-K66)										
Black, female										
2002.....	*	*	*	*	*	*	*	*	8.0	11.1
2001.....	*	*	*	*	*	*	*	*	8.2	10.5
2000.....	*	*	*	*	*	*	*	*	*	10.7
1999.....	*	*	*	*	*	*	4.5	*	*	12.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of liver (K70-K76)										
All races 3/, both sexes										
2002.....	12.3	20.7	25.7	27.5	31.2	36.0	40.3	41.9	41.9	35.7
2001.....	11.9	21.1	25.0	26.4	31.7	36.6	39.7	42.2	40.2	36.1
2000.....	12.4	20.4	23.3	26.9	32.3	36.3	39.7	41.5	40.7	36.7
1999.....	12.1	19.9	23.0	26.6	32.9	38.0	40.9	42.4	41.9	37.2
All races 3/, male										
2002.....	16.3	29.8	38.7	39.7	43.1	47.8	51.6	51.0	52.8	47.3
2001.....	16.0	30.6	37.2	38.3	44.4	49.4	51.0	51.2	50.4	49.4
2000.....	17.2	30.5	34.8	38.2	45.2	48.4	51.1	52.8	48.1	52.2
1999.....	17.4	30.2	34.9	39.4	45.5	52.4	52.6	51.1	53.5	52.3
All races 3/, female										
2002.....	8.4	11.8	13.3	15.9	20.4	25.8	31.3	35.5	35.2	30.7
2001.....	7.9	11.8	13.3	15.3	20.3	25.5	30.7	35.9	34.1	30.5
2000.....	7.7	10.6	12.4	16.3	20.7	25.9	30.8	33.6	36.4	30.4
1999.....	6.9	10.0	11.6	14.6	21.5	25.8	31.9	36.3	35.0	31.1
White, both sexes										
2002.....	12.6	20.5	25.3	27.2	31.4	37.0	41.9	43.6	42.9	36.6
2001.....	11.8	20.9	24.2	25.6	32.1	37.4	40.6	43.4	41.4	37.0
2000.....	12.3	20.2	22.3	26.2	32.3	36.8	40.7	42.9	41.6	37.5
1999.....	11.7	19.0	22.5	25.6	32.3	38.4	41.6	43.4	43.2	37.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of liver (K70-K76)										
White, male										
2002.....	16.9	30.0	38.2	39.3	43.2	48.8	53.3	53.0	53.9	48.5
2001.....	15.8	30.6	36.1	36.9	44.5	50.2	52.1	52.3	51.8	50.8
2000.....	17.2	30.7	33.3	37.4	45.3	49.2	52.0	54.3	49.8	53.3
1999.....	17.1	29.1	34.1	38.0	45.2	52.7	53.4	52.3	55.7	53.8
White, female										
2002.....	8.3	11.1	12.8	15.6	20.4	26.6	32.5	36.9	36.1	31.4
2001.....	7.8	11.3	12.6	14.9	20.7	26.2	31.4	37.2	35.1	31.2
2000.....	7.3	9.8	11.6	15.6	20.3	26.0	31.5	34.8	36.8	31.1
1999.....	6.4	9.1	11.1	13.8	20.5	26.0	32.3	37.1	35.8	31.2
Black, both sexes										
2002.....	11.2	24.4	33.4	34.0	33.9	32.6	30.4	30.1	28.8	27.6
2001.....	13.6	25.5	34.4	37.6	33.0	36.0	34.0	33.0	29.0	21.8
2000.....	14.2	25.2	34.8	34.5	36.0	36.7	35.6	29.1	28.2	26.0
1999.....	15.6	29.0	30.9	36.9	41.6	38.1	36.5	34.5	28.0	31.2
Black, male										
2002.....	13.4	32.7	50.1	51.5	49.9	47.1	41.7	36.7	39.0	36.5
2001.....	18.6	36.9	53.4	56.8	50.6	53.4	48.1	44.1	37.7	27.2
2000.....	18.5	34.6	53.4	50.7	50.5	51.0	50.5	41.1	29.5	35.8
1999.....	21.2	43.5	47.7	55.8	56.4	55.8	50.5	42.3	33.1	37.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of liver (K70-K76)										
Black, female										
2002.....	9.4	17.1	19.3	19.6	21.4	21.8	22.9	26.1	23.5	24.2
2001.....	9.2	15.7	18.3	21.8	19.0	23.1	24.7	26.3	24.5	19.8
2000.....	10.4	17.0	18.8	21.2	24.5	26.1	25.8	21.9	27.6	22.3
1999.....	10.6	16.4	16.7	21.5	30.0	25.0	27.2	29.8	25.5	28.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)										
All races 3/, both sexes										
2002.....	5.9	9.4	11.4	11.4	11.8	10.7	9.5	6.7	4.4	2.1
2001.....	6.1	10.2	11.3	11.1	12.0	11.1	9.0	6.9	3.9	2.4
2000.....	6.6	10.0	10.7	11.7	12.9	11.4	9.1	7.2	4.4	2.2
1999.....	6.4	10.0	10.9	11.5	12.6	11.6	9.4	7.3	5.2	2.8
All races 3/, male										
2002.....	8.0	13.8	17.7	17.6	19.2	17.4	15.4	10.9	8.0	4.7
2001.....	8.4	15.3	17.0	18.0	19.0	17.9	15.2	12.1	7.0	5.5
2000.....	9.4	15.3	16.9	18.0	20.3	18.7	15.7	13.8	7.7	5.3
1999.....	9.4	15.5	17.2	18.7	20.0	19.7	16.2	12.1	9.7	6.1
All races 3/, female										
2002.....	3.9	5.1	5.4	5.6	5.1	4.9	4.8	3.7	2.2	1.0
2001.....	3.9	5.3	5.8	4.7	5.6	5.3	4.0	3.3	2.1	1.0
2000.....	3.9	4.8	4.8	5.8	6.2	5.1	3.9	2.6	2.4	1.0
1999.....	3.5	4.6	4.8	4.9	5.9	4.6	4.0	3.9	2.7	1.5
White, both sexes										
2002.....	6.2	9.6	11.5	11.4	12.0	11.0	9.8	6.8	4.5	2.1
2001.....	6.2	10.5	11.0	11.0	12.2	11.2	9.2	7.1	4.0	2.3
2000.....	6.6	10.0	10.6	11.5	13.0	11.6	9.3	7.3	4.3	2.3
1999.....	6.2	9.8	10.8	11.2	12.4	11.5	9.4	7.2	5.3	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)										
White, male										
2002.....	8.3	14.3	17.9	17.9	19.4	17.7	15.8	11.1	8.2	4.5
2001.....	8.5	15.9	16.8	17.6	19.4	17.8	15.4	12.5	7.2	5.4
2000.....	9.4	15.6	16.8	18.0	20.6	19.0	15.9	14.1	7.8	5.4
1999.....	9.2	15.4	17.1	18.2	20.0	19.5	16.2	11.9	9.8	5.8
White, female										
2002.....	4.0	5.0	5.2	5.3	5.1	5.0	4.9	3.7	2.2	1.1
2001.....	3.9	5.2	5.4	4.6	5.7	5.4	4.2	3.3	2.1	1.0
2000.....	3.8	4.4	4.5	5.3	6.0	5.1	4.0	2.6	2.3	1.0
1999.....	3.3	4.3	4.5	4.7	5.5	4.6	4.0	3.9	2.6	1.5
Black, both sexes										
2002.....	4.2	8.2	12.1	12.5	12.0	9.6	7.4	6.2	*	*
2001.....	5.9	9.0	14.7	14.1	11.5	13.0	7.8	5.7	*	*
2000.....	6.9	11.0	13.3	13.9	12.6	11.8	8.4	6.2	*	*
1999.....	7.2	12.8	12.8	14.8	15.6	12.9	10.4	8.0	*	*
Black, male										
2002.....	5.6	11.5	18.7	17.4	20.4	16.5	13.0	10.6	*	*
2001.....	8.2	12.9	22.3	23.5	18.7	22.9	15.1	*	*	*
2000.....	9.6	15.0	20.6	20.4	18.0	19.3	16.7	11.3	*	*
1999.....	10.1	19.2	20.2	24.9	23.3	24.1	18.7	14.2	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)										
Black, female										
2002.....	3.0	5.3	6.6	8.4	5.4	4.4	*	*	*	*
2001.....	3.9	5.5	8.2	6.4	5.8	5.6	*	*	*	*
2000.....	4.5	7.5	7.1	8.5	8.3	6.3	*	*	*	*
1999.....	4.7	7.2	6.6	6.5	9.5	4.6	5.0	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic hepatitis, not elsewhere classified (K73)

All races 3/, both sexes

2002.....	0.1	0.2	0.2	0.2	0.4	0.7	0.8	1.0	1.0	0.8
2001.....	*	0.2	0.2	0.3	0.3	0.6	0.8	0.8	0.8	0.7
2000.....	0.1	0.1	0.2	0.3	0.3	0.7	0.6	0.9	0.8	0.8
1999.....	*	0.2	0.2	0.3	0.4	0.4	0.7	0.8	0.9	0.6

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2002.....	*	0.2	*	0.4	0.6	1.0	1.1	1.3	1.3	0.9
2001.....	*	0.2	0.2	0.4	0.5	0.9	1.0	1.1	1.0	0.7
2000.....	*	*	*	0.4	0.6	1.1	0.9	1.2	1.0	0.9
1999.....	*	*	*	0.4	0.6	0.6	0.9	1.0	1.2	0.7

White, both sexes

2002.....	*	0.2	0.2	0.2	0.4	0.7	0.9	1.1	0.9	0.8
2001.....	*	0.2	0.2	0.3	0.3	0.6	0.7	0.8	0.9	0.8
2000.....	*	*	0.2	0.3	0.4	0.7	0.6	0.9	0.8	0.9
1999.....	*	0.2	0.2	0.3	0.4	0.4	0.7	0.8	0.9	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Fibrosis and cirrhosis of liver (K74)										
All races 3/, both sexes										
2002.....	3.6	6.4	8.7	10.1	12.2	16.5	20.9	24.8	24.5	18.5
2001.....	3.6	6.7	8.6	9.7	12.4	17.2	21.4	23.5	23.9	19.1
2000.....	3.5	6.7	7.7	9.9	12.9	16.8	21.1	23.9	24.4	20.0
1999.....	3.4	6.3	7.5	9.6	13.5	17.7	21.6	24.6	24.6	19.8
All races 3/, male										
2002.....	4.8	9.3	13.1	14.3	15.1	19.6	24.1	27.9	28.5	23.2
2001.....	4.7	9.8	12.9	13.3	16.2	21.5	25.2	25.2	27.8	25.7
2000.....	4.9	10.1	11.5	13.6	17.3	20.4	24.8	26.6	26.4	28.4
1999.....	5.1	9.6	11.3	13.6	17.0	21.9	24.9	27.5	28.5	26.6
All races 3/, female										
2002.....	2.5	3.6	4.5	6.1	9.6	13.9	18.3	22.6	22.1	16.5
2001.....	2.4	3.7	4.5	6.3	9.1	13.4	18.5	22.4	21.6	16.4
2000.....	2.1	3.5	4.1	6.5	8.9	13.7	18.2	21.9	23.2	16.6
1999.....	1.8	3.0	3.8	5.8	10.2	14.1	19.0	22.5	22.4	17.0
White, both sexes										
2002.....	3.8	6.3	8.5	10.1	12.4	17.3	22.0	26.2	25.5	19.4
2001.....	3.5	6.5	8.3	9.4	12.7	17.9	22.4	24.7	25.1	20.1
2000.....	3.5	6.8	7.4	9.9	13.1	17.2	21.9	25.2	25.4	20.8
1999.....	3.4	5.9	7.4	9.3	13.5	18.3	22.5	25.7	25.8	20.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Fibrosis and cirrhosis of liver (K74)										
White, male										
2002.....	5.1	9.3	12.8	14.2	15.1	20.2	25.3	29.8	29.5	24.4
2001.....	4.6	9.5	12.3	12.8	16.2	22.2	26.2	26.2	29.5	27.3
2000.....	5.0	10.2	10.9	13.4	17.5	21.0	25.7	27.8	27.7	29.4
1999.....	5.0	9.1	11.1	13.2	17.1	22.6	25.9	28.7	30.0	27.9
White, female										
2002.....	2.5	3.3	4.4	6.2	9.9	14.7	19.3	23.7	23.0	17.2
2001.....	2.4	3.5	4.4	6.1	9.6	14.1	19.3	23.6	22.4	17.1
2000.....	2.0	3.4	4.0	6.5	9.0	13.8	18.9	23.3	24.1	17.3
1999.....	1.7	2.8	3.7	5.7	10.2	14.6	19.7	23.6	23.4	17.5
Black, both sexes										
2002.....	3.4	8.8	12.4	12.0	13.0	12.6	12.3	13.9	11.0	9.2
2001.....	4.5	9.9	12.0	13.5	12.0	13.5	14.1	14.1	11.4	7.4
2000.....	3.9	8.0	12.2	11.8	13.2	14.8	15.9	11.9	12.7	11.6
1999.....	4.6	9.5	9.8	13.1	14.5	14.1	14.1	14.6	12.4	12.1
Black, male										
2002.....	4.0	11.5	19.2	19.5	18.6	18.2	16.0	12.4	*	*
2001.....	6.2	15.1	19.8	20.3	19.5	19.8	18.5	18.1	*	*
2000.....	4.8	11.3	19.8	17.9	19.3	18.5	20.7	16.1	*	*
1999.....	7.0	15.0	16.1	19.2	19.1	19.2	18.0	17.6	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Fibrosis and cirrhosis of liver (K74)										
Black, female										
2002.....	2.8	6.4	6.7	5.9	8.7	8.5	9.9	14.8	9.5	9.0
2001.....	2.9	5.4	5.3	7.8	6.1	8.8	11.2	11.6	12.7	*
2000.....	3.1	5.1	5.8	6.7	8.3	12.1	12.7	9.4	13.2	10.3
1999.....	2.5	4.8	4.5	8.2	10.8	10.4	11.5	12.9	11.7	12.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of liver (K71-K72,K75-K76)										
All races 3/, both sexes										
2002.....	2.7	4.7	5.4	5.7	6.8	8.1	9.2	9.4	12.0	14.3
2001.....	2.1	3.9	4.9	5.3	7.0	7.7	8.6	10.9	11.5	13.8
2000.....	2.2	3.6	4.7	5.0	6.2	7.4	8.9	9.5	11.2	13.6
1999.....	2.2	3.5	4.5	5.1	6.4	8.4	9.3	9.8	11.0	14.0
All races 3/, male										
2002.....	3.4	6.5	7.7	7.7	8.7	10.5	11.6	11.6	15.9	18.9
2001.....	2.8	5.4	7.1	6.8	9.1	9.8	10.2	13.5	15.0	17.4
2000.....	2.8	5.1	6.2	6.4	7.4	9.1	10.4	12.1	13.6	17.8
1999.....	2.9	4.9	6.1	6.9	8.2	10.6	11.0	11.0	14.9	19.2
All races 3/, female										
2002.....	2.0	2.9	3.2	3.8	5.0	6.0	7.2	7.9	9.6	12.3
2001.....	1.5	2.6	2.8	3.9	5.1	6.0	7.2	9.0	9.4	12.3
2000.....	1.6	2.2	3.2	3.7	5.1	6.0	7.7	7.8	9.7	11.9
1999.....	1.6	2.1	2.9	3.5	4.7	6.4	7.9	8.9	8.8	11.9
White, both sexes										
2002.....	2.6	4.4	5.1	5.4	6.6	8.1	9.2	9.4	11.9	14.3
2001.....	2.0	3.7	4.7	4.9	6.8	7.8	8.4	10.8	11.4	13.8
2000.....	2.1	3.4	4.2	4.6	5.8	7.4	8.8	9.5	11.1	13.6
1999.....	2.1	3.2	4.1	4.8	6.0	8.2	9.1	9.7	11.2	13.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of liver (K71-K72,K75-K76)										
White, male										
2002.....	3.3	6.2	7.3	7.1	8.5	10.6	11.7	11.3	15.8	18.9
2001.....	2.6	5.1	6.8	6.3	8.9	10.1	10.0	13.2	14.6	17.3
2000.....	2.8	4.8	5.5	5.9	7.0	8.9	10.3	12.0	13.9	17.8
1999.....	2.8	4.5	5.7	6.5	7.8	10.5	10.9	11.2	15.4	19.6
White, female										
2002.....	1.8	2.6	3.0	3.7	4.9	5.9	7.2	8.0	9.6	12.3
2001.....	1.4	2.4	2.6	3.6	4.9	5.8	7.0	9.1	9.4	12.4
2000.....	1.4	2.0	2.9	3.3	4.7	6.0	7.6	7.6	9.4	11.9
1999.....	1.3	1.8	2.6	3.1	4.3	6.2	7.6	8.7	8.7	11.6
Black, both sexes										
2002.....	3.3	7.0	8.5	9.3	8.4	9.7	10.5	10.0	12.8	15.4
2001.....	3.0	6.4	7.6	9.9	9.3	9.1	11.4	12.0	14.4	11.4
2000.....	3.3	6.0	8.9	8.6	10.0	9.2	10.9	10.8	10.7	12.2
1999.....	3.7	6.4	8.0	8.7	10.9	10.4	11.2	11.2	9.2	14.0
Black, male										
2002.....	3.6	9.4	11.9	14.6	10.9	12.2	12.7	13.8	17.6	*
2001.....	4.1	8.7	11.1	13.0	12.2	10.2	14.5	17.2	23.3	*
2000.....	3.9	8.2	12.8	12.2	13.3	13.0	13.0	13.7	*	*
1999.....	4.1	9.0	11.1	11.7	14.0	12.5	13.2	10.4	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of liver (K71-K72,K75-K76)										
Black, female										
2002.....	3.0	4.9	5.5	5.0	6.5	7.8	9.0	7.7	10.3	13.6
2001.....	2.1	4.4	4.5	7.3	7.0	8.3	9.4	8.9	9.8	10.1
2000.....	2.7	4.0	5.5	5.7	7.5	6.3	9.5	9.1	11.0	10.7
1999.....	3.3	4.2	5.3	6.2	8.5	8.9	9.9	11.7	9.9	13.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)										
All races 3/, both sexes										
2002.....	0.2	0.2	0.4	0.6	1.1	2.0	3.6	5.7	10.8	22.9
2001.....	0.1	0.3	0.4	0.7	1.3	2.0	3.8	5.8	10.0	23.7
2000.....	0.1	0.2	0.4	0.6	1.0	1.9	3.0	5.7	10.3	24.8
1999.....	0.2	0.3	0.4	0.7	1.0	1.8	3.4	5.7	10.9	23.9
All races 3/, male										
2002.....	0.2	0.3	0.5	0.8	1.1	2.3	4.0	7.5	13.5	24.1
2001.....	*	0.4	0.4	0.9	1.5	2.2	4.4	7.6	12.1	26.9
2000.....	*	0.2	0.3	0.7	1.2	2.3	3.6	7.4	12.9	27.6
1999.....	*	0.3	0.4	0.9	1.3	2.0	4.2	6.9	13.3	27.7
All races 3/, female										
2002.....	*	*	0.3	0.4	1.0	1.8	3.2	4.4	9.1	22.4
2001.....	*	0.3	0.3	0.5	1.1	1.9	3.3	4.6	8.7	22.4
2000.....	*	0.2	0.4	0.5	0.8	1.6	2.5	4.5	8.7	23.7
1999.....	*	0.3	0.5	0.6	0.7	1.6	2.8	4.8	9.5	22.4
White, both sexes										
2002.....	0.2	0.2	0.3	0.6	1.0	2.0	3.5	5.5	10.6	23.5
2001.....	*	0.3	0.3	0.6	1.3	1.8	3.7	5.7	9.8	23.9
2000.....	0.1	0.2	0.3	0.6	0.9	1.8	2.7	5.6	10.1	25.6
1999.....	0.1	0.3	0.4	0.6	0.9	1.6	3.4	5.7	11.0	24.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Black, female

2002.....	*	*	*	*	*	*	5.4	7.7	11.1	16.0
2001.....	*	*	*	*	*	*	6.3	6.9	13.1	16.8
2000.....	*	*	*	*	*	*	4.9	6.3	11.5	19.3
1999.....	*	*	*	*	*	*	*	*	12.1	15.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of biliary tract and pancreas (K83-K86)										
All races 3/, both sexes										
2002.....	1.0	1.3	1.5	2.3	2.7	4.0	5.8	8.1	12.0	21.7
2001.....	1.0	1.4	1.7	2.1	3.1	4.2	6.1	7.6	11.9	22.9
2000.....	1.0	1.3	1.5	2.3	2.7	3.7	5.5	7.9	11.5	22.0
1999.....	1.0	1.3	1.6	2.0	2.8	3.6	5.4	7.2	10.3	22.2
All races 3/, male										
2002.....	1.3	1.9	2.2	3.0	4.0	5.5	7.4	10.3	14.3	21.2
2001.....	1.3	1.9	2.2	2.8	3.9	5.1	8.0	8.8	12.5	22.4
2000.....	1.5	1.8	2.1	2.9	3.1	5.2	6.7	10.2	13.3	22.9
1999.....	1.4	1.8	2.1	2.8	3.6	4.6	6.7	9.1	11.7	23.1
All races 3/, female										
2002.....	0.7	0.7	0.9	1.7	1.6	2.7	4.4	6.6	10.5	21.9
2001.....	0.7	1.0	1.2	1.4	2.4	3.3	4.5	6.8	11.6	23.2
2000.....	0.6	0.9	1.0	1.7	2.3	2.5	4.6	6.3	10.5	21.6
1999.....	0.7	0.8	1.1	1.4	2.2	2.8	4.4	5.9	9.5	21.8
White, both sexes										
2002.....	0.9	1.1	1.3	2.2	2.5	3.9	5.7	8.2	12.1	22.0
2001.....	0.9	1.3	1.4	1.9	3.0	4.1	6.0	7.6	12.0	23.0
2000.....	0.9	1.1	1.4	2.0	2.5	3.4	5.4	7.8	11.5	21.9
1999.....	1.0	1.0	1.4	1.8	2.6	3.5	5.5	7.1	10.2	22.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of biliary tract and pancreas (K83-K86)										
White, male										
2002.....	1.2	1.7	1.9	2.8	3.6	5.5	7.2	10.5	14.5	21.2
2001.....	1.2	1.6	1.9	2.6	3.7	5.0	8.0	8.7	12.9	22.2
2000.....	1.3	1.5	1.8	2.7	2.9	4.7	6.6	10.2	13.2	22.0
1999.....	1.3	1.4	1.8	2.4	3.3	4.6	6.8	9.0	11.7	22.0
White, female										
2002.....	0.5	0.6	0.8	1.7	1.4	2.6	4.5	6.6	10.6	22.3
2001.....	0.6	0.9	1.0	1.3	2.3	3.3	4.3	6.8	11.5	23.4
2000.....	0.6	0.8	1.0	1.5	2.2	2.2	4.5	6.2	10.4	21.9
1999.....	0.6	0.6	1.1	1.3	1.9	2.6	4.4	5.7	9.3	22.0
Black, both sexes										
2002.....	1.8	2.9	3.2	3.8	5.3	5.0	7.1	6.7	11.5	21.4
2001.....	2.2	3.2	4.1	3.9	4.4	6.0	7.5	8.0	11.6	22.1
2000.....	2.0	3.1	2.8	4.5	4.6	6.7	7.7	8.7	12.1	23.5
1999.....	1.7	3.3	3.3	4.1	5.5	5.6	5.3	8.4	12.4	23.0
Black, male										
2002.....	2.1	4.5	4.7	5.7	7.3	6.0	11.7	*	*	25.7
2001.....	2.7	4.7	5.5	5.9	5.9	8.4	9.1	11.2	*	*
2000.....	3.0	3.9	4.4	6.3	5.6	10.4	9.7	10.9	*	32.3
1999.....	2.5	5.0	5.4	6.6	6.4	6.8	*	10.4	*	32.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of biliary tract and pancreas (K83-K86)										
Black, female										
2002.....	1.5	1.5	1.9	*	3.7	4.3	*	5.5	11.5	19.7
2001.....	1.8	2.0	2.9	*	3.2	4.3	6.5	6.1	13.5	22.3
2000.....	*	2.4	*	3.1	3.8	4.0	6.4	7.4	11.0	20.2
1999.....	*	1.8	*	*	4.8	4.8	5.2	7.2	13.0	19.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pancreatitis (K85,K86.0-K86.1)										
All races 3/, both sexes										
2002.....	0.9	1.2	1.3	1.9	2.2	2.9	4.0	5.7	7.8	11.9
2001.....	0.9	1.2	1.4	1.8	2.5	3.2	4.5	5.4	7.7	13.2
2000.....	0.9	1.1	1.3	1.8	2.3	2.8	4.0	5.4	7.6	12.1
1999.....	0.9	1.1	1.4	1.6	2.2	2.9	4.0	5.2	6.8	12.8
All races 3/, male										
2002.....	1.1	1.8	1.9	2.5	3.2	4.2	5.1	7.1	8.8	11.2
2001.....	1.2	1.7	1.8	2.5	3.1	3.9	6.1	6.1	7.8	12.9
2000.....	1.4	1.4	1.7	2.4	2.6	4.1	5.0	6.8	8.3	11.9
1999.....	1.3	1.6	1.8	2.2	2.9	3.8	5.1	6.4	7.5	12.7
All races 3/, female										
2002.....	0.6	0.6	0.8	1.4	1.2	1.8	3.1	4.7	7.1	12.2
2001.....	0.6	0.8	1.0	1.1	1.9	2.7	3.3	4.9	7.7	13.4
2000.....	0.5	0.8	0.8	1.2	1.9	1.8	3.2	4.4	7.2	12.2
1999.....	0.6	0.7	0.9	1.0	1.6	2.2	3.2	4.3	6.5	12.8
White, both sexes										
2002.....	0.8	1.0	1.1	1.9	1.9	2.9	4.0	5.8	7.9	12.1
2001.....	0.8	1.1	1.1	1.7	2.4	3.2	4.5	5.5	7.7	13.3
2000.....	0.8	0.9	1.2	1.6	2.1	2.6	3.9	5.3	7.7	12.2
1999.....	0.9	0.9	1.2	1.4	2.0	2.9	4.0	5.0	6.8	12.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pancreatitis (K85,K86.0-K86.1)										
White, male										
2002.....	1.1	1.5	1.6	2.4	2.8	4.3	5.1	7.2	8.9	11.0
2001.....	1.1	1.5	1.5	2.3	2.9	3.8	6.2	6.1	7.8	12.9
2000.....	1.2	1.2	1.5	2.1	2.4	3.7	4.9	6.7	8.3	11.5
1999.....	1.2	1.2	1.5	1.9	2.7	3.7	5.2	6.5	7.4	12.0
White, female										
2002.....	0.5	0.5	0.6	1.4	1.1	1.8	3.2	4.8	7.2	12.6
2001.....	0.5	0.7	0.8	1.1	1.9	2.6	3.1	5.0	7.7	13.4
2000.....	0.5	0.7	0.8	1.0	1.8	1.6	3.2	4.4	7.3	12.5
1999.....	0.5	0.6	0.9	1.0	1.3	2.1	3.1	4.1	6.4	13.1
Black, both sexes										
2002.....	1.4	2.5	2.8	3.0	4.7	3.5	3.9	4.3	6.5	11.3
2001.....	2.1	2.8	3.8	3.1	3.8	4.2	5.1	5.0	8.7	13.2
2000.....	1.9	2.7	2.4	3.9	3.8	5.2	5.6	5.3	7.6	11.9
1999.....	1.6	3.0	3.1	3.3	4.3	4.5	4.2	6.4	8.1	12.8
Black, male										
2002.....	1.8	3.9	4.2	4.5	6.9	*	*	*	*	*
2001.....	2.6	4.2	4.9	5.0	5.5	5.6	*	*	*	*
2000.....	3.0	3.2	3.8	5.4	4.8	8.3	7.7	*	*	*
1999.....	2.4	4.5	5.1	5.4	4.9	6.0	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pancreatitis (K85,K86.0-K86.1)										
Black, female										
2002.....	*	*	*	*	*	*	*	*	*	9.4
2001.....	1.6	1.7	2.9	*	*	*	4.7	*	9.4	13.9
2000.....	*	2.2	*	2.7	*	*	*	*	*	9.9
1999.....	*	1.6	*	*	3.8	*	*	6.0	*	10.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)

All races 3/, both sexes

2002.....	0.1	0.1	0.2	0.4	0.6	1.1	1.8	2.5	4.2	9.8
2001.....	*	0.2	0.3	0.3	0.6	1.0	1.5	2.2	4.2	9.7
2000.....	0.1	0.2	0.3	0.5	0.4	0.9	1.5	2.5	3.9	9.9
1999.....	0.1	0.1	0.2	0.5	0.6	0.7	1.4	2.0	3.5	9.4

All races 3/, male

2002.....	*	*	0.3	0.5	0.8	1.3	2.3	3.3	5.5	10.0
2001.....	*	0.2	0.4	0.3	0.7	1.3	1.9	2.6	4.7	9.5
2000.....	*	0.3	0.4	0.6	0.5	1.1	1.6	3.5	5.0	11.0
1999.....	*	0.2	0.3	0.6	0.7	0.8	1.6	2.7	4.2	10.5

All races 3/, female

2002.....	*	*	*	0.3	0.3	0.9	1.4	1.9	3.4	9.8
2001.....	*	0.2	*	0.3	0.5	0.7	1.2	1.9	3.9	9.8
2000.....	*	*	*	0.4	0.4	0.7	1.4	1.9	3.2	9.4
1999.....	*	*	*	0.4	0.6	0.6	1.2	1.6	3.1	9.0

White, both sexes

2002.....	*	0.1	0.2	0.4	0.5	1.0	1.7	2.4	4.2	9.9
2001.....	*	0.2	0.3	0.2	0.6	0.9	1.5	2.1	4.3	9.7
2000.....	*	0.2	0.2	0.5	0.4	0.8	1.5	2.5	3.8	9.7
1999.....	*	0.1	0.3	0.4	0.6	0.7	1.4	2.0	3.4	9.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of digestive system (K90,K92)										
All races 3/, both sexes										
2002.....	0.4	0.8	1.2	1.7	2.6	4.7	7.6	14.1	27.6	76.2
2001.....	0.4	0.7	1.0	1.8	3.1	4.8	7.7	14.1	28.2	77.8
2000.....	0.4	0.7	1.1	1.5	2.8	4.1	7.9	15.0	29.0	80.6
1999.....	0.4	0.7	1.0	1.7	2.8	4.7	7.7	14.1	27.8	77.2
All races 3/, male										
2002.....	0.6	1.2	1.6	2.3	3.8	6.1	9.6	17.8	31.0	76.2
2001.....	0.6	1.0	1.6	2.5	3.9	6.3	9.7	17.5	33.0	81.4
2000.....	0.6	1.0	1.6	2.0	3.8	5.2	9.2	18.3	36.5	83.1
1999.....	0.5	1.0	1.4	2.3	3.6	6.1	10.0	17.3	34.3	80.8
All races 3/, female										
2002.....	0.2	0.4	0.7	1.2	1.6	3.4	6.0	11.5	25.5	76.2
2001.....	0.2	0.4	0.5	1.1	2.4	3.6	6.1	11.8	25.3	76.3
2000.....	0.2	0.4	0.6	1.0	1.9	3.0	6.8	12.7	24.6	79.6
1999.....	0.2	0.4	0.6	1.2	2.1	3.6	6.0	11.8	24.0	75.8
White, both sexes										
2002.....	0.3	0.7	1.1	1.5	2.4	4.5	7.3	13.6	27.6	77.8
2001.....	0.3	0.6	0.9	1.6	2.8	4.5	7.5	13.7	28.4	79.1
2000.....	0.3	0.6	0.9	1.3	2.4	3.7	7.6	14.4	28.8	81.7
1999.....	0.3	0.6	0.8	1.6	2.5	4.4	7.5	13.6	27.6	78.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of digestive system (K90,K92)										
White, male										
2002.....	0.5	1.0	1.6	2.1	3.5	5.8	9.1	17.3	31.2	77.3
2001.....	0.5	0.9	1.3	2.1	3.5	5.8	9.3	16.7	33.5	82.8
2000.....	0.5	0.8	1.4	1.7	3.4	4.9	8.8	17.5	36.2	84.9
1999.....	0.5	0.9	1.2	2.0	3.4	5.7	9.6	16.6	34.4	82.1
White, female										
2002.....	*	0.4	0.6	1.0	1.4	3.3	5.8	11.0	25.4	78.1
2001.....	*	0.4	0.5	1.0	2.1	3.4	6.0	11.5	25.3	77.5
2000.....	*	0.3	0.5	0.8	1.6	2.7	6.6	12.2	24.4	80.5
1999.....	*	0.3	0.5	1.1	1.8	3.3	5.8	11.4	23.6	77.6
Black, both sexes										
2002.....	0.8	1.6	2.0	3.6	5.3	7.8	12.2	21.6	33.5	65.3
2001.....	0.9	1.3	2.2	4.2	6.4	8.5	11.2	22.1	31.4	73.1
2000.....	0.9	1.6	2.5	3.9	6.3	7.2	12.9	22.6	36.7	77.9
1999.....	*	1.2	2.2	3.1	5.5	7.8	12.2	22.0	33.2	66.0
Black, male										
2002.....	*	2.7	2.5	4.5	7.3	11.2	17.0	26.2	36.7	70.8
2001.....	*	2.2	3.7	6.3	7.9	11.4	16.5	30.2	36.1	71.5
2000.....	*	2.4	3.3	4.9	8.7	7.8	17.4	29.3	47.2	78.4
1999.....	*	*	3.3	5.3	6.2	9.9	16.6	29.0	38.4	71.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of digestive system (K90,K92)										
Black, female										
2002.....	*	*	*	2.8	3.7	5.2	9.0	18.9	31.8	63.3
2001.....	*	*	*	*	5.3	6.4	7.6	17.2	29.0	73.7
2000.....	*	*	*	3.0	4.4	6.7	10.0	18.5	31.4	77.7
1999.....	*	*	*	*	5.0	6.3	9.2	17.7	30.7	64.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XII. Diseases of the skin and subcutaneous tissue (L00-L98)										
All races 3/, both sexes										
2002.....	0.4	0.5	0.7	1.1	1.4	2.0	4.2	6.5	11.8	32.8
2001.....	0.3	0.5	0.7	1.0	1.4	2.2	3.5	6.5	12.0	32.2
2000.....	0.3	0.4	0.5	0.9	1.5	2.1	4.0	6.4	12.7	34.7
1999.....	0.3	0.5	0.6	0.8	1.3	2.2	3.5	6.6	12.7	35.0
All races 3/, male										
2002.....	0.3	0.5	0.7	1.1	1.2	2.0	4.5	6.5	11.9	28.6
2001.....	0.2	0.6	0.7	1.0	1.4	2.5	3.6	6.7	10.8	24.9
2000.....	0.3	0.4	0.4	1.0	1.6	1.6	4.0	6.1	12.3	29.7
1999.....	0.2	0.5	0.6	0.8	1.1	2.3	3.5	6.1	12.2	31.7
All races 3/, female										
2002.....	0.4	0.4	0.7	1.1	1.5	2.1	3.9	6.6	11.7	34.7
2001.....	0.3	0.5	0.6	1.0	1.5	1.9	3.4	6.4	12.7	35.2
2000.....	0.3	0.5	0.5	0.9	1.3	2.4	3.9	6.6	13.0	36.7
1999.....	0.4	0.5	0.6	0.8	1.4	2.2	3.6	7.0	12.9	36.3
White, both sexes										
2002.....	0.3	0.4	0.5	0.9	1.0	1.7	3.3	5.6	10.0	28.3
2001.....	0.2	0.4	0.6	0.8	1.2	1.8	3.0	5.6	10.4	28.1
2000.....	0.2	0.3	0.4	0.8	1.2	1.7	3.1	5.2	11.0	29.7
1999.....	0.2	0.4	0.4	0.6	1.0	1.8	2.9	5.3	10.8	29.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XII. Diseases of the skin and subcutaneous tissue (L00-L98)										
White, male										
2002.....	0.3	0.5	0.5	1.0	0.9	1.6	3.6	5.7	10.7	26.1
2001.....	*	0.5	0.5	0.9	1.2	2.2	3.3	5.5	9.4	21.8
2000.....	*	0.3	0.4	0.8	1.4	1.2	3.1	4.9	10.4	25.8
1999.....	*	0.3	0.4	0.6	0.9	1.9	2.8	4.7	10.4	27.8
White, female										
2002.....	0.3	0.3	0.5	0.9	1.1	1.8	3.1	5.5	9.6	29.3
2001.....	0.2	0.4	0.6	0.8	1.1	1.5	2.8	5.6	11.0	30.8
2000.....	0.2	0.2	0.4	0.7	1.1	2.1	3.2	5.5	11.4	31.2
1999.....	0.3	0.4	0.4	0.6	1.0	1.7	3.0	5.6	11.0	30.0
Black, both sexes										
2002.....	1.1	1.0	2.0	3.2	4.4	5.1	12.5	19.1	35.3	92.9
2001.....	0.8	1.3	1.5	2.5	3.8	5.9	9.0	19.1	33.9	88.4
2000.....	0.8	1.4	1.1	2.5	3.7	6.0	12.5	19.9	36.1	97.3
1999.....	1.0	1.7	2.0	2.9	4.2	6.7	11.5	23.2	38.7	105.2
Black, male										
2002.....	*	*	2.2	*	*	6.0	15.4	18.4	32.9	63.3
2001.....	*	*	*	*	*	6.6	8.1	24.6	34.5	71.5
2000.....	*	*	*	*	*	6.2	15.0	20.3	41.3	81.9
1999.....	*	2.2	*	*	*	6.8	12.2	24.2	39.2	85.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XII. Diseases of the skin and subcutaneous tissue (L00-L98)

Black, female

2002.....	1.4	*	1.8	3.7	5.1	4.4	10.5	19.5	36.6	104.3
2001.....	*	*	*	3.0	4.2	5.4	9.6	15.8	33.5	94.7
2000.....	*	2.0	*	2.7	3.6	5.8	10.9	19.7	33.5	103.0
1999.....	*	*	*	2.9	4.7	6.7	11.0	22.6	38.5	112.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Infections of skin and subcutaneous tissue (L00-L08)										
All races 3/, both sexes										
2002.....	0.1	0.2	0.3	0.5	0.6	0.8	1.3	2.2	3.6	9.3
2001.....	0.1	0.3	0.3	0.4	0.7	0.9	1.2	2.0	3.4	9.4
2000.....	0.1	0.1	0.2	0.4	0.6	0.7	1.4	2.0	3.4	9.3
1999.....	*	0.2	0.2	0.3	0.5	0.7	1.1	1.5	2.9	8.3
All races 3/, male										
2002.....	0.2	0.3	0.4	0.6	0.6	0.9	1.5	2.2	4.1	9.1
2001.....	*	0.4	0.4	0.5	0.8	1.1	1.4	2.2	2.9	8.1
2000.....	*	*	*	0.6	0.7	0.7	1.6	2.1	4.0	8.9
1999.....	*	0.3	0.3	0.4	0.4	0.7	1.0	1.3	3.3	9.1
All races 3/, female										
2002.....	*	*	0.2	0.4	0.5	0.8	1.2	2.2	3.2	9.4
2001.....	*	*	0.3	0.4	0.6	0.7	1.0	1.9	3.8	10.0
2000.....	*	*	*	0.3	0.5	0.7	1.2	1.9	3.1	9.5
1999.....	*	*	*	*	0.5	0.7	1.1	1.6	2.7	8.0
White, both sexes										
2002.....	0.1	0.2	0.3	0.5	0.6	0.7	1.2	2.1	3.6	9.7
2001.....	*	0.2	0.3	0.4	0.6	0.9	1.2	2.0	3.4	9.6
2000.....	*	0.1	0.2	0.4	0.6	0.7	1.3	1.9	3.4	9.3
1999.....	*	0.2	0.2	0.3	0.4	0.7	1.1	1.5	3.0	8.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)										
All races 3/, both sexes										
2002.....	0.2	0.3	0.4	0.6	0.8	1.2	2.8	4.3	8.2	23.5
2001.....	0.2	0.3	0.3	0.6	0.8	1.3	2.3	4.5	8.5	22.7
2000.....	0.2	0.3	0.3	0.5	0.8	1.4	2.6	4.4	9.3	25.4
1999.....	0.2	0.3	0.4	0.5	0.8	1.5	2.5	5.1	9.7	26.6
All races 3/, male										
2002.....	*	0.2	0.3	0.5	0.5	1.1	3.0	4.3	7.8	19.4
2001.....	*	0.2	0.3	0.5	0.6	1.4	2.2	4.5	7.9	16.8
2000.....	0.2	*	*	0.4	0.9	0.9	2.4	4.0	8.3	20.8
1999.....	*	0.2	0.3	0.4	0.7	1.6	2.4	4.8	8.9	22.6
All races 3/, female										
2002.....	0.3	0.3	0.5	0.7	1.0	1.3	2.7	4.3	8.5	25.3
2001.....	0.2	0.3	0.4	0.6	0.9	1.2	2.4	4.5	8.9	25.2
2000.....	0.2	0.4	0.4	0.6	0.8	1.7	2.7	4.7	9.9	27.3
1999.....	0.3	0.3	0.4	0.6	0.8	1.5	2.5	5.4	10.2	28.3
White, both sexes										
2002.....	0.2	0.2	0.2	0.4	0.5	1.0	2.1	3.5	6.4	18.7
2001.....	0.1	0.2	0.3	0.4	0.6	1.0	1.8	3.5	7.0	18.5
2000.....	0.1	0.2	0.2	0.3	0.6	1.0	1.8	3.3	7.6	20.4
1999.....	0.1	0.2	0.2	0.3	0.5	1.1	1.8	3.8	7.7	21.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)

White, male

2002.....	*	*	*	0.4	*	0.8	2.2	3.6	6.4	16.7
2001.....	*	*	*	0.4	0.4	1.1	1.8	3.3	6.4	13.3
2000.....	*	*	*	*	0.7	0.6	1.5	2.9	6.5	17.0
1999.....	*	*	*	*	0.5	1.1	1.7	3.6	7.0	18.9

White, female

2002.....	0.2	*	0.3	0.5	0.6	1.1	2.0	3.4	6.5	19.6
2001.....	*	0.2	0.3	0.5	0.7	0.9	1.8	3.6	7.4	20.6
2000.....	*	*	0.3	0.4	0.6	1.4	2.0	3.6	8.3	21.7
1999.....	0.2	0.3	0.3	0.4	0.6	1.1	1.9	3.9	8.1	21.9

Black, both sexes

2002.....	0.7	*	1.4	2.1	3.6	3.3	10.3	15.3	31.7	85.8
2001.....	*	*	1.0	1.7	2.6	4.5	8.2	16.8	29.3	79.2
2000.....	*	1.1	*	1.8	2.9	4.6	10.9	17.2	32.2	87.3
1999.....	0.8	0.9	1.5	1.9	3.1	5.9	10.1	22.0	36.4	96.3

Black, male

2002.....	*	*	*	*	*	*	12.7	15.2	29.8	56.8
2001.....	*	*	*	*	*	5.3	7.4	21.4	32.1	65.8
2000.....	*	*	*	*	*	*	12.7	18.0	36.2	72.7
1999.....	*	*	*	*	*	6.8	11.2	21.4	35.8	73.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)

Black, female

2002.....	*	*	*	2.7	4.0	*	8.7	15.4	32.6	96.9
2001.....	*	*	*	*	*	3.9	8.7	14.1	27.8	84.2
2000.....	*	1.7	*	*	*	4.8	9.8	16.8	30.1	92.7
1999.....	*	*	*	*	*	5.2	9.5	22.3	36.7	104.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)										
All races 3/, both sexes										
2002.....	1.5	2.3	3.0	4.3	7.0	10.1	15.9	25.6	40.2	94.9
2001.....	1.7	2.1	3.1	4.2	6.6	10.1	15.8	25.4	42.0	93.4
2000.....	1.7	2.3	3.0	4.3	6.7	10.3	15.2	25.9	41.8	94.0
1999.....	1.6	2.4	3.1	4.3	6.7	10.1	15.5	25.5	39.3	91.2
All races 3/, male										
2002.....	1.0	1.7	2.3	3.0	5.3	7.8	13.0	19.4	31.3	63.5
2001.....	1.1	1.5	2.4	3.2	4.7	7.1	12.3	20.1	32.4	63.2
2000.....	1.2	1.6	1.8	3.4	4.8	7.7	12.1	20.6	33.0	67.5
1999.....	1.1	1.5	2.1	3.1	4.7	6.9	12.5	20.2	30.9	63.7
All races 3/, female										
2002.....	2.1	3.0	3.8	5.6	8.6	12.2	18.2	30.1	45.6	108.5
2001.....	2.2	2.8	3.7	5.2	8.3	12.7	18.5	29.1	47.8	106.0
2000.....	2.2	3.0	4.1	5.2	8.4	12.4	17.6	29.6	46.9	104.8
1999.....	2.1	3.3	4.1	5.5	8.5	12.8	18.0	29.3	44.3	102.3
White, both sexes										
2002.....	1.1	1.8	2.7	3.9	6.6	9.9	15.8	25.9	40.7	98.2
2001.....	1.2	1.7	2.6	3.9	6.1	9.9	15.5	25.8	43.0	95.8
2000.....	1.2	1.7	2.5	3.8	6.3	10.2	15.0	26.0	42.5	96.9
1999.....	1.1	1.8	2.6	3.8	6.4	9.7	15.6	25.8	40.6	94.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)										
White, male										
2002.....	0.8	1.4	2.1	2.7	5.0	7.3	12.7	19.4	31.0	64.3
2001.....	0.8	1.2	2.1	2.8	4.4	6.8	11.9	20.3	32.7	64.3
2000.....	0.9	1.3	1.6	3.2	4.5	7.6	11.3	20.2	32.9	67.3
1999.....	0.8	1.2	1.9	2.7	4.6	6.6	12.6	19.4	31.9	64.4
White, female										
2002.....	1.4	2.3	3.2	5.0	8.1	12.2	18.4	30.6	46.6	112.9
2001.....	1.6	2.2	3.1	4.8	7.6	12.6	18.4	29.6	49.3	109.0
2000.....	1.5	2.0	3.4	4.4	7.9	12.6	17.9	30.1	48.2	108.9
1999.....	1.4	2.5	3.3	4.9	8.0	12.4	18.1	30.2	45.7	106.5
Black, both sexes										
2002.....	4.5	6.0	6.5	8.9	11.4	13.5	18.8	25.6	39.0	67.4
2001.....	4.9	5.5	7.5	8.0	11.2	14.1	19.6	25.0	36.0	75.5
2000.....	5.4	7.2	6.9	8.9	11.1	12.4	18.7	26.8	37.6	68.2
1999.....	5.4	6.8	7.3	8.9	10.5	15.3	17.5	27.0	27.2	60.3
Black, male										
2002.....	2.5	3.9	4.3	6.7	9.1	12.5	18.7	21.6	39.7	61.1
2001.....	3.4	4.2	5.7	6.6	9.2	11.7	17.2	22.8	36.9	59.0
2000.....	3.6	4.2	3.7	6.3	8.3	12.0	21.7	25.5	39.6	78.4
1999.....	3.4	4.0	4.2	6.4	6.8	12.2	14.2	33.7	20.9	63.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

Black, female

2002.....	6.3	7.8	8.4	10.6	13.2	14.2	18.8	28.0	38.6	69.8
2001.....	6.3	6.6	8.9	9.1	12.8	16.0	21.3	26.3	35.6	81.7
2000.....	7.1	9.8	9.6	11.1	13.2	12.7	16.6	27.6	36.5	64.4
1999.....	7.2	9.3	10.0	10.9	13.5	17.5	19.6	22.9	30.2	59.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)										
All races 3/, both sexes										
2002.....	*	0.2	0.3	0.7	1.4	2.5	4.2	8.0	10.6	14.2
2001.....	*	0.1	0.3	0.7	1.4	2.5	4.3	7.6	10.5	12.7
2000.....	*	0.2	0.3	0.6	1.4	2.7	4.4	7.5	11.6	12.9
1999.....	*	0.2	0.3	0.8	1.6	2.5	4.5	8.7	10.6	13.9
All races 3/, male										
2002.....	*	*	*	0.5	1.0	1.7	2.6	4.5	6.6	7.6
2001.....	*	*	*	0.6	0.9	1.5	2.6	4.6	5.8	5.5
2000.....	*	*	*	0.4	0.9	1.7	3.1	4.4	6.6	6.9
1999.....	*	*	*	0.5	1.1	1.6	3.1	5.0	5.5	7.2
All races 3/, female										
2002.....	*	0.2	0.4	0.9	1.8	3.2	5.4	10.4	13.1	17.0
2001.....	*	0.2	0.5	0.9	1.8	3.3	5.5	9.7	13.3	15.7
2000.....	*	0.3	0.5	0.8	1.9	3.5	5.4	9.7	14.6	15.3
1999.....	*	0.3	0.3	1.0	2.0	3.3	5.6	11.3	13.5	16.7
White, both sexes										
2002.....	*	0.2	0.3	0.7	1.4	2.6	4.3	8.2	10.9	14.8
2001.....	*	0.1	0.3	0.7	1.4	2.7	4.4	8.0	10.9	12.8
2000.....	*	0.2	0.3	0.6	1.5	2.8	4.7	7.8	12.2	13.4
1999.....	*	0.2	0.3	0.8	1.7	2.6	4.8	9.0	11.0	14.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)

Black, female

2002.....	*	*	*	*	*	*	*	7.1	9.2	9.9
2001.....	*	*	*	*	*	*	*	*	*	14.7
2000.....	*	*	*	*	*	*	*	6.0	*	9.4
1999.....	*	*	*	*	*	*	*	6.3	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Systemic lupus erythematosus (M32)										
All races 3/, both sexes										
2002.....	0.5	0.6	0.6	0.7	1.0	0.9	1.0	1.3	1.4	1.3
2001.....	0.5	0.6	0.7	0.7	0.9	0.9	1.2	1.4	1.6	1.2
2000.....	0.6	0.6	0.7	0.8	1.0	1.0	0.9	1.4	1.5	1.2
1999.....	0.6	0.7	0.7	0.7	0.8	1.0	1.2	1.3	1.3	1.4
All races 3/, male										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	0.2	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	0.8	1.1	0.9	1.2	1.7	1.4	1.5	1.7	1.9	1.7
2001.....	0.8	1.1	1.1	1.1	1.5	1.5	1.8	2.2	2.0	1.3
2000.....	1.1	1.1	1.2	1.3	1.5	1.5	1.2	2.0	2.0	1.4
1999.....	1.0	1.2	1.2	1.2	1.4	1.7	1.8	1.8	1.7	1.6
White, both sexes										
2002.....	0.3	0.4	0.4	0.5	0.7	0.7	1.0	1.3	1.4	1.4
2001.....	0.2	0.4	0.5	0.6	0.7	0.8	1.1	1.4	1.7	1.2
2000.....	0.4	0.3	0.5	0.6	0.8	0.9	0.8	1.4	1.6	1.3
1999.....	0.3	0.5	0.5	0.6	0.7	0.8	1.1	1.3	1.3	1.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Systemic lupus erythematosus (M32)										
Black, female										
2002.....	3.1	4.0	2.7	3.5	5.1	4.4	*	*	*	*
2001.....	3.4	3.3	3.2	2.7	3.5	*	*	*	*	*
2000.....	3.6	4.4	3.4	3.6	3.8	*	*	*	*	*
1999.....	3.9	3.9	3.3	3.2	*	4.8	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)										
All races 3/, both sexes										
2002.....	0.7	0.9	1.4	1.7	2.9	3.9	5.6	8.1	11.9	27.2
2001.....	0.7	0.9	1.3	1.7	2.7	4.1	5.9	8.2	13.0	27.7
2000.....	0.7	1.0	1.3	1.8	2.8	4.0	6.0	8.7	13.1	28.0
1999.....	0.7	1.1	1.5	1.8	2.9	4.2	6.0	7.9	11.9	27.6
All races 3/, male										
2002.....	0.4	0.7	0.9	1.3	2.0	2.5	4.2	6.3	9.6	18.6
2001.....	0.5	0.8	1.0	1.1	2.0	2.3	4.6	6.5	10.6	20.5
2000.....	0.6	0.7	0.8	1.4	1.9	2.8	4.0	7.3	11.2	21.1
1999.....	0.5	0.7	1.1	1.3	1.9	2.8	4.4	6.5	10.0	22.5
All races 3/, female										
2002.....	0.9	1.2	1.9	2.2	3.6	5.1	6.7	9.4	13.3	31.0
2001.....	1.0	1.1	1.5	2.2	3.3	5.6	7.0	9.4	14.4	30.7
2000.....	0.9	1.2	1.7	2.2	3.5	5.1	7.5	9.6	14.3	30.8
1999.....	0.8	1.4	1.9	2.3	3.8	5.4	7.2	8.8	13.0	29.7
White, both sexes										
2002.....	0.5	0.7	1.2	1.6	2.8	4.0	5.5	8.3	12.1	28.0
2001.....	0.5	0.7	1.0	1.6	2.5	3.9	5.9	8.5	13.4	28.6
2000.....	0.5	0.7	1.0	1.6	2.6	4.1	5.9	8.8	13.4	28.8
1999.....	0.5	0.7	1.1	1.6	2.6	4.1	5.9	8.0	12.2	28.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)										
White, male										
2002.....	0.3	0.5	0.8	1.1	1.9	2.5	4.0	6.5	9.7	19.1
2001.....	0.3	0.5	0.8	1.0	1.9	2.2	4.5	6.7	10.8	21.1
2000.....	0.4	0.6	0.7	1.3	1.7	2.7	3.8	7.2	11.1	20.6
1999.....	0.3	0.5	0.9	1.1	1.8	2.7	4.4	6.4	10.4	23.3
White, female										
2002.....	0.6	0.9	1.6	2.0	3.7	5.3	6.8	9.6	13.6	31.9
2001.....	0.8	0.9	1.2	2.1	3.1	5.5	7.0	9.8	14.9	31.7
2000.....	0.6	0.8	1.4	1.9	3.4	5.3	7.5	10.0	14.7	32.1
1999.....	0.6	1.0	1.4	2.1	3.4	5.3	7.2	9.2	13.3	30.4
Black, both sexes										
2002.....	2.1	2.5	3.0	3.6	4.1	4.4	7.0	7.4	10.7	21.7
2001.....	2.2	2.7	3.6	2.7	3.7	6.0	6.1	6.4	11.6	22.4
2000.....	2.3	3.0	3.3	3.6	4.4	4.0	7.9	7.5	12.1	21.9
1999.....	2.4	3.3	4.1	3.7	5.9	5.3	7.0	7.3	8.7	20.4
Black, male										
2002.....	*	2.0	2.2	3.6	4.2	*	*	*	*	*
2001.....	1.6	2.6	2.9	*	*	*	*	*	*	*
2000.....	1.8	*	*	*	4.1	*	*	*	*	27.7
1999.....	2.0	2.3	2.5	*	*	*	*	9.5	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)										
Black, female										
2002.....	2.8	3.0	3.8	3.5	4.0	4.8	7.4	8.8	11.9	24.6
2001.....	2.7	2.8	4.2	3.3	4.5	7.3	7.2	6.7	12.7	24.4
2000.....	2.8	4.2	4.5	4.3	4.6	4.0	8.9	7.1	10.6	19.7
1999.....	2.7	4.2	5.5	4.3	7.3	6.3	8.1	6.0	10.4	22.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Dorsopathies (M40-M54)										
All races 3/, both sexes										
2002.....	*	0.1	0.2	0.3	0.4	0.6	1.2	1.7	2.9	6.0
2001.....	*	*	0.2	0.3	0.3	0.5	1.0	1.7	3.3	7.4
2000.....	*	0.1	0.1	0.2	0.4	0.7	0.9	1.7	3.3	6.1
1999.....	*	0.1	0.2	0.2	0.3	0.6	0.9	1.5	2.5	5.4
All races 3/, male										
2002.....	*	*	0.3	0.3	0.4	0.5	1.3	2.0	3.1	5.5
2001.....	*	*	*	0.3	0.4	0.6	0.9	1.9	3.5	6.4
2000.....	*	*	*	*	0.4	0.8	1.2	1.5	3.8	6.5
1999.....	*	*	*	*	0.5	0.7	1.0	1.8	2.8	5.9
All races 3/, female										
2002.....	*	*	*	0.3	0.4	0.7	1.1	1.5	2.8	6.2
2001.....	*	*	*	*	*	0.5	1.1	1.6	3.2	7.8
2000.....	*	*	*	*	*	0.5	0.7	1.9	3.1	6.0
1999.....	*	*	0.2	*	*	0.5	0.9	1.3	2.4	5.2
White, both sexes										
2002.....	*	0.1	0.2	0.3	0.4	0.7	1.2	1.7	3.0	6.2
2001.....	*	*	0.2	0.2	0.3	0.5	1.0	1.7	3.3	7.8
2000.....	*	0.1	*	0.2	0.4	0.7	0.8	1.7	3.3	6.4
1999.....	*	*	0.2	0.2	0.3	0.5	0.9	1.5	2.7	5.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Soft tissue disorders (M60-M79)										
All races 3/, both sexes										
2002.....	0.2	0.3	0.4	0.5	0.6	0.9	1.3	1.9	3.1	5.3
2001.....	0.2	0.2	0.4	0.5	0.7	0.9	1.2	2.0	2.8	4.3
2000.....	0.2	0.4	0.4	0.5	0.6	0.8	1.2	2.0	2.3	4.2
1999.....	0.2	0.2	0.3	0.5	0.5	0.8	1.2	1.7	2.7	4.3
All races 3/, male										
2002.....	0.3	0.4	0.4	0.4	0.8	1.3	1.6	2.0	4.3	7.8
2001.....	0.3	0.3	0.6	0.6	0.6	1.1	1.6	2.7	3.9	6.2
2000.....	0.2	0.4	0.3	0.6	0.5	0.9	1.6	2.6	3.5	6.7
1999.....	0.3	0.3	0.3	0.6	0.5	0.8	1.7	2.5	4.0	6.8
All races 3/, female										
2002.....	*	*	0.3	0.6	0.5	0.6	1.1	1.8	2.4	4.2
2001.....	*	*	0.3	0.5	0.7	0.6	0.9	1.4	2.1	3.5
2000.....	*	0.3	0.4	0.4	0.6	0.8	0.9	1.6	1.7	3.2
1999.....	*	0.2	0.3	0.4	0.5	0.8	0.7	1.2	2.0	3.3
White, both sexes										
2002.....	0.2	0.2	0.4	0.4	0.6	0.9	1.2	1.8	3.1	5.0
2001.....	0.2	0.2	0.4	0.5	0.5	0.8	1.1	1.8	2.7	4.2
2000.....	0.1	0.3	0.3	0.5	0.5	0.8	1.1	1.7	2.3	4.0
1999.....	0.2	0.2	0.3	0.4	0.4	0.7	1.1	1.7	2.6	4.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)										
All races 3/, both sexes										
2002.....	*	0.1	0.2	0.4	0.8	1.2	2.6	4.7	10.2	40.8
2001.....	*	0.2	0.2	0.3	0.7	1.3	2.2	4.4	10.8	40.1
2000.....	*	*	0.2	0.3	0.6	1.1	1.8	4.6	9.8	41.6
1999.....	*	0.1	0.2	0.4	0.5	1.0	1.8	4.4	10.3	38.6
All races 3/, male										
2002.....	*	*	0.3	0.4	0.9	1.4	2.9	4.1	7.2	23.6
2001.....	*	*	0.2	0.5	0.7	1.3	2.1	4.0	7.7	23.6
2000.....	*	*	*	0.4	0.6	1.2	1.8	4.2	7.1	25.6
1999.....	*	*	*	0.3	0.5	0.8	1.8	3.7	7.8	20.4
All races 3/, female										
2002.....	*	*	*	0.4	0.7	1.1	2.4	5.2	12.0	48.3
2001.....	*	*	*	*	0.7	1.3	2.2	4.8	12.7	47.0
2000.....	*	*	*	*	0.5	0.9	1.9	4.8	11.3	48.1
1999.....	*	*	*	0.4	0.6	1.2	1.7	4.8	11.8	45.9
White, both sexes										
2002.....	*	0.1	0.2	0.4	0.8	1.1	2.7	4.6	10.1	42.8
2001.....	*	0.1	0.2	0.3	0.7	1.2	2.0	4.5	11.0	41.3
2000.....	*	*	0.2	0.3	0.5	1.0	1.8	4.6	9.8	43.0
1999.....	*	0.1	0.2	0.3	0.5	0.9	1.7	4.2	10.7	40.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)										
Black, female										
2002.....	*	*	*	*	*	*	*	6.3	10.3	23.4
2001.....	*	*	*	*	*	*	4.7	*	9.4	31.2
2000.....	*	*	*	*	*	*	*	*	10.2	25.8
1999.....	*	*	*	*	*	*	*	5.7	8.6	20.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Osteoporosis (M80-M81)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	0.2	0.6	1.9	5.4	29.0
2001.....	*	*	*	*	*	0.3	0.5	1.7	5.6	26.8
2000.....	*	*	*	*	*	0.3	0.4	2.1	5.5	28.5
1999.....	*	*	*	*	*	0.3	0.5	2.0	5.8	27.1
All races 3/, male										
2002.....	*	*	*	*	*	*	*	0.7	1.7	9.9
2001.....	*	*	*	*	*	*	*	*	1.6	9.3
2000.....	*	*	*	*	*	*	*	0.9	1.9	9.9
1999.....	*	*	*	*	*	*	*	0.7	2.2	7.0
All races 3/, female										
2002.....	*	*	*	*	*	*	1.0	2.9	7.6	37.2
2001.....	*	*	*	*	*	0.4	0.6	2.4	8.0	34.1
2000.....	*	*	*	*	*	0.4	0.6	2.9	7.7	36.0
1999.....	*	*	*	*	*	0.4	0.7	2.9	7.9	35.1
White, both sexes										
2002.....	*	*	*	*	*	*	0.7	2.0	5.8	31.3
2001.....	*	*	*	*	*	0.3	0.5	1.8	6.0	28.8
2000.....	*	*	*	*	*	0.3	0.4	2.3	5.9	30.7
1999.....	*	*	*	*	*	0.3	0.5	2.2	6.2	29.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)										
All races 3/, both sexes										
2002.....	*	0.1	0.2	0.4	0.7	1.0	2.0	2.8	4.8	11.8
2001.....	*	0.2	0.2	0.3	0.6	1.0	1.7	2.8	5.3	13.3
2000.....	*	*	0.2	0.3	0.5	0.8	1.4	2.5	4.3	13.1
1999.....	*	0.1	0.2	0.3	0.4	0.7	1.3	2.4	4.6	11.5
All races 3/, male										
2002.....	*	*	0.3	0.4	0.9	1.3	2.6	3.4	5.5	13.7
2001.....	*	*	0.2	0.4	0.7	1.2	1.8	3.3	6.1	14.2
2000.....	*	*	*	0.4	0.6	1.2	1.6	3.4	5.2	15.6
1999.....	*	*	*	0.3	0.5	0.7	1.7	3.0	5.7	13.3
All races 3/, female										
2002.....	*	*	*	0.4	0.6	0.7	1.4	2.3	4.4	11.1
2001.....	*	*	*	*	0.5	0.8	1.7	2.4	4.8	12.9
2000.....	*	*	*	*	0.4	0.5	1.3	1.9	3.7	12.0
1999.....	*	*	*	0.3	0.4	0.7	1.0	1.9	3.9	10.8
White, both sexes										
2002.....	*	0.1	0.2	0.3	0.7	0.8	2.0	2.5	4.4	11.5
2001.....	*	0.1	0.2	0.3	0.5	0.9	1.5	2.6	5.0	12.5
2000.....	*	*	0.1	0.3	0.4	0.7	1.3	2.3	3.8	12.4
1999.....	*	0.1	0.2	0.3	0.5	0.6	1.2	2.0	4.5	11.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XIV. Diseases of the genitourinary system (N00-N98)										
All races 3/, both sexes										
2002.....	2.6	4.5	7.1	12.2	20.5	36.5	64.2	115.2	206.9	469.4
2001.....	2.6	4.5	7.1	12.1	20.6	39.2	64.4	113.7	202.0	476.4
2000.....	2.6	4.3	6.8	11.5	20.7	36.4	60.5	111.8	202.9	471.4
1999.....	2.5	3.9	6.4	11.2	19.8	35.5	61.7	109.5	197.7	471.1
All races 3/, male										
2002.....	2.7	5.1	8.1	12.9	22.3	40.4	73.7	134.0	246.8	553.0
2001.....	2.7	5.0	7.8	12.9	21.7	44.0	71.6	129.8	242.7	564.2
2000.....	2.9	5.1	7.6	12.6	22.1	39.6	67.3	131.0	243.1	567.5
1999.....	2.7	4.5	6.9	12.1	22.0	38.3	70.3	130.4	241.8	558.3
All races 3/, female										
2002.....	2.4	4.0	6.2	11.6	18.9	33.1	56.7	102.0	182.6	433.2
2001.....	2.4	3.9	6.3	11.4	19.6	35.0	58.6	102.4	177.6	439.6
2000.....	2.2	3.4	6.0	10.4	19.4	33.7	55.2	98.5	179.1	432.3
1999.....	2.3	3.3	5.9	10.4	17.9	33.1	54.9	95.0	171.9	436.0
White, both sexes										
2002.....	1.9	3.3	5.0	9.4	16.4	30.9	56.8	106.8	198.9	458.5
2001.....	1.9	3.1	5.1	9.2	16.2	32.5	57.2	105.3	192.3	465.1
2000.....	1.9	3.0	4.9	8.7	16.5	30.2	52.8	103.9	193.4	460.5
1999.....	1.7	2.7	4.6	8.8	15.8	29.3	54.0	101.0	187.8	456.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XIV. Diseases of the genitourinary system (N00-N98)										
White, male										
2002.....	2.0	3.6	5.6	9.8	17.5	34.7	65.9	125.5	240.2	543.2
2001.....	2.0	3.4	5.8	9.8	17.5	37.1	64.8	122.6	233.9	556.8
2000.....	2.1	3.5	5.4	9.5	17.9	33.7	60.4	123.0	234.8	561.2
1999.....	1.8	3.0	4.9	9.9	17.5	32.0	62.2	120.6	230.8	542.8
White, female										
2002.....	1.8	2.9	4.4	9.0	15.4	27.5	49.3	93.5	173.5	421.7
2001.....	1.7	2.8	4.5	8.6	14.9	28.5	51.0	93.0	167.1	426.7
2000.....	1.6	2.6	4.4	7.9	15.3	27.0	46.7	90.5	168.8	419.6
1999.....	1.6	2.3	4.2	7.7	14.1	27.0	47.5	87.2	162.4	421.1
Black, both sexes										
2002.....	7.6	13.6	24.1	37.9	58.7	91.9	150.1	225.4	332.3	662.5
2001.....	7.6	14.5	23.0	36.9	60.3	102.5	146.3	223.5	347.5	664.5
2000.....	7.5	14.0	22.6	36.6	58.1	96.6	145.9	219.5	343.1	637.5
1999.....	7.9	13.5	22.7	32.7	56.2	94.5	147.3	219.9	337.2	679.1
Black, male										
2002.....	8.3	16.4	29.0	44.4	70.1	101.7	177.3	262.7	382.1	794.4
2001.....	7.8	18.3	25.9	42.9	63.3	118.4	158.5	241.6	404.5	755.7
2000.....	8.9	18.8	27.3	42.8	63.2	102.3	155.8	259.4	398.7	716.2
1999.....	9.2	17.3	26.4	32.9	65.8	102.0	172.3	270.7	433.4	802.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XIV. Diseases of the genitourinary system (N00-N98)										
Black, female										
2002.....	7.0	11.1	20.0	32.6	49.9	84.5	132.0	203.1	306.4	612.0
2001.....	7.4	11.3	20.5	32.1	57.9	90.7	138.1	212.7	318.5	630.6
2000.....	6.3	9.9	18.6	31.6	54.1	92.4	139.4	195.4	315.0	608.3
1999.....	6.7	10.2	19.5	32.5	48.8	88.9	130.7	189.4	289.5	633.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of urinary system (N00-N39)										
All races 3/, both sexes										
2002.....	2.5	4.4	7.0	12.1	20.3	36.1	63.5	113.9	203.4	460.8
2001.....	2.5	4.3	6.9	11.9	20.4	38.7	63.3	112.0	198.9	467.4
2000.....	2.5	4.2	6.6	11.3	20.4	35.9	59.7	110.2	199.3	462.5
1999.....	2.4	3.8	6.3	11.0	19.5	35.0	60.8	107.8	194.1	462.6
All races 3/, male										
2002.....	2.7	5.0	8.0	12.8	21.9	39.8	72.4	132.0	239.9	533.8
2001.....	2.7	5.0	7.7	12.7	21.5	43.5	70.2	127.2	237.2	542.9
2000.....	2.9	5.1	7.5	12.4	21.7	39.0	66.0	128.5	236.8	546.9
1999.....	2.6	4.5	6.8	11.9	21.6	37.6	69.0	127.7	234.6	537.0
All races 3/, female										
2002.....	2.3	3.8	6.1	11.4	18.8	32.9	56.3	101.2	181.1	429.1
2001.....	2.3	3.7	6.1	11.2	19.4	34.6	57.8	101.3	175.9	435.8
2000.....	2.1	3.3	5.8	10.3	19.3	33.3	54.7	97.5	177.2	428.1
1999.....	2.1	3.1	5.8	10.2	17.6	32.8	54.4	93.9	170.4	432.6
White, both sexes										
2002.....	1.8	3.2	4.9	9.2	16.2	30.5	56.1	105.6	195.3	449.9
2001.....	1.8	3.0	5.0	9.0	16.0	32.1	56.1	103.6	189.1	456.0
2000.....	1.8	2.9	4.8	8.6	16.3	29.7	51.9	102.4	189.9	451.6
1999.....	1.6	2.6	4.5	8.6	15.5	28.9	53.3	99.4	184.2	447.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of urinary system (N00-N39)										
White, male										
2002.....	2.0	3.6	5.6	9.7	17.2	34.2	64.7	123.7	233.4	523.8
2001.....	2.0	3.3	5.7	9.7	17.4	36.5	63.4	119.9	228.5	535.0
2000.....	2.1	3.5	5.4	9.4	17.5	33.2	59.3	120.6	228.4	540.8
1999.....	1.8	3.0	4.8	9.8	17.2	31.4	61.1	117.9	223.8	521.3
White, female										
2002.....	1.7	2.8	4.3	8.8	15.2	27.3	49.0	92.6	172.0	417.8
2001.....	1.6	2.7	4.4	8.4	14.8	28.2	50.2	92.0	165.3	423.0
2000.....	1.5	2.4	4.2	7.8	15.2	26.7	46.1	89.5	167.0	415.4
1999.....	1.5	2.2	4.1	7.5	13.9	26.7	47.1	86.3	160.9	417.7
Black, both sexes										
2002.....	7.5	13.3	23.8	37.8	58.0	90.9	148.1	222.8	328.9	652.1
2001.....	7.3	14.2	22.5	36.7	59.1	101.9	144.9	220.6	343.2	655.3
2000.....	7.2	13.9	22.5	36.1	57.6	95.6	144.5	217.0	339.1	628.5
1999.....	7.7	13.1	22.3	32.2	55.4	93.3	145.0	216.7	332.3	669.3
Black, male										
2002.....	8.3	16.4	28.8	44.4	68.9	100.5	173.6	256.7	373.7	776.2
2001.....	7.6	18.1	25.4	42.5	61.8	117.9	156.5	237.9	395.7	736.4
2000.....	8.8	18.8	27.0	42.0	62.1	101.2	153.5	254.2	392.0	695.4
1999.....	9.2	17.2	25.7	32.2	64.5	100.5	168.9	268.3	421.2	779.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of urinary system (N00-N39)										
Black, female										
2002.....	6.7	10.7	19.6	32.4	49.4	83.8	131.1	202.5	305.6	604.6
2001.....	7.0	10.8	19.9	31.8	57.0	90.1	137.2	210.2	316.4	625.2
2000.....	5.8	9.6	18.6	31.3	54.1	91.4	138.5	194.5	312.5	603.5
1999.....	6.4	9.7	19.3	32.3	48.3	87.9	129.2	185.7	288.2	628.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Glomerular and renal tubulo-interstitial diseases (N00-N15)										
All races 3/, both sexes										
2002.....	0.1	0.2	0.3	0.5	0.6	1.1	2.0	3.9	6.2	13.9
2001.....	0.1	0.2	0.3	0.4	0.6	1.2	2.0	3.8	5.8	13.7
2000.....	0.2	0.2	0.3	0.5	0.7	1.2	1.9	3.7	6.5	15.7
1999.....	0.2	0.2	0.5	0.6	0.8	1.3	2.2	4.2	7.2	16.4
All races 3/, male										
2002.....	*	*	0.3	0.5	0.6	1.2	2.3	4.7	6.7	18.1
2001.....	*	*	0.3	0.4	0.5	1.3	2.4	3.9	7.6	17.2
2000.....	*	0.2	0.4	0.5	0.7	1.2	1.6	4.3	7.2	20.0
1999.....	0.2	0.2	0.4	0.6	0.8	1.2	2.0	4.8	9.8	22.6
All races 3/, female										
2002.....	*	0.3	0.3	0.4	0.7	1.1	1.7	3.3	6.0	12.0
2001.....	*	0.2	0.4	0.4	0.7	1.2	1.6	3.7	4.7	12.3
2000.....	*	0.3	0.3	0.5	0.7	1.2	2.2	3.3	6.1	13.9
1999.....	*	0.3	0.5	0.6	0.7	1.3	2.5	3.8	5.6	13.9
White, both sexes										
2002.....	0.2	0.2	0.3	0.4	0.6	1.2	2.1	3.8	6.2	13.9
2001.....	0.1	0.1	0.3	0.4	0.5	1.1	2.0	3.8	5.5	13.9
2000.....	0.1	0.2	0.2	0.5	0.6	1.1	1.8	3.5	6.3	15.2
1999.....	0.1	0.3	0.4	0.5	0.6	1.1	2.1	4.2	7.1	16.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Glomerular diseases (N00-N07)										
All races 3/, both sexes										
2002.....	*	0.1	0.1	0.2	0.2	0.4	0.7	1.3	2.2	3.3
2001.....	*	*	0.2	0.2	0.2	0.6	0.9	1.2	1.8	3.3
2000.....	0.1	0.1	0.2	0.2	0.3	0.6	0.8	1.2	2.4	3.6
1999.....	*	0.1	0.2	0.2	0.4	0.5	0.9	1.7	2.3	3.7
All races 3/, male										
2002.....	*	*	*	0.3	*	0.5	0.9	1.6	2.4	4.3
2001.....	*	*	*	*	*	0.7	1.1	1.5	2.6	4.6
2000.....	*	*	0.3	*	0.4	0.7	0.8	1.7	2.6	5.6
1999.....	*	*	*	*	0.5	0.5	0.8	2.2	3.4	5.6
All races 3/, female										
2002.....	*	*	*	*	*	*	0.6	1.1	2.0	2.8
2001.....	*	*	*	*	*	0.6	0.7	1.0	1.3	2.8
2000.....	*	*	*	*	*	0.5	0.8	0.8	2.3	2.8
1999.....	*	*	*	*	0.4	0.5	1.0	1.4	1.7	2.9
White, both sexes										
2002.....	*	*	*	0.2	0.2	0.4	0.7	1.2	2.0	3.0
2001.....	*	*	*	*	*	0.5	0.9	1.2	1.7	3.3
2000.....	*	*	*	0.2	0.3	0.5	0.8	1.1	2.4	3.3
1999.....	*	*	0.1	0.2	0.3	0.4	0.9	1.7	2.1	3.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nephrotic syndrome (N04)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	*	*	*	0.6	0.7
2001.....	*	*	*	*	*	*	*	0.3	*	0.7
2000.....	*	*	*	*	*	*	*	*	0.6	1.0
1999.....	*	*	*	*	*	*	0.3	0.4	0.5	0.7
All races 3/, male										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	0.7
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	0.6	0.8
1999.....	*	*	*	*	*	*	*	0.5	*	*
White, both sexes										
2002.....	*	*	*	*	*	*	*	*	0.5	0.6
2001.....	*	*	*	*	*	*	*	0.3	*	0.6
2000.....	*	*	*	*	*	*	*	*	0.6	0.9
1999.....	*	*	*	*	*	*	0.3	0.4	*	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified glomerular diseases (N02-N03,N05-N07)

All races 3/, both sexes

2002.....	*	0.1	0.1	0.2	*	0.3	0.6	1.1	1.6	2.4
2001.....	*	*	0.1	*	0.2	0.5	0.7	0.9	1.4	2.5
2000.....	*	*	0.1	0.2	0.2	0.4	0.6	1.0	1.7	2.5
1999.....	*	*	0.2	0.2	0.3	0.4	0.6	1.2	1.8	2.9

All races 3/, male

2002.....	*	*	*	*	*	*	0.8	1.5	1.8	3.3
2001.....	*	*	*	*	*	0.5	1.0	1.2	2.2	3.5
2000.....	*	*	*	*	*	0.5	0.7	1.4	1.7	3.8
1999.....	*	*	*	*	*	*	0.5	1.7	2.4	4.4

All races 3/, female

2002.....	*	*	*	*	*	*	0.4	0.8	1.4	1.9
2001.....	*	*	*	*	*	0.5	0.4	0.6	1.0	2.1
2000.....	*	*	*	*	*	*	0.6	0.7	1.7	1.9
1999.....	*	*	*	*	*	0.4	0.7	0.9	1.4	2.3

White, both sexes

2002.....	*	*	*	0.2	*	0.3	0.6	1.0	1.5	2.2
2001.....	*	*	*	*	*	0.4	0.7	0.9	1.4	2.5
2000.....	*	*	*	0.2	*	0.4	0.6	0.9	1.6	2.1
1999.....	*	*	*	*	0.3	0.3	0.6	1.2	1.7	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal tubulo-interstitial diseases (N10-N15)										
All races 3/, both sexes										
2002.....	*	0.1	0.2	0.2	0.4	0.7	1.2	2.6	4.1	10.6
2001.....	*	0.1	0.2	0.2	0.4	0.6	1.1	2.5	4.0	10.4
2000.....	*	0.1	0.2	0.3	0.4	0.6	1.1	2.5	4.1	12.0
1999.....	0.1	0.1	0.3	0.3	0.4	0.8	1.3	2.5	4.9	12.7
All races 3/, male										
2002.....	*	*	*	*	0.4	0.7	1.5	3.1	4.3	13.8
2001.....	*	*	*	*	*	0.6	1.3	2.3	5.0	12.5
2000.....	*	*	*	*	*	0.5	0.8	2.6	4.6	14.4
1999.....	*	*	*	0.3	0.4	0.8	1.1	2.6	6.4	17.0
All races 3/, female										
2002.....	*	*	*	*	0.5	0.7	1.1	2.2	3.9	9.2
2001.....	*	*	0.3	0.3	0.5	0.5	0.9	2.7	3.4	9.5
2000.....	*	*	*	0.3	0.5	0.7	1.4	2.5	3.8	11.1
1999.....	*	*	0.3	0.4	*	0.8	1.5	2.4	4.0	11.0
White, both sexes										
2002.....	*	*	0.2	0.2	0.4	0.8	1.3	2.5	4.2	10.9
2001.....	*	*	0.2	0.2	0.3	0.5	1.1	2.6	3.8	10.6
2000.....	*	0.1	*	0.3	0.4	0.6	1.0	2.4	3.9	12.0
1999.....	*	0.2	0.3	0.4	0.3	0.7	1.2	2.5	5.0	12.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)										
All races 3/, both sexes										
2002.....	2.0	3.6	5.7	9.8	16.5	29.0	49.1	83.9	139.8	284.8
2001.....	2.0	3.5	5.7	9.7	16.6	31.1	48.4	80.9	133.8	284.0
2000.....	2.0	3.3	5.3	9.2	16.6	28.9	46.2	78.1	130.0	273.7
1999.....	1.9	3.0	4.9	8.7	15.3	27.7	45.7	75.7	124.8	264.8
All races 3/, male										
2002.....	2.2	4.2	6.7	10.6	18.4	33.1	58.8	103.9	180.0	380.7
2001.....	2.2	4.2	6.5	10.7	17.9	35.9	55.8	98.9	173.9	388.7
2000.....	2.4	4.2	6.1	10.4	18.1	32.1	53.3	97.9	170.1	374.2
1999.....	2.2	3.6	5.6	9.7	17.3	30.5	53.9	96.7	165.1	364.6
All races 3/, female										
2002.....	1.8	2.9	4.8	9.0	14.9	25.4	41.4	69.9	115.3	243.2
2001.....	1.8	2.8	4.8	8.8	15.5	27.0	42.5	68.3	109.8	240.1
2000.....	1.6	2.4	4.5	8.0	15.2	26.1	40.6	64.4	106.4	232.8
1999.....	1.6	2.3	4.3	7.9	13.5	25.3	39.3	61.1	101.2	224.5
White, both sexes										
2002.....	1.4	2.5	3.8	7.1	12.9	23.9	42.3	76.8	133.2	275.3
2001.....	1.4	2.3	4.0	7.1	12.6	25.2	42.1	73.8	126.0	274.6
2000.....	1.4	2.1	3.7	6.6	12.8	23.3	39.6	71.7	122.6	265.0
1999.....	1.2	1.8	3.4	6.4	11.9	22.5	39.4	69.0	116.2	251.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)										
White, male										
2002.....	1.5	2.9	4.5	7.7	14.0	27.9	52.0	97.3	175.3	372.7
2001.....	1.6	2.7	4.7	8.0	14.1	30.0	50.1	93.0	166.1	383.9
2000.....	1.7	2.7	4.2	7.6	14.5	27.2	47.7	91.6	163.8	370.4
1999.....	1.4	2.2	3.8	7.6	13.6	25.5	48.0	89.1	156.2	354.9
White, female										
2002.....	1.2	2.1	3.2	6.6	11.8	20.3	34.4	62.1	107.4	232.9
2001.....	1.1	1.9	3.3	6.2	11.2	21.1	35.6	60.2	101.8	228.8
2000.....	1.1	1.6	3.1	5.6	11.3	19.9	33.2	57.8	98.1	222.1
1999.....	1.1	1.5	2.9	5.3	10.3	19.9	32.5	54.8	92.5	210.6
Black, both sexes										
2002.....	6.6	11.2	21.1	33.3	50.1	78.8	125.7	175.4	238.1	434.2
2001.....	6.5	12.3	19.3	32.2	52.5	87.1	119.0	170.6	245.9	425.6
2000.....	6.1	12.3	18.9	32.0	50.1	82.1	118.6	163.1	238.9	400.4
1999.....	6.7	11.7	18.9	28.5	46.1	76.5	115.7	161.0	241.6	425.3
Black, male										
2002.....	7.4	14.4	26.1	39.4	61.2	88.8	148.2	203.5	278.2	555.3
2001.....	6.8	16.1	22.1	37.2	54.5	99.1	129.9	187.7	307.4	508.3
2000.....	7.7	17.1	22.7	37.0	53.2	84.8	127.0	196.6	287.5	480.9
1999.....	8.0	15.3	22.7	28.7	53.5	80.5	128.9	204.7	320.9	513.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)										
Black, female										
2002.....	5.9	8.5	16.8	28.2	41.5	71.4	110.7	158.6	217.2	387.7
2001.....	6.4	9.0	16.9	28.1	50.9	78.3	111.7	160.3	214.6	394.9
2000.....	4.7	8.2	15.7	27.8	47.6	80.1	113.0	143.0	214.4	370.4
1999.....	5.6	8.5	15.8	28.4	40.3	73.5	106.9	134.7	202.2	392.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Urolithiasis (N20-N23)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	0.2	0.3	0.5	0.9	1.8
2001.....	*	*	*	*	*	*	0.3	0.6	0.7	1.8
2000.....	*	*	*	*	*	*	0.2	0.7	0.9	1.8
1999.....	*	*	*	*	0.2	*	0.3	0.3	0.9	1.9
All races 3/, male										
2002.....	*	*	*	*	*	*	*	*	*	2.0
2001.....	*	*	*	*	*	*	*	*	*	2.0
2000.....	*	*	*	*	*	*	*	*	1.1	1.7
1999.....	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	*	0.5	0.9	1.7
2001.....	*	*	*	*	*	*	*	0.7	0.6	1.7
2000.....	*	*	*	*	*	*	*	0.8	0.8	1.8
1999.....	*	*	*	*	*	*	*	*	0.8	2.0
White, both sexes										
2002.....	*	*	*	*	*	0.2	0.3	0.5	0.9	1.9
2001.....	*	*	*	*	*	*	0.3	0.6	0.8	1.9
2000.....	*	*	*	*	*	*	*	0.7	1.0	1.8
1999.....	*	*	*	*	*	*	0.3	0.3	1.0	1.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Urinary tract infection, site not specified (N39.0)										
All races 3/, both sexes										
2002.....	0.2	0.4	0.7	1.4	2.4	4.8	10.8	23.7	52.2	152.2
2001.....	0.2	0.4	0.6	1.4	2.3	5.3	11.4	24.5	54.9	160.4
2000.....	0.2	0.4	0.7	1.2	2.5	4.9	9.9	25.6	57.7	163.4
1999.....	0.2	0.4	0.6	1.2	2.6	5.1	11.2	25.4	57.2	172.0
All races 3/, male										
2002.....	0.2	0.4	0.5	1.2	2.2	4.3	9.5	20.5	47.1	123.5
2001.....	*	0.3	0.5	1.2	2.1	5.0	10.1	21.6	50.1	125.8
2000.....	*	0.4	0.6	1.0	2.1	4.5	9.2	23.4	52.4	139.8
1999.....	0.2	0.4	0.4	1.1	2.6	4.4	10.9	23.3	53.5	137.5
All races 3/, female										
2002.....	0.2	0.4	0.8	1.6	2.6	5.3	11.9	26.0	55.3	164.7
2001.....	0.2	0.5	0.7	1.5	2.5	5.6	12.4	26.6	57.8	174.9
2000.....	*	0.4	0.7	1.3	2.8	5.1	10.5	27.1	60.8	173.0
1999.....	0.3	0.5	0.8	1.3	2.5	5.6	11.4	26.9	59.3	185.9
White, both sexes										
2002.....	0.2	0.4	0.6	1.3	2.2	4.4	10.2	22.8	50.9	150.8
2001.....	0.2	0.4	0.5	1.2	2.1	4.8	10.6	23.3	53.2	158.2
2000.....	0.1	0.4	0.5	1.1	2.2	4.5	9.1	24.4	56.0	161.9
1999.....	0.2	0.4	0.5	1.2	2.2	4.4	10.3	24.0	56.3	169.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Urinary tract infection, site not specified (N39.0)										
White, male										
2002.....	*	0.4	0.5	1.1	2.0	3.9	8.6	19.3	45.6	121.1
2001.....	*	0.3	0.4	0.9	1.9	4.4	9.2	20.4	49.3	122.1
2000.....	*	0.3	0.5	0.9	1.8	4.0	8.3	22.1	50.2	137.8
1999.....	*	0.3	0.4	1.1	2.2	3.6	9.4	21.3	52.1	132.1
White, female										
2002.....	*	0.4	0.7	1.6	2.3	4.7	11.5	25.3	54.2	163.8
2001.....	0.2	0.5	0.5	1.5	2.3	5.1	11.8	25.3	55.6	173.3
2000.....	*	0.4	0.6	1.3	2.5	4.9	9.7	26.0	59.4	171.7
1999.....	0.2	0.4	0.6	1.3	2.3	5.1	10.9	25.9	58.9	185.0
Black, both sexes										
2002.....	*	*	1.4	2.8	5.0	9.0	18.6	38.7	76.9	191.5
2001.....	*	1.0	1.4	2.5	4.4	9.6	20.3	41.7	83.7	208.5
2000.....	*	*	1.7	2.2	5.6	9.1	19.3	43.4	85.8	196.8
1999.....	*	0.9	1.4	1.8	6.0	11.9	21.6	45.7	74.3	213.0
Black, male										
2002.....	*	*	*	3.0	4.2	8.2	20.4	40.9	77.9	188.7
2001.....	*	*	*	3.1	4.5	10.7	20.2	41.4	72.2	204.2
2000.....	*	*	*	*	6.0	9.9	20.1	45.4	91.0	175.3
1999.....	*	*	*	*	7.0	13.0	29.8	51.8	80.2	215.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Urinary tract infection, site not specified (N39.0)										
Black, female										
2002.....	*	*	*	2.6	5.6	9.6	17.5	37.3	76.4	192.6
2001.....	*	*	*	*	4.3	8.8	20.4	41.9	89.5	210.1
2000.....	*	*	*	*	5.2	8.4	18.9	42.2	83.2	204.8
1999.....	*	*	*	*	5.2	11.2	16.2	42.1	71.3	212.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of urinary system (N28-N36,N39.1-N39.9)										
All races 3/, both sexes										
2002.....	0.1	0.2	0.3	0.3	0.6	0.9	1.2	1.7	3.9	7.5
2001.....	0.1	0.2	0.3	0.4	0.6	0.9	1.2	2.1	3.5	7.1
2000.....	0.2	0.2	0.3	0.4	0.6	0.8	1.2	2.0	3.9	7.5
1999.....	*	0.1	0.2	0.4	0.6	0.8	1.3	1.8	3.8	7.1
All races 3/, male										
2002.....	*	0.2	0.4	0.4	0.7	1.0	1.4	2.1	4.9	9.0
2001.....	*	*	0.3	0.4	0.7	1.1	1.4	2.4	4.4	8.9
2000.....	0.2	*	0.3	0.5	0.6	1.1	1.5	2.3	5.6	10.4
1999.....	*	0.2	0.2	0.5	0.6	1.2	1.8	2.2	4.7	9.9
All races 3/, female										
2002.....	*	*	*	*	0.4	0.8	1.0	1.4	3.4	6.9
2001.....	*	*	*	0.4	0.5	0.7	1.0	1.9	2.9	6.3
2000.....	*	*	0.2	0.4	0.5	0.6	1.0	1.8	2.9	6.3
1999.....	*	*	*	0.3	0.5	0.4	0.9	1.5	3.2	6.0
White, both sexes										
2002.....	*	*	0.2	0.2	0.4	0.8	1.1	1.6	3.8	7.5
2001.....	*	0.1	0.2	0.3	0.5	0.8	1.0	2.1	3.4	7.2
2000.....	0.2	0.1	0.3	0.3	0.5	0.7	1.0	1.9	3.8	7.3
1999.....	*	*	0.2	0.3	0.5	0.7	1.2	1.7	3.4	7.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of male genital organs (N40-N50)										
All races 3/, both sexes										
2002.....	*	*	*	*	0.2	0.2	0.6	0.8	2.6	5.8
2001.....	*	*	*	*	*	0.2	0.6	1.1	2.1	6.3
2000.....	*	*	*	*	*	0.3	0.5	1.0	2.3	5.9
1999.....	*	*	*	*	0.2	0.3	0.6	1.1	2.7	6.1
All races 3/, male										
2002.....	*	*	*	*	0.4	0.5	1.3	2.1	6.9	19.2
2001.....	*	*	*	*	*	0.5	1.4	2.6	5.5	21.2
2000.....	*	*	*	*	*	0.5	1.2	2.5	6.3	20.5
1999.....	*	*	*	*	0.4	0.7	1.3	2.6	7.2	21.2
All races 3/, female										
2002.....
2001.....
2000.....
1999.....
White, both sexes										
2002.....	*	*	*	*	*	*	0.5	0.7	2.6	5.9
2001.....	*	*	*	*	*	0.2	0.6	1.1	2.1	6.4
2000.....	*	*	*	*	*	0.2	0.5	0.9	2.4	5.9
1999.....	*	*	*	*	*	0.3	0.5	1.1	2.6	6.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hyperplasia of prostate (N40)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	*	0.4	0.6	2.2	4.8
2001.....	*	*	*	*	*	*	0.4	0.8	1.6	5.5
2000.....	*	*	*	*	*	*	0.3	0.9	1.9	5.2
1999.....	*	*	*	*	*	*	0.4	0.8	2.1	5.0
All races 3/, male										
2002.....	*	*	*	*	*	*	0.8	1.6	5.8	15.8
2001.....	*	*	*	*	*	*	0.8	2.0	4.2	18.6
2000.....	*	*	*	*	*	*	0.7	2.1	5.2	18.1
1999.....	*	*	*	*	*	*	0.8	2.0	5.6	17.5
All races 3/, female										
2002.....
2001.....
2000.....
1999.....
White, both sexes										
2002.....	*	*	*	*	*	*	0.3	0.6	2.2	4.9
2001.....	*	*	*	*	*	*	0.4	0.8	1.6	5.7
2000.....	*	*	*	*	*	*	0.3	0.9	2.0	5.2
1999.....	*	*	*	*	*	*	0.4	0.8	2.1	5.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of male genital organs (N41-N50)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*	*	0.4	1.0
2001.....	*	*	*	*	*	*	0.3	0.3	0.5	0.8
2000.....	*	*	*	*	*	*	0.2	*	0.4	0.7
1999.....	*	*	*	*	*	*	*	0.3	0.6	1.1

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	1.1	3.4
2001.....	*	*	*	*	*	*	0.6	0.7	1.3	2.6
2000.....	*	*	*	*	*	*	0.5	*	1.1	2.4
1999.....	*	*	*	*	*	*	*	0.7	1.6	3.7

All races 3/, female

2002.....
2001.....
2000.....
1999.....

White, both sexes

2002.....	*	*	*	*	*	*	*	*	*	1.0
2001.....	*	*	*	*	*	*	*	*	0.5	0.7
2000.....	*	*	*	*	*	*	*	*	*	0.7
1999.....	*	*	*	*	*	*	*	*	0.6	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Disorders of female genital organs (N70-N98)

All races 3/, both sexes

2002.....	*	0.1	*	*	*	*	*	0.4	0.9	2.4
2001.....	*	*	*	*	*	*	0.4	0.5	0.9	2.1
2000.....	*	*	*	*	*	0.2	0.3	0.5	1.0	2.5
1999.....	*	*	*	*	*	*	0.2	0.6	0.8	2.0

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	*	0.2	*	*	*	*	*	0.8	1.4	3.4
2001.....	*	*	*	*	*	*	0.6	0.9	1.5	3.0
2000.....	*	*	*	*	*	0.4	0.5	0.9	1.6	3.6
1999.....	*	*	*	*	*	*	0.4	1.0	1.3	2.9

White, both sexes

2002.....	*	*	*	*	*	*	*	0.5	0.9	2.3
2001.....	*	*	*	*	*	*	0.3	0.5	0.9	2.1
2000.....	*	*	*	*	*	*	0.3	0.5	1.0	2.6
1999.....	*	*	*	*	*	*	*	0.5	0.9	2.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Noninflammatory disorders of female genital tract (N80-N98)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*	*	0.6	1.6
2001.....	*	*	*	*	*	*	0.2	0.3	0.6	1.2
2000.....	*	*	*	*	*	*	*	0.3	0.8	1.9
1999.....	*	*	*	*	*	*	*	0.4	0.5	1.6

All races 3/, male

2002.....
2001.....
2000.....
1999.....

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	1.0	2.3
2001.....	*	*	*	*	*	*	0.4	0.6	0.9	1.8
2000.....	*	*	*	*	*	*	*	0.5	1.2	2.7
1999.....	*	*	*	*	*	*	*	0.6	0.7	2.3

White, both sexes

2002.....	*	*	*	*	*	*	*	*	0.7	1.5
2001.....	*	*	*	*	*	*	*	0.3	0.6	1.2
2000.....	*	*	*	*	*	*	*	0.3	0.7	2.0
1999.....	*	*	*	*	*	*	*	0.3	0.5	1.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
All races 3/, both sexes										
2002.....	1.3	1.6	1.9	2.3	2.4	2.3	2.5	3.1	4.8	6.6
2001.....	1.4	1.5	1.8	2.1	2.5	2.2	2.7	3.3	4.3	7.2
2000.....	1.3	1.4	1.8	2.2	2.5	2.6	2.4	3.3	5.0	7.5
1999.....	1.2	1.7	1.8	2.3	2.4	2.6	2.5	3.4	5.0	7.8
All races 3/, male										
2002.....	1.4	1.9	2.0	2.3	2.7	2.4	2.7	3.4	4.8	7.0
2001.....	1.6	1.5	1.9	2.1	2.5	2.5	3.1	3.6	4.8	7.7
2000.....	1.5	1.6	2.2	2.0	2.6	2.7	2.6	3.3	5.3	10.6
1999.....	1.3	1.9	2.0	2.5	2.5	2.4	2.6	3.4	5.7	8.5
All races 3/, female										
2002.....	1.2	1.4	1.8	2.3	2.2	2.3	2.3	2.9	4.9	6.5
2001.....	1.3	1.4	1.8	2.1	2.6	2.0	2.4	3.1	4.0	7.0
2000.....	1.1	1.3	1.5	2.4	2.5	2.5	2.3	3.3	4.8	6.3
1999.....	1.2	1.6	1.7	2.2	2.3	2.8	2.3	3.4	4.5	7.6
White, both sexes										
2002.....	1.3	1.7	2.0	2.4	2.5	2.5	2.5	3.2	4.8	6.9
2001.....	1.5	1.4	1.9	2.2	2.7	2.3	2.7	3.4	4.4	7.4
2000.....	1.3	1.5	2.0	2.2	2.6	2.6	2.4	3.4	5.2	7.8
1999.....	1.3	1.8	1.8	2.4	2.4	2.7	2.4	3.3	5.1	8.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congenital malformations of heart (Q20-Q24)										
All races 3/, both sexes										
2002.....	0.5	0.6	0.5	0.6	0.5	0.6	0.7	1.0	1.6	2.3
2001.....	0.6	0.5	0.6	0.5	0.5	0.6	0.7	1.1	1.4	2.6
2000.....	0.6	0.5	0.5	0.6	0.6	0.7	0.8	1.1	2.0	2.5
1999.....	0.6	0.6	0.6	0.6	0.7	0.9	0.9	1.2	1.7	3.3
All races 3/, male										
2002.....	0.5	0.6	0.5	0.6	0.5	0.5	0.7	1.0	1.3	1.4
2001.....	0.6	0.6	0.6	0.5	0.5	0.7	1.0	1.0	1.6	1.7
2000.....	0.6	0.5	0.6	0.6	0.6	0.8	0.8	1.1	2.0	2.7
1999.....	0.6	0.7	0.7	0.6	0.7	0.7	0.9	1.0	1.8	3.1
All races 3/, female										
2002.....	0.5	0.6	0.6	0.6	0.4	0.6	0.8	1.0	1.8	2.6
2001.....	0.6	0.5	0.6	0.4	0.5	0.6	0.5	1.2	1.3	3.0
2000.....	0.5	0.5	0.5	0.7	0.5	0.5	0.8	1.0	1.9	2.4
1999.....	0.6	0.6	0.5	0.7	0.6	1.0	0.8	1.3	1.7	3.4
White, both sexes										
2002.....	0.5	0.6	0.5	0.6	0.4	0.6	0.8	1.0	1.6	2.4
2001.....	0.6	0.6	0.6	0.4	0.5	0.6	0.7	1.2	1.5	2.6
2000.....	0.6	0.5	0.6	0.6	0.6	0.7	0.7	1.1	2.1	2.7
1999.....	0.6	0.7	0.6	0.7	0.6	0.9	0.9	1.3	1.8	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other congenital malformations of circulatory system (Q25-Q28)										
All races 3/, both sexes										
2002.....	0.1	0.2	0.2	0.3	0.3	0.4	0.6	0.8	1.3	1.5
2001.....	0.2	0.2	0.2	0.3	0.3	0.4	0.8	0.6	0.9	1.7
2000.....	0.2	0.2	0.2	0.3	0.4	0.3	0.6	0.7	1.2	2.1
1999.....	0.1	0.3	0.3	0.3	0.3	0.6	0.5	0.7	1.2	1.7
All races 3/, male										
2002.....	*	0.3	0.2	0.3	0.4	0.5	0.8	0.7	1.3	1.8
2001.....	0.2	0.3	*	0.4	*	*	0.8	0.9	1.2	2.1
2000.....	0.2	0.3	0.4	*	0.5	*	0.8	0.8	1.4	3.0
1999.....	*	0.3	0.4	0.4	*	0.6	*	0.9	1.4	2.2
All races 3/, female										
2002.....	*	0.2	0.2	0.3	*	0.4	0.4	0.9	1.3	1.4
2001.....	*	*	*	*	*	0.4	0.7	*	0.7	1.6
2000.....	*	0.2	*	0.3	0.4	*	0.4	0.7	1.1	1.7
1999.....	*	0.3	0.2	*	0.4	0.6	0.5	0.5	1.1	1.5
White, both sexes										
2002.....	*	0.2	0.2	0.2	0.3	0.5	0.6	0.8	1.3	1.6
2001.....	0.1	0.2	0.2	0.3	0.3	0.4	0.7	0.7	1.0	1.7
2000.....	0.2	0.2	0.2	0.2	0.4	0.3	0.6	0.8	1.2	2.2
1999.....	0.1	0.3	0.3	0.3	0.3	0.6	0.4	0.6	1.2	1.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)										
All races 3/, both sexes										
2002.....	0.3	0.4	0.4	0.5	0.6	0.6	1.0	1.1	1.7	2.7
2001.....	0.3	0.3	0.3	0.4	0.6	0.7	0.9	1.3	1.8	2.5
2000.....	0.3	0.3	0.3	0.5	0.6	0.8	0.8	1.3	1.6	2.8
1999.....	0.2	0.3	0.3	0.4	0.6	0.7	0.9	1.3	1.8	2.6
All races 3/, male										
2002.....	0.4	0.5	0.5	0.4	0.7	0.7	1.2	1.4	1.8	3.6
2001.....	0.4	0.3	0.3	0.4	0.6	0.8	1.0	1.4	1.8	3.7
2000.....	0.3	0.4	0.4	0.4	0.6	0.9	0.8	1.1	1.8	4.6
1999.....	0.3	0.4	0.4	0.5	0.5	0.7	1.1	1.4	2.1	3.0
All races 3/, female										
2002.....	0.2	0.2	0.3	0.5	0.5	0.5	0.8	0.9	1.7	2.3
2001.....	0.2	0.3	0.2	0.4	0.6	0.6	0.8	1.2	1.8	2.0
2000.....	0.2	0.2	*	0.6	0.6	0.8	0.9	1.3	1.5	2.1
1999.....	0.2	0.3	0.2	0.4	0.6	0.8	0.7	1.2	1.7	2.5
White, both sexes										
2002.....	0.3	0.4	0.4	0.5	0.6	0.6	1.0	1.1	1.7	2.7
2001.....	0.3	0.3	0.3	0.4	0.6	0.7	0.9	1.3	1.7	2.6
2000.....	0.3	0.3	0.3	0.5	0.6	0.8	0.9	1.3	1.7	2.8
1999.....	0.2	0.3	0.3	0.5	0.6	0.8	0.9	1.2	1.8	2.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Down's syndrome (Q90)										
All races 3/, both sexes										
2002.....	0.2	0.3	0.7	0.9	1.0	0.6	*	*	*	*
2001.....	0.3	0.3	0.6	0.9	1.0	0.4	*	*	*	*
2000.....	0.2	0.3	0.7	0.8	0.8	0.6	*	*	*	*
1999.....	0.2	0.4	0.5	0.8	0.7	0.3	*	*	*	*
All races 3/, male										
2002.....	0.2	0.4	0.7	0.9	1.1	0.6	*	*	*	*
2001.....	0.3	0.3	0.6	0.8	1.0	0.5	*	*	*	*
2000.....	*	0.2	0.8	0.8	0.7	0.5	*	*	*	*
1999.....	*	0.4	0.4	0.8	0.8	*	*	*	*	*
All races 3/, female										
2002.....	0.2	0.3	0.6	0.9	0.9	0.6	*	*	*	*
2001.....	0.3	0.3	0.7	1.0	1.0	*	*	*	*	*
2000.....	*	0.3	0.6	0.7	0.9	0.7	*	*	*	*
1999.....	*	0.3	0.6	0.8	0.6	0.4	*	*	*	*
White, both sexes										
2002.....	0.2	0.3	0.7	1.0	1.1	0.6	*	*	*	*
2001.....	0.3	0.3	0.7	1.0	1.1	0.5	*	*	*	*
2000.....	0.2	0.3	0.8	0.8	0.9	0.7	*	*	*	*
1999.....	0.2	0.4	0.5	0.9	0.8	0.3	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
All races 3/, both sexes										
2002.....	6.0	6.4	6.9	7.0	8.3	10.4	16.1	28.3	53.3	208.0
2001.....	7.7	9.1	8.8	8.4	9.9	12.7	17.8	29.8	54.3	206.4
2000.....	7.6	8.6	8.0	8.6	10.2	14.2	19.0	30.1	56.5	208.1
1999.....	5.1	5.9	5.8	6.9	8.9	11.9	17.6	26.5	50.5	187.7
All races 3/, male										
2002.....	7.6	8.5	9.0	9.0	11.2	12.7	19.7	32.9	60.6	175.5
2001.....	9.9	11.9	11.8	11.0	13.2	15.5	20.9	33.2	60.6	175.0
2000.....	10.2	11.7	11.2	11.4	13.4	18.9	22.4	36.1	62.3	182.7
1999.....	6.7	8.2	7.9	9.1	11.9	15.1	21.8	31.5	54.4	166.3
All races 3/, female										
2002.....	4.5	4.4	5.0	5.0	5.6	8.4	13.3	25.1	48.9	222.1
2001.....	5.5	6.3	5.9	5.9	6.9	10.2	15.4	27.4	50.4	219.5
2000.....	5.0	5.6	5.0	5.9	7.3	10.1	16.4	26.0	53.0	218.5
1999.....	3.4	3.7	3.9	4.9	6.2	9.2	14.2	23.0	48.3	196.4
White, both sexes										
2002.....	5.7	5.8	6.1	6.4	7.9	9.8	15.9	27.9	53.3	211.6
2001.....	7.6	8.8	8.3	7.9	9.4	11.8	17.2	29.5	53.9	210.0
2000.....	7.4	8.2	7.3	7.9	9.5	13.0	18.5	29.5	56.0	212.2
1999.....	4.4	5.2	5.0	6.2	8.2	11.1	17.0	25.5	49.5	190.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
White, male										
2002.....	7.2	7.6	7.7	8.1	10.7	11.9	19.2	31.6	60.0	178.2
2001.....	10.0	11.6	11.2	10.1	12.2	14.3	20.3	33.2	60.6	177.0
2000.....	10.1	11.2	10.1	10.2	12.4	17.1	21.5	35.4	62.0	186.8
1999.....	5.9	7.0	6.7	8.0	10.9	13.9	21.0	30.3	52.6	167.3
White, female										
2002.....	4.2	4.0	4.4	4.7	5.3	7.9	13.1	25.3	49.2	226.1
2001.....	5.3	6.0	5.5	5.7	6.7	9.7	14.6	26.8	49.9	223.8
2000.....	4.6	5.2	4.6	5.6	6.7	9.4	16.0	25.4	52.4	222.5
1999.....	2.9	3.4	3.4	4.5	5.7	8.6	13.8	22.1	47.6	199.4
Black, both sexes										
2002.....	9.5	12.1	15.7	13.8	13.6	17.8	23.1	39.5	64.9	196.0
2001.....	9.8	13.2	14.6	14.3	15.4	21.7	27.9	38.6	66.6	191.6
2000.....	11.2	13.8	15.9	16.0	19.0	28.3	29.7	44.2	73.4	186.4
1999.....	10.3	13.1	13.3	14.9	17.0	22.3	27.2	42.7	72.5	182.4
Black, male										
2002.....	12.4	17.1	22.1	19.5	20.0	22.9	31.0	57.0	87.1	181.2
2001.....	11.7	17.3	18.9	21.7	25.4	29.5	31.6	40.9	70.6	184.9
2000.....	13.5	18.2	23.6	25.0	25.5	41.4	38.1	54.8	83.5	161.5
1999.....	13.6	19.7	19.1	22.4	24.0	30.1	35.6	54.1	88.1	184.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, female

2002.....	7.0	7.7	10.2	9.0	8.7	14.1	17.7	29.1	53.3	201.7
2001.....	8.1	9.7	11.0	8.2	7.5	16.0	25.3	37.2	64.6	194.1
2000.....	9.1	9.9	9.4	8.6	13.9	18.6	24.2	37.9	68.4	195.7
1999.....	7.3	7.3	8.5	8.9	11.5	16.5	21.6	35.8	64.8	181.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Senility (R54)										
All races 3/, both sexes										
2002.....	*	*	*	*	*	0.4	1.2	3.7	9.8	72.0
2001.....	*	*	*	*	*	0.4	1.3	3.3	10.8	73.0
2000.....	*	*	*	*	*	0.4	1.3	3.6	10.3	67.2
1999.....	*	*	*	*	*	0.3	1.1	2.9	9.0	56.5
All races 3/, male										
2002.....	*	*	*	*	*	0.5	1.2	3.6	10.3	56.5
2001.....	*	*	*	*	*	0.5	1.5	3.2	11.4	57.2
2000.....	*	*	*	*	*	*	1.3	3.8	10.5	51.7
1999.....	*	*	*	*	*	*	1.5	3.1	8.9	45.8
All races 3/, female										
2002.....	*	*	*	*	*	*	1.2	3.8	9.5	78.8
2001.....	*	*	*	*	*	*	1.2	3.3	10.4	79.6
2000.....	*	*	*	*	*	*	1.2	3.4	10.2	73.6
1999.....	*	*	*	*	*	*	0.8	2.8	9.0	60.8
White, both sexes										
2002.....	*	*	*	*	*	0.4	1.2	3.8	10.1	74.2
2001.....	*	*	*	*	*	0.3	1.4	3.3	11.0	75.1
2000.....	*	*	*	*	*	0.3	1.2	3.5	10.2	69.0
1999.....	*	*	*	*	*	0.4	1.1	2.8	8.8	58.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Senility (R54)										
Black, female										
2002.....	*	*	*	*	*	*	*	*	8.8	57.9
2001.....	*	*	*	*	*	*	*	*	10.6	56.4
2000.....	*	*	*	*	*	*	*	*	14.4	63.5
1999.....	*	*	*	*	*	*	*	*	11.7	41.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ill-defined and unknown causes of mortality (R95-R99)										
All races 3/, both sexes										
2002.....	5.0	5.3	5.5	5.5	5.8	6.6	8.7	13.4	20.8	58.3
2001.....	6.6	8.0	7.6	6.8	7.6	8.8	10.4	14.9	22.9	62.1
2000.....	6.5	7.4	6.8	6.9	7.6	10.0	11.9	16.2	26.9	73.0
1999.....	4.2	4.9	4.6	5.2	6.3	7.9	10.5	13.5	22.9	66.7
All races 3/, male										
2002.....	6.5	7.2	7.4	7.3	8.1	8.4	11.4	16.4	25.5	55.0
2001.....	8.6	10.5	10.2	9.1	10.2	11.2	13.2	18.1	27.6	58.1
2000.....	8.8	10.3	9.6	9.3	10.4	13.8	14.3	20.2	31.2	72.1
1999.....	5.7	6.9	6.3	6.9	8.8	10.5	13.8	17.0	26.4	64.4
All races 3/, female										
2002.....	3.6	3.4	3.6	3.8	3.8	5.0	6.6	11.3	17.9	59.8
2001.....	4.7	5.5	5.0	4.8	5.2	6.8	8.1	12.6	20.0	63.8
2000.....	4.3	4.7	4.1	4.7	5.1	6.8	10.1	13.4	24.3	73.3
1999.....	2.8	3.1	3.0	3.6	4.0	5.7	7.9	11.2	20.9	67.7
White, both sexes										
2002.....	4.9	4.8	4.8	5.1	5.6	6.3	8.7	13.3	20.8	59.3
2001.....	6.8	7.8	7.3	6.6	7.4	8.6	10.1	14.8	22.6	63.2
2000.....	6.4	7.2	6.3	6.5	7.3	9.3	11.9	16.0	27.1	74.5
1999.....	3.7	4.4	4.1	4.7	5.9	7.4	10.3	13.2	22.4	67.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ill-defined and unknown causes of mortality (R95-R99)										
White, male										
2002.....	6.3	6.5	6.4	6.7	7.8	7.9	11.3	16.0	24.9	56.2
2001.....	8.9	10.3	9.9	8.5	9.9	10.8	13.0	18.1	27.2	59.3
2000.....	8.8	10.0	8.8	8.7	9.9	12.5	13.9	19.9	31.4	73.2
1999.....	5.1	6.0	5.4	6.1	8.2	9.7	13.4	16.5	25.3	63.1
White, female										
2002.....	3.5	3.2	3.3	3.6	3.7	4.8	6.5	11.4	18.3	60.6
2001.....	4.6	5.3	4.7	4.7	5.1	6.7	7.9	12.4	19.8	64.9
2000.....	4.0	4.4	3.8	4.4	4.9	6.5	10.2	13.2	24.5	75.0
1999.....	2.4	2.8	2.7	3.3	3.7	5.4	7.8	10.9	20.7	68.5
Black, both sexes										
2002.....	7.4	9.5	11.9	9.9	9.1	10.9	11.0	17.9	24.3	56.1
2001.....	7.4	10.9	11.2	10.4	10.2	11.9	14.0	18.6	28.2	55.6
2000.....	9.2	11.0	12.6	11.6	12.6	18.6	15.2	21.5	30.5	66.0
1999.....	8.2	10.2	10.1	11.0	11.5	13.7	13.8	19.3	32.4	71.4
Black, male										
2002.....	9.9	13.8	17.3	14.6	14.1	15.5	16.0	25.7	40.5	53.6
2001.....	8.4	14.0	14.4	15.8	16.1	17.0	18.5	22.8	35.3	49.9
2000.....	11.2	14.8	18.7	17.0	18.2	28.6	21.4	29.3	38.8	70.3
1999.....	10.8	15.4	15.0	16.4	17.4	20.0	20.4	25.6	41.9	89.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ill-defined and unknown causes of mortality (R95-R99)										
Black, female										
2002.....	5.1	5.7	7.2	6.1	5.3	7.6	7.6	13.2	15.9	57.1
2001.....	6.6	8.3	8.5	5.9	5.6	8.1	11.0	16.1	24.5	57.7
2000.....	7.4	7.8	7.4	7.2	8.2	11.1	11.1	16.8	26.3	64.4
1999.....	6.0	5.7	6.0	6.5	6.8	9.0	9.5	15.4	27.7	64.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other ill-defined and unknown causes of mortality (R96-R99)										
All races 3/, both sexes										
2002.....	5.0	5.3	5.5	5.5	5.8	6.6	8.7	13.4	20.8	58.3
2001.....	6.6	8.0	7.6	6.8	7.6	8.8	10.4	14.9	22.9	62.1
2000.....	6.5	7.4	6.8	6.9	7.6	10.0	11.9	16.2	26.9	73.0
1999.....	4.2	4.9	4.6	5.2	6.3	7.9	10.5	13.5	22.9	66.7
All races 3/, male										
2002.....	6.5	7.2	7.4	7.3	8.1	8.4	11.4	16.4	25.5	55.0
2001.....	8.6	10.5	10.2	9.1	10.2	11.2	13.2	18.1	27.6	58.1
2000.....	8.8	10.3	9.6	9.3	10.4	13.8	14.3	20.2	31.2	72.1
1999.....	5.7	6.9	6.3	6.9	8.8	10.5	13.8	17.0	26.4	64.4
All races 3/, female										
2002.....	3.6	3.4	3.6	3.8	3.8	5.0	6.6	11.3	17.9	59.8
2001.....	4.7	5.5	5.0	4.8	5.2	6.8	8.1	12.6	20.0	63.8
2000.....	4.3	4.7	4.1	4.7	5.1	6.8	10.1	13.4	24.3	73.3
1999.....	2.8	3.1	3.0	3.6	4.0	5.7	7.9	11.2	20.9	67.7
White, both sexes										
2002.....	4.9	4.8	4.8	5.1	5.6	6.3	8.7	13.3	20.8	59.3
2001.....	6.8	7.8	7.3	6.6	7.4	8.6	10.1	14.8	22.6	63.2
2000.....	6.4	7.2	6.3	6.5	7.3	9.3	11.9	16.0	27.1	74.5
1999.....	3.7	4.4	4.1	4.7	5.9	7.4	10.3	13.2	22.4	67.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other ill-defined and unknown causes of mortality (R96-R99)										
White, male										
2002.....	6.3	6.5	6.4	6.7	7.8	7.9	11.3	16.0	24.9	56.2
2001.....	8.9	10.3	9.9	8.5	9.9	10.8	13.0	18.1	27.2	59.3
2000.....	8.8	10.0	8.8	8.7	9.9	12.5	13.9	19.9	31.4	73.2
1999.....	5.1	6.0	5.4	6.1	8.2	9.7	13.4	16.5	25.3	63.1
White, female										
2002.....	3.5	3.2	3.3	3.6	3.7	4.8	6.5	11.4	18.3	60.6
2001.....	4.6	5.3	4.7	4.7	5.1	6.7	7.9	12.4	19.8	64.9
2000.....	4.0	4.4	3.8	4.4	4.9	6.5	10.2	13.2	24.5	75.0
1999.....	2.4	2.8	2.7	3.3	3.7	5.4	7.8	10.9	20.7	68.5
Black, both sexes										
2002.....	7.4	9.5	11.9	9.9	9.1	10.9	11.0	17.9	24.3	56.1
2001.....	7.4	10.9	11.2	10.4	10.2	11.9	14.0	18.6	28.2	55.6
2000.....	9.2	11.0	12.6	11.6	12.6	18.6	15.2	21.5	30.5	66.0
1999.....	8.2	10.2	10.1	11.0	11.5	13.7	13.8	19.3	32.4	71.4
Black, male										
2002.....	9.9	13.8	17.3	14.6	14.1	15.5	16.0	25.7	40.5	53.6
2001.....	8.4	14.0	14.4	15.8	16.1	17.0	18.5	22.8	35.3	49.9
2000.....	11.2	14.8	18.7	17.0	18.2	28.6	21.4	29.3	38.8	70.3
1999.....	10.8	15.4	15.0	16.4	17.4	20.0	20.4	25.6	41.9	89.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other ill-defined and unknown causes of mortality (R96-R99)

Black, female

2002.....	5.1	5.7	7.2	6.1	5.3	7.6	7.6	13.2	15.9	57.1
2001.....	6.6	8.3	8.5	5.9	5.6	8.1	11.0	16.1	24.5	57.7
2000.....	7.4	7.8	7.4	7.2	8.2	11.1	11.1	16.8	26.3	64.4
1999.....	6.0	5.7	6.0	6.5	6.8	9.0	9.5	15.4	27.7	64.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)										
All races 3/, both sexes										
2002.....	1.0	1.1	1.5	1.4	2.3	3.4	6.2	11.2	22.8	77.7
2001.....	1.0	1.1	1.2	1.5	2.2	3.4	6.1	11.6	20.6	71.3
2000.....	1.1	1.1	1.3	1.6	2.5	3.8	5.9	10.4	19.3	67.9
1999.....	0.8	1.0	1.2	1.7	2.5	3.7	6.0	10.0	18.7	64.5
All races 3/, male										
2002.....	1.1	1.3	1.6	1.7	3.0	3.8	7.1	12.9	24.8	64.0
2001.....	1.3	1.4	1.6	1.9	2.9	3.8	6.2	11.8	21.6	59.7
2000.....	1.4	1.4	1.7	2.1	2.9	4.7	6.8	12.1	20.7	58.9
1999.....	1.1	1.3	1.5	2.2	2.9	4.2	6.4	11.5	19.2	56.1
All races 3/, female										
2002.....	0.9	1.0	1.3	1.2	1.7	3.1	5.5	10.1	21.5	83.6
2001.....	0.7	0.8	0.9	1.1	1.6	3.1	6.0	11.4	20.1	76.2
2000.....	0.7	0.9	0.9	1.2	2.1	3.0	5.1	9.2	18.5	71.6
1999.....	0.6	0.7	0.8	1.3	2.1	3.3	5.6	9.0	18.3	67.9
White, both sexes										
2002.....	0.8	1.0	1.2	1.2	2.1	3.1	6.0	10.8	22.4	78.2
2001.....	0.9	1.0	1.0	1.3	1.9	3.0	5.6	11.4	20.3	71.6
2000.....	1.0	1.0	1.0	1.3	2.1	3.3	5.4	10.0	18.8	68.8
1999.....	0.7	0.8	1.0	1.5	2.2	3.3	5.6	9.4	18.2	64.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)										
White, male										
2002.....	0.9	1.1	1.3	1.3	2.7	3.5	6.8	12.0	24.3	64.0
2001.....	1.1	1.3	1.3	1.5	2.3	3.1	5.8	11.8	21.8	59.4
2000.....	1.3	1.1	1.3	1.5	2.5	4.2	6.2	11.8	20.1	59.8
1999.....	0.9	1.0	1.3	1.9	2.6	3.8	6.1	10.8	18.7	56.8
White, female										
2002.....	0.7	0.8	1.1	1.1	1.5	2.8	5.3	10.0	21.2	84.3
2001.....	0.6	0.7	0.7	1.0	1.6	2.8	5.5	11.1	19.4	76.7
2000.....	0.6	0.8	0.8	1.1	1.8	2.6	4.7	8.7	18.0	72.4
1999.....	0.5	0.5	0.7	1.2	1.9	2.9	5.2	8.4	17.9	67.6
Black, both sexes										
2002.....	2.2	2.5	3.8	3.8	4.5	6.4	10.3	18.0	33.0	84.3
2001.....	2.4	2.3	3.3	3.8	4.9	8.5	12.9	16.0	27.6	79.2
2000.....	2.0	2.8	3.3	4.5	6.2	8.8	12.3	17.8	28.8	64.8
1999.....	2.0	2.9	3.2	4.0	5.1	8.5	11.9	18.9	26.9	71.1
Black, male										
2002.....	2.5	3.2	4.8	4.9	5.9	7.0	12.7	26.6	41.3	78.3
2001.....	3.3	3.3	4.2	5.8	8.9	11.4	12.8	14.9	24.1	77.2
2000.....	2.3	3.5	4.9	8.1	7.0	11.7	15.4	19.4	31.2	56.5
1999.....	2.9	4.3	3.9	5.9	5.9	10.1	12.9	23.3	29.6	58.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

Black, female

2002.....	1.9	1.9	3.0	2.9	3.4	5.9	8.7	12.9	28.6	86.7
2001.....	1.6	*	2.5	*	*	6.4	13.0	16.6	29.4	80.0
2000.....	1.7	2.1	2.0	*	5.6	6.7	10.2	16.8	27.6	67.8
1999.....	*	1.6	2.5	*	4.5	7.3	11.3	16.3	25.5	75.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XX. External causes of mortality (*U01-*U03,V01-Y89)										
All races 3/, both sexes										
2002.....	66.2	64.5	56.6	51.0	50.3	56.5	72.1	104.7	161.6	308.3
2001 4/.....	64.3	62.5	54.8	50.2	49.5	56.0	69.9	106.1	161.2	310.1
2000.....	61.0	57.9	51.4	48.3	49.1	52.5	69.9	100.7	155.4	310.5
1999.....	59.9	56.7	49.0	48.3	48.0	57.0	72.3	104.8	164.4	318.2
All races 3/, male										
2002.....	96.2	94.6	83.1	76.8	73.3	82.4	104.9	150.3	226.1	414.8
2001 4/.....	95.4	94.2	81.1	74.0	71.9	80.8	100.2	152.7	231.8	429.1
2000.....	91.6	87.5	76.8	71.2	70.9	74.9	100.4	145.6	225.1	438.2
1999.....	89.5	85.9	73.2	71.9	71.0	82.8	104.4	149.4	240.9	454.8
All races 3/, female										
2002.....	36.6	35.3	31.1	26.9	29.5	34.1	45.9	72.5	122.4	262.0
2001 4/.....	33.7	31.8	29.6	27.8	29.3	34.6	45.8	73.4	118.9	260.3
2000.....	30.9	29.1	27.0	26.8	29.3	33.2	45.9	69.5	114.3	258.5
1999.....	30.7	28.5	25.9	26.4	27.3	35.0	47.3	73.8	119.6	263.2
White, both sexes										
2002.....	67.2	64.6	56.2	51.0	50.4	56.0	73.1	106.1	166.2	319.7
2001 4/.....	64.3	61.7	54.4	49.9	48.9	55.9	69.6	107.6	164.7	319.5
2000.....	60.0	56.8	50.0	47.6	48.9	51.8	69.4	101.5	158.6	320.4
1999.....	59.0	55.2	47.8	47.4	46.6	56.5	71.4	104.5	166.6	326.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XX. External causes of mortality (*U01-*U03,V01-Y89)										
White, male										
2002.....	97.4	93.9	81.5	75.5	73.0	80.7	105.8	151.4	231.7	427.4
2001 4/.....	95.0	92.0	79.4	72.5	70.4	80.4	99.5	154.4	236.3	441.2
2000.....	89.7	84.9	73.6	69.1	69.8	72.8	99.3	146.2	228.6	451.7
1999.....	87.8	82.5	70.6	69.2	68.6	81.4	102.7	148.6	244.1	465.8
White, female										
2002.....	36.9	35.5	31.5	27.5	29.5	34.2	46.4	73.7	126.0	272.8
2001 4/.....	33.4	31.7	29.9	28.4	29.2	34.3	45.4	74.5	121.4	268.6
2000.....	30.2	28.9	26.9	27.0	29.7	33.4	45.3	70.1	116.9	267.0
1999.....	30.1	28.2	25.6	26.6	26.3	34.7	46.6	73.6	120.7	270.9
Black, both sexes										
2002.....	73.4	78.0	69.9	59.7	57.4	61.7	70.5	98.6	118.5	199.6
2001 4/.....	78.0	80.6	69.3	60.7	60.8	63.5	79.8	98.6	129.2	216.8
2000.....	79.5	76.2	70.5	60.6	56.3	64.2	82.0	97.2	127.1	212.4
1999.....	77.5	78.8	65.8	61.7	62.0	68.3	85.8	115.0	147.9	239.5
Black, male										
2002.....	109.0	122.1	112.6	102.1	92.1	100.7	110.8	157.1	182.6	307.7
2001 4/.....	119.6	131.2	111.9	100.3	96.8	97.8	120.5	151.0	199.8	306.4
2000.....	124.9	123.6	116.0	101.8	91.3	106.9	123.7	150.3	199.8	318.3
1999.....	122.3	130.6	108.6	104.1	97.6	108.0	134.0	175.2	215.4	367.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XX. External causes of mortality (*U01-*U03,V01-Y89)										
Black, female										
2002.....	42.1	39.7	33.7	24.9	30.2	32.7	43.5	63.7	85.1	158.1
2001 4/.....	41.3	36.8	32.9	28.2	32.5	38.1	52.7	67.4	93.2	183.6
2000.....	39.4	35.0	31.8	26.9	28.6	32.6	54.4	65.2	90.4	173.0
1999.....	38.1	34.1	29.4	27.0	34.1	38.9	53.8	78.7	114.5	191.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
All races 3/, both sexes										
2002.....	39.4	38.7	34.3	30.4	32.6	38.3	50.8	80.0	131.1	275.4
2001.....	36.7	36.0	31.9	29.3	31.5	36.8	49.3	80.4	130.4	276.4
2000.....	35.4	34.2	30.8	29.9	32.2	35.4	48.8	75.5	124.6	273.5
1999.....	35.4	34.1	29.2	30.1	31.2	38.7	50.9	78.6	133.3	282.4
All races 3/, male										
2002.....	56.8	56.6	49.6	44.8	45.7	53.4	68.9	106.4	170.1	346.6
2001.....	54.1	54.0	46.9	41.8	44.5	49.9	65.6	108.3	173.2	358.5
2000.....	53.1	51.6	46.0	43.1	45.3	48.1	64.6	99.8	164.8	359.7
1999.....	53.1	51.7	43.6	43.6	44.5	53.1	68.2	102.8	181.8	375.4
All races 3/, female										
2002.....	22.4	21.3	19.6	16.8	20.7	25.3	36.3	61.4	107.3	244.6
2001.....	19.6	18.5	17.5	17.6	19.6	25.5	36.3	60.9	104.8	242.0
2000.....	18.0	17.3	16.1	17.5	20.4	24.5	36.4	58.5	100.9	238.4
1999.....	18.1	17.0	15.5	17.6	19.2	26.4	37.3	61.8	105.0	244.8
White, both sexes										
2002.....	40.5	38.5	33.6	29.6	32.1	37.3	51.0	81.1	134.6	286.1
2001.....	36.8	34.9	31.1	28.5	30.5	36.4	48.9	81.2	133.3	285.5
2000.....	35.2	33.2	29.2	28.9	31.8	34.5	48.1	75.7	127.3	282.8
1999.....	35.2	32.8	27.8	28.9	29.5	37.7	49.5	78.0	135.0	290.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
White, male										
2002.....	58.2	56.4	47.8	42.8	44.5	50.9	68.6	106.5	173.0	356.6
2001.....	54.2	51.8	45.1	40.0	42.6	48.9	64.6	108.2	175.2	368.1
2000.....	52.5	49.8	43.2	40.8	43.8	45.9	63.3	98.7	166.4	369.8
1999.....	52.5	49.1	41.1	41.1	42.0	51.1	65.7	100.7	183.4	383.6
White, female										
2002.....	22.8	20.7	19.7	16.9	20.7	25.3	36.7	62.9	111.2	255.4
2001.....	19.3	18.1	17.4	17.6	19.3	25.4	36.1	62.0	107.9	251.0
2000.....	17.8	16.8	15.5	17.5	20.7	24.6	35.8	59.6	104.0	247.4
1999.....	17.9	16.6	14.9	17.4	18.0	26.1	36.6	62.0	106.5	253.0
Black, both sexes										
2002.....	40.0	48.3	45.1	42.0	40.6	47.3	51.9	74.9	96.8	173.4
2001.....	44.3	50.1	44.8	40.6	43.7	45.5	58.0	78.2	104.5	183.6
2000.....	43.8	47.0	47.0	42.0	39.4	46.1	60.2	75.3	100.0	181.1
1999.....	43.6	50.4	44.4	43.4	46.3	50.7	66.6	90.3	124.2	207.6
Black, male										
2002.....	57.9	72.0	72.6	71.2	65.3	75.8	78.4	117.6	149.8	260.5
2001.....	66.6	81.7	71.8	65.9	69.8	69.1	82.5	119.4	164.5	255.3
2000.....	68.8	74.5	76.5	71.5	65.4	74.2	86.3	119.5	156.0	275.6
1999.....	69.0	82.7	73.2	71.5	70.6	77.9	101.8	135.3	177.0	316.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
Black, female										
2002.....	24.2	27.7	21.8	18.1	21.2	26.1	34.1	49.4	69.2	140.1
2001.....	24.5	22.7	21.8	19.9	23.2	28.0	41.7	53.5	74.0	157.0
2000.....	21.7	23.1	21.9	17.8	18.8	25.3	42.8	48.7	71.8	145.9
1999.....	21.3	22.5	20.1	20.4	27.3	30.6	43.2	63.2	98.1	166.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
All races 3/, both sexes										
2002.....	17.1	16.9	15.8	14.9	16.7	16.7	20.2	25.0	29.7	28.8
2001.....	17.3	16.3	15.6	15.2	15.4	16.4	19.5	26.2	30.4	28.9
2000.....	16.8	16.2	15.5	15.7	16.4	16.8	19.4	25.1	30.0	31.5
1999.....	16.6	15.9	14.9	16.3	15.7	18.1	20.9	25.9	32.1	32.0
All races 3/, male										
2002.....	24.5	25.1	22.7	22.2	22.8	23.4	27.5	34.5	42.6	53.0
2001.....	25.4	24.0	22.5	21.4	22.0	22.1	26.8	36.2	45.5	52.7
2000.....	24.8	23.6	22.3	22.4	22.6	22.6	26.3	34.5	44.0	59.8
1999.....	24.0	23.1	21.5	23.0	21.8	24.8	27.5	36.2	48.3	63.5
All races 3/, female										
2002.....	9.9	8.8	9.2	7.9	11.1	10.9	14.3	18.3	21.8	18.3
2001.....	9.4	8.8	8.9	9.4	9.3	11.6	13.8	19.2	21.3	18.9
2000.....	9.0	8.9	8.9	9.5	10.7	11.7	13.9	18.5	21.7	19.9
1999.....	9.4	8.9	8.6	10.0	10.2	12.4	15.7	18.8	22.6	19.3
White, both sexes										
2002.....	17.5	16.9	15.9	14.8	16.7	16.5	20.5	25.5	30.3	29.6
2001.....	17.5	15.9	15.7	15.1	15.2	16.4	19.5	26.6	31.2	29.9
2000.....	16.8	16.0	15.1	15.6	16.4	16.6	19.2	25.3	30.7	32.2
1999.....	16.7	15.6	14.6	16.1	15.4	18.0	20.8	26.0	32.4	32.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
White, male										
2002.....	24.9	25.3	22.5	21.9	22.8	22.7	27.5	34.4	42.8	53.8
2001.....	25.4	23.2	22.6	21.0	21.5	21.7	26.6	36.2	46.6	54.1
2000.....	24.6	23.4	21.7	21.8	22.2	21.9	25.8	34.3	44.5	61.6
1999.....	24.0	22.5	21.0	22.5	21.1	24.6	27.1	35.9	48.4	63.7
White, female										
2002.....	10.0	8.6	9.3	8.1	11.0	11.0	14.8	19.1	22.6	19.1
2001.....	9.5	8.7	8.9	9.5	9.3	11.6	13.8	19.8	21.8	19.7
2000.....	9.0	8.7	8.6	9.6	11.1	12.0	13.9	19.0	22.6	20.3
1999.....	9.4	8.8	8.4	9.9	10.1	12.3	15.8	19.1	23.0	19.7
Black, both sexes										
2002.....	17.1	18.9	16.5	16.8	17.4	17.3	16.9	20.1	24.1	18.4
2001.....	19.4	20.5	16.8	17.8	18.6	18.6	19.0	22.1	19.8	17.5
2000.....	19.3	19.4	19.3	18.1	16.4	18.2	21.3	19.4	20.9	21.6
1999.....	18.0	19.6	18.3	18.7	17.7	18.0	21.0	24.6	28.3	27.7
Black, male										
2002.....	25.2	29.1	26.3	29.0	27.3	28.7	28.4	38.1	44.3	46.1
2001.....	30.6	33.8	25.7	28.6	30.9	28.2	26.3	39.0	31.3	35.2
2000.....	31.0	29.3	30.0	31.0	27.5	31.2	33.4	34.5	39.6	45.0
1999.....	28.3	31.2	29.1	29.3	28.4	27.3	31.5	40.4	48.8	68.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
Black, female										
2002.....	10.0	10.0	8.2	6.8	9.6	8.9	9.2	9.3	13.5	*
2001.....	9.4	8.9	9.3	8.9	9.0	11.5	14.1	11.9	13.9	10.9
2000.....	9.0	10.8	10.2	7.6	7.7	8.6	13.3	10.3	11.5	12.9
1999.....	9.0	9.6	9.1	10.1	9.3	11.2	14.0	15.2	18.1	12.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Railway accidents (V05,V15,V80.6,V81.2-V81.9)										
All races 3/, both sexes										
2002.....	0.3	0.2	0.2	0.2	*	*	*	*	*	*
2001.....	0.3	0.3	0.2	0.1	*	*	*	*	*	*
2000.....	0.3	0.2	0.2	*	*	*	*	*	*	*
1999.....	0.3	0.2	0.2	*	*	*	*	*	*	*
All races 3/, male										
2002.....	0.6	0.5	0.4	0.3	*	*	*	*	*	*
2001.....	0.5	0.4	0.4	*	*	*	*	*	*	*
2000.....	0.6	0.4	0.3	*	*	*	*	*	*	*
1999.....	0.5	0.4	0.3	*	*	*	*	*	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.3	0.2	0.2	*	*	*	*	*	*	*
2001.....	0.3	0.2	0.2	*	*	*	*	*	*	*
2000.....	0.3	0.2	0.1	*	*	*	*	*	*	*
1999.....	0.3	0.2	0.1	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
All races 3/, both sexes										
2002.....	15.7	15.2	14.4	13.4	15.2	15.3	18.9	23.8	28.4	28.0
2001.....	15.5	14.5	13.9	13.7	13.7	14.8	18.0	24.6	28.9	27.8
2000.....	15.1	14.6	13.8	14.2	14.7	15.1	18.1	23.8	28.5	30.4
1999.....	14.8	14.0	13.1	14.4	13.9	16.4	19.3	24.1	30.1	30.4
All races 3/, male										
2002.....	22.1	22.2	20.3	19.6	20.2	20.8	25.0	32.3	39.9	51.7
2001.....	22.5	21.2	19.7	18.9	19.2	19.2	24.2	33.6	43.1	51.1
2000.....	21.9	21.1	19.6	19.9	19.7	19.8	23.9	32.2	41.4	57.5
1999.....	21.0	20.0	18.6	19.8	18.7	22.0	24.7	33.6	45.2	59.8
All races 3/, female										
2002.....	9.4	8.4	8.7	7.5	10.6	10.6	14.1	17.7	21.4	17.7
2001.....	8.7	8.0	8.3	8.8	8.8	11.0	13.1	18.2	20.4	18.0
2000.....	8.4	8.3	8.2	9.0	10.2	11.0	13.5	17.9	20.9	19.3
1999.....	8.6	8.1	8.0	9.4	9.5	11.6	15.0	17.5	21.3	18.5
White, both sexes										
2002.....	16.0	15.2	14.3	13.3	15.1	15.1	19.2	24.3	29.0	28.8
2001.....	15.6	14.2	13.9	13.6	13.5	14.7	17.9	25.0	29.7	28.8
2000.....	15.0	14.5	13.4	14.1	14.7	14.9	17.9	24.0	29.2	31.1
1999.....	14.8	13.7	12.9	14.1	13.5	16.3	19.2	24.3	30.6	30.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
White, male										
2002.....	22.5	22.3	20.0	19.3	20.0	20.1	25.0	32.4	40.1	52.4
2001.....	22.5	20.5	19.7	18.7	18.5	19.0	23.9	33.6	44.3	52.6
2000.....	21.6	20.9	18.9	19.3	19.2	19.1	23.4	32.0	41.7	59.2
1999.....	20.9	19.4	18.1	19.2	18.1	21.8	24.4	33.3	45.5	60.3
White, female										
2002.....	9.5	8.2	8.8	7.6	10.5	10.7	14.5	18.5	22.2	18.5
2001.....	8.8	8.0	8.2	8.8	8.8	11.0	13.0	19.0	20.9	18.9
2000.....	8.4	8.1	7.9	9.1	10.6	11.3	13.5	18.4	21.8	19.7
1999.....	8.6	8.0	7.7	9.3	9.3	11.6	15.1	18.0	21.8	18.8
Black, both sexes										
2002.....	15.9	16.9	15.1	15.0	16.2	15.9	15.8	18.4	22.5	18.1
2001.....	17.6	18.1	15.3	15.9	16.9	16.5	18.3	20.1	19.0	16.6
2000.....	17.9	17.6	18.3	16.7	15.4	16.6	20.1	18.1	20.0	20.3
1999.....	16.6	17.8	16.3	17.1	15.8	16.6	19.6	22.9	26.0	25.5
Black, male										
2002.....	23.0	25.6	23.6	25.3	24.9	25.4	25.7	33.5	40.5	45.0
2001.....	27.5	29.9	23.2	24.7	28.5	23.9	24.6	36.2	30.5	35.2
2000.....	28.4	26.3	28.1	28.2	25.7	27.8	31.4	32.1	38.8	43.8
1999.....	25.6	28.2	25.5	26.3	24.6	24.9	28.5	38.9	45.3	60.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
Black, female										
2002.....	9.6	9.4	8.0	6.6	9.4	8.9	9.2	9.3	13.1	*
2001.....	8.9	8.0	8.7	8.7	7.8	11.1	14.1	10.5	13.1	9.7
2000.....	8.6	10.0	9.9	7.3	7.4	8.3	12.7	9.7	10.6	11.6
1999.....	8.7	8.9	8.5	9.6	8.8	10.4	13.7	13.2	16.4	12.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pedestrian involved in collision with motor vehicle (V02-V04)										
All races 3/, both sexes										
2002.....	1.4	1.4	1.3	1.4	1.4	1.9	2.1	2.4	2.8	3.3
2001.....	1.5	1.4	1.4	1.4	1.3	1.2	2.0	2.4	3.8	3.0
2000.....	1.4	1.4	1.3	1.5	1.5	1.5	1.7	2.5	3.3	3.6
1999.....	1.3	1.3	1.2	1.6	1.5	1.4	1.9	2.3	2.8	3.7
All races 3/, male										
2002.....	2.0	2.2	2.0	2.0	1.9	2.6	2.8	3.0	3.9	5.8
2001.....	2.2	2.0	1.9	2.0	2.0	1.4	2.9	3.2	5.6	5.9
2000.....	2.0	2.2	1.9	2.3	2.1	2.1	2.3	3.4	4.5	7.0
1999.....	1.9	2.0	1.9	2.3	2.1	2.0	2.5	3.2	4.7	6.5
All races 3/, female										
2002.....	0.9	0.7	0.7	0.8	1.0	1.3	1.5	2.0	2.1	2.2
2001.....	0.7	0.8	0.8	0.8	0.7	1.0	1.3	1.8	2.7	1.8
2000.....	0.8	0.6	0.8	0.8	0.9	1.0	1.3	1.9	2.5	2.3
1999.....	0.7	0.6	0.5	0.9	0.9	0.9	1.4	1.7	1.7	2.6
White, both sexes										
2002.....	1.3	1.3	1.2	1.3	1.3	1.6	2.0	2.2	2.6	3.2
2001.....	1.2	1.3	1.2	1.3	1.2	1.1	1.8	2.2	3.5	2.9
2000.....	1.2	1.2	1.2	1.3	1.4	1.4	1.6	2.3	3.2	3.4
1999.....	1.0	1.2	1.1	1.4	1.4	1.3	1.7	2.1	2.7	3.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pedestrian involved in collision with motor vehicle (V02-V04)										
White, male										
2002.....	1.8	2.0	1.8	1.9	1.8	2.3	2.7	2.7	3.6	5.4
2001.....	1.9	1.9	1.7	1.9	1.7	1.3	2.6	2.9	5.5	5.5
2000.....	1.7	1.9	1.7	2.0	1.9	1.8	2.1	3.0	4.1	6.2
1999.....	1.5	1.7	1.7	2.0	2.0	1.9	2.1	2.8	4.4	6.0
White, female										
2002.....	0.8	0.6	0.7	0.8	0.9	1.1	1.5	1.8	2.0	2.2
2001.....	0.6	0.7	0.7	0.8	0.7	0.9	1.1	1.7	2.4	1.9
2000.....	0.6	0.5	0.7	0.7	0.9	1.0	1.3	1.8	2.6	2.2
1999.....	0.6	0.6	0.5	0.8	0.8	0.7	1.4	1.6	1.7	2.6
Black, both sexes										
2002.....	2.0	2.6	2.3	1.9	2.2	2.7	*	*	*	*
2001.....	2.7	2.5	2.6	2.0	2.5	*	*	*	*	*
2000.....	2.8	2.7	2.2	3.0	1.9	*	*	*	*	*
1999.....	3.1	2.4	2.1	3.4	*	*	3.1	*	*	*
Black, male										
2002.....	2.9	4.4	3.7	3.4	*	*	*	*	*	*
2001.....	4.4	3.7	3.9	3.6	5.1	*	*	*	*	*
2000.....	4.0	4.4	3.3	5.3	*	*	*	*	*	*
1999.....	5.1	4.2	3.9	5.9	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)

All races 3/, both sexes

2002.....	0.3	0.4	0.2	0.2	0.2	*	*	*	*	*
2001.....	0.3	0.3	0.2	0.2	0.2	0.2	*	*	*	*
2000.....	0.3	0.3	0.2	0.3	*	*	*	*	*	*
1999.....	0.4	0.3	0.2	0.2	*	0.2	*	*	*	*

All races 3/, male

2002.....	0.5	0.7	0.3	0.3	*	*	*	*	*	*
2001.....	0.6	0.6	0.5	0.5	0.4	0.5	*	*	*	*
2000.....	0.5	0.6	0.4	0.5	*	*	*	*	*	*
1999.....	0.6	0.5	0.4	*	*	0.5	*	*	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.3	0.3	0.2	0.2	*	*	*	*	*	*
2001.....	0.3	0.3	0.2	0.2	0.2	*	*	*	*	*
2000.....	0.2	0.3	0.2	0.3	*	*	*	*	*	*
1999.....	0.4	0.2	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)

All races 3/, both sexes

2002.....	1.8	1.7	1.6	1.2	0.8	0.5	0.3	*	*	*
2001.....	1.7	1.5	1.3	0.8	0.6	0.4	0.3	*	*	*
2000.....	1.6	1.4	1.1	0.9	0.6	0.5	0.3	*	*	*
1999.....	1.4	1.1	0.9	0.8	0.6	0.3	*	*	*	*

All races 3/, male

2002.....	3.2	3.1	2.9	2.1	1.6	1.0	0.6	*	*	*
2001.....	3.0	2.7	2.3	1.6	1.2	0.8	0.6	*	*	*
2000.....	2.9	2.5	2.0	1.6	1.1	0.9	0.7	*	*	*
1999.....	2.3	2.0	1.6	1.5	1.1	0.7	*	*	*	*

All races 3/, female

2002.....	0.4	0.4	0.3	0.3	*	*	*	*	*	*
2001.....	0.4	0.3	0.3	*	*	*	*	*	*	*
2000.....	0.4	0.4	*	*	*	*	*	*	*	*
1999.....	0.4	0.3	*	*	*	*	*	*	*	*

White, both sexes

2002.....	2.0	1.9	1.9	1.3	0.9	0.5	0.3	*	*	*
2001.....	1.9	1.7	1.5	0.9	0.7	0.4	0.3	*	*	*
2000.....	1.9	1.7	1.2	1.0	0.7	0.5	0.3	*	*	*
1999.....	1.5	1.3	1.0	0.9	0.6	0.4	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51, V56-V57,V60-V61,V66-V67,V70-V71,V76-V77)										
All races 3/, both sexes										
2002.....	1.6	1.5	1.3	1.2	1.4	1.2	1.2	1.5	1.7	1.5
2001.....	1.4	1.3	1.1	0.9	1.0	1.1	1.1	1.7	1.6	1.7
2000.....	1.2	1.1	1.0	0.9	1.0	1.0	1.3	1.3	1.5	1.7
1999.....	1.2	1.1	1.0	1.0	1.0	1.3	1.1	1.4	1.6	1.2
All races 3/, male										
2002.....	2.4	2.1	1.8	1.8	2.3	1.7	1.8	2.3	2.1	3.2
2001.....	2.2	2.0	1.6	1.6	1.5	1.6	1.5	2.5	2.6	3.1
2000.....	1.8	1.5	1.5	1.3	1.5	1.5	1.8	2.0	2.2	3.0
1999.....	1.8	1.6	1.6	1.4	1.5	1.9	1.6	2.1	2.8	2.5
All races 3/, female										
2002.....	0.9	0.9	0.7	0.6	0.6	0.8	0.8	0.9	1.5	0.8
2001.....	0.6	0.6	0.6	0.3	0.5	0.7	0.8	1.1	0.9	1.2
2000.....	0.7	0.7	0.5	0.5	0.6	0.6	1.0	0.8	1.0	1.2
1999.....	0.6	0.7	0.4	0.7	0.6	0.8	0.7	1.0	0.9	0.7
White, both sexes										
2002.....	1.8	1.5	1.2	1.1	1.4	1.1	1.2	1.6	1.8	1.6
2001.....	1.4	1.2	1.1	0.9	0.9	1.1	1.1	1.7	1.5	1.9
2000.....	1.2	1.0	0.9	0.8	1.0	1.0	1.3	1.2	1.5	1.8
1999.....	1.2	1.1	0.9	0.9	1.0	1.3	1.1	1.3	1.6	1.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)										
All races 3/, both sexes										
2002.....	3.4	3.2	3.4	3.4	4.4	4.2	6.2	8.3	10.4	9.9
2001.....	3.3	3.1	3.1	3.5	3.6	4.6	5.7	7.8	9.1	10.1
2000.....	3.0	3.2	3.1	3.6	4.0	4.2	5.9	7.9	10.5	10.1
1999.....	3.3	3.2	3.2	3.8	3.7	4.5	6.1	7.7	10.4	9.2
All races 3/, male										
2002.....	4.2	4.0	4.2	4.5	4.9	4.9	7.3	10.7	14.5	16.9
2001.....	4.1	3.9	3.7	4.1	4.2	5.5	6.8	9.6	13.0	18.1
2000.....	3.7	4.2	3.9	4.4	4.5	4.6	7.2	9.5	15.0	18.2
1999.....	4.1	4.0	3.9	4.2	4.4	5.2	7.3	9.7	14.9	17.8
All races 3/, female										
2002.....	2.6	2.4	2.7	2.3	3.9	3.6	5.3	6.7	7.9	6.8
2001.....	2.5	2.4	2.5	2.9	3.1	3.8	4.8	6.5	6.9	6.8
2000.....	2.3	2.3	2.4	2.8	3.6	3.8	4.9	6.9	7.8	6.8
1999.....	2.4	2.4	2.5	3.4	3.1	4.0	5.2	6.4	7.8	5.7
White, both sexes										
2002.....	3.5	3.1	3.4	3.4	4.4	4.3	6.4	8.7	10.9	10.4
2001.....	3.4	3.0	3.1	3.6	3.5	4.6	5.7	8.2	9.5	10.6
2000.....	3.0	3.2	3.0	3.6	4.0	4.3	6.0	8.2	10.9	10.6
1999.....	3.2	3.2	3.1	3.8	3.7	4.6	6.2	8.0	10.7	9.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)										
White, male										
2002.....	4.2	3.9	4.2	4.4	4.9	4.9	7.3	10.8	15.0	17.9
2001.....	4.0	3.6	3.6	4.3	4.0	5.4	6.9	9.9	13.4	18.9
2000.....	3.6	4.1	3.7	4.4	4.1	4.7	7.3	9.6	15.6	19.3
1999.....	4.0	4.0	3.7	4.1	4.3	5.0	7.3	9.8	15.2	18.1
White, female										
2002.....	2.7	2.3	2.7	2.3	4.0	3.7	5.7	7.1	8.4	7.1
2001.....	2.7	2.4	2.6	3.0	3.1	3.9	4.8	6.9	7.2	7.2
2000.....	2.3	2.3	2.3	2.9	3.8	4.0	5.0	7.2	8.2	7.1
1999.....	2.5	2.4	2.6	3.5	3.2	4.2	5.3	6.7	8.1	5.9
Black, both sexes										
2002.....	3.7	3.5	3.5	3.3	4.3	3.4	4.8	5.5	6.0	*
2001.....	3.7	4.1	3.6	3.1	4.7	5.4	6.5	4.3	6.2	*
2000.....	3.6	4.0	4.4	3.5	4.7	3.4	5.5	4.8	6.8	*
1999.....	3.7	3.5	3.7	4.1	3.9	4.5	5.7	5.2	7.8	7.7
Black, male										
2002.....	4.8	4.6	4.9	5.1	5.5	*	7.0	9.2	*	*
2001.....	5.6	6.6	4.9	3.8	6.1	7.6	*	*	*	*
2000.....	5.4	5.7	5.8	5.6	8.1	*	*	*	*	*
1999.....	5.5	4.8	5.1	5.4	5.3	6.8	8.8	9.5	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

Black, female

2002.....	2.8	2.6	2.4	*	3.3	*	*	*	*	*
2001.....	2.0	1.9	2.6	2.6	3.7	3.8	6.5	*	*	*
2000.....	2.1	2.5	3.3	*	*	*	4.9	*	*	*
1999.....	2.1	2.4	2.5	3.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)										
All races 3/, both sexes										
2002.....	1.5	1.4	1.2	1.1	1.3	0.9	1.0	1.0	1.0	0.9
2001.....	1.4	1.3	1.2	1.0	1.2	0.9	1.0	1.1	1.0	0.7
2000.....	1.3	1.2	1.1	0.9	1.0	0.8	0.8	0.9	0.9	1.0
1999.....	1.4	1.2	1.0	1.1	1.0	1.1	0.8	0.8	0.8	1.0
All races 3/, male										
2002.....	2.2	2.2	1.7	1.6	1.7	1.4	1.5	1.5	1.6	1.8
2001.....	2.0	2.0	1.8	1.5	1.7	1.3	1.5	1.5	1.4	*
2000.....	1.9	1.9	1.6	1.3	1.4	1.1	1.3	1.4	1.3	2.2
1999.....	2.0	1.7	1.5	1.6	1.5	1.6	0.9	1.3	*	2.0
All races 3/, female										
2002.....	0.8	0.7	0.7	0.6	0.9	0.6	0.6	0.7	0.6	*
2001.....	0.8	0.6	0.7	0.6	0.7	0.6	0.6	0.8	0.7	*
2000.....	0.7	0.6	0.5	0.6	0.6	0.6	*	0.5	0.7	*
1999.....	0.8	0.7	0.5	0.6	0.7	0.6	0.7	0.5	*	*
White, both sexes										
2002.....	1.5	1.5	1.2	1.1	1.3	1.0	1.0	1.0	1.0	0.8
2001.....	1.5	1.2	1.3	0.9	1.1	1.0	1.0	1.1	1.0	0.7
2000.....	1.3	1.3	1.0	0.8	0.9	0.9	0.8	0.9	0.9	1.0
1999.....	1.4	1.2	1.0	1.1	1.0	1.1	0.8	0.8	0.8	1.0

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of special-use motor vehicle involved in any accident (V83-V86)

All races 3/, both sexes

2002.....	0.3	0.3	0.2	0.3	0.3	0.3	0.5	0.4	0.5	*
2001.....	0.3	0.2	0.3	0.4	0.4	0.4	0.4	0.6	0.5	*
2000.....	0.3	0.3	0.3	0.4	0.5	0.5	0.4	0.5	0.4	*
1999.....	0.3	0.3	0.4	0.4	0.5	0.6	0.6	0.6	0.6	0.7

All races 3/, male

2002.....	0.5	0.7	0.4	0.5	0.6	0.7	0.9	0.9	1.2	*
2001.....	0.6	0.4	0.5	0.7	0.8	0.9	0.8	1.5	1.3	*
2000.....	0.6	0.5	0.5	0.8	1.1	1.0	0.8	1.1	1.1	*
1999.....	0.6	0.6	0.7	0.7	0.9	1.3	1.2	1.4	1.3	2.2

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.3	0.4	0.3	0.3	0.4	0.3	0.5	0.5	0.5	*
2001.....	0.4	0.2	0.3	0.4	0.4	0.5	0.4	0.7	0.6	*
2000.....	0.4	0.3	0.3	0.5	0.5	0.6	0.4	0.5	0.4	*
1999.....	0.4	0.3	0.4	0.4	0.5	0.7	0.6	0.6	0.6	0.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)										
All races 3/, both sexes										
2002.....	5.3	5.1	4.9	4.6	5.2	5.9	7.3	9.6	11.4	11.4
2001.....	5.4	5.1	5.1	5.3	5.2	5.8	7.2	10.4	12.4	11.7
2000.....	5.7	5.4	5.5	5.6	5.8	6.2	7.3	10.2	11.5	13.1
1999.....	5.4	5.2	5.1	5.3	5.2	6.8	8.1	10.8	13.5	13.9
All races 3/, male										
2002.....	6.9	7.0	6.5	6.5	6.5	8.0	9.5	13.0	15.3	21.7
2001.....	7.5	7.2	7.1	6.8	6.9	7.0	9.4	14.3	18.2	21.2
2000.....	8.0	7.4	7.5	7.4	7.6	7.9	9.1	13.6	16.6	24.7
1999.....	7.2	7.2	6.7	7.3	6.7	8.8	9.9	15.0	19.6	27.5
All races 3/, female										
2002.....	3.7	3.3	3.4	2.8	3.9	4.1	5.5	7.2	9.0	6.9
2001.....	3.4	3.1	3.2	3.9	3.7	4.8	5.4	7.7	9.0	7.7
2000.....	3.4	3.5	3.6	3.9	4.2	4.8	5.8	7.8	8.6	8.4
1999.....	3.6	3.3	3.6	3.4	3.9	5.2	6.7	7.8	9.9	8.5
White, both sexes										
2002.....	5.3	5.0	4.8	4.5	5.1	5.9	7.4	9.8	11.6	11.8
2001.....	5.4	5.0	5.1	5.1	5.1	5.7	7.2	10.6	13.0	12.2
2000.....	5.7	5.2	5.3	5.5	5.9	6.1	7.1	10.3	11.9	13.5
1999.....	5.4	5.0	4.9	5.2	5.0	6.8	8.2	10.8	13.6	14.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)										
White, male										
2002.....	6.9	6.8	6.3	6.2	6.4	7.8	9.7	12.9	15.4	21.9
2001.....	7.4	6.9	7.1	6.4	6.7	6.9	9.3	14.3	18.8	22.3
2000.....	7.8	7.1	7.2	7.1	7.5	7.5	8.7	13.8	16.8	25.8
1999.....	7.2	6.8	6.5	7.2	6.3	8.7	10.1	14.7	19.6	28.3
White, female										
2002.....	3.7	3.1	3.4	2.8	3.9	4.2	5.5	7.6	9.3	7.4
2001.....	3.4	3.2	3.2	3.9	3.7	4.7	5.6	8.0	9.4	7.9
2000.....	3.5	3.4	3.6	3.9	4.4	4.8	5.8	7.9	8.9	8.5
1999.....	3.6	3.2	3.5	3.4	3.8	5.1	6.7	8.1	10.1	8.7
Black, both sexes										
2002.....	5.4	6.2	5.8	5.7	5.6	6.4	6.0	8.4	9.7	7.1
2001.....	6.4	6.7	5.6	7.2	6.1	6.9	6.9	10.4	7.0	7.1
2000.....	6.7	7.5	7.3	5.9	5.4	7.4	8.7	7.3	8.5	8.8
1999.....	5.4	7.9	6.7	5.9	6.9	7.2	8.3	11.2	13.3	10.2
Black, male										
2002.....	7.5	8.5	8.2	9.5	8.1	10.2	9.0	16.1	*	21.4
2001.....	9.0	10.8	8.2	11.1	10.0	8.4	10.1	18.1	*	*
2000.....	10.9	10.7	11.2	10.0	8.3	11.7	14.4	11.3	16.9	*
1999.....	7.7	12.0	10.1	9.0	10.6	10.1	8.8	19.9	24.4	23.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

Black, female

2002.....	3.6	4.2	3.7	2.6	3.6	*	*	*	*	*
2001.....	4.1	3.0	3.4	4.0	*	5.8	4.7	5.8	*	*
2000.....	3.0	4.8	4.0	*	*	4.2	4.9	*	*	*
1999.....	3.4	4.3	3.9	3.5	4.0	5.0	7.9	6.0	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)										
All races 3/, both sexes										
2002.....	0.2	0.3	0.3	0.2	0.3	0.2	0.4	0.4	0.5	*
2001.....	0.4	0.4	0.4	0.3	0.4	0.6	0.5	0.6	0.6	0.6
2000.....	0.5	0.4	0.4	0.4	0.4	0.5	0.5	0.6	0.7	0.7
1999.....	0.6	0.7	0.6	0.6	0.6	0.8	0.7	1.2	1.2	1.2
All races 3/, male										
2002.....	0.3	0.6	0.5	0.4	0.5	0.5	0.6	0.7	1.0	*
2001.....	0.6	0.6	0.6	0.5	0.7	1.0	0.7	0.9	*	*
2000.....	0.7	0.5	0.5	0.5	0.7	0.7	0.8	0.9	1.1	*
1999.....	0.9	1.0	0.9	1.0	0.9	1.1	1.0	1.6	1.8	2.7
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	0.3	*	*	*	*	*	0.5	*	*
2000.....	0.2	0.3	0.3	*	*	*	*	*	*	*
1999.....	0.3	0.4	0.2	0.3	*	0.4	0.4	0.9	0.9	*
White, both sexes										
2002.....	0.2	0.3	0.3	0.2	0.3	*	0.4	0.3	0.5	*
2001.....	0.3	0.4	0.3	0.3	0.4	0.5	0.5	0.6	0.6	0.5
2000.....	0.4	0.4	0.5	0.3	0.4	0.4	0.4	0.6	0.7	0.7
1999.....	0.6	0.7	0.6	0.7	0.6	0.7	0.6	1.1	1.2	1.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Water transport accidents (V90-V94)										
All races 3/, both sexes										
2002.....	0.3	0.2	0.3	0.3	0.2	0.4	0.3	*	*	*
2001.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	*	*	*
2000.....	0.3	0.2	0.3	0.3	0.2	0.3	0.2	*	*	*
1999.....	0.3	0.4	0.3	0.3	0.3	0.3	*	*	*	*
All races 3/, male										
2002.....	0.5	0.4	0.5	0.5	0.5	0.7	0.6	*	*	*
2001.....	0.5	0.5	0.5	0.5	0.5	0.5	0.6	*	*	*
2000.....	0.6	0.5	0.6	0.5	0.4	0.6	*	*	*	*
1999.....	0.5	0.7	0.5	0.6	0.6	0.6	*	*	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.3	0.2	0.2	0.2	0.3	0.3	0.3	*	*	*
2001.....	0.3	0.3	0.3	0.2	0.3	*	0.3	*	*	*
2000.....	0.4	0.2	0.3	0.3	0.2	0.3	0.3	*	*	*
1999.....	0.3	0.4	0.3	0.3	0.3	0.3	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Air and space transport accidents (V95-V97)

All races 3/, both sexes

2002.....	0.3	0.4	0.4	0.5	0.5	0.4	0.4	*	*	*
2001.....	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.3	*	*
2000.....	0.3	0.4	0.5	0.5	0.6	0.5	0.2	*	*	*
1999.....	0.4	0.4	0.4	0.5	0.5	0.4	0.4	*	*	*

All races 3/, male

2002.....	0.5	0.8	0.7	0.9	0.9	0.8	0.8	*	*	*
2001.....	0.8	0.7	0.9	0.8	0.9	0.9	0.7	*	*	*
2000.....	0.6	0.7	0.9	1.0	0.9	0.9	*	*	*	*
1999.....	0.5	0.7	0.7	1.0	0.9	0.6	0.8	*	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	0.3	0.2	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.4	0.5	0.5	0.6	0.6	0.4	0.4	*	*	*
2001.....	0.6	0.6	0.6	0.6	0.6	0.6	0.4	0.3	*	*
2000.....	0.4	0.5	0.6	0.6	0.6	0.5	0.3	*	*	*
1999.....	0.4	0.5	0.5	0.6	0.5	0.4	0.4	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified transport accidents and their sequelae (V98-V99,Y85)

All races 3/, both sexes

2002.....	0.3	0.5	0.3	0.3	0.4	0.3	*	0.4	0.5	*
2001.....	0.4	0.3	0.3	0.2	0.3	0.2	0.3	0.5	0.4	*
2000.....	0.3	0.2	0.2	0.2	0.3	0.3	0.3	*	*	*
1999.....	0.2	0.2	0.3	0.3	0.3	*	0.3	*	*	*

All races 3/, male

2002.....	0.4	0.7	0.4	0.5	0.6	0.5	*	*	*	*
2001.....	0.6	0.6	0.5	0.4	0.5	*	*	0.7	*	*
2000.....	0.4	0.4	0.4	*	0.6	0.5	0.5	*	*	*
1999.....	0.4	0.3	0.5	0.5	0.4	*	*	*	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.3	0.5	0.3	0.3	0.3	0.2	*	0.3	0.5	*
2001.....	0.4	0.3	0.3	0.2	0.3	*	0.3	0.4	*	*
2000.....	0.4	0.2	0.2	0.2	0.3	0.4	*	*	*	*
1999.....	0.3	0.2	0.3	0.2	0.3	*	0.3	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)										
All races 3/, both sexes										
2002.....	1.8	2.3	2.8	3.4	4.9	7.9	13.9	27.1	54.0	130.4
2001.....	1.6	2.5	2.7	3.3	4.9	7.8	12.4	25.2	49.9	121.5
2000.....	1.6	2.0	2.6	3.3	4.6	6.2	12.0	22.1	44.5	112.6
1999.....	1.7	2.3	2.2	3.0	4.6	6.9	11.3	22.3	45.5	110.2
All races 3/, male										
2002.....	2.9	3.6	4.3	5.2	7.1	10.7	18.9	35.1	67.9	158.9
2001.....	2.8	4.2	4.4	5.0	7.1	10.6	16.0	34.0	64.1	155.7
2000.....	2.7	3.4	4.2	5.0	6.9	8.6	15.7	29.4	57.3	145.4
1999.....	2.8	3.8	3.6	4.7	7.0	9.8	15.3	27.9	61.6	142.5
All races 3/, female										
2002.....	0.7	1.0	1.4	1.8	3.0	5.5	9.9	21.4	45.5	118.1
2001.....	0.6	0.9	1.2	1.6	2.9	5.4	9.5	19.0	41.3	107.2
2000.....	0.6	0.8	1.2	1.8	2.6	4.2	9.1	17.1	36.9	99.2
1999.....	0.6	0.8	0.9	1.5	2.4	4.5	8.1	18.4	36.0	97.2
White, both sexes										
2002.....	1.8	2.3	2.8	3.4	5.1	8.1	14.4	28.3	57.1	138.0
2001.....	1.7	2.5	2.8	3.2	4.9	8.0	12.8	26.3	52.4	128.0
2000.....	1.7	2.1	2.5	3.2	4.6	6.2	12.3	23.1	46.6	119.1
1999.....	1.7	2.2	2.1	2.9	4.3	6.9	11.4	23.0	47.7	115.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)										
White, male										
2002.....	3.0	3.7	4.2	5.1	7.2	10.8	19.3	36.4	71.2	166.4
2001.....	2.8	4.1	4.4	4.9	7.0	10.8	16.5	35.1	66.6	162.3
2000.....	2.7	3.5	3.9	4.8	6.9	8.4	16.2	30.3	59.7	152.8
1999.....	2.7	3.7	3.3	4.4	6.6	9.6	15.3	28.3	64.2	149.1
White, female										
2002.....	0.7	1.0	1.5	1.8	3.1	5.8	10.3	22.6	48.5	125.6
2001.....	0.5	0.9	1.2	1.6	2.9	5.6	9.9	20.0	43.9	113.6
2000.....	0.6	0.7	1.2	1.8	2.6	4.3	9.3	18.1	38.8	105.4
1999.....	0.6	0.8	0.9	1.5	2.2	4.6	8.3	19.3	37.9	102.6
Black, both sexes										
2002.....	1.6	2.5	3.2	3.8	4.7	6.1	10.9	14.8	18.1	49.3
2001.....	2.0	2.6	3.3	3.9	4.9	7.9	9.7	16.3	21.9	44.5
2000.....	1.6	2.0	3.6	4.5	5.1	7.1	9.7	11.6	20.3	38.8
1999.....	1.7	3.2	3.6	4.5	6.7	9.0	9.5	13.4	23.1	45.3
Black, male										
2002.....	2.6	4.1	5.6	6.3	7.3	9.2	15.4	22.0	28.3	77.2
2001.....	3.3	4.9	6.0	6.4	7.7	11.9	13.5	24.2	36.1	68.1
2000.....	2.6	3.4	6.3	7.7	8.7	10.7	12.4	18.4	26.1	61.1
1999.....	3.3	5.6	6.0	7.6	10.2	14.8	14.6	20.4	31.4	58.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Falls (W00-W19)

Black, female

2002.....	*	*	*	*	*	3.7	7.8	10.4	12.7	38.6
2001.....	*	*	*	*	*	4.9	7.2	11.6	14.7	35.8
2000.....	*	*	*	*	*	4.4	8.0	7.4	17.4	30.5
1999.....	*	*	*	*	4.0	4.6	6.1	9.2	19.0	40.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Fall on same level (W00-W09,W18)										
All races 3/, both sexes										
2002.....	0.3	0.5	0.7	0.9	1.3	2.2	4.2	8.9	17.8	50.8
2001.....	0.2	0.5	0.6	0.8	1.2	1.9	3.3	6.8	14.4	37.3
2000.....	0.2	0.3	0.5	0.6	1.0	1.4	2.7	5.5	11.9	33.0
1999.....	0.1	0.3	0.3	0.4	0.5	1.1	1.9	4.2	8.2	21.4
All races 3/, male										
2002.....	0.5	0.8	1.0	1.2	1.7	2.9	5.7	11.0	21.6	60.8
2001.....	0.3	0.8	0.8	1.0	1.7	2.4	3.7	9.0	18.5	45.3
2000.....	0.4	0.4	0.8	0.8	1.2	1.7	3.4	6.6	15.2	39.6
1999.....	0.2	0.5	0.5	0.5	0.6	1.5	2.5	5.3	10.5	26.6
All races 3/, female										
2002.....	*	0.2	0.4	0.7	0.9	1.6	2.9	7.3	15.5	46.4
2001.....	*	0.3	0.4	0.5	0.8	1.4	2.9	5.2	11.9	33.9
2000.....	*	*	0.3	0.5	0.8	1.1	2.2	4.7	10.0	30.4
1999.....	*	*	*	*	0.4	0.8	1.4	3.5	6.8	19.4
White, both sexes										
2002.....	0.3	0.5	0.7	0.9	1.4	2.3	4.3	9.2	18.8	53.7
2001.....	0.2	0.6	0.6	0.7	1.3	2.0	3.4	6.9	15.1	39.3
2000.....	0.3	0.3	0.5	0.6	1.0	1.4	2.8	5.8	12.5	35.0
1999.....	0.2	0.3	0.3	0.3	0.4	1.1	2.0	4.3	8.4	22.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Fall from one level to another (W10-W17)										
All races 3/, both sexes										
2002.....	0.8	1.0	0.9	1.3	1.7	2.3	3.4	5.1	7.5	10.9
2001.....	0.8	1.1	1.1	1.3	1.5	2.3	3.0	5.1	7.6	11.3
2000.....	0.9	1.0	1.0	1.2	1.5	1.9	3.0	4.2	6.5	8.8
1999.....	0.9	1.0	0.9	1.3	1.9	2.3	2.9	4.8	7.3	10.4
All races 3/, male										
2002.....	1.5	1.7	1.5	2.2	2.8	3.5	5.3	8.2	12.0	17.4
2001.....	1.4	2.0	1.9	2.2	2.6	3.9	4.6	8.4	11.4	18.2
2000.....	1.5	1.7	1.8	2.1	2.4	2.9	4.9	7.0	10.8	16.4
1999.....	1.5	1.7	1.5	2.1	3.1	4.0	4.2	6.9	12.7	17.8
All races 3/, female										
2002.....	0.2	0.2	0.4	0.4	0.7	1.2	1.9	3.0	4.8	8.1
2001.....	*	0.2	0.3	0.4	0.5	1.0	1.8	2.8	5.3	8.3
2000.....	0.2	0.3	0.3	0.5	0.6	1.0	1.5	2.2	3.9	5.7
1999.....	0.2	0.2	0.2	0.5	0.7	0.9	1.8	3.4	4.2	7.4
White, both sexes										
2002.....	0.9	1.0	1.0	1.3	1.8	2.4	3.6	5.4	7.9	11.2
2001.....	0.8	1.1	1.1	1.3	1.5	2.4	3.2	5.3	8.1	11.7
2000.....	0.9	1.0	1.0	1.3	1.5	1.8	3.1	4.4	6.8	9.2
1999.....	0.9	1.0	0.8	1.3	2.0	2.4	3.0	5.0	7.7	10.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified fall (W19)										
All races 3/, both sexes										
2002.....	0.6	0.8	1.2	1.3	2.0	3.5	6.3	13.1	28.6	68.8
2001.....	0.6	0.9	1.1	1.3	2.1	3.6	6.1	13.4	27.9	73.0
2000.....	0.5	0.8	1.1	1.4	2.2	3.0	6.3	12.5	26.1	70.7
1999.....	0.7	1.0	1.0	1.4	2.2	3.5	6.5	13.2	30.0	78.4
All races 3/, male										
2002.....	0.9	1.1	1.8	1.9	2.6	4.3	7.8	15.9	34.3	80.7
2001.....	1.0	1.3	1.7	1.8	2.8	4.3	7.6	16.6	34.2	92.1
2000.....	0.8	1.2	1.6	2.1	3.3	4.0	7.5	15.9	31.3	89.4
1999.....	1.0	1.6	1.5	2.1	3.2	4.3	8.5	15.7	38.4	98.1
All races 3/, female										
2002.....	0.3	0.6	0.6	0.7	1.4	2.8	5.1	11.2	25.2	63.6
2001.....	0.3	0.4	0.5	0.7	1.5	3.0	4.8	11.1	24.2	64.9
2000.....	0.3	0.3	0.6	0.8	1.2	2.2	5.5	10.1	23.1	63.1
1999.....	0.3	0.4	0.5	0.8	1.3	2.7	4.9	11.5	25.0	70.4
White, both sexes										
2002.....	0.6	0.8	1.1	1.2	1.9	3.4	6.4	13.7	30.4	73.0
2001.....	0.6	0.9	1.1	1.2	2.1	3.6	6.3	14.0	29.2	77.0
2000.....	0.5	0.8	1.0	1.4	2.1	3.0	6.4	12.9	27.3	74.9
1999.....	0.6	0.9	1.0	1.3	1.9	3.4	6.4	13.7	31.6	82.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified fall (W19)										
White, male										
2002.....	0.9	1.1	1.6	1.8	2.3	4.1	7.9	16.6	36.1	85.1
2001.....	1.0	1.3	1.7	1.7	2.8	4.2	7.9	17.3	35.3	95.7
2000.....	0.7	1.3	1.5	1.9	3.1	3.9	7.6	16.2	32.6	94.0
1999.....	1.0	1.5	1.4	1.8	2.8	4.1	8.4	15.9	40.3	101.9
White, female										
2002.....	0.3	0.6	0.6	0.7	1.4	2.9	5.3	11.7	27.0	67.8
2001.....	0.2	0.4	0.5	0.7	1.5	3.0	5.0	11.7	25.5	69.2
2000.....	0.3	0.3	0.6	0.8	1.2	2.2	5.5	10.5	24.2	67.2
1999.....	0.3	0.3	0.5	0.7	1.1	2.8	4.8	12.1	26.4	74.4
Black, both sexes										
2002.....	*	1.1	2.0	1.8	3.0	3.6	5.6	7.6	10.5	25.2
2001.....	0.8	1.0	1.2	1.8	2.5	4.8	4.7	8.9	13.8	26.4
2000.....	0.9	0.9	1.5	2.1	2.8	3.8	6.3	8.0	13.3	25.7
1999.....	0.8	1.9	1.6	2.6	4.5	4.9	6.4	8.6	15.0	33.5
Black, male										
2002.....	*	1.8	3.7	3.0	5.1	5.2	8.0	*	*	38.6
2001.....	*	1.9	*	*	*	6.1	*	12.5	20.9	46.5
2000.....	*	*	2.7	3.6	5.2	5.2	7.4	11.8	18.5	38.1
1999.....	*	3.2	3.0	4.4	6.8	7.5	9.2	12.8	19.2	46.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified fall (W19)										
Black, female										
2002.....	*	*	*	*	*	*	*	7.1	8.4	20.1
2001.....	*	*	*	*	*	3.8	*	6.7	10.2	18.9
2000.....	*	*	*	*	*	*	5.5	5.7	10.6	21.0
1999.....	*	*	*	*	*	*	4.5	6.0	13.0	28.5

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental discharge of firearms (W32-W34)										
All races 3/, both sexes										
2002.....	0.3	0.2	0.2	0.2	0.2	*	*	*	*	*
2001.....	0.3	0.3	0.2	0.3	0.2	*	*	0.3	*	*
2000.....	0.3	0.2	0.2	0.3	*	0.3	*	*	*	*
1999.....	0.3	0.3	0.2	0.2	0.2	*	*	*	*	*
All races 3/, male										
2002.....	0.5	0.4	0.4	0.4	*	*	*	*	*	*
2001.....	0.6	0.5	0.4	0.5	0.4	*	*	0.7	*	*
2000.....	0.6	0.3	0.4	0.5	*	0.5	*	*	*	*
1999.....	0.5	0.5	0.4	0.3	*	*	*	*	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.3	0.2	0.2	0.2	*	*	*	*	*	*
2001.....	0.3	0.3	0.3	0.3	0.3	*	*	0.3	*	*
2000.....	0.4	0.2	0.2	0.3	*	0.3	*	*	*	*
1999.....	0.3	0.3	0.3	0.2	0.2	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental drowning and submersion (W65-W74)										
All races 3/, both sexes										
2002.....	1.1	1.0	0.9	0.8	0.8	0.8	1.0	1.2	1.5	1.7
2001.....	1.1	1.0	0.8	0.8	0.9	0.8	0.9	1.1	1.4	1.7
2000.....	1.0	0.9	1.0	0.9	0.9	0.8	1.1	1.0	1.7	1.5
1999.....	1.0	1.0	1.0	0.9	0.9	0.9	1.1	1.2	1.4	1.6
All races 3/, male										
2002.....	1.9	1.6	1.5	1.3	1.2	1.4	1.8	1.8	2.5	3.6
2001.....	1.7	1.8	1.3	1.2	1.4	1.4	1.6	1.8	2.4	3.5
2000.....	1.7	1.6	1.7	1.6	1.5	1.4	1.8	1.2	2.7	2.8
1999.....	1.7	1.7	1.8	1.5	1.5	1.5	1.7	2.1	2.7	2.3
All races 3/, female										
2002.....	0.4	0.4	0.3	0.3	0.4	*	0.4	0.7	0.9	0.9
2001.....	0.4	0.3	0.3	0.4	0.4	*	*	0.7	0.8	1.0
2000.....	0.3	0.3	0.3	*	0.4	*	0.6	0.8	1.1	1.0
1999.....	0.3	0.3	0.3	0.4	0.4	0.4	0.7	0.6	0.7	1.4
White, both sexes										
2002.....	1.2	0.9	0.8	0.8	0.8	0.8	0.9	1.1	1.4	1.6
2001.....	1.0	1.0	0.8	0.7	0.9	0.8	0.9	1.1	1.3	1.7
2000.....	1.0	0.9	0.9	0.9	0.9	0.8	1.1	0.9	1.8	1.5
1999.....	1.0	1.1	1.0	0.9	0.9	0.9	1.0	1.1	1.4	1.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)										
All races 3/, both sexes										
2002.....	0.4	0.5	0.8	1.1	1.3	2.6	3.6	7.3	12.6	28.4
2001.....	0.5	0.8	1.0	1.2	1.5	2.3	3.6	7.2	12.8	30.0
2000.....	0.4	0.7	0.9	1.2	1.6	2.3	4.0	7.7	13.4	32.6
1999.....	0.5	0.5	0.8	1.0	1.4	2.2	4.0	6.6	12.1	30.3
All races 3/, male										
2002.....	0.6	0.7	0.9	1.5	1.7	3.4	4.4	9.6	16.0	37.0
2001.....	0.6	1.0	1.2	1.6	1.6	2.9	4.3	9.2	17.5	39.8
2000.....	0.5	1.0	1.2	1.4	1.9	2.8	4.5	9.7	18.9	41.0
1999.....	0.6	0.6	1.2	1.1	1.6	2.5	5.2	7.9	15.9	39.2
All races 3/, female										
2002.....	0.3	0.4	0.6	0.8	1.0	2.0	2.9	5.6	10.6	24.6
2001.....	0.3	0.6	0.8	0.8	1.3	1.8	3.1	5.8	9.9	26.0
2000.....	0.4	0.4	0.6	0.9	1.4	1.8	3.5	6.3	10.1	29.1
1999.....	0.3	0.5	0.4	0.8	1.2	2.0	3.0	5.6	9.9	26.7
White, both sexes										
2002.....	0.4	0.5	0.7	1.1	1.2	2.3	3.4	7.1	12.5	28.3
2001.....	0.4	0.7	1.0	1.1	1.4	2.3	3.3	6.8	12.4	29.7
2000.....	0.5	0.7	0.8	1.1	1.5	2.1	3.7	7.6	13.3	32.5
1999.....	0.5	0.5	0.8	0.9	1.2	2.1	3.7	6.2	11.7	30.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

White, male

2002.....	0.6	0.7	0.8	1.4	1.6	2.9	4.3	9.4	15.6	37.2
2001.....	0.6	0.9	1.1	1.3	1.5	2.8	3.9	8.6	16.4	39.3
2000.....	0.6	0.9	1.1	1.4	1.8	2.5	4.2	9.5	18.4	40.2
1999.....	0.6	0.6	1.1	1.0	1.3	2.4	4.9	7.3	15.4	39.7

White, female

2002.....	0.3	0.4	0.6	0.8	0.9	1.8	2.7	5.6	10.6	24.5
2001.....	0.3	0.6	0.9	0.8	1.2	1.8	2.9	5.6	10.0	25.7
2000.....	0.4	0.5	0.6	0.9	1.3	1.7	3.3	6.3	10.2	29.4
1999.....	0.3	0.5	0.4	0.7	1.0	1.9	2.7	5.3	9.5	26.7

Black, both sexes

2002.....	*	*	1.4	1.9	2.2	5.8	7.1	11.0	16.0	33.0
2001.....	*	1.3	1.0	2.3	2.8	3.6	7.5	11.5	18.7	36.8
2000.....	*	0.8	1.5	1.9	2.6	5.0	7.2	10.8	16.7	35.3
1999.....	*	1.0	1.2	2.1	3.0	3.6	7.7	12.3	18.2	33.2

Black, male

2002.....	*	*	2.2	*	*	8.7	8.0	16.5	23.7	41.8
2001.....	*	*	*	3.6	*	*	9.8	15.8	36.1	51.1
2000.....	*	*	2.4	*	*	6.5	8.4	15.6	27.8	51.9
1999.....	*	*	*	*	*	*	10.5	17.6	25.3	43.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

Black, female

2002.....	*	*	*	*	*	3.7	6.5	7.7	11.9	29.6
2001.....	*	*	*	*	*	*	6.1	8.9	9.8	31.6
2000.....	*	*	*	*	*	3.8	6.4	8.0	11.0	29.2
1999.....	*	*	*	*	*	*	5.9	9.2	14.7	29.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other accidental and unspecified threats to breathing (W75-W77,W81-W84)										
All races 3/, both sexes										
2002.....	0.3	0.3	0.3	0.3	0.2	0.3	0.2	0.4	*	1.0
2001.....	0.4	0.4	0.2	0.2	0.2	0.2	0.4	0.4	0.4	0.9
2000.....	0.3	0.2	0.3	0.4	0.3	0.3	0.4	0.5	0.5	1.2
1999.....	0.4	0.4	0.3	0.3	0.3	0.5	0.8	1.4	1.8	4.7
All races 3/, male										
2002.....	0.5	0.5	0.5	0.4	*	*	*	*	*	*
2001.....	0.6	0.6	0.4	0.3	*	*	*	*	*	*
2000.....	0.6	0.4	0.6	0.6	0.5	*	*	*	*	*
1999.....	0.6	0.6	0.6	0.6	*	0.7	1.1	1.9	1.8	5.4
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	1.1
2001.....	*	*	*	*	*	*	*	*	*	0.9
2000.....	*	*	*	*	*	*	*	*	*	1.1
1999.....	*	*	*	*	*	*	0.6	1.1	1.7	4.5
White, both sexes										
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	*	0.4	*	1.0
2001.....	0.4	0.4	0.3	0.2	*	0.3	0.4	0.5	0.4	0.9
2000.....	0.4	0.3	0.4	0.4	0.3	0.3	0.3	0.5	0.6	1.2
1999.....	0.4	0.4	0.4	0.3	0.3	0.5	0.8	1.3	1.8	4.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental exposure to electric current (W85-W87)

All races 3/, both sexes

2002.....	0.3	0.2	0.2	*	*	*	*	*	*	*
2001.....	0.3	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	0.2	0.1	*	*	*	*	*	*	*
1999.....	0.3	0.2	*	0.2	*	*	*	*	*	*

All races 3/, male

2002.....	0.5	0.4	0.3	*	*	*	*	*	*	*
2001.....	0.5	0.4	*	*	*	*	*	*	*	*
2000.....	0.4	0.5	0.2	*	*	*	*	*	*	*
1999.....	0.5	0.4	*	0.4	*	*	*	*	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.3	0.2	0.2	*	*	*	*	*	*	*
2001.....	0.3	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	0.2	*	*	*	*	*	*	*	*
1999.....	0.3	0.2	*	0.2	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)										
All races 3/, both sexes										
2002.....	0.9	1.0	1.3	1.2	1.4	1.8	2.4	2.8	4.2	4.8
2001.....	1.1	1.0	1.1	1.5	1.5	2.0	2.5	3.0	3.8	5.7
2000.....	1.0	1.1	1.3	1.4	1.7	2.0	2.4	3.5	4.3	6.2
1999.....	1.0	1.0	1.0	1.2	1.5	2.0	2.5	3.5	4.7	5.9
All races 3/, male										
2002.....	1.3	1.4	1.6	1.5	2.0	2.6	3.2	3.5	6.1	7.0
2001.....	1.6	1.5	1.5	2.1	1.8	2.5	3.0	3.5	5.1	8.3
2000.....	1.2	1.5	1.8	1.9	2.2	2.4	3.2	4.5	5.9	9.4
1999.....	1.5	1.5	1.3	1.6	1.9	2.5	3.4	4.3	6.6	9.3
All races 3/, female										
2002.....	0.6	0.6	0.9	0.9	0.8	1.1	1.8	2.4	3.0	3.9
2001.....	0.7	0.6	0.7	1.0	1.2	1.5	2.1	2.7	3.1	4.6
2000.....	0.7	0.7	0.7	0.9	1.2	1.6	1.8	2.8	3.4	4.9
1999.....	0.5	0.5	0.7	0.9	1.2	1.6	1.8	3.0	3.6	4.5
White, both sexes										
2002.....	0.9	0.9	1.1	1.1	1.1	1.6	2.1	2.5	3.7	4.0
2001.....	1.0	0.9	1.0	1.3	1.3	1.7	2.1	2.6	3.2	4.8
2000.....	0.9	0.9	1.2	1.1	1.4	1.7	2.0	2.8	3.5	5.1
1999.....	0.9	0.9	0.9	1.1	1.2	1.6	2.0	2.8	3.9	4.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)										
White, male										
2002.....	1.3	1.3	1.4	1.4	1.6	2.2	2.7	3.1	5.8	5.8
2001.....	1.4	1.2	1.2	1.8	1.5	2.1	2.4	2.8	4.4	7.0
2000.....	1.1	1.3	1.7	1.6	1.8	2.0	2.7	3.6	4.8	8.2
1999.....	1.4	1.3	1.2	1.4	1.7	2.0	2.6	3.3	5.8	8.0
White, female										
2002.....	0.6	0.6	0.9	0.8	0.6	1.0	1.6	2.1	2.5	3.2
2001.....	0.6	0.5	0.7	0.8	1.1	1.3	1.9	2.4	2.5	3.8
2000.....	0.6	0.6	0.7	0.7	1.1	1.4	1.5	2.3	2.7	3.9
1999.....	0.5	0.5	0.6	0.9	0.8	1.3	1.5	2.5	2.8	3.3
Black, both sexes										
2002.....	1.1	1.6	2.8	2.6	4.2	4.0	6.3	7.2	10.7	15.7
2001.....	2.1	2.3	2.5	3.7	3.3	5.2	7.3	8.5	11.6	17.8
2000.....	2.0	2.2	2.6	3.9	4.5	5.0	7.1	11.4	16.1	20.3
1999.....	1.7	2.0	2.3	2.6	4.3	5.7	8.3	12.0	15.3	21.7
Black, male										
2002.....	1.8	2.1	4.0	3.4	6.3	6.7	9.3	9.2	*	25.7
2001.....	2.9	3.7	4.2	5.0	4.9	7.4	11.1	12.1	*	27.2
2000.....	2.7	3.2	4.0	5.8	7.0	7.5	10.4	16.5	21.9	27.7
1999.....	2.5	3.0	3.3	4.1	4.2	7.8	13.2	17.1	20.1	29.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)										
Black, female										
2002.....	*	*	*	*	*	*	*	6.0	9.2	11.9
2001.....	1.3	*	*	2.7	*	*	4.7	6.4	10.2	14.3
2000.....	1.3	*	*	*	*	*	4.9	8.3	13.2	17.6
1999.....	*	*	*	*	4.3	4.2	5.0	8.9	13.0	18.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)										
All races 3/, both sexes										
2002.....	14.8	13.8	9.2	5.0	3.0	2.3	1.9	2.1	2.7	3.9
2001.....	12.0	10.9	7.0	3.7	2.4	1.7	1.5	2.1	2.4	3.5
2000.....	11.2	9.9	6.1	3.3	2.2	1.5	1.5	1.6	2.6	4.3
1999.....	11.1	9.8	5.4	3.4	2.0	1.7	1.6	1.9	2.6	3.4
All races 3/, male										
2002.....	20.3	18.7	12.8	6.8	3.6	2.8	2.1	2.3	3.2	5.1
2001.....	16.8	15.7	9.9	5.0	3.0	2.0	2.1	2.7	2.8	3.8
2000.....	16.4	14.7	8.9	4.0	2.7	1.8	1.7	1.6	2.9	6.2
1999.....	16.6	15.1	7.9	4.7	3.0	2.2	1.9	2.1	2.9	4.4
All races 3/, female										
2002.....	9.4	9.1	5.8	3.4	2.4	1.8	1.7	1.9	2.3	3.4
2001.....	7.3	6.2	4.1	2.6	1.9	1.4	1.0	1.6	2.2	3.3
2000.....	6.0	5.3	3.4	2.7	1.8	1.3	1.3	1.6	2.4	3.6
1999.....	5.7	4.7	3.1	2.2	1.2	1.2	1.4	1.7	2.4	3.0
White, both sexes										
2002.....	15.4	13.7	8.7	4.6	2.7	2.2	1.9	2.1	2.6	3.9
2001.....	12.1	10.5	6.1	3.4	2.1	1.6	1.4	2.0	2.3	3.3
2000.....	11.0	9.4	5.3	3.0	2.1	1.5	1.4	1.7	2.6	4.1
1999.....	10.8	9.1	4.8	3.1	1.7	1.5	1.4	1.9	2.6	3.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)										
White, male										
2002.....	21.0	18.3	11.7	5.8	2.9	2.6	2.0	2.3	3.0	5.2
2001.....	17.1	14.9	8.5	4.3	2.3	1.9	2.0	2.4	2.7	3.7
2000.....	16.2	13.7	7.6	3.3	2.4	1.6	1.6	1.7	2.9	5.9
1999.....	16.2	13.8	6.6	4.0	2.3	2.0	1.7	2.1	3.0	4.5
White, female										
2002.....	9.8	9.0	5.8	3.4	2.5	1.9	1.8	1.9	2.5	3.4
2001.....	7.1	6.1	3.8	2.6	1.9	1.4	1.0	1.7	2.0	3.1
2000.....	5.8	5.1	3.1	2.7	1.8	1.4	1.3	1.7	2.5	3.4
1999.....	5.4	4.5	3.0	2.3	1.2	1.1	1.2	1.8	2.4	2.8
Black, both sexes										
2002.....	15.7	19.4	16.1	10.3	6.0	3.0	*	*	*	*
2001.....	15.2	17.7	15.4	7.4	5.4	2.6	*	*	*	6.4
2000.....	15.3	16.5	14.2	6.8	3.9	2.7	*	*	*	8.1
1999.....	16.1	17.9	12.3	6.5	5.5	2.9	3.2	*	*	*
Black, male										
2002.....	21.6	27.2	25.6	17.7	11.3	5.5	*	*	*	*
2001.....	20.3	28.0	24.8	12.7	9.6	*	*	*	*	*
2000.....	22.8	26.3	23.1	11.5	7.0	*	*	*	*	*
1999.....	24.2	29.7	20.9	11.9	10.0	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)										
All races 3/, both sexes										
2002.....	14.2	13.2	8.7	4.4	2.5	1.8	1.3	1.5	1.9	3.1
2001.....	11.4	10.4	6.4	3.3	2.0	1.4	1.2	1.5	1.7	2.5
2000.....	10.6	9.3	5.6	2.8	1.7	1.2	1.0	1.2	1.7	3.3
1999.....	10.5	9.2	4.9	2.9	1.6	1.3	1.1	1.4	1.9	2.8
All races 3/, male										
2002.....	19.2	17.7	12.0	5.8	2.9	2.0	1.4	1.6	2.1	3.7
2001.....	15.8	14.9	9.1	4.4	2.4	1.5	1.5	1.8	1.7	1.8
2000.....	15.6	13.9	8.2	3.2	2.0	1.5	1.2	1.1	1.5	4.4
1999.....	15.5	14.2	7.1	4.0	2.2	1.6	1.3	1.3	1.7	3.3
All races 3/, female										
2002.....	9.2	8.9	5.6	3.2	2.1	1.5	1.3	1.4	1.8	2.8
2001.....	7.0	6.0	3.9	2.3	1.7	1.2	0.9	1.2	1.7	2.7
2000.....	5.8	4.9	3.2	2.4	1.4	1.0	0.9	1.3	1.9	2.8
1999.....	5.5	4.5	2.9	1.9	1.1	1.0	1.0	1.4	2.0	2.6
White, both sexes										
2002.....	14.8	13.1	8.2	4.1	2.2	1.7	1.4	1.5	1.9	3.2
2001.....	11.5	9.9	5.7	3.0	1.8	1.3	1.1	1.4	1.7	2.3
2000.....	10.5	8.8	4.9	2.5	1.5	1.2	1.0	1.3	1.7	3.1
1999.....	10.2	8.6	4.3	2.7	1.3	1.2	1.0	1.4	1.9	2.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

White, male

2002.....	19.9	17.4	10.9	5.0	2.2	1.8	1.3	1.5	1.9	3.8
2001.....	16.1	14.0	7.8	3.7	1.8	1.4	1.4	1.6	1.8	1.7
2000.....	15.3	12.9	6.9	2.6	1.6	1.2	1.0	1.2	1.4	4.1
1999.....	15.1	12.9	5.9	3.4	1.6	1.4	1.2	1.4	1.7	3.2

White, female

2002.....	9.5	8.8	5.6	3.2	2.3	1.6	1.4	1.5	2.0	2.9
2001.....	6.8	5.9	3.6	2.3	1.7	1.2	0.9	1.3	1.7	2.6
2000.....	5.6	4.7	2.9	2.5	1.4	1.1	0.9	1.3	1.9	2.6
1999.....	5.2	4.3	2.8	2.0	1.1	1.0	0.8	1.4	2.0	2.3

Black, both sexes

2002.....	15.0	18.5	15.4	9.1	5.5	2.4	*	*	*	*
2001.....	14.6	17.2	14.3	7.0	4.7	2.2	*	*	*	*
2000.....	14.6	15.8	13.5	5.9	3.4	2.3	*	*	*	6.3
1999.....	15.7	17.1	11.4	5.6	4.7	2.4	*	*	*	*

Black, male

2002.....	20.5	25.8	24.4	15.3	10.5	*	*	*	*	*
2001.....	19.2	27.2	23.1	12.2	8.1	*	*	*	*	*
2000.....	21.8	25.2	22.3	9.9	6.2	*	*	*	*	*
1999.....	23.5	28.3	19.5	10.3	8.5	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Black, female

2002.....	10.1	12.1	7.7	3.9	*	*	*	*	*	*
2001.....	10.4	8.5	6.8	2.8	*	*	*	*	*	*
2000.....	8.3	7.8	6.0	*	*	*	*	*	*	*
1999.....	8.8	7.4	4.6	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)

All races 3/, both sexes

2002.....	0.4	0.3	0.3	0.3	0.2	*	*	*	*	*
2001.....	0.4	0.3	0.3	0.2	0.2	*	*	*	*	*
2000.....	0.3	0.3	0.2	0.3	0.2	*	*	*	*	*
1999.....	0.4	0.3	0.2	0.3	0.2	*	*	*	*	*

All races 3/, male

2002.....	0.6	0.5	0.5	0.5	0.4	*	*	*	*	*
2001.....	0.5	0.4	0.5	*	*	*	*	*	*	*
2000.....	0.5	0.5	0.4	0.4	*	*	*	*	*	*
1999.....	0.6	0.5	0.4	0.3	*	*	*	*	*	*

All races 3/, female

2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.4	0.3	0.3	0.2	*	*	*	*	*	*
2001.....	0.4	0.3	0.3	0.2	*	*	*	*	*	*
2000.....	0.3	0.3	0.2	0.3	0.2	*	*	*	*	*
1999.....	0.4	0.3	0.2	0.2	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning by and exposure to other gases and vapors (X47)										
All races 3/, both sexes										
2002.....	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.5	0.5	0.6
2001.....	0.2	0.3	0.2	0.2	0.2	*	0.2	0.5	0.6	0.9
2000.....	0.2	0.3	0.2	0.3	0.3	*	0.3	0.3	0.7	0.8
1999.....	0.3	0.2	0.2	0.3	0.2	0.2	0.3	0.3	0.5	0.5
All races 3/, male										
2002.....	0.5	0.4	0.4	0.4	0.4	0.6	0.6	*	1.0	*
2001.....	0.4	0.4	0.4	0.3	*	*	*	0.8	*	1.7
2000.....	0.4	0.4	0.4	0.4	0.4	*	*	*	1.2	*
1999.....	0.5	0.4	0.4	0.4	0.4	*	*	*	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.3	0.3	0.3	0.2	0.3	0.4	0.4	0.5	0.5	0.7
2001.....	0.2	0.3	0.2	0.2	*	*	*	0.4	0.5	0.8
2000.....	0.2	0.3	0.2	0.2	0.3	*	0.3	*	0.7	0.7
1999.....	0.3	0.2	0.2	0.3	*	*	*	0.3	0.5	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)										
All races 3/, both sexes										
2002.....	2.2	2.2	2.7	3.2	3.9	5.5	7.0	13.5	25.6	75.8
2001.....	2.0	2.5	2.9	2.9	4.3	4.9	8.0	14.4	28.4	83.4
2000.....	2.3	2.6	2.7	3.1	4.1	5.0	7.6	13.6	26.9	82.6
1999.....	2.4	2.5	2.9	3.2	4.3	5.9	8.1	15.3	32.3	92.8
All races 3/, male										
2002.....	3.5	3.8	4.2	5.1	6.1	7.9	9.8	17.8	30.3	79.9
2001.....	3.3	4.2	4.7	4.4	6.6	7.1	10.6	19.0	33.9	92.4
2000.....	3.9	4.4	4.5	5.2	6.1	7.1	10.0	17.5	31.1	91.7
1999.....	3.9	4.1	4.9	5.2	6.6	8.2	11.1	19.5	40.3	107.0
All races 3/, female										
2002.....	0.8	0.7	1.2	1.4	1.8	3.3	4.7	10.4	22.7	74.0
2001.....	0.7	0.9	1.3	1.4	2.2	3.1	5.8	11.2	25.1	79.6
2000.....	0.7	0.8	0.9	1.2	2.2	3.2	5.6	10.9	24.3	78.9
1999.....	0.9	0.9	1.0	1.4	2.2	3.9	5.8	12.4	27.6	87.1
White, both sexes										
2002.....	2.2	2.2	2.6	3.1	3.8	5.1	7.1	13.4	25.9	79.1
2001.....	1.9	2.4	2.8	2.9	4.1	4.8	7.9	14.6	29.2	86.4
2000.....	2.2	2.4	2.6	3.0	4.0	4.8	7.5	13.4	27.5	86.2
1999.....	2.3	2.2	2.8	3.0	4.0	5.6	7.8	15.0	32.6	96.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)										
White, male										
2002.....	3.6	3.8	4.1	4.8	6.0	7.3	9.9	17.6	30.7	82.5
2001.....	3.1	4.1	4.4	4.4	6.3	6.9	10.5	19.3	34.4	95.8
2000.....	3.7	4.0	4.3	4.7	5.8	6.7	10.0	16.9	31.4	95.0
1999.....	3.7	3.6	4.6	4.8	6.3	7.7	10.5	19.0	40.5	108.9
White, female										
2002.....	0.7	0.6	1.2	1.5	1.8	3.1	4.8	10.4	23.0	77.6
2001.....	0.6	0.8	1.3	1.5	2.0	3.1	5.8	11.3	26.0	82.4
2000.....	0.6	0.7	0.9	1.3	2.3	3.2	5.4	11.0	25.3	82.6
1999.....	0.9	0.9	1.0	1.4	1.9	3.8	5.7	12.2	27.9	91.0
Black, both sexes										
2002.....	2.4	3.0	3.3	4.7	4.7	9.3	6.3	15.6	22.8	46.9
2001.....	2.8	3.5	4.3	2.8	6.8	6.4	9.6	15.1	23.0	58.3
2000.....	3.1	4.2	3.6	5.3	5.4	6.8	9.3	17.8	21.2	51.0
1999.....	3.4	4.7	4.6	5.7	7.1	9.4	13.1	20.7	32.6	67.0
Black, male										
2002.....	3.2	4.6	5.5	8.6	7.9	13.5	9.0	22.5	29.8	57.9
2001.....	4.7	5.4	7.3	5.0	10.0	10.4	13.1	18.6	32.9	62.4
2000.....	5.4	7.8	7.1	10.4	9.3	10.9	11.7	27.9	32.9	69.2
1999.....	5.9	8.4	7.8	9.8	10.8	15.1	20.4	28.0	41.0	106.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

Black, female

2002.....	1.8	1.6	*	*	*	6.3	4.5	11.5	19.1	42.7
2001.....	*	1.7	*	*	4.3	*	7.2	13.0	18.0	56.8
2000.....	*	*	*	*	*	3.8	7.8	11.7	15.3	44.2
1999.....	*	*	*	*	4.2	5.2	8.3	16.3	28.5	52.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
All races 3/, both sexes										
2002.....	16.1	16.3	15.1	14.6	12.3	12.5	14.6	16.6	19.3	18.0
2001 4/.....	15.2	15.7	14.6	14.0	12.0	12.7	13.9	16.4	18.9	17.5
2000.....	15.4	14.9	13.9	12.7	11.4	11.3	13.8	16.5	19.3	19.6
1999.....	14.3	14.3	13.4	12.8	11.4	12.4	14.5	17.4	19.2	19.3
All races 3/, male										
2002.....	24.7	24.9	23.8	23.5	20.4	22.0	27.9	34.3	44.0	50.7
2001 4/.....	23.6	24.3	22.4	22.3	19.5	22.7	26.7	33.8	44.2	51.1
2000.....	23.9	23.3	21.4	20.1	18.7	19.8	26.1	34.9	44.8	57.5
1999.....	22.1	22.3	20.6	20.3	19.1	22.1	27.7	35.6	44.1	57.1
All races 3/, female										
2002.....	7.7	8.0	6.7	6.2	5.0	4.3	3.9	4.1	4.4	3.8
2001 4/.....	6.9	7.4	7.0	6.2	5.2	4.1	3.7	4.3	3.8	3.4
2000.....	7.0	6.7	6.8	5.8	4.9	4.0	4.1	3.8	4.2	4.2
1999.....	6.7	6.7	6.5	5.8	4.4	4.2	4.1	4.8	4.6	4.1
White, both sexes										
2002.....	18.2	18.4	16.8	16.1	13.6	13.7	15.7	17.6	20.9	19.1
2001 4/.....	16.9	17.6	16.2	15.3	13.1	13.8	14.9	17.6	20.0	18.6
2000.....	17.0	16.5	15.3	13.9	12.6	12.3	14.7	17.6	20.4	21.0
1999.....	16.0	15.9	14.8	14.0	12.3	13.3	15.6	18.6	20.4	20.7

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
White, male										
2002.....	27.6	27.6	26.4	25.7	22.5	23.8	30.1	36.3	47.2	53.9
2001 4/.....	26.1	27.1	24.7	24.3	21.0	24.5	28.3	36.0	46.8	55.0
2000.....	26.2	25.6	23.2	21.8	20.4	21.3	27.6	37.1	47.7	61.6
1999.....	24.5	24.4	22.3	21.9	20.5	23.6	29.5	37.9	46.8	61.5
White, female										
2002.....	8.7	9.3	7.4	6.9	5.3	4.8	3.9	4.2	4.7	4.0
2001 4/.....	7.8	8.1	7.9	6.8	5.8	4.3	4.0	4.6	3.8	3.4
2000.....	7.9	7.5	7.5	6.4	5.5	4.4	4.4	3.9	4.2	4.5
1999.....	7.5	7.4	7.4	6.4	4.8	4.4	4.5	5.0	4.8	4.3
Black, both sexes										
2002.....	7.1	5.9	5.9	4.6	5.0	3.5	6.6	6.7	*	5.9
2001 4/.....	6.8	6.5	5.9	5.8	4.4	4.6	6.5	5.2	6.5	*
2000.....	8.0	6.6	6.4	4.6	3.7	4.3	6.1	5.3	6.5	*
1999.....	6.8	6.4	4.7	5.2	4.7	6.4	5.1	6.2	*	*
Black, male										
2002.....	11.5	11.0	9.2	8.6	8.1	7.7	12.4	15.6	*	*
2001 4/.....	10.8	10.1	9.6	10.2	8.3	8.1	14.1	12.1	16.1	*
2000.....	13.4	11.8	10.5	8.4	6.4	9.4	13.4	11.8	*	*
1999.....	10.8	10.9	8.7	9.3	8.9	12.2	11.2	14.2	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

All races 3/, both sexes

2002.....	2.9	3.1	2.6	2.0	1.4	1.0	1.0	1.0	1.1	0.8
2001.....	2.7	2.9	2.4	1.7	1.2	1.0	0.9	0.9	0.6	0.9
2000.....	2.6	2.5	2.3	1.8	1.0	0.7	0.9	0.8	0.9	1.0
1999.....	2.5	2.4	2.2	1.6	1.1	1.0	0.9	1.1	1.1	0.8

All races 3/, male

2002.....	2.8	2.7	2.6	1.7	1.2	0.9	0.9	1.1	1.3	*
2001.....	2.8	2.8	2.0	1.4	0.8	1.2	0.8	0.8	*	*
2000.....	2.5	2.5	2.0	1.6	0.7	0.5	0.8	0.9	1.3	*
1999.....	2.5	2.5	2.0	1.4	1.0	1.1	0.9	0.7	1.2	*

All races 3/, female

2002.....	3.1	3.5	2.5	2.3	1.6	1.1	1.1	0.9	0.9	0.7
2001.....	2.6	3.1	2.7	1.9	1.5	0.9	1.0	0.9	0.6	0.6
2000.....	2.8	2.5	2.5	1.9	1.3	0.7	1.1	0.8	0.7	0.9
1999.....	2.5	2.4	2.4	1.8	1.2	0.9	0.9	1.3	1.0	*

White, both sexes

2002.....	3.4	3.6	2.9	2.3	1.5	1.1	1.0	1.1	1.1	0.9
2001.....	3.1	3.3	2.7	1.9	1.3	1.1	1.0	1.0	0.7	0.9
2000.....	3.0	2.8	2.5	2.0	1.2	0.7	1.1	0.9	1.0	1.1
1999.....	2.8	2.8	2.5	1.8	1.2	1.1	1.0	1.1	1.2	0.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

All races 3/, both sexes

2002.....	0.9	1.0	0.9	0.8	0.4	0.4	0.4	0.6	0.7	0.8
2001.....	0.9	1.0	0.8	0.9	0.5	0.5	0.4	0.6	0.7	0.8
2000.....	0.9	1.1	0.8	0.8	0.5	0.4	0.4	0.5	0.7	0.8
1999.....	1.0	1.0	0.9	0.7	0.6	0.3	0.4	0.8	0.9	1.2

All races 3/, male

2002.....	1.5	1.7	1.3	1.3	0.7	0.7	0.6	1.1	1.5	2.2
2001.....	1.4	1.6	1.2	1.3	0.7	0.8	0.7	1.0	1.5	1.9
2000.....	1.4	1.7	1.2	1.3	0.7	0.6	0.6	1.0	1.6	2.3
1999.....	1.5	1.5	1.2	1.0	0.8	0.6	0.6	1.3	1.7	3.9

All races 3/, female

2002.....	0.4	0.4	0.4	0.3	*	*	*	*	*	*
2001.....	0.4	0.4	0.4	0.6	*	*	*	*	*	*
2000.....	0.4	0.4	0.5	0.4	*	*	*	*	*	*
1999.....	0.4	0.5	0.5	0.4	0.4	*	*	*	*	*

White, both sexes

2002.....	1.1	1.2	1.0	0.9	0.4	0.5	0.5	0.7	0.8	0.9
2001.....	1.1	1.1	1.0	1.0	0.6	0.6	0.5	0.7	0.8	0.9
2000.....	1.1	1.2	0.9	0.9	0.6	0.4	0.4	0.5	0.8	0.9
1999.....	1.1	1.1	1.0	0.8	0.7	0.4	0.4	0.8	0.9	1.3

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)										
All races 3/, both sexes										
2002.....	3.4	3.0	2.4	1.9	1.4	1.4	1.3	1.5	1.8	2.3
2001.....	3.2	2.7	2.1	1.7	1.4	1.3	1.3	1.4	2.1	2.1
2000.....	3.1	2.5	1.9	1.5	1.3	1.1	1.2	1.7	2.0	2.9
1999.....	2.8	2.1	1.9	1.4	1.3	1.4	1.3	1.8	2.0	2.8
All races 3/, male										
2002.....	5.6	5.0	3.8	3.3	2.4	2.4	2.2	2.6	3.0	5.2
2001.....	5.5	4.7	3.6	2.9	2.1	2.2	2.4	2.3	4.3	5.2
2000.....	5.4	4.3	3.2	2.4	2.0	1.5	2.1	3.0	3.8	7.2
1999.....	4.7	3.6	3.0	2.5	2.1	2.1	2.2	2.9	3.6	6.6
All races 3/, female										
2002.....	1.1	1.0	1.0	0.6	0.6	0.6	0.6	0.7	1.0	1.0
2001.....	0.9	0.8	0.7	0.6	0.7	0.6	*	0.7	0.8	0.8
2000.....	0.9	0.8	0.7	0.7	0.7	0.8	0.5	0.8	1.0	1.2
1999.....	0.8	0.7	0.7	0.4	0.6	0.8	0.5	1.0	1.1	1.3
White, both sexes										
2002.....	3.7	3.2	2.6	2.0	1.6	1.3	1.3	1.5	1.8	2.3
2001.....	3.5	3.0	2.2	1.8	1.3	1.3	1.2	1.4	2.1	2.1
2000.....	3.3	2.7	2.0	1.6	1.4	1.1	1.2	1.6	2.0	2.9
1999.....	3.0	2.3	1.9	1.5	1.3	1.4	1.3	1.7	2.1	2.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
All races 3/, both sexes										
2002.....	7.4	7.8	7.9	8.6	8.1	8.8	10.7	12.4	14.3	12.0
2001.....	7.0	7.5	8.0	8.4	8.0	9.0	10.3	12.4	14.1	12.2
2000.....	7.4	7.4	7.8	7.4	7.7	8.3	10.3	12.4	14.0	13.2
1999.....	6.7	7.4	7.2	8.1	7.5	8.7	10.9	12.9	13.5	12.6
All races 3/, male										
2002.....	12.6	13.3	14.0	15.4	14.8	17.0	22.5	27.8	35.9	37.8
2001.....	12.0	12.8	13.8	14.9	14.2	17.3	21.4	27.7	35.5	38.9
2000.....	12.6	12.8	13.3	13.1	13.9	15.9	21.3	28.1	35.4	42.9
1999.....	11.2	12.6	12.4	14.0	14.0	16.9	22.4	29.1	34.6	41.6
All races 3/, female										
2002.....	2.3	2.4	2.1	2.3	2.0	1.8	1.3	1.5	1.2	0.8
2001.....	2.2	2.3	2.4	2.4	2.3	1.9	1.4	1.7	1.3	1.0
2000.....	2.4	2.2	2.5	2.1	2.1	1.8	1.6	1.4	1.4	1.1
1999.....	2.2	2.4	2.1	2.6	1.6	1.7	1.8	1.6	1.2	0.9
White, both sexes										
2002.....	8.5	8.9	8.9	9.7	9.0	9.9	11.7	13.3	15.6	12.9
2001.....	7.9	8.5	9.0	9.4	8.9	10.0	11.1	13.4	15.0	13.2
2000.....	8.4	8.3	8.7	8.2	8.5	9.2	11.1	13.4	15.2	14.3
1999.....	7.5	8.3	8.0	8.9	8.2	9.5	11.8	13.9	14.4	13.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
White, male										
2002.....	14.3	15.0	15.5	16.9	16.3	18.8	24.4	29.6	38.8	40.5
2001.....	13.3	14.5	15.4	16.3	15.7	19.0	22.9	29.7	37.8	42.1
2000.....	14.0	14.2	14.7	14.2	15.3	17.3	22.7	30.4	38.1	46.5
1999.....	12.5	14.0	13.7	15.1	15.2	18.3	24.0	31.2	36.8	44.9
White, female										
2002.....	2.7	2.8	2.4	2.7	2.3	2.1	1.4	1.6	1.3	0.9
2001.....	2.4	2.6	2.8	2.7	2.7	2.0	1.5	1.8	1.3	1.1
2000.....	2.7	2.5	2.8	2.4	2.3	2.1	1.8	1.5	1.5	1.2
1999.....	2.5	2.7	2.5	3.0	1.8	1.8	2.1	1.8	1.3	1.0
Black, both sexes										
2002.....	3.0	2.9	2.9	2.7	2.9	2.3	4.6	5.3	*	*
2001.....	3.2	2.6	3.0	3.0	2.4	2.9	4.7	4.5	6.2	*
2000.....	3.3	3.3	3.3	2.9	3.0	3.1	4.5	*	*	*
1999.....	2.7	3.4	2.4	3.5	3.4	4.0	3.9	5.0	*	*
Black, male										
2002.....	5.2	5.7	5.3	5.5	5.5	5.0	10.0	13.3	*	*
2001.....	5.4	4.4	5.4	6.1	5.1	5.6	10.4	10.7	*	*
2000.....	6.1	6.0	5.6	5.8	5.6	7.0	11.0	*	*	*
1999.....	5.1	6.0	4.9	7.1	6.8	7.8	9.8	11.9	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by jumping from a high place (X80)										
All races 3/, both sexes										
2002.....	0.4	0.3	0.3	0.3	0.2	*	0.3	*	0.5	0.6
2001.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.4	*	*
2000.....	0.2	0.3	0.3	0.2	0.2	*	0.2	*	0.5	0.6
1999.....	0.4	0.4	0.3	0.2	*	0.3	0.3	0.3	0.6	0.5
All races 3/, male										
2002.....	0.5	0.4	0.4	0.5	0.4	*	*	*	*	*
2001.....	0.4	0.4	0.4	0.5	0.4	*	*	0.7	*	*
2000.....	0.3	0.5	0.4	*	*	*	*	*	*	*
1999.....	0.6	0.5	0.4	*	*	*	0.5	*	*	*
All races 3/, female										
2002.....	0.2	0.2	0.2	*	*	*	*	*	*	*
2001.....	0.2	0.2	0.3	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.4	0.3	0.3	0.3	0.2	*	0.3	*	0.5	0.6
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	*	*
2000.....	0.2	0.4	0.3	0.2	0.2	*	*	*	0.5	0.5
1999.....	0.4	0.3	0.3	0.2	*	0.3	0.3	0.3	0.5	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)										
All races 3/, both sexes										
2002.....	1.0	1.0	1.0	0.9	0.7	0.6	0.8	0.8	0.9	1.4
2001 4/.....	0.8	1.1	0.9	0.8	0.6	0.6	0.7	0.8	1.0	1.0
2000.....	0.9	0.9	0.8	0.9	0.7	0.6	0.7	0.8	1.0	1.2
1999.....	0.9	0.9	0.9	0.6	0.6	0.7	0.7	0.6	1.1	1.3
All races 3/, male										
2002.....	1.6	1.6	1.6	1.3	1.0	0.9	1.2	1.3	1.5	3.1
2001 4/.....	1.2	1.8	1.3	1.2	1.1	0.9	1.0	1.2	1.8	2.5
2000.....	1.4	1.4	1.1	1.4	0.9	1.0	1.0	1.4	1.8	2.4
1999.....	1.3	1.3	1.3	1.0	0.9	1.0	1.0	1.1	1.9	2.5
All races 3/, female										
2002.....	0.5	0.4	0.4	0.5	0.5	*	0.5	*	*	0.7
2001 4/.....	0.5	0.5	0.5	0.5	*	*	0.4	0.5	*	*
2000.....	0.4	0.5	0.5	0.4	0.4	*	0.4	*	*	0.7
1999.....	0.6	0.4	0.5	0.3	0.4	0.5	0.5	*	*	0.8
White, both sexes										
2002.....	1.1	1.1	1.0	0.9	0.8	0.6	0.8	0.8	0.9	1.5
2001 4/.....	0.9	1.2	0.9	0.9	0.7	0.5	0.7	0.8	1.0	1.1
2000.....	0.9	1.0	0.8	0.9	0.7	0.6	0.7	0.9	1.0	1.2
1999.....	1.0	0.9	0.9	0.6	0.6	0.7	0.7	0.7	1.1	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)										
All races 3/, both sexes										
2002.....	6.5	5.4	4.0	3.5	2.7	2.4	2.2	2.3	2.4	2.1
2001 4/.....	8.6	7.0	5.5	4.4	3.5	3.3	2.5	2.5	2.5	2.4
2000.....	6.6	5.3	4.0	3.4	2.6	2.4	2.4	2.3	2.6	2.4
1999.....	6.6	5.1	4.0	3.2	2.8	2.7	2.4	2.5	2.4	2.4
All races 3/, male										
2002.....	9.6	8.0	6.0	5.3	4.2	3.4	3.2	3.3	2.9	3.0
2001 4/.....	12.6	10.6	8.3	6.7	5.0	4.6	3.3	3.2	3.0	3.2
2000.....	9.7	7.8	5.9	5.2	3.9	3.3	3.4	2.8	3.8	3.3
1999.....	9.5	7.7	5.8	5.1	4.3	4.2	3.1	3.4	3.4	3.4
All races 3/, female										
2002.....	3.5	2.9	2.2	1.8	1.4	1.5	1.4	1.6	2.1	1.7
2001 4/.....	4.7	3.6	2.7	2.3	2.2	2.1	1.9	1.9	2.2	2.0
2000.....	3.6	2.8	2.1	1.7	1.5	1.7	1.6	2.0	1.9	2.0
1999.....	3.6	2.5	2.2	1.4	1.5	1.5	1.9	1.9	1.9	2.0
White, both sexes										
2002.....	4.5	3.8	2.9	2.9	2.2	1.9	1.9	1.8	2.1	1.8
2001 4/.....	6.8	5.6	4.4	3.8	3.0	2.7	2.2	2.1	2.2	2.2
2000.....	4.4	3.7	2.9	2.7	2.0	1.8	2.0	2.1	2.0	2.2
1999.....	4.5	3.6	3.0	2.6	2.3	2.3	2.0	2.0	2.1	2.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)										
White, male										
2002.....	6.5	5.1	4.0	4.1	3.2	2.7	2.6	2.4	2.3	2.7
2001 4/.....	9.8	8.2	6.5	5.5	4.2	3.7	2.6	2.7	2.6	2.8
2000.....	6.4	5.1	4.0	4.0	2.8	2.2	2.6	2.4	2.5	2.8
1999.....	6.3	5.3	4.3	3.8	3.3	3.4	2.5	2.6	2.9	2.7
White, female										
2002.....	2.5	2.5	1.8	1.7	1.3	1.3	1.4	1.4	1.9	1.5
2001 4/.....	3.8	3.0	2.4	2.1	1.9	1.9	1.8	1.7	1.9	2.0
2000.....	2.5	2.3	1.8	1.4	1.3	1.4	1.5	1.8	1.7	1.9
1999.....	2.6	1.9	1.7	1.3	1.3	1.4	1.7	1.6	1.6	1.9
Black, both sexes										
2002.....	20.6	17.6	13.0	8.9	7.5	5.8	5.6	8.2	6.0	*
2001 4/.....	21.7	18.1	13.6	9.9	7.5	7.7	6.5	6.9	6.2	*
2000.....	21.8	16.8	12.4	9.1	7.4	8.2	6.8	5.3	9.6	*
1999.....	21.2	16.2	11.2	8.8	7.4	6.6	6.0	8.2	6.1	*
Black, male										
2002.....	32.4	30.8	22.5	16.8	13.5	10.0	11.3	15.2	*	*
2001 4/.....	34.2	30.0	23.1	17.4	12.0	13.2	11.8	11.6	*	*
2000.....	34.1	28.7	22.4	16.0	13.0	15.1	12.4	*	19.4	*
1999.....	34.1	27.3	18.5	17.0	12.7	13.0	10.2	13.8	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)										
Black, female										
2002.....	10.1	6.1	4.9	*	*	*	*	*	*	*
2001 4/.....	10.6	7.7	5.6	3.8	4.0	*	*	*	*	*
2000.....	10.9	6.5	3.8	3.5	*	*	*	*	*	*
1999.....	9.8	6.7	5.0	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by hanging, strangulation and suffocation (X91)

All races 3/, both sexes

2002.....	0.3	0.3	0.2	*	0.2	*	*	*	*	*
2001.....	0.3	0.3	0.2	0.1	0.2	0.2	0.3	*	*	*
2000.....	0.3	0.2	0.1	0.2	*	*	*	*	*	*
1999.....	0.3	0.2	0.2	*	*	0.3	*	*	*	*

All races 3/, male

2002.....	*	0.3	*	*	*	*	*	*	*	*
2001.....	0.2	0.2	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2002.....	0.4	0.3	0.3	*	*	*	*	*	*	*
2001.....	0.5	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	0.3	*	*	*	*	*	*	*	*
1999.....	0.4	0.2	*	*	*	*	*	*	*	*

White, both sexes

2002.....	0.2	0.2	0.2	*	*	*	*	*	*	*
2001.....	0.2	0.2	0.2	*	*	*	0.3	*	*	*
2000.....	0.2	0.2	0.1	*	*	*	*	*	*	*
1999.....	0.2	0.1	0.1	*	*	0.3	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
All races 3/, both sexes										
2002.....	3.9	2.8	2.2	1.8	1.3	1.0	0.8	1.0	0.8	*
2001.....	3.7	2.7	2.1	1.5	1.4	1.2	0.9	0.8	0.7	*
2000.....	3.8	2.9	2.1	1.7	1.2	1.0	1.0	0.7	0.5	0.7
1999.....	3.7	2.8	2.0	1.6	1.6	1.1	1.0	0.8	0.7	0.6
All races 3/, male										
2002.....	6.2	4.2	3.4	2.7	1.9	1.4	1.0	1.4	1.0	*
2001.....	5.6	4.2	3.2	2.3	2.0	1.8	1.1	1.2	*	*
2000.....	5.8	4.5	3.3	2.6	1.8	1.3	1.3	0.8	*	*
1999.....	5.5	4.4	3.0	2.6	2.5	1.5	1.4	1.1	*	*
All races 3/, female										
2002.....	1.7	1.5	1.1	0.9	0.7	0.6	0.6	0.6	0.7	*
2001.....	1.7	1.1	1.0	0.7	0.8	0.7	0.7	0.5	0.7	*
2000.....	1.7	1.3	1.0	0.9	0.7	0.8	0.7	0.7	*	*
1999.....	1.8	1.3	1.1	0.6	0.7	0.7	0.7	0.6	0.6	*
White, both sexes										
2002.....	2.6	1.9	1.6	1.5	0.9	0.8	0.7	0.8	0.8	*
2001.....	2.6	1.9	1.5	1.2	1.2	1.0	0.7	0.7	0.7	*
2000.....	2.5	2.0	1.5	1.4	1.0	0.8	0.8	0.7	*	0.6
1999.....	2.5	1.9	1.5	1.2	1.2	1.0	0.9	0.7	0.7	0.6

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
White, male										
2002.....	4.0	2.6	2.4	2.1	1.4	1.0	0.8	1.2	*	*
2001.....	3.7	2.8	2.3	1.6	1.7	1.3	0.8	1.0	*	*
2000.....	3.7	3.0	2.2	2.1	1.4	1.0	0.9	0.7	*	*
1999.....	3.6	2.8	2.3	1.9	1.8	1.3	1.1	0.7	*	*
White, female										
2002.....	1.3	1.3	0.9	0.9	0.5	0.5	0.6	0.6	0.7	*
2001.....	1.4	0.9	0.8	0.8	0.7	0.8	0.7	0.5	0.7	*
2000.....	1.3	1.1	0.9	0.8	0.7	0.7	0.7	0.7	*	*
1999.....	1.4	1.0	0.7	0.6	0.7	0.7	0.7	0.6	*	*
Black, both sexes										
2002.....	13.1	9.6	7.1	4.2	4.0	2.8	*	*	*	*
2001.....	11.6	8.4	6.5	4.3	3.0	2.9	*	*	*	*
2000.....	12.6	9.4	7.0	4.2	2.6	2.8	2.7	*	*	*
1999.....	12.2	9.7	5.9	4.5	4.2	2.2	*	*	*	*
Black, male										
2002.....	22.7	17.4	12.3	8.3	6.5	5.2	*	*	*	*
2001.....	20.3	15.3	11.8	8.6	5.1	6.4	*	*	*	*
2000.....	21.6	16.7	13.1	7.1	5.4	5.2	*	*	*	*
1999.....	20.5	17.3	9.5	9.3	8.3	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, female

2002.....	4.6	2.8	2.8	*	*	*	*	*	*	*
2001.....	3.9	2.4	1.9	*	*	*	*	*	*	*
2000.....	4.7	3.2	*	*	*	*	*	*	*	*
1999.....	4.8	3.2	2.8	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by sharp object (X99)										
All races 3/, both sexes										
2002.....	1.1	0.9	0.6	0.6	0.4	0.4	0.5	0.3	0.4	*
2001.....	0.9	0.8	0.6	0.5	0.3	0.6	0.3	0.5	*	*
2000.....	1.0	0.7	0.5	0.5	0.4	0.4	0.3	0.4	0.5	*
1999.....	1.0	0.8	0.6	0.5	0.4	0.3	0.3	0.4	*	*
All races 3/, male										
2002.....	1.5	1.3	1.0	1.0	0.5	0.5	0.8	*	*	*
2001.....	1.3	1.2	0.9	0.8	*	0.7	*	*	*	*
2000.....	1.4	1.0	0.6	0.8	0.5	0.5	0.6	0.7	*	*
1999.....	1.6	1.1	0.9	0.7	0.6	0.5	*	*	*	*
All races 3/, female										
2002.....	0.7	0.4	0.3	0.3	*	*	*	*	*	*
2001.....	0.6	0.4	0.3	0.3	0.3	0.5	*	0.5	*	*
2000.....	0.6	0.4	0.4	*	*	*	*	*	*	*
1999.....	0.4	0.4	0.3	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.7	0.5	0.4	0.4	0.3	0.3	0.4	*	*	*
2001.....	0.6	0.5	0.4	0.4	0.3	0.5	*	0.4	*	*
2000.....	0.6	0.4	0.4	0.3	0.3	*	*	0.3	*	*
1999.....	0.6	0.5	0.4	0.4	0.3	0.3	0.3	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by sharp object (X99)										
White, male										
2002.....	1.0	0.8	0.5	0.6	*	*	*	*	*	*
2001.....	0.8	0.7	0.6	0.6	*	0.5	*	*	*	*
2000.....	0.8	0.5	0.4	0.6	*	*	*	*	*	*
1999.....	1.0	0.7	0.6	0.4	*	*	*	*	*	*
White, female										
2002.....	0.5	0.3	*	*	*	*	*	*	*	*
2001.....	0.4	0.3	0.3	*	*	*	*	*	*	*
2000.....	0.3	0.3	0.3	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002.....	3.9	3.1	2.6	2.4	*	*	*	*	*	*
2001.....	3.5	3.0	1.7	1.7	*	*	*	*	*	*
2000.....	3.8	2.8	1.6	1.8	*	*	*	*	*	*
1999.....	3.4	2.8	1.9	1.6	*	*	*	*	*	*
Black, male										
2002.....	5.4	5.3	4.7	4.3	*	*	*	*	*	*
2001.....	5.3	5.4	3.2	*	*	*	*	*	*	*
2000.....	5.1	4.7	2.4	*	*	*	*	*	*	*
1999.....	6.0	4.1	3.1	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

All races 3/, both sexes

2002.....	1.1	1.2	0.8	0.8	0.8	0.8	0.7	0.8	0.7	0.9
2001 4/.....	3.4	3.1	2.5	2.1	1.4	1.1	0.9	0.8	1.0	1.1
2000.....	1.3	1.2	1.0	0.9	0.7	0.7	0.8	0.9	1.3	0.9
1999.....	1.3	1.1	1.0	0.8	0.7	0.8	0.7	0.9	1.0	1.1

All races 3/, male

2002.....	1.6	1.8	1.2	1.3	1.4	1.2	1.1	1.2	1.1	*
2001 4/.....	5.1	4.5	3.8	3.2	2.2	1.7	1.4	1.1	1.4	1.6
2000.....	1.9	1.8	1.6	1.3	1.3	1.2	1.3	1.1	2.0	*
1999.....	1.9	1.7	1.5	1.3	1.0	1.4	0.9	1.4	1.8	*

All races 3/, female

2002.....	0.6	0.6	0.4	0.4	*	0.4	*	0.5	*	0.8
2001 4/.....	1.7	1.6	1.2	1.1	0.8	0.6	0.4	0.6	0.8	0.9
2000.....	0.7	0.7	0.4	0.5	*	*	*	0.7	0.8	0.7
1999.....	0.8	0.6	0.5	0.4	*	*	0.5	0.6	*	0.9

White, both sexes

2002.....	0.9	0.9	0.6	0.7	0.7	0.6	0.6	0.6	0.6	0.7
2001 4/.....	3.2	2.8	2.2	2.0	1.3	0.9	0.7	0.7	0.9	1.0
2000.....	0.9	0.9	0.7	0.7	0.5	0.5	0.6	0.7	1.0	0.8
1999.....	0.9	0.9	0.7	0.7	0.6	0.7	0.6	0.8	0.9	0.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

White, male

2002.....	1.3	1.3	0.8	1.1	1.2	0.9	1.0	0.8	*	*
2001 4/.....	4.9	4.2	3.3	3.1	1.8	1.5	1.1	0.9	1.2	*
2000.....	1.4	1.1	1.1	1.0	0.9	0.8	1.0	0.9	1.3	*
1999.....	1.4	1.3	1.0	1.1	0.9	1.1	0.7	1.3	1.5	*

White, female

2002.....	0.4	0.5	0.4	0.4	*	*	*	*	*	0.7
2001 4/.....	1.5	1.4	1.0	1.0	0.7	*	*	0.5	0.7	0.9
2000.....	0.5	0.6	0.4	0.4	*	*	*	0.6	0.7	*
1999.....	0.5	0.4	0.5	*	*	*	0.5	*	*	0.9

Black, both sexes

2002.....	2.5	3.5	2.5	1.7	2.3	*	*	*	*	*
2001 4/.....	4.9	5.4	4.6	3.1	2.9	2.6	*	*	*	*
2000.....	3.6	3.7	3.1	2.2	2.7	2.2	*	*	*	*
1999.....	4.2	3.0	2.7	1.8	*	*	*	*	*	*

Black, male

2002.....	3.7	6.1	4.5	3.6	4.6	*	*	*	*	*
2001 4/.....	6.8	7.8	7.0	4.7	5.3	*	*	*	*	*
2000.....	5.5	6.5	6.2	4.3	5.4	*	*	*	*	*
1999.....	6.2	4.7	5.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2002.....	3.4	3.3	2.3	1.4	1.0	0.8	0.7	0.6	1.0	1.4
2001.....	3.2	3.0	2.0	1.1	0.7	0.6	0.5	0.7	1.1	1.0
2000.....	3.1	2.7	1.7	1.1	0.7	0.6	0.6	0.6	0.7	1.3
1999.....	2.9	2.7	1.6	1.0	0.8	0.8	0.5	0.7	0.9	1.4
All races 3/, male										
2002.....	4.3	4.1	2.9	1.8	1.3	1.0	1.0	0.8	1.1	2.2
2001.....	4.4	4.2	2.5	1.5	0.8	0.9	0.6	0.9	1.5	1.5
2000.....	4.2	3.8	2.3	1.6	0.9	0.9	0.8	1.0	1.1	1.7
1999.....	4.0	3.7	2.1	1.5	1.1	1.1	0.6	1.0	1.3	2.1
All races 3/, female										
2002.....	2.6	2.6	1.7	1.0	0.7	0.7	0.5	0.5	0.9	1.1
2001.....	2.0	1.8	1.5	0.6	0.5	*	*	0.6	0.8	0.8
2000.....	1.9	1.7	1.1	0.7	0.5	*	*	*	*	1.2
1999.....	1.9	1.7	1.0	0.6	0.6	0.5	*	0.5	0.7	1.1
White, both sexes										
2002.....	3.5	3.3	2.2	1.4	0.9	0.8	0.7	0.6	0.9	1.4
2001.....	3.2	2.9	1.9	1.1	0.6	0.6	0.5	0.7	1.0	1.0
2000.....	2.9	2.6	1.6	1.0	0.7	0.5	0.5	0.6	0.8	1.3
1999.....	2.8	2.5	1.4	1.0	0.8	0.8	0.4	0.7	0.9	1.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
White, male										
2002.....	4.3	4.0	2.5	1.8	1.2	0.9	0.8	*	1.1	2.3
2001.....	4.2	4.0	2.3	1.4	0.7	0.8	0.6	0.9	1.5	*
2000.....	4.0	3.5	2.1	1.3	0.9	0.8	0.7	1.0	*	*
1999.....	3.8	3.2	1.8	1.3	1.0	1.1	*	1.0	1.3	2.1
White, female										
2002.....	2.6	2.6	1.9	1.1	0.7	0.7	0.5	*	0.7	1.1
2001.....	2.1	1.9	1.6	0.7	0.6	*	*	0.6	0.7	0.8
2000.....	1.9	1.7	1.1	0.7	0.5	*	*	*	*	1.1
1999.....	1.7	1.8	1.0	0.7	0.7	0.5	*	*	0.8	1.1
Black, both sexes										
2002.....	4.3	4.5	3.8	1.5	*	*	*	*	*	*
2001.....	4.0	4.1	3.3	1.5	*	*	*	*	*	*
2000.....	4.8	4.4	2.8	2.3	*	*	*	*	*	*
1999.....	4.6	4.8	3.3	1.5	*	*	*	*	*	*
Black, male										
2002.....	5.5	6.2	6.5	*	*	*	*	*	*	*
2001.....	6.6	7.0	5.4	*	*	*	*	*	*	*
2000.....	7.2	7.0	4.4	4.0	*	*	*	*	*	*
1999.....	6.5	8.3	5.7	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

All races 3/, both sexes

2002.....	2.7	2.6	1.7	0.8	0.5	0.3	*	*	*	*
2001.....	2.5	2.2	1.4	0.6	0.3	*	*	*	*	*
2000.....	2.4	2.1	1.1	0.5	0.3	*	*	*	*	*
1999.....	2.3	2.0	1.0	0.6	0.3	0.3	*	*	*	*

All races 3/, male

2002.....	3.2	3.0	1.9	0.9	0.6	*	*	*	*	*
2001.....	3.3	3.2	1.7	0.7	*	*	*	*	*	*
2000.....	3.3	2.8	1.5	0.7	*	*	*	*	*	*
1999.....	3.1	2.6	1.3	0.7	*	*	*	*	*	*

All races 3/, female

2002.....	2.2	2.2	1.5	0.7	0.4	*	*	*	*	*
2001.....	1.7	1.3	1.1	0.5	0.3	*	*	*	*	*
2000.....	1.6	1.4	0.8	0.4	*	*	*	*	*	*
1999.....	1.6	1.4	0.7	0.4	*	*	*	*	*	*

White, both sexes

2002.....	2.7	2.6	1.6	0.8	0.5	0.3	*	*	*	*
2001.....	2.5	2.2	1.4	0.6	0.3	*	*	*	*	*
2000.....	2.3	2.0	1.1	0.5	0.3	*	*	*	*	*
1999.....	2.2	1.9	0.9	0.5	0.3	0.3	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

White, male

2002.....	3.3	3.0	1.7	0.8	0.6	*	*	*	*	*
2001.....	3.2	3.0	1.6	0.7	*	*	*	*	*	*
2000.....	3.1	2.7	1.4	0.6	*	*	*	*	*	*
1999.....	2.9	2.2	1.1	0.6	*	*	*	*	*	*

White, female

2002.....	2.2	2.2	1.6	0.7	0.5	*	*	*	*	*
2001.....	1.9	1.5	1.2	0.6	0.4	*	*	*	*	*
2000.....	1.5	1.3	0.8	0.4	*	*	*	*	*	*
1999.....	1.5	1.5	0.7	0.5	*	*	*	*	*	*

Black, both sexes

2002.....	3.4	3.4	2.9	*	*	*	*	*	*	*
2001.....	3.3	3.0	2.4	*	*	*	*	*	*	*
2000.....	3.8	3.3	1.9	*	*	*	*	*	*	*
1999.....	3.6	3.6	2.3	*	*	*	*	*	*	*

Black, male

2002.....	4.0	4.2	4.8	*	*	*	*	*	*	*
2001.....	5.6	5.2	3.7	*	*	*	*	*	*	*
2000.....	5.6	4.9	2.8	*	*	*	*	*	*	*
1999.....	4.8	6.0	3.7	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2002.....	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.6	1.1
2001.....	0.5	0.5	0.4	0.3	0.2	0.3	0.2	0.4	0.7	0.7
2000.....	0.4	0.5	0.4	0.5	0.3	0.3	0.4	0.3	0.5	0.9
1999.....	0.4	0.5	0.4	0.4	0.3	0.4	0.2	0.4	0.5	1.0
All races 3/, male										
2002.....	0.7	0.8	0.7	0.7	0.6	0.7	0.5	*	*	1.7
2001.....	0.7	0.7	0.6	0.6	*	0.5	*	*	1.1	*
2000.....	0.6	0.8	0.6	0.6	0.5	0.5	0.5	*	*	*
1999.....	0.6	0.8	0.6	0.7	0.5	0.5	*	*	*	*
All races 3/, female										
2002.....	0.4	0.3	*	*	*	*	*	*	*	0.8
2001.....	0.2	0.3	0.3	*	*	*	*	*	*	*
2000.....	0.3	0.2	0.3	*	*	*	*	*	*	0.9
1999.....	0.2	*	*	*	*	*	*	*	*	0.8
White, both sexes										
2002.....	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.4	0.6	1.1
2001.....	0.4	0.5	0.4	0.3	*	0.3	*	0.4	0.7	0.7
2000.....	0.4	0.4	0.4	0.4	0.2	0.3	0.3	0.3	0.6	0.8
1999.....	0.3	0.4	0.3	0.4	0.3	0.3	*	0.4	0.5	0.9

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Legal intervention (Y35,Y89.0)										
All races 3/, both sexes										
2002.....	0.2	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	0.2	*	*	*	*	*	*	*
All races 3/, male										
2002.....	0.4	0.4	*	*	*	*	*	*	*	*
2001.....	0.4	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	0.4	*	*	*	*	*	*	*	*
1999.....	0.4	*	0.4	*	*	*	*	*	*	*
All races 3/, female										
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002.....	0.2	0.2	*	*	*	*	*	*	*	*
2001.....	0.2	0.1	*	*	*	*	*	*	*	*
2000.....	0.1	0.2	*	*	*	*	*	*	*	*
1999.....	0.2	*	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)										
All races 3/, both sexes										
2002.....	0.4	0.5	0.8	1.1	1.6	2.3	3.9	5.1	7.7	11.3
2001.....	0.4	0.7	0.8	1.3	1.9	2.6	3.6	5.9	8.3	12.8
2000.....	0.4	0.6	0.9	1.2	2.0	2.8	4.3	5.8	8.0	13.6
1999.....	0.4	0.5	0.7	1.1	1.7	2.4	4.0	5.5	8.4	12.7
All races 3/, male										
2002.....	0.4	0.6	0.7	1.2	1.5	2.5	3.8	5.4	7.7	12.3
2001.....	0.4	0.7	0.8	1.3	2.0	2.6	3.8	6.3	9.9	14.5
2000.....	0.4	0.6	1.0	1.2	2.0	2.8	5.3	7.0	10.4	16.0
1999.....	0.4	0.4	0.7	1.2	1.7	2.3	4.5	6.4	10.1	16.5
All races 3/, female										
2002.....	0.4	0.5	0.9	1.0	1.7	2.2	3.9	5.0	7.7	10.9
2001.....	0.5	0.6	0.8	1.2	1.7	2.6	3.5	5.7	7.3	12.1
2000.....	0.4	0.7	0.9	1.1	2.0	2.7	3.6	4.9	6.7	12.7
1999.....	0.4	0.5	0.7	0.9	1.6	2.5	3.6	4.8	7.4	11.2
White, both sexes										
2002.....	0.4	0.5	0.7	1.0	1.5	2.2	3.8	5.0	7.6	11.2
2001.....	0.4	0.6	0.7	1.1	1.6	2.4	3.2	5.8	8.2	12.1
2000.....	0.3	0.6	0.9	1.0	1.8	2.6	4.1	5.4	8.0	13.1
1999.....	0.3	0.4	0.6	0.9	1.5	2.3	3.8	5.2	8.0	12.1

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)										
White, male										
2002.....	0.4	0.5	0.6	1.0	1.5	2.2	3.7	5.3	7.8	11.8
2001.....	0.3	0.6	0.6	1.1	1.7	2.5	3.2	6.2	10.0	13.6
2000.....	0.4	0.5	1.0	1.1	1.8	2.5	5.0	6.8	10.6	15.8
1999.....	0.3	0.3	0.6	1.0	1.6	2.3	4.2	6.2	9.7	15.6
White, female										
2002.....	0.3	0.4	0.7	0.9	1.5	2.1	3.9	4.8	7.5	10.9
2001.....	0.4	0.5	0.7	1.1	1.6	2.4	3.2	5.6	7.0	11.5
2000.....	0.3	0.6	0.9	0.9	1.8	2.7	3.3	4.5	6.4	12.0
1999.....	0.4	0.5	0.6	0.7	1.5	2.4	3.4	4.5	7.1	10.7
Black, both sexes										
2002.....	1.0	1.2	2.1	2.6	2.8	4.0	5.2	7.6	9.4	14.0
2001.....	1.0	1.6	1.5	2.7	4.2	4.6	8.5	8.0	11.1	23.3
2000.....	*	1.1	1.7	2.6	4.7	4.6	7.9	10.5	10.4	21.0
1999.....	0.8	0.9	1.8	2.6	2.6	3.5	7.3	9.5	12.4	20.1
Black, male										
2002.....	*	*	*	*	*	5.0	*	*	*	*
2001.....	*	1.7	*	3.4	5.1	5.3	11.1	*	*	29.5
2000.....	*	*	*	*	4.6	6.2	9.4	10.4	*	*
1999.....	*	*	*	*	*	*	9.2	10.4	*	23.4

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Complications of medical and surgical care (Y40-Y84,Y88)

Black, female

2002.....	*	*	2.5	2.4	3.1	*	4.9	8.2	9.9	11.5
2001.....	*	*	*	*	3.5	4.1	6.7	8.0	11.0	21.0
2000.....	*	*	*	3.1	4.7	*	6.9	10.5	10.6	21.9
1999.....	*	*	*	*	*	*	6.1	8.9	10.8	18.8

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)

All races 3/, both sexes

2002.....	*	*	*	*	*	*	*	0.3	0.4	0.5
2001.....	*	*	*	*	*	0.2	*	0.4	0.4	0.5
2000.....	*	*	*	*	*	0.2	0.3	0.3	*	0.6
1999.....	*	*	*	*	*	*	0.3	0.3	0.5	*

All races 3/, male

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	0.5	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, both sexes

2002.....	*	*	*	*	*	*	*	*	0.4	*
2001.....	*	*	*	*	*	*	*	0.3	0.5	0.5
2000.....	*	*	*	*	*	*	*	0.3	*	0.6
1999.....	*	*	*	*	*	*	*	0.3	0.5	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)										
All races 3/, both sexes										
2002.....	0.3	0.4	0.7	0.8	1.4	2.0	3.3	4.6	6.8	10.4
2001.....	0.3	0.5	0.6	1.1	1.6	2.2	3.2	5.2	7.4	11.4
2000.....	0.3	0.5	0.8	0.9	1.8	2.4	3.8	5.0	7.0	12.2
1999.....	0.3	0.3	0.6	0.8	1.4	2.0	3.5	4.9	7.5	11.8
All races 3/, male										
2002.....	0.3	0.4	0.6	0.9	1.3	2.1	3.3	4.9	7.0	11.7
2001.....	0.2	0.6	0.6	1.2	1.6	2.3	3.4	5.5	8.8	13.2
2000.....	0.3	0.5	0.8	1.0	1.8	2.5	4.7	6.0	8.9	14.3
1999.....	0.3	0.3	0.6	1.0	1.5	2.0	4.0	5.8	9.1	15.4
All races 3/, female										
2002.....	0.4	0.4	0.7	0.8	1.5	1.9	3.3	4.4	6.7	9.9
2001.....	0.4	0.5	0.6	1.1	1.6	2.1	3.0	4.9	6.5	10.6
2000.....	0.3	0.5	0.7	0.9	1.8	2.3	3.1	4.4	5.9	11.4
1999.....	0.3	0.4	0.5	0.7	1.3	2.1	3.1	4.3	6.6	10.4
White, both sexes										
2002.....	0.3	0.4	0.6	0.7	1.3	1.8	3.3	4.5	6.7	10.2
2001.....	0.3	0.5	0.5	1.0	1.4	2.0	2.8	5.1	7.2	10.7
2000.....	0.3	0.5	0.8	0.8	1.7	2.3	3.6	4.7	7.0	11.8
1999.....	0.3	0.3	0.5	0.7	1.3	2.0	3.4	4.7	7.2	11.2

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

White, male

2002.....	0.3	0.4	0.5	0.8	1.3	1.8	3.2	4.9	7.1	11.2
2001.....	*	0.6	0.5	1.0	1.4	2.2	2.9	5.5	8.9	12.3
2000.....	0.3	0.4	0.8	0.9	1.6	2.3	4.5	5.8	9.1	14.0
1999.....	0.2	*	0.5	0.7	1.5	1.9	3.8	5.6	8.7	14.5

White, female

2002.....	0.2	0.4	0.6	0.7	1.4	1.8	3.4	4.3	6.4	9.8
2001.....	0.4	0.4	0.6	1.0	1.5	1.9	2.8	4.9	6.2	10.0
2000.....	*	0.5	0.7	0.7	1.7	2.2	2.9	3.9	5.7	10.8
1999.....	0.3	0.4	0.5	0.6	1.2	2.0	3.0	4.0	6.2	9.9

Black, both sexes

2002.....	0.9	0.9	1.7	1.8	2.3	3.8	4.2	6.7	9.2	13.4
2001.....	0.8	1.0	1.3	2.5	3.6	3.8	7.3	6.4	10.3	22.1
2000.....	*	0.8	1.2	2.4	3.6	4.0	6.7	9.8	9.0	19.1
1999.....	*	*	1.5	2.2	2.5	2.9	6.0	8.4	11.0	19.1

Black, male

2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	3.3	4.7	*	9.4	*	*	28.4
2000.....	*	*	*	*	*	5.2	7.7	9.9	*	*
1999.....	*	*	*	*	*	*	8.1	9.5	*	*

Worktable 292R. Death rates for 358 selected causes by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)										
Black, female										
2002.....	*	*	1.8	*	*	*	*	7.1	9.5	11.1
2001.....	*	*	*	*	*	*	5.8	6.4	10.2	19.8
2000.....	*	*	*	2.8	3.4	*	6.0	9.7	8.9	19.3
1999.....	*	*	*	*	*	*	4.5	7.7	9.5	18.0

* Figure does not meet standards of reliability and precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Includes races other than white and black.

4/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.