

2 Bi-layer Tablets

NEW
Mucinex^{600 mg}
Guaifenesin Extended-Release Tablets
EXPECTORANT

Drug Facts

Active ingredient (in each extended-release tablet) Purpose
Guaifenesin 600 mg Expectorant

Uses helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Do not use ■ for children under 12 years of age

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Stop use and ask a doctor if ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for the timing of meals
- adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if neckband is broken
- store between 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

Inactive ingredients carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

NDC 63824-008-82

NEW
Mucinex^{600 mg}
Guaifenesin Extended-Release Tablets
EXPECTORANT



2 Bi-layer Tablets



Mucinex^{600 mg}
Guaifenesin Extended-Release Tablets
EXPECTORANT




Please visit our Web site:
www.adamslabs.com

PHYSICIAN SAMPLE-
NOT TO BE SOLD



Fort Worth, Texas 76155 022802
© 2002 Adams Laboratories, Inc. 2000802

3-63824-008-82-0



NDC 63824-008-82

Mucinex^{600mg}
Guaifenesin Extended-Release Tablets

EXPECTORANT

2 Bi-layer Tablets

SEE CARTON FOR COMPLETE INFORMATION

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for the timing of meals
- adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years of age: do not use

**PHYSICIAN
SAMPLE-
NOT TO BE SOLD**

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155

3100802 022802

3-6382400882-0

NDC 63824-008-10

Mucinex^{600mg}
Guaifenesin Extended-Release Tablets
EXPECTORANT

3-6382400810-3

100 Bi-layer Tablets

Adams[®]
LABORATORIES, INC. Fort Worth, Texas 76155
3000810 022802

Drug Facts

Active ingredient (in each extended-release tablet) Purpose
Guaifenesin 600 mg Expectorant

Uses helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Do not use ■ for children under 12 years of age

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use. ▶

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for the timing of meals
- adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if neckband is broken
- store between 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

Inactive ingredients carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

20 BI-LAYER TABLETS

Mucinex
600mg
NEW
Guaifenesin Extended-Release Tablets
EXPECTORANT

Drug Facts

Active ingredient Purpose
(in each extended-release tablet)
Guaifenesin 600 mg Expectorant

Uses helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Do not use ■ for children under 12 years of age

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Stop use and ask a doctor if ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

NEW

Mucinex
600mg
Guaifenesin Extended-Release Tablets
EXPECTORANT

Drug Facts (continued)

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for the timing of meals
- adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if neckband is broken
- store between 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

Inactive ingredients carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

NDC 63824-008-20

NEW

Mucinex
600mg
Guaifenesin Extended-Release Tablets
EXPECTORANT



20 BI-LAYER TABLETS

NEW

Mucinex
600mg
Guaifenesin Extended-Release Tablets
EXPECTORANT

Please visit our Web site:
www.adamslabs.com

Adams
LABORATORIES, INC.


Fort Worth, Texas 76155

© 2002 Adams Laboratories, Inc.

030802

2000820

3-63824-008-20-2

<p>NDC 63824-008-20</p> <p>Mucinex^{600mg}</p> <p>Guafenesin Extended-Release Tablets</p> <p>EXPECTORANT</p>  <p>20 Bi-layer Tablets</p> <p>Adams[®]</p> <p><small>LABORATORIES, INC.</small> Fort Worth, Texas 76155 3000820 030802</p>	<p>Drug Facts</p> <p>Active ingredient (in each extended-release tablet) Purpose Guafenesin 600 mg Expectorant</p> <p>Uses helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive</p> <p>Warnings Do not use ■ for children under 12 years of age</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none">■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema■ cough accompanied by too much phlegm (mucus) <p>Stop use and ask a doctor if ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.</p> <p>If pregnant or breast-feeding, ask a health professional before use. ▶</p>	<p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none">■ do not crush, chew, or break tablet■ take with a full glass of water■ this product can be administered without regard for the timing of meals■ adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.■ children under 12 years of age: do not use <p>Other information</p> <ul style="list-style-type: none">■ tamper evident: do not use if neckband is broken■ store between 20-25°C (68-77°F)■ see bottom of bottle for lot code and expiration date <p>Inactive ingredients carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF</p>

40 BI-LAYER TABLETS

Mucinex
600mg
NEW
Guaifenesin Extended-Release Tablets
EXPECTORANT

Drug Facts

Active ingredient Purpose
(in each extended-release tablet)
Guaifenesin 600 mg Expectorant

Uses helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings
Do not use ■ for children under 12 years of age
Ask a doctor before use if you have
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough accompanied by too much phlegm (mucus)

Stop use and ask a doctor if ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

NEW
Mucinex^{600mg}
Guaifenesin Extended-Release Tablets
EXPECTORANT

Drug Facts (continued)
Directions
■ do not crush, chew, or break tablet
■ take with a full glass of water
■ this product can be administered without regard for the timing of meals
■ adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
■ children under 12 years of age: do not use

Other information
■ tamper evident: do not use if neckband is broken
■ store between 20-25°C (68-77°F)
■ see bottom of bottle for lot code and expiration date

Inactive ingredients carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

NEW NDC 63824-008-40
Mucinex^{600mg}
Guaifenesin Extended-Release Tablets
EXPECTORANT




40 BI-LAYER TABLETS

NEW
Mucinex^{600mg}
Guaifenesin Extended-Release Tablets
EXPECTORANT

Please visit our Web site:
www.adamslabs.com

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155
© 2002 Adams Laboratories, Inc.
022802
2000840

3-63824-008-40-0

<p>NDC 63824-008-40</p> <p>Mucinex[®] 600mg</p> <p>Guafenesin Extended-Release Tablets</p> <p>EXPECTORANT</p>  <p>40 Bi-layer Tablets</p> <p>Adams[®] LABORATORIES, INC. Fort Worth, Texas 76155 3000840 022802</p>	<p>Drug Facts</p> <p>Active ingredient (in each extended-release tablet) Purpose Guafenesin 600 mg Expectorant</p> <p>Uses helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive</p> <p>Warnings Do not use ■ for children under 12 years of age</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none">■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema■ cough accompanied by too much phlegm (mucus) <p>Stop use and ask a doctor if ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.</p> <p>If pregnant or breast-feeding, ask a health professional before use. ▶</p>	<p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none">■ do not crush, chew, or break tablet■ take with a full glass of water■ this product can be administered without regard for the timing of meals■ adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.■ children under 12 years of age: do not use <p>Other information</p> <ul style="list-style-type: none">■ tamper evident: do not use if neckband is broken■ store between 20-25°C (68-77°F)■ see bottom of bottle for lot code and expiration date <p>Inactive ingredients carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF</p>

NDC 63824-008-50

Mucinex^{600mg}

Guaifenesin Extended-Release Tablets

EXPECTORANT

3-6382400850-9



500 Bi-layer Tablets

Adams
LABORATORIES, INC. Fort Worth, Texas 76155

Adams
LABORATORIES, INC.

3000650 022802

Drug Facts

Active ingredient (in each extended-release tablet) Purpose
Guaifenesin 600 mg Expectorant

Uses helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Do not use ■ for children under 12 years of age

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Stop use and ask a doctor if ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for the timing of meals
- adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if neckband is broken
- store between 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

Inactive ingredients carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

**This is a representation of an electronic record that was signed electronically and
this page is the manifestation of the electronic signature.**

/s/

Badrul Chowdhury
7/12/02 03:44:27 PM

Linda Katz
7/12/02 03:49:12 PM
Linda M. Katz, M.D. signing for Charles J. Ganley, M.D.