


**PROJECT DESCRIPTION**  
This lidar-derived topographic map was produced as a collaborative effort between the U.S. Geological Survey (USGS) Coastal and Marine Geology Program, the Northeast Coastal and Barrier Network of the National Park Service (NPS) Inventory and Monitoring Program, the South Florida/Caribbean Network of the NPS Inventory and Monitoring Program, and the National Aeronautics and Space Administration (NASA) Wallops Flight Facility. The aims of the partnership that created this product are to develop advanced survey techniques for mapping barrier island geomorphology and habitats, and to enable the monitoring of ecological and geological change within National Seashores. This product is based on data from an innovative airborne lidar instrument under development at the NASA Wallops Flight Facility, the NASA Experimental Airborne Advanced Research Lidar (EAARL).

**DATA DESCRIPTION**  
The laser soundings used to create this map were collected during November 2002 by the NASA EAARL system mounted on a Cessna 310 aircraft. The EAARL uses a "waveform-resolving" green laser capable of mapping submarine and subaerial (land) topography in a single overflight. The EAARL system is typically flown at 300 m altitude AGL, resulting in a 240 m swath for each flightline. Data collection occurred with approximately 50% overlap between flightlines, resulting in about one laser sounding per square meter. The data were processed by the USGS Center for Coastal and Watershed Studies to produce 1-meter resolution raster images that can be easily ingested into a Geographic Information System (GIS). The data were organized as 2 km by 2 km data tiles in 32-bit floating-point integer (GeoTIFF) format. Contour lines and hillshade layers were generated from the lidar data tile and incorporated into this map product.

**FURTHER READING**  
Brock, J.C., and Sallenger, Ashbury, 2001, Airborne topographic lidar mapping for coastal science and resource management: U.S. Geological Survey Open File Report 01-46, p. 4.  
Brock, J.C., Wright, C.W., Nayegandhi, Amar, Clayton, Tonya, Hansen, Mark, Longrecker, John, Gesch, Dean, and Crane, Michael, 2002, Initial results from a test of the NASA EAARL lidar in the Tampa Bay Region: Transactions of the Gulf Coast Association of Geologists, v. 52, p. 89-98.  
Wright, C.W. and Brock, J.C., 2002, EAARL: A lidar for mapping shallow coral reefs and other coastal environments, in the Proceedings of the Seventh International Conference on Remote Sensing for Marine and Coastal Environments, Miami, May 20-22, 2002. Ann Arbor, MI, Veridian International Conferences, 1 computer optical disc.



**Fire Island National Seashore  
USGS-NPS-NASA EAARL First Return Topography  
Map Tile 68600e\_451600n**

By  
**John C. Brock<sup>1</sup>, C. Wayne Wright<sup>2</sup>, Matt Patterson<sup>3</sup>, Amar Nayegandhi<sup>1</sup>, and Judd Patterson<sup>3</sup>,**  
<sup>1</sup>USGS Center for Coastal and Watershed Studies, St. Petersburg, FL  
<sup>2</sup>NASA Wallops Flight Facility, Wallops Is., VA  
<sup>3</sup>NPS South Florida/Caribbean Network Inventory and Monitoring Program, Miami, FL  
<sup>4</sup>ETI Professionals, Lakewood, CO

2006

**NAVIG 88 Elevation (meters)**  
-1 0 1 2 3 4 5 6 7 8 9 10 11 12

NO DATA

Universal Transverse Mercator, 1983 North American Datum  
Zone 18 South  
This map is not intended for use in navigation.  
Topography mapped using NASA Experimental  
Advanced Airborne Research Lidar (EAARL)  
November 2002

**USGS**  
science for a changing world

Cooperator:  
NATIONAL PARK SERVICE  
NPS

**NASA**  
science for a changing world

LAND  
AREA OF THIS MAP