



Descriptive and Grade-Tonnage Models and Database for Iron Oxide Cu-Au Deposits

By Dennis P. Cox¹ and Donald A. Singer²

Open-File Report 2007-1155

2007

**U.S. Department of the Interior
U.S. Geological Survey**

¹ Menlo Park, Calif. (retired)

¹ Menlo Park, Calif.

U.S. Department of the Interior
DIRK KEMPTHORNE, Secretary

U.S. Geological Survey
Mark D. Myers, Director

U.S. Geological Survey, Reston, Virginia 2007

For product and ordering information:
World Wide Web: <http://www.usgs.gov/pubprod>
Telephone: 1-888-ASK-USGS

For more information on the USGS—the Federal source for science about the Earth, its natural and living resources, natural hazards, and the environment:
World Wide Web: <http://www.usgs.gov>
Telephone: 1-888-ASK-USGS

Suggested citation:
Cox, D.P. and Singer, D.A., 2007, Descriptive and grade-tonnage models and database for iron oxide Cu-Au deposits: U.S. Geological Survey Open-File Report 2007-1155
[<http://pubs.usgs.gov/of/2007/1155/>].

Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Although this report is in the public domain, permission must be secured from the individual copyright owners to reproduce any copyrighted material contained within this report.

DESCRIPTIVE AND GRADE-TONNAGE MODELS AND DATABASE FOR IRON OXIDE Cu-Au DEPOSITS

By Dennis P. Cox and Donald A. Singer

APPROXIMATE SYNONYMS Ironstone Cu-Au. The acronym is IOCG. (Replaces Olympic Dam Cu-Au-U, Cox, 1986b)

DESCRIPTION Iron oxide Cu-Au deposits are veins and breccia-hosted bodies of hematite and/or magnetite with disseminated Cu +Au ± Ag ± Pd ± Pt ± Ni ± U ± LREE minerals formed in sedimentary or volcano-sedimentary basins intruded by igneous rocks. Deposits are associated with broad redox boundaries and feature sodic alteration of source rocks and potassic alteration of host rocks.

GENERAL REFERENCES Williams and others, 2005; Barton, 2001; Haynes, 2000; Hitzman, 2000; Warren, 1999.

GEOLOGICAL ENVIRONMENT

Rock Types A type or I type, magnetite-series intrusions are commonly associated with IOCG deposits (Hitzman, 2000). As pointed out by Barton and Johnson (2000, p. 47) there is a wide variation in composition of the associated igneous rocks. Of the 31 copper-gold deposits in the accompanying database, 10 deposits are associated with gabbro, diorite or mafic dikes, 2 with felsic porphyry or dacite dikes, one deposit, each for monzonite, tonalite and granodiorite, and 7 deposits with granite. One prospect (visited by Cox in western Mongolia) was associated with syenite. Nine deposits have no record of an associated intrusion.

IOCG deposits are also associated with redbeds and evaporites. These rocks are older than, but close in age to, the intrusive rocks. They are commonly altered to albite-rich assemblages. Of 36 iron and copper-gold deposits in the accompanying database, 21 have associated albite alteration, and six have evidence of an evaporite in the sedimentary section.

Host rocks are faulted and deformed volcanic and sedimentary rocks with bedding-parallel permeability, and volcanic, sedimentary, and tectonic breccias. Less commonly faults

and breccias in intrusive rocks can host ore. The size and degree of permeability of fault and breccia structures in the host rocks directly controls the tonnage of the contained deposit.

Age Range Lower Proterozoic deposits are known in the Carajas region of Brazil and in Mauritania (2.5-2.3 Ga), Australia (1.9-1.5 Ga), and the Khetri Copper Belt of India (1.7-1.5 Ga). See figure 1. Mesozoic deposits occur in the Coast Range of northern Chile (119-90 Ma). Miocene deposits are known in the Andes of Argentina (Dow and Hitzman, 2002). Deposits older than Late Archean are unlikely because an oxygen-rich earth atmosphere is required.

Depositional Environment and Tectonic Setting Hitzman (2000) described two permissive environments for IOCG deposits:

1. Continental margin subduction complexes with local extensional features; Resulting rifts contain oxidized rocks including subaerial volcanic deposits, conglomerates, redbeds and evaporites. Sediments deposited in these environments are sources for oxidizing, hematite-stable, NaCl-rich fluids capable of leaching and transporting copper. Magmatic belts coextensive with these rifts provide heat sources for driving hydrothermal circulation. This environment is represented by the La Candelaria and Punta del Cobre district in Chile.
2. Compression, folding and magmatism of intracratonic basins; Granitic rocks intrude rift-related assemblages, similar to those described above, that have been folded and metamorphosed. Brines released from the rift sediments are mobilized to leach metals from source rocks and deposit them in faults and breccias. The Cloncurry district in Queensland, Australia, provides examples of this environment.

Associated Deposit Types Redbed, Revett, and reduced facies sedimentary copper deposits indicate permissive areas for IOCG. Chilean manto Fe-Cu-Ag deposits (Maksev and Zentilli, 2002), albite-scapolite iron ores (Sokolov and Grigor'ev, 1977) and volcanic-hosted magnetite deposits (Cox, 1986 a) are formed in the same environment.

Iron oxide hosted copper-gold-bismuth deposits in the Tennant Creek Inlier west of the Cloncurry District (Skirrow, 2000) are similar to IOCG deposits. They have smaller tonnages (the median tonnage of 10 deposits is 3.5 million tons) and higher gold grades than IOCG deposits. Bismuth is an important byproduct and gold is locally found as a selenide. Na-Ca

alteration, prominent in IOCG deposits, is lacking in the Tennant Creek District. Because of these differences, Tennant Creek deposits are not included in the IOCG model.

DEPOSIT DESCRIPTION

Mineralogy Principal minerals are magnetite, hematite, and siderite with biotite, calc-silicate minerals and minor apatite. Iron oxides have Fe:Ti ratio greater than iron oxides in most igneous rocks (Williams and others, 2005). Fe oxide Cu-Au deposits contain chalcopyrite, bornite, and gold and, less commonly, chalcocite. Pyrite or pyrrhotite are present in most deposits. Less common minerals include bastnaesite, uraninite, monazite, allanite, Pt group minerals, molybdenite, sphalerite, galena, bismuthinite, scheelite, arsenopyrite, cobaltite, and Ni-Co arsenides. Gangue minerals are quartz, biotite, calc-silicate minerals, scapolite (marialite variety, $3\text{NaAlSi}_3\text{O}_8\cdot\text{NaCl}$), albite, fluorite, fluorapatite, calcite, barite and tourmaline. Quartz vein stockworks, common to porphyry copper deposits, are not present in IOCG deposits (Sillitoe, 2003).

Texture/Structure Ore minerals form veins and disseminations in lenticular, elongate iron oxide bodies. Open space filling in faults, tension gashes, and breccias are common (Michael Evans, Phelps Dodge Corp., written commun., 2002). Mineralized replacement features may follow bedding and other sedimentary structures. Iron oxide and copper sulfide minerals fill spaces in the matrix of breccia pipes and irregular breccia bodies of sedimentary, volcanic, or tectonic origin.

Alteration Extensive albite-oligoclase-chlorite-actinolite alteration with Na-scapolite is present in sedimentary, volcanic, and plutonic rocks near the deposits. The albite zone is depleted in K, Fe, U, REE, S, and most base metals. Albite-oligoclase alteration is also referred to as Na-Ca alteration (Carten, 1986).

Host rocks exhibit potassic (biotite-K-feldspar) alteration and may contain chlorite, pyroxene, amphibole, epidote, garnet, Na-scapolite, and anhydrite. This alteration cuts or overprints albite-oligoclase alteration.

Calcite-dolomite-pyrite veins and disseminations occur outboard of the iron oxide bodies and cut ore-related alteration (G. McKelvey, Phelps Dodge Corp., oral commun., 2002).

Ore and alteration mineral facies are asymmetric (Sillitoe, 2003) reflecting formation on a redox boundary with bornite, chalcocite, magnetite and biotite on the oxidized side and chalcopyrite and distal pyrite and calcite on the reduced side. At Olympic Dam and Prominent Hill in Australia, mineral zonation is vertical with chalcocite at the top grading downward to bornite, chalcopyrite, and pyrite at depth.

Ore Controls Pre-ore permeability of host rocks and faults and shear zones combined with redox fronts are the major ore controls of ore deposition. The largest deposits are hosted by breccias of tectonic sedimentary, volcanic or phreatomagmatic origin. Major shear zones are also important loci of IOCG deposition. Small deposits form as fault-controlled veins.

Geochemical Signature The elements Cu + Au ± Bi ± U ± Ni ± Co ± PGE ± REE are present in K-rich rocks. All of these elements can form chlorides or chloride complexes in the presence of NaCl fluids. In rocks with albite alteration, the above elements have abundances close to the limit of detection.

Deposit Tonnage and Grade Ore tonnages vary widely with more than 3.8 billion tons at Olympic Dam, Australia (Williams and others, 2005), and 470 million tons at La Candelaria, Chile. Tonnages of iron-only deposits range from 700 million to 2 billion tons or more.

Within the 33 IOCG deposits in the accompanying file, two distinct tonnage-grade populations can be recognized (Table 1): 12 consist of deposits hosted in faults or veins (labeled as fault-/vein-hosted type in the database) and 21 consist of deposits hosted in breccias or fault-zones (labeled as fault-zone-/breccia-hosted). The difference in tonnage (median 3.5 Mmt vs. 120 Mmt for fault zone-/breccia-hosted) reflects the type and extent of permeability in the host rock and the amount of open space available for mineral deposition. The differences in grade (2.3 percent Cu versus 1.1 percent for fault-zone-/breccia-hosted) may reflect the economies of scale, such that large tonnage deposits with lower grade ores can be mined more cheaply by bulk mining methods.

Of the twelve fault-vein-hosted deposits, six are in Queensland, Australia and have no recognized associated intrusion, and six are in northern Chile and are associated with diorite intrusions.

GENETIC OVERVIEW

Although some authors have shown evidence for a magmatic origin of the ore fluids responsible for mineralization (Pollard, 2000; Kendrick and others, 2007), the present authors believe that the accompanying database shows that the primary role of magmas was the production of heat and that their contribution of metals to the ore fluid was minor. Evidence for this conclusion are the wide variation in composition of associated igneous rocks, and the presence of albite alteration (in 21 deposits) and Na-scapolite (17 deposits) suggesting a widespread reaction with NaCl brines. Brine influx of such a magnitude is more likely to have originated from sedimentary sources than from igneous intrusions. In our database we have attempted to identify the sedimentary, and/or volcanic rocks surrounding the deposit as the source of the brine and ore-metals. We were successful in 19 of the 36 cases studied.

Mafic to granitic magmas with low volatile content activate oxidized NaCl brines from evaporites and other sedimentary sources. These brines, heated by the intrusions, leach K, Fe, Cu, Au and other metals from the surrounding rocks and form metal chloride complexes at near magmatic temperatures (Barton, 2001). The oxidized brines migrate from the source rocks into surrounding country rocks. Ore deposition occurs where the brines mix with reduced fluids in permeable breccias and in fault zones. Skarn-like mineral assemblages in the ore zone are produced in response to heat from the associated magmas. Potassic alteration results from reaction of KCl in the brine with mafic silicates in the host rock. Reduced fluids deposit pyrite and calcite outboard of the ore zone. The highly variable metal composition of the ores depends in large part on the composition of different source rocks leached by the NaCl brine.

Iron-only deposits (albite-scapolite iron deposits) may reflect ore deposition in environments where reduced sulfur is unavailable. Haynes (2002) pointed out that districts with large iron-only deposits (Sokolovskaya, Kachar) generally do not contain significant IOCG deposits. Conversely, in districts with large IOCG deposits (i.e. Punta del Cobre district, Chile) iron-only deposits (Marcona, Peru and El Romeral, Chile) are smaller.

Iron oxide Cu-Au deposits are similar to sediment-hosted copper deposits in the alteration mineralogy of source rocks and host rocks. Hayes (1990) described albite-chlorite alteration of redbed source rocks, and distal pyrite-calcite outboard of copper deposits in the Revett Quartzite of Montana and Idaho.

EXAMPLES

La Candelaria, Chile (Marschik and Fontboté, 2001)

Starra, Australia (Rotherham and others, 1998)

Ernest Henry, Australia (Mark and others, 2000)

ACKNOWLEDGEMENTS

The authors thank Dan Mosier, David John, and Tom Frost of the USGS for thoughtful reviews of this report.

REFERENCES CITED

- Adshead-Bell, N.D., Voulgaris, P., and Muscio, V.N., 1998, Osbourne copper-gold deposit, *in* Berkman, D.A., and MacKenzie, D.H., eds., *Geology of Australian and Papua New Guinean mineral deposits: Australasian Institute of Mining and Metallurgy*, 22, p. 793-799.
- Baker, T., 1998, Alteration, mineralization, and fluid evolution at the Eloise Cu-Au deposit. Cloncurry district, northwest Queensland, Australia: *Economic Geology*, v. 93, no. 8, p. 1213-1236.
- Barton, M.D., and Johnson D.A., 2000, Alternative brine sources for Fe-oxide(Cu-Au) systems: Implications for hydrothermal alteration and metals, *in* Porter, T.M., ed., *Hydrothermal iron oxide copper-gold and related deposits: a global perspective*, v. 1: PGC Publishing, Linden Park, Australia, p. 43-60.
- Barton, M.D., 2001, Sodic alteration and Fe-oxide-rich hydrothermal systems [abs.]: *Geological Society of America Abstracts with Programs*, v.33, no. 6, p. 4.
- Belperino, Antonio, and Freeman, Hamish, 2004, Common geological characteristics of Prominent Hill and Olympic Dam—implications for iron oxide copper-gold exploration models: *Australian Institute of Mining and Metallurgy Bull.* no. 6, Nov.-Dec. 2004, p. 67-75.
- Carten, R.B., 1986, Sodium-calcium metasomatism; chemical, temporal, and spatial relationships at the Yerington, Nevada, porphyry copper deposit: *Economic Geology*, v. 81, no. 6, p. 1495-1519.
- Cox, D.P., 1986a, Descriptive model of volcanic-hosted magnetite, *in* Cox, D.P., and Singer, D.A., eds., *Mineral Deposit Models: U.S. Geological Survey Bulletin 1693*, p. 172.
- Cox, D.P., 1986b, Descriptive model of Olympic Dam Cu-Au-U, *in* Cox, D.P., and Singer, D.A., eds., *Mineral Deposit Models: U.S. Geological Survey Bulletin 1693*, p. 200.
- Davidson, G.J., Davis, B.K., and Garner, Andrew, 2002, Structural and geochemical constraints on the emplacement of the Monakoff oxide Cu-Au (-Co-U-REE-Ag-Zn-Pb) deposit, Mount Isa Inlier, Australia, *in* Porter, T.M. ed., *Hydrothermal iron oxide copper-gold and related deposits: a global perspective*, v. 2 PGC Publishing, Linden Park, Australia, p. 49-75.

- Dow, R.J., and Hitzman, M.W., 2002 Geology of the Arizaro and Lindero prospects, Salta province, northwest Argentina: Mid Miocene hydrothermal Fe-Ox copper-gold mineralization, *in* Porter, T.M. ed. Hydrothermal iron oxide copper-gold and related deposits: a global perspective, v. 2 PGC Publishing, Linden Park, Australia, p. 153-161.
- Hawkes, Nicholas, Clark, A.H., and Moody, T.C., 2002. Marcona and Pampa de Pongo giant Mesozoic Fe-(Cu-Au) deposits in the Peruvian Coastal Belt, *in* Porter, T.M. ed., Hydrothermal Iron oxide copper-gold and related deposits: a global perspective, v. 2 PGC Publishing, Linden Park, Australia, p. 115-130.
- Hayes, T.S, 1990, A preliminary study of thermometry and metal sources of the Spar Lake stratabound copper-silver deposit, Belt Supergroup, Montana: U.S. Geological Survey Open File Report 90-0484, 30 p.
- Haynes, D.W., 2000, Iron oxide copper gold deposits: their position in the ore deposit spectrum and modes of origin, *in* Porter, T.M., ed., Hydrothermal iron oxide copper-gold and related deposits: a global perspective: Australian Mineral Foundation, Adelaide, p. 71-90.
- Haynes, D.W., 2002, Giant iron oxide-copper-gold deposits: are they in distinctive geologic settings? *in* Cooke, D.R. and Pongratz, June, Giant ore deposits: Characteristics, genesis, and exploration: Centre for Ore Deposits Research Univ. of Tasmania, Hobart, p. 57-78.
- Hitzman, M. W, 2000, Iron oxide-Cu-Au deposits: what, where, when, and why, *in* Porter, T.M., ed., Hydrothermal iron oxide copper-gold and related deposits: a global perspective: Australian Mineral Foundation, Adelaide, p. 9-25.
- Hopper, David, and Correa, Arturo, 2000, The Panucillo and Teresa de Colmo copper deposits: two contrasting examples of Fe-Ox Cu-Au mineralization from the Coastal Cordillera, Central Chile, *in* Porter, T.M. ed., Hydrothermal iron oxide copper-gold and related deposits: a global perspective, v. 1, PGC Publishing, Linden Park, Australia, p. 177-189.
- Huston, D.L., Bolger, Chris, and Cozens, Greg, 1993, A comparison of mineral deposits at Gecko, and White Devil deposits: implications for ore genesis in the Tennant Creek district, Northern Territory, Australia: Economic Geology, v.88, p. 1198-1225.
- Kendrick, M.A., Mark, G., and Phillips D., 2007, Mid-crustal fluid mixing in a Proterozoic Fe oxide-Cu-Au deposit, Earnest Henry, Australia: Evidence from Ar, Kr, Xe, Cl, Br, and I: Earth and Planetary Science Letters, 256, p. 238-343.
- Knight, Joe, Joy, Sojen, Lowe, Jon, Cameron, John, Merilees, James, Nag, Sudipta, Shah, Nalin, Dua, Gaurav, and Jhala, Khamalendra, 2002, The Khetri Copper Belt, Rajasthan: iron oxide copper gold terrane in the Proterozoic of northwest India, *in* Porter, T.M. ed., Hydrothermal Iron oxide copper-gold and related deposits: a global perspective, v. 2, PGC Publishing, Linden Park, Australia, p. 321-341.
- Kolb, Jochen, Sakellaris, G.A., and Meyer, F.M., 2006, Controls on hydrothermal Fe oxide-Cu-Au-Co mineralization at the Gelb Moghrein deposit, Akjoujt area, Mauritania: Mineralium Deposita, v. 41, p. 68-81.
- Kremarov, R.L., and Stewart, J.I., 1995, Geology and mineralization of the Greenmount Cu-Au-Co deposit, southeastern Marimo Basin: Australian Journal of Earth Science, v. 45, p. 463-482.
- Maksaev, Victor, and Zentilli, Marcos, 2002, Chilean strata-bound Cu-(Ag) deposits: an overview system, *in* Porter, T.M., ed., Hydrothermal iron oxide copper-gold and related deposits: a global perspective, V. 2: PGC Publishing, Linden Park, Australia, p. 185-205.
- Mark, Geordie, Oliver, N.H.S., Williams, P.J., Valenta, R.K., and Crookes, R.A., 2000, The evolution of the Ernest Henry Fe-oxide-(Cu-Au) hydrothermal system, *in* Porter, T.M., ed., Hydrothermal iron oxide copper-gold and related deposits: a global perspective, v. 1: PGC Publishing, Linden Park, Australia, p. 123-136.

- Marschik, Robert, 2001, Iron oxide Cu-Au deposits in South America: Candelaria Chile and Sossego, Brazil, [abs.] Geological Society of America Abstracts with Programs, v. 33, no. 6, p. 2.
- Marschik, Robert, and Fontboté, Lluís, 1996, Copper (-iron) mineralization and superposition of alteration events in the Punta del Cobre belt, northern Chile, *in* Camus, F., Sillitoe, R.M., Peterson, R., and Sheahan, P., eds., Andean copper deposits: new discoveries, mineralization styles, and metallogeny: Economic Geology, Special Publication 5, p. 171-190.
- Marschik, Robert, and Fontboté, Lluís, 2001, The Candelaria—Punta del Cobre iron oxide Cu-Au (-Zn-Ag) deposits, Chile: Economic Geology, v. 96, p. 1799-1826.
- Marschick, R., Leveille, R.A., and Martin, Walter, 2000, La Candelaria and the Punta del Cobre district: Early Cretaceous iron-oxide Cu-Au (-Zn-Ag) mineralization, *in* Porter, T.M. ed., Hydrothermal iron oxide copper-gold and related deposits: a global perspective, v. 1 PGC Publishing, Linden Park, Australia, p. 49-75.
- McLean, R.N., 2002, The Sin Quyen iron oxide-copper-gold-rare earth oxide mineralization of North Vietnam, *in* Porter, T.M. ed. Hydrothermal iron oxide copper-gold and related deposits: a global perspective, v. 2, PGC Publishing, Linden Park, Australia, p. 293-301.
- Meyer, F.M., Kolb, Jochen, Sakellaris, G. A., and Gerds, Axel, 2006, New ages from the Mauritanides Belt: recognition of Archean IOCG mineralization at Gelb Moghreïn, Mauritania: Terra Nova, v. 18, no. 5, p. 345-352.
- Oreskes, Naomi, and Einaudi, M.T., 1990, Origin of rare-earth element-enriched hematite breccias at the Olympic Dam Cu-U-Au-Ag deposit, Roxby Downs, South Australia: Society of Economic Geologists, v.85, p. 1-28.
- Oreskes, Naomi, and Einaudi, M.T., 1992, Origin of hydrothermal fluids at Olympic Dam: preliminary results from fluid inclusions and stable isotopes: Society of Economic Geologists, v. 87, no. 1, p. 64-90.
- Pollard, P.J., 2000, Evidence of a magmatic fluid and metal source for Fe-oxide Cu-Au mineralization, *in* Porter, T.M. ed., Hydrothermal iron oxide copper-gold and related deposits: a global perspective, v. 1 PGC Publishing, Linden Park, Australia, p. 27-42.
- Ray, G.E., and Lawrence A.D., 2002, The Productora prospect in north-central Chile: an example of an intrusion-related, Candelaria type Fe-Cu-Au hydrothermal system, *in* Porter, T.M. ed., Hydrothermal iron oxide copper-gold and related deposits: a global perspective, v. 2, PGC Publishing, Linden Park, Australia, p. 131-151.
- Requia, Karin, 2001, The Salobo iron oxide copper-gold hydrothermal system, Carajas Mineral Province, Brazil [abs.]: Geological Society of America Abstracts with Programs, v. 33, no. 6, p. 2.
- Requia, Karin, and Fontboté, Lluís, 1999, Hydrothermal alkali metasomatism in the Salobo iron oxide Cu (-Au) deposit, Carajás mineral province, northern Brazil, *in* Stanley, C.J. (ed.), Mineral deposits: processes to processing, Balkema, Amsterdam, p.1025-1028.
- Rotherham, J.F., Blake, K.L., Cartwright, Ian, and Williams, P.J., 1998, Stable isotope evidence for the origin of the Mesoproterozoic Starra Au-Cu deposit, Cloncurry district, northwest Queensland: Economic Geology, v. 93, no. 8, p. 1435-1449.
- Ryan, P.J., Lawrence, A.L., Jenkins, R.A., Matthews, J.P., Zamora, J.C., Marino, E.W., and Diaz, I.U., 1995, The Candelaria copper-gold deposit, Chile, *in* Pierce, F.W., and Bolm. J.G., eds., Porphyry copper deposits of the American Cordillera: Arizona Geological Society Digest v. 20, p. 625–645.
- Sillitoe, R.H., 2003, Iron oxide copper-gold deposits: an Andean view: Mineralium Deposita, v. 38, no. 7, p. 787-812.

- Skirrow, R.G., 2000, Gold-copper-bismuth deposits of the Tennant Creek District, Australia: a reappraisal of diverse high-grade systems, *in* Porter, T.M. ed., Hydrothermal iron oxide copper-gold and related deposits: a global perspective, v. 1, PGC Publishing, Linden Park, Australia, p. 149-160.
- Sokolov, G.A., and Grigor'ev, V.M., 1977, Deposits of iron, *in* Smirnov, V.I., ed., Ore deposits of the USSR, v. 1, Pitman Publishing, London, p. 7-113.
- Tallarico, F.H.B., Feigueiredo, B.R., Groves, D.I., Kositcin, N., Mc Naughton, I.R., Fletcher, I.R., and Rego, J.L., 2005, Geology and SHRIMP geochronology of the Igarape Bahia deposit, Carajas Copper Gold Belt, Brazil: an Archean (2.75 Ga) example of iron oxide Cu-Au-(U-REE) mineralization: *Economic Geology*, v. 100, no. 7, p. 7-28.
- Vila, Tomás, Lindsay, Nicholas, and Zamora, Richard, 1996, Geology of the Manto Verde copper deposit, northern Chile: a specularite-rich hydrothermal-tectonic breccia related to the Atacama fault zone, *in* Camus, Francisco, Sillitoe, R.M., Peterson Richard, and Sheahan, Patricia, eds., Andean copper deposits: new discoveries, mineralization, styles and metallogeny: Society of Economic Geology Special Publication no. 5, p.158-170.
- Warren, John, 1999, Evaporites, their evolution and economics, Blackwell Science, Oxford, 438 p.
- Williams, P.J., Barton, M.D, Johnson, D.A., Fontboté, Lluís, De Haller, Antoine, Mark, Geordie, Oliver, N.H.S., and Marschik, Robert, 2005, Iron oxide copper-gold deposits: geology, space-time distribution and possible modes of origin, *in* Hedenquist, J.W., Thompson, J.F.H., Goldfarb, R.J., and Richards, J.P., eds., *Economic Geology 100th anniversary volume, 1902-2005*, p. 371-405.
- Williams, P.J., and Skirrow, R.G. 2000, Overview of iron oxide-copper-gold deposits in the Curnamona Province and Cloncurry district (Eastern Mount Isa Block), Australia, *in* Porter, T.M. ed., Hydrothermal iron oxide copper-gold and related deposits: a global perspective, v. 1: PGC Publishing, Linden Park, Australia, p. 105-122.
- Zang, W., and Fyfe, W.S, 1995, Chloritization of the hydrothermally altered bedrock at the Igarapé Bahia gold deposit, Carajás, Brazil: *Mineralium Deposita* v. 30, no. 1, p. 30-38.

TABLE

Table 1. Percentile distribution of ore tonnage and copper and gold grades in two subtypes of IOCG deposits.

	Fault-/vein-hosted	Fault-zone-/breccia-hosted
<u>number</u>	<u>12</u>	<u>21 deposits</u>
90th Tons	0.65	25 million mt
50th Tons	3.5	120 million mt
10th Tons	15	1,100 million mt
90th Cu	1.2	0.41 percent
50th Cu	2.3	1.1 percent
10th Cu	4.9	1.7 percent
90th Au	0.14	0.13 grams/ton
50th Au	1	0.48 grams/ton
10th Au	5.8	2.1 grams/ton




Figure 1 World distribution of 33 iron oxide Cu-Au deposits and 3 albite-scapolite iron deposits, numbers 19, 20, 21.