

# η(1295)

$$I^G(J^{PC}) = 0^+(0^-+)$$

See also the mini-review under non- $q\bar{q}$  candidates. (See the index for the page number.)

## η(1295) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1297.0±2.8 OUR AVERAGE</b>				
1299 ±4	2100	ALDE	97B GAM4	100 π <sup>-</sup> p → ηπ <sup>0</sup> π <sup>0</sup> n
1295 ±4		FUKUI	91C SPEC	8.95 π <sup>-</sup> p → ηπ <sup>+</sup> π <sup>-</sup> n
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 1275		STANTON	79 CNTR	8.4 π <sup>-</sup> p → nη2π

## η(1295) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>53±6</b>				
		FUKUI	91C SPEC	8.95 π <sup>-</sup> p → ηπ <sup>+</sup> π <sup>-</sup> n
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<40	2100	ALDE	97B GAM4	100 π <sup>-</sup> p → ηπ <sup>0</sup> π <sup>0</sup> n
~ 70		STANTON	79 CNTR	8.4 π <sup>-</sup> p → nη2π

## η(1295) DECAY MODES

Mode	Fraction (Γ <sub>i</sub> /Γ)
Γ <sub>1</sub> ηπ <sup>+</sup> π <sup>-</sup>	seen
Γ <sub>2</sub> a <sub>0</sub> (980)π	seen
Γ <sub>3</sub> γγ	
Γ <sub>4</sub> ηπ <sup>0</sup> π <sup>0</sup>	seen
Γ <sub>5</sub> η(ππ) <sub>S-wave</sub>	seen

## η(1295) Γ(i)Γ(γγ)/Γ(total)

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	Γ <sub>1</sub> Γ <sub>3</sub> /Γ
<b>Γ(ηπ<sup>+</sup>π<sup>-</sup>) × Γ(γγ)/Γ<sub>total</sub></b>					
<b>&lt;0.3</b>					
		ANTREASYAN	87 CBAL	e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> ηππ	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<0.6	90	AIHARA	88C TPC	e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> ηπ <sup>+</sup> π <sup>-</sup>	

## $\eta(1295)$ BRANCHING RATIOS

### $\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	BERTIN	97 OBLX	0.0 $\bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
seen	BIRMAN	88 MPS	8 $\pi^-p \rightarrow K^+\bar{K}^0\pi^-n$
large	ANDO	86 SPEC	8 $\pi^-p \rightarrow \eta\pi^+\pi^-n$
large	STANTON	79 CNTR	8.4 $\pi^-p \rightarrow n\eta 2\pi$

### $\Gamma(a_0(980)\pi)/\Gamma(\eta\pi^0\pi^0)$ $\Gamma_2/\Gamma_4$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.65 ± 0.10</b>	<sup>1</sup> ALDE	97B GAM4	100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$

<sup>1</sup> Assuming that  $a_0(980)$  decays only to  $\eta\pi$ .

### $\Gamma(\eta(\pi\pi)_{S\text{-wave}})/\Gamma(\eta\pi^0\pi^0)$ $\Gamma_5/\Gamma_4$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.35 ± 0.10</b>	ALDE	97B GAM4	100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$

## $\eta(1295)$ REFERENCES

ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)
BERTIN	97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.)
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP
ANTREASYAN	87	PR D36 2633	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+) IJP
STANTON	79	PRL 42 346	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+) JP