

## SUNYAEV-ZELDOVICH EFFECT IMAGING OF MASSIVE CLUSTERS OF GALAXIES AT REDSHIFT $z > 0.8$

MARSHALL JOY,<sup>1</sup> SAMUEL LAROQUE,<sup>2</sup> LAURA GREGO,<sup>3</sup> JOHN E. CARLSTROM,<sup>2</sup> KYLE DAWSON,<sup>4</sup> HARALD EBELING,<sup>5</sup>  
WILLIAM L. HOLZAPFEL,<sup>4</sup> DAISUKE NAGAI,<sup>2</sup> AND ERIK D. REESE<sup>2</sup>  
*Received 2000 December 4; accepted 2001 March 2; published 2001 March 28*

### ABSTRACT

We present Sunyaev-Zeldovich effect (SZE) imaging observations of three distant ( $z > 0.8$ ) and highly X-ray luminous clusters of galaxies, Cl J1226.9+3332, Cl J0152.7–1357, and MS 1054.4–0321. Two of the clusters, Cl J1226.9+3332 and Cl J0152.7–1357, were recently discovered in deep *ROSAT* X-ray images. Their high X-ray luminosity suggests that they are massive systems, which, if confirmed, would provide strong constraints on the cosmological parameters of structure formation models. Our SZE data provide confirmation that they are massive clusters similar to the well-studied cluster MS 1054.4–0321. Assuming the clusters have the same gas mass fraction as that derived from SZE measurements of 18 known massive clusters, we are able to infer their mass and electron temperature from the SZE data. The derived electron temperatures are  $9.8^{+4.7}_{-1.9}$ ,  $8.7^{+4.1}_{-1.8}$ , and  $10.4^{+5.0}_{-2.0}$  keV, respectively, and we infer total masses of  $\sim 2 \times 10^{14} h_{100}^{-1} M_{\odot}$  within a radius of  $65''$  ( $340 h_{100}^{-1}$  kpc) for all three clusters. For Cl J0152.7–1357 and MS 1054.4–0321, we find good agreement between our SZE-derived temperatures and those inferred from X-ray spectroscopy. No X-ray-derived temperatures are available for Cl J1226.9+3332, and thus the SZE data provide the first confirmation that it is indeed a massive system. The demonstrated ability to determine cluster temperatures and masses from SZE observations without access to X-ray data illustrates the power of using deep SZE surveys to probe the distant universe.

*Subject headings:* cosmic microwave background — cosmology: observations —  
galaxies: clusters: individual (Cl J0152.7–1357, Cl J1226.9+3332, MS 1054.4–0321) —  
techniques: interferometric

*On-line material:* color figure

### 1. INTRODUCTION

The existence of galaxy clusters at high redshift can place powerful constraints on the physical and cosmological parameters of structure formation models (Bahcall & Cen 1992; Lupino & Gioia 1995; Oukbir & Blanchard 1997; Donahue et al. 1998; Eke et al. 1998; Haiman, Mohr, & Holder 2001). The greatest leverage is provided by the most massive and distant clusters (e.g., Viana & Liddle 1996). Two distant and highly X-ray luminous clusters were recently discovered in deep *ROSAT* X-ray images: Cl J1226.9+3332, a cluster at redshift  $z = 0.89$ , was discovered in the Wide-Angle *ROSAT* Pointed Survey (WARPS; Ebeling et al. 2001; Scharf et al. 1997). Cl J0152.7–1357, at redshift  $z = 0.83$ , was detected in the *ROSAT* Deep Cluster Survey, the Serendipitous High-Redshift Archival *ROSAT* Cluster survey, and WARPS (Della Ceca et al. 2000; Romer et al. 2000; Ebeling et al. 2000). Based on their X-ray luminosities [ $L_X$  (0.5–2 keV)  $\geq 2 \times 10^{44} h_{100}^{-2}$  ergs  $s^{-1}$ ], these clusters are thought to be highly massive, which, if confirmed, will provide significant constraints on cosmological models (Bahcall & Fan 1998).

In this Letter, we present interferometric imaging of the Sunyaev-Zeldovich effect (SZE) in these clusters, which provides a measure of the gas pressure integrated along the line of sight (Sunyaev & Zeldovich 1972; Birkinshaw 1999). The change in the observed brightness temperature of the cosmic

microwave background (CMB) radiation that results from passage through the thermally ionized gas permeating a galaxy cluster is given by

$$\frac{\Delta T_{\text{thermal}}}{T_{\text{CMB}}} = f(\nu) \frac{k_B \sigma_T}{m_e c^2} \int n_e T_e dl,$$

where  $T_{\text{CMB}}$  is the microwave background temperature (2.7 K),  $\sigma_T$  is the Thomson scattering cross section, and  $m_e$ ,  $n_e$ , and  $T_e$  are the electron mass, density, and temperature. The frequency dependence of the SZE is represented by  $f(\nu)$ ; in the Rayleigh-Jeans limit,  $f(\nu) = -2$ . We use the SZE to determine the mass of Cl J1226.9+3332 and Cl J0152.7–1357 using the methods developed by Grego et al. 2001; in addition, we present interferometric SZE data on MS 1054.4–0321, a cluster of known temperature and mass at  $z = 0.83$  (Gioia & Luppino 1994; Donahue et al. 1998; Hoekstra, Franx, & Kuijken 2000), which provides a standard against which the newly discovered clusters can be compared.

The SZE observations and data analysis are described in § 2, and the conclusions drawn from these data will be found in § 3. Throughout this Letter, we parameterize the Hubble constant in terms of  $h_{100}$ , where  $H_0 \equiv 100 h_{100} \text{ km s}^{-1} \text{ Mpc}^{-1}$ , and we use the cosmological parameters  $\Omega_M = 0.3$  and  $\Omega_{\Lambda} = 0.7$  unless otherwise stated. Uncertainties are reported at the 68% confidence level.

### 2. INTERFEROMETRIC IMAGING OF THE SUNYAEV-ZELDOVICH EFFECT

#### 2.1. Observations

To image the SZE in these distant clusters, we outfitted the Owens Valley Radio Observatory (OVRO) and Berkeley-Illinois-Maryland Association (BIMA) millimeter interferome-

<sup>1</sup> Department of Space Science, SD50, NASA Marshall Space Flight Center, Huntsville, AL 35812.

<sup>2</sup> Department of Astronomy and Astrophysics, University of Chicago, 5640 South Ellis Avenue, Chicago, IL 60637.

<sup>3</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138.

<sup>4</sup> Physics Department, University of California at Berkeley, 366 LeConte Hall, Berkeley, CA 94720.

<sup>5</sup> Institute for Astronomy, 2680 Woodlawn Drive, Honolulu, HI 96822.

TABLE 1  
CLUSTER OBSERVATION LOG

CLUSTER NAME	REDSHIFT	POINTING CENTER (J2000)		ON-SOURCE INTEGRATION TIME (hr)
		$\alpha$	$\delta$	
Cl J1226.9+3332 .....	0.89	12 26 58.0	33 32 45 <sup>a</sup>	41.6
Cl J0152.7-1357 .....	0.83	01 52 43.0	-13 57 29 <sup>b</sup>	27.8
MS 1054.4-0321 .....	0.83	10 56 59.5	-03 37 28 <sup>c</sup>	43.0

NOTE.—Units of right ascension are hours, minutes, and seconds, and units of declination are degrees, arcminutes, and arcseconds.

<sup>a</sup> Pointing center coincident with X-ray cluster center from Ebeling et al. 2001.

<sup>b</sup> Pointing center coincident with X-ray cluster center from Ebeling et al. 2000.

<sup>c</sup> Pointing center coincident with optical cluster center from Gioia & Luppino 1994. The ROSAT/High Resolution Imager X-ray center is at 10<sup>h</sup>56<sup>m</sup>58<sup>s</sup>.6, -03<sup>o</sup>37'36" (Neumann & Arnaud 2000).

ters with sensitive centimeter-wave receivers optimized for SZE measurements (Carlstrom, Joy, & Grego 1996). Our receivers use cryogenically cooled 26–36 GHz high electron mobility transistor amplifiers (Pospieszalski et al. 1995), with characteristic receiver temperatures of  $T_{\text{rx}} \sim 11\text{--}20$  K at the 28.5 GHz frequency used for these observations. The cluster pointing centers and on-source integration times are given in Table 1.

The interferometric measurements of Cl J1226.9+3332 and Cl J0152.7-1357 were made at the BIMA interferometer in 1998 and 2000 with nine 6.1 m antennas in a closely packed configuration to maximize sensitivity to the SZE, with a 6'6 FWHM primary beam and baselines ranging from 0.6 to 14.3 k $\lambda$  (6–140 m). Typical system temperatures, scaled to above the atmosphere, are  $\sim 40\text{--}45$  K in an 800 MHz band centered at 28.5 GHz. Observations of a bright phase calibrator were interleaved with cluster measurements every 25 minutes, and Mars was used for amplitude calibration (Rudy 1987; Grego et al. 2000). The MIRIAD software package (Sault, Teuben, & Wright 1995) was used to calibrate and edit the visibility data and to output the reduced data in UVFITS format for subsequent analysis.

The interferometric measurements of MS 1054.4-0321 were made at the OVRO millimeter array in 1996 June with six 10.4 m antennas in a closely packed configuration, with a 4'2 FWHM primary beam and baselines ranging from 1.0 to 12.0 k $\lambda$  (10–120 m). Typical system temperatures, scaled to above the atmosphere, are  $\sim 45$  K in two 1 GHz channels centered at 28.5 and 30.0 GHz (2 GHz total bandwidth). Observations of a bright phase calibrator were interleaved with cluster measurements every 24 minutes. The MMA software package (Scoville et al. 1993) was used to calibrate and edit the visibility data and to output the reduced data in UVFITS format for subsequent analysis.

We flagged data from baselines when one of the telescopes was shadowed by another telescope in the array, cluster data that

were not bracketed in time by phase calibrator data (mainly at the beginning or end of a track), data for which the phase calibrator indicated poor atmospheric coherence, and, rarely, data with spurious correlations.

## 2.2. Data Analysis

In order to properly model the cluster, we must account for any point sources in the field. To identify these point sources, we used DIFMAP (Pearson et al. 1994) to produce a high-resolution image, using only data from baselines longer than 20 m. The resulting synthesized beam sizes, rms noise levels, and point-source detections are given in Table 2.

We perform a quantitative analysis of the observed SZE profiles by fitting isothermal  $\beta$  models (Cavaliere & Fusco-Femiano 1976, 1981) and point-source profiles to the interferometric data directly in the Fourier ( $u, v$ )-plane, where the noise characteristics and the spatial filtering of the interferometer are well understood. The spherical isothermal  $\beta$  model density is described by

$$n_e(r) = n_{e0} \left( 1 + \frac{r^2}{r_c^2} \right)^{-3\beta/2}, \quad (1)$$

where the core radius  $r_c$  and  $\beta$  are shape parameters, and  $n_{e0}$  is the central electron number density. With this model, the SZE temperature decrement is

$$\Delta T(\theta) = \Delta T_0 \left( 1 + \frac{\theta^2}{\theta_c^2} \right)^{1/2-3\beta/2}, \quad (2)$$

where  $\theta = r/D_A$ ,  $\theta_c = r_c/D_A$ ,  $D_A$  is the angular diameter distance, and  $\Delta T_0$  is the temperature decrement at zero projected radius.

We determine the best-fit point-source positions and fluxes, as well as the cluster centroid, using a simplex algorithm that minimizes the  $\chi^2$  statistic (Reese et al. 2000). We fix the cluster

TABLE 2  
RADIO POINT SOURCES

Cluster Name	Synthesized Beam ( $r_{\text{av}} > 2$ k $\lambda$ ) (arcsec)	$\Delta\alpha$ (arcsec)	$\Delta\delta$ (arcsec)	Observed Flux (mJy) <sup>a</sup>	rms Noise (mJy beam <sup>-1</sup> )
Cl J1226.9+3332 .....	14.1 $\times$ 16.1	260.0	-39.3	1.73	0.152
Cl J0152.7-1357 .....	13.7 $\times$ 23.1	...	...	...	0.221
MS 1054.4-0321 .....	17.4 $\times$ 23.3	0.1	1.4	0.98	0.092
	17.4 $\times$ 23.3	-161.0	1.9	0.63	0.092
	17.4 $\times$ 23.3	-25.5	-86.9	0.35	0.092

<sup>a</sup> Uncorrected for primary-beam attenuation.




FIG. 1.—Synthesized images of the SZE decrement in CL J1226.9+3332, CL J0152.7–1357, and MS 1054.4–0321. *Left panel:* Synthesized image of CL J1226.9+3332, obtained by applying a Gaussian taper with a half-power radius of  $1 \text{ k}\lambda$  to the  $u$ - $v$  data, yielding a resolution of  $99''.3 \times 87''.4$  at position angle (P.A.) =  $32^\circ$ . Contours are multiples of  $290 \mu\text{Jy beam}^{-1}$  ( $1.5 \sigma$ ), and the rms is  $190 \mu\text{Jy beam}^{-1}$ . *Center panel:* Synthesized image of CL J0152.7–1357, obtained by applying a Gaussian taper with a half-power radius of  $1 \text{ k}\lambda$  to the  $u$ - $v$  data, yielding a resolution of  $151'' \times 87''.9$  at P.A. =  $4^\circ$ . Contours are multiples of  $480 \mu\text{Jy beam}^{-1}$  ( $1.5 \sigma$ ), and the rms is  $320 \mu\text{Jy beam}^{-1}$ . *Right panel:* Synthesized image of MS 1054.4–0321, obtained by applying a Gaussian taper with a half-power radius of  $2 \text{ k}\lambda$  to the  $u$ - $v$  data, yielding a resolution of  $73''.1 \times 45''.5$  at P.A. =  $2^\circ$ . Contours are multiples of  $120 \mu\text{Jy beam}^{-1}$  ( $1.5 \sigma$ ), and the rms is  $80 \mu\text{Jy beam}^{-1}$ . [See the electronic edition of the Journal for a color version of this figure.]

centroid and the point-source positions and fluxes at their best-fit values and calculate the  $\chi^2$  statistic over a large range of  $\theta_c$ ,  $\beta$ , and  $\Delta T_0$ -values. For a given electron temperature  $T_e$ , the  $\beta$  model then yields the gas density profile  $n_e(\theta)$  at each  $(\theta_c, \beta, \Delta T_0)$  point. The gas mass and the total cluster mass can be calculated directly from  $n_e(\theta)$ , assuming that the intracluster medium is in hydrostatic equilibrium with the cluster potential. Following the methods outlined in Grego et al. (2001), we calculate the gas mass, total mass, and gas mass fraction over the  $(\theta_c, \beta, \Delta T_0)$ -grid and report the total mass values for which the  $\chi^2$  statistic is within the 68% confidence interval ( $\Delta\chi^2 = 1$ ). We evaluate these quantities at an angular radius of  $65''$ , where our observational techniques best constrain the cluster gas mass fraction (Grego et al. 2001). For clusters at  $z \sim 0.8$ , this angular radius corresponds to a physical radius of  $340 h_{100}^{-1} \text{ kpc}$  for an  $\Omega_M = 0.3$ ,  $\Omega_\Lambda = 0.7$  cosmology.

We can estimate the electron temperature directly from the SZE data by finding the range of  $T_e$ -values that yield a cluster gas mass fraction,  $f_g$ , consistent with the mean value measured by Grego et al. (2001) for a sample of 18 clusters. To determine the gas mass fraction at  $r_{500}$  for CL J1226.9+3332, CL J0152.7–1357, and MS 1054.4–0321, we scale the gas mass fractions measured at  $65''$  to  $r_{500}$  using relations derived from numerical simulations (Evrard, Metzler, & Navarro 1996; Evrard 1997), as discussed in Grego et al. (2001). This calculation is repeated for a number of different temperatures ranging from 4–18 keV, and we report the  $T_e^{\text{SZ}}$ -values that are consistent with a mean value of  $f_g(r_{500}) = 0.081$  (Grego et al. 2001) within the sample standard deviation of 0.04. To determine how sensitive the derived mass and temperature are to the adopted cosmological model, we repeated the calculations above for  $\Omega_M = 0.3$  and  $\Omega_\Lambda = 0.0$ . The SZE derived mass and temperature are reduced

by  $\sim 3\%$ ; the overall change is small because the effects of decreased distance are offset by a compensating change in the  $f_g$  scaling relation.

### 3. RESULTS AND CONCLUSIONS

Synthesized images of the SZE toward CL J1226.9+3332, CL J0152.7–1357, and MS 1054.4–0321 are shown in Figure 1. The SZE decrement is detected with high significance in all of these distant clusters, and the locations of the SZE and X-ray centroids are consistent (Tables 1 and 3). Using the SZE data and the analysis techniques described in § 2.2, we determine the temperature and mass of each cluster (Table 3). X-ray temperature measurements for CL J0152.7–1357 (Della Ceca et al. 2000) and MS 1054.4–0321 (Donahue et al. 1998) are also shown in Table 3, and we find that these X-ray temperature measurements are consistent with the values inferred from the SZE data within the stated uncertainties.

From the SZE data, we infer a total mass of  $\geq 2 \times 10^{14} h_{100}^{-1} M_\odot$  within a radius of  $65''$  ( $340 h_{100}^{-1} \text{ kpc}$ ) for each of the clusters shown in Table 3. These mass calculations can be checked against values derived from gravitational lensing, X-ray, and optical observations of MS 1054.4–0321 (Donahue et al. 1998; van Dokkum et al. 2000; Hoekstra et al. 2000). Hoekstra et al. (2000) infer a total mass of  $(5.4 \pm 0.6) \times 10^{14} h_{100}^{-1} M_\odot$  within an aperture of radius  $94''$ . To compare the SZE and gravitational lensing results, we calculate the gas mass within a  $94''$  radius aperture, for an electron temperature of 10.4 keV:  $M_{\text{gas}}^{\text{SZ}}(<94'') = (3.7 \pm 0.6) \times 10^{13} h_{100}^{-2} M_\odot$ , where the uncertainties reflect the statistical 68% confidence interval in modeling the SZE data. The total mass is estimated by scaling this value by the mean gas mass fraction (§ 2.2), from which

TABLE 3  
CLUSTER PROPERTIES DERIVED FROM SZE MEASUREMENTS

CLUSTER NAME	CLUSTER POSITION <sup>a</sup>		$T_e^{\text{X}}$ (keV)	$T_e^{\text{SZ}}$ (keV)	$M_{\text{total}}(<65'')$ ( $\times 10^{14} h^{-1} M_\odot$ )
	$\Delta\alpha$ (arcsec)	$\Delta\delta$ (arcsec)			
CL J1226.9+3332 .....	0.2	12.3	...	$9.8_{-1.9}^{+4.7}$	$2.7 \pm 0.5$
CL J0152.7–1357 .....	–1.8	–9.2	$6.5_{-1.2}^{+1.7}$	$8.7_{-1.8}^{+4.1}$	$2.1 \pm 0.7$
MS 1054.4–0321 .....	–7.8	–5.3	$12.3_{-2.2}^{+3.1}$	$10.4_{-2.0}^{+3.0}$	$2.3 \pm 0.3$

<sup>a</sup> Offsets from radio pointing center (Table 1).

we obtain an SZE estimate of the total mass within a  $94''$  radius:  $M_{\text{total}}^{\text{SZ}}(< 94'') = (4.6 \pm 0.8) \times 10^{14} h_{100}^{-1} M_{\odot}$ . We find that the total mass estimated from the SZE data is consistent with the lensing measurements of Hoekstra et al. (2000).

Based on the SZE data, we conclude that the newly discovered clusters CI J1226.9+3332 and CI J0152.7–1357 are highly massive, with a total mass of  $M_{\text{total}}^{\text{SZ}} \geq 2 \times 10^{14} h_{100}^{-1} M_{\odot}$  within a radius of  $65''$  ( $340 h_{100}^{-1}$  kpc). These values are comparable to the mass inferred from SZE imaging of the  $z = 0.83$  cluster MS 1054.4–0321, which has been confirmed by X-ray, optical, and gravitational lensing studies. These results demonstrate the ability to determine cluster temperatures and masses from SZE data without access to X-ray data and illustrate the power of using deep SZE surveys to probe the distant universe. More precise measures of the temperature and mass of CI J1226.9+3332 and CI J0152.7–1357 will be possible with deep X-ray imaging and spectroscopy, which will be obtained within the coming year by the *Chandra* and *XMM* X-ray observatories; with these data in hand, the SZE measurements can be used to measure the distance to each cluster (Reese et al. 2000) and to further constrain their density, mass, and gas mass fraction (Grego et al. 2001). Additional, independent constraints on the mass distribution in CI

J1226.9+3332 will be obtained from a weak-lensing analysis of wide-field imaging data taken with the *Hubble Space Telescope* and the Subaru 8.3 m telescope.

We dedicate this Letter to the memory of our friend and colleague Mark Warnock, who freely gave of his expertise and time and made great contributions to the interferometric SZE imaging experiment. We also thank Cheryl Alexander, Rick Forster, Steve Padin, Dick Plambeck, Steve Scott, David Woody, and the staff of the BIMA and OVRO observatories for their outstanding support and Laurence Jones, Eric Perlman, and Caleb Scharf for providing data on CI J1226.9+3332 prior to publication. E. D. R. gratefully acknowledges support from NASA GSRP Fellowship NGT5-50173. This work is supported by NASA LTSA grants NAG5-7986 (J. E. C., M. J., W. L. H.) and NAG5-8253 (H. E.). Radio astronomy at the BIMA millimeter array is supported by NSF grant AST 96-13998. The OVRO millimeter array is supported by NSF grant AST 96-13717. The funds for the additional hardware for the SZE experiment were from a NASA CDDF grant, an NSF-YI Award, and the David and Lucile Packard Foundation.

#### REFERENCES

- Bahcall, N. A., & Cen, R. 1992, *ApJ*, 398, L81  
 Bahcall, N. A., & Fan, X. 1998, *ApJ*, 504, 1  
 Birkinshaw, M. 1999, *Phys. Rep.*, 310, 97  
 Carlstrom, J. E., Joy, M., & Grego, L. 1996, *ApJ*, 456, L75  
 Cavaliere, A., & Fusco-Femiano, R. 1976, *A&A*, 49, 137  
 ———. 1981, *A&A*, 100, 194  
 Della Ceca, R., Scaramella, R., Gioia, I. M., Rosati, P., Fiore, F., & Squires, G. 2000, *A&A*, 353, 498  
 Donahue, M., Voit, G. M., Gioia, I., Lupino, G., Hughes, J. P., & Stocke, J. T. 1998, *ApJ*, 502, 550  
 Ebeling, H., Jones, L. R., Fairley, B. W., Perlman, E., Scharf, C., & Horner, D. 2001, *ApJ*, 548, L23  
 Ebeling, H., et al. 2000, *ApJ*, 534, 133  
 Eke, V. R., Cole, S., Frenk, C. S., & Henry, J. P. 1998, *MNRAS*, 298, 1145  
 Evrard, A. E. 1997, *MNRAS*, 292, 289  
 Evrard, A. E., Metzler, C. A., & Navarro, J. F. 1996, *ApJ*, 469, 494  
 Gioia, I. M., & Luppino, G. A. 1994, *ApJS*, 94, 583  
 Grego, L., Carlstrom, J. E., Joy, M. K., Reese, E. D., Holder, G. P., Patel, S., Cooray, A. R., & Holzzapfel, W. L. 2000, *ApJ*, 539, 39  
 Grego, L., Carlstrom, J. E., Reese, E. D., Holder, G. P., Holzzapfel, W. L., Joy, M. K., Mohr, J., & Patel, S. 2001, *ApJ*, in press  
 Haiman, Z., Mohr, J. J., & Holder, G. P. 2001, *ApJ*, in press  
 Hoekstra, H., Franx, M., & Kuijken, K. 2000, *ApJ*, 532, 88  
 Luppino, G. A., & Gioia, I. M. 1995, *ApJ*, 445, L77  
 Neumann, D. M., & Arnaud, M. 2000, *ApJ*, 542, 35  
 Oukbir, J., & Blanchard, A. 1997, *A&A*, 317, 1  
 Pearson, T. J., Shepherd, M. C., Taylor, G. B., & Myers, S. T. 1994, *BAAS*, 185, 08.08  
 Pospieszalski, M. W., Lakatos, W. J., Nguyen, L. D., Lui, M., Liu, T., Le, M., Thompson, M. A., & Delaney, M. J. 1995, *IEEE MTT-S Digest*, 3, 1121  
 Reese, E. D., et al. 2000, *ApJ*, 533, 38  
 Romer, A. K., et al. 2000, *ApJS*, 126, 209  
 Rudy, D. J. 1987, Ph.D. thesis, Caltech  
 Sault, R. J., Teuben, P. J., & Wright, M. C. H. 1995, in *ASP Conf. Ser. 77, Astronomical Data Analysis Software and Systems IV*, ed. R. A. Shaw, H. E. Payne, & J. J. E. Hayes (San Francisco: ASP), 433  
 Scharf, C. A., Jones, L. R., Ebeling, H., Perlman, E., Malkan, M., & Wegner, G. 1997, *ApJ*, 477, 79  
 Scoville, N. Z., Carlstrom, J. E., Chandler, C. J., Phillips, J. A., Scott, S. L., Tilanus, R. P. J., & Wang, Z. 1993, *PASP*, 105, 1482  
 Sunyaev, R. A., & Zeldovich, Y. B. 1972, *Comments Astrophys. Space Phys.*, 4, 173  
 van Dokkum, P. G., Franx, M., Fabricant, D., Illingworth, G. D., & Kelson, D. D. 2000, *ApJ*, 541, 95  
 Viana, P. T. P., & Liddle, A. R. 1996, *MNRAS*, 281, 323