

**Table 9. Estimated numbers of persons living with HIV infection (not AIDS), by race/ethnicity, sex, transmission category, and age category, 2001–2004—35 areas with confidential name-based HIV infection reporting**

Transmission category	2001	2002	2003	2004
<b>White, not Hispanic</b>				
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	38,419	41,283	43,726	46,608
Injection drug use	4,591	4,752	4,811	4,896
Male-to-male sexual contact and injection drug use	4,150	4,255	4,323	4,450
High-risk heterosexual contact <sup>a</sup>	2,393	2,571	2,707	2,820
Other <sup>b</sup>	459	458	454	450
Subtotal	50,011	53,320	56,022	59,224
<b>Female adult or adolescent</b>				
Injection drug use	3,636	3,773	3,883	3,955
High-risk heterosexual contact <sup>a</sup>	7,042	7,585	8,028	8,420
Other <sup>b</sup>	153	159	156	154
Subtotal	10,832	11,516	12,067	12,529
<b>Total adults and adolescents</b>	<b>60,843</b>	<b>64,836</b>	<b>68,089</b>	<b>71,753</b>
<b>Child (&lt;13 yrs)</b>				
Perinatal	434	440	459	458
Other <sup>c</sup>	159	146	138	133
Subtotal	593	586	597	591
<b>Total</b>	<b>61,435</b>	<b>65,422</b>	<b>68,686</b>	<b>72,344</b>
<b>Black, not Hispanic</b>				
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	24,764	26,934	28,772	30,842
Injection drug use	11,924	12,211	12,221	12,207
Male-to-male sexual contact and injection drug use	3,495	3,519	3,481	3,463
High-risk heterosexual contact <sup>a</sup>	11,853	12,876	13,656	14,337
Other <sup>b</sup>	397	398	403	402
Subtotal	52,433	55,937	58,533	61,251
<b>Female adult or adolescent</b>				
Injection drug use	8,003	8,205	8,241	8,195
High-risk heterosexual contact <sup>a</sup>	26,012	28,508	30,266	31,875
Other <sup>b</sup>	488	516	545	553
Subtotal	34,503	37,229	39,052	40,623
<b>Total adults and adolescents</b>	<b>86,935</b>	<b>93,166</b>	<b>97,585</b>	<b>101,874</b>
<b>Child (&lt;13 yrs)</b>				
Perinatal	2,139	2,224	2,286	2,323
Other <sup>c</sup>	335	349	349	363
Subtotal	2,473	2,573	2,635	2,686
<b>Total</b>	<b>89,409</b>	<b>95,739</b>	<b>100,221</b>	<b>104,560</b>

**Table 9. Estimated numbers of persons living with HIV infections (not AIDS), by race/ethnicity, sex, transmission category, and age category, 2001–2004—35 areas with confidential name-based HIV infection reporting (cont)**

Transmission category	2001	2002	2003	2004
<b>Hispanic</b>				
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	9,440	10,848	12,072	13,425
Injection drug use	4,519	4,626	4,678	4,587
Male-to-male sexual contact and injection drug use	993	1,042	1,089	1,074
High-risk heterosexual contact <sup>a</sup>	2,531	2,847	3,126	3,356
Other <sup>b</sup>	92	96	100	96
Subtotal	17,574	19,459	21,065	22,537
<b>Female adult or adolescent</b>				
Injection drug use	1,997	2,066	2,070	2,007
High-risk heterosexual contact <sup>a</sup>	4,853	5,389	5,725	5,965
Other <sup>b</sup>	99	111	121	120
Subtotal	6,949	7,567	7,916	8,092
<b>Total adults and adolescents</b>	<b>24,524</b>	<b>27,026</b>	<b>28,981</b>	<b>30,629</b>
<b>Child (&lt;13 yrs)</b>				
Perinatal	733	757	768	788
Other <sup>c</sup>	121	118	121	123
Subtotal	854	875	889	911
<b>Total</b>	<b>25,378</b>	<b>27,901</b>	<b>29,870</b>	<b>31,540</b>
<b>Asian/Pacific Islander</b>				
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	432	507	615	698
Injection drug use	76	89	92	93
Male-to-male sexual contact and injection drug use	21	24	30	36
High-risk heterosexual contact <sup>a</sup>	82	105	121	134
Other <sup>b</sup>	24	26	26	23
Subtotal	634	750	885	984
<b>Female adult or adolescent</b>				
Injection drug use	32	37	45	50
High-risk heterosexual contact <sup>a</sup>	165	197	223	259
Other <sup>b</sup>	12	13	12	13
Subtotal	209	247	279	322
<b>Total adults and adolescents</b>	<b>843</b>	<b>998</b>	<b>1,164</b>	<b>1,306</b>
<b>Child (&lt;13 yrs)</b>				
Perinatal	13	16	18	18
Other <sup>c</sup>	4	4	4	4
Subtotal	17	20	22	22
<b>Total</b>	<b>860</b>	<b>1,018</b>	<b>1,186</b>	<b>1,328</b>

**Table 9. Estimated numbers of persons living with HIV infection (not AIDS), by race/ethnicity, sex, transmission category, and age category, 2001–2004—35 areas with confidential name-based HIV infection reporting (cont)**

Transmission category	2001	2002	2003	2004
<b>American Indian/Alaska Native</b>				
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	386	413	424	464
Injection drug use	94	94	96	95
Male-to-male sexual contact and injection drug use	91	91	95	89
High-risk heterosexual contact <sup>a</sup>	47	56	62	73
Other <sup>b</sup>	2	2	1	1
Subtotal	620	656	679	722
<b>Female adult or adolescent</b>				
Injection drug use	80	90	96	96
High-risk heterosexual contact <sup>a</sup>	146	167	186	201
Other <sup>b</sup>	2	2	3	2
Subtotal	228	259	286	299
<b>Total adults and adolescents</b>	<b>848</b>	<b>915</b>	<b>965</b>	<b>1,021</b>
<b>Child (&lt;13 yrs)</b>				
Perinatal	8	10	10	10
Other <sup>c</sup>	2	1	1	1
Subtotal	10	11	11	11
<b>Total</b>	<b>858</b>	<b>926</b>	<b>976</b>	<b>1,032</b>
<b>Total cases<sup>d</sup></b>	<b>179,678</b>	<b>192,801</b>	<b>202,722</b>	<b>212,572</b>

*Note.* These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

Since 2000, the following 35 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, Guam, and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting for new diagnoses only.

<sup>a</sup> Includes heterosexual contact with a person who is HIV infected, a male who engages in male-to-male sex, a person who injects drugs, has hemophilia or coagulation disorder, or is a transplant or transfusion recipient.

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>c</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

<sup>d</sup> Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.