Agricultural Marketing Service, USDA

- (1) Callous growth cracks (other than callous apex-end cracks) which, singly or in the aggregate on a prune, exceed % inch in length; and callous apex-end cracks which, singly or in the aggregate on a prune, exceed ½ inch in length.
- (2) Splits or skin breaks not having callous edges when the flesh is mashed out beyond the protecting skin so as to affect materially the normal appearance of the prune;
- (3) Any cracks, splits, or skin breaks open to the pit;
- (4) Any skin damage from multiple short skin breaks giving a very "rough" appearance to the prune such as may result from over-dipping, rain damage, processing, or other causes which in Type I dried prunes affect materially the appearance or edibility of the prune, or which in other types or other mixtures affect markedly the appearance or edibility of the prune.
- (e) Fermentation. "Fermentation" means that the prune is fermented as evidenced by a distinct sour taste or odor or by darkening in color characteristic of fermentation or souring.
 - (f) Scars. "Scars" mean:
- (1) Tough or thick scab which, singly or in combination on a prune, exceeds the area of a circle % inch in diameter such as may result from thrip injury, mildew, leaf chafing, limb rubs, or other means; or
- (2) Scab which is not tough or thick and which, singly or in combination on a prune, exceeds the area of a circle ³/₄ inch in diameter and which is inconsequential but unsightly though practically blending in color with the skin on the portion of the prune not affected.
- (g) Heat damage. "Heat damage" means burning or scorching from the sun or in dehydration so as to damage

- materially the skin or flesh, or both, of the prune.
- (h) Insect injury. "Insect injury" means healed or unhealed surface blemishes and healed or unhealed blemishes extending into the flesh which affect materially the appearance, edibility, or keeping quality of the prune but which do not possess evidence of insect infestation.
- (i) Other means. "Other means" includes damage by any injury or defect or group of defects not specifically mentioned in this section which materially affects the appearance, edibility, or keeping quality of the fruit, but "other means" does not include defects of a nature such as defined in paragraph (j), (k), (l), (m), or (n) of this section.
- (j) *Mold.* "Mold" means a characteristic fungus growth as evidenced by a moldy or smutty condition and which, singly or in the aggregate on a prune, is equal to or exceeds the area of a circle %6 inch in diameter.
- (k) *Dirt*. "Dirt" means the presence of any quantity of such substance, whether imbedded or adhering to the prune, which gives the prune a dirty, smudgy appearance and which may not be removed readily by washing.
- (1) Foreign material. "Foreign material" means leaves, twigs, pieces of wood, and similar extraneous materials which are objectionable.
- (m) Insect infestation. "Insect infestation" means the presence of dead insects, insect fragments, or insect remains. (No live insects are permitted).
- (n) Decay. "Decay" means a state of decomposition, wholly or in part, of the prune.
- [21 FR 8177, Oct. 25, 1956, as amended at 30 FR 11596, Sept. 10, 1965. Redesignated at 42 FR 32514, June 27, 1977, and further redesignated at 46 FR 63203, Dec. 31, 1981]

WORK SHEET

$\S 52.3188$ Work sheet for dried prunes.

Size and kind of container.
Container mark or identification.
Label or brand.
Varietal type.
Style.
Size:
Count per pound (Average).
Uniformity.

§ 52.3751

) Extra large) Large.) Medium.) Small. Moisture content percent; Uniformity. Varietal characteristics: (). Similar. () Dissimilar.

Defects and summary of allowances 1	Grade A max- imum	Grade B max- imum	Grade C max- imum	Substandard max- imum
Total of all defects, including off-color	10 percent	15 percent		No limit except as indicated below.
Total of all defects, including off-color and poor texture.			20 percent.	
Poor texture, end cracks, skin or flesh damage, ³ fermentation, scars, heat damage, insect injury, other means, mold, dirt, foreign	But no more than 6 percent.	But no more than 8 percent.		
material, insect infestation, decay. End cracks, ² skin or flesh damage, ³ fermenta- tion, scars, heat damage, insect injury, other means, mold, dirt, foreign material, in-			10 percent ² .	
sect infestation, decay. Skin or flesh damage, 3 fermentation, scars, heat damage, insect injury, other means, mold, dirt, foreign material, insect infestation, decay.			But no more than 8 percent.	
Mold, dirt, foreign material, insect infestation, decay.	3 percent	4 percent	5 percent	5 percent.
Decay	But no more than 1 percent.	But no more than 1 percent.	But no more than 1 percent.	But no more than 1 percent.

Total

U.S. Grade (including all factors)

[21 FR 8177, Oct. 25, 1956, as amended at 30 FR 11596, Sept. 10, 1965. Redesignated at 42 FR 32514, June 27, 1977 and at 46 FR 63203, Dec. 31, 1981]

Subpart—United States Standards for Grades of Canned Ripe Olives 1

SOURCE: 42 FR 38585, July 29, 1977, unless otherwise noted. Redesignated at 46 FR 63203, Dec. 31, 1981.

PRODUCT DESCRIPTION, TYPES, STYLES, AND GRADES

§52.3751 Product description.

Canned ripe olives are prepared from properly matured olives which have first been properly treated to remove characteristic bitterness; packed in a solution of sodium chloride, with or without spices, and are sufficiently processed by heat in hermetically sealed containers. Canned olives which are not oxidized in processing and which possess a tan to light bronze color indicative of preparation from olives of advanced maturity and commonly referred to as "tree-ripened" or "home-cured" are not covered by the standards in this subpart.

§ 52.3752 Types of canned ripe olives.

Canned ripe olives are processed as two distinct types. Unless a specific type is stated in this subpart, "canned ripe olives" refers to olives of either "ripe-type" or "green-ripe type."

- (a) Ripe type. "Ripe type" olives are those which have been treated and oxidized in processing to produce a typical dark brown to black color.
- (b) Green-ripe type. "Green-ripe type" olives are those which have not been oxidized in processing; which range in color from yellow-green; green-yellow or other greenish casts; and which may be mottled.

Percentages of defects are "by weight."
 Except that each 1 percent of end cracks to, and including 8 percent, by weight, shall be considered as ½ percent damaged by end cracks; and any additional end cracks shall be calculated as true percentage, by weight.
 Allowances for "skin or flesh damage" apply only to "Whole Unpitted" style.

¹Compliance with the provisions of these standards shall not excuse failure to comply with the provisions of the Federal Food, Drug and Cosmetic Act, or with applicable State laws and regulations.