

# SHORELINE CHANGES IN THE CAMINADA-MOREAU HEADLAND AND GRAND ISLE - 1887 TO 1996 LAFOURCHE AND JEFFERSON PARISHES, LOUISIANA

Shea Penland<sup>1</sup>, Chris Zganjar<sup>1</sup>, Karen A. Westphal<sup>2</sup>, Paul Connor<sup>1</sup>  
Jeff List<sup>3</sup> and S. Jeffress Williams<sup>3</sup>

## INTRODUCTION

The U.S. Geological Survey (USGS), in cooperation with the Coastal Research Laboratory in the Department of Geology and Geophysics at the University of New Orleans (UNO) and the Center for Coastal Energy and Environmental Resources at Louisiana State University (LSU), is investigating the processes of coastal erosion in Louisiana (Sallenger and others, 1987; Sallenger and Williams 1989; Penland and others, 1992). Building on the USGS Louisiana Barrier Island Study (Williams and others, 1992), this USGS Open-File Report depicts shoreline changes between 1887 and 1996, which provides an 8.9-year update of McBride and others (1992). In order to quantify shoreline changes since January 21, 1988, new vertical aerial mapping photography was acquired on December 9, 1996. The methods and transects used by McBride and others (1992) were used to insure data compatibility of the new measurements and analysis (Plate 7). Tables 1 and 2 present the transect measurements of shoreline change for the Caminada-Moreau Headland and Grand Isle. For gulfside change measurements, a negative (-) sign signifies landward movement or erosion and a positive (+) sign signifies a seaward movement or progradation. For bayside change measurements, a negative (-) sign signifies a seaward movement or erosion and a positive (+) sign signifies a landward movement or accretion.

The Caminada-Moreau Headland and Grand Isle are located approximately 90km south of New Orleans (Figure 1) and is bordered by Raccoon Pass in the west and Barataria Pass in the east. Since 1887 the Caminada-Moreau Headland has experienced some of the highest rates of shoreline movement along the Louisiana coastline. Conversely, Grand Isle, east of Caminada-Moreau and separated by the Caminada Pass, has experienced stationary or accretionary rates of shoreline movement over most of its coastal shoreline.

The Caminada-Moreau Headland primarily consists of cohesive deltaic sediment and a large, sandy beach ridge plain with no back-barrier lagoon or bay. Bell Pass, Pass Fouchon, and Bayou Moreau separate the central headland area. With the exception of Bayou Moreau, Bell Pass and Pass Fouchon have seen extensive man-made influences in the form of canal widening and dredging. By 1956, the Bell Pass Jetties had begun to interrupt the large quantities of erosional downdrift sediment from the central headland area causing the magnitude of downdrift offset to increase west of Bell Pass. As a result of this highly erosional shoreline, Bayou Moreau now intersects the shoreline in three different locations. The interior marsh of the Caminada-Moreau Headland is also

---

<sup>1</sup> Department of Geology and Geophysics, University of New Orleans

<sup>2</sup> Center for Coastal, Energy and Environmental Resources, Louisiana State University

<sup>3</sup> U.S. Geological Survey, Coastal and Marine Geology Program

experiencing high rates of loss primarily as a result of numerous dredge canals and levees.

To the east of Caminada Pass lies one of Louisiana's most unique barrier islands. Grand Isle is one of the few barrier islands, which has sustained an accretionary growth since the early 20<sup>th</sup> century. Because of this morphological stability, it is the only barrier island in Louisiana commercially and residentially developed (Meyer-Arendt, 1987). For 101 years, the gulf shoreline has experienced retreat along its western end while remaining relatively stationary at its midsections and accreting seaward on its eastern end. These trends show that Grand Isle is slowly rotating clockwise around a stable midpoint (McBride and others, 1992). The accretion on the eastern end of Grand Isle can be attributed to Louisiana's deepest tidal inlet (> 40m in 1989) bordering its eastern end, known as Barataria Pass. The Barataria Pass tidal inlet system is a large sediment sink storing most of its sand as a large ebb-tidal delta. Shoreline advance at the eastern end of Grand Isle is directly related to this ebb-tidal delta (Shamban, 1982).

## **SHORELINE MOVEMENT**

Magnitude and rate of change for the Caminada-Moreau Headland and Grand Isle were derived from 91 shore-normal transects along the gulf and bay shorelines (Transects Map, Tables 1&2). Comparisons of shoreline positions are made for the periods 1887 vs. 1988, 1988 vs. 1996, and 1887 vs. 1996. The overlay maps illustrate land loss and quantitative changes for the Caminada-Moreau Headland and Grand Isle. Because of the unique nature of this region, this study divided the above mentioned barrier system into two sections: Caminada-Moreau Headland, between Raccoon Pass in the west and Caminada Pass in the east, and Grand Isle, between Caminada Pass in the west and Barataria Pass in the east.

## **GULFSIDE SHORELINE CHANGES**

In terms of the long-term gulfside shoreline change history for the 109 year period between 1887 and 1996, the Caminada-Moreau Headland shoreline transects measured between -150m and -2068m (Table 1, 1887-1996, transects 1-38). The average shoreline change was determined to be 1373.7m, which yields an average yearly rate of -12.6m/yr (Table 3). During the same period, Grand Isle's gulfside transects measured between 674m and -262m (Table 1, 1887-1996, transects 39-62) resulting in an average shoreline change of 128.8m. This positive, or gulfward shoreline change measurement reveals a long-term yearly accretion rate of 1.2m/yr (Table 3), which predominately occurs along the eastern two thirds of Grand Isle's gulf shoreline.

For the short-term shoreline change analysis of the 8.9-year period between 1988 and 1996, the Caminada-Moreau Headland gulfside transects ranged from 148m to -100m (Table 1, 1988-1996, transects 1-38) for an average shoreline change of -28.4m or -3.2 m/yr (Table 3). The short-term yearly rate of -3.2m/yr. could be as much as 4.0m/yr. if it were not for the influence of highly variable spit movement at the west end of the Caminada-Moreau Headland (Caminada spit). In terms of the short-term shoreline

change history, Grand Isle's gulfside transects were measured from 149m to -69m (Table 1, 1988-1996, transects 39-62) resulting in an average shoreline movement of 36.8m or 4.1m/yr (Table 3).

Previous work by McBride and others (1992) documents long-term shoreline change between years 1887 and 1988 (101 years) and short-term shoreline change between 1978 and 1988 (10 years). For the Caminada-Moreau Headland, the earlier analysis reported a long-term gulf shoreline movement rate of 13.3m/yr (Table 3) and an average short-term shoreline movement rate of 13.6m/yr (Table 3). The Grand Isle gulf shoreline experienced an average long-term advance of 0.9m/yr (Table 3) and a short-term gulfward movement of 5.2m/yr (Table 3).

The comparison between the new long-term shoreline change rate and the McBride and others (1992) long-term change rate shows only a slight difference. The long-term gulf shoreline rate changed from -133m/yr (1887-1988) to -12.6m/yr (1887-1996) (Table 3), for the Caminada-Moreau Headland, indicating a reduction in erosion by 0.7m/yr. In both studies Grand Isle's rate of long-term shoreline movement has remained nearly the same. The long-term shoreline movement increased toward the Gulf of Mexico by 0.3m/yr, from 0.9m/yr. (1887-1988) to 1.2m/yr. (1887-1996) (Table 3). The Caminada- Moreau Headland experienced a dramatic shift in short-term gulfside erosion rates between the McBride and others (1992) study and this update. For the years between 1978 and 1988, the Caminada-Moreau Headland experienced a gulfside rate of change of -13.6m/yr (Table 3). Compared to this study, which reported a short-term gulfside change rate of -3.2m/yr. (Table 3), a reduction of shoreline erosion by 10.4m/yr. was observed.

## **BAYSIDE SHORELINE CHANGES**

The Caminada-Moreau Headland consists predominately of cohesive deltaic sediment and a large, sandy beach ridge plain with no back barrier lagoon or bay. Therefore, the bay shoreline change measurements pertain only to the Caminada spit located at the eastern end of the Caminada-Moreau Headland.

In terms of the long-term bay shoreline change history for the 109-year period between 1887 and 1996, the Caminada spit shoreline transects measured between 203m and 594m (Table 2, 1887-1996, transects 33- 38). The average change was determined to be 394m or 3.6m/yr (Table 3). Grand Isle experienced long-term bay shoreline movement between 326m and -333m (Table 2, 1887-1996, transects 39-62) yielding an average measurement of -108.1m or -1.0m/yr (Table 4).

For the 8.9-year period between 1988 and 1996, Caminada spit underwent very little change in its short-term bay shoreline movement. The Caminada spit bayside movement ranged from 12m to -11m (Table 2, 1988-1996, transects 33-38) for an average of 2.6m at a rate of 0.0m/yr (Table 4). Grand Isle also saw modest movement of its bay shoreline between the years 1988 and 1996. Bayside movement ranged from -47m to 42m, providing an average of -3.6m or -0.6m/yr (Table 4).

McBride and others (1992) studied long-term change rates between years 1884 and 1988 (104 years) and short-term change rates between years 1978 and 1988 (10 years). In this earlier study, long-term bay shoreline change rates for Caminada spit were calculated as 4.1m/yr (Table 4), and short-term shoreline changes were -1.8m/yr. Grand Isle's long-term and short-term bayside shoreline change rates were determined to be -1.0m/yr. and -3.2m/yr. respectively (Table 4).

Drawing on comparisons between the bay shoreline movement rate of the McBride and others (1992) analysis and this study, long-term change rates have remained virtually unchanged. McBride and others (1992) reported a long-term change rate of Caminada spit as 4.1m/yr. (Table 4). This recent study calculated a rate of 3.6m/yr (Table 4), indicating a 0.5m/yr. reduction in landward migration. Grand Isle's long-term bay shoreline change rate was identical in both the McBride and others (1992) and this study, at -1.0m/yr (Table 4).

The short-term bay movement rates revealed slightly more variability between this study and the previous one. McBride and others (1992) observed Caminada spit to have a change rate of -1.8m/yr (Table 4). The short-term rate in this new study is 0.0m/yr (Table 4) indicating the bay shoreline of Caminada spit is entering a period of stability or possible landward migration.

Similar to Caminada spit, the bay shoreline of Grand Isle is also entering a period of transition. Based on the comparison between the short-term movement rates of 3.2m/yr (McBride and others, 1992) (Table 4) and -0.6m/yr (Table 4) calculated in the new study, one could observe an erosional shift of 3.8m/yr. on the bay shoreline of Grand Isle.

## **AREA CHANGES**

Between 1887 and 1996, the total area of Grand Isle has decreased by 170 acres. What is not seen in this long-term comparison is since 1956 the area of Grand Isle has steadily increased. Based on this study's new short-term (1988-1996) area rate calculations, Grand Isle could achieve its 1887 area in a little over sixteen years.

Between 1887 and 1996, Grand Isle's area decreased from 2617 acres to 2447 acres (Table 5). This represents a 6% loss of island area at a rate of -1.6 acres/yr (Table 6). This long-term rate of loss, acting on Grand Isle's area in 1996, forecasts a disappearance date of 3525 (Table 6). Previously, in McBride and others (1992), Grand Isle decreased in area at a rate of 2.4 acres/yr. between 1887 and 1988 suggesting a long-term disappearance date of 2968 (Table 6). This seemingly small rate decrease in area loss will extend the life of Grand Isle by 557 years. For the short-term area analysis, between the years 1988 and 1996, Grand Isle's area increased from 2372 acres to 2447 acres (Table 5). This indicates a 3% gain at a rate of +8.4 acres/yr (Table 6). McBride and others (1992) reported a 3% gain in island area between the years 1978 and 1988, indicating a net short-term gain of 5.9 acres/yr. Comparing the two studies, the recently observed short-term (1988-1996) rate shows an increase of 2.5 acres/yr in the growth of Grand Isle.

## ACKNOWLEDGEMENTS

This analysis of Louisiana's barrier islands was funded by the U.S. Geological Survey Coastal and Marine Geology Program.

## DISCLAIMER


This poster is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards (and stratigraphic nomenclature). Any use of trade names is for descriptive purposes only and does not imply endorsement by the USGS.

## BIBLIOGRAPHY


- McBride, R.A., Penland, S., Hiland, M., Williams, S.J., Westphal, K.A., Jaffe, B., and Sallenger, A.H., Jr., 1992. Louisiana barrier shoreline change analysis- 1853 to 1989: methodology, database, and results. *in*: Williams, S.J., Penland, S., and Sallenger, A.H., (editors), *Atlas of Shoreline Changes in Louisiana from 1853 to 1989*. US Geological Survey, Reston, Virginia.
- Penland, S., Williams, S.J., Davis, D.W., Sallenger, A.H., Jr., and Groat, C.G., 1992. Barrier island erosion and wetland loss in Louisiana, *in* Williams, S.J., Penland, S., and Sallenger, A.H., Jr., eds., *Louisiana Barrier island erosion study--atlas of barrier shoreline changes in Louisiana from 1853 to 1989*: U.S. Geological Survey Miscellaneous Investigations Series I-2150\_A, p.2-7.
- Sallenger, A.H., Jr, Penland, S., Williams, S.J., and Suter, J.R., 1987. Louisiana barrier island erosion study: Coastal Sediments '87, American Society of Civil Engineers, p. 1503-1516.
- Sallenger, A.H., Jr., and Williams, S.J., 1989. U.S. Geological Survey studies of Louisiana barrier island erosion and wetlands loss: An interim report on status and results: U.S. Geological Survey Open- File Report 89-372, 17 p.
- Shamban, A., 1982, Coastal processes and geomorphology, Barataria Pass, Louisiana: Baton Rouge, Louisiana State University unpublished M.S. theses, 121p
- Mayer-Arendt, K. J., 1985, The Grand Isle, Louisiana resort cycle: *Annals of Tourism Research*, V. 12, p. 449-465
- Williams, S.J., Penland, S., and Sallenger, A.H., Jr., eds., 1992, Louisiana Barrier island erosion study--atlas of barrier shoreline changes in Louisiana from 1853 to 1989: U.S. Geological Survey Miscellaneous Investigations Series I-2150\_A, 103 p.




## FIGURES

**Figure 1:** The Caminada-Moreau Headland and Grand Isle located 90 km south of New Orleans



# MAPS







## TABLES

**Table 1**

**Table 2**

Caminada-Moreau and Grand Isle gulfside magnitude of change (meters)						Caminada-Moreau and Grand Isle bayside magnitude of change (meters)					
1887-1988		1988-1996		1887-1996		1887-1988		1988-1996		1887-1996	
1	-1969	1	-3	1	-1974	1	n.a.	1	n.a.	1	n.a.
2	-1962	2	-4	2	-1959	2	n.a.	2	n.a.	2	n.a.
3	-1912	3	-17	3	-1932	3	n.a.	3	n.a.	3	n.a.
4	-1837	4	-30	4	-1895	4	n.a.	4	n.a.	4	n.a.
5	-1906	5	15	5	-1894	5	n.a.	5	n.a.	5	n.a.
6	-2021	6	4	6	-2020	6	n.a.	6	n.a.	6	n.a.
7	-1930	7	-7	7	-1947	7	n.a.	7	n.a.	7	n.a.
8	-2025	8	-3	8	-2033	8	n.a.	8	n.a.	8	n.a.
9	-2020	9	-39	9	-2068	9	n.a.	9	n.a.	9	n.a.
10	-1969	10	-44	10	-2068	10	n.a.	10	n.a.	10	n.a.
11	-1980	11	-54	11	-2031	11	n.a.	11	n.a.	11	n.a.
12	-1935	12	-73	12	-2009	12	n.a.	12	n.a.	12	n.a.
13	-1887	13	-94	13	-1982	13	n.a.	13	n.a.	13	n.a.
14	-1863	14	-66	14	-1927	14	n.a.	14	n.a.	14	n.a.
15	-1837	15	-45	15	-1884	15	n.a.	15	n.a.	15	n.a.
16	-1717	16	-48	16	-1770	16	n.a.	16	n.a.	16	n.a.
17	-1675	17	-34	17	-1716	17	n.a.	17	n.a.	17	n.a.
18	-1734	18	121	18	-1612	18	n.a.	18	n.a.	18	n.a.
19	-1396	19	-42	19	-1435	19	n.a.	19	n.a.	19	n.a.
20	-1219	20	-55	20	-1275	20	n.a.	20	n.a.	20	n.a.
21	-1172	21	-86	21	-1256	21	n.a.	21	n.a.	21	n.a.
22	-1111	22	-85	22	-1194	22	n.a.	22	n.a.	22	n.a.
23	-1034	23	-77	23	-1115	23	n.a.	23	n.a.	23	n.a.
24	-1001	24	-57	24	-1060	24	n.a.	24	n.a.	24	n.a.
25	-939	25	-61	25	-995	25	n.a.	25	n.a.	25	n.a.
26	-959	26	-37	26	-999	26	n.a.	26	n.a.	26	n.a.
27	-921	27	-18	27	-938	27	n.a.	27	n.a.	27	n.a.
28	-856	28	-12	28	-872	28	n.a.	28	n.a.	28	n.a.
29	-841	29	-42	29	-888	29	n.a.	29	n.a.	29	n.a.
30	-814	30	-94	30	-908	30	n.a.	30	n.a.	30	n.a.
31	-741	31	-100	31	-837	31	n.a.	31	n.a.	31	n.a.
32	-740	32	-83	32	-824	32	n.a.	32	n.a.	32	n.a.
33	-664	33	-43	33	-706	33	n.a.	33	n.a.	33	n.a.
34	-625	34	-13	34	-637	34	218	34	12	34	231
35	-543	35	9	35	-559	35	417	35	5	35	413
36	-493	36	24	36	-469	36	531	36	-4	36	529
37	-428	37	66	37	-362	37	707	37	-11	37	594
38	-296	38	148	38	-150	38	193	38	11	38	203
39	n.a.	39	n.a.	39	n.a.	39	283	39	42	39	326

40	-206	40	102	40	-105	40	-4	40	0	40	-2
41	-328	41	104	41	-194	41	-124	41	-1	41	-12
42	-341	42	115	42	-230	42	-152	42	-10	42	-16
43	-329	43	65	43	-262	43	-230	43	-3	43	-234
44	-306	44	65	44	-240	44	-274	44	-38	44	-318
45	-242	45	57	45	-188	45	-287	45	-47	45	-333
46	-117	46	89	46	-30	46	-211	46	-30	46	-239
47	-179	47	50	47	-128	47	-194	47	-29	47	-227
48	-20	48	-95	48	-96	48	-142	48	-15	48	-158
49	-46	49	78	49	31	49	-97	49	37	49	-61
50	1	50	82	50	82	50	-59	50	-22	50	-90
51	56	51	21	51	76	51	-145	51	-9	51	-153
52	63	52	-6	52	88	52	-153	52	9	52	-145
53	158	53	-18	53	138	53	-214	53	23	53	-184
54	194	54	-6	54	189	54	-205	54	8	54	-196
55	237	55	-1	55	236	55	-35	55	7	55	-27
56	284	56	19	56	304	56	-23	56	1	56	-25
57	413	57	-5	57	407	57	-21	57	1	57	-20
58	510	58	-47	58	463	58	-26	58	9	58	-17
59	616	59	-69	59	554	59	-5	59	13	59	8
60	624	60	53	60	674	60	147	60	1	60	149
61	584	61	44	61	629	61	-90	61	-28	61	-118
62	418	62	149	62	565	62	-239	62	-3	62	-243

**Table 3:** Average Gulfside Erosion Rates for the Caminada-Moreau Headlands and Grand Isle: 1887 – 1996

Years	Average Gulfside Erosion Rates (m/yr)		Caminada Moreau Headland and Grand Isle
	Caminada Moreau Headland	Grand Isle	
Previous Analysis <sup>1</sup>	-13.6	-5.2	-4.2
1978 - 1988 (short-term)			
1887 - 1988 (long-term)	-13.3	0.9	-6.2
New Analysis <sup>2</sup>	-3.2	4.1	0.5
1988 - 1996 (short-term)			
1887 - 1996 (long-term)	-12.6	1.2	-5.7

<sup>1</sup> McBride and others (1992)

<sup>2</sup> This USGS Open-File Report

**Table 4:** Average Bayside Erosion Rates for the Caminada-Moreau Headlands and Grand Isle: 1887 – 1996

Years	Average Bayside Erosion Rates (m/yr)		Caminada Moreau Headland and Grand Isle
	Caminada Moreau Headland (Caminada spit)	Grand Isle	
Previous Analysis <sup>1</sup>	-1.8	3.2	0.7
1978 - 1988 (short-term)			
1887 - 1988 (long-term)	4.1	1.0	2.6
New Analysis <sup>2</sup>	0.0	-0.6	-0.3
1988 - 1996 (short-term)			
1887 - 1996 (long-term)	3.6	-1.0	1.3

<sup>1</sup> McBride and others (1992)

<sup>2</sup> This USGS Open-File Report

**Table 5:** Grand Isle Area Measurements by Island (acres)

	Previous <sup>1</sup>			Update <sup>2</sup>
	1887	1978	Jan 1988	Dec 1996
Grand Isle	2617	2313	2372	2447

<sup>1</sup> McBride and others (1992)

<sup>2</sup> This USGS Open-File Report

**Table 6:** Grand Isle Area Change Rate and Predicted Disappearance Dates: 1887 – 1996<sup>1</sup>

	Long-Term Rate		Short-Term Rate		Previous Long-Term DD <sup>5</sup>	Previous Short-Term DD <sup>6</sup>	New Long-Term DD <sup>3</sup>	New Short-Term DD <sup>4</sup>
	Previous 1887 - 1988 <sup>2</sup>	New 1887 - 1996	Previous 1978 - 1988 <sup>2</sup>	New 1988 - 1996				
Grand Isle	-2.4	-1.6	5.9	8.4	2968	NA <sup>7</sup>	3525	NA <sup>7</sup>

<sup>1</sup> Area Change Rate in acres per year (a/yr)

<sup>2</sup> Data Converted from hectares to acres from McBride and others (1992)

<sup>3</sup> New Long-term disappearance date: 1887 – 1996 [109 yrs]

<sup>4</sup> New Short-term disappearance date: 1988 – 1996 [8.9 yrs]

- <sup>5</sup> Previous Long-Term disappearance date: 1887 – 1988 (McBride and others, 1992)
- <sup>6</sup> Previous Short-Term disappearance date: 1978 – 1988 (McBride and others, 1992)
- <sup>7</sup> Not Applicable because of increase in area