


NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)

Amikacin




↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)

Amoxicillin/Clavulanic Acid




↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)

Ampicillin




↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)

Cefoxitin




↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)

Ceftiofur




↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)


Ceftriaxone



↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)




↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)

Ciprofloxacin




↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)


Gentamicin



↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)




↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)

Nalidixic Acid




↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)

Streptomycin




↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)

Sulfizoxazole




↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)

Tetracycline




↓ Breakpoint

NARMS – EB 2004 Veterinary Isolates

Fig. 31 Minimum Inhibitory Concentrations ($\mu\text{g/ml}$) by Antimicrobial Agent for Major Serotypes from Turkey (Diagnostic)

Trimethoprim/Sulfamethoxazole



↓ Breakpoint