

10017
Ilmenite Basalt (high K)
973 grams



Figure 1: Close-up photo of dusted surface of 10017, 81 N1 face. Sample is 4 cm across. NASA #76-25451.

Introduction

10017 is the largest rock sample returned by Apollo 11. It is a fine-grained, vesicular, ilmenite basalt of the high-K variety (figure 1). The original crystallization age was determined as 3.6 b.y.

10017 was used for the study of solar-ray and cosmic-ray interaction with the lunar surface (Shedlovsky et al. 1970, SHRELLDALFF 1970). Shedlovsky et al. determined the orientation by radiation counting and found that *“most photographs made in the preliminary examination were in what we now consider a position inverted relative to its orientation on the moon.”*

Overall, 10017 was found to have a high level of cosmic-ray-induced radioactivity with an exposure age of 480 m.y.

Photos of 10017 taken before it was broken show that all sides are rounded and covered with micrometeorite craters and patina indicating that this sample “tumbled” while on the lunar surface (figures 2 and 3). However, one side is flatter and turned out to have been the surface that was most recently exposed on the Moon (from ^{56}Co activity). Depth profiles of cosmic-ray-induced nuclear tracks were found on both the top and



Figure 2: Original photo of rounded, dust-covered, surface of 10017,0. NASA S69-47573. Note micrometeorite craters and vesicles. This was originally thought to be the “top” surface until 56Co was found on the other side (figure 3).

bottom sides of the sample. The sample has not always been in the orientation that it was found, and it is not the most suitable sample for the separation of the nuclear effects caused by solar-ray and cosmic-ray radiation.

Petrography

There is a variability to the texture of 10017, even within a given thin section, which is due to apparent mineral segregations and differences in grain size (French et al. 1970, Dence et al. 1970). For this reason, and because petrologists were not familiar with high Ti basalts, a wide variety of descriptions and names were applied to Apollo 11 basalts. All investigators were surprised by the lack of any alteration. Mesostasis is glassy.

Schmitt et al. (1970) originally described 10017 as a “poikiloblastic olivine basalt (very fine-grained, vesicular, vuggy, glass-bearing).” James and Jackson (1970) listed

10017 as “hornfelsic, intersertal basalt” where ilmenite and pyroxene “occur in mosaics of equant interlocking grains and plagioclase forms coarse poikiloblastic crystals” (figure 4). Dence et al. (1970) describe the texture of 10017 as “intergranular”, Brown et al. (1970) describe the texture as “hypidiomorphic granular” and McGee et al. (1977) described the texture as “intersertal.”

Adler et al. (1970), French et al. (1970), Brown et al. (1970), Kushiro and Nakamura (1970), Dence et al. (1970), McGee et al. (1977), Beatty and Albee (1978) and others have provided detailed petrographic descriptions of 10017. 10017 is a fine- to medium-grained basalt composed primarily of clinopyroxene, plagioclase and ilmenite, with minor mesostasis including high-K glass. Plagioclase megacrysts are typically poikilitic and enclose anhedral pyroxene and ilmenite crystals. The rock is vesicular, with both circular and irregular vesicles about 0.2 to 0.5 mm in size, which make up only a few percent of the sample.




Figure 3: Original photo taken in F-201 of “flatter” side of 10017,0 showing numerous, well-developed micrometeorite craters. This side was eventually determined to be the most recent top surface, because it had abundant short-lived radionuclides (see text). NASA S69-47560. Sample is 16 cm long.

In 10017, pyroxenes are often granular, whereas plagioclase is anhedral to subhedral (Kushiro and Nakamura 1970). The grain size is variable with pyroxene and ilmenite from 50 to 150 microns, but plagioclase is up to 2 - 3 mm in length (French et al. 1970). This high Ti, lunar basalt is unusual in that the ilmenite is equant rather than tabular.

Thin sections of 10017 exhibit regions with mineralogical segregations and differences in grain sizes. Plagioclase forms concentrations of large lath-shaped crystals, while adjacent areas are enriched in smaller grains of pyroxene and ilmenite (see figure 1 in French et al. 1970).

Mineralogical Mode for 10017

	Beaty and Albee 1978	French et al. 1970	Brown et al. 1970	Kushiro and Nakamura 70	James and Jackson 70
Olivine	absent				
Pyroxene	50.6 vol. %	49.7	59.4	51	47.6
Plagioclase	23.6	18	25.1	21.5	26.9
Ilmenite	15.1	23.9	14.5	20.2	15
mesostasis	8.5	8.3		6.1	8.5
silica	1.4		tr	1.1	1.6
troilite	0.5		0.36		
phosphate	0.2				




Figure 4: Thin section photomicrographs of 10017,20 (reflected, transmitted and crossed-Nicols of same area). Scale is 2.5 mm across. NASA S79-27113 - 27115.

Mineralogy

Olivine: none

Pyroxene: The chemical compositions of fine-grained pyroxenes in 10017 were determined by French et al. (1970), Brown et al. (1970), Dence et al. (1970) and Beatty and Albee (1978) (figure 5).




Figure 5: Pyroxene composition of 10017 (note lack of extreme iron enrichment).

Plagioclase: The composition of plagioclase in 10017 is $An_{78}Ab_{20}Or_2$.

Ilmenite: Ilmenite in 10017 occurs as equant grains (French et al. 1970).

Apatite: French et al. give an analysis of fluoroapatite ($F = 3.1\%$).

Silica: Kushiro and Nakamura (1970) report relatively large cristobalite crystals in 10017.

Mesostasis: French et al. (1970) and Dence et al. (1970) report K-rich mesostasis.

Chemistry

Compston et al. (1970), Gast et al. (1970) and others observed that Apollo 11 basalts could be divided in two groups based on the higher K and Rb contents of Group 1 as compared with that of Group 2 (low-K). 10017 belongs to the high-K group and also has higher REE content (figure 6). Anders et al. (1971) found that the meteoritic siderophiles (Ir and Au) were low in this igneous rock. Gibson et al. (1970) determined the gas released from 10017 (figure 11).

Radiogenic age dating

Papanastassiou et al. (1970), DeLaeter et al. (1973) and Gopalan et al. (1970) determined the Rb/Sr isochron age for 10017 (figures 8, 9). Turner (1970) determined the Ar/Ar age. Eberhardt et al. (1974) determined 2.35 b.y. by whole rock K/Ar.




Figure 6: Normalized rare-earth-element composition for high-K basalt 10017 (the line) compared with that of low-K basalt 10020 and high-K basalt 10049 (the dots) (data from Wiesmann et al. 1975).




Figure 7: Composition of 10017 compared with that of other lunar basalts.

Cosmogenic isotopes and exposure ages

Perkins et al. (1970) determined the cosmic ray induced activity of ^{22}Na (45 dpm/kg), ^{26}Al (80 dpm/kg), ^{46}Sc (13 dpm/kg) and ^{54}Mn (46 dpm/kg). O'Kelley et al. (1970) determined of ^{22}Na (39 dpm/kg), ^{26}Al (73 dpm/kg), ^{46}Sc (13 dpm/kg) and ^{54}Mn (33 dpm/kg) and ^{56}Co (26 dpm/kg). Wrigley and Quaide (1970) also determined ^{22}Na (47 dpm/kg) and ^{26}Al (83 dpm/kg) in general agreement if one considers the counting errors (see references).

Eberhardt et al. (1970) observed that high-K basalts had distinctly different exposure ages from low-K Apollo 11 basalts. Eberhardt et al. (1974) determined the cosmic ray exposure age as 480 ± 25 m.y. by the reliable ^{83}Kr method and 371 m.y. by ^{126}Xe (as calculated by Srinivasan 1974) while Marti et al. (1970) determined 509 ± 29 m.y. by ^{83}Kr and 308 m.y. by ^{126}Xe (calculated). Turner (1970) and Hintenberger et al. (1971) reported exposure ages of 440 and 510 m.y. respectively by ^{38}Ar .

Shedlovsky et al. (1970) and O'Kelley et al. (1970) reported significant ^{56}Co (77 days) on the "top" surface of 10017 (125 and 44 dpm/kg respectively), thus establishing its most recent lunar surface orientation.

Other Studies

A detailed cosmic ray depth profile was attempted by Shredldaff (1970), but only ^{22}Na showed significant variation with depth (table 3). The problem is that this sample was irradiated on both sides, and it was not thick enough to stop high-energy cosmic rays. Additionally, the analytical techniques in 1969 required

large subsamples. 10017 was the first of these studies, later carried out with more success on 14310, 68815 and 74275.

Fleischer et al. (1970), Lal et al. (1970) and Crozaz et al. (1970) studied solar flare and cosmic ray tracks as a function of depth in 10017 (figures 15 and 16). The near surface regions were dominated by solar flare tracks and the erosion rate from micrometeorite bombardment could be determined.

Oxygen isotopes were reported for mineral separates of 10017 by Onuma et al. (1970) and Taylor and Epstein (1970).

O'Hara et al. (1970) and Ringwood and Essene (1970) experimentally determined the phase diagram for 10017 (figures 12 and 13). These experiments showed that Apollo 11 basalts probably formed at depths of 200 to 400 km by a small degree of partial melting from pyroxenitic source material.

Eugster et al. (1970) reported a significant isotopic anomaly in ^{158}Gd proving that 10017 had received a high dosage of low energy neutrons over a long period of time (burial?).

Bogard et al. (1971) and Eberhardt et al. (1974) reported the abundance and isotopic ratios of noble gases in 10017.

The elastic properties for a lunar basalt (compressibility, sound velocity) were determined on 10017 by Anderson et al. (1970). The magnetic




Figure 8: Rb/Sr isochron for 10017 (from Gopalan et al. 1970).

properties of 10017 were determined by Runcorn et al. (1970) and the remanent magnetization was found to be weak.

Processing

10017 was returned in ALSRC #1004 and originally processed in the "Vac Lab". It was one of the rocks in the F-201 at the time of the glove rupture (with exposure to Houston air).

10017 originally broke in two large pieces (.5 and .6) along a penetrating fracture (figure 19). Surface pieces .74 and .42 from opposite ends of this fractured surface were allocated to Arnold.

Shredliff (1970) illustrated the cutting plan as it pertained to their depth profile studies (figure 14). A cube (.180) and a column (.64) were cut from the sample by sawing (figure 18). *All the King's men couldn't put 10017 together again!*




Figure 9: Rb/Sr isochron for 10017 (from Papanastassiou and Wasserburg 1970).




Figure 10: Original PET photo of side of 10017,0 in F-201. The sample is sitting on its top surface (flatter side). NASA S69-45375.

Summary of Age Data for 10017

	Rb/Sr	Nd/Sm	Ar/Ar	K/Ar
Papanastassiou et al. 1970	3.59 ± 0.05 b.y.			
DeLaeter et al. 1973	3.71 ± 0.11			
Gopalan et al. 1970		3.575 ± 0.215		
Turner 1970			3.23 ± 0.06	
Eberhardt et al. 1974				2.35 ± 0.06 whole rock

Caution: These ages have not been corrected for new decay constants.

Table 1a. Chemical composition of 10017.

reference weight	LSPET69	Maxwell70		Maxwell70	Gast70	Wiesmann75	Perkins70	Wrigley70	O'Kelley70
		GSC	GSF	Abbey		187 mg	115 g	115 g	971 g
SiO2 %	40	(a) 40.78	40.77	(b) 40.14	(c)				
TiO2	11	(a) 11.71	11.82	(b) 11.16	(c)				
Al2O3	10	(a) 8.12	7.92	(b) 8.17	(c)				
FeO	19	(a) 19.82	19.79	(b) 19.38	(c)				
MnO	0.35	(a) 0.22	0.22	(b) 0.25	(c)				
MgO	8.5	(a) 7.65	7.74	(b) 7.72	(c)				
CaO	10	(a) 10.55	10.58	(b) 10.99	(c)				
Na2O	0.65	(a) 0.51	0.51	(b) 0.51	(c) 0.51	(e) 0.51	(e)		
K2O	0.22	(a) 0.3	0.29	(b) 0.3	(c) 0.31	(d) 0.31	(d) 0.31	(h)	0.29 (h)
P2O5			0.13	(b) 0.15	(c)				
S %			0.22	(b)					
sum									
Sc ppm	55	(a) 52	77	(b)					
V	30	(a) 49	70	(b)					
Cr	4600	(a) 2200	2260	(b) 2463	(c)				
Co	10	(a) 20	30	(b)					
Ni									
Cu			20	(b)					
Zn		49	47	(b)					
Ga									
Ge ppb									
As									
Se									
Rb	6	(a)			5.63 (d)	5.63 (d)			
Sr	55	(a)			175 (d)	175 (d)			
Y	310	(a) 160	160	(b)					
Zr	1250	(a) 410	430	(b)					
Nb									
Mo									
Ru									
Rh									
Pd ppb									
Ag ppb									
Cd ppb									
In ppb									
Sn ppb									
Sb ppb									
Te ppb									
Cs ppm					0.16 (d)	0.155 (d)			
Ba	120	(a)			309 (d)	309 (d)			
La					26.6 (d)	26.6 (d)			
Ce					77.3 (d)	77.3 (d)			
Pr									
Nd					59.5 (d)	59.5 (d)			
Sm			25	(b)	20.9 (d)	20.9 (d)			
Eu			2.1	(b)	2.14 (d)	2.14 (d)			
Gd					27.4 (d)	27.4 (d)			
Tb			4	(b)					
Dy					31.7 (d)	31.7 (d)			
Ho									
Er					20 (d)	20 (d)			
Tm									
Yb			19	(b)	19.2 (d)	17.1 (d)			
Lu			5	(b)	2.66 (d)	2.66 (d)			
Hf			12	(b)					
Ta			5	(b)					
W ppb									
Re ppb									
Os ppb									
Ir ppb									
Pt ppb									
Au ppb									
Th ppm			4.6	(b)			3.2 (h)	2.9 (h)	3.25 (h)
U ppm							0.86 (h)	0.9 (h)	0.83 (h)

technique: (a) OES, (b) mixed, (c) atomic absorption spec., colorimetry, (d) idms, (e) AA, (f) wet, (g) XRF, (h) radiation counting

Table 1b. Chemical composition of 10017.

reference weight	Tera70	Wanke70	Dickinson89	Wiik70	Philpotts69 Philpotts70 125 mg	Haramura in Kushiro 70	Compston70	Wakita70 637 mg	343 mg
SiO2 %		41.9		40.77	(b)	40.03	(f) 40.69	(g) 42.8	44.5 (a)
TiO2		11.7		11.82	(b)	11.61	(f) 11.92	(g)	13 (a)
Al2O3		8.3		7.92	(b)	8.04	(f) 7.78	(g) 7.94	7.6 (a)
FeO		18.8		19.79	(b)	20.46	(f) 19.59	(g) 21.9	23.3 (a)
MnO		0.19		0.22	(b)	0.24	(f) 0.28	(g) 0.24	0.25 (a)
MgO		7.96		7.74	(b)	8.17	(f) 7.51	(g) 7.96	7.63 (a)
CaO	10.3	(d) 11.5	8.8	13.4	10.58	(b) 10.26	(f) 10.76	(g)	10.6 (a)
Na2O	0.47	0.42	0.4	0.43	0.51	(b) 0.49	(f) 0.51	0.49	0.5 (a)
K2O	0.29	(d) 0.25		0.29	(b) 0.29	(d) 0.3	(f) 0.3	(g) 0.19	0.26 (a)
P2O5				0.18	(b)	0.17	(f) 0.18	(g)	
S %							0.23	(g)	
sum									
Sc ppm		86	(a) 73	80	(a) 77	(b)		87	93 (a)
V				820	(a) 70	(b)		49	43 (a)
Cr		2310	(a) 1500	2600	(a) 2258	(b) 2326	(f)	2450	2710 (a)
Co		24.5	(a) 24	26	(a) 30	(b)		35	37 (a)
Ni					<5	(b)			
Cu		7.7	(a)		20	(b)			
Zn					47	(b)			
Ga		4.2	(a)	56	(a)				
Ge ppb			5.5	6	(a)				
As									
Se									
Rb	5.57	(d) 4.2	(a)			5.7 (d)	5.72	(g) 6	(a)
Sr	156	(d) 151	(a) 151			165 (d)	164	(g)	
Y				160	(b)		159	(g) 184	(a)
Zr			418	430	(b)		476	(g) 1200	1570 (a)
Nb									
Mo									
Ru									
Rh									
Pd ppb		1	(a)						
Ag ppb									
Cd ppb								44	(a)
In ppb		138	(a)					1	(a)
Sn ppb									
Sb ppb									
Te ppb									
Cs ppm	0.159	(d) 0.12	(a)					0.26	(a)
Ba	280	(d) 256	(a) 254	258	(a) 287	(d)		350	410 (a)
La		21	(a) 22.1	23.8	(a)			24	28 (a)
Ce		64	(a) 62	62	(a) 75.5	(d)		84	(a)
Pr		7.3	(a)					12.9	(a)
Nd		58	(a) 56	54	(a) 66.1	(d)		74	(a)
Sm		18	(a) 15	19	(a) 25	(b) 23.4	(d)	23	27 (a)
Eu		1.89	(a) 1.9	2.1	(a) 2.1	(b) 2.26	(d)	2.3	2.9 (a)
Gd		19	(a)			28.4	(d)	30	(a)
Tb		4.6	(a) 4.7	5	(a) 4	(b)		5.5	(a)
Dy		19	(a)			32.8	(d)	36	(a)
Ho		3.8	(a)					8.3	(a)
Er		13	(a)	11.5	(a) 19.1	(d)		21	(a)
Tm				2	(a)			3	(a)
Yb		15.6	(a) 15.8	16.4	(a) 19	(b) 18.1	(d)	17.8	23 (a)
Lu		2.12	(a) 2.4	2.7	(a) 5	(b) 2.63	(d)	2.5	3 (a)
Hf		16.5	(a) 13.2	14.9	(a) 12	(b)		23	26 (a)
Ta		2.2	(a) 1.9	2.1	(a) 5	(b)			
W ppb		0.4	(a)						
Re ppb									
Os ppb									
Ir ppb									
Pt ppb									
Au ppb		8.1	(a)						
Th ppm		3.94	(a) 2.9	3.1	(a) 4.6	(b)	3.7	(g) 4.3	6.4 (a)
U ppm		0.69	(a)						

technique: (a) INAA & RNAA, (b) mixed, (c) atomic absorption spec., colorimetry, (d) idms, (e) AA, (f) wet, (g) XRF

Table 1c. Chemical composition of 10017.

reference	Anders71	
weight		
SiO2 %		
TiO2		
Al2O3		
FeO		
MnO		
MgO		
CaO		
Na2O		
K2O		
P2O5		
S %		
sum		
Sc ppm		
V		
Cr		
Co	31	(a)
Ni		
Cu		
Zn	18	(a)
Ga	5.1	(a)
Ge ppb		
As		
Se	0.215	(a)
Rb	6.6	(a)
Sr		
Y		
Zr		
Nb		
Mo		
Ru		
Rh		
Pd ppb		
Ag ppb	16	(a)
Cd ppb	68	(a)
In ppb	70	(a)
Sn ppb		
Sb ppb		
Te ppb		
Cs ppm	0.186	(a)
Ba		
La		
Ce		
Pr		
Nd		
Sm		
Eu		
Gd		
Tb		
Dy		
Ho		
Er		
Tm		
Yb		
Lu		
Hf		
Ta		
W ppb		
Re ppb		
Os ppb		
Ir ppb	0.02	(a)
Pt ppb		
Au ppb	0.72	(a)
Th ppm		
U ppm		
technique:	(a) RNAA	




Figure 11: Gas released as function of increasing temperature for 10017 (Gibson et al. 1971).




Figure 12: Phase diagram determined for 10017 (and 10084?) by O'Hara et al. 1970.




Figure 13: Phase diagram for Apollo 11 basalt from Ringwood and Essene 1970.




Figure 17 : Group photo of 10017. NASA S75-30215. Cube is 1 cm.




Figure 18: Group photo of 10017. NASA S76-21150. Cube is 1 cm.




Figure 19: Photo showing original break of 10017. S69-49220.



Figure 20: Photo of display sample 10017,37 showing micrometeorite craters and vesicles. NASA photo #S84-40064.

List of Photo #s for 10017

- S69-45375 – 45379 PET B&W
- S69-47560
- S69-47575
- S69-49221 – 222
- S69-49235 – 236
- S69-49243 – 244
- S69-53982 TS
- S69-54058 TS
- S70-48930 – 931 color TS
- S70-49222 – 223
- S70-49230 – 231
- S70-49868 – 869
- S70-49872 – 873
- S70-49878 – 879
- S70-49973 – 974
- S76-26302 – 303 reflected
- S76-20792
- S76-20805
- S78-27113 – 115 fig 3
- S75-30209 – 215 processing
- S76-21149 – 150 processing
- S76-25453 – 458 ,81
- S84-40064 – 469 ,37

