

Table 5. Coretop counting categories

Taxon	Comments
<i>Candeina nitida</i> d'Orbigny	
<i>Globierinella aequilateralis</i> (Brady)	
<i>Globigerina bulloides</i> d'Orbigny	
<i>Globigerina calida</i> (Parker)	
<i>Globigerina conglomerata</i> Schwager	
<i>Globigerina digitata</i> Brady	
<i>Globigerina falconensis</i> Blow	
<i>Globigerina rubescens</i> Hofker	
<i>Globigerinita glutinata</i> (Egger)	
<i>Globigerinoides conglobatus</i> (Brady)	
<i>Globigerinoides ruber</i> (d'Orbigny)	pink and white varieties combined with and without sacklike final chamber
<i>Globigerinoides sacculifer</i> (Brady)	
<i>Globigerinoides tenellus</i> Parker	
<i>Globorotalia hirsuta</i> (d'Orbigny)	
<i>Globorotalia crassaformis</i> (Galloway and Wissler)	
<i>Globorotalia inflata</i> (d'Orbigny)	
<i>Globorotalia menardii</i> (Parker, Jones, and Brady)	including all subspecies of <i>G. tumida</i>
<i>Globorotalia pumilio</i> Parker	
<i>Globorotalia scitula</i> (Brady)	
<i>Globorotalia truncatulinoides</i> (d'Orbigny)	
<i>Globorotaloides hexagona</i> (Natland)	
<i>Neogloboquadrina dutertrei</i> (d'Orbigny)	
<i>Neogloboquadrina pachyderma</i> (Ehrenberg)	sinistral coiling dextral coiling includes 4.5 chambered dextral forms transitional between <i>N. pachyderma</i> and <i>N. dutertrei</i>
<i>Neogloboquadrina pachyderma</i> (Ehrenberg)	
"dupac"	
<i>Orbulina universa</i> d'Orbigny	
<i>Pulleniatina obliquiloculata</i> (Parker and Jones)	
<i>Turborotalita quinqueloba</i> (Natland)	