

Prevalence and Consequences of Fluoroquinolone-Resistant *Campylobacter* Infections: NARMS 1997-2000

J. McClellan, S. Rossiter, K.
Joyce, K. Stamey, A.D. Anderson,
and the NARMS Working Group

Centers for Disease Control and Prevention



Campylobacter

- In the United States,
 - Approximately 2.4 million infections annually
 - 150,000 physician visits
 - 13,000 hospitalizations
 - 100 deaths
 - Mild, moderate, severe infections
 - Antimicrobials usually not essential, but may be life-saving
 - Chickens and turkeys are primary reservoir

Fluoroquinolones (FQs)

- Use in humans:
 - Approved in 1986
 - Most commonly used antimicrobials for *Campylobacter* infections
 - Reduce severity and duration of symptoms associated with infection
- Use in animals:
 - Chickens and turkeys
 - Approved in 1995
 - Added to drinking water

Human FQ-resistant *Campylobacter* infections

- CDC survey:
 - 1990
 - No FQ-resistant *Campylobacter*
- MN study
 - 1996-1998
 - Increase from 0.8% to 3%

What is the current prevalence of FQ-resistant *Campylobacter* in humans in the United States?



NARMS

- National Antimicrobial Resistance Monitoring System
 - Receives 1 *Campylobacter* isolate per week from 9 NARMS sites
 - *Campylobacter* isolates speciated and susceptibility tested
 - Ciprofloxacin MIC ≥ 4 $\mu\text{g/ml}$ were considered resistant

NARMS

Sites

Connecticut

Georgia

Minnesota

Oregon


California

Colorado


Maryland

New York

Tennessee



Human *Campylobacter* isolates, 1997-2000 (N=1205)



What are the clinical consequences associated with FQ-resistant *Campylobacter* infections?



Steps

- Step One:
 - How many people took FQs for their illness?
- Step Two:
 - Among FQ users, difference in diarrhea duration between FQ-resistant and FQ-susceptible infections?

Epidemiological Study

- FoodNet/NARMS (Jan 1998 – Jan 1999)
 - 858 culture-confirmed *Campylobacter* infections
 - Isolates tested at CDC, CT, MN, or NY
- Questionnaire
 - Antimicrobial use (including FQs)
 - Antidiarrheal use
 - Duration of diarrhea



Medications

- 54% (326/609) of culture-confirmed patients took FQs
 - 39% (128/326) FQ use and no other medications
- Wide use of FQs and high prevalence of FQ-resistant infections
 - Differences in outcome?

Duration of Diarrhea (N=128)

- Average duration of diarrhea
 - FQ users:
 - FQ-resistant (N=17)
 - 8 days (range: 2-14 days)
 - FQ-susceptible (N=111)
 - 6 days (range: 2-31 days)
- People with FQ-resistant infections, who are treated with FQs, had 2 days longer duration, $p=0.08$

Intriguing Finding

- Non-treated patients (N=67)
 - FQ-resistant (N=6):
 - 12 days (range: 8-20 days)
 - FQ-susceptible (N=61):
 - 6 days (range: 2-21 days)
- Among non-treated, FQ-resistance may be associated with longer duration of diarrhea, $p < 0.01$

Multivariate Analysis

- N=858
 - Controlling for factors potentially associated with duration of diarrhea
 - FQ-resistant infections associated with 1 day longer duration of diarrhea, $p=0.05$

Summary

- FQ-resistance is prevalent
 - 14% in 2000
- Over 50% of culture-confirmed *Campylobacter* infections receive FQs
- FQ-resistance associated with longer duration of diarrhea
- Association was significant, after adjusting for covariates
- FQ-resistant infections have an adverse health consequence
 - A longer duration of diarrhea

Other Investigations and Actions

- Investigations:
 - Additional FoodNet *Campylobacter* case-control study
 - Eating poultry outside home was a risk factor for domestic acquired FQ-resistant infections
 - FDA quantitative risk assessment
 - FQ use in chickens and turkeys resulted in FQ-resistant *Campylobacter* infections each year in humans
- Actions:
 - Proposed FQ withdrawal
 - Supported by CDC, ASM, and others
 - Opposed by a pharmaceutical company, AVMA, and others



Acknowledgments

CDC

- Fred Angulo
- Alicia Anderson
- Felicia Hardnett
- Mike Hoekstra
- Kevin Joyce
- Nina Marano
- Shannon Rossiter
- Karen Stamey
- California Department of Health Services
- Connecticut Department of Public Health
- Georgia Division of Public Health
- Maryland Department of Health and Mental Hygiene
- Minnesota Department of Health
- New York State Department of Health
- Oregon Department of Human Resources

NARMS website: www.cdc.gov/narms

